



Contents

Chromatography DC

| SAFETY IN THE LABORATORY | | ORGANIC SOLVENTS | | |
|--|--|---|----------------------------------|---|
| Poisons information centres Proper conduct in the laboratory Notes on the handling of ethers and other peroxide-forming solvents Risk phrases Safety phrases Hazard symbols Incompatible chemicals | p. 4 p. 8 p. 10 p. 12 p. 18 p. 24 p. 26 | Organic solvents - Properties and drying Ethanol-water mixtures Drying agents Vapour pressure of water LiChrosolv - Solvents for chromatography | p. p. p. p. | 83 86 87 87 88 |
| | | PHYSICAL METHODS FOR THE DETERMINATION OF ELEMENTS | | |
| CHEMICAL AND PHYSICAL PROPERTIES OF ELEMENTS AND INORGANIC COMPOUNDS Hardness scale according to MOHS Electrochemical series of the elements Table of elements Electrochemical series of some nonmetals Covalent single-bond radiuses | p. 29 p. 29 p. 30 p. 34 p. 34 | Flame photometry Wavelength and wave number Photometry: Transmission rate and absorbance Calculation of the standard deviation Direct-current polarography Cathode ray polarography | p. p. p. p. p. | 91 91 91 91 92 93 |
| | | MASS AND WEIGHT | | |
| General mixing formulas for liquids Conversion table for water hardness Mixture rule Preparation of dilute solutions Solubility of inorganic compounds in water in relation to temperature Solubility products of slightly soluble inorganic compounds Sample preparation Acids: Sulfuric acid | p. 37 p. 38 p. 38 p. 39 p. 40 p. 48 p. 51 p. 52 | Energy dimensions - conversion factors Pressure dimensions - conversion factors Decimal multiples and parts of units Concentration values Basic units Derived units American and English measures | p. p. p. p. p. p. | 95 96 96 96 97 102 |
| Phosphoric acid Hydrochloric acid Nitric acid Sodium hydroxide solution Potassium hydroxide solution Ammonia Commercially available concentrations of some acids and alkalis | p. 54 p. 55 p. 56 p. 57 p. 59 p. 61 | OTHER USEFUL TABLES Physical constants Establishment of constant humidity in closed vessels Greek alphabet Greek numerals / Roman numbers Freezing mixtures Extran Particle sizes NMR: Carbon (13 C) chemical shifts NMR: Proton chemical shifts | p. p. p. p. p. p. | 105 106 107 108 109 110 112 114 118 |
| INDICATORS AND BUFFERS | | Miscibility table Heating baths | p. | 122 124 |
| PH indicators Acid-base indicators Redox indicators Substances for the preparation of buffer solutions Buffer solutions Chromatography | p. 65 p. 66 p. 71 p. 73 p. 74 p. 80 | reduing battle | р. | 124 |

p. 81



POISONS INFORMATION CENTRES

Several cities of the Federal Republic of Germany have poisons information centres. These centres, open to anybody, are on call day and night to give information over the phone about countermeasures to be taken in intoxication accidents of any type.

| | body, are on call day and night to give information over easures to be taken in intoxication accidents of any type. |
|----------------|--|
| Austria | Vergiftungsinformationszentrale · Allgemeines Krankenhaus Währinger Gürtel 18-20 · 1090 Wien Telephone: (+43) (0)1 4 06 43 43 Languages: German, English, (French) |
| Belgium | Centre Antipoisons c/o Hôpital Militaire Reine Astrid Rue Bruyn · 1120 Bruxelles Telephone: (+32) (070) 24 52 45 Languages: French, Flemish, English, (German) |
| Bulgaria | National Center of Clinical Toxicology Emergency Medical Institute "Pirogov" · 1606 Sofia Telephone: (+359) (0)2 5 15 32 34 / 34 09 / 35 46 Languages: Bulgarian, English, German, (French) |
| Czech Republic | Poison Information Centre Clinic for Occupational Diseases Vysehradska 49 · 12821 Prague 2 Telephone/Telefax: (+42) (0)2 29 38 68 Languages: Czech, English, German |
| Denmark | Poisons Information Centre Rigshospitalet, 7122 Tagensvej 20 · 2200 Copenhagen Ø Telephone: (+45) 35 45 35 45 Telefax: (+45) 35 45 73 49 Languages: Danish, English, (German) |
| Denmark | Arbejdsmiljøinstituttet · Produktregistret Postbox 850 Lers Parkalle 105 · 2100 Copenhagen Ø Telephone: (+45) 39 29 97 11 Telefax: (+45) 39 29 97 12 Languages: Danish |
| Finland | Poison Information Centre Department of Clinical Pharmacology University Central Hospital Haartmaninkatu 4 · 00290 Helsinki Telephone: (+358) (0)9 41 43 92 (+358) (0)9 47 11/27 88 Telefax: (+358) (0)9 47 14 70 2 Languages: Finnish, Swedish, English |
| France | Centre Anti-Poisons de Lille · C.H.R. 5 Avenue Oscar Lambret · 59037 Lille Cedex Telephone: (+33) 03 20 44 44 44 Telefax: (+33) 03 20 44 56 28 Languages: French |
| France | Centre Anti-Poisons · Service de Pharmaco Toxicovigilance et Centre Anti-Poisons · Hópital Eduoard Herriot 5 Place d'Arsonval · 69003 Lyon Telephone: (+33) (0)72 11 69 11 Telefax: (+33) (0)72 11 69 85 Languages: French |
| France | Centre Anti-Poisons de Marseille Hôpital Salvator 249, bd Ste-Marguerite · 13009 Marseille Telephone: (+33) 91/75 25 25 Telefax: (+33) 91/74 41 68 Languages: French, English, (Italian), (Spanish) |

| rrance | 29 Avenue Maréchal-de-Lattre-de-Tassigny 54035 Nancy Cedex Telephone: (+33) 83 85 20 28 Telefax: (+33) 83 85 20 15 Languages: French |
|-------------|---|
| Greece | Poison Information Centre Childrens Hospital "Aglaia Kyriakou" · 11527 Athens Telephone: (+30) (0)1 7 79 37 77 Telefax: (+30) (0)1 7 48 61 14 Languages: Greek, English |
| Hungary | Department of Clinical Toxicology Hospital Elizabeth Alsoerdosor 7 - Budapest VII Telephone: (+36) (0)1 21 52 15 Telefax: (+36) (0)1 22 94 60 Languages: Hungarian, German, English |
| Ireland | Poisons Information Centre Royal Victoria Hospital Grosvenor Road Belfast BT 12 6BA Northern Ireland Telephone: (+44) (0) 12 32 24 05 03 Telefax: (+44) (0) 12 32 24 80 30 Languages: English |
| Ireland | Poisons Information Centre Beaumont Hospital P.O. Box 1297 Beaumont Road · Dublin 9 Republic of Ireland Telephone: (+353) 0 10 35 31/8 37 99 64 Telefax: (+353) 0 10 35 31/8 37 69 82 Languages: English |
| Italy | Centro Antiveleni Ospedale Niguarda "Ca Grande" Piazza Ospedale Maggiore 3 · 20162 Milano Telephone: (+39) (02) 66 10 10 29 Telefax: (+39) (02) 64 44 27 68 Languages: Italian, (French), (English) |
| Italy | Centro Antiveleni Università di Floma Policlinico Umberto I Viale del Policlinico 155 · 00161 Roma Telephone: (+39) (06) 49 06 63 Telefax: (+39) (06) 49 97 06 98 Languages: Italian, French, English |
| Italy | Centro Antiveleni Università di Torino Corso A.M Dogliotti, 14 · 10126 Torino Telephone: (+39) (011) 6 63 76 37 Languages: Italian, French |
| Netherlands | Nationaal Vergiftingen Informatie Centrum Rijksinstiuut voor Volksgezondheid en Milieuhygiene Antonie van Leewenhoeklaan 9 Postbus 1 · 3720 BA Blithoven Telephone: (+31) (0)30 2 54 15 11 Telefax: (+31) (0)30 2 74 88 88 Languages: Dutch, (French), (English), (German) |

Centre Anti-Poisons de Nancy · Hôpital Central

POISONS INFORMATION CENTRES

| POISONS INF | ORIVIATION CENTRES | | |
|--------------------|---|-------------------------|---|
| Norway | Giftinformasjonssentralen National Poison Information Centre P.O. Box 8189 Dep. · 0034 Oslo 1 Telephone: (+47) (0)22 59 13 00 Telefax: (+47) (0)22 60 85 75 Languages: Norwegian, English | Spain | Instituto Nacional de Toxicologia Servicio de Informacion Toxicoligica ctra. San Jeronimo, Km 1,4 P.O. Box 663 • 41800 Sevilla Telephone: (+34) (5)/4 37 12 33 Telefax: (+34) (5)/4 37 02 62 Languages: Spanish |
| Poland | National Poison Information Centre and Clinical Department of Toxicology Institute of Occupational Medicine ul. Sw. Teresy B, P.O. Box 199 90-950 Lodz Telephone: (+48) (0)42 57 99 00 (+48) (0)42 57 42 95/ (+48) (0)42 31 47 24/25/52/67 Telefax: 448) (0)42 34 83 31/55 61 02 Languages: Polish, English, German | Schweden United Kingdom | Giftinformationscentralen Swedish Poisons Information Centre 17176 Stockholm Telephone: (+46) (0)8 7 38 03 84 Telefax: (+46) (0)8 32 75 84 Languages: Swedish, English, (German) England |
| Poland | Posion Centre of Warsaw PL Weteranas 4 · 03-401 Warszawa Telephone: (+48) (0)28 19 08 97 | United Kingdom | National Poisons Information Service West Midlands Poisons Unit City Hospital NHS Trust Dudley Road · Birmingham B18 7QH Telephone: (444) (0)1 21 5 07 55 88/9 Telefax: (444) (0)1 21 5 07 55 80 Languages: English England Leeds Poisons Information Service Pharmacy Department |
| Portugal | Centro de Informacao Antivenenos Instituto Nacional de Emergencia Medica Rua Infante D. Pedro, 8 · 1799 Lisboa Cedex Telephone: (+351) (0)21 7 95 01 43 (+351) (0)21 7 95 01 44 (+351) (0)21 7 95 01 46 Telefax: (+351) (0)21 7 93 71 24 Languages: Portuguese, French, (English) | United Kingdom | The General Hospital Great George Street · Leeds LS1 3EX Telephone: (+44) (0)1 13 2 43 07 15 Telefax: (+44) (0)1 13 2 44 58 49 Languages: English England National Poisons Information Service (L) Medical Toxicology Unit |
| Rumania | Centre anti-Poisons National Institute of Legal Medicine: Sos. Vitan Birzesti 9, Sector 4 · 75669 Bucharest Telephone: (+40) (0)1 6 34 38 90/1 35 Telefax: (+40) (0)1 32 10 260 Languages: Rumanian, French, English, German | 11 St. 1 MO. | Avonley Road London SE14 5ER Telephone: (+44) (01 71) 6 35 91 91 Telefax: (+44) (01 71) 7 71 53 09 Languages: English |
| Schweiz | Schweizerisches Toxikologisches Informationszentrum Freiestr. 16 · CH-8028 Zürich Telephone: (+41) (0)1 2 51 66 66 Telefax: (+41) (0)1 2 52 88 33 Languages: German, (English) | United Kingdom | England Regional Drugs and Therapeutics Centre Wolfson Unit Claremont Place · Newcastle-upon-Tyne Telephone: (+44) (0)1 91 2 32 15 25 (direct line, normal office hours) (+44) (0)1 91 2 32 25 13 1 |
| Slovakian Republic | Poison Information Centre Clinic of Occupational Diseases and Toxicology Dumbierska 3 · 83101 Bratislava Telephone: (+42) (7) 37 41 66 Telefax: (+42) (7) 37 46 05 Languages: Slovakian, German, English | United Kingdom | (24-hour service) Telefax: (+44) (0) 1 91 2 61 57 33 Languages: English Scotland Scottish Poisons Information Bureau The Royal Infirmary |
| Spain | Instituto Nacional de Toxicologia Servicio Nacional de Informacion Toxicoligica c/Mercé, 1 · 08002 Barcelona Telephone: (+34) (3) 3 17 44 00 Telefax: (+34) (3) 3 18 25 30 Languages: Spanish | United Kingdom | Lauriston Place · Edinburgh EH3 9YW Telephone: (+44) (0)1 31 5 36 22 98 Telefax: (+44) (0)1 31 5 36 23 04 Languages: English Wales Welsh National Poisons Unit |
| Spain | Servicio Nacional de Informacion Toxicolígica · Instituto Nacional de Toxicologia c/Luis Cabrera, 9 · 28002 Madrid Telephone: (+34) (1) 5 63 69 24 Languages: Spanish, English | | Ward West 5 · Llandough Hospital Penarth Cardiff CF6 1FF · South Glamorgan Telephone: (+44) (0)12 22 70 99 01 Telefax: (+44) (0)12 22 70 43 57 Languages: English |

PROPER CONDUCT IN THE LABORATORY

Chemistry is a fascinating thing for many beginners in related jobs.

Handling chemicals is, however, not only fascinating, but also risky, especially if processes are not performed adequately due to insufficient knowledge of the properties of the substances used. It is, therefore, absolutely necessary to inform oneself prior to its first use about any possible hazards involved in the handling of a certain chemical.

Every manufacurer of laboratory reagents is obliged to label these with the respective safety data. Should there be no reliable data on the toxicity of a preparation, Merck reagents are labeled with a corresponding warning note.

Laboratory chemicals are neither intended nor approved for use in

humans or animals! Specifically tested pharmaceutical chemicals are available for such purpose.

The handling of hazardous chemicals demands special know-how, carefulness and adequate precautions. All tasks are to be performed such that hazardous gases, vapors or suspended particles are - as far as technically possible - not released.

If necessary, measures are to be taken to assure their uptake at the point of discharge/origin. When reaching the MAC specified in TRGS 900, special safety measures become necessary such as safe and practical personal protective gear. The following rules apply to provide safety during the working process:

- When working in the laboratory or warehouse wear safety glasses and, where necessary, gloves.
- 2. In any case, avoid contact with skin, eyes and mucous membranes.

- Rinse off any splashes on the skin with plenty of cold water; in the case of lipophilic substances rinse off with polyglycol. Due to the danger of absorption, never use other organic solvents.
- 4. Thoroughly rinse chemically burned eyes under a gentle stream of water or with a special eye shower. Rinse with your eyes wide open and roll your eyes in all directions. Subsequently, an ophthalmologic examination must be performed. It is absolutely necessary to inform the ophthalmologist of the chemical used. It is also recommended to state its hazardous properties in order to decide on adequate treatment.
- 5. Take off immediately any clothing contaminated with chemicals.
- In case of accidents or if you feel unwell, consult a doctor and state the cause of accident including the name of the chemical involved.
- 7. Do not smoke, eat and drink in laboratory rooms.

What is a hazardous material?

In accordance with the German Chemicals Act (ChemG) such substances and preparations are classified hazardous materials that have toxic (acute and/or chronic) and/or dangerous physical-chemical and/or environmentally harmful properties.



NOTES ON HANDLING ETHERS AND OTHER PEROXIDE-FORMING SOLVENTS

Ether peroxides are generally known to be hazardous.

Some people still often underestimate the risk involved thus occasionally causing explosions when distilling off ethers. Such highly powerful explosions may also occur when containers filled with peroxide-containing liquids are shaken or when upon opening of a bottle the peroxide contained in the bottle neck is ignited by the resultant friction. These hazards apply not only to ethers but also to other liquids tending to form peroxides such as unsaturated hydrocarbons, aldehydes, ketones and tetralin. In order to prevent accidents, such liquids should be tested regularly for peroxides.

Determination of peroxides 1.10011 Merckoquant® Peroxide Test Box with 100 strips

Ether peroxide can be easily and reliably detected and determined quantitatively with the test strips.

1.16206 Perex Test®

Information about the respective present peroxide content is the precondition for possibly quantitative and at the same time possibly economical elimination of peroxides. The methods described so far all have the same disadvantage of not being universally employable due to e.g. formation of 2 phases in aqueous reagents, no detection of polymeric peroxides, only qualitative results etc.

The Perex Test® provides a detection system for peroxides which fulfills the following demands:

- Detection of peroxides in aqueous and organic environments.
- Determination of peroxides in organic solvents in one phase.
- Detection of hydroperoxides and polymeric peroxides.
- Semiquantitative determination possible.

Not detectable: Dialkyl peroxides, dibenzoyl peroxides, peracid esters.



RISK (R) PHRASES:

| R1 | Explosive when dry. |
|------------|---|
| R2 | Risk of explosion by shock, friction, fire or other |
| | sources of ignition. |
| R3 | Extreme risk of explosion by shock, friction, fire or other |
| | sources of ignition. |
| R4 | Forms very sensitive explosive metallic compounds. |
| R5 | Heating may cause an explosion. |
| R6 | Explosive with or without air contact. |
| R7 | May cause fire |
| R8 | Contact with combustible material may cause fire. |
| R9 | Explosive when mixed with combustible material. |
| R10 | Flammable. |
| R11 | Highly flammable. |
| R12 | Extremely flammable. |
| R14 | Reacts violently with water. |
| R15 | Contact with water liberates extremely |
| | flammable gases. |
| R16 | Explosive when mixed with oxidizing substances. |
| R17 | Spontaneously flammable in air. |
| R18 | In use, may form flammable/explosive |
| | vapour-air mixture. |
| R19 | May form explosive peroxides. |
| R20 | Harmful by inhalation. |
| R21 | Harmful in contact with skin. |
| R22 | Harmful if swallowed. |
| R23 | Toxic by inhalation. |
| R24 | Toxic in contact with skin. |
| R25 | Toxic if swallowed. |
| R26 | Very toxic by inhalation. |
| R27 | Very toxic in contact with skin. |
| R28 | Very toxic if swallowed. |
| R29 | Contact with water liberates toxic gas. |
| R30 | Can become highly flammable in use. |
| R31 | Contact with acids liberates toxic gas. |
| R32 | Contact with acids liberates very toxic gas. |
| R33 | Danger of cumulative effects. |
| R34 | Causes burns. |
| R35 | Causes severe burns. |
| R36 R37 | Irritating to eyes. |
| R38 | Irritating to respiratory system. |
| R39 | Irritating to skin. Danger of very serious irreversible effects. |
| R40 | Possible risk of irreversible effects. |
| R41 | Risk of serious damage to eyes. |
| R42 | May cause sensitization by inhalation. |
| R42 | May cause sensitization by finialation. May cause sensitization by skin contact. |
| R44 | Risk of explosion if heated under confinement. |
| R45 | May cause cancer. |
| R46 | May cause heritable genetic damage. |
| R48 | Danger of serious damage to health by |
| | prolonged exposure. |
| R49 | May cause cancer by inhalation. |
| R50 | Very toxic to aquatic organisms. |
| | |

| B | l 🕳 a company of the |
|-----|---|
| R51 | Toxic to aquatic organisms. |
| R52 | Harmful to aquatic organisms. |
| R53 | May cause long-term adverse effects in the aquatic environment. |
| R54 | Toxic to flora. |
| R55 | Toxic to fauna. |
| R56 | Toxic to soil organisms. |
| R57 | Toxic to bees. |
| R58 | May cause long-term adverse effects in |
| | the environment. |
| R59 | Dangerous for the ozone layer. |
| R60 | May impair fertility. |
| R61 | May cause harm to the unborn child. |
| R62 | Possible risk of impaired fertility. |
| R63 | Possible risk of harm to the unborn child. |
| R64 | May cause harm to breastfed babies. |
| R65 | Harmful: May cause lung damage if swallowed. |
| R66 | Repeated exposure may cause skin dryness and dizziness. |
| R67 | Vapours may cause tiredness and lightheadedness. |
| R68 | Possible risks of irreversible effects. |
| | |

COMBINATION OF RISK PHRASES:

Harmful: possible risk of irreversible effects in contact with skin

Harmful: possible risk of irreversible effects if swallowed.

| | COMBINATION OF KISK FIRASES | | |
|------------------|--|--|--|
| R14/15 | Reacts violently with water, liberating extremely flammable gases. | | |
| R14/15 R15/29 | Contact with water liberates toxic, highly flammable gas. | | |
| R20/21 | Harmful by inhalation and in contact with skin. | | |
| | | | |
| R20/21/22 | Harmful by inhalation, in contact with skin and if swallowed. | | |
| R20/22 | Harmful by inhalation and if swallowed. | | |
| R21/22 | Harmful in contact with skin and if swallowed. | | |
| R23/24 | Toxic by inhalation and in contact with skin. | | |
| R23/24/25 | Toxic by inhalation, in contact with skin | | |
| | and if swallowed. | | |
| R23/25 | Toxic by inhalation and if swallowed. | | |
| R24/25 | Toxic in contact with skin and if swallowed. | | |
| R26/27 | Very toxic by inhalation and in contact with skin. | | |
| R26/27/28 | Very toxic by inhalation, in contact with skin and if swallowed. | | |
| R26/28 | Very toxic by inhalation and if swallowed. | | |
| R27/28 | Very toxic in contact with skin and if swallowed. | | |
| R36/37 | Irritating to eyes and respiratory system. | | |
| R36/37/38 | Irritating to eyes, respiratory system and skin. | | |
| R36/38 | Irritating to eyes and skin. | | |
| R68/20 | Harmful: possible risk of irreversible effects through inhalation. | | |
| R68/20/21 | Harmful: possible risk of irreversible effects through inhalation | | |
| | and in contact with skin. | | |
| R68/20/21/22 | Harmful: possible risk of irreversible effects through inhalation, | | |
| | in contact with skin and if swallowed. | | |
| R68/20/22 | Harmful: possible risk of irreversible effects through inhalation | | |
| | and if swallowed. | | |
| R68/21 | Harmful: possible risk of irreversible effects in contact with skin. | | |

and if swallowed.

R68/21/22

R68/22

RISK (R) PHRASES:

| R37/38 | Irritating to respiratory system and skin. |
|------------------|---|
| R39/23 | Toxic: danger of very serious irreversible effects |
| | through inhalation. |
| R39/23/24 | Toxic: danger of very serious irreversible effects through |
| | inhalation and in contact with skin. |
| R39/23/25 | Toxic: danger of very serious irreversible effects |
| Dan/aa/ | through inhalation and if swallowed. |
| R39/23/ 24/25 | Toxic: danger of very serious irreversible effects effects through inhalation, in contact with |
| 24/23 | skin and if swallowed. |
| R39/24 | Toxic: danger of very serious irreversible effects |
| , | in contact with skin. |
| R39/24/25 | Toxic: danger of very serious irreversible effects |
| | in contact with skin and if swallowed. |
| R39/25 | Toxic: danger of very serious irreversible effects |
| Baarlaa | if swallowed. |
| R39/26 | Very toxic: danger of very serious irreversible |
| R39/26/27 | effects through inhalation. Very toxic: danger of very serious irreversible effects |
| 1133/20/27 | through inhalation and in contact with skin. |
| R39/26/ | Very toxic: danger of very serious irreversible |
| 27/28 | effects through inhalation, in contact with skin |
| | and if swallowed. |
| R39/26/28 | Very toxic: danger of very serious irreversible effects |
| | through inhalation and if swallowed. |
| R39/27 | Very toxic: danger of very serious irreversible effects |
| R39/27/28 | in contact with skin. Very toxic: danger of very serious effects in contact with |
| N39/2//20 | skin and if swallowed. |
| R39/28 | Very toxic: danger of very serious irreversible effects |
| | if swallowed. |
| R40/20 | Harmful: possible risk of irreversible effects |
| | through inhalation. |
| R40/20/21 | Harmful: possible risk of irreversible effects through |
| | inhalation and in contact with skin. |
| R40/20/ | Harmful: possible risk of irreversible effects |
| 21/22 | through inhalation, in contact with skin and if swallowed. |
| R40/20/22 | Harmful: possible risk of irreversible effects through |
| ,20,22 | inhalation and if swallowed. |
| R40/21 | Harmful: possible risk of irreversible effects in contact |
| | with skin. |
| R40/21/22 | Harmful: possible risk of irreversible effects in |
| D / | contact with skin and if swallowed. |
| R40/22 | Harmful: possible risk of irreversible effects if swallowed. |
| R42/43 R48/20 | May cause sensitization by inhalation and skin contact. Harmful: danger of serious damage to health by |
| 11-13/20 | prolonged exposure through inhalation. |
| R48/20/21 | Harmful: danger of serious damage to health by prolonged |
| | exposure through inhalation and in contact with skin. |
| R48/20/ | Harmful: danger of serious damage to health by |
| 21/22 | prolonged exposure through inhalation, in contact |
| | with skin and if swallowed. |

| 48/20/22 | Harmful: danger of serious damage to health by |
|----------|---|
| 40/04 | prolonged exposure through inhalation and if swallowed. |
| 48/21 | Harmful: danger of serious damage to health by |
| | prolonged exposure in contact with skin. |
| 48/21/22 | Harmful: danger of serious damage to |
| | health by prolonged exposure in contact with |
| | skin and if swallowed. |
| 48/22 | Harmful: danger of serious damage to health by |
| | prolonged exposure if swallowed. |
| 48/23 | Toxic: danger of serious damage to health by |
| | prolonged exposure through inhalation. |
| 48/23/24 | Toxic: danger of serious damage to health by |
| | prolonged exposure through inhalation and |
| | in contact with skin. |
| 48/23/ | Toxic: danger of serious damage to health by prolonged |
| 4/25 | exposure through inhalation, in contact with skin and |
| | if swallowed. |
| 48/23/25 | Toxic: danger of serious damage to health by |
| | prolonged exposure through inhalation |
| | and if swallowed. |
| 48/24 | Toxic: danger of serious damage to health by prolonged |
| ,- | exposure in contact with skin. |
| 48/24/25 | Toxic: danger of serious damage to health by |
| .0,2.,20 | prolonged exposure in contact with skin and |
| | if swallowed. |
| 48/25 | Toxic: danger of serious damage to health by prolonged |
| 40/23 | exposure if swallowed. |
| 50/53 | Very toxic to aquatic organisms, may |
| 30/33 | cause long-term adverse effects in the |
| | aquatic environment. |
| 51/53 | Toxic to aquatic organisms, may cause long-term adverse |
| 31/33 | effects in the aquatic environment. |
| 52/53 | Harmful to aquatic organisms, may cause long-term |
| 52/53 | |
| | adverse effects in the aquatic environment. |

RISK PHRASES WITH SUPPLEMENTS

| R E20 | Also harmful by inhalation. |
|-------------|--|
| R E20/21 | Also harmful by inhalation and in contact with skin. |
| R E20/21/22 | Also harmful by inhalation, in contact with skin |
| | and if swallowed. |
| R E20/22 | Also harmful by inhalation and if swallowed. |
| R E21 | Also harmful in contact with skin. |
| R E21/22 | Also harmful in contact with skin and if swallowed. |
| R E22 | Also harmful if swallowed. |
| R E23 | Also toxic by inhalation. |
| R E23/24 | Also toxic by inhalation and in contact with skin. |
| R E23/24/25 | Also toxic by inhalation, in contact with skin and |
| | if swallowed. |

RISK (R) PHRASES:

| R E23/25 | Also toxic by inhalation and if swallowed. | R E40/20/ | Also harmful: possible risk of irreversible effects through |
|--------------|--|-------------|---|
| R E24 | Also toxic in contact with skin. | 21/22 | inhalation, in contact with skin and if swallowed. |
| R E24/25 | Also toxic in contact with skin and if swallowed. | R E40/20/22 | Also harmful: possible risk of irreversible effects |
| R E25 | Also toxic if swallowed. | | through inhalation and if swallowed. |
| R E26 | Also very toxic by inhalation. | R E40/21 | Also harmful: possible risk of very serious irreversible |
| R E26/27 | Also very toxic by inhalation and in contact with skin. | | effects in contact with skin. |
| R E26/27/28 | Also very toxic by inhalation, in contact with skin | R E40/21/22 | Also harmful: possible risk of irreversible effects in |
| | and if swallowed. | | contact with skin and if swallowed. |
| R E26/28 | Also very toxic by inhalation and if swallowed. | R E40/22 | Also harmful: possible risk of irreversible effects |
| R E27 | Also very toxic in contact with skin. | N L40/22 | if swallowed. |
| R E27/28 | Also very toxic in contact with skin and if swallowed. R E28 | R E42/43 | May cause sensitization by inhalation and |
| | Also very toxic if swallowed. | 11 142/43 | skin contact. |
| R E39/23 | Also toxic: danger of very serious irreversible | R E48/20 | Also harmful: danger of serious damage to health |
| | effects through inhalation. | N L40/20 | by prolonged exposure through inhalation. |
| R E39/23/24 | Also toxic: danger of very serious irreversible effects | D E40/20/21 | Also harmful: danger of serious damage to health |
| | through inhalation and in contact with skin. | R E48/20/21 | by prolonged exposure through inhalation and |
| R E39/23/ | Also toxic: danger of very serious irreversible | | in contact with skin. |
| 24/25 | effects through inhalation, in contact with skin | D. Facileal | |
| • | and if swallowed. | R E48/20/ | Also harmful: danger of serious damage to health |
| R E39/23/25 | Also toxic: danger of very serious irreversible | 21/22 | by prolonged exposure through inhalation, in contact |
| ,, | effects through inhalation and if swallowed. | D = / / | with skin and if swallowed. |
| R E39/24 | Also toxic: danger of very serious irreversible | R E48/20/22 | Also harmful: danger of serious damage to health |
| 200/2 . | effects in contact with skin. | | by prolonged exposure through inhalation and |
| R E39/24/25 | Also toxic: danger of very serious irreversible effects | D =/ | if swallowed. |
| 200/2 1/20 | in contact with skin and if swallowed. | R E48/21 | Also harmful: danger of serious damage to health |
| R E39/25 | Also toxic: danger of very serious irreversible | | by prolonged exposure in contact with skin. |
| N L33/23 | effects if swallowed. | R E48/21/22 | Also harmful: danger of serious damage to health |
| R E39/26 | Also very toxic: danger of very serious irreversible | | by prolonged exposure in contact with skin and |
| 11 133/20 | effects through inhalation. | D =/ | if swallowed. |
| R E39/26/27 | Also very toxic: danger of very serious irreversible | R E48/22 | Also harmful: danger of serious damage to health |
| 11 133/20/27 | effects through inhalation and in contact with skin. | D = 40/00 | by prolonged exposure if swallowed. |
| R E39/26/ | Also very toxic: danger of very serious irreversible | R E48/23 | Also toxic: danger of serious damage to health |
| 27/28 | effects through inhalation, in contact with skin | D F40/22/24 | by prolonged exposure through inhalation. |
| 21/20 | and if swallowed. | R E48/23/24 | Also toxic: danger of serious damage to health |
| R E39/26/28 | Also very toxic: danger of very serious irreversible | | by prolonged exposure through inhalation and in contact with skin. |
| N E39/20/20 | effects through inhalation and if swallowed. | R E48/23/ | Also toxic: danger of serious damage to |
| R E39/27 | Also very toxic: danger of very serious irreversible | 24/25 | health by prolonged exposure through inhalation, |
| N E39/27 | effects in contact with skin. | 24/25 | in contact with skin and if swallowed. |
| D F20/27/20 | | R E48/23/25 | Also toxic: danger of serious damage to health |
| R E39/27/28 | Also very toxic: danger of very serious irreversible effects | N L40/23/23 | 3 |
| D 500/00 | in contact with skin and if swallowed. | | by prolonged exposure through inhalation and if swallowed. |
| R E39/28 | Also very toxic: danger of very serious irreversible | R E48/24 | Also toxic: danger of serious damage to health |
| D 540/00/04 | effects if swallowed. | n £40/24 | by prolonged exposure in contact with skin. |
| R E40/20/21 | Also harmful: possible risk of irreversible effects through | R E48/24/25 | Also harmful: danger of serious damage to health |
| | inhalation and in contact with skin. | n £40/24/25 | by prolonged exposure in contact with skin and |
| | | | if swallowed. |
| | | R E48/25 | Also toxic: danger of serious damage to health |
| | | 11 240/23 | by prolonged exposure if swallowed. |
| | | | by protonged exposure it swallowed. |
| | | | |

SAFETY (S) PHRASES:

| S 1 | Keep locked up. | S 28.2 | After contact with skin, wash immediately with |
|---------|--|--------|---|
| S 2 | Keep out of reach of children. | 3 20.2 | plenty of soap and water. |
| S 3 | Keep in a cool place. | S 28.3 | After contact with skin, wash immediately with |
| S 4 | Keep away from living quarters. | 3 20.3 | plenty of soap and water, if possible also with |
| S 5 | Keep contents under | | polyethylene glycol 400. |
| S 6 | Keep contents under (inert gas to be specified | S 28.4 | After contact with skin, wash immediately with plenty of |
| | by the manufacturer). | 5 2011 | polyethylene glycol 300 and ethanol (2:1) followed by |
| S 6.1 | Keep under nitrogen. | | plenty of soap and water. |
| S 7 | Keep container tightly closed. | S 28.5 | After contact with skin, wash immediately with plenty of |
| S 8 | Keep container dry. | 5 20.0 | polyethylene glycol 400. |
| S 9 | Keep container in a well-ventilated place. | S 28.6 | After contact with skin, wash immediately with plenty of |
| S 12 | Do not keep the container sealed. | 5 20.0 | polyethylene glycol 400, then rinse with plenty of water. |
| S 13 | Keep away from food, drink and animal feeding | S 28.7 | After contact with skin, wash immediately with plenty |
| | stuffs. | | of water and acidic soap. |
| S 14 | Keep away from | S 29 | Do not empty into drains. |
| S 14.1 | Keep away from reducing agents, heavy metal | S 30 | Never add water to this product. |
| | compounds, acids and alkalis. | S 33 | Take precautionary measures against static discharges. |
| S 14.10 | Keep away from acids, reducing agents and | S 35 | This material and its container must be disposed of |
| | flammable material. | | in a safe way. |
| S 14.11 | Keep away from flammable material. | S 36 | Wear suitable protective clothing. |
| S 14.2 | Keep away from oxidizing and acidic substances as | S 37 | Wear suitable gloves. |
| | well as heavy metal compounds. | S 38 | In case of insufficient ventilation, wear suitable |
| S 14.3 | Keep away from iron. | | respiratory equipment. |
| S 14.4 | Keep away from water and alkalis. | S 39 | Wear eye/face protection. |
| S 14.5 | Keep away from acids. | S 40 | To clean the floor and all objects contaminated by this |
| S 14.6 | Keep away from alkalis. | | material use (to be specified by the manufacturer). |
| S 14.7 | Keep away from metals. | S 41 | In case of fire and/or explosion do not breathe fumes. |
| S 14.8 | Keep away from oxidizing and acidic substances. | S 42 | During fumigation/spraying with wear suitable |
| S 14.9 | Keep away from flammable organic substances. | | respiratory equipment. |
| S 15 | Keep away from heat. | S 43 | In case of fire, use (indicate the precise type of |
| S 16 | Keep away from sources of ignition - No smoking. | | fire-fighting equipment. If water increases risk, |
| S 17 | Keep away from combustible material. | | add -'Never use water'. |
| S 18 | Handle and open container with care. | S 43.1 | In case of fire, use water. |
| S 20 | When using do not eat or drink. | S 43.2 | In case of fire, use water or powder extinguisher. |
| S 21 | When using do not smoke. | S 43.3 | In case of fire, use powder extinguisher. |
| S 22 | Do not breathe dust. | | Never use water. |
| S 23 | Do not breathe gas/fumes/vapour/spray | S 43.4 | In case of fire, use CO2 – never use water. |
| S 23.1 | Do not breathe gas. | S 43.6 | In case of fire, use sand - never use water. |
| S 23.2 | Do not breathe vapour. | S 43.7 | In case of fire, use metal fire powder - never use water. |
| S 23.3 | Do not breathe spray | S 43.8 | In case of fire, use sand, CO2 or powder extinguisher, |
| S 23.4 | Do not breathe fumes. | | never use water. |
| S 23.5 | Do not breathe fumes/spray. | S 45 | In case of accident or if you feel unwell, seek |
| S 24 | Avoid contact with skin. | | medical advice immediately (show the label where |
| S 25 | Avoid contact with eyes. | | possible). |
| S 26 | In case of contact with eyes, rinse immediately with | S 46 | If swallowed, seek medical advice immediately and |
| C 07 | plenty of water and seek medical advice. | 6.47 | show this container or label. |
| S 27 | Take off immediately all contaminated clothing. | S 47 | Keep at temperature no exceeding°C (to be |
| S 28 | After contact with skin, wash immediately with | S 47.1 | specified by the manufacturer). |
| C 20 1 | plenty of | 5 47.1 | Keep at temperature no exceeding 20°C. |
| S 28.1 | After contact with skin, wash immediately with | | |
| | plenty of water. | | |

SAFETY (S) PHRASES:

| S 48 | Keep wet with (appropriate material to be specified by the manufacturer). |
|--------|---|
| S 49 | Keep only in the original container. |
| S 50 | Do not mix with (to be specified by the manufacturer). |
| S 50.1 | Do not mix with acids. |
| S 50.2 | Do not mix with alkalis. |
| S 50.3 | Do not mix with strong acids, strong bases, non-ferrous |
| | metals or their salts. |
| S 51 | Use only in well-ventilated areas. |
| S 52 | Not recommended for interior use on large surface areas. |
| S 53 | Avoid exposure - obtain special instructions before use. |
| | Restricted to professional users. |
| S 56 | Dispose of this material and its container at hazardous |
| | or special waste collection point. |
| S 57 | Use appropriate container to avoid environmental |
| | contamination! |
| S 59 | Refer to manufacturer/supplier for information on |
| | recovery/recycling. |
| S 60 | This material and its container must be disposed of |
| | as hazardous waste. |
| S 61 | Avoid release to the environment. |
| | Refer to special instructions / Safety data sheet. |
| S 62 | If swallowed, do not induce vomiting. Seek medical |
| | advice immediately and show this container or label. |
| S 63 | In case of accident through inhalation: remove casually |
| | to fresh air and kept at rest. |
| S 64 | In case of swallowed, rinse mouth with water (only if |
| | the person is conscious). |
| | |
| | |

COMBINATION OF SAFETY PHRASES

| S 1/2 S 3/7 | Keep locked up and out of the reach of children. Keep container tightly closed in a cool place. |
|----------------|---|
| | |
| S 3/9 | Keep in a cool, well-ventilated place. |
| S 3/9/14 | Keep in a cool, well-ventilated place away from |
| | (incompatible substances are to be specified |
| | by the manufacturer). |
| S 3/9/14.1 | Keep in a cool, well-ventilated place away from |
| | reducing agents, heavy metal compounds, |
| | acids and alkalis. |
| S 3/9/14.2 | Keep in a cool, well-ventilated place away |
| | from oxidizing agents and acidic substances as |
| | well as heavy metal compounds. |
| S 3/9/14.3 | Keep in a cool, well-ventilated place away from iron. |
| S 3/9/14.4 | Keep in a cool, well-ventilated place away from |
| , ., | water and alkalis. |
| S 3/9/14.5 | Keep in a cool, well-ventilated place away from acids. |
| S 3/9/14.6 | Keep in a cool, well-ventilated place away from alkalis. |
| S 3/9/14.7 | Keep in a cool, well-ventilated place away from metals. |
| 3 3/3/14.7 | recep in a coof, wen ventuated place away from metals. |
| | |

| S 3/9/14.8 | Keep in a cool, well-ventilated place away from oxidizing and acidic substances. |
|---------------|--|
| S 3/9/14/49 | Keep only in the original container in a cool, well-ventilated place away from (incompatible |
| S 3/9/14.1/49 | materials to be indicated by the manufacturer). Keep in the original container in a cool, well-ventilate place away from reducing agents, heavy metal compounds, acids and alkalis. |
| S 3/9/14.2/49 | Keep only in the original container in a cool, well-ventilated place away from oxidizing and acidic substances as well as heavy metal compounds. |
| S 3/9/14.3/49 | Keep only in the original container in a cool, well-ventilated place away from iron. |
| S 3/9/14.4/49 | Keep only in the original container in a cool, well-ventilated place away from water and alkalis. |
| S 3/9/14.5/49 | Keep only in the original container in a cool, well-ventilated place away from acids. |
| S 3/9/14.6/49 | Keep only in the original container in a cool, well-ventilated place away from alkalis. |
| S 3/9/14.7/49 | Keep only in the original container in a cool, well-ventilated place away from metals. |
| S 3/9/14.8/49 | Keep only in the original container in a cool, well-ventilated place away from oxidizing and acidic substances. |
| S 3/9/49 | Keep only in the original container in a cool, well-ventilated place. |
| S 3/14 | Keep in a cool place away from (incompatible materials to be indicated by the manufacturer). |
| S 3/14.1 | Keep in a cool place away from reducing agents, heavy metal compounds, acids and alkalis. |
| S 3/14.2 | Keep in a cool place away from oxidizing and acidic substances as well as heavy metal compounds. |
| S 3/14.3 | Keep in a cool place away from iron. |
| S 3/14.4 | Keep in a cool place away from water and alkalis. |
| S 3/14.5 | Keep in a cool place away from acids. |
| S 3/14.6 | Keep in a cool place away from alkalis. |
| S 3/14.7 | Keep in a cool place away from metals. |
| S 3/14.8 | Keep in a cool place away from oxidizing and |
| | acidic substances. |

SAFETY (S) PHRASES:

S 7/8 S 7/9 S 7/47 S 20/21 S 24/25

S 27/28

Keep container tightly closed and dry.
Keep container tightly closed and in a well-ventilated place.
Keep container tightly closed and at temperature no exceeding ... "C (to be specified by the manufacturer).
When using do not eat, drink or smoke.
Avoid contact with skin and eyes.
After contact with skin, take off immediately all contaminated clothing and wash skin with plenty of ... (to be specified by the manufacturer).

| S 29/56 | Do not empty into drains; dispose of this material and its container at hazardous or special waste collection point. |
|------------|--|
| S 36/37 | Wear suitable protective clothing and gloves. |
| S 36/37/39 | Wear suitable protective clothing, gloves and eye/face protection. |
| S 36/39 | Wear suitable protective clothing and eye/face protection. |
| S 37/39 | Wear suitable gloves and eye/face protection. |
| S 47/49 | Keep only in the original container at temperature ne exceeding *C (to be specified by the manufacturer). |
| | |

Copyright © 1999 [Merck KGaA]. All rights reserved. Last update: 18.05.01

















E: Explosive

Criteria: Chemicals and preparations which may react exothermically without atmos-

pheric oxygen and which under defined test conditions detonate, quickly deflagrate or upon heating explode when partially confined.

Precaution: Avoid impact, knocks, friction, sparks, fire, and heat.



F+: Extremely flammable.

Criteria: Liquids with a flash point below 0 °C and a boiling point of max. 35 °C.

Gases and gas mixtures which are flammable in air at normal pressure and average temperatures.

Precaution: Keep away from naked flames, sparks, and sources of heat.



C: Corrosive.

Criteria: Total damage to living tissues or when this result can be predicted.

Precaution: Take special measures to protect eyes, skin, and clothes. Do not inhale vapors! In case of accident or if you feel unwell, seek medical advice immediately.



Xi: Irritating.

Criteria: Without being corrosive, immediate, prolonged, or repeated contact with skin

or mucous membranes may cause inflammations. Risk of sensitization by skin contact (classification with R43).

Precaution: Avoid contact with eyes and skin, do not inhale vapors.



and smoulder.

sparks, and sources of heat.

O: Oxidizing.

Criteria: Organic peroxides which are combustible even if not in contact with com-

bustible materials. Other chemicals and preparations which as a rule are not combustible themselves, but which in contact with combustible materials, mainly through oxygen evolution, considerably increase the fire hazard and the intensity of a fire.

Precaution: Avoid all contact with combustible substances.

Risk of ignition: The substance promotes fires once started and impedes fire fighting.

F: Highly flammable.

exposure to a source of ignition may be

easily inflamed and then continue to burn

Precaution: Keep away from naked flames,

Criteria: Liquids with a flash point below 21 °C that are not extremely flammable. Solid substances and preparations which on brief



T: Toxic.

Criteria: Inhalation, swallowing, or absorption through the skin in small amounts

can cause considerable damage to health, and may sometimes be lethal. In the event of serious evidence of severe, possibly irreversible damage to health by single, repeated, or prolonged absorption, especially carcinogenic, mutagenic, and reproduction-toxic effects.

Precaution: All contact with the human body must be avoided. If you feel unwell, seek medical advice immediately. Particular attention is drawn to the carcinogenic, teratogenic, or mutagenic risks associated with certain substances.

Observe special regulations when handling these substances!



T+: Verv toxic.

Criteria: Inhalation, swallowing, or absorption through the skin in very small amounts

can cause considerable damage to health, and may sometimes be lethal. In the event of serious evidence of severe, possibly irreversible damage to health by single, repeated, or prolonged absorption.

Criteria: All contact with the human body must be avoided. If you feel unwell, seek medical advice immediately.



Xn: Harmful.

Criteria: Inhalation, swallowing, or absorption through the skin can cause acute or

chronic damage to health. In the event of evidence of severe, possibly irreversible damage to health by single, repeated, or prolonged absorption, especially in suspected carcinogenic, mutagenic, and reproduction-toxic effects.

Risk of sensitization by inhalation (classification with R42).

Precaution: All contact with the human body must be avoided. Particular attention is drawn to substances which are suspected to have a carcinogenic, mutagenic, or reproduction-toxic effect.



N: Dangerous for the environment.

Criteria: Liberation into the aquatic and non-aquatic envi-

ronments can have an immediate or delayed detrimental effect upon the ecosystem through alteration of the natural balance. Some substances or their conversion products may simultaneously affect various constituents of the ecosystem.

Precaution: Depending on the risk potential do not allow to enter sewerage systems. Observe special disposal regulations!

INCOMPATIBLE CHEMICALS


The chemicals listed below may react violently with one another. They must be kept apart and must never come into contact with one another. The objective of this list is to give information on how to avoid accidents in the laboratory.

Due to the great number of hazardous materials, this list includes only the most important examples.

| Substance | Incompatible with |
|--|--|
| Acetylene Acetic acid | Chlorine, bromine, fluorine, copper, silver, mercury Chromium (VI) oxide, nitric acid, alcohols, ethylene glycol, perchloric acid, peroxides, permanganates |
| Activated carbon | Calcium hypochlorite, oxidizing agents |
| Alkali metals | Water, carbon tetrachloride and other halogenated alkanes, carbon dioxide, halogens |
| Aluminum alkyls | Water |
| Ammonia (laborato- ry gas or solutions) | Mercury (e.g. in pressure gauges), chlorine, calcium hypochlorite, iodine, bromine, hydrogen fluoride |
| Ammonium nitrate | Acids, powdered metals, flammable liquids, chlorates, nitrates, sulfur, fine-particulate organic or combustible materials |
| Aniline | Nitric acid, hydrogen peroxide |
| Bromine | See chlorine |
| Chlorine | Ammonia, acetylene, butadiene, butane, methane, propane, hydrogen, petroleum benzine, benzene, powdered metals |
| Chlorates | Ammonium salts, acids, powdered metals, sulfur, fine-particulate organic or combustible substances |
| Chromium (VI) oxide | Acetic acid, naphthalene, camphor, glycerol, petroleum benzine, alcohols, flammable liquids |
| Copper | Acetylene, hydrogen peroxide |
| Cumene hydroperoxide | Acids, both organic and inorganic |
| Cyanides | Acids |
| Flammable liquids | Ammonium nitrate, chromium (VI) oxide, hydrogen peroxide, nitric acid, sodium peroxide, halogens |
| Fluorine | Extremely aggressive; store separately! |



| Hydrocarbon | Fluorine, chlorine, bromine, |
|-------------------------------|---|
| (butane, propane, | chromium (VI) oxide, sodium peroxide |
| benzene etc.) | |
| Hydrogen fluoride | Ammonia (laboratory gas or solutions) |
| Hydrogen peroxide | Copper, chromium, iron, metals and metal salts, alcohols, acetone, organic substances, aniline, nitromethane, combustible substances (solid or liquid) |
| Hydrogen sulfide | Fuming nitric acid, oxidizing gases |
| lodine | Acetylene, ammonia (laboratory gas or solutions) |
| Mercury | Acetylene, ammonia |
| Nitric acid (concentrated) | Acetic acid, aniline, chromium (VI) oxide, prussic acid, hydrogen sulfide, flammable liquids and gases |
| Oxalic acid | Silver, mercury |
| Perchloric acid | Acetic anhydride, bismuth and its alloys, alcohols, paper, wood |
| Phosphorus | Sulfur, compounds containing oxygen, e.g. chlorates |
| Potassium | See alkali metals |
| Potassium chlorate | See chlorate |
| Potassium perchlorate | See chlorate |
| Potassium | Glycerol, ethylene glycol, benzaldehyde, sulfuric acid |
| permanganate | |
| Silver | Acetylene, oxalic acid, tartaric acid, ammonium compounds |
| Sodium | See alkali metals |
| Sodium peroxide | Methanol, ethanol, glacial acetic acid, acetic anhydride, benzaldehyde, carbon disulfide, glycerol, ethylene glycol, ethyl acetate, methyl acetate, furfural |
| Sulfuric acid | Potassium chorate, potassium perchlorate, potassium permangangate |



HARDNESS SCALE ACC. TO MOHS

| Hardness | Mineral | Formula |
|----------|-----------|--|
| 1 | Talcum | Mg ₃ [(OH) ₂ / Si ₄ O ₁₀] |
| 2 | Gypsum | CaSO ₄ · 2H ₂ O |
| 3 | Calcite | CaCO ₃ |
| 4 | Fluorspar | CaF ₂ |
| 5 | Apatite | Ca_5 [(F, CI, OH) / $(PO_4)_3$] |
| 6 | Feldspar | KAISi ₃ O ₈ |
| 7 | Quartz | SiO ₂ |
| 8 | Topaz | $AI_2[F_2/SiO_4]$ |
| 9 | Corundum | Al_2O_3 |
| 10 | Diamond | С |

ELECTROCHEMICAL SERIES OF THE ELEMENTS (ACIDIC SOLUTION)

 ϵ^o under standard conditions [25 °C, 10 5 Pa (1 bar)] measured against the hydrogen electrode as reference electrode

| $Red \rightleftarrows Ox + e$ | e° (Volt) | Red ⇄ Ox +e | e° (Volt) |
|--|-----------|---|-----------|
| Li ⇄ Li+ 1e | - 3.045 | $Fe \rightleftharpoons Fe^2 + 2e$ | - 0.41 |
| Rb ← Rb+ + 1e | - 2.98 | $Cd \stackrel{\longrightarrow}{\leftarrow} Cd^{2+} + 2e$ | - 0.403 |
| $Cs \rightleftharpoons Cs^+ + 1e$ | - 2.92 | TI ⇄ TI+ +1e | - 0.335 |
| $K \rightleftharpoons K^+ + 1e$ | - 2.92 | $Co \stackrel{\textstyle \longrightarrow}{\longleftarrow} Co^{2+} + 2e$ | - 0.277 |
| $Ba \rightleftharpoons Ba^{2+} + 2e$ | - 2.90 | $Ni \rightleftharpoons Ni^{2+} + 2e$ | - 0.25 |
| $Sr \rightleftharpoons Sr^{2+} + 2e$ | - 2.89 | $Sn ightleftharpoons Sn^{2+} 2e$ | - 0.136 |
| Ca ⇌ Ca ²⁺ + 2e | - 2.866 | $Pb \rightleftharpoons Pb^{2+} + 2e$ | - 0.126 |
| Na → Na ⁺ + 1e | - 2.713 | H ₂ 2 2H+ + 2e | ± 0.000 |
| $Mg ightleftharpoons Mg^{2+} + 2e$ | - 2.36 | $Cu^{2} \rightleftharpoons Cu^{2+} + 2e$ | + 0.337 |
| $Be \stackrel{\longrightarrow}{\leftarrow} Be^{2+} + 2e$ | - 1.85 | Cu ⇄ Cu+ + 1e | + 0.52 |
| $AI \rightleftharpoons AI^{3+} + 3e$ | - 1.66 | Te $ ightleftharpoons$ Te ⁴⁺ + 4e | + 0.56 |
| $Mn \rightleftharpoons Mn^{2+} + 2e$ | - 1.18 | $Ag \stackrel{\longrightarrow}{\leftarrow} Ag^+ + 1e$ | + 0.799 |
| $Zn \rightleftharpoons Zn^{2+} + 2e$ | - 0.763 | $Hg ightharpoonup Hg^{2+} + 2e$ | + 0.854 |
| $Cr \rightleftharpoons Cr^{3+} + 3e$ | - 0.744 | $Pd \stackrel{\longrightarrow}{\rightleftharpoons} Pd^{2+} + 2e$ | + 0.987 |
| $Co \rightleftharpoons Co^{3+} + 3e$ | - 0.74 | $Pt \stackrel{\textstyle \longrightarrow}{\leftarrow} Pt^{2+} + 2e$ | ~ + 1.2 |
| $Ga \rightleftharpoons Ga^{3+} + 3e$ | - 0.53 | $Au \rightleftharpoons Au^{3+} + 3e$ | + 1.498 |

*** Earth's crust

| Element name | Sym- bol | Ordinal- number | Atomic mass** | Density D ^{20°} 4° | Melting- point [°C] | Boiling- point [°C] | Occurrence of the elements*** [%] | Atomic radiuses [pm] | lonic radiuses [pm] | Electro- negativity |
|-------------------|-------------|--------------------|---------------|-----------------------------------|------------------------|------------------------|-----------------------------------|----------------------|--------------------------------------|------------------------|
| Actinium | Ac | 89 | 227.028 | 10.1 | 1050 | 3200 ± 300 | _ | - | 118 (III) | 1.00 |
| Aluminium | Al | 13 | 26.98154 | 2.70 | 660.37 | 2467 | 8.1 | 143 | 51 (III) | 1.47 |
| Americium | Am | 95 | (243) | 11.7 | 994 ± 4 | 2607 | _ | - | 107 (III), 92 (IV) | ~ 1.2 |
| Antimony | Sb | 51 | 121.76 | 6.68 | 630.74 | 1750 | 0.0001 | 145 | 76 (III), 62 (V) | 1.82 |
| Argon | Ar | 18 | 39.948 | *1.784 | -189.2 | -185.7 | - | 191 | - | _ |
| Arsenic | As | 33 | 74.9216 | 5.73 | 817 (28 bar) | | 0.0005 | 125 | 58 (III), 46 (V) | 2.20 |
| Astatine | At | 85 | (210) | - | 302 | 337 | - | - | 62 (VII) | 1.96 |
| Barium | Ba | 56 | 137.33 | 3.7 | 725 | 1640 | 0.025 | 217 | 134 (II) | 0.97 |
| Berkelium | Bk | 97 | (247) | - | - | - | - | - | - | ~ 1.2 |
| Beryllium | Be | 4 | 9.01218 | 1.86 | 1278 ± 5 | 2970 (5 mm) | 0.0006 | 112 | 35 (II) | 1.47 |
| Bismuth | Bi | 83 | 208.9804 | 9.80 | 271.3 | 1560 ± 5 | 0.00002 | 155 | 96 (III), 74 (V) | 1.67 |
| Boron | В | 5 | 10.81 | 2.34 | 2300 | 2550 (sub.) | 0.0003 | 97 | 23 (III) | 2.01 |
| Bromine | Br | 35 | 79.904 | 3.14 | -7.2 | 58.78 | 0.00016 | 119 | 196 (-I), 47 (V), 39 (VII) | 2.74 |
| Cadmium | Cd | 48 | 112.41 | 8.64 | 320.9 | 765 | 0.000015 | 149 | 97 (II) | 1.46 |
| Cesium | Cs | 55 | 132.9054 | 1.90 | 28.40 ± 0.01 | 678.4 | 0.0007 | 262 | 167 (I) | 0.86 |
| Calcium | Ca | 20 | 40.078 | 1.55 | 839 ± 2 | 1484 | 3.6 | 196 | 99 (II) | 1.04 |
| Californium | Cf | 98 | (251) | - | - | - | - | - | _ | ~ 1.2 |
| Carbon | С | 6 | 12.011 | 2.25 | ~ 3550 | 4827 | 0.03 | 77 | 16 (IV) | 2.50 |
| Cerium | Ce | 58 | 140.115 | 6.8 | 798 ± 3 | 3257 | 0.0046 | 182 | 107 (III), 94 (IV) | 1.06 |
| hlorine | CI | 17 | 35.4527 | *3.214 | -100.98 | -34.6 | 0.031 | 107 | 181 (-I), 34 (V), 27 (VII) | 2.83 |
| hromium | Cr | 24 | 51.996 | 7.19 | 1857 ± 20 | 2672 | 0.02 | 125 | 63 (III), 52 (VI) | 1.56 |
| Cobalt | Co | 27 | 58. 9332 | 8.83 | 1495 | 2870 | 0.0023 | 125 | 72 (II), 63 (III) | 1.70 |
| Copper | Cu | 29 | 63.546 | 8.93 | 1083.4 ± 0.2 | 2567 | 0.007 | 128 | 96 (I), 72 (II) | 1.75 |
| Curium | Cm | 96 | (247) | _ | 1340 ± 40 | - | - | - | - | ~ 1.2 |
| ysprosium | Dy | 66 | 162.50 | 8.54 | 1409 | 2335 | 0.00045 | - | 92 (III) | 1.10 |
| insteinium | Es | 99 | (254) | _ | - | - | - | - | - | ~ 1.2 |
| rbium | Er | 68 | 167.26 | 9.05 | 1522 | 2510 | 0.00025 | - | 89 (III) | 1.11 |
| uropium | Eu | 63 | 151.96 | 5.26 | 822 ± 5 | 1597 | 0.00011 | - | 124 (II), 98 (III) | 1.01 |
| ermium | Fm | 100 | (257) | - | - | - | - | - | - | ~ 1.2 |
| luorine | F | 9 | 18.9984 | *1.70 | -219.62 | -188.14 | 0.03 | 71 | 133 (-I), 8 (VII) | 4.10 |
| rancium | Fr | 87 | (223) | - | (27) | (677) | _ | - | 180 (I) | 0.86 |
| iadolinium | Gd | 64 | 157.25 | 7.90 | 1311 ± 1 | 3233 | 0.00064 | - | 97 (III) | 1.11 |
| iallium | Ga | 31 | 69.723 | 6.0 | 29.78 | 2403 | 0.0015 | - | 62 (III) | 1.82 |
| Germanium | Ge | 32 | 72.61 | 5.36 | 937.4 | 2830 | 0.0007 | - | 73 (II), 53 (IV) | 2.02 |
| iold | Au | 79 | 196.966 | 19.3 | 1064.4 | 2807 | 0.00000005 | 144 | 137 (I), 85 (III) | 1.42 |
| lafnium | Hf | 72 | 178.49 | 13.3 | 2227 ± 20 | 4602 | 0.00045 | - | 78 (IV) | 1.23 |
| lelium | He | 2 | 4.00260 | *0.178 | -272.2 | -268.934 | 0.00000003 | 145 | - | - |
| lolmium | Ho | 67 | 164.93 | 8.80 | 1470 | 2720 | 0.00012 | - | 91 (III) | 1.10 |
| lydrogen | Н | 1 | 1.00794 | *0.0899 | - 259.14 | - 252.87 | 0.14 | 46 | 154 (-I) | 2.20 |
| ndium | In | 49 | 114.82 | 7.31 | 156.61 | 2080 | 0.00001 | - | 81 (III) | 1.49 |
| od | 1 | 53 | 126.9045 | 4.94 | 113.5 | 184.35 | 0.00003 | 136 | 220 (-), 62 (V), 50 (VII) | 2.21 |
| idium | lr | 77 | 192.22 | 22.6 | 2410 | 4130 | 0.0000001 | - | 68 (IV) | 1.55 |
| on | Fe | 26 | 55.847 | 7.86 | 1535 | 2750 | 5.0 | 124 | 74 (II), 64 (III) | 1.64 |
| Trypton | Kr | 36 | 83.80 | *3.708 | -156.6 | -152(3) | _ | - | - | - |
| anthanum | La | 57 | 138.9055 | 6.1 | 920 ± 5 | 3454 | 0.0018 | - | 114 (III) | 1.08 |
| awrencium | Lr | 103 | (260) | - | - | - | - | - | - | - |
| ead | Pb | 82 | 207.2 | 11.4 | 327.5 | 1740 | 0.0016 | 175 | 215 (-II), 120 (II), 84 (IV) | 1.55 |
| ithium | Li | 3 | 6.941 | 0.53 | 180.54 | 1347 | 0.0065 | 152 | 68 (I) | 0.97 |
| utetium | Lu | 71 | 174.967 | 9.84 | 1656 ± 5 | 3315 | 0.00008 | _ | 85 (III) | 1.14 |
| N agnesium | Mg | 12 | 24.305 | 1.74 | 648.8 ± 0,5 | 1090 | 0.21 | 160 | 66 (II) | 1.23 |
| Manganese | Mn | 25 | 54.93805 | 7.3 | 1244 ± 3 | 1962 | 0.1 | 118 | 80 (II), 66 (III), 60 (IV), 46 (VII) | 1.60 |

normal

°C and

at

[g/ I]

Gases in

CAL

TABLE OF FLEMENTS Density Occurrence of Atomic Melting-Boiling-**Flement** Ordinal-Atomic Flectro-Svm D 20 Ionic radiuses the elements* radiuse point [°C] name bol number mass** point [°C] [pm] negativity [%] [ma] Md (258)Mendelevium 101 ~ 1.2 Ha 80 200 59 13.55 - 38.87 356 58 0.00005 150 110 (II) 1.44 Mercury Mo 70 (IV), 62 (VI) Molybdenum 42 95.94 10.2 2617 4612 0.0015 1.30 _ 104 (III) Neodymium Nd 60 144.24 7.0 1010 3127 0.0024 1.07 Ne 10 20 1797 *0.90 - 248 7 - 246.05 Neon _ 93 640+1 _ 1.22 Neptunium Nn 237.0482 19.5 3902 110 (III), 95 (IV), 71 (VII) Nickel Ni 28 58.69 8.90 1453 2732 0.008 124 69 (II) 1.75 Niobium Nb 41 92.9064 8.5 2468 ± 10 4742 0.0024 74 (IV), 69 (VI) 1.23 N *1.251 3.07 Nitrogen 7 14.0067 - 209.86 - 195.8 0.0046 71 16 (III), 13 (V) No 102 Nobelium (259)**O**smium 0s 76 190.23 22.5 3045 ± 30 0.00000001 67 (IV), 69 (VI) 1.52 5027+100 0 8 Oxygen 15.9994 *1.429 -218.4- 182.962 46.6 65 132 (-II), 10 (VI) 3.50 Pd Palladium 46 106.42 12.0 1552 3140 0.0000001 80 (II), 65 (IV) 1.35 Р 15 30.97376 1.83 44.1 280 12 44 (III), 35 (V) 2.06 Phosphorous, white Platinum Pt 78 195.08 21.45 1.772 3827 0.00000005 138 80 (II), 65 (IV) 1.44 Pu 94 Plutonium (244)19.7 641 3232 108 (III), 93 (IV) 1.22 Polonium Po 84 (209)9 32 254 962 1.76 K 19 Potassium 39.0983 0.86 63.65 774 2.6 231 133 (I) 0.91 59 931 ± 4 Praseodymium Pr 140.908 6.7 3212 0.00055 106 (III), 92 (IV) 1.07 Pm 61 (145)~ 1080 106 (III) 1.07 Promethium Pa Protactinium 91 231.036 < 1600 113 (III), 98 (IV), 89 (V) 1.14 Radium Ra 88 226.0254 ~6 700 1140 143 (II) 0.97 Radon Rn 86 *9 96 - 71 - 61.8 Re 1.46 Rhenium 75 186,207 20.9 3180 0.00000001 72 (IV), 56 (VII) Rhodium Rh 45 102 905 12.4 1966 ± 3 3727±100 0.00000001 68 (III) 1.45 Rb 37 85,4678 1.53 38.89 688 0.03 243 147 (I) 0.89 Rubidium Ru 44 67 (IV) Ruthenium 101.07 12.4 2310 3000 0.00000001 1.42 Samarium Sm 62 150.36 7.5 1072 + 51778 0.00065 100 (III) 1.07 Scandium Sc 21 44.9559 3.0 1539 2832 0.0005 81 (III) 1.20 Selenium Se 34 78.96 4.8 217 684.9+1.0 0.000009 _ 191 (-II), 83 (III), 50 (IV), 42 (VI) 2.48 Silver Ag 47 107.8682 10.5 961 93 2212 0.00001 144 126 (I), 89 (II) 1.42 Si Silicium 14 28.0855 2.4 1410 2355 27.7 117 221 (-IV), 42 (IV) 1.74 Sodium Na 11 22.98977 0.97 97.81 + 0.03882.9 2.8 186 97 (I) 1.01 38 Strontium Sr 87.62 2.6 769 1384 0.03 112 (II) 0.99 Sulphur S 16 32.066 2.0 112.8 444.674 0.05 104 174 (-II), 37 (IV), 30 (VI) 2.44 Ta Tantalum 73 180.9479 16.7 2996 5425±100 0.00021 68 (V) 1.33 Technetium Tc 43 (97)11.5 2172 4877 56 (VII) 1.36 52 6.2 2.01 Tellurium Te 127.60 449.5 + 0.3989.8+3.8 0.00000002 211 (-II), 70 (IV), 56 (VI) Terbium Tb 65 158.92534 8.3 1360 ± 4 3041 0.00009 _ 93 (III), 89 (IV) 1.10 Thallium TI 81 204.3833 11.85 303.5 1457+10 0.00006 147 (I), 95 (III) 1.44 Thorium Th 90 232.0381 11.7 1750 ca. 4790 0.0012 102 (IV) 1.11 Thulium Tm 69 168.9342 9.33 1545 ± 15 1727 0.00002 _ 87 (III) 1.11 Sn 50 118.71 7.3 231.9681 2270 0.004 140 294 (-IV), 93 (II), 71 (IV) 1.72 Tin Ti 22 47.88 4.51 1660 + 103287 0.45 80 (II), 76 (III), 68 (IV) 1.32 Titanium Tungston W 74 183.84 19.30 3410 ± 20 5660 0.007 136 70 (IV), 62 (VI) 1.40 92 Uranium U 238.029 19.1 1132.3+0.8 3818 0.0004 138 97 (IV), 80 (VI) 1.22 Vanadium V 23 50.9415 6.1 1890 + 103380 0.015 88 (II), 74 (III), 63 (IV), 59 (V) 1.45 Xenon Xe 54 131.29 *5.89 -111.9- 107.1±3 Ytterhium Yh 70 173.04 6.5 824 ± 5 1193 86 (III) 1.06 0.00027 _ Υ 39 92 (III) 88.90585 4.5 1523 + 83337 0.0028 _ 1.11 Yttrium 74 (II) Zinc 7n 30 65.39 7.2 419.58 907 0.013 133 1.66 Zirkonium 7r 40 91.224 6.5 1852 + 24377 0.022 79 (IV) 1.22


ELECTROCHEMICAL SERIES OF SOME NONMETALS (ALKALINE SOLUTION)

| Red $\rightleftharpoons^{\mathbb{R}}$ Ox + e | e° (Volt) | Red $\rightleftharpoons^{\mathbb{R}}$ Ox + e | e° (Volt) |
|--|-----------|---|-----------|
| Te²- ~ Te + 2e | - 1.14 | 2 l⁻⇄ l₂ + 2e | + 0.54 |
| $Se^{2-} \rightleftharpoons Se + 2e$ | - 0.92 | $2 \text{ Br} \xrightarrow{-} \text{Br}_2 + 2e$ | + 1.07 |
| $S^{2-} \rightleftharpoons S + 2e$ | - 0.48 | $2 \text{ CI} \xrightarrow{\longrightarrow} \text{CI}_2 + 2 \text{e}$ | + 1.36 |
| | | 2 F-→ F ₊ + 2e | + 287 |

COVALENT SINGLE-BOND RADIUSES (IN PM)

| H | * | 28 | 0 | 66 |
|---|----------------|-----|----|-----|
| | | | | |
| C | | 77 | S | 104 |
| S | ii | 117 | Se | 117 |
| G | ie | 122 | Te | 137 |
| S | in | 140 | F | 64 |
| N | 1 | 70 | CI | 99 |
| P |) | 110 | Br | 114 |
| А | \s | 121 | 1 | 133 |
| c | ih | 141 | | |

^{*} Determined from H-X bond distances





GENERAL FORMULAS FOR MIXING LIQUIDS

$$A = C - B$$

$$B = \frac{C(a - c)}{a - b}$$

$$C = B \frac{(a - b)}{a - c}$$

With:

A = weight of the original liquid a = its content in % by weight B = weight of the diluent C = weight of the prepared mixture

For water as diluent: b = 0

b = its content in % by weight

c = its content in % by weight

Example:

10 I of battery sulfuric acid with a density of

 $D^{20^{\circ}}_{4^{\circ}} = 1.28$. 1.28 is required. Available: concentrated

sulfuric acid with a density of $D_{4^{\circ}}^{20^{\circ}} = 1.84$ (= 97.5 weight%).

How much sulfuric acid and how much water are needed to prepare 10 l (= 12.8 kg) of battery sulfuric acid?

Calculation:

In the table 'Sulfuric acid' on page 52 we find:

 $D_{4^{\circ}}^{20^{\circ}} = 1.28$ equivalent to 37.36 weight%.

$$B = \frac{C (a - c)}{a - b} = \frac{12.80 (97.50 - 37.36)}{97.50 - 0} = 7.895 \text{ kg diluent}$$
(water)

Consequently, 4.905 kg (= 2.666 l) of concentrated sulfuric acid with a density of $D_{4^{\circ}}^{20^{\circ}}$ = 1.84 must be added to 7.895 kg (= I) of water to yield 10 I of battery acid with a density of $D_{4^{\circ}}^{20^{\circ}} = 1.28$.

CONVERSION TABLE FOR WATER HARDNESS UNITS

| | Alkaline earth ions mmol/l | Alkaline earth ions mval/l | German degree °d | ppm CaCO ₃ | English degree °e | French degree °f |
|---------------------------------|----------------------------------|----------------------------------|------------------------|--------------------------|-------------------------|------------------------|
| 1 mmol/l Alkaline earth ions | 1.00 | 2.00 | 5.60 | 100.00 | 7.02 | 10.00 |
| 1 mval/l Alkaline earth ions | 0.50 | 1.00 | 2.80 | 50.00 | 3.51 | 5.00 |
| 1 German degree | 0.18 | 0.357 | 1.00 | 17.80 | 1.25 | 1.78 |
| 1 ppm CaCO | 0.01 | 0.020 | 0.056 | 1.00 | 0.0702 | 0.10 |
| 1 English degree | 0.14 | 0.285 | 0.798 | 14.30 | 1.00 | 1.43 |
| 1 French degree | 0.10 | 0.200 | 0.560 | 10.00 | 0.702 | 1.00 |

MIXTURE RULE

Example:

Sulfuric acid with a density of $D_{4^{\circ}}^{20^{\circ}} = 1.520$ is to be prepared from sulfuric acid with a density of $D_{4^{\circ}}^{20^{\circ}} = 1.435$ and sulfuric acid of $D_{4^{\circ}}^{20^{\circ}} = 1.824$.

Calculation:

The table 'Sulfuric acid' (p. 52) informs that sulfuric acid with a density of $D_{A^\circ}^{20^\circ}=1.435=54.00$ weight% H_2SO_4 contains sulfuric acid with a density of $D_{A^\circ}^{20^\circ}=1.824=92.00$ weight% H_2SO_4 and that of $D_{A^\circ}^{20^\circ}=1.520=62.00$ weight% H_2SO_4 .

From this, form the mixing cross:

i.e. 30 parts by weight of 54.00 % sulfuric acid must be mixed with 8 parts by weight of 92.00 % sulfuric acid to yield sulfuric acid of 62.00 weight% $\rm H_2SO_{4^9}$ equivalent to D $^{2}\rm Q^{9^{\circ}}_{4^{\circ}}=$ 1.520.

PREPARATION OF DILUTE SOLUTIONS

Slowly stir the stated quantity of concentrated solution or solid KOH or NaOH, respectively, into water.

Caution: Strong development of heat may occur! Cool to room temperature, then make up to 1 liter with water. Store alkaline solutions in polyethylene bottles, because they attack glass.

As a rule of thumb, more concentrated solutions can be prepared by taking a multiple of the stated quantity.

Example:

6 mol/l HNO₃ from $6/2 \times 140 \text{ ml} = 420 \text{ ml} 65 \% \text{ HNO}_3$.

| | Solution to | be prepa | red | Original quantity to prepare 1 I of dilute solution | | |
|---|-------------|----------|-------|--|--------|--|
| | Weight% | Density | mol/l | Weight% | ml | |
| Acetic acid | 12 | 1.01 | 2 | 100 | 115 | |
| Nitric acid | 12 | 1.07 | 2 | 65 | 140 | |
| Hydrochloric acid | 7 | 1.03 | 2 | 36 | 165 | |
| Sulfuric acid | 9.5 | 1.06 | 1 | 96 | 56 | |
| Ammonia | 3.5 | 0.98 | 1 | 30 | 115 | |
| Potassium | 10.5 | 1.09 | 2 | 113 g solid KOH | (85%) | |
| hydroxide solution Sodium hydroxide solution | 7.5 | 1.08 | 2 | 80 g solid NaOH | (100%) | |

| | | | 6 | | | | | | Content of | |
|----------|--|---|----------|----------------|-----------|--------|---------|------------------|-----------------------|-----------------------|
| Cat. No. | Name | Formula | Solubili | ty in g/10 | 0 g H₂O a | t °C | | | the total solution at | the total solution at |
| | | Tormana | 0 | 20 | 40 | 60 | 80 | 100 | 20 °C. in % | |
| 1.01031 | Aluminum ammonium sulfate dodecahydrate | AINH ₄ (SO ₄) ₂ · 12 H ₂ O | 2.6 | 6.6 | 12.4 | 21.1 | 35.2 | 109.2 (95 °C) | 6.2 | 1.0459 (15.5 °C) |
| 1.01084 | Aluminum chloride hexahydrate | AICI ₃ · 6H ₂ O | 44.9 | 45.6 | 46.3 | 47.7 | 47.7 | - | 31.3 | - |
| 1.01063 | Aluminum nitrate nonahydrate | $AI(NO_3)_3 \cdot ^29H_2O$ | 61.0 | 75.4 | 89.0 | 108.0 | - | - | 43.0 | - |
| 1.01047 | Aluminum potassium sulfate dodecahydrate | $AIK(SO_4)_2 \cdot 12H_2O$ | 2.96 | 6.01 | 13.6 | 33.3 | 72.0 | 109.0 (90 °C) | 5.67 | 1.053 |
| 1.01102 | Aluminum sulfate octadecahydrate | Al ₂ (SO ₄) ₃ · 18H ₂ O | 31.2 | 36.4 | 45.6 | 58.0 | 73.0 | 89.0 | 26.7 | 1.308 |
| 1.01125 | Ammonium bromide | NĤ₄Br | 60.6 | 75.5 | 91.1 | 107.8 | 126.7 | 145.6 | 43.9 | - |
| 1.01145 | Ammonium chloride | NH₄CI | 29.7 | 37.6 | 46.0 | 55.3 | 65.6 | 77.3 | 27.3 | 1.075 |
| 1.01126 | Ammonium dihydrogen phosphate | NH ₄ H ₂ PO ₄ | 22.7 | 36.8 | 56.7 | 82.9 | 120.7 | 174.0 | 26.9 | - |
| 1.01131 | Ammonium hydrogen carbonate | NH4HCO3 | 11.9 | 21.2 | 36.6 | 59.2 | 109.2 | 355.0 | 17.5 | 1.07 |
| 1.01207 | di-Ammonium hydrogen phosphate | (NH ₄) ₂ HPO ₄ | 57.5 | 68.6 | 81.8 | 97.6 | (115.5) | - | 40.70 | 1.3436 (14.5 °C) |
| 1.01173 | Ammonium iodide | NH₄ I | 154.2 | 172.3 | 190.5 | 208.9 | 228.8 | 250.3 | 63.3 | - |
| 1.03792 | Ammonium iron(II) sulfate hexahydrate | $(NH_4)_2Fe(SO_4)_2 \cdot 6H_2O$ | 17.8 | 26.9 | 38.5 | 53.4 | 72.0 | _ | 21.2 | 1.18 |
| 1.01226 | Ammonium monovanadate | NH,VO, | - | 4.8 | 13.2 | - | - | - | _ | - |
| 1.01188 | Ammonium nitrate | NH ₄ NO ₂ | 118.5 | 187.7 | 283.0 | 415.0 | 610.0 | 1000.0 | 65.0 | 1.308 |
| 1.01217 | Ammonium sulfate | (NH ₄) ₂ SO ₄ | 70.4 | 75.4 | 81.2 | 87.4 | 94.1 | 102.0 | 43.0 | 1.247 |
| 1.01213 | Ammonium thiocyanate | NH,SCN | 115.0 | 163.0 | 235.0 | 347.0 | - | _ | 62.0 | - |
| 1.07838 | Antimony(III) chloride | SbCl ₂ | 601.6 | 931.5 | 1368.0 | 4531.0 | - | _ | 90.3 | _ |
| 1.01704 | Barium acetate | Ba(CH ₂ COO) ₂ | 58.0 | 72.0 | 79.0 | 74.0 | 74.0 | 74.0 | _ | - |
| 1.01719 | Barium chloride dihydrate | BaCl ₂ · 2H ₂ O | 30.7 | 35.7 | 40.8 | 46.4 | 52.5 | 58.7 | 26.3 | 1.28 |
| 1.01737 | Barium hydroxide octahydrate | Ba(OH) ₂ · 8H ₂ O | 1.5 | 3.5 | 8.2 | 21.0 | - | _ | 3.4 | 1.04 |
| 1.01729 | Barium nitrate | Ba(NO ₂) ₂ | 5.0 | 9.1 | 14.4 | 20.3 | 27.2 | 34.2 | 8.3 | 1.069 |
| 1.00163 | di-Boron trioxide | B ₂ O ₂ | 1.1 | 2.2 | 4.0 | 6.2 | 9.5 | 15.7 | 2.15 | _ |
| 1.00165 | Boric acid | H,BO, | 2.7 | 5.04 | 8.7 | 14.8 | 23.6 | 39.7 | 4.8 | 1.015 |
| 1.02019 | Cadmium nitrate tetrahydrate | Cd(NO ₂) ₂ · 4H ₂ O | - | 153.0 | 199.0 | - | - | - | 60.5 | - |
| 1.02027 | Cadmium sulfate hydrate | 3CdS0, 8H,0 | 75.5 | 76.7 | 79.3 | 82.0 | 84.6 | _ | 43.4 | 1.616 |
| 1.09325 | Calcium acetate | Ca(CH ₃ COO) ₂ | 37.4 | 34.7 | 33.2 | 32.7 | 33.5 | 29.7 | _ | - |
| 1.02382 | Calcium chloride dihydrate | CaCl ₂ · 2H ₂ O | - | - | 128.1 | 136.8 | 147.0 | 159.0 | _ | - |
| 1.02121 | Calcium nitrate tetrahydrate | Ca(NO ₃) ₂ · 4H ₂ O | 101.0 | 129.4 | 196.0 | - | - | - | 56.4 | - |
| 1.02161 | Calcium sulfate dihydrate | CaSO ₄ · 2H ₂ O | 0.18 | 0.20 | 0.21 | 0.20 | 0.19 | 0.16 | 0.20 | 1.001 |
| 1.02038 | Cesium chloride | CsCl | 161.0 | 187.0 | 208.0 | 230.0 | 250.0 | 271.0 | - | - |
| 1.02856 | Cesium nitrate | CsNO ₃ | 9.3 | 23.0 | 47.2 | 83.8 | 134.0 | 197.0 | _ | _ |
| 1.00229 | Chromium(VI) oxide | CrO ₃ | 163.0 | 166.7 | 171.0 | 176.0 | 189.0 | 199.0 | 62.50 | 1.7100 (16.5 °C) |
| 8.02540 | Cobalt chloride | CoCl ₂ | 74.5 | 91.9 | - | - | - | - | 47.9 | 1.52 |
| 1.02539 | Cobalt chloride hexahydrate | CoCl ₂ · 6H ₂ O | - | 62.35 | 68.6 | 78.3 | - | - | 38.4 | 1.49 |
| 1.02536 | Cobalt nitrate hexahydrate | Co(NO ₃) ₂ · 6H ₂ O | - | - | - | - | 525.1 | 537.0 | - | - |
| 1.02556 | Cobalt sulfate heptahydrate | CoSO ₄ · 7H ₂ O | 41.9 | 53.6 | 69.5 | - | - | - | 34.9 | - |
| 1.02739 | Copper(I) chloride | CuCl | - | 1.5 (25 °C) | - | - | - | - | 1,497 (25 °C) | - |
| 1.02733 | Copper(II) chloride dihydrate | CuCl ₂ · 2H ₂ O | 70.65 | 77.0 | 83.8 | 91.2 | 99.2 | 107.9 | 43.5 | 1. 55 |
| 1.02753 | Copper(II) nitrate trihydrate | Cu(NO ₃) ₂ · 3H ₂ O | - | - | 160.0 | 179.0 | 208.0 | (257.0) | - | - |
| 1.02790 | Copper(II) sulfate pentahydrate | CuSO ₄ · 5H ₂ O | 14.8 | 20.8 | 29.0 | 39.1 | 53.6 | 73.6 | 17.2 | 1.1965 |
| 1.02791 | Copper sulfate | CuSO ₄ | 25.5 | 36.2 | 48.0 | 60.0 | 70.0 | 83.0 | _ | - |
| 8.03945 | Iron(III) chloride | FeCl ₃ | 25.5 | 36.3 | 49.9 | - | - | - | 26.6 | - |
| 1.03943 | Iron(III) chloride hexahydrate | FeCl ₃ · 6H ₂ O | 83.5 | 100.0 | 126.0 | 169.5 | - | _ | 50.0 | - |

| | | | Solubility in g/100 g H _s O at °C | | | | | | Content of the total | Density of the total |
|----------|--|---|--|-----------------|--------|---------------------|-------------------|---------------------|----------------------------|----------------------|
| Cat. No. | Name | Formula | 0 | 20 | 40 | 60 | 80 | 100 | solution at 20 °C. in % | solution a |
| 1.03861 | Iron(II) chloride tetrahydrate | FeCl ₂ · 4H ₂ O | | _ | - | (90.5) | 100.0 | 107.5 | _ | _ |
| 1.03001 | mon(n) emonde tetranydrate | reci ₂ · 4n ₂ 0 | _ | _ | _ | (56 °C) | 100.0 | 107.5 | _ | _ |
| 1.03965 | Iron(II) sulfate heptahydrate | FeSO ₄ · 7H ₂ O | 15.6 | 26.6 | 40.3 | 47.6 | - | - | 21.0 | 1.225 |
| 1.03967 | Iron(II) sulfate monohydrate | FeSO, · H₃Ó | - | - | - | _ | 43.8 | (31.6) | _ | - |
| 3.07383 | Lead chloride | PbCl ₂ 2 | 0.67 | 0.99 | 1.45 | 1.98 | 2.6 | 3.3 | 0.98 | 1.007 |
| .07398 | Lead nitrate | Pb(NO ₂) ₂ | 36.4 | 52.2 | 69.4 | 88.0 | 107.5 | 127.3 | 34.3 | 1.40 |
| .05669 | Lithium bromide | LiBr | 143.0 | 177.0 | 205.0 | 224.0 | 245.0 | 266.0 | - | - |
| .05680 | Lithium carbonate | Li ₂ CO ₃ | - | 1.3 | - | - | - | - | 1.31 | - |
| 1.05677 | Lithium chloride monohydrate | LiĈI · H₂O | - | 82.8 | 90.4 | 100.0 | 113.0 | (127.5) | 45.3 | 1.29 |
| 1.11652 | Lithium hydroxide monohydrate | LiOH · Ĥ₃O | 12.0 | 12.4 | - | 13.4 | 14.9 | 17.9 | 11.0 | - |
| 3.18287 | Lithium iodide | Lil | 151.0 | 165.0 | 180.0 | - | - | 480.0 | - | - |
| .12230 | Lithium nitrate | LiNO ₂ | 48.0 | 76.0 | - | _ | _ | 227.0 | _ | _ |
| 1.05694 | Lithium sulfate monohydrate | LiSO, · H ₂ O | 36.2 | 34.8 | 33.5 | 32.3 | 31.5 | 31.0 | 25.6 | 1.23 |
| 1.05833 | Magnesium chloride hexahydrate | MqCl ₂ · 6H ₂ O | 52.8 | 54.6 | 57.5 | 60.7 | 65.9 | 72.7 | 35.3 | 1.331 |
| 1.05853 | Magnesium nitrate hexahydrate | $Mg(NO_3)_2 \cdot 6H_2O$ | 63.9 | 70.1 | 81.8 | 93.7 | - | - | 41.2 | 1.388 (25 °C) |
| .05886 | Magnesium sulfate heptahydrate | MqSO ₄ · 7H ₂ O | _ | 35.6 | 45.4 | _ | _ | _ | 26.25 | 1.31 |
| .05927 | Manganese(II) chloride tetrahydrate | MnCl ₂ · 4H ₂ 0 | 63.6 | 73.6 | 88.7 | (106.0) (58.1°C) | - | - | 42.4 | 1.499 |
| .05934 | Manganese(II) chloride dihydrate | MnCl ₂ · 2H ₂ O | _ | _ | _ | - | 110.5 | 115.0 | _ | _ |
| .05941 | Manganese(II) sulfate monohydrate | MnSO ₄ · H ₂ O | _ | _ | 60.0 | 58.6 | 45.5 | 35.5 | _ | _ |
| .04421 | Mercury(II) bromide | HgBr ₂ | - | 0.62 (25 °C) | (0.96) | 1.7 | 2.8 | 4.9 | 0.62 (25 °C) | - |
| .04419 | Mercury(II) chloride | HqCl ₂ | 4.29 | 6.6 | 9.6 | 13.9 | 24.2 | 54.1 | 6.2 | 1.052 |
| .06717 | Nickel chloride hexahydrate | NiCl ₂ · 6H ₂ O | 51.7 | 55.3 | - | - | - | _ | 35.6 | 1.46 |
| .06721 | Nickel nitrate hexahydrate | Ni(NO ₂) ₂ · 6H ₂ O | 79.2 | 94.1 | 118.8 | - | - | - | 48.5 | _ |
| .06727 | Nickel sulfate hexahydrate | Ni ₂ SO ₄ · 6H ₂ O | _ | _ | _ | 57.0 | _ | _ | _ | _ |
| .04820 | Potassium acetate | KCH,C00 | 217.0 | 256.0 | 323.0 | 350.0 | - | 380.0 | _ | - |
| .04912 | Potassium bromate | KBrO ₂ | 3.1 | 6.8 | 13.1 | 22.0 | 33.9 | 49.7 | 6.4 | 1.048 |
| 1.04905 | Potassium bromide | KBr | 54.0 | 65.8 | 76.1 | 85.9 | 95.3 | 104.9 | 39.7 | 1.370 |
| .04928 | Potassium carbonate | K,CO, | 106.0 | 110.0 | 117.0 | 127.0 | 140.0 | 156.0 | _ | - |
| .04944 | Potassium chlorate | KČIO ₃ | 3.3 | 7.3 | 14.5 | 25.9 | 39.7 | 56.2 | 6.8 | 1.042 |
| .04936 | Potassium chloride | KCI | 28.2 | 34.2 | 40.3 | 45.6 | 51.0 | 56.2 | 25.5 | 1.174 |
| .04952 | Potassium chromate | K ₂ CrO ₄ | 59.0 | 63.7 | 67.0 | 70.9 | 75.1 | 79.2 | 38.9 | 1.378 |
| 1.04967 | Potassium cyanide | KĆN [↑] | (63.0) | 71.6 (25 °C) | - | 81.0 (50 °C) | (95.0) (75 °C) | 122.0 (103.3 °C) | 41.73 (25 °C) | - |
| .04864 | Potassium dichromate | K ₂ Cr ₂ O ₇ | 4.7 | 12.5 | 26.3 | 45.6 | 73.0 | 103.0 | 11.1 | 1.077 |
| 1.04873 | Potassium dihydrogen phosphate | KĤ ₃ PÔ, | 14.3 | 22.7 | 33.9 | 48.6 | 68.0 | - | 18.5 | - |
| .05057 | Potassium disulfite | K ₂ S ₂ O ₂ | 27.5 | 44.9 | 63.9 | 85.0 | 108.0 | 133.0 | 30.99 | - |
| 1.19238 | Potassium hexachloroplatinate(IV) | K ₂ [Pt(CI) ₆] | 0.74 | 1.1 | 1.7 | 2.6 | 3.8 | 5.2 | _ | _ |
| 1.04984 | Potassium hexacyanoferrate(II) trihydrate | K_4 [Fe(CN) ₆] · 3H ₂ O | 15.0 | 28.9 | 42.7 | 56.0 | 68.9 | (82.7) | 22.4 | 1.16 |
| 1.04973 | Potassium hexacyanoferrate(III) | K ₃ [Fe(CN) ₆] | 29.9 | 46.0 | 59.5 | 70.9 | 81.8 | 91.6 | 31.5 | 1.18 |
| .04854 | Potassium hydrogen carbonate | KHCO. | 22.6 | 33.3 | 45.3 | 60.0 | - | - | 24.98 | 1.18 |
| .05099 | di-Potassium hydrogen phosphate trihydrate | K ₂ HPO ₄ · 3H ₂ O | - | 159.0 | 212.5 | - (50 °C) | – (75 °C) | - | 61.4 | - |
| .05104 | di-Potassium hydrogen phosphate | K₂HPO₄ | - | - | - | 266.0 | - | - | - | - |
| 1.04885 | Potassium hydrogen sulfate | KHSO, | 36.3 | 51.4 | 67.3 | - | - | 121.6 | 33.95 | - |
| 1.05002 | Potassium hydroxide monohydrate | KOH · H₂O | | _ | 136.4 | 147.0 | 160.0 | 178.0 | _ | _ |

| COIVII | OUNDS IN WAILI | | 1 -11 | /// LIV | AIOIII | _ | | | | |
|----------|--|---|-------|---------|----------|-----------------|---------|---------|----------------------------------|--|
| Cat. No. | Name | Formula | | , , | 00 g H₂O | | | | Content of the total solution at | Density of the total solution at |
| | | | 0 | 20 | 40 | 60 | 80 | 100 | 20 °C. in % | 20 °C. in % |
| 1.05051 | Potassium iodate | KIO ₂ | 4.7 | 8.1 | 12.9 | 18.5 | 24.8 | 32.3 | 7.5 | 1.064 |
| 1.05051 | Potassium iodide | KI | 127.8 | 144.5 | 161.0 | 176.2 | 191.5 | 208.0 | 59.1 | 1.71 |
| 1.05043 | Potassium nitrate | KNO. | 13.3 | 31.7 | 63.9 | 109.9 | 169.0 | 245.2 | 24.1 | 1.16 |
| 1.05003 | di-Potassium oxalate monohydrate | K ₂ C ₂ O ₄ · H ₂ O | - | 35.9 | - 03.3 | - | - | - | 26.4 | - |
| 1.05073 | Potassium perchlorate | KCIO ₄ | 0.76 | 1.7 | 3.6 | 7.2 | 13.4 | 22.2 | 1.7 | 1.008 |
| 1.05076 | Potassium periodate | KIO ₄ | 0.76 | 0.4 | 0.9 | 2.2 | 4.4 | 7.9 | 0.418 | 1.008 |
| 1.05079 | Potassium periodate Potassium permanganate | KMnO, | 2.8 | 6.4 | 12.6 | 22.4 | 4.4 | 7.9 | 6.0 | 1.04 |
| 1.05082 | Potassium permanganate Potassium peroxodisulfate | K ₂ S ₂ O ₈ | 0.18 | 0.5 | 1.1 | | _ | _ | 0.468 | - |
| | | | | 11.1 | | | _ | 24.1 | | |
| 1.05153 | Potassium sulfate | K₂SO₄ KSCN | 7.3 | 218.0 | 14.8 | 18.2 | 21.3 | 24.1 | 10.0 | 1.0807 |
| 1.05125 | Potassium thiocyanate | | 177.0 | | | _ | _ | | 68.55 | 1.42 |
| 1.07615 | Rubidium chloride | RbCl | 70.6 | 83.6 | - | | - | 128.0 | | - |
| 1.06267 | Sodium acetate trihydrate | NaCH ₃ COO · 3H ₂ O | 36.3 | 46.4 | 65.4 | 138.0 (58°C) | _ | - | 31.7 | 1.17 |
| 1.06363 | Sodium bromide | NaBr | - | - | - | 118.0 | 118.3 | 121.2 | - | - |
| 1.06391 | Sodium carbonate decahydrate | $Na_2CO_3 \cdot 10H_2O$ | 6.86 | 21.7 | - | _ | - | - | 17.8 | 1.1941 |
| 1.06386 | Sodium carbonate monohydrate | $Na_2CO_3 \cdot H_2O$ | - | - | 48.9 | 46.2 | 44.5 | 44.5 | - | - |
| 1.06392 | Sodium carbonate | Na ₂ CO ₃ | 7.1 | 21.4 | 48.5 | 46.5 | 45.8 | 45.5 | _ | - |
| 1.06423 | Sodium chlorate | NaClO ₃ | 80.5 | 98.8 | 115.2 | (138.0) | (167.0) | 204.0 | 49.7 | - |
| 1.06404 | Sodium chloride | NaCl | - | 35.9 | 36.4 | 37.1 | 38.1 | 39.2 | 26.4 | 1.201 |
| 1.06425 | Sodium chromate tetrahydrate | $Na_2CrO_4 \cdot 4H_2O$ | - | - | 95.3 | 115.1 | - | - | - | - |
| 1.06427 | Sodium chromate | Na ₂ CrO ₄ | - | - | - | _ | 124.0 | 125.9 | _ | - |
| 1.06336 | Sodium dichromate dihydrate | Na ₂ Cr ₂ O ₋₇ · 2H ₂ O | 163.2 | 180.2 | 220.5 | 283.0 | 385.0 | - | 64.3 | - |
| 1.06345 | Sodium dihydrogen phosphate dihydrate | $NaH_2PO_4 \cdot 2H_2O$ | 57.7 | 85.2 | 138.2 | - | - | - | 46.0 | - |
| 1.06370 | Sodium dihydrogen phosphate | NaH ₂ PO ₄ | - | - | - | 179.3 | 207.3 | 284.4 | - | - |
| 1.06591 | tetra-Sodium diphosphate decahydrate | $Na_4P_2O_7 \cdot 10H_2O$ | 2.7 | 5.5 | 12.5 | 21.9 | 30.0 | 40.3 | 5.2 | 1.05 |
| 1.06528 | Sodium disulfite | $Na_2S_2O_5$ | - | 65.3 | 71.1 | 79.9 | 88.7 | (100.0) | 39.5 | - |
| 1.06449 | Sodium fluoride | NaF | (3.6) | 4.1 | - | - | - | - | 3.94 | 1.04 |
| 1.06329 | Sodium hydrogen carbonate | NaHCO ₃ | 6.89 | 9.6 | 12.7 | 16.0 | 19.7 | 23.6 | 8.76 | 1.08 |
| 1.06579 | di-Sodium hydrogen phosphate dodecahydrate | Na ₂ HPO ₄ · 12H ₂ O | 1.63 | 7.7 | - | - | - | - | 7.2 | 1.08 |
| 1.06575 | di-Sodium hydrogen phosphate heptahydrate | Na ₂ HPO ₄ · 7H ₂ O | - | - | 55.0 | - | - | - | - | - |
| 1.06580 | di-Sodium hydrogen phosphate dihydrate | Na ₂ HPO ₄ · 2H ₂ O | - | - | - | 83.0 | 92.4 | - | - | - |
| 1.06586 | di-Sodium hydrogen phosphate | Na ₂ HPO ₄ | - | - | - | - | - | 104.1 | - | - |
| 1.06466 | Sodium hydroxide monohydrate | NaOH · H ₂ O | - | 109.2 | 126.0 | 178.0 | - | - | 52.2 | 1.55 |
| 1.06498 | Sodium hydroxide | NaOH | - | - | - | - | 313.7 | 341.0 | - | - |
| 1.06525 | Sodium iodate | NalO ₂ | 2.5 | 9.1 | - | 23.0 | 27.0 | 32.8 | - | - |
| 1.06523 | Sodium iodide | Nal | - | _ | - | _ | 295.0 | 303.0 | - | _ |
| 1.06537 | Sodium nitrate | NaNO ₂ | 70.7 | 88.3 | 104.9 | 124.7 | 148.0 | 176.0 | 46.8 | 1.38 |
| 1.06549 | Sodium nitrite | NaNO ₂ | 73.0 | 84.5 | 95.7 | 112.3 | 135.5 | 163.0 | 45.8 | 1.33 |
| 1.06564 | Sodium perchlorate monohydrate | NaClO ₄ · H ₂ O | 167.0 | 181.0 | 243.0 | - | - | - | 64.4 | 1.757 |
| 1.06578 | tri-Sodium phosphate dodecahydrate | Na ₂ PO ₄ · 12H ₂ O | 1.5 | 12.1 | 31.0 | 55.0 | 81.0 | 108.0 | 10.8 | 1.106 |
| 1.06648 | Sodium sulfate decahydrate | Na ₃ SO ₄ · 10H ₂ O | 4.56 | 19.2 | - | - | - | - | 16.1 | 1.150 |
| 1.06649 | Sodium sulfate | Na ₂ SO ₄ | - | - | 48.1 | 45.3 | 43.1 | 42.3 | _ | - |
| 1.06638 | Sodium sulfide nonahydrate | Na ₂ S · 9H ₂ O | 12.4 | 18.8 | 29.0 | - | - | - | 15.8 | 1.18 |
| 1.06657 | Sodium sulfite | Na ₂ SO ₂ | - | - | 37.0 | 33.2 | 29.0 | 26.6 | - | - |
| 1.06310 | di-Sodium tetraborate | Na ₂ B ₄ O ₇ | 1.2 | 2.7 | 6.0 | 20.3 | 31.5 | 52.5 | _ | _ |
| 1.06516 | Sodium thiosulfate pentahydrate | Na ₂ S ₂ O ₂ · 5H ₂ O | 52.5 | 70.1 | 102.6 | - | - | - | 41.2 | 1.39 |
| 1.01512 | Silver nitrate | AqNO ₂ | 115.0 | 219.2 | 334.8 | 471.0 | 652.0 | 1024.0 | 68.6 | 2.18 |
| | | Ag ₂ SO ₄ | 0.57 | 0.79 | 0.98 | 1.15 | 1.3 | 1.5 | 0.75 | |

SOLUBILITY OF INORGANIC COMPOUNDS IN WATER

| Cat. No. | Name | Formula | Solubili | ty in g/10 | 00 g H₂O | at °C | | | Content of the total solution at | the total |
|---------------|------------------------------|---|----------|------------|----------|-------|-------|-------|----------------------------------|-----------|
| Cat. No. Name | Name | Torniula | 0 | 20 | 40 | 60 | 80 | 100 | 20 °C. in % | |
| 1.07865 | Strontiumchlorid-Hexahydrat | SrCl ₂ · 6H ₂ O | 44.1 | 53.9 | 66.6 | 85.2 | - | - | 35.0 | 1.39 |
| 1.07876 | Strontiumhydroxid-Octahydrat | Sr(0H), · 8H,0 | 0.35 | 0.7 | 1.5 | 3.1 | 7.0 | 24.2 | 0.69 | _ |
| 1.07872 | Strontiumnitrat | Sr(NO ₃) ₂ | - | - | 91.2 | 94.2 | 97.2 | 101.2 | - | - |
| 8.18150 | Tin(II) chloride | SnCl ₂ | 83.9 | 269.8 | - | - | - | - | 72.96 | 2.07 |
| | | - | | (15°C) | | | | | (15°C) | |
| 1.08144 | Thallium(I) nitrate | TINO ₃ | 3.81 | 9.5 | 20.9 | 46.2 | 111.0 | 413.0 | 8.7 | - |
| 8.18631 | Zinc bromide | ZnBr ₂ | 390.0 | 440.0 | - | 620.0 | 640.0 | 670.0 | - | _ |
| 1.08816 | Zinc chloride | ZnCl ₂ | - | - | 453.0 | 488.0 | 541.0 | - | - | - |
| 1.08836 | Zinc nitrate hexahydrate | $Zn(NO_3)_3 \cdot 6H_2O$ | 92.7 | 118.3 | - | - | - | - | 54.2 | 1.67 |
| 1.08833 | Zinc nitrate tetrahydrate | Zn(NO ₃) ₂ · 4H ₂ 0 | - | - | 211.5 | - | - | - | - | - |
| 1.08883 | Zinc sulfate heptahydrate | ZnSO₄ · 7H₂O Î | 41.6 | 53.8 | - | - | - | - | 35.0 | 1.47 |
| 1 00000 | Zina sulfata manahudrata | 7×50 . U 0 | | | | 7C E | CC 7 | CO E | | |



SOLUBILITY PRODUCTS OF SLIGHTLY SOLUBLE INORGANIC COMPOUNDS

| SOLOBLE INORGAN | ic colvii ooli | |
|------------------------|--|--|
| Substance | Formula | Solubility product at given temperature, in [mol/l] |
| Aluminum hydroxide | AI(OH) ₃ | 4.00 x 10 ⁻¹³ (15°) 1.50 x 10 ⁻¹⁵ (18°) 3.70 x 10 ⁻¹⁵ (25°) |
| Arsenic(III) sulfide | As_2S_3 | 4.00 x 10 ⁻²⁹ (18°) |
| Barium carbonate | BaCO ₃ | 7.00 x 10 ⁻⁹ (16°) 8.10 x 10 ⁻⁹ (25°) |
| Barium chromate | BaCrO ₄ | 1.60 x 10 ⁻¹⁰ (18°) 2.40 x 10 ⁻¹⁰ (28°) |
| Barium fluoride | BaF ₂ | 1.60 x 10 ⁻⁶ (10°) 1.70 x 10 ⁻⁶ (18°) |
| Barium oxalate | $BaC_2O_4 \cdot 2H_2O$ | 1.20 x 10 ⁻⁷ (18°) |
| Barium sulfate | BaSO ₄ | 8.70 x 10 ⁻¹¹ (18°) 1.08 x 10 ⁻¹⁰ (25°) 1.98 x 10 ⁻¹⁰ (50°) |
| Beryllium hydroxide | Be(OH) ₂ | 2.70 x 10 ⁻¹⁹ (25°) |
| Bismuth hydroxide | Bi(OH) ₃ | 4.30 x 10 ⁻³¹ (18°) |
| Bismuth oxide chloride | BiOCI | 1.60 x 10 ⁻³¹ (25°) |
| Bismuth sulfide | Bi ₂ S ₃ | 1.60 x 10 ⁻⁷² (18°) |
| Cadmium carbonate | CdCO ₃ | 2.50 x 10 ⁻¹⁴ (25°) |
| Cadmium oxalate | $CdC_2O_4 \cdot 3H_2O$ | 1.53 x 10 ⁻⁸ (18°) |
| Cadmium sulfide | CdS | 3.60 x 10 ⁻²⁹ (18°) |
| Calcium carbonate | CaCO ₃ | 4.80 x 10 ⁻⁹ (25°) |
| Calcium fluoride | CaF ₂ | 3.40 x 10 ⁻¹¹ (18°) 3.95 x 10 ⁻¹¹ (26°) |
| Calcium hydroxide | Ca(OH) ₂ | 5.47 x 10 ⁻⁶ (18°) |
| Calcium oxalate | CaC₂O₄ · H₂O | 1.78 x 10 ⁻⁹ (18°) 2.57 x 10 ⁻⁹ (25°) |
| Calcium phosphate | Ca ₃ (PO ₄) ₂ | 1.00 x 10 ⁻²⁵ (25°) |
| Calcium sulfate | CaSO ₄ | 6.10 x 10 ⁻⁵ (10°) 2.45 x 10 ⁻⁵ (25°) |
| Calcium tartrate | CaC ₄ H ₄ O ₆ · 2H ₂ O | 7.70 x 10 ⁻⁷ (25°) |
| Cobalt(II) carbonate | CoCO ₃ | 1.00 x 10 ⁻¹² (25°) |
| Cobalt(II) sulfide | CoS | 1.90 x 10 ⁻²⁷ (20°) |
| Copper(I) bromide | CuBr | 4.15 x 10 ⁻⁸ (18–20°) |
| Copper(II) carbonate | CuCO₃ | 1.37 x 10 ⁻¹⁰ (25°) |
| Copper(I) chloride | CuCl | 1.02 x 10 ⁻⁶ (18-20°) |
| Copper(II) hydroxide | Cu(OH) ₂ | 5.60 x 10 ⁻²⁰ (25°) 5.06 x 10 ⁻¹² (18–20°) |
| Copper(I) iodide | Cul | 5.06 x 10 ⁻¹² (18–20°) |

SOLUBILITY PRODUCTS OF SLIGHTLY SOLUBLE INORGANIC COMPOUNDS

| SOLOBLE INONGANIC COMI OUNDS | | | | | | | |
|------------------------------|-----------------------------------|--|----------------------------|--|--|--|--|
| Substance | Formula | Solubility pro at given temperature, | | | | | |
| Copper(I) sulfide | Cu ₂ S | 2.00 x 10 ⁻⁴⁷ | (18°) | | | | |
| Copper(II) sulfide | CuS | 8.00 x 10 ⁻⁴⁵ | (18°) | | | | |
| Copper(I) thiocyanate | CuSCN | 1.60 x 10 ⁻¹¹ | (18°) | | | | |
| Iron(II) carbonate | FeCO ₃ | 2.50 x 10 ⁻¹¹ | (20°) | | | | |
| Iron(II) hydroxide | Fe(OH) ₂ | 1.64 x 10 ⁻¹⁴ | (18°) | | | | |
| Iron(III) hydroxide | Fe(OH) ₃ | 1.10 x 10 ⁻³⁶ | (18°) | | | | |
| Lanthanum hydroxide | La(OH) ₃ | ~ 10 ⁻²⁰ | (25°) | | | | |
| Lead bromide | PbBr ₂ | 3.90 x 10 ⁻⁵ | (25°) | | | | |
| Lead carbonate | PbCO ₃ | 3.30 x 10 ⁻¹⁴ | (18°) | | | | |
| Lead chloride | PbCl ₂ | 2.12 x 10 ⁻⁵ | (25°) | | | | |
| Lead chromate | PbCrO ₄ | 1.77 x 10 ⁻¹⁴ | (25°) | | | | |
| Lead fluoride | PbF ₂ | 2.70 x 10 ⁻⁸ 3.20 x 10 ⁻⁸ | (9°) (18°) | | | | |
| Lead iodate | Pb(IO ₃) ₂ | 5.30 x 10 ⁻¹⁴ 1.20 x 10 ⁻¹³ 2.60 x 10 ⁻¹³ | (9.2°) (18°) (25.8°) | | | | |
| Lead iodide | Pbl ₂ | 7.50 x 10 ⁻⁹ 1.40 x 10 ⁻⁹ | (15°) (25°) | | | | |
| Lead oxalate | PbC ₂ O ₄ | 2.74 x 10 ⁻¹¹ | (18°) | | | | |
| Lead sulfate | PbSO₄ | 1.06 x 10 ⁻⁸ | (18°) | | | | |
| Lead sulfide | PbS | 3.40 x 10 ⁻²⁸ | (18°) | | | | |
| Lithium carbonate | Li ₂ CO ₃ | 1.70 x 10 ⁻³ | (25°) | | | | |
| Magnesium ammonium phosphate | MgNH ₄ PO ₄ | 2.50 x 10 ⁻¹³ | (25°) | | | | |
| Magnesium carbonate | MgCO ₃ | 2.60 x 10 ⁻⁵ | (12°) | | | | |
| Magnesium fluoride | MgF ₂ | 7.10 x 10 ⁻⁹ | (18°) | | | | |
| Magnesium hydroxide | Mg(OH) ₂ | 1.20 x 10 ⁻¹¹ | (18°) | | | | |
| Manganese carbonate | MnCO ₃ | 8.80 x 10 ⁻¹⁰ | (18°) | | | | |
| Manganese sulfide | MnS | 7.00 x 10 ⁻¹⁶ | (18°) | | | | |
| Mercury(I) bromide | Hg ₂ Br ₂ | 1.30 x 10 ⁻²¹ | (25°) | | | | |
| Mercury(I) chloride | Hg ₂ Cl ₂ | 2.00 x 10 ⁻¹⁸ | (25°) | | | | |
| Mercury(I) chromate | Hg₂CrO₄ | 2.00 x 10 ⁻⁹ | (25°) | | | | |
| Mercury(I) cyanide | Hg ₂ (CN) ₂ | 5.00 x 10 ⁻⁴⁰ | (25°) | | | | |
| Mercury(I) iodide | Hg ₂ l ₂ | 1.20 x 10 ⁻²⁸ | (25°) | | | | |
| Mercury(II) iodide | Hgl ₂ | 3.20 x 10 ⁻²⁹ | (25°) | | | | |
| Mercury(I) oxide | Hg ₂ O | 1.60 x 10 ⁻²³ | (25°) | | | | |
| Mercury(II) oxide | HgO | 1.70 x 10 ⁻²⁶ | (25°) | | | | |

SOLUBILITY PRODUCTS OF SLIGHTLY SOLUBLE INORGANIC COMPOUNDS

| Substance | Formula | Solubility product at given temperature, in [mol/l] |
|------------------------------------|--|--|
| Mercury(I) sulfide | Hg₂S | 1.00 x 10 ⁻⁴⁷ (18°) |
| Mercury(II) sulfide | HgS | 3.00 x 10 ⁻⁵⁴ (18°) |
| Nickel(II) carbonate | NiCO ₃ | 1.35 x 10 ⁻⁷ (25°) |
| Nickel(II) hydroxide | Ni(OH) ₂ | 1.60 x 10 ⁻¹⁴ (25°) |
| Nickel(II) sulfide | NiS | 1.00 x 10 ⁻²⁶ (20°) |
| Potassium hexachloroplatinate (IV) | K ₂ PtCl ₆ | 1.10 x 10 ⁻⁵ (18°) |
| Potassium hydrogen tartrate | KHC ₄ H ₄ O ₆ | 3.80 x 10 ⁻⁴ (18°) |
| Potassium perchlorate | KCIO₄ | 1.07 x 10 ⁻² (25°) |
| Silver arsenate | Ag ₃ AsO ₄ | 1.00 x 10 ⁻¹⁹ (25°) |
| Silver bromide | AgBr | 4.10 x 10 ⁻¹³ (18°) 7.70 x 10 ⁻¹³ (25°) |
| Silver chloride | AgCl | 0.21 x 10 ⁻¹⁰ (4.7°) 0.37 x 10 ⁻¹⁰ (9.7°) 1.56 x 10 ⁻¹⁰ (25°) 13.2 x 10 ⁻¹⁰ (50°) 215 x 10 ⁻¹⁰ (100°) |
| Silver chromate | Ag ₂ CrO ₄ | 1.20 x 10 ⁻¹² (14.8°) 9.00 x 10 ⁻¹² (25°) |
| Silver iodide | Agl | 0.32×10^{-16} (13°) 1.50 × 10^{-16} (25°) |
| Silver sulfide | Ag_2S | 1.60 x 10 ⁻⁴⁹ (18°) |
| Silver thiocyanate | AgSCN | 0.49×10^{-12} (18°) 1.16 x 10^{-12} (25°) |
| Strontium carbonate | SrCO ₃ | 1.60 x 10 ⁻⁹ (25°) |
| Strontium fluoride | SrF ₂ | 2.80 x 10 ⁻⁹ (18°) |
| Strontium oxalate | SrC ₂ O ₄ | 5.60 x 10 ⁻⁸ (18°) |
| Strontium sulfate | SrSO ₄ | 2.80 x 10 ⁻⁷ (2.9°) 3.80 x 10 ⁻⁷ (17.4°) |
| Thallium(I) bromide | TIBr | 3.90 x 10 ⁻⁶ (25°) |
| Thallium(I) chloride | TICI | 1.90 x 10 ⁻⁴ (25°) |
| Thallium(I) iodide | TII | 5.80 x 10 ⁻⁸ (25°) |
| Thallium(III) hydroxide | TI(OH) ₃ | 1.40 x 10 ⁻⁵³ (25°) |
| Thallium(II) sulfide | TI ₂ S | 9.00 x 10 ⁻²³ (25°) |
| Thallium(I) thiocyanate | TISCN | 2.30 x 10 ⁻⁴ (25°) |
| Zinc carbonate | ZnCO ₃ | 6.00 x 10 ⁻¹¹ (25°) |
| Zinc hydroxide | Zn(OH) ₂ | 1.00 x 10 ⁻¹⁷ (25°) |
| Zinc sulfide, alpha | ZnS | 6.90 x 10 ⁻²⁶ (20°) |
| Zinc sulfide, beta | ZnS | 1.10 x 10 ⁻²⁴ (25°) |

SAMPLE PREPARATION

The more complicated the matrix, the more important the sample preparation! The better the sample preparation, the simpler the subsequent chromatographic separation!

The consequence of these two statements is: Sample preparation is a must!

Best performance with Merck products:

- Digestion systems and ultrapure digestion media for more 'difficult' dissolutions or processes to be performed in accordance with DIN* standards
- An extensive range of different reagents and solvents for improved precipitation, distribution and extraction
- The Extrelut® product range a porous kieselguhr and corresponding prepacked columns provide not only quicker but also better results than the conventional liquid-liquid extraction of aqueous matrices in the separation funnel
- LiChroLut® extraction columns for rapid, convenient and efficient extractions, enrichments and selective elutions
- LiChroCART® range of precolumns, which are an excellent alternative for online sample preparation for HPLC
- Inorganic membrane filters avoiding clogging of your HPLC columns by particles
- Derivatization substances for gas chromatography, i.e. for samples which are volatile by definition

Sample preparation with Merck for reliable performance!

*DIN = Deutsches Institut für Normung e.V.(German Institute of Standardization)

SULFURIC ACID H₂SO₄ , M = 98.08 g/mol

| 112504 , 1 | VI — 30 | Goo grilloi | | | |
|---------------------|---|------------------|---------------------|---|------------------|
| Density d 20°/4° | H ₂ SO ₄ weight% | content mol/l | Density d 20°/4° | H ₂ SO ₄ weight% | content mol/l |
| 1 000 | 0.3600 | 0.0266 | 1.245 | 33.22 | 4.216 |
| 1.000 | 0.2609 0.9855 | 0.0266 | 1.245 | 33.22 | 4.216 |
| 1.010 | 1.731 | 0.101 | 1.255 | 34.42 | 4.404 |
| 1.015 | 2.485 | 0.2595 | 1.260 | 35.01 | 4.498 |
| 1.020 | 3.242 | 0.2333 | 1.265 | 35.60 | 4.592 |
| 1.025 | 4.000 | 0.4180 | 1.270 | 36.19 | 4.686 |
| 1.030 | 4.746 | 0.4983 | 1.275 | 36.78 | 4.781 |
| 1.035 | 5.493 | 0.5796 | 1.280 | 37.36 | 4.876 |
| 1.040 | 6.237 | 0.6613 | 1.285 | 37.95 | 4.972 |
| 1.045 | 6.956 | 0.7411 | 1.290 | 38.53 | 5.068 |
| 1.050 | 7.704 | 0.8250 | 1.295 | 39.10 | 5.163 |
| 1.055 | 8.415 | 0.9054 | 1.300 | 39.68 | 5.259 |
| 1.060 | 9.129 | 0.9865 | 1.305 | 40.25 | 5.356 |
| 1.065 | 9.843 | 1.066 | 1.310 | 40.82 | 5.452 |
| 1.070 | 10.56 | 1.152 | 1.315 | 41.39 | 5.549 |
| 1.075 | 11.26 | 1.235 | 1.320 | 41.95 | 5.646 |
| 1.080 | 11.96 | 1.317 | 1.325 | 42.51 | 5.743 |
| 1.085 | 12.66 | 1.401 | 1.330 | 43.07 | 5.840 |
| 1.090 | 13.36 | 1.484 | 1.335 | 43.62 | 5.938 |
| 1.095 | 14.04 | 1.567 | 1.340 | 44.17 | 6.035 |
| 1.100 | 14.73 | 1.652 | 1.345 | 44.72 | 6.132 |
| 1.105 | 15.41 | 1.735 | 1.350 | 45.26 | 6.229 |
| 1.110 | 16.08 | 1.820 | 1.355 | 45.80 | 6.327 |
| 1.115 | 16.76 | 1.905 | 1.360 | 46.33 | 6.424 |
| 1.120 | 17.43 | 1.990 | 1.365 | 46.86 | 6.522 |
| 1.125 | 18.09 | 2.075 | 1.370 | 47.39 | 6.620 |
| 1.130 | 18.76 | 2.161 | 1.375 | 47.92 | 6.718 |
| 1.135 | 19.42 | 2.247 | 1.380 | 48.45 | 6.817 |
| 1.140 | 20.08 | 2.334 | 1.385 | 48.97 | 6.915 |
| 1.145 | 20.73 | 2.420 | 1.390 1.395 | 49.48 | 7.012 7.110 |
| 1.150 1.155 | 21.38 22.03 | 2.507 2.594 | 1.400 | 49.99 50.50 | 7.110 |
| 1.160 | 22.03 | 2.594 | 1.405 | 51.01 | 7.208 |
| 1.165 | 23.31 | 2.768 | 1.410 | 51.52 | 7.406 |
| 1.170 | 23.95 | 2.857 | 1.415 | 52.02 | 7.505 |
| 1.175 | 24.58 | 2.945 | 1.420 | 52.51 | 7.603 |
| 1.180 | 25.21 | 3.033 | 1.425 | 53.01 | 7.702 |
| 1.185 | 25.84 | 3.122 | 1.430 | 53.50 | 7.801 |
| 1.190 | 26.47 | 3.211 | 1.435 | 54.00 | 7.901 |
| 1.195 | 27.10 | 3.302 | 1.440 | 54.49 | 8.000 |
| 1.200 | 27.72 | 3.302 | 1.445 | 54.97 | 8.099 |
| 1.205 | 28.33 | 3.481 | 1.450 | 55.45 | 8.198 |
| 1.210 | 28.95 | 3.572 | 1.455 | 55.93 | 8.297 |
| 1.215 | 29.57 | 3.663 | 1.460 | 56.41 | 8.397 |
| 1.220 | 30.18 | 3.754 | 1.465 | 56.89 | 8.497 |
| 1.225 | 30.79 | 3.846 | 1.470 | 57.36 | 8.598 |
| 1.230 | 31.40 | 3.938 | 1.475 | 57.84 | 8.699 |
| 1.235 | 32.01 | 4.031 | 1.480 | 58.31 | 8.799 |
| 1.240 | 32.61 | 4.123 | 1.485 | 58.78 | 8.899 |

SULFURIC ACID H_2SO_4 , M = 98.08 g/mol

| Density | H ₂ SO ₄ co | ntent | Density | H ₂ SO ₄ cor | ntent |
|----------|-----------------------------------|--------|----------|------------------------------------|-------|
| d 20°/4° | weight% | mol/l | d 20°/4° | weight% | mol/ |
| 1.490 | 59.24 | 9.000 | 1.735 | 80.70 | 14.28 |
| 1.495 | 59.70 | 9.100 | 1.740 | 81.16 | 14.40 |
| 1.500 | 60.17 | 9.202 | 1.745 | 81.62 | 14.52 |
| 1.505 | 60.62 | 9.303 | 1.750 | 82.09 | 14.65 |
| 1.510 | 61.08 | 9.404 | 1.755 | 82.57 | 14.78 |
| 1.515 | 61.54 | 9.506 | 1.760 | 83.06 | 14.90 |
| 1.520 | 62.00 | 9.608 | 1.765 | 83.57 | 15.04 |
| 1.525 | 62.45 | 9.711 | 1.770 | 84.08 | 15.17 |
| 1.530 | 62.91 | 9.8136 | 1.775 | 84.61 | 15.31 |
| 1.535 | 63.36 | 9.916 | 1.780 | 85.16 | 15.46 |
| 1.540 | 63.81 | 10.02 | 1.785 | 85.74 | 15.61 |
| 1.545 | 64.26 | 10.12 | 1.790 | 86.35 | 15.76 |
| 1.550 | 64.71 | 10.23 | 1.795 | 86.99 | 15.92 |
| 1.555 | 65.15 | 10.33 | 1.800 | 87.69 | 16.09 |
| 1.560 | 65.59 | 10.43 | 1.805 | 88.43 | 16.27 |
| 1.565 | 66.03 | 10.54 | 1.810 | 89.23 | 16.47 |
| 1.570 | 66.47 | 10.64 | 1.815 | 90.12 | 16.68 |
| 1.575 | 66.91 | 10.74 | 1.820 | 91.11 | 16.91 |
| 1.580 | 67.35 | 10.85 | 1.821 | 91.33 | 16.96 |
| 1.585 | 67.79 | 10.96 | 1.822 | 91.56 | 17.01 |
| 1.590 | 68.23 | 11.06 | 1.823 | 91.78 | 17.06 |
| 1.595 | 68.66 | 11.16 | 1.824 | 92.00 | 17.11 |
| 1.600 | 69.09 | 11.27 | 1.825 | 92.25 | 17.17 |
| 1.605 | 69.53 | 11.38 | 1.826 | 92.51 | 17.17 |
| 1.610 | 69.96 | 11.48 | 1.827 | 92.77 | 17.28 |
| 1.615 | 70.39 | 11.59 | 1.828 | 93.03 | 17.34 |
| 1.620 | 70.82 | 11.70 | 1.829 | 93.33 | 17.40 |
| 1.625 | 71.25 | 11.80 | 1.830 | 93.64 | 17.47 |
| 1.630 | 71.67 | 11.91 | 1.831 | 93.94 | 17.54 |
| 1.635 | 72.09 | 12.02 | 1.832 | 94.32 | 17.62 |
| 1.640 | 72.52 | 12.13 | 1.833 | 94.72 | 17.70 |
| 1.645 | 72.95 | 12.24 | | | |
| 1.650 | 73.37 | 12.43 | | | |
| 1.655 | 73.80 | 12.45 | | 1900 | |
| 1.660 | 74.22 | 12.56 | | 100 | |
| 1.665 | 74.64 | 12.67 | | ALC: U | |
| 1.670 | 75.07 | 12.78 | | ALC: NO. | |
| 1.675 | 75.49 | 12.89 | | | |
| 1.680 | 75.92 | 13.00 | | | |
| 1.685 | 76.34 | 13.12 | - 9 | | |
| 1.690 | 76.77 | 13.23 | | | |
| 1.695 | 77.20 | 13.34 | - 6 | | |
| 1.700 | 77.63 | 13.46 | | | |
| 1.705 | 78.06 | 13.57 | | | |
| 1.710 | 78.49 | 13.69 | | | |
| 1.715 | 78.93 | 13.80 | - 0 | | |
| 1.720 | 79.37 | 13.92 | | | |
| 1.725 | 79.81 | 14.04 | | | |
| 1.730 | 80.25 | 14.16 | | | |

PHOSPHORIC ACID H₃PO₄, M = 97.99/mol

| 1131 041 1 | . 07. | 00/1110 |
|------------|----------|----------------|
| Density | H₃PO₄ co | ntent |
| d 20°/4° | weight% | mol/l |
| | | |
| 1.0038 | 1 | 0.102 |
| 1.0092 | 2 | 0.206 |
| 1.0146 | 3 | 0.312 |
| 1.0200 | 4 | 0.416 |
| 1.0255 | 5 | 0.523 |
| 1.0309 | 6 | 0.631 |
| 1.0365 | 7 | 0.740 |
| 1.0420 | 8 | 0.851 |
| 1.0476 | 9 | 0.962 |
| 1.0532 | 10 | 1.074 |
| 1.0590 | 11 | 1.189 |
| 1.0647 | 12 | 1.304 |
| 1.0705 | 13 14 | 1.420 |
| 1.0764 | | 1.538 |
| 1.0824 | 15 16 | 1.657 |
| 1.0884 | 17 | 1.777 |
| 1.1008 | 18 | 1.899 2.021 |
| 1.1071 | 19 | 2.147 |
| 1.1134 | 20 | 2.272 |
| 1.1199 | 21 | 2.400 |
| 1.1263 | 22 | 2.529 |
| 1.1329 | 23 | 2.659 |
| 1.1395 | 24 | 2.791 |
| 1.1462 | 25 | 2.924 |
| 1.1529 | 26 | 3.059 |
| 1.1597 | 27 | 3.195 |
| 1.1665 | 28 | 3.333 |
| 1.1735 | 29 | 3.473 |
| 1.1805 | 30 | 3.614 |
| 1.216 | 35 | 4.333 |
| 1.254 | 40 | 5.118 |
| 1.293 | 45 | 5.938 |
| 1.335 | 50 | 6.811 |
| 1.379 | 55 | 7.740 |
| 1.426 | 60 | 8.731 |
| 1.476 | 65 | 9.784 |
| 1.526 | 70 | 10.90 |
| 1.579 | 75 | 12.08 |
| 1.633 | 80 | 13.33 |
| 1.689 | 85 | 14.65 |
| 1.746 | 90 | 16.03 |
| 1.770 | 92 | 16.61 |
| 1.794 | 94 | 17.20 |
| 1.819 | 96 | 17.82 |
| 1.844 | 98 | 18.44 |
| 1.870 | 100 | 19.08 |



HYDROCHLORIC ACID HCI, M = 36.47 g/mol

| | | 9/ |
|----------------|--------------------|--------------------|
| Density | HCI co | ntent |
| d 20°/4° | weight% | mol/l |
| | _ | |
| 1.000 | 0.3600 | 0.09872 |
| 1.005 | 1.360 | 0.3748 |
| 1.010 | 2.364 | 0.6547 |
| 1.015 | 3.374 | 0.9391 |
| 1.020 | 4.388 | 1.227 |
| 1.025 | 5.408 | 1.520 |
| 1.030 | 6.433 | 1.817 |
| 1.035 | 7.464 | 2.118 |
| 1.040 | 8.490 | 2.421 |
| 1.045 | 9.510 | 2.725 |
| 1.050 | 10.52 | 3.029 |
| 1.055 | 11.52 | 3.333 |
| 1.060 | 12.51 | 3.638 |
| 1.065 | 13.50 | 3.944 |
| 1.070 | 14.49 ₅ | 4.253 |
| 1.075 | 15.48 ₅ | 4.565 |
| 1.080 | 16.47 | 4.878 |
| 1.085 | 17.45 | 5.192 |
| 1.090 | 18.43 | 5.509 ₅ |
| 1.095 | 19.41 | 5.829 |
| 1.100 | 20.39 | 6.150 |
| 1.105 | 21.36 | 6.472 |
| 1.110 | 22.33 | 6.796 |
| 1.115 | 23.29 | 7.122 |
| 1.120 | 24.25 | 7.449 |
| 1.125 | 25.22 | 7.782 |
| 1.130 | 26.20 | 8.118 |
| 1.135 | 27.18 | 8.459 |
| 1.140 | 28.18 | 8.809 |
| 1.145 | 29.17 | 9.159 |
| 1.150 1.155 | 30.14 | 9.505 |
| | 31.14 | 9.863 10.225 |
| 1.160 | 32.14 | |
| 1.165 1.170 | 33.16 34.18 | 10.595 10.97 |
| 1.175 | 35.20 | 11.34 |
| 1.180 | 36.23 | 11.73 |
| 1.185 | 37.27 | 12.11 |
| 1.185 | 38.32 | 12.11 |
| 1.195 | 39.37 | 12.90 |
| 1.198 | 40.00 | 13.14 |
| 1.130 | 70.00 | 13.14 |

NITRIC ACID HNO₃, M = 63.02 g/mol

| HNO_3 , $M = 63.02 \text{ g/mol}$ | | | | | |
|-------------------------------------|---------|---------|----------|----------------------|-------|
| Density HNO3 content | | | Density | HNO ₃ cor | itent |
| d 20°/4° | weight% | mol/l | d 20°/4° | weight% | mol/l |
| | | | | | |
| 1.000 | 0.3333 | 0.05231 | 1.260 | 42.14 | 8.426 |
| 1.005 | 1.255 | 0.2001 | 1.265 | 42.92 | 8.616 |
| 1.010 | 2.164 | 0.3468 | 1.270 | 43.70 | 8.808 |
| 1.015 | 3.073 | 0.4950 | 1.275 | 44.48 | 9.001 |
| 1.020 | 3.982 | 0.6445 | 1.280 | 45.27 | 9.195 |
| 1.025 | 4.883 | 0.7943 | 1.285 | 46.06 | 9.394 |
| 1.030 | 5.784 | 0.9454 | 1.290 | 46.85 | 9.590 |
| 1.035 | 6.661 | 1.094 | 1.295 | 47.63 | 9.789 |
| 1.040 | 7.530 | 1.243 | 1.300 | 48.42 | 9.990 |
| 1.045 | 8.398 | 1.393 | 1.305 | 49.21 | 10.19 |
| 1.050 | 9.259 | 1.543 | 1.310 | 50.00 | 10.39 |
| 1.055 | 10.12 | 1.694 | 1.315 | 50.85 | 10.61 |
| 1.060 | 10.97 | 1.845 | 1.320 | 51.71 | 10.83 |
| 1.065 | 11.81 | 1.997 | 1.325 | 52.56 | 11.05 |
| 1.070 | 12.65 | 2.148 | 1.330 | 53.41 | 11.27 |
| 1.075 | 13.48 | 2.301 | 1.335 | 54.27 | 11.49 |
| 1.080 | 14.31 | 2.453 | 1.340 | 55.13 | 11.72 |
| 1.085 | 15.13 | 2.605 | 1.345 | 56.04 | 11.96 |
| 1.090 | 15.95 | 2.759 | 1.350 | 56.95 | 12.20 |
| 1.095 | 16.76 | 2.913 | 1.355 | 57.87 | 12.44 |
| 1.100 | 17.58 | 3.068 | 1.360 | 58.78 | 12.68 |
| 1.105 | 18.39 | 3.224 | 1.365 | 59.69 | 12.93 |
| 1.110 | 19.19 | 3.381 | 1.370 | 60.67 | 13.19 |
| 1.115 | 20.00 | 3.539 | 1.375 | 61.69 | 13.19 |
| 1.120 | 20.79 | 3.696 | 1.380 | 62.70 | 13.73 |
| 1.125 | 21.59 | 3.854 | 1.385 | 63.72 | 14.01 |
| 1.130 | 22.38 | 4.012 | 1.390 | 64.74 | 14.29 |
| 1.135 | 23.16 | 4.171 | 1.395 | 65.84 | 14.23 |
| 1.140 | 23.10 | 4.330 | 1.400 | 66.97 | 14.88 |
| 1.145 | 24.71 | 4.489 | 1.405 | 68.10 | 15.18 |
| 1.145 | 25.48 | 4.469 | 1.410 | 69.23 | 15.16 |
| | 26.24 | 4.810 | 1.415 | 70.39 | 15.49 |
| 1.155 | | | | 70.39 | 16.14 |
| 1.160 | 27.00 | 4.970 | 1.420 | | |
| 1.165 | 27.76 | 5.132 | 1.425 | 72.86 | 16.47 |
| 1.170 | 28.51 | 5.293 | 1.430 | 74.09 | 16.81 |
| 1.175 | 29.25 | 5.455 | 1.435 | 75.35 | 17.16 |
| 1.180 | 30.00 | 5.618 | 1.440 | 76.71 | 17.53 |
| 1.185 | 30.74 | 5.780 | 1.445 | 78.07 | 17.90 |
| 1.190 | 31.47 | 5.943 | 1.450 | 79.43 | 18.28 |
| 1.195 | 32.21 | 6.107 | 1.455 | 80.88 | 18.68 |
| 1.200 | 32.94 | 6.273 | 1.460 | 82.39 | 19.09 |
| 1.205 | 33.68 | 6.440 | 1.465 | 83.91 | 19.51 |
| 1.210 | 34.41 | 6.607 | 1.470 | 85.50 | 19.95 |
| 1.215 | 35.16 | 6.778 | 1.475 | 87.29 | 20.43 |
| 1.220 | 35.93 | 6.956 | 1.480 | 89.07 | 20.92 |
| 1.225 | 36.70 | 7.135 | 1.485 | 91.13 | 21.48 |
| 1.230 | 37.48 | 7.315 | 1.490 | 93.49 | 22.11 |
| 1.235 | 38.25 | 7.497 | 1.495 | 95.46 | 22.65 |
| 1.240 | 39.02 | 7.679 | 1.500 | 96.73 | 23.02 |
| 1.245 | 39.80 | 7.863 | 1.501 | 96.98 | 23.10 |
| 1.250 | 40.58 | 8.049 | 1.502 | 97.23 | 23.18 |
| 1.255 | 41.36 | 8.237 | 1.503 | 97.49 | 23.25 |

NITRIC ACID HNO₃, M = 63.02 g/mol

| Density | HNO ₃ cor | ntent |
|----------|----------------------|-------|
| d 20°/4° | weight% | mol/l |
| 1.504 | 97.74 | 23.33 |
| | | |
| 1.505 | 97.99 | 23.40 |
| 1.506 | 98.25 | 23.48 |
| 1.507 | 98.50 | 23.56 |
| 1.508 | 98.76 | 23.63 |
| 1.509 | 99.01 | 23.71 |
| 1.510 | 99.26 | 23.79 |
| 1.511 | 99.52 | 23.86 |
| 1.512 | 99.77 | 23.94 |
| 1 513 | 100.0 | 24.01 |

SODIUM HYDROXIDE SOLUTION NaOH, M = 40.01 g/mol

| Density | Density NaOH content | | | NaOH content | |
|----------|----------------------|--------|----------|--------------------|-------|
| d 20°/4° | weight% | mol/l | d 20°/4° | weight% | mol/l |
| 1.000 | 0.159 | 0.0398 | 1.105 | 9.64 | 2.664 |
| 1.005 | 0.602 | 0.151 | 1.110 | 10.10 ⁵ | 2.802 |
| 1.010 | 1.0455 | 0.264 | 1.115 | 10.55 | 2.942 |
| 1.015 | 1.49 | 0.378 | 1.120 | 11.01⁵ | 3.082 |
| 1.020 | 1.94 | 0.494 | 1.125 | 11.46 | 3.224 |
| 1.025 | 2.39 | 0.611 | 1.130 | 11.92 | 3.367 |
| 1.030 | 2.84 | 0.731 | 1.135 | 12.37 | 3.510 |
| 1.035 | 3.29 | 0.851 | 1.140 | 12.83 | 3.655 |
| 1.040 | 3.745 | 0.971 | 1.145 | 13.28 | 3.801 |
| 1.045 | 4.20 | 1.097 | 1.150 | 13.73 | 3.947 |
| 1.050 | 4.655 | 1.222 | 1.155 | 14.18 | 4.095 |
| 1.055 | 5.11 | 1.347 | 1.160 | 14.64 | 4.244 |
| 1.060 | 5.56 | 1.474 | 1.165 | 15.09 | 4.395 |
| 1.065 | 6.02 | 1.602 | 1.170 | 15.54 | 4.545 |
| 1.070 | 6.47 | 1.731 | 1.175 | 15.99 | 4.697 |
| 1.075 | 6.93 | 1.862 | 1.180 | 16.44 | 4.850 |
| 1.080 | 7.38 | 1.992 | 1.185 | 16.89 | 5.004 |
| 1.085 | 7.83 | 2.123 | 1.190 | 17.34 | 5.160 |
| 1.090 | 8.28 | 2.257 | 1.195 | 17.80⁵ | 5.317 |
| 1.095 | 8.74 | 2.391 | 1.200 | 18.25 | 5.476 |
| 1.100 | 9.19 | 2.527 | 1.205 | 18.71 ⁵ | 5.636 |


SODIUM HYDROXIDE SOLUTION NaOH, M = 40.01 g/mol

| 3, | | | | | |
|----------|---------|--------|----------|--------------|-------|
| Density | NaOH co | ontent | Density | NaOH content | |
| d 20°/4° | weight% | mol/l | d 20°/4° | weight% | mol/l |
| | | | | | |
| 1.210 | 19.16 | 5.796 | 1.370 | 34.03 | 11.65 |
| 1.215 | 19.62 | 5.958 | 1.375 | 34.52 | 11.86 |
| 1.220 | 20.07 | 6.122 | 1.380 | 35.01 | 12.08 |
| 1.225 | 20.53 | 6.286 | 1.385 | 35.505 | 12.29 |
| 1.230 | 20.98 | 6.451 | 1.390 | 36.00 | 12.51 |
| 1.235 | 21.44 | 6.619 | 1.395 | 36.495 | 12.73 |
| 1.240 | 21.90 | 6.788 | 1.400 | 36.99 | 12.95 |
| 1.245 | 22.36 | 6.958 | 1.405 | 37.49 | 13.17 |
| 1.250 | 22.82 | 7.129 | 1.410 | 37.99 | 13.39 |
| 1.255 | 23.275 | 7.302 | 1.415 | 38.49 | 13.61 |
| 1.260 | 23.73 | 7.475 | 1.420 | 38.99 | 13.84 |
| 1.265 | 24.19 | 7.650 | 1.425 | 39.495 | 14.07 |
| 1.270 | 24.645 | 7.824 | 1.430 | 40.00 | 14.30 |
| 1.275 | 25.10 | 8.000 | 1.435 | 40.515 | 14.53 |
| 1.280 | 25.56 | 8.178 | 1.440 | 41.03 | 14.77 |
| 1.285 | 26.02 | 8.357 | 1.445 | 41.55 | 15.01 |
| 1.290 | 26.48 | 8.539 | 1.450 | 42.07 | 15.25 |
| 1.295 | 26.94 | 8.722 | 1.455 | 42.59 | 15.49 |
| 1.300 | 27.41 | 8.906 | 1.460 | 43.12 | 15.74 |
| 1.305 | 27.87 | 9.092 | 1.465 | 43.64 | 15.98 |
| 1.310 | 28.33 | 9.278 | 1.470 | 44.17 | 16.23 |
| 1.315 | 28.80 | 9.466 | 1.475 | 44.695 | 16.48 |
| 1.320 | 29.26 | 9.656 | 1.480 | 45.22 | 16.73 |
| 1.325 | 29.73 | 9.875 | 1.485 | 45.75 | 16.98 |
| 1.330 | 30.20 | 10.04 | 1.490 | 46.27 | 17.23 |
| 1.335 | 30.67 | 10.23 | 1.495 | 46.80 | 17.49 |
| 1.340 | 31.14 | 10.43 | 1.500 | 47.33 | 17.75 |
| 1.345 | 31.62 | 10.63 | 1.505 | 47.85 | 18.00 |
| 1.350 | 32.10 | 10.83 | 1.510 | 48.38 | 18.26 |
| 1.355 | 32.58 | 11.03 | 1.515 | 48.905 | 18.52 |
| 1.360 | 33.06 | 11.24 | 1.520 | 49.44 | 18.78 |
| 1.365 | 33.54 | 11.45 | 1.525 | 49.97 | 19.05 |
| | | | 1.530 | 50.50 | 19.31 |

POTASSIUM HYDROXIDE SOLUTION KOH, M = 56.11 g/mol

| KOH content | | Density | KOH con | tent |
|-------------|---|---|--|--|
| weight% | mol/l | d 20°/4° | weight% | mol/l |
| | | | | |
| | | | | 3.58 |
| | | | | 3.70 |
| 1.295 | 0.233 | 1.170 | 18.32 | 3.82 |
| 1.84 | 0.333 | 1.175 | 18.84 | 3.945 |
| | 0.4355 | 1.180 | | 4.07 |
| | 0.536 | | | 4.195 |
| | 0.6395 | 1.190 | 20.37 | 4.32 |
| | 0.774 | 1.195 | 20.88 | 4.45 |
| | | 1.200 | 21.38 | 4.57 |
| 5.12 | 0.954 | 1.205 | 21.88 | 4.70 |
| 5.66 | 1.06 | 1.210 | 22.38 | 4.83 |
| 6.20 | 1.17 | 1.215 | 22.88 | 4.955 |
| 6.74 | 1.27 | 1.220 | 23.38 | 5.08 |
| 7.28 | 1.38 | 1.225 | 23.87 | 5.21 |
| 7.82 | 1.49 | 1.230 | 24.37 | 5.34 |
| 8.36 | 1.60 | 1.235 | 24.86 | 5.47 |
| 8.89 | 1.71 | 1.240 | 25.36 | 5.60 |
| 9.43 | 1.82 | 1.245 | 25.85 | 5.74 |
| 9.96 | 1.94 | 1.250 | 26.34 | 5.87 |
| 10.49 | 2.05 | 1.255 | 26.83 | 6.00 |
| 11.03 | 2.16 | 1.260 | 27.32 | 6.135 |
| 11.56 | 2.28 | 1.265 | 27.80 | 6.27 |
| 12.08 | 2.39 | 1.270 | 28.29 | 6.40 |
| 12.61 | 2.51 | 1.275 | 28.77 | 6.54 |
| 13.14 | 2.62 | 1.280 | 29.25 | 6.67 |
| 13.66 | 2.74 | 1.285 | 29.73 | 6.81 |
| 14.19 | 2.86 | 1.290 | 30.21 | 6.95 |
| 14.705 | 2.975 | 1.295 | 30.68 | 7.08 |
| 15.22 | 3.09 | 1.300 | 31.15 | 7.22 |
| 15.74 | 3.21 | 1.305 | 31.62 | 7.36 |
| 16.26 | 3.33 | 1.310 | 32.09 | 7.49 |
| 16.78 | 3.45 | 1.315 | 32.56 | 7.63 |
| | weight% 0.197 0.743 1.295 1.84 2.38 2.93 3.48 4.03 4.58 5.12 5.66 6.20 6.74 7.28 7.82 8.36 8.89 9.43 9.96 10.49 11.03 11.56 12.08 12.61 13.14 13.66 14.19 14.705 15.22 15.74 | weight% mol/l 0.197 0.0351 0.743 0.133 1.295 0.233 1.84 0.333 2.38 0.4355 2.93 0.536 3.48 0.6395 4.03 0.774 4.58 0.848 5.12 0.954 5.66 1.06 6.20 1.17 6.74 1.27 7.28 1.38 7.82 1.49 8.36 1.60 8.89 1.71 9.43 1.82 9.96 1.94 10.49 2.05 11.03 2.16 11.56 2.28 12.08 2.39 12.61 2.51 13.14 2.62 13.66 2.74 14.19 2.86 14.705 2.975 15.22 3.09 15.74 3.21 16.26 3.33 | weight% mol/l d 20"/4" 0.197 0.0351 1.160 0.743 0.133 1.165 1.295 0.233 1.175 2.38 0.4355 1.180 2.93 0.536 1.185 3.48 0.6395 1.190 4.03 0.774 1.195 4.58 0.848 1.200 5.12 0.954 1.205 5.66 1.06 1.210 6.20 1.17 1.215 6.74 1.27 1.220 7.28 1.49 1.230 8.36 1.60 1.235 8.89 1.71 1.240 9.43 1.82 1.245 9.96 1.94 1.250 10.49 2.05 1.255 11.03 2.16 1.260 11.56 2.28 1.265 12.08 2.39 1.270 12.61 2.51 1.275 13.14 | weight% mol/l d 20°/4° weight% 0.197 0.0351 1.160 17.29 0.743 0.133 1.165 17.81 1.295 0.233 1.170 18.32 1.84 0.333 1.175 18.84 2.38 0.4355 1.180 19.35 2.93 0.536 1.185 19.86 3.48 0.6395 1.190 20.37 4.03 0.774 1.195 20.88 4.58 0.848 1.200 21.38 5.12 0.954 1.205 21.88 5.66 1.06 1.210 22.38 6.20 1.17 1.215 22.88 6.74 1.27 1.220 23.38 7.82 1.49 1.230 24.37 8.36 1.60 1.235 24.86 8.89 1.71 1.240 25.36 9.43 1.82 1.245 25.85 9.96 1.94 |





POTASSIUM HYDROXIDE SOLUTION KOH, M = 56.11 g/mol

| | | 3, | | | |
|---------------------|--------------------|----------------|---------------------|-------------------|----------------|
| Density d 20°/4° | KOH con weight% | itent mol/l | Density d 20°/4° | KOH co weight% | ntent mol/l |
| | | | | | |
| 1.320 | 33.03 | 7.77 | 1.425 | 42.60 | 10.82 |
| 1.325 | 33.50 | 7.91 | 1.430 | 43.04 | 10.97 |
| 1.330 | 33.97 | 8.05 | 1.435 | 43.48 | 11.12 |
| 1.335 | 34.43 | 8.19 | 1.440 | 43.92 | 11.28 |
| 1.340 | 34.90 | 8.335 | 1.445 | 44.36 | 11.42 |
| 1.345 | 35.36 | 8.48 | 1.450 | 44.79 | 11.58 |
| 1.350 | 35.82 | 8.62 | 1.455 | 45.23 | 11.73 |
| 1.355 | 36.28 | 8.76 | 1.460 | 45.66 | 11.88 |
| 1.360 | 36.735 | 8.905 | 1.465 | 46.095 | 12.04 |
| 1.365 | 37.19 | 9.05 | 1.470 | 46.53 | 12.19 |
| 1.370 | 37.65 | 9.19 | 1.475 | 46.96 | 12.35 |
| 1.375 | 38.105 | 9.34 | 1.480 | 47.39 | 12.50 |
| 1.380 | 38.56 | 9.48 | 1.485 | 47.82 | 12.66 |
| 1.385 | 39.01 | 9.63 | 1.490 | 48.25 | 12.82 |
| 1.390 | 39.46 | 9.78 | 1.495 | 48.675 | 12.97 |
| 1.395 | 39.92 | 9.93 | 1.500 | 49.10 | 13.13 |
| 1.400 | 40.37 | 10.07 | 1.505 | 49.53 | 13.29 |
| 1.405 | 40.82 | 10.22 | 1.510 | 49.95 | 13.45 |
| 1.410 | 41.26 | 10.37 | 1.515 | 50.38 | 13.60 |
| 1.415 | 41.71 | 10.52 | 1.520 | 50.80 | 13.76 |
| 1.420 | 42.155 | 10.67 | 1.525 | 51.22 | 13.92 |
| | | | 1.530 | 51.64 | 14.08 |

Ammonia NH₃, M = 17.03 g/mol

| 141131 111 | | 9/11101 | | | |
|------------|-------------------------|---------|----------|---------------------|-------|
| Density | NH ₃ content | | Density | NH ₃ cor | ntent |
| d 20°/4° | weight% | mol/l | d 20°/4° | weight% | mol/l |
| | | | | | |
| 0.998 | 0.0465 | 0.0273 | 0.932 | 17.24 | 9.44 |
| 0.996 | 0.512 | 0.299 | 0.930 | 17.85 | 9.75 |
| 0.994 | 0.977 | 0.570 | 0.928 | 18.45 | 10.06 |
| 0.992 | 1.43 | 0.834 | 0.926 | 19.06 | 10.37 |
| 0.990 | 1.89 | 1.10 | 0.924 | 19.67 | 10.67 |
| 0.988 | 2.35 | 1.365 | 0.922 | 20.27 | 10.97 |
| 0.986 | 2.82 | 1.635 | 0.920 | 20.88 | 11.28 |
| 0.984 | 3.30 | 1.91 | 0.918 | 21.50 | 11.59 |
| 0.982 | 3.78 | 2.18 | 0.916 | 22.125 | 11.90 |
| 0.980 | 4.27 | 2.46 | 0.914 | 22.75 | 12.21 |
| 0.978 | 4.76 | 2.73 | 0.912 | 23.39 | 12.52 |
| 0.976 | 5.25 | 3.01 | 0.910 | 24.03 | 12.84 |
| 0.974 | 5.75 | 3.29 | 0.908 | 24.68 | 13.16 |
| 0.972 | 6.25 | 3.57 | 0.906 | 25.33 | 13.48 |
| 0.970 | 6.75 | 3.84 | 0.904 | 26.00 | 13.80 |
| 0.968 | 7.26 | 4.12 | 0.902 | 26.67 | 14.12 |
| 0.966 | 7.77 | 4.41 | 0.900 | 27.33 | 14.44 |
| 0.964 | 8.29 | 4.69 | 0.898 | 28.00 | 14.76 |
| 0.962 | 8.82 | 4.98 | 0.896 | 28.67 | 15.08 |
| 0.960 | 9.34 | 5.27 | 0.894 | 29.33 | 15.40 |
| 0.958 | 9.87 | 5.55 | 0.892 | 30.00 | 15.71 |
| 0.956 | 10.405 | 5.84 | 0.890 | 30.685 | 16.04 |
| 0.954 | 10.95 | 6.13 | 0.888 | 31.37 | 16.36 |
| 0.952 | 11.49 | 6.42 | 0.886 | 32.09 | 16.69 |
| 0.950 | 12.03 | 6.71 | 0.884 | 32.84 | 17.05 |
| 0.948 | 12.58 | 7.00 | 0.882 | 33.595 | 17.40 |
| 0.946 | 13.14 | 7.29 | 0.880 | 34.35 | 17.75 |
| 0.944 | 13.71 | 7.60 | | | |
| 0.942 | 14.29 | 7.91 | | | |
| 0.940 | 14.88 | 8.21 | | | |
| 0.938 | 15.47 | 8.52 | | | |
| 0.936 | 16.06 | 8.83 | | | |
| 0.934 | 16.65 | 9.13 | | | |
| | | | | | |



COMMERCIALLY AVAILABLE CONCENTRATIONS OF SOME ACIDS AND ALKALIS

| Name | Weight % | Density d ²⁰ ° | Density (mol/l*) |
|-------------------------------------|----------|------------------------------|---------------------|
| Acetic acid | 96 | 1.06 | 17 |
| Acetic acid (glacial acetic acid) | 99 - 100 | 1.06 | 18 |
| Acetic acid, dilute | 30 | 1.04 | 5 |
| Ammonia solution | 35 | 0.88 | 18 |
| Ammonia solution | 30 | 0.88 | 15.5 |
| Ammonia solution | 25 | 0.91 | 13.5 |
| Formic acid | 98 - 100 | 1.22 | 26 |
| Hydriodic acid | 57 | 1.7 | 7.5 |
| Hydrobromic acid | 40 | 1.38 | 7 |
| Hydrochloric acid | 25 | 1.12 | 8 |
| Hydrochloric acid, concentr. (1.16) | 32 | 1.16 | 10 |
| Hydrochloric acid, concentr. (1.18) | 36 | 1.18 | 12 |
| Hydrochloric acid, fuming | 37 | 1.19 | 12.5 |
| Hydrofluoric acid | 48 | 1.16 | 28 |
| Hydrofluoric acid | 40 | 1.13 | 23 |
| Nitric acid, concentr. | 65 | 1.40 | 14 |
| Nitric acid, fuming | 100 | 1.52 | 21 |
| Perchloric acid | 70 | 1.67 | 12 |
| Perchloric acid | 60 | 1.53 | 9 |
| Phosphoric acid, concentr. (1.71) | 85 | 1.71 | 15 |
| Phosphoric acid, concentr. (1.75) | 89 | 1.75 | 16 |
| Potassium hydroxide solution | 47 | 1.5 | 12.5 |
| Potassium hydroxide solution | 30 | 1.3 | 7 |
| Sodium hydroxide solution | 33 | 1.36 | 11 |
| Sulfuric acid, concentr. | 95 - 97 | 1.84 | 18 |
| Sulfuric acid, dilute | 25 | 1.18 | 3 |

* rounded off

Baumé degrees (°Bé) and density °Be = 145 - $\frac{145}{\text{density}}$


Example:

Sodium hydroxide solution 40 % with a density of 1.430 g/cm³:

 $145 - \frac{145}{143} = 43.60$ °Be



pH-INDICATORS



The pH ranges and color shades shown are approximations

ACID-BASE INDICATORS (SORTED ACCORDING TO INCREASING PH TRANSITION RANGE)

| (SORT | ED ACCORDING I |) INCRE | ASING PH | TRANSITION RANGE |
|-------------|---|--------------------|------------------------------|---|
| Cat. No. | Indicator | Color cha interval | nge | Indicator solution (preparation) |
| 1.01398 | Malachite green oxalate (C.I. No. 42000) | 0.0 – yellow | 2.0 green-blue | 0.1 g in 100 ml of water |
| 1.01310 | Brilliant green (C.I. No. 42040) | 0.0 – yellow | 2.6 green | 0.1 g in 100 ml of water |
| 1.15935 | Eosin yellowish (C.I. No. 45380) | 0.0 – yellow | 3.0 green fluorescence | 0.1 g in 100 ml of water |
| 1.15936 | Erythrosin B (C.I. No. 45430) | 0.0 – orange | 3.6 red | 0.1 g in 100 ml of water |
| 1.15944 | Methyl green (C.I. No. 42590) | 0.1 – yellow | 2.3 blue | 0.1 g in 100 ml of water |
| 1.15945 | Methyl violet (C.I. No. 42535) | 0.1 – yellow | 2.7 violet | 0,1 g in 100 ml of ethanol (20 %) |
| 1.00623 | Picric acid (C.I. No. 10305) | 0.2 – colorless | 1.0 yellow | 0,1 g in 100 ml of ethanol (20 %) |
| 1.05225 | Cresol red | 0.2 – red | 1.8 yellow | Dissolve 0.1 g in 100 ml of ethanol (20 %) or 0.04 g in 1.05 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.01408 | Crystal violet (C.I. No. 42555) | 0.8 – yellow | 2.6 blue-violet | 0,1 g in 100 ml of ethanol (70%) |
| 1.05228 | m-Cresol purple | 1.2 – red | 2.8 yellow | Dissolve 0.04 g in 100 ml of ethanol (20 %) or 0.04 g in 1.05 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.08176 | Thymol blue | 1.2 - red | 2.8 yellow | Dissolve 0.04 g in 100 ml of ethanol (20 %) or 0.04 g in 0.86 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.08682 | p-Xylenol blue | 1.2 - red | 2.8 yellow | Dissolve 0.1 g in 100 ml of ethanol (50%) or 0.04 g in 0.98 ml of sodium hydroxide solution 0.1 mol/l and make up to 0.1 ml with water. |
| | | | | |

ACID-BASE INDICATORS (SORTED ACCORDING TO INCREASING PH TRANSITION RANGE)

| Cat. | Indicator | Color cha | nge | Indicator solution |
|---------|--|--------------------|-----------------------------|---|
| No. | mulcator | interval | | (preparation) |
| 1.11473 | 2.2', 2", 4.4'- Pentamethoxytriphenyl carbinol | 1.2 - red | 3.2 colorless | 0.1 g in 100 ml of ethanol (96%) |
| 1.15934 | Eosin bluish (C.I. No. 45400) | 1.4 – colorless | 2.4 pink fluorescence | 0.1 g in 100 ml of water |
| 1.02282 | Quinaldine red | 1.4 – colorless | 3.2 pink | 0.1 g in 100 ml of ethanol (60%) |
| 1.03464 | 2,4-Dinitrophenol | 2.8 – colorless | 4.7 yellow | 0.1 g in 100 ml of ethanol (70%) |
| 1.03055 | 4-(Dimethylamino) azobenzene (C.I. No. 11020) | 2.9 - red | 4.0 yellow-orange | 0.1 - 0.5 g in 100 ml of ethanol (90%) |
| 1.03022 | Bromochlorophenol blue | 3.0 – yellow | 4.6 blue-violet | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 0.69 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.08122 | Bromophenol blue | 3.0 – yellow | 4.6 blue-violet | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 0.6 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.01340 | Congo red (C.I. No. 22120) | 3.0 – blue | 5.2 yellow-orange | 0.2 g in 100 ml of water |
| 1.01322 | Methyl orange (C.I. No. 13025) | 3.1 - blue | 4.4 yellow-orange | 0.04 g in 100 ml of ethanol (20%) or 0.4 g in 100 ml of water |
| 1.01323 | Methyl orange solution | 3.1 – red | 4.4 yellow-orange | 0.04 g in 100 ml of ethanol (20%) or 0.4 g in 100 ml of water |
| 1.01359 | Mixed indicator 4.5 acc. to Mortimer | 4.3 – red | 5.2 blue | |
| 1.08121 | Bromocresol green | 3.8 - yellow | 5.4 blue | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 0.58 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.03465 | 2.5-Dinitrophenol | 4.0 – colorless | 5.8 yellow | 0.05 -0.1 g in 100 ml ethanol (70%) |

ACID-BASE INDICATORS (SORTED ACCORDING TO INCREASING PH TRANSITION RANGE)

| (SORTI | ED ACCORDING TO | DINCRE | ASING PH | TRANSITION RANGE) |
|-------------|--|----------------------------|----------------------|---|
| Cat. No. | Indicator | Color cha interval | nge | Indicator solution (preparation) |
| 1.06279 | Alizarinsulfonic acid sodium salt (C.I. No. 58005) | 4.3 - yellow | 6.3 violet | 0.1 g in 100 ml of ethanol (50%) or 0.1 g in 100 ml of water |
| 1.06076 | Methyl red (C.I. No. 13020) | 4.4 – red | 6.2 yellow-orange | 0.1 g in 100 ml of ethanol (96%) |
| 1.06078 | Methylred sodium salt (C.I. No. 13020) | 4.4 – red | 6.2 yellow-orange | 0.1 g in 100 ml of water |
| 1.06130 | Mixed indicator 5 | 4.4 – red-violet | 5.8 green | |
| 1.04795 | Cacotheline | + 0.525 yellow | 18.1 red-violet | saturated (irreversibly in water colorless in too strong reduction) |
| 1.03024 | Chlorophenol red | 4.8 – yellow | 6.4 purple | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 0.94 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.05312 | Litmus (C.I. No. 1242) | 5.0 – red | 8.0 blue | 4 g in 100 ml of water |
| 1.03025 | Bromocresol purple | 5.2 – yellow | 6.8 purple | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.4 g in 0.74 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.03023 | Bromophenol red | 5.2 – orange- yellow | 6.8 purple | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 0.94 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.06798 | 4-Nitrophenol | 5.4 – colorless | 7.5 yellow | 0.2 g in 100 ml of ethanol (96%) or 0.08 g in 100 ml of water |
| 1.03033 | Bromoxylenol blue | 5.7 – yellow | 7.5 blue | 0.1 g in 100 ml of ethanol (96%) |
| 1.01016 | Alizarin (C.I. No. 58000) | 5.8 – yellow | 7.2 red | 0.5 g in 100 ml Ethanol (96%) |
| 1.03026 | Bromothymol blue | 6.0 - | 7.6 | 0.1 g in 100 ml Ethanol (20%) yellow blue 0.04 g in 0.64 ml Natronlauge 0.1 mol/l lösen und mit Wasser auf 100 ml auffüllen |

ACID-BASE INDICATORS

| Cat. No. | Indicator | Color cha interval | nge | Indicator solution (preparation) |
|-------------|--|--------------------|----------------------|--|
| 1.07241 | Phenol red | 6.4 – yellow | 8.2 red-violet | Dissolve 0.1 g in 100 ml of ethanol (20%) or 0.04 g in 1.13 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.06794 | 3-Nitrophenol | 6.6 – colorless | 8.6 yellow-orange | 0.3 g in 100 ml of ethanol (96%) or 0.08 g 100 ml of water |
| 1.01369 | Neutral red (C.I. No. 50040) | 6.8 – blue-red | 8.0 orange-yellow | 0.3 g in 100 ml of ethanol (70%) |
| 1.13857 | 4,5,6,7-Tetrabromo- phenolphthalein | 7.0 – colorless | 8.0 purple | 0.1 g in 100 ml of ethanol (96%) |
| 1.05225 | Cresol red | 7.0 – orange | 8.8 purple | Dissolve 0.1 g in 100 ml of ethanol (50%) or 0.04 g in 1.05 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.06246 | 1-Naphtolphthalein | 7.1 – brownish | 8.3 blue-green | 0.1 g in 100 ml of ethanol (96%) |
| 1.05228 | Cresol purple | 7.4 – yellow | 9.0 purple | Dissolve 0.04 g in 100 ml of ethanol (20%) or 0.04 g in 1.05 ml of sodium hydroxide solution 0.1 mol/l and make p to 100 ml with water. |
| 1.08176 | Thymol blue | 8.0 – yellow | 9.6 blue | Dissolve 0.04 g in 100 ml of ethanol (20%) or 0.04 g in 0.86 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.08682 | p-Xylenol blue | 8.0 – yellow | 9.6 blue | Dissolve 0.1 g in 100 ml of ethanol (50%) or 0.04 g in 0.98 ml of sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water. |
| 1.07233 | Phenolphthalein | 8.2 – colorless | 9.8 red-violet | 0.1 g in 100 ml of ethanol (96%) |

ACID-BASE INDICATORS (SORTED ACCORDING TO INCREASING PH TRANSITION RANGE)

| (JUNII) | LD ACCONDING I | JINCIL | ASING III | TRANSITION RANGE) |
|-------------|---|---------------------------|-----------------------------|---|
| Cat. No. | Indicator | Color cha interval | nge | Indicator solution (preparation) |
| 1.07227 | Phenolphthalein solution (1% in ethanol) | 8.2 – colorless | 9.8 red-violet | |
| 1.07238 | Phenolphthalein solution (0.375% in methanol) | 8.2 – colorless | 9.8 red-violet | |
| 1.08175 | Thymolphthalein | 9.3 – colorless | 10.5 blue | 0.1 g in 100 ml of ethanol (50%) |
| 1.01331 | Alkali blue (C.I. No. 42765) | 9.4 – violet | 14.0 pink | 0.1 g in 100 ml of ethanol (96%) |
| 1.06776 | Alizarin yellow GG (C.I. No. 14025) | 10.0 – light yellow | 12.1 brownish- yellow | 0.1 g in 100 ml of water |
| 1.01016 | Alizarin (C.I. No. 58000) | 10.1 – red | 12.1 purple | 0.5 g in 100 ml of ethanol (96%)) |
| 1.01307 | Titan yellow (C.I. No. 19540) | 12.0 – yellow | 13.0 red | 0.1 g in 100 ml of ethanol (20 %) |
| 1.04724 | Indigo carmine (C.I. No. 73015) | 11.5 – blue | 13.0 yellow | 0.25 g in 100 ml of ethanol (50%) or 1 g in 100 ml of water |
| 1.06810 | Epsilon blue | 11.6 – orange | 13.0 violet | 0.1 g in 100 ml of water |

REDOX INDICATORS (IN ALPHABETICAL ORDER)

| (| | | | | | | |
|-------------|---|--------------------------------|--|------------|-------------------------|----------------|---|
| Cat. No. | Indicator | E, [v] | otential E _m [v] (pH=7) 30°C | rH | Color cha oxidized f | | Indicator solution (preparation) |
| 1.01167 | Amido black 10 B (C.I. No. 20470) | + 0.57 | + 0.84 | | yellowish- brown | blue | 0.2 g in 100 ml of water |
| 1.03098 | 2.2'-Bipyridin (iron(II) complex) | + 1.03 | | | pale blue | red | $0.695~g$ of Fe SO_4 \cdot 7H $_2O$ + 1.171 g of 2.2'-bipyridin in 100 ml of water |
| 1.01368 | Brillant cresyl blue (C.I. No. 51010) | + 0.58 | + 0.05 | | blue | colorless | 0.5 g in 100 ml of water or ethanol (96%) |
| 1.03028 | 2.6-Dichlorophenolin- dophenol sodium salt (dihydrate) | + 0.67 | + 0.23 | 22 | blue | colorless | 0.02 g in 100 ml of water |
| 1.03122 | 3.3'-Dimethylnaphthidine (4.4'-Diamino-3.3'- dimethyl-1.1'- binaphthaline) | + 0.78 | | | purple- red | colorless | 1.0 g in 100 ml of glacial acetic acid |
| 1.03067 | N,N-Dimethyl- 1.4-phenylenediam- monium dichloride | + 0.751 | | | dark- blue | colorless | 0.2 g in 100 ml of water |
| 1.03086 | Diphenylamine | + 0.76 | | | blue- violet | colorless | 1.0 g in 100 ml of sulfuric acid (96%) |
| 1.00255 | Diphenylamine-4- sulfonic acid barium salt | + 0.84 i Schwefe 1 mol/l | | 28.5 | red- violet | colorless | 0.2 g in 100 ml of water |
| 8.20530 | Diphenylbenzidine | + 0.76 i Schwefe 1 mol/l | | 26 ±3.0 | violet | colorless | 1.0 g in 100 ml of sulfuric acid (96%) |
| 1.09161 | Ferroin indicator soluti mol/l for waste water | . , . | | | | | |
| 1.09193 | Ferroin solution 1/40 mol/l | | | | | | |
| | Ferroin (1.10-Phenanthroline iron(II) complex) | + 1.06 | | 40 | blue | orange- red | |
| 1.04724 | Indigo carmine (Indigo disulfonate disodium salt) (C.I. No. 73015) | + 0.29 | - 0.11 | 10 | blue | yellowish | 0.05 g in 100 ml of water |

REDOX INDICATORS (IN ALPHABETICAL ORDER)

| Cat. No. | Indicator | E ₀ [v] | otential E _m [v] (pH=7) 30°C | rH | Color cha oxidized f | | Indicator solution (preparation) |
|-------------|---|--------------------|--|------|---|-----------|---|
| 1.15943 | Methylene blue (C.I. No. 52015) | + 0.53 (30 °C) | + 0.01 | 14.5 | blue | colorless | 0.1 – 0.5 g in 100 ml of water |
| 1.01369 | Neutral red (C.I. No. 50040) | + 0.24 | - 0.29 | 3 | violet- red | colorless | 0.05 g in 100 ml of ethanol (96%) |
| 1.15946 | Nile blue (sulfate) (C.I. No. 51180) | + 0.41 | - 0.12 | | blue- red | colorless | 0.1 g in 100 ml of water |
| 1.07225 | 1.10-Phenanthroline (monohydrate) | + 1.14 | + 1.06 | | pale blue | ted | 0.695 g FeSO_4 \cdot 7H ₂ O + 1.487 g 1.10-Phenanthroli in 100 ml of wate |
| 8.20979 | N-Phenylanthranilic acid | + 0.89 | | | purple- red | colorless | Dissolve 0.1 g in 5 ml sodium hydroxide solution 0.1 mol/l and make up to 100 ml with water or dissolve 0.107 g in 20 ml of sodium carbonate solution (5%) and make up to 100 ml with water |
| 1.15498 | Safranin (C.I. No. 50240) | + 0.24 | - 0.29 | 4 | blue- violet (acidic) brown (alkaline) | colorless | 0.05 g in 100 ml of water |
| 1.13759 | 2.2':6'2"-Terpyridine (iron(II) complex) | + 1.25 | | | pale blue | rot | 0.232 g FeSO_4 $\cdot 7\text{H}_2\text{O} + 0.389 \text{ g}$ $2.2' \cdot 6'2$ -terpyridine in 100 ml of water |
| 1.15929 | Thionine (C.I. No. 52000) | + 0.56 | + 0.06 | 16 | violet | colorless | 0.05 g in 100 ml of ethanol (60%) |
| 1.08507 | Variamine blue salt B (C.I. No. 37255) | + 0.712 | + 0.31 | | blue- violet (acidic) yellow (alkaline) | colorless | 1.0 g in 100 ml of water or grind with sodium chloride or sodium sulfate anhydrous trituration |

SUBSTANCES FOR THE PREPARATION OF BUFFER STOCK SOLUTIONS

| Cat. No. | Name |
|----------|--|
| | |
| 1.00165 | Boric acid cryst. GR for analysis |
| 1.00244 | Citric acid monohydrate GR |
| 1.06318 | 5.5-Diethyl barbituric acid sodium salt buffer substance |
| 1.04201 | Glycine GR for analysis |
| 1.04716 | Imidazole buffer substance |
| 1.04873 | Potassium dihydrogen phosphate GR for analysis |
| 1.04874 | Potassium hydrogen phthalate GR |
| 1.06268 | Sodium acetate anhydrous GR for analysis |
| 1.06267 | Sodium acetate trihydrate GR for analysis |
| 1.06392 | Sodium carbonate anhydrous GR |
| 1.06404 | Sodium chloride GR for analysis |
| 1.06329 | Sodium hydrogen carbonate GR for analysis |
| 1.06580 | di-Sodium hydrogen phosphate dihydrate GR for analysis |
| 1.08418 | Titriplex®III GR for analysis |
| 1.08379 | Triethanolamine GR for analysis |
| 1.08382 | Tris(hydroxymethyl) aminomethane GR for analysis |



BUFFER SOLUTIONS

Prepare stock and buffer solutions with distilled, boiled, $\rm CO_2$ -free water. The volumes x (in ml) of the stock solutions required to obtain the different pH values, are listed on the next page.


| | er - Stock solutions and their con | tent of buffer substance | Composition |
|--------------|---|--|---|
| solut No. | tion A | В | of buffer solution |
| 1 | Glycine 0.1 mol/l + NaCl 0.1 mol/l [Glycine: 7.507 g/l + NaCl: 5.844 g/l] | HCI 0.1 mol/I | x parts A + (100-x) parts B |
| 2 | di-Sodium citrate 0.1 mol/l [Citric acid monohydrate: 21.014 g/l + 200 ml NaOH 1 mol/l] | HCI 0.1 mol/I | x parts A + (100-x) parts B |
| 3 | Potassium hydrogen phthalate I 0.1 mol/[C ₈ H ₅ KO ₄ : 20.42 g/l | HCI 0.1 mol/I | 50 ml A + x ml B, make up to 100 ml* |
| 4 | As No. 3 | NaOH 0.1 mol/l make up to 100 ml* | 50 ml A + x ml B, |
| 5 | As No. 2 | NaOH 0.1 mol/l (100-x) parts B | x parts A + |
| 6 | Potassium dihydrogen phosphate 1/15 mol/l [KH ₂ PO ₄ : 9.073 g/l] [Na ₂ HPO ₄ · 2 H ₂ O: 11.87 g/l] | di-Sodium hydrogen phosphate 1/15 mol/l | x parts A + (100-x) parts B |
| 7 | 5.5-Diethylbarbituric acid sodium salt 0.1 mol/I [Barbital-Na: 20.62 g/I] | HCI 0.1 mol/I | x parts A + (100-x) parts B |
| 8 | Borax solution 0.05 mol/l [H ₃ BO ₃ : 12.37 g/l + 100 ml NaOH 1 mol/l] | HCI 0.1 mol/I | x parts A + (100-x) parts B |
| 9 | As No. 1 | NaOH 0.1 mol/l (100-x) parts B | x parts A + |
| 10 | Citric acid 0.1 mol/l [Citric acid monohydrate: 21.014 g/l] [Na ₂ HPO ₄ \cdot 2 H ₂ O: 35.60 g/l] | di-Sodium hydrogen phosphate 0.2 mol/l | x parts A + (100-x) parts B |
| 11 | Sodium acetate 0.1 mol/l [$C_2H_3O_2Na: 8.204 \text{ g/l} \text{ or } C_2H_3O_2Na: 3 H_2O: 13.61 \text{ g/l}]$ | Acetic acid 0.1 mol/l | x parts A + (100-x) parts B |
| 12 | Imidazole 0.2 mol/l [C ₃ H ₄ N ₂ : 13.62 g/l] | HCI 0.1 mol/I | 25 ml A + x ml B, make up to 100 ml* |
| 13 | $ \begin{array}{l} \text{Triethanolamine 0.5 mol/l +} \\ \text{Titriplex}^{\$} \text{ III} \\ \left[C_{e} H_{15} N Q_{3} \cdot 74.60 \text{ g/l +} \\ \text{Titriplex}^{\$} \text{ III: 20 g/l]} \end{array} $ | HCI 0.05 mol/I | 10 ml A + x ml B, make up to 100 ml* |
| 14 | Tris(hydroxymethyl)aminomethane 0.2 mol/l [TRIS: 24.23 g/l] | HCI 0.1 mol/I | 25 ml A + x ml B, make up to 100 ml* |
| 15 | Sodium carbonate 0.1 mol/l (10.60 g/l) $[Na_2CO_3: 10.60 \text{ g/l}]$ $[NaHCO_3: 8.401 \text{ g/l}]$ | Sodium hydrogen carbonate 0.1 mol/l | x parts A + (100-x) parts B |

BUFFER SOLUTIONS



Merck KGaA · D-64271 Darmstadt Germany


* Lösung auffüllen



pKa VALUES OF SELECTED BIOLOGICAL BUFFERS

| Buffer | pK _a (4°C) | pK _a (20 °C) | pK _a (25 °C) | pK _a (37 °C) | ∆pK _a /°C |
|----------------------------|-----------------------|-------------------------|-------------------------|-------------------------|----------------------|
| ACES | 7.22 | 6.90 | 6.80 | 6.56 | -0.020 |
| ADA | 6.80 | 6.62 | 6.56 | 6.43 | -0.011 |
| BES | 7.41 | 7.15 | 7.07 | 6.88 | -0.016 |
| BICIN | 8.64 | 8.35 | 8.26 | 8.04 | -0.018 |
| BIS-TRIS | 6.88 | 6.56 | 6.46 | 6.22 | -0.020 |
| CHES | 9.73 | 9.55 | 9.50 | 9.36 | -0.011 |
| Citrat pK _{a2} | 4.79 | 4.77 | 4.76 | 4.74 | -0.0016 |
| Glycin pK _{a2} | 10.32 | 9.91 | 9.78 | 9.47 | -0.026 |
| Gly-Gly | 8.85 | 8.40 | 8.26 | 7.92 | -0.028 |
| HEPES | 7.77 | 7.55 | 7.48 | 7.32 | -0.014 |
| HEPPS | 8.18 | 8.00 | 7.95 | 7.8 | -0.011 |
| Imidazole | 7.37 | 7.05 | 6.95 | 6.71 | -0.020 |
| MES | 6.33 | 6.15 | 6.10 | 5.97 | -0.011 |
| MOPS | 7.41 | 7.20 | 7.14 | 6.98 | -0.013 |
| PIPES | 6.94 | 6.80 | 6.76 | 6.66 | -0.0085 |
| Phosphate pK _{a2} | 7.26 | 7.21 | 7.20 | 7.17 | -0.0028 |
| TAPS | 8.02 | 8.31 | 8.40 | 8.62 | +0.018 |
| TES | 7.82 | 7.50 | 7.40 | 7.16 | -0.020 |
| TRICIN | 8.49 | 8.15 | 8.05 | 7.79 | -0.021 |
| TRIS | 8.75 | 8.30 | 8.08 | 7.82 | -0.028 |

BUFFER RANGES



CHROMATOGRAPHY

For every application the matching stationary phase - only MERCK offers this:

- Coarse-granular for industrial chromatography such as e.g. aluminum oxides, cellulose and, of course, all types of silica gels
- Finely classified for the demanding requirements of preparative chromatography, e.g. our LiChroprep® range
- Finely classified for all kinds of analytical HPLC such as our LiChrosorb®, LiChrospher®, Superspher® etc.
- Organic polymer resins specially designed for the requirements of biochromatography.

From the coarsest to the most finely granulated materials the selectivities are identical. MERCK provides you with a problem-free scale-up. For every separation problem we have the most appropriate prepacked column:

- ☐ Prepbar® stainless-steel prepacked columns for use in industrial production
- ☐ Lobar® glass prepacked columns to solve preparative separation problems in the laboratory
- ☐ Hibar® or LiChroCART® stainless-steel columns for analytical HPLC in all dimensions including online sample preparation
- ☐ CGC glass cartridges for HPLC problems where chemical inertness is the key.
- ☐ Superformance® glass cartridges and glass prepacked columns represent the inert, biocompatible and pressure-resistant system for analytical and preparative biochromatography

To satisfy all demands: for all our columns we provide a complete range of auxiliaries and the matching instrument components such as our Prepbar® system for industrial separation problems or our LiChroGraph® system for HPLC.

Should you have a chromatographical separation problem, ask MERCK first - we surely have the solution.

By the way, if you require special solvents for HPLC ask about our LiChrosolv® range – you will have difficulty finding better!


PREPARATIVE AND ANALYTICAL THIN-LAYER CHROMATOGRAPHY

First of all, ask MERCK


For example: The format 2.5 cm x 7.5 cm for orientating separations (reaction process control).



For example: Chiral plates for exact separations of enantiomers.



For example: Coded precoated plates for GMP/GLP-conform applications.



For example: 0.5...1...2 mm layers for efficient preparative scale separations.



Detailed information on our range of ready-to-use products for thin-layer chromatography can be found in our "Chromatography Catalogue" which we shall be pleased to send you on request.



ORGANIC SOLVENTS PROPERTIES AND DRYING

| Solvent | Boiling pt. [°C] | D ^{20°} | n D | Flash pt [°C] | MA(| C (2) mg/m ³ | Drying agent (1) |
|---|---------------------|------------------|-------|------------------|-----------|----------------------------|--|
| Acetone | 56 | 0.791 | 1.359 | - 18 | 500 | 1200 | K₂CO₃ Mole- cular sieve 0.3 nm |
| Acetic acid | 118 | 1.049 | 1.372 | + 40 | 10 | 25 | P ₂ O ₅ ; CuSO ₄ |
| Acetic anhydride | 136 | 1.082 | 1.390 | + 49 | 5 | 20 | CaCl ₂ |
| Acetonitrile | 82 | 0.782 | 1.344 | + 6 | 40 | 69 | CaCl ₂ ; P ₂ O ₅ ; K ₂ CO ₃ Molecular sieve 0.3 nr |
| Aniline | 184 | 1.022 | 1.586 | + 76 | | 8 | KOH; BaO |
| Anisole | 154 | 0.995 | 1.518 | + 51 | | | CaCl ₂ ; distillation; Na |
| Benzene | 80 | 0.879 | 1.501 | - 10 | H.A | | distillation CaCl ₂ ; Na; Pb/Na Molecular sieve 0.4 n |
| 1-Butanol | 117 | 0.810 | 1.399 | + 29 | 100 | 310 | K ₂ CO ₃ ; distillation |
| 2-Butanol | 100 | 0.808 | 1.398 | + 24 | 100 | 310 | K ₂ CO ₃ ; distillation |
| tert-Butanol | 82 | 0.786 | 1.384 | + 11 | 100 | 310 | CaO; freezing |
| n-Butyl acetate | 127 | 0.882 | 1.394 | + 33 | 100 | 480 | MgSO ₄ |
| Carbon disulfide | 46 | 1.263 | 1.626 | - 30 | H 5 | 16 | CaCl ₂ ; P ₂ O ₅ |
| Carbon tetrachloride | 77 | 1.594 | 1.460 | non flammable | H 10 B | 65 | Distillation; CaCl ₂ ; P ₂ O ₅ ; Pb/Na; Molecular sieve 0.4 n |
| Chlorobenzene | 132 | 1.106 | 1.525 | + 29 | 10 | 47 | CaCl ₂ ; distillation; P ₂ O ₅ |
| Chloroform | 62 | 1.486 | 1.448 | non flammable | B.10 | 50 | CaCl ₂ ; P ₂ O ₅ ; Pb/Na Molecular sieve 0.4 n |
| Cyclohexane | 81 | 0.779 | 1.426 | - 17 | 200 | 700 | Na; Na/Pb; LiAIH4 Molecular sieve 0.4 n |
| Decahydronaphthalene (Dekalin) | 189/ 191 | 0.886 | 1.48 | < 54 | - | - | CaCl ₂ ; Na; Pb/Na |
| Dichloromethane (Methylene chloride) | 40 | 1.325 | 1.424 | non flammable | B.100 | 350 | CaCl ₂ ; Pb/Na Molecular sieve 0.4 n |
| Diethyl carbonate | 126 | 0.975 | 1.384 | + 25 | - | - | K ₂ CO ₃ ; Na ₂ SO ₄ |
| Diethylene glycoldibutyl ether | 255 | 0.885 | 1.423 | + 118 | - | - | CaCl ₂ ; Na |
| Diethylene glycoldiethyl ether | 188 | 0.906 | 1.412 | + 82.5 | - | - | CaCl ₂ ; Na |
| Diethylene glycol dimethyl ether | 155 165 | 0.945 | 1.407 | + 70 | - | - | CaCl ₂ ; Na |
| Diethyl ether | 34 | 0.714 | 1.353 | - 40 | 400 | 1200 | CaCl ₂ ; Na; Pb/Na; LiAIH ₄ Molecular sieve 0.4 n |
| Diisopropyl ether | 68 | 0.726 | 1.368 | - 23 | 500 | 2100 | CaCl ₂ ; Na Molecular sieve 0.4 n |
| Dimethyl formamide | 153 | 0.950 | 1.430 | + 62 | H. 10 | 30 | Distillation Molecular sieve 0.4 n |
| Dimethyl sulfoxide | 189 | 1.101 | 1.478 | + 95 | - | - | Distillation Molecular sieve 0.3 r |

ORGANIC SOLVENTS PROPERTIES AND DRYING

| FROFERIL | 3 A | ND L | ווזחי | NG | | | |
|-------------------------------------|---------------------|---------------------|--------|------------------|------------|----------------------------|--|
| Solvent | Boiling pt. [°C] | D ^{20°} 4° | n D | Flash pt [°C] | MA(ppm | C (2) mg/m ³ | Drying agent (1) |
| 1.4-Dioxane | 101 | 1.034 | 1.422 | + 11.8 | H.B20 | 73 | CaCl ₂ ; Na Molecular sieve 0.4 nm |
| Ethanol | 79 | 0.791 | 1.361 | + 12 | 1000 | 1900 | CaO; Mg; MgO. Molecular sieve 0.3 nm |
| Ethyl acetate | 77 | 0.901 | 1.372 | - 4 | 400 | 1500 | K ₂ CO ₃ ; P ₂ O ₅ ; Na ₂ SO ₄ . Molecular sieve 0.4 nm |
| Ethylene glycol | 197 | 1.109 | 1.432 | + 111 | - | - | Distillation; Na ₂ SO ₄ |
| Ethylene glycol monoethyl ether | 135 | 0.930 | 1.408 | + 41 | H 5 | 20 | Distillation |
| Ethylene glycol monomethyl | 125 | 0.965 | 1.402 | + 52 | H 5 | 15 | Distillation |
| Ethyl formate | 54 | 0.924 | 1.360 | - 20 | 100 | 300 | MgSO ₄ ; Na ₂ SO ₄ |
| Formamide | 211 | 1.134 | 1.447 | 155 | - | - | Na ₂ SO ₄ ; CaO |
| Glycerol | 290 | 1.260 | 1.475 | + 176 | | | Distillation |
| Hexafluoroacetone (sesqui-hydrate) | | 1.685 | | non flammable | | | |
| n-Hexane | 69 | 0.659 | 1.375 | - 23 | 50 | 180 | Na; Pb/Na; LiAIH ₄ . Molecular sieve 0.4 nn |
| Isobutanol | 108 | 0.803 | 1.396 | + 28 | 100 | 300 | K ₂ CO ₃ ; CaO; Mg; Ca |
| Isobutyl methyl ketone | 117 | 0.801 | 1.396 | + 15.5 | 20 | 83 | K ₂ CO ₃ |
| Methanol | 65 | 0.792 | 1.329 | + 11 | H 200 | 270 | Mg; CaO. Molecular sieve 0.3 nn |
| Methyl acetate | 57 | 0.933 | 1.362 | - 10 | 5 | 20 | K ₂ CO ₃ ; CaO |
| 1-Methyl-2-pyrrolidon | 202 | 1.0260 | 1.4684 | + 95 | 20 | 80 | Distillation; Na ₂ SO ₄ ; Molecular sieve 0.4 nn |
| Methyl ethyl ketone | 80 | 0.806 | 1.379 | - 4.4 | 200 | 600 | K ₂ CO ₃ |
| Nitrobenzene | 211 | 1.204 | 1.556 | + 92 | H 1 | 5 | CaCl ₂ ; P ₂ O ₅ ; Distillation |
| n-Pentane | 36 | 0.626 | 1.358 | - 49 | 1000 | 3000 | Na; Pb/Na |
| 1-Propanol | 97 | 0.804 | 1.385 | + 15 | - | - | CaO; Mg |
| 2-Propanol | 82 | 0.785 | 1.378 | + 12 | 200 | 500 | CaO; Mg; Molecular sieve 0.3 nm |
| Pyridine | 116 | 0.982 | 1.510 | + 20 | 5 | 15 | KOH; BaO; Molecular sieve 0.4 nm |
| Tetrahydrofuran | 66 | 0.887 | 1.405 | - 17.5 | 50 | 150 | Molecular sieve 0.4 nm |
| Tetrahydronaphthalene (Tetralin) | 208 | 0.973 | 1.541 | + 78 | - | - | CaCl ₂ ; Na |
| Toluene | 111 | 0.867 | 1.496 | + 4 | 50 | 190 | Distillation; Ca; CaCl ₂ Molecular sieve 0.4 nn |
| Trichloroethylene | 87 | 1.462 | 1.477 | non flammable | В. – | - | Distillation; Na ₂ SO ₄ ; K ₂ CO ₃ |
| Xylene (isomeric mixture) | 137/ 140 | ~ 0.86 | ~ 1.50 | + 25 | 100 | 440 | Distillation; Na; CaCl ₂ Molecular sieve 0.4 nm |

ORGANIC SOLVENTS PROPERTIES AND DRYING

- (1) For details of drying methods please refer to the brochure "Drying in the laboratory and pilot plant"
- (2) MAC values
 - S = Danger of absorption through the skin
 - A = This substance is definitely known to be a carcinogenic; no MAC values can be quoted.
 - **B** = There are grounds to suppose that this substance has carcinogenic potential.

Substances for which no MAC value is given have not been classified by the German Senate Commission on hazardous materials, though this fact is not to be construed as meaning that the substances carry no risk.

ETHANOL-WATER MIXTURES

| | ETHANO | L-VVAID | L IVIIVI | UNE3 | | |
|---|--------------------|----------------------|----------------------|--------------------|----------------------|----------------------|
| ١ | Density | % | 0/0 | Density | % | 0/0 |
| | D 20° | by weight ethanol | by volume ethanol | D 20° | by weight ethanol | by volume ethanol |
| i | 1.00000 | 0 | 0 | 0.91546 | 50 | 57.8 |
| ı | 0.99813 | 1 | 1.3 | 0.91322 | 51 | 58.8 |
| ı | 0.99629 | 2 | 2.5 | 0.91097 | 52 | 59.8 |
| | 0.99451 | 3 | 3.8 | 0.90872 | 53 | 60.8 |
| | 0.99279 | 4 | 5.0 | 0.90645 | 54 | 61.8 |
| | 0.99113 | 5 | 6.2 | 0.90418 | 55 | 62.8 |
| | 0.98955 | 6 | 7.5 | 0.90191 | 56 | 63.8 |
| | 0.98802 | 7 | 8.7 | 0.89962 | 57 | 64.8 |
| | 0.98653 | 8 | 10.0 | 0.89733 | 58 | 65.8 |
| | 0.98505 | 9 | 11.2 | 0.89502 | 59 | 66.8 |
| | 0.98361 | 10 | 12.4 | 0.89271 | 60 | 67.7 |
| | 0.98221 | 11 | 13.6 | 0.89040 | 61 | 68.6 |
| | 0.98084 | 12 | 14.8 | 0.88807 | 62 | 69.9 |
| | 0.97948 | 13 | 16.1 | 0.88574 | 63 | 70.5 |
| | 0.97560 | 14 | 17.3 | 0.88339 | 64 | 71.5 |
| | 0.97687 | 15 | 18.5 | 0.88104 | 65 | 72.4 |
| | 0.97687 | 16 | 19.7 | 0.87869 | 66 | 73.3 |
| ı | 0.97431 | 17 | 20.9 | 0.87632 | 67 | 74.2 |
| | 0.97301 | 18 | 22.1 | 0.87396 | 68 | 75.1 |
| ı | 0.97169 | 19 | 23.3 | 0.87158 | 69 | 76.0 |
| | 0.97036 | 20 | 24.5 | 0.86920 | 70 | 76.9 |
| ı | 0.96901 | 21 | 25.7 | 0.86680 | 71 | 77.8 |
| ı | 0.96763 | 22 | 26.9 | 0.86440 | 72 | 78.6 |
| ı | 0.96624 | 23 | 28.1 | 0.86200 | 73 | 79.5 |
| | 0.96483 | 24 | 29.2 | 0.85958 | 74 | 80.4 |
| ı | 0.96339 0.96190 | 25 26 | 30.4 31.6 | 0.85716 0.85473 | 75 76 | 81.2 82.1 |
| | 0.96190 | 26 | 31.6 | 0.85473 | 76 | 83.0 |
| ì | 0.95880 | 28 | 33.9 | 0.84985 | 78 | 83.8 |
| | 0.95717 | 29 | 35.1 | 0.84740 | 79 | 84.6 |
| ì | 0.95551 | 30 | 36.2 | 0.84494 | 80 | 85.4 |
| ı | 0.95381 | 31 | 37.4 | 0.84245 | 81 | 86.2 |
| ì | 0.95207 | 32 | 38.5 | 0.83997 | 82 | 87.1 |
| ı | 0.95028 | 33 | 39.6 | 0.83747 | 83 | 87.9 |
| ì | 0.94847 | 34 | 40.7 | 0.83496 | 84 | 88.7 |
| ı | 0.94662 | 35 | 41.9 | 0.83242 | 85 | 89.5 |
| ı | 0.94432 | 36 | 43.0 | 0.82987 | 86 | 90.2 |
| | 0.94281 | 37 | 44.1 | 0.82729 | 87 | 91.0 |
| | 0.94086 | 38 | 45.2 | 0.82469 | 88 | 91.8 |
| | 0.93886 | 39 | 46.3 | 0.82207 | 89 | 92.5 |
| | 0.93648 | 40 | 47.4 | 0.81942 | 90 | 93.2 |
| | 0.93479 | 41 | 48.43 | 0.81674 | 91 | 94.0 |
| | 0.93272 | 42 | 49.51 | 0.81401 | 92 | 94.7 |
| | 0.93062 | 43 | 50.6 | 0.81127 | 93 | 95.4 |
| | 0.92849 | 44 | 51.6 | 0.80848 | 94 | 96.1 |
| | 0.92636 | 45 | 52.6 | 0.80567 | 95 | 96.7 |
| | 0.92421 | 46 | 53.7 | 0.80280 | 96 | 97.4 |
| ı | 0.92204 | 47 | 54.7 | 0.79988 | 97 | 98.1 |
| | 0.91986 | 48 | 55.8 | 0.79688 | 98 | 98.7 |
| ı | 0.91766 | 49 | 56.8 | 0.79383 | 99 | 99.3 |
| | | | | 0.79074 | 100 | 100.0 |

DRYING AGENTS

| Name | Formula | Water content of air in equilibrium, in mg/l (at 25°C) |
|------------------------|----------------------------------|---|
| Aluminium oxide | Al_2O_3 | 0.003 |
| Calcium chloride | CaCl ₂ | < 0.00001 |
| Calcium hydride | CaH ₂ | 0.14 |
| Calcium oxide | Ca0 | 0.003 |
| Calcium sulfate | CaSO ₄ | 0.004 - 0.07 |
| Copper sulfate | CuSO ₄ | 1.4 |
| Dessicant sachets | SiO ₂ | 0.003 |
| Magnesium oxide | Mg0 | 0.008 |
| Magnesium perchlorate | $Mg(CIO_4)_2$ | 0.0005 - 0.002 |
| Magnesium sulfate | MgSO ₄ | 1.0 |
| Molecular sieves | - | 0.0001 - 0.5 |
| Phosphorus pentoxide | $P_{2}O_{5}$ | 0.00002 |
| Potassium hydroxide | КОН | 0.002 |
| Sicacide® | H ₂ SO ₄ * | 0.003 - 0.3 |
| Sicapent® | P ₂ O ₅ * | < 0.000025 |
| Silica gel, blue gel | (SiO ₂) _x | 0.003 |
| Silica gel, orange gel | SiO ₂ | 0.003 |
| Sodium hydroxide | NaOH | 0.002 |
| Sodium sulfate | Na ₂ SO ₄ | 1.0 |
| Sulfuric acid | H ₂ SO ₄ | 0.005 - 0.3 |

^{*}on siliceous supporting material

VAPOUR PRESSURE OF WATER OVER H2SO4 (AT 20 °C)

| % H ₂ SO ₄ | 10 | 20 | 30 | 40 | 50 | 55 | 60 |
|----------------------------------|------|------|------|------|------|-------|-----|
| p [mbar] | 22,9 | 20,5 | 17,6 | 13,0 | 8,2 | 5,9 | 3,7 |
| % H ₂ SO ₄ | 65 | 70 | 75 | 80 | 85 | 90 | |
| p [mbar] | 2,1 | 1,1 | 0,4 | 0,1 | 0,04 | 0,007 | |

LICHROSOLV® SOLVENTS FOR CHROMATOGRAPHY

| Cat. No. | Elutropic series | Polarity index acc. to Snyder (1) | to | Molar mass | Refrac- tive index | point | Vapor pres- sure | Dynan viscos | | Surface tension against air or vapor | MAC value 1998 mg | Dielectric constant | Dipole moment acc. to Debye | ε°against Al ₂ O ₃ (1) acc. to Snyder | x [mr (silica | coeffion ² /s] (gel 6 ated p | DC- 0 |
|----------|------------------------------------|-----------------------------------|--|---------------|--------------------------|-------|------------------------|-----------------|----------------|--|----------------------------|------------------------|--------------------------------------|--|-----------------------------------|--|---------------|
| | | (1) | | [g/mol] | n ^{20°} D | [°C] | [mbar] (20°C) | [mPa (22°C) | · s] (40°C) | [mN/m] (20 °C) | [ml/m³] oder [ppm] | (20 or 25 °C) | | | migration dist [mm] 50 75 1 | | stance 100 |
| 1.04390 | n-Heptane | - | C ₇ H ₁₆ | 100.21 | 1.388 | 98.4 | 48 | 0.40 | 0.33 | 20.4 | 500 | 1.9 | 0 | 0.01 | 9.2 | 10.6 | 11.4 |
| 1.04391 | n-Hexane | 0.0 | C ₆ H ₁₄ | 86.18 | 1.375 | 68.9 | 160 | 0.31 | 0.26 | 18.4 | 50 | 1.9 | 0 | - | 12.5 | 13.9 | 14.6 |
| 1.02827 | Cyclohexane | 0.0 | C ₆ H ₁₂ | 84.16 | 1.427 | 80.7 | 104 | 0.94 | 0.71 | 25.5 | 200 | 2.0 | 0 | 0.04 | 5.4 | 6.3 | 6.7 |
| 1.04717 | Isooctane | 0.4 | C ₈ H ₁₈ | 114.23 | 1.392 | 99.2 | 51 | 0.51 | 0.50 | - | 500 | 1.9 | 0 | 0.01 | 7.9 | 8.3 | 8.7 |
| 1.08250 | 1, 1.2-Trichlor- trifluorethane | - | Cl ₂ FCCCIF ₂ | 187.38 | 1.356 | 47.7 | 368 | 0.70 | - | - | 500 | 2.4 | - | - | 5.2 | 5.8 | 5.9 |
| 1.02223 | Carbon- tetrachloride | 1.7 | CCI ₄ | 153.82 | 1.460 | 76.5 | 120 | 0.97 | 0.74 | 27.0 | 10 | 2.2 | 0 | 0.18 | 6.1 | 6.7 | 7.0 |
| 1.08327 | Toluene | 2.3 | C ₆ H ₅ CH ₃ | 92.14 | 1.496 | 110.6 | 29 | 0.58 | 0.47 | 28.5 | 50 | 2.4 | 0.36 | 0.29 | 8.3 | 9.8 | 11.0 |
| 1.02444 | Chloroform | 4.4 | CHCI ₃ | 119.38 | 1.946 | 61.7 | 210 | 0.56 | 0.47 | 27.1 | 10 | 4.8 | 1.01 | 0.40 | 9.0 | 10.5 | 11.6 |
| 1.13713 | Dichloro- ethane | 3.7 | CICH ₂ CH ₂ CI | 98.97 | 1.445 | 83.4 | 87 | 0.80 | 0.65 | 24.2 | A ⁽³⁾ | 10.6 | 1.75 | 0.44 | 7.6 | 8.4 | 8.9 |
| 1.06044 | Dichlor- methane | 3.4 | CH ₂ Cl ₂ | 84.93 | 1.424 | 40.0 | 453 | 0.43 | 0.36 | 26.5 | 100 | 9.1 | 1.60 | 0.42 | 10.1 | 11.8 | 13.2 |
| 1.01988 | 1-Butanol | 3.9 | $CH_3(CH_2)_3OH$ | 74.12 | 1.399 | 117.2 | 6.7 | 2.95 | 1.78 | 24.6 | 100 | 17.8 | 1.66 | - | - | - | - |
| 1.00030 | Acetonitrile | 6.2 | CH ₃ CN | 41.05 | 1.344 | 81.6 | 97 | 0.39 | - | 29.3 | 40 | 37.5 | 3.44 | 0.65 | 12.6 | 14.0 | 15.4 |
| 1.01040 | 2Propanol | 4.3 | CH ₃ CH(OH)CH ₃ | 60.10 | 1.378 | 82.4 | 43 | 2.27 | 1.35 | 21.7 | 200 | 18.3 | 1.66 | 0.82 | 2.1 | 2.3 | 2.5 |
| 1.00868 | Ethyl acetate | 4.3 | CH ₃ COOC ₂ H ₅ | 88.10 | 1.372 | 77.1 | 97 | 0.44 | 0.36 | 23.9 | 400 | 6.0 | 1.78 | 0.58 | 9.2 | 10.9 | 12.1 |
| 1.00020 | Acetone | 5.4 | CH ₃ COCH ₃ | 58.08 | 1.359 | 56.2 | 233 | 0.32 | 0.27 | 23.7 | 500 | 20.7 | 2.70 | 0.56 | 12.7 | 14.7 | 16.2 |
| 1.11727 | Ethanol | 5.2 | C ₂ H ₅ OH | 46.07 | 1.361 | 78.5 | 59 | 1.20 | 0.83 | 22.8 | 1000 | 24.3 | 1.70 | 0.88 | 3.4 | 3.9 | 4.2 |
| 1.03132 | 1.4-Dioxane | 4.8 | $C_4H_8O_2$ | 88.11 | 1.422 | 101.0 | 41 | 1.21 | 0.92 | 33.7 | 20 | 2.2 | 0.40 | 0.56 | 5.2 | 6.0 | 6.5 |
| 1.08101 | Tetra- hydrofuran | 4.2 | C ₄ H ₈ O | 72.11 | 1.405 | 66.0 | 200 | 0.47 | 0.38 | - | 50 | 7.4 | 1.63 | 0.57 | 10.9 | 11.9 | 12.6 |
| 1.06007 | Methanol | 6.6 | CH ₃ OH | 32.04 | 1.329 | 65.0 | 128 | 0.52 | 0.45 | 22.6 | 200 | 32.6 | 1.70 | 0.95 | 5.6 | 6.5 | 7.1 |
| 1.15333 | Water | 9.0 | H ₂ 0 | 18.01 | 1.333 | 100.0 | 23 | 0.95 | 0.65 | 72.8 | - | 80.2 | 1.85 | - | 5.1 | 5.7 | 5.8 |

(1) acc. to L. R. Snyder, Journal of Chromatography 92, 233, (1974)

(2) Detailed solvent tables acc. to H. Halpaap can be found in: Einführung in HPDC,

ed. R. E. Kaiser, IfC-Verlag Bad Dürkheim 1976, p. 232–233; HPTLC, ed. A. Zlatkis, R. E. Kaiser Elsevier and IfC 1977, p. 126–127.

(3) A = This substance is definitely known to be a carcinogenic; no MAC values can be quoted.

PHYSICAL METHODS FOR THE DETERMINATION OF ELEMENTS



FLAME PHOTOMETRY:

IMPORTANT EMISSION LINES IN THE FLAME SPECTRA OF SOME ELEMENTS

| Element | Wavelength I [nm] | | | | | | |
|---------|-------------------|-----------|-----------|--|--|--|--|
| Aq | 328.1 | 338.3 | | | | | |
| Ba | 553.6 | 744 (B) | 873 (B) | | | | |
| В | 452 (B) | 548 (B) | 345 (B) | | | | |
| Ca | 422.7 | 554 (B) | 622 (B) | | | | |
| Co | 346.6 (G) | 353.0 | 387.4 | | | | |
| Cr | 360.5 | 427.5 (G) | 425.5 | | | | |
| Cs | 455.5 | 852.1 | 894.3 | | | | |
| Cu | 324.8 | 327.4 | 520 (B) | | | | |
| Fe | 373.7 (G) | 386.0 (G) | 385.6 (G) | | | | |
| K | 404.7 (D) | 766.5 (D) | 344.6 (D) | | | | |
| Li | 670.8 | 460.3 | 323.3 | | | | |
| Mg | 285.2 | 371 (B) | 383 (B) | | | | |
| Mn | 403.3 (G) | 543.3 | 279.5 | | | | |
| Na | 330.3 (D) | 589.3 (D) | 818.3 (D) | | | | |
| Ni | 341.5 (G) | 352.5 (G) | 385.8 (G) | | | | |
| Pb | 368.4 | 405.8 | 261.4 (D) | | | | |
| Rb | 420.2 (D) | 780.0 | 794.8 | | | | |
| Sr | 460.7 | 821 (B) | 407.8 | | | | |
| Ti | 377.6 | 535.0 | 276.8 | | | | |

(B) = Band of the oxide

(D) = Dual line, the center point between the two lines is given

(G) = Group of lines in the region of the stated wavelength

WAVELENGTH AND WAVE NUMBER

Wavelength l [nm] and wave number n $[cm^{\scriptscriptstyle -1}]$

 $\overline{v} = \frac{1}{\lambda}$; 400 nm \triangleq 25000 cm⁻¹

PHOTOMETRY: TRANSMISSION RATE AND ABSORBANCE

A = -IgT(e.g. $A = 23.6\% = 0,236 \rightarrow T = 0.627$)

CALCULATION OF THE STANDARD DEVIATION

$$A \,=\, \sqrt{\sum \, F^2}$$

DIRECT-CURRENT POLAROGRAPHY HALF-WAVE POTENTIALS OF SOME IMPORTANT METALS


| Metal | Half-wave potential [V] | Support electrolyte /concentration |
|------------------|-------------------------|--|
| Cu ²⁺ | - 0.42 | NH ₄ CH ₃ COO 0.85 mol/l |
| Pb ²⁺ | - 0.48 | KSCN 0.0025 mol/l |
| Cd ²⁺ | - 0.64 | и |
| Ni ²⁺ | - 1.00 | и |
| Zn ²⁺ | - 1.06 | ш |
| Co ²⁺ | - 1.30 | и |
| Fe ²⁺ | - 1.41 | и |
| Mn ²⁺ | -1.55 | и |
| Cu ²⁺ | - 0.32 | Saturated CaCl ₂ solution |
| | | about 10-12 mol/l |
| Pb ²⁺ | - 0.52 | ш |
| Co ²⁺ | - 0.86 | и |
| Zn ²⁺ | - 1.08 | и |
| Mn ²⁺ | - 1.40 | и |

CATHODE RAY POLAROGRAPHY PEAK POTENTIALS OF SOME IMPORTANT METALS (1)

| Metal | Peak potential [V] | Support electrolyte / co | oncentration | Interference by |
|------------------|-----------------------|--------------------------|--------------|------------------------------------|
| Zn ²⁺ | - 1.03 | Pyridine hydrochloride | 0.1 mol/l | Co ²⁺ |
| Cd ²⁺ | - 0.63 | HCI | 0.2 mol/l | |
| Cu ²⁺ | - 0.15 | HCI | 0.2 mol/l | Sb ³⁺ |
| Pb ²⁺ | - 0.40 | HCI | 0.2 mol/l | Sn ²⁺ |
| Ni ²⁺ | - 0.80 | Pyridine hydrochloride | 0.1 mol/l | |
| Bi ³⁺ | - 0.08 | HCI | 0.5 mol/l | Cu ²⁺ |
| Co ²⁺ | - 1.05 | Pyridine hydrochloride | 0.1 mol/l | Zn ²⁺ |
| Sb ³⁺ | - 0.13 | HCI | 0.5 mol/l | Cu ²⁺ |
| Sn ²⁺ | - 0.50 | HCI | 5 mol/l | Pb ²⁺ , TI ⁺ |
| Cr ⁶⁺ | - 0.75 | LiOH | 0.5 mol/l | |
| Cr ³⁺ | - 1.10 | LiCI | 0.5 mol/l | Zn ²⁺ |

(1) measured against a silver/silver chloride reference electrode





| ENERGY DIMENSIONS – CONVERSION FACTORS | | | | | | | |
|--|-----|-------------------------|---------------------------|------------------------|----------------------------|---------------------------|-------------------------|
| Given dimension with conversion factor (2) | | | | | | | |
| Unit | | J | kWh | MeV | mkp | kcal _{15°} | erg |
| 1 J | (1) | 1 | 2.77778·10 ⁻⁷ | 6.242·10 ¹² | 0.1019716 | 2.38920·10 ⁻⁴ | 10 ⁷ |
| 1 kWh | (1) | 3600000 | 1 | 2.247·10 ¹⁹ | 367097.8 | 860.11 | 3.6·10 ¹³ |
| 1 MeV | (1) | 1.602·10 ⁻¹³ | 4.45·10 ⁻²⁰ | 1 | 1.634·10 ⁻¹⁴ | 3.827·10 ⁻¹⁷ | 1.602·10 ⁻⁶ |
| 1 mkp | | 9.80665 | 2.72407·10 ⁻⁶ | 6.124·10 ¹³ | 1 | 2.34301·10 ⁻³ | 9.80665-10 |
| kcal _{15°} | | 4185.5 | 1.16264·10 ⁻³ | 2.613·10 ¹⁶ | 426.80 | 1 | 4.1855·10 ¹⁰ |
| l erg | | 10 ⁻⁷ | 2.77778·10 ⁻¹⁴ | 6.242·10 ⁵ | 0.1019716·10 ⁻⁷ | 2.38920·10 ⁻¹¹ | 1 |

- (1) Legal measurement units
- (2) Examples: $1 J = 2.38920 \cdot 10 4 \text{ kcal } 1 \text{ MeV} = 1.602 \cdot 10 13J$

PRESSURE DIMENSIONS – CONVERSION FACTORS

| Given dimension | Required dimension with conversion factor (2) | | | | | | |
|---------------------------------|---|---------------------------|--------------------------|---------------------------|--------------------------|--------------------------|--|
| Unit | N·m⁻² (Pa) | bar | atm | kp·m ^{−2} | Torr (mm Hg) | lbs/sq.in. (psi) | |
| 1 N·m ⁻² 1 Pa (1) | 1 | 10-5 | 9.8692·10 ⁻⁶ | 1.019710·10 ⁻¹ | 7.50062·10 ⁻³ | 1.45038·10 ⁻⁴ | |
| 1 bar (1) | 10 ⁵ | 1 | 0.98692 | 10197.16 | 750.062 | 14.5038 | |
| 1 atm | 101325 | 1.01325 | 1 | 10332.27 | 759.9988 | 14.6960 | |
| 1 kp·m² | 9.80665 | 9.80665·10 ⁻⁵ | 9.67841·10 ⁻⁵ | 1 | 7.35559·10 ⁻² | 1.42234·10 ⁻³ | |
| 1 Torr (1 mm Hg) | 133.3224 | 1.333224·10 ⁻³ | 1.31579·10 ⁻³ | 13.5951 | 1 | 1.93368·10 ⁻² | |
| 10 lbs/sq.in. (psi) | 68948 | 0.68948 | 0.68046 | 7030.68 | 517.148 | 10 | |

- (1) Legal measurement units
- (2) Examples: 1 Pa = 7.50062·10-3Torr 10 psi = 0.68046 atm

DECIMAL UNIT MULTIPLES AND SUBDIVISIONS

| Prefix | | Symbol | Prefix | | Symbol |
|------------------|-------|--------|-------------------|-------|--------|
| 10 ¹⁸ | Exa | E | 10-1 | Deci | d |
| 10 ¹⁵ | Peta | Р | 10-2 | Centi | С |
| 10 ¹² | Tera | T | 10-3 | Milli | m |
| 10 ⁹ | Giga | G | 10 ⁻⁶ | Micro | μ |
| 10 ⁶ | Mega | M | 10-9 | Nano | n |
| 10 ³ | Kilo | k | 10 ⁻¹² | Piko | р |
| 10 ² | Hecto | h | 10-15 | Femto | f |
| 10 | Deca | da | 10-18 | Atto | а |

CONCENTRATIONS

| Proportion | Potency | % | g/kg mg/g µg/mg | ppm mg/kg μg/g ng/mg | ppb µg/kg ng/g pg/mg | ppt ng/kg pg/g fg/mg |
|------------------|-----------------------|-------------|-----------------------|-------------------------------|-------------------------------|-------------------------------|
| 1:100 | 1 x 10 ⁻² | 1 | 10 | 10 000 | | |
| 1:1000 | 1 x 10 ⁻³ | 0.1 | 1 | 1 000 | | |
| 1:10 000 | 1 x 10 ⁻⁴ | 0.01 | 0.1 | 100 | | |
| 1:100 000 | 1 x 10 ⁻⁵ | 0.001 | 0.01 | 10 | | |
| 1:1million | 1 x 10 ⁻⁶ | 0.000 1 | 0.001 | 1 | 1 000 | |
| 1:10 million | 1 x 10 ⁻⁷ | 0.000 01 | 0.000 1 | 0.1 | 100 | |
| 1:100 Millionen | 1 x 10 ⁻⁸ | 0.000 001 | 0.000 01 | 0.01 | 10 | |
| 1:1 Milliarde | 1 x 10 ⁻⁹ | 0.000 000 1 | 0.000 001 | 0.001 | 1 | 1 000 |
| 1:10 Milliarden | 1 x 10 ⁻¹⁰ | | | | 0.1 | 100 |
| 1:100 Milliarden | 1 x 10 ⁻¹¹ | | | | 0.01 | 10 |
| 1:1 Billion | 1 x 10 ⁻¹² | | | | 0.001 | 1 |

The tables on pages 96 to 101 were reprinted with permission of the Deutsches Institut für Normung e.V. (DIN, German Institute of Standardization). They contain excerpts from DIN 1301. Always use the most current version of the norm to be obtained from Beuth Verlag GmbH, Burggrafenstraße 4–10, 10772 Berlin, Germany.

BASIC UNITS

| Dimension | Basic unit Name | Symbol |
|---------------------|--------------------|--------|
| Length | Meter | m |
| Mass | Kilogram | kg |
| Time | Second | S |
| Electric current | Ampere | Α |
| Temperature | Kelvin | K |
| Luminous intensity | Candela | cd |
| Amount of substance | Mole | mol |

DERIVED UNITSWITH CONVERSION OF PREVIOUS UNITS

| Size | SI unit Name | Symbol | Further units Name Symbol | | Relationship |
|-------------------------|-----------------|----------------|------------------------------|---|---------------------|
| LENGTH, SURFACE, VOLUME | | | | | |
| Length | Meter | m | | | |
| Surface | Square meter | m ² | | | |
| Volume | Cubic meter | m^3 | Liter | 1 | $1 I = 10^{-3} m^3$ |

MASS

| Mass | Kilogram | kg | | | |
|----------|-----------------|----------------------------------|-----------|---|-----------------------|
| | | | Metric | | |
| | | | ton | t | $1 t = 10^3 kg$ |
| | | | Atomic | | 1 u = 1,66053 |
| | | | mass unit | u | ·10 ⁻²⁷ kg |
| Density | Kilogram per | | | | |
| | cubic meter | kg·m⁻³ | | | |
| Specific | Cubic meter | | | | |
| volume | per kilogram | m ³ ·kg ⁻¹ | | | |

AMOUNT OF SUBSTANCE

| Amount of substance Mole mol Molar Mass mass per amount |
|---|
| |
| of substance kg·mol ⁻¹ g·mol ⁻¹ |
| Concentration of a substance in given volume of solvent mol·m ⁻³ mol·l ⁻¹ |
| Molality Amount of substance per mass |
| of solvent mol·kg ⁻¹ mol·g ⁻¹ |

DERIVED UNITSWITH CONVERSION OF PREVIOUS UNITS

| Size | SI unit | | Further uni | 4. | Dalatianskin |
|-------------------|------------------------|-------------------|-----------------------|--------------------|---|
| Size | Name | Symbol | Name | ts Symbol | Relationship |
| | | | | | |
| TEMPERATURE | | | | | |
| Temperature | Kelvin | K | Degree centigrade | °C | |
| | | | | | |
| TIME | | | | | |
| Time | second | S | | | |
| Time interval | | | minute hour day | min h d | 1 min = 60 s 1 h = 60 min 1 d = 24 h |
| Frequency | Hertz | Hz | | | 1 Hz = 1s ⁻¹ |
| Velocity | Meter per second | m·s ^{−1} | | | |
| | | | Kilometer | | |
| | | | per hour | km·h ^{−1} | $1 \text{ km} \cdot \text{h}^{-1} \\ = \frac{1}{3.6} \text{ m} \cdot \text{s}^{-1}$ |
| | | | | | |
| VISCOSITY | | | | | |
| Dynamic viscosity | Pascal- second | Pa·s | | | 1 Pa·s = 1 N·s·m ⁻² = 1 kg·m ⁻¹ ·s ⁻¹ |

| Dynamic viscosity | Pascal- second | Pa·s | | | 1 Pa·s = 1 N·s·m ⁻² = 1 kg·m ⁻¹ ·s ⁻¹ |
|-------------------|-------------------|--------------------|------------|-----|---|
| | | | Poise | P | 1 P |
| | | | | _ | = 0.1 Pa·s |
| | | | Centipoise | cP | 1 cP = 0.01 P |
| | | | | | = 0.01 P = 0.001 Pa·s |
| | | | | | = mPa·s |
| Kinematic | Square | $m^2 \cdot s^{-1}$ | Stokes | St | 1 St |
| viscosity | meters | | | | $= 1 \text{ cm}^2 \cdot \text{s}_{-1}$ |
| | per | | Centistoke | cSt | 1 cSt |
| | second | | | | $= 1 \text{ mm}^{2} \cdot \text{s}^{-1}$ |

DERIVED UNITSWITH CONVERSION OF PREVIOUS UNITS

| Size | SI unit Name | Symbol | Further un Name | its Symbol | Relationship |
|-------------------------------|----------------------------------|-------------------|--------------------|---------------|---|
| FORCE, ENERGY, | , POWER | | | | |
| Force | Newton | N | | | 1 N = 1 kg·m·s ⁻² |
| Pressure | Newton per square meter | N·m ^{−2} | | | |
| | Pascal | Pa | Bar | bar | 1 Pa = 1 N·m ⁻² 1 bar = 10 ⁵ Pa |
| Energy, work heat quantity | Joule | J | | | 1 J = 1 N·m = 1 W·s = 1 kg·m ² ·s ⁻² |
| | | | Kilowatt- hour | kW∙h | 1 kW·h = 3.6 MJ |
| Power | Watt | W | | | 1 W = 1 J·s ⁻¹ = 1 Nm·s ⁻¹ = 1 VA |
| | | | | | |

DEDIVED LIMITS

| DERIVED UNITS | | | | | | | |
|-----------------------|-----------------|--------|------------------------------|--------------|--|--|--|
| Size | SI unit Name | Symbol | Further units Name Symbol | Relationship | | | |
| ELECTRICAL MEASURES | | | | | | | |
| Electric current | Ampere | Α | | | | | |
| Electric potential | Volt | V | | | | | |

Ampere-

hour

conductance Electric Ohm ٧ resistance Electric Coulomb С

Siemens

S

LUMINOUS INTENSITY

Electric

| Luminous intensity | Candela | cd |
|--------------------|---------|----|
| Luminous flux | Lumen | lm |
| Illuminance | Lux | lx |
| | | |

1 Im = 1 cd·sr $1 \text{ Ix} = 1 \text{ Im·m}^{-2}$ = 1 cd·sr·m⁻²

1 S $= 1 \text{ A} \cdot \text{V}^{-1}$

1 V

1 C

 $A \times h$

= 1 A·s

= 3600 A·s

= 1 C·V⁻¹

1 A·h

1 F

 $= 1 \text{ V} \cdot \text{A}^{-1}$ $= 1 S^{-1}$

DERIVED UNITS

| Size | SI unit Name | Symbol | Further unit Name | ts Symbol | Relationship | | | |
|-----------------------|-----------------|--------|----------------------|--------------|--|--|--|--|
| IONIZING RADIATION | | | | | | | | |
| Actitvity | Becquerel | Bq | Curie | Ci | 1 Bq = 1 s ⁻¹ 1 Ci = 37 G Bq | | | |
| ENZYMATIC ACT | TIVITY | | | | | | | |
| Enzymatic activity | Katal | kat | Enzyme unit | U | | | | |

US AND BRITISH MEASURING UNITS CONVERSION TO METRIC UNITS

VOLUME

| Liters | Х | 0.2642 | = US gallons |
|------------------|---|---------|--------------------|
| US gallons | х | 3.785 | = Liters |
| Imperial gallons | Х | 1.201 | = US gallons |
| Imperial gallons | х | 4.546 | = Liters |
| US gallons | Х | 0.8327 | = Imperial gallons |
| Cubic meters | х | 35.31 | = Cubic feet |
| Cubic feet | Х | 0.0283 | = Cubic meters |
| Cubic meters | х | 264.2 | = US gallons |
| US gallons | Х | 0.00379 | = Cubic meters |

LENGTH

| Meters | X | 3.281 | = Feet |
|-------------|---|--------|---------------|
| Feet | х | 0.3048 | = Meters |
| Meters | Х | 39.37 | = Inches |
| Inches | Х | 0.0254 | = Meters |
| Centimeters | X | 0.3937 | = Inches |
| Inches | х | 2.540 | = Centimeters |
| Millimeters | Х | 0.0394 | = Inches |
| Inches | х | 25.4 | = Millimeters |

WEIGHT

| Kilograms | X | 2.2046 | = Pounds |
|----------------|---|-----------|---------------|
| Pounds | Х | 0.4536 | = Kilograms |
| Tons (long) | X | 1016.05 | = Kilograms |
| Tons (long) | Х | 2240 | = Pounds |
| Tonnes (metr.) | X | 1000 | = Kilograms |
| Tonnes (metr.) | Х | 2204.6 | = Pounds |
| Tons (short) | X | 907.185 | = Kilograms |
| Tons (short) | Х | 2000 | = Pounds |
| Grams | X | 15.432653 | = Grains |
| Grains | Х | 0.0647989 | = Grams |
| Grams | X | 0.0352740 | = Ounces (US) |
| Ounces (US) | Х | 28.349527 | = Grams |
| Ounces (troy) | V | 31 1035 | - Grams |

US AND BRITISH MEASURING UNITS CONVERSION TO METRIC UNITS

TEMPERATURE

| given in | °Centigrade | required in °Réaumur | °Fahrenheit |
|----------------|--------------------|-------------------------|------------------------------|
| a ° Centigrade | - | <u>a · 8</u> 10 | $\frac{a \cdot 18}{10} + 32$ |
| b ° Réaumur | <u>b ⋅ 10</u> 8 | - | $\frac{b\cdot 18}{8} + 32$ |
| c ° Fahrenheit | (c - 32) · 10 | (c - 32) · 8 | _ |

| ° C | ° F | ° C | ° F | ° C | ° F | ° C | ° F | ° C | ° F | ° C | ° F |
|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| - 40 | - 40 | + 5 | 41 | 50 | 122 | 95 | 203 | 140 | 284 | 185 | 365 |
| - 35 | - 31 | 10 | 50 | 55 | 131 | 100 | 212 | 145 | 293 | 190 | 374 |
| - 30 | - 22 | 15 | 59 | 60 | 140 | 105 | 221 | 150 | 302 | 195 | 383 |
| - 25 | - 13 | 20 | 68 | 65 | 149 | 110 | 230 | 155 | 311 | 200 | 392 |
| - 20 | - 4 | 25 | 77 | 70 | 158 | 115 | 239 | 160 | 320 | 210 | 410 |
| - 15 | + 5 | 30 | 86 | 75 | 167 | 120 | 248 | 165 | 329 | | |
| - 10 | 14 | 35 | 95 | 80 | 176 | 125 | 257 | 170 | 338 | | |
| - 5 | 23 | 40 | 104 | 85 | 185 | 130 | 266 | 175 | 347 | | |
| Ω | 32 | 45 | 113 | 90 | 194 | 135 | 275 | 180 | 356 | | |



PHYSICAL CONSTANTS

| Constant | Symbol | Value | | | |
|---|--------------------------------------|----------|---|-------------------|-------------------------------------|
| Atomic mass unit | $m_{\scriptscriptstyle \cup}$ | 1.660540 | | 10 ⁻²⁷ | kg |
| Avogadro constant | N _A | 6.022137 | | 10 ²³ | mol ⁻¹ |
| Bohr magneton | $\mu_{\scriptscriptstyle B}$ | 9.274015 | | 10 ⁻²⁴ | JT ⁻¹ |
| Bohr radius | a_{0} | 5.291771 | | 10 ⁻¹¹ | m |
| Boltzmann constant | k _B | 1.380662 | | 10 ⁻²³ | JK ⁻¹ |
| Compton wavelength (e) | $\lambda_{	ext{Ce}}$ | 2.426311 | | 10-12 | m |
| Compton wavelength (n) | $\lambda_{	extsf{Cn}}$ | 1.319591 | | 10 ⁻¹⁵ | m |
| Compton wavelength (p) | λ_{Cp} | 1.321410 | | 10 ⁻¹⁵ | m |
| Electric field constant in vacuo | $\mathcal{E}_{\scriptscriptstyle 0}$ | 8.854188 | | 10 ⁻¹² | Fm ⁻¹ |
| Electron radius | $	au_{ m e}$ | 2.817941 | | 10 ⁻¹⁵ | m |
| Elementary charge | e | 1.602177 | | 10 ⁻¹⁹ | С |
| Faraday constant | F | 9.648531 | | 10 ⁴ | Cmol ⁻¹ |
| Fine structure constant | α | 7.297353 | | 10-3 | |
| Gas constant | R | 8.31451 | | | J mol ⁻¹ K ⁻¹ |
| Gravitation constant | f | 6.672590 | | 10 ⁻¹¹ | Nm ² kg ⁻² |
| Intrinsic impedance | Γ | 3.767301 | | 10 ² | Ω |
| Light velocity in vacuo | с | 2.997924 | | 10 ⁸ | ms ⁻¹ |
| Loschmidt constant | $N_{\scriptscriptstyle L}$ | 2.686763 | | 10 ²⁵ | m ⁻³ |
| Magnetic field constant in vacuo | μ_0 | 1.256637 | | 10 ⁻⁷ | Hm ⁻¹ |
| Molar volume of ideal gases 298 K, 101.325 kPa | υ | 2.445294 | ٠ | 10-2 | m ³ mol ⁻¹ |
| Normal acceleration of fall | g | 9.80665 | | | ms ⁻² |
| Planck constant | h | 6.626075 | | 10-34 | Js |
| Rest mass of the electron | m _e | 9.109390 | | 10 ⁻³¹ | kg |
| Rest mass of the neutron | m _n | 1.674929 | | 10-27 | kg |
| Rest mass of the proton | $m_{_{\mathrm{p}}}$ | 1.672623 | | 10 ⁻²⁷ | kg |
| Rotational quantum | $h/(2\pi)$ | 1.054588 | | 10-34 | Js |
| ydberg constant | R _∞ | 1.097373 | | 10 ⁷ | m-1 |

CREATING CONSTANT AIR HUMIDITY IN CLOSED VESSELS

| IN CEOSED VESSEES | | | | | | | |
|---|--|--|--|--|--|--|--|
| Saturated aqueous solution with considerable precipitates | | % relative air humidity above the solution (at 20°C) | | | | | |
| di-Sodium hydrogen phosphate | $\mathrm{Na_2HPO_4} \cdot 12\mathrm{H_2O}$ | 95 | | | | | |
| Sodium carbonate | Na ₂ CO ₃ · 10 H ₂ O | 92 | | | | | |
| Zinc sulfate | $ZnSO_4 \cdot 7 H_2O$ | 90 | | | | | |
| Potassium chloride | KCI | 86 | | | | | |
| Ammonium sulfate | (NH ₄) ₂ SO ₄ | 80 | | | | | |
| Sodium chloride | NaCl | 76 | | | | | |
| Sodium nitrite | NaNO ₂ | 65 | | | | | |
| Ammonium nitrate | NH ₄ NO ₃ | 63 | | | | | |
| Calcium nitrate | $\mathrm{Ca} \; \mathrm{(NO_3)_2} \cdot 4 \; \mathrm{H_2O}$ | 55 | | | | | |
| Potassium carbonate | K ₂ CO ₃ | 45 | | | | | |
| Zinc nitrate | $\mathrm{Zn}\; \mathrm{(NO_3)_2} \cdot \mathrm{6}\; \mathrm{H_2O}$ | 42 | | | | | |
| Calcium chloride | CaCl ₂ · 6 H ₂ O | 32 | | | | | |
| Lithium chloride | LiCl · H₂O | 15 | | | | | |



GREEK ALPHABET

| Letter | | Name | Pronunciation | | | | |
|--------|--------|---------|---------------|--|--|--|--|
| | | | | | | | |
| A o | χ | álpha | а | | | | |
| Вβ | 3 | béta | ь | | | | |
| Γ γ | 1 | gámma | g | | | | |
| Δ δ | 5 | délta | d | | | | |
| Ε ε | : | épsilon | e (short) | | | | |
| Zζ | | zéta | Z | | | | |
| Н п | 1 | éta | e (long) | | | | |
| Θ ΰ | 9 | théta | th | | | | |
| Ι ι | | ióta | i | | | | |
| К к | c | káppa | k | | | | |
| Λ λ | l | lámbda | 1 | | | | |
| Μ μ | ı | mü | m | | | | |
| N v | , | nü | n | | | | |
| Ξ ξ | | xi | х | | | | |
| 0 0 |) | ómicron | o (short) | | | | |
| Π π | τ | pi | р | | | | |
| Ρ ρ |) | rho | r | | | | |
| Σ σ | 51, ç2 | sigma | S | | | | |
| Τ τ | : | tau | t | | | | |
| Y |) | ýpsilon | у | | | | |
| Φ φ | Р | phi | ph | | | | |
| Χ χ | c | chi | ch | | | | |
| ΨΨ | ų | psi | ps | | | | |
| Ω σ | 0 | ómega | o (long) | | | | |

¹ At the beginning and in the middle of a word.

2 At the end of a word.

GREEK NUMERALS / ROMAN NUMBERS

| 1/2 | hemi – | |
|------|-------------------|-------|
| 1 | mono – | 1 |
| 11/2 | sesqui – | |
| 2 | di –, bi – | H |
| 21/2 | hemipenta – | |
| 3 | tri – | III |
| 4 | tetra – | IV |
| 5 | penta – | V |
| 6 | hexa – | VI |
| 7 | hepta – | VII |
| 8 | octa – | VIII |
| 9 | nona –, ennea – | IX |
| 10 | deca – | Χ |
| 11 | hendeca –, undeca | XI |
| 12 | dodeca – | XII |
| 13 | trideca | XIII |
| 14 | tetradeca – | XIV |
| 15 | pentadeca – | XV |
| 16 | hexadeca – | XVI |
| 17 | heptadeca – | XVII |
| 18 | octadeca – | XVIII |
| 19 | nonadeca – | XIX |
| 20 | eicosa – | XX |
| 40 | tetraconta – | XL |
| 50 | pentaconta – | L |
| 60 | hexaconta – | LX |
| 90 | nonaconta – | XC |
| 99 | | IC |
| 100 | hecta – | С |
| 200 | | CC |
| 400 | | CD |
| 500 | | D |
| 600 | | DC |
| 900 | | CM |
| 990 | | XM |
| 1000 | | M |

FREEZING MIXTURES

| The numbers represe | Lower tempe from | ing rature [°C] to | |
|---------------------|---|--------------------------|-------|
| 4 water | + 1 potassium chloride | + 10 | - 12 |
| 1 water | + 1 ammonium nitrate | + 10 | - 15 |
| 1 water | + 1 sodium nitrate + 1 ammonium chloride | + 8 | - 24 |
| 3 ice ground | + 1 sodium chloride | 0 | - 21 |
| 1,2 ice ground | + 1 magnesium chloride (MgCl $_2 \cdot$ 7 H $_2$ 0) | 0 | - 34 |
| 1,2 ice ground | + 2 calcium chloride (CaCl ₂ · 6 H ₂ 0) | 0 | - 39 |
| 1,4 ice ground | + 2 calcium chloride ($CaCl_2 \cdot 6 H_20$) | 0 | - 55 |
| Methanol or acetone | + dry ice | + 15 | - 77 |
| Diethyl ether | + dry ice | + 15 | - 100 |

EXTRAN® LABORATORY CLEANING AGENTS

| LXIII | IN LADOIN | ATONT CLLANING AGENTS |
|----------|---|--|
| Cat. No. | Name | Notes on use |
| 1.07555 | Extran® MA 01 alkaline / liquid | Universal cleaning agent for heavy contamination. For hard water even up to 40° d. For cleaning tables, tiles, and floors in the laboratory. Tested for radioactive decontamination. Suitable for ultrasonic cleaning. |
| 1.07553 | Extran® MA 02 neutral / liquid | Special cleaner for precision instruments of glass, quartz and sensitive metals. Suitable for ultrasonic cleaning. |
| 1.07550 | Extran® MA 03 phosphate-free liquid | Universal cleaning agent for heavy contamination. With very hard water also usable without restric tions. Environmentally friendly as it contains no phosphate. Suitable for ultrasonic cleaning. |
| 1.07551 | Extran® MA 04 disinfecting | For manual instrument disinfection. |
| 1.07558 | Extran® AP 11 mildly alkaline/ powder | Gentle cleaning action; e.g. in the analytical laboratory. Cleaning action equivalent to that of AP 14 liquid. |
| 1.07563 | Extran® AP 12 alkaline/powder | Powerful cleaning action. Particularly with starch and protein residues. Cleaning action equivalent to that of AP 15 liquid. |
| 1.07565 | Extran® AP 13 alkaline with detergents/ powder | Powerful cleaning action. Particularly with fat residues. |
| 1.07573 | Extran® AP 14 mildly alkaline/ liquid | Gentle cleaning action for machines with liquid liquid dosing; e.g. in the analytical laboratory. Environmentally friendly as it contains no phosphate. Cleaning action equivalent to that of AP 11 powder. |
| 1.07575 | Extran [®] AP 15 alkaline/liquid | Powerful cleaning for machines with liquid dosing. Environmentally friendly as it contains no phosphate. Cleaning action equivalent to that of AP 12 powder. |
| 1.07559 | Extran® AP 21 acidic with phosphoric acid liquid | First rinse in the presence of residues of carbonates, hydroxides, proteins, amines, etc. Neutralising Neutralising final rinse. Also for a gentle main wash. Prevents the formation of calcareous deposits. |
| 1.07561 | Extran® AP 22 acidic with citric acid liquid | Gentle prerinse or final rinse with neutralizing action. Prevents the formation of calcareous deposits. Environmentally friendly as it contains no phosphate. |
| 1.07560 | Extran® AP 31 antifoam/ liquid | Additive for foam-forming residues: proteins, fats, emulsifiers of all types. |
| 1.07556 | Extran® AP 32 clear-rinse/ liquid | Final rinse: gives spotless surfaces even with hard water. Environmentally friendly as it contains no phosphate. |
| 1.07570 | Extran® AP 41 enzymatic/ powder | For medical and dental practices, anaesthetic equipment. For the removal of mucus, saliva, blood etc. Temperature: 55–65 °C. |

EXTRAN® LABORATORY CLEANING AGENT

| Cat. No. | | Notes on use |
|----------|--|---|
| 1.07571 | Extran® AP 42 fat emulsion/ liquid | Machine washing: in the final rinse for restoring the lubrication of surgical instruments. Manually: 5-minute immersion in 100 ml/l |
| 1.07584 | Extran® AP 43 bactericidal/ liquid | For disinfection after the main wash; e.g. operation shoes. Temperature: 40–60 °C/15 minutes. Environmentally friendly as it contains no phosphate. |



PARTICLE SIZES

Mesh size w

| Mesh size w | | | | |
|--|-----------------------------------|----------------------------------|----------------------------------|-----------------|
| DIN 4188 (D) [mm] | ASTM E11-70 (USA) [mesh] | ASTM E161-70 (USA) [μm] | BS 410 : 1969 (GB) [μm] | Tyler [mesh] |
| | | 5 | | |
| | | 10 15 | | |
| 0.02 0.022 0.025 | | 22 | | |
| 0.023 | | 27 | | |
| 0.028 0.032 0.036 | | 32 | | |
| | 400 | 38 | 38 | 400 |
| 0.04 0.045 | 325 | 45 | 45 | 325 |
| 0.045 0.05 0.056 0.063 0.071 | 323 | 43 | 45 | 323 |
| | 270 | 53 | 53 | 270 |
| | 230 | 63 | 63 | 250 |
| | 200 | 75 | 75 | 200 |
| 0.08 0.09 0.1 | 170 | 90 | 90 | 170 |
| 0.1 | 140 | 106 | 106 | 150 |
| 0.112 0.125 0.14 | 120 | 125 | 125 | 115 |
| | 100 | 150 | 150 | 100 |
| 0.16 | | | | |
| 0.18 | 80 | | 180 | 80 |
| 0.2 | 70 | | 212 | C.F. |
| | 70 | | 212 | 65 |
| 0.224 0.25 0.28 | 60 | | 250 | 60 |
| | 50 | | 300 | 48 |
| 0.315 | | | | |
| 0.355 0.4 | 45 | | 355 | 42 |

PARTICLE SIZES

Mesh size w

| Mesh size w | | | | |
|-------------------------|-----------------------------------|----------------------------------|--------------------------------|-----------------|
| DIN 4188 (D) [mm] | ASTM E11-70 (USA) [mesh] | ASTM E161–70 (USA) [μm] | BS 410:1969 (GB) [μm] | Tyler [mesh] |
| | 40 | | 425 | 35 |
| 0.45 0.5 0.56 | 35 | | 500 | 32 |
| | 30 | | 600 | 28 |
| 0.63 0.71 0.8 | 25 | | 710 | 24 |
| | 20 | | 850 | 20 |
| 0.9 1 1.12 | 18 | | 1000 | 16 |
| 1.18 | 16 | | 1180 | 14 |
| 1.25 1.4 1.6 | 14 | | 1400 | 12 |
| | 12 | | 1700 | 10 |
| 1.8 2 2.24 | 10 | | 2000 | 9 |
| | 8 | | 2360 | 8 |
| 2.5 2.8 3.15 | 7 | | 2800 | 7 |
| | 6 | | 3350 | 6 |
| 3.55 4 4.5 | 5 | | 4000 | 5 |
| | 4 | | 4750 | 4 |
| 5 5.6 | 31/2" | | 5600 | 31/2 |

NMR: CARBON (13C) CHEMICAL SHIFTS

| TMS = 0 ppm | | 22 | 20 2 | 00 1 | 80 1 | 60 1 | 40 1 | 20 1 | 00 8 | 80 6 | 60 4 | 10 2 | .0 (|) -2 | 20 -40 |
|-----------------------------|--------------|----|------|------|------|------|------|------|------|------|------|------|-----------|----------|---------|
| H ₃ C − C € | C primary | | | | | | | | | | | | | | |
| H ₃ C – Halogen | | | | | | | | | | | | CI | | | I |
| H ₃ C - 0 - | | | | | | | | | | | | | | | |
| H ₃ C - N< | | | | | | | | | | | | | | | |
| H ₃ C - S - | | | | | | | | | | | | | | | |
| - CH ₂ - C € | C secondary | | | | | | | | | | | | aliphates | — cyclop | ropanes |
| - CH ₂ - Halogen | | | | | | | | | | | CI | | | ı | |
| - CH ₂ - 0 - | | | | | | | | | | | | | | | |
| - CH ₂ - N < | | | | | | | | | | | | | | | |
| - CH ₂ - S - | | | | | | | | | | | | | | | |
| > CH - C € | C tertiary | | | | | | | | | | | | | | |
| >CH - Halogen | | | | | | | | | | CI | | -1 | | | |
| >CH - 0 - | | | | | | | | | | | | | | | |
| >CH - N < | | | | | | | | | | | | | | | |
| >CH - S - | | | | | | | | | | | | | | | |
| ≥C - C € | C quaternary | | | | | | | | | | | | | | |
| ∋C – Halogen | | | | | | | | | | CI | | ı | | | |
| ≥C - O - | | | | | | | | | | | | | | | |



NMR: CARBON (13C) CHEMICAL SHIFTS


| TMS = 0 ppm | | 2 | 20 2 | 00 1 | 80 | 160 1 | 40 1 | 20 1 | 00 8 | 30 (| 60 4 | 10 2 | 20 | 0 - | 20 |
|------------------------|-----------------|---|------|------|----|-------|------|------|------|------|------|------|----|-----|----|
| ≥C - S - | | | | | | | | | | | | | | | |
| ≥C - C € | Alkanes | | | | | | | | | | | | | | |
| - C <u>=</u> C - | Alkynes | | | | | | | | | | | | | | |
| > C = C < | Alkenes | | | | | | | | | | | | | | |
| > C = C < | Aromatics | | | | | | | | | | | | | | |
| > C = C < | Heteroaromatics | | | | | | | | | | | | | | |
| - 0 - C≡N | Cyanates | | | | | | | | | | | | | | |
| - S - C ≡ N | Thiocyanates | | | | | | | | | | | | | | |
| - C ≡ N | Cyanides | | | | | | | | | | | | | | |
| > C = N - | Azomethines | | | | | | | | | | | | | | |
| (- CO) ₂ O | Anhydrides | | | | | | | | | | | | | | |
| - COOR | Esters | | | | | | | | | | | | | | |
| (- CO) ₂ NR | Imides | | | | | l | | | | | | | | | |
| - CONHR | Amides | | | 1 | | | | | | | | | | | |
| - COOH | Acids | | | | | | | | | | | | | | |
| - COCI | Acid chlorides | | | _ | | ı | | | | | | | | | |
| - C = 0 | Aldehydes | | | | | | | | | | | | | | |
| >C = 0 | Ketones | | | | | | | | | | | | | | |



NMR: PROTON CHEMICAL SHIFTS

| TMS = 0 ppm | | 3 1 | 2 1 | 1 1 | 0 ! | 9 8 | 3 | 7 (| 6 ! | 5 4 | 1 3 | 3 : | 2 1 | l (| 0 |
|-------------------------------------|-------------------|-----|-----|-----|-----|-----|---|-----|-----|-----|-----|-----|-----|-----|---|
| H ₃ C - CR ₃ | Methyl protons | | | | | | | | | | | | | | |
| H ₃ C – Halogen | | | | | | | | | | F = | | | 1 | | |
| H ₃ C - 0 - | | | | | | | | | | | | | | | |
| H ₃ C - N < | | | | | | | | | | | | | | | |
| H ₃ C >C = C < | | | | | | | | | | | | | | | |
| $H_3C > C = 0$ | | | | | | | | | | | | | | | |
| H ₃ C – Ar | | | | | | | | | | | | | | | Г |
| - CH ₂ - CR ₃ | Methylene protons | | | | | | | | | | | | | | |
| – CH ₂ – Halogen | | | | | | | | | | F | | | 1 | | Г |
| - CH ₂ - 0 - | | | | | | | | | | | | | | | |
| - CH ₂ - N < | | | | | | | | | | | | | | | Г |
| - CH ₂ >C = C < | | | | | | | | | | | | | | | |
| - CH ₂ >C = 0 | | | | | | | | | | | | | | | Г |
| - CH ₂ - Ar | | | | | | | | | | | | | | | |
| >CH - CR ₃ | Methine protons | | | | | | | | | | | | | • | |
| >CH - Halogen | | | | | | | | F | | | ı. | | | | |
| >CH - 0 - | | | | | | | | | | | | | | | |
| >CH - N < | | | | | | | | | | | | | | | |

NMR: PROTON CHEMICAL SHIFTS



*The chemical shifts of these groups are concentration-dependent and are shifted to

lower ppm values in more dilute solutions.


(ppm)



HEATING BATHS

| | Media | Melting point (°C) | Boiling point (°C) | Working range (°C) | Flash point (°C) | Remarks |
|--------|------------------------------|-----------------------|-----------------------|-----------------------|---------------------|---|
| | Water | 0 | 100 | 0-80 | - within the | ldeal stated range |
| 822329 | Ethylene glycol | -12 | 198 | -10-180 | 111 | Flammable, difficult to remove from equipment |
| 107742 | Silicon oil | -60 | - | 20-250 | 300 | Becomes viscid at lower temperatures – noncorrosive |
| 808245 | Triethylene glycol | -7 | 287 | 0-250 | 156 | Water-soluble, stable |
| 818709 | Glycerol | 18 | 290 | 20-260 | 160 | Water-soluble, nontoxic, viscid |
| 107162 | Paraffin, liquid | - | 300-500 | 20-200 | 230 | Flammable |
| 107160 | Paraffin viscious | - | 300-500 | 20-200 | 230 | Flammable |
| 106900 | Oil bath filling | -5 | - | 0-250 | 300 | Flammable |
| 115265 | Heating bath fluid | -40 | - | 0-170 | 280 | Miscible with water, nontoxic |
| 105061 | Potassium nitrate (51.3%) | 219 | - | 230-500 | - | Stable in air, strong oxidizing agent, nonflammable |
| 106535 | Sodium nitrate (48.7%) | | | | | |
| 106482 | Sodium hydroxide (40%) | 167 | - | 200-800 | - | Corrosive and harmful |
| 105032 | Potassium hydroxide (60%) | | - | | - | |

PERIODIC TABLE OF THE FLEMENTS



PERIONIC TABLE OF THE FLEMEN

We provide information and advice to our customers to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

Merck KGaA 64271 Darmstadt, Germany Fax 0049 (0) 61 51 72-60 80 E-mail: reagents@merck.de Internet: www.merck.de www.reagents.merck.de