

PURIFICATION
OF
LABORATORY CHEMICALS
Fifth Edition

Wilfred L. F. Armarego

Visiting Fellow

Division of Biomolecular Science

The John Curtin School of Medical Research

Australian National University, Canberra

A. C. T., Australia

Christina Li Lin Chai

Reader in Chemistry

Department of Chemistry

Australian National University, Canberra

A. C. T., Australia

BUTTERWORTH
HEINEMANN

An Imprint of Elsevier Science

www.bh.com

Amsterdam Boston London New York Oxford Paris
San Diego San Francisco Singapore Sydney Tokyo

Butterworth-Heinemann is an imprint of Elsevier Science.

Copyright © 2003, Elsevier Science (USA). All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher.



Recognizing the importance of preserving what has been written, Elsevier-Science prints its books on acid-free paper whenever possible.

Library of Congress Cataloging-in-Publication Data

A catalog record for this book is available from the Library of Congress.

ISBN: 0-7506-7571-3

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

The publisher offers special discounts on bulk orders of this book.

For information, please contact:

Manager of Special Sales
Elsevier Science
200 Wheeler Road
Burlington, MA 01803
Tel: 781-313-4700
Fax: 781-313-4882

For information on all Butterworth-Heinemann publications available, contact our World Wide Web home page at: <http://www.bh.com>

10 9 8 7 6 5 4 3 2 1

Printed and bound in Great Britain by MPG Books Ltd, Bodmin, Cornwall

Preface to the Fifth Edition

THE DEMAND for **Purification of Laboratory Chemicals** has not abated since the publication of the fourth edition as evidenced by the number of printings and the sales. The request by the Editor for a fifth edition offered an opportunity to increase the usefulness of this book for laboratory purposes. It is with deep regret that mention should be made that Dr Douglas D. Perrin had passed away soon after the fourth edition was published. His input in the first three editions was considerable and his presence has been greatly missed. A fresh, new and young outlook was required in order to increase the utility of this book and it is with great pleasure that Dr Christina L.L. Chai, a Reader in Chemistry and leader of a research group in organic and bio-organic chemistry, has agreed to coauthor this edition. The new features of the fifth edition have been detailed below.

Chapters 1 and 2 have been reorganised and updated in line with recent developments. A new chapter on the 'Future of Purification' has been added. It outlines developments in syntheses on solid supports, combinatorial chemistry as well as the use of ionic liquids for chemical reactions and reactions in fluorous media. These technologies are becoming increasingly useful and popular so much so that many future commercially available substances will most probably be prepared using these procedures. Consequently, a knowledge of their basic principles will be helpful in many purification methods of the future.

Chapters 4, 5 and 6 (3, 4 and 5 in the 4th edn) form the bulk of the book. The number of entries has been increased to include the purification of many recent commercially available reagents that have become more and more popular in the syntheses of organic, inorganic and bio-organic compounds. Several purification procedures for commonly used liquids, e.g. solvents, had been entered with excessive thoroughness, but in many cases the laboratory worker only requires a simple, rapid but effective purification procedure for immediate use. In such cases a **Rapid purification** procedure has been inserted at the end of the respective entry, and should be satisfactory for most purposes. With the increased use of solid phase synthesis, even for small molecules, and the use of reagents on solid support (e.g. on polystyrene) for reactions in liquid media, compounds on solid support have become increasingly commercially available. These have been inserted at the end of the respective entry and have been listed in the General Index together with the above rapid purification entries.

A large number of substances are ionisable in aqueous solutions and a knowledge of their ionisation constants, stated as pK (pKa) values, can be of importance not only in their purification but also in their reactivity. Literature values of the pK's have been inserted for ionisable substances, and where values could not be found they were estimated (pK_{Est}). The estimates are usually so close to the true values as not to affect the purification process or the reactivity seriously. The book will thus be a good compilation of pK values for ionisable substances.

Almost all the entries in Chapters 4, 5 and 6 have CAS (Chemical Abstract Service) Registry Numbers to identify them, and these have been entered for each substance. Unlike chemical names which may have more than one synonymous name, there is only one CAS Registry Number for each substance (with only a few exceptions, e.g. where a substance may have another number before purification, or before determination of absolute configuration). To simplify the method for locating the purification of a substance, a CAS Registry Number Index with the respective page numbers has been included after the General Index at the end of the book. This will also provide the reader with a rapid way to see if the purification of a particular

substance has been reported in the book. The brief General Index includes page references to procedures and equipment, page references to abbreviations of compounds, e.g. TRIS, as well as the names of substances for which a Registry Number was not found.

Website references for distributors of substances or/and of equipment have been included in the text. However, since these may be changed in the future we must rely on the suppliers to inform users of their change in website references.

We wish to thank readers who have provided advice, constructive criticism and new information for inclusion in this book. We should be grateful to our readers for any further comments, suggestions, amendments and criticisms which could, perhaps, be inserted in a second printing of this edition. In particular, we thank Professor Ken-chi Sugiura (Graduate School of Science, Tokyo Metropolitan University, Japan) who has provided us with information on the purification of several organic compounds from his own experiences, and Joe Papa BS MS (EXAXOL in Clearwater, Florida, USA) who has provided us not only with his experiences in the purification of many inorganic substances in this book, but also gave us his analytical results on the amounts of other metal impurities at various stages of purification of several salts. We thank them graciously for permission to include their reports in this work. We express our gratitude to Dr William B. Cowden for his generous advice on computer hardware and software over many years and for providing an Apple LaserWriter (16/600PS) which we used to produce the master copy of this book. We also extend our sincere thanks to Dr Bart Eschler for advice on computer hardware and software and for assistance in setting up the computers (iMac and eMac) used to produce this book.

We thank Dr Pauline M. Armarego for assistance in the painstaking task of entering data into respective files, for many hours of proofreading, correcting typographical errors and checking CAS Registry Numbers against their respective entries.

One of us (W.L.F.A) owes a debt of gratitude to Dr Desmond (Des) J. Brown of the Research School of Chemistry, ANU, for unflinching support and advice over several decades and for providing data that was difficult to acquire not only for this edition but also for the previous four editions of this book.

One of us (C.L.L.C) would specially like to thank her many research students (past and present) for their unwavering support, friendship and loyalty, which enabled her to achieve what she now has. She wishes also to thank her family for their love, and would particularly like to dedicate her contribution towards this book to the memory of her brother Andrew who had said that he should have been a scientist.

We thank Mrs Joan Smith, librarian of the Research School of Chemistry, ANU, for her generous help in many library matters which have made the tedious task of checking references more enduring.

W.L.F. Armarego & C.L.L. Chai
November 2002

Preface to the First Edition

WE BELIEVE that a need exists for a book to help the chemist or biochemist who wishes to purify the reagents she or he uses. This need is emphasised by the previous lack of any satisfactory central source of references dealing with individual substances. Such a lack must undoubtedly have been a great deterrent to many busy research workers who have been left to decide whether to purify at all, to improvise possible methods, or to take a chance on finding, somewhere in the chemical literature, methods used by some previous investigators.

Although commercially available laboratory chemicals are usually satisfactory, as supplied, for most purposes in scientific and technological work, it is also true that for many applications further purification is essential.

With this thought in mind, the present volume sets out, firstly, to tabulate methods, taken from the literature, for purifying some thousands of individual commercially available chemicals. To help in applying this information, two chapters describe the more common processes currently used for purification in chemical laboratories and give fuller details of new methods which appear likely to find increasing application for the same purpose. Finally, for dealing with substances not separately listed, a chapter is included setting out the usual methods for purifying specific classes of compounds.

To keep this book to a convenient size, and bearing in mind that its most likely users will be laboratory-trained, we have omitted manipulative details with which they can be assumed to be familiar, and also detailed theoretical discussion. Both are readily available elsewhere, for example in Vogel's very useful book **Practical Organic Chemistry** (Longmans, London, 3rd ed., 1956), or Fieser's **Experiments in Organic Chemistry** (Heath, Boston, 3rd ed, 1957).

For the same reason, only limited mention is made of the kinds of impurities likely to be present, and of the tests for detecting them. In many cases, this information can be obtained readily from existing monographs.

By its nature, the present treatment is not exhaustive, nor do we claim that any of the methods taken from the literature are the best possible. Nevertheless, we feel that the information contained in this book is likely to be helpful to a wide range of laboratory workers, including physical and inorganic chemists, research students, biochemists, and biologists. We hope that it will also be of use, although perhaps to only a limited extent, to experienced organic chemists.

We are grateful to Professor A. Albert and Dr D.J. Brown for helpful comments on the manuscript.

D.D.P., W.L.F.A. & D.R.P.
1966

Preface to the Second Edition

SINCE the publication of the first edition of this book there have been major advances in purification procedures. Sensitive methods have been developed for the detection and elimination of progressively lower levels of impurities. Increasingly stringent requirements for reagent purity have gone hand-in-hand with developments in semiconductor technology, in the preparation of special alloys and in the isolation of highly biologically active substances. The need to eliminate trace impurities at the micro- and nanogram levels has placed greater emphasis on ultrapurification technique. To meet these demands the range of purities of laboratory chemicals has become correspondingly extended. Purification of individual chemicals thus depends more and more critically on the answers to two questions - Purification from what, and to what permissible level of contamination. Where these questions can be specifically answered, suitable methods of purification can usually be devised.

Several periodicals devoted to ultrapurification and separations have been started. These include "Progress in Separation and Purification" Ed. (vol. 1) E.S. Perry, Wiley-Interscience, New York, vols. 1-4, 1968-1971, and **Separation and Purification Methods** Ed. E.S.Perry and C.J.van Oss, Marcel Dekker, New York, vol. 1- , 1973-. Nevertheless, there still remains a broad area in which a general improvement in the level of purity of many compounds can be achieved by applying more or less conventional procedures. The need for a convenient source of information on methods of purifying available laboratory chemicals was indicated by the continuing demand for copies of this book even though it had been out of print for several years.

We have sought to revise and update this volume, deleting sections that have become more familiar or less important, and incorporating more topical material. The number of compounds in Chapters 3 and 4 have been increased appreciably. Also, further details in purification and physical constants are given for many compounds that were listed in the first edition.

We take this opportunity to thank users of the first edition who pointed out errors and omissions, or otherwise suggested improvements or additional material that should be included. We are indebted to Mrs S.Schenk who emerged from retirement to type this manuscript.

D.D.P., W.L.F.A. & D.R.P.
1980

Preface to the Third Edition

THE CONTINUING demand for this monograph and the publisher's request that we prepare a new edition, are an indication that **Purification of Laboratory Chemicals** fills a gap in many chemists' reference libraries and laboratory shelves. The present volume is an updated edition which contains significantly more detail than the previous editions, as well as an increase in the number of individual entries and a new chapter.

Additions have been made to Chapters 1 and 2 in order to include more recent developments in techniques (e.g. Schlenk-type, *cf* p. 10), and chromatographic methods and materials. Chapter 3 still remains the core of the book, and lists in alphabetical order relevant information on *ca* 4000 organic compounds. Chapter 4 gives a smaller listing of *ca* 750 inorganic and metal-organic substances, and makes a total increase of *ca* 13% of individual entries in these two chapters. Some additions have also been made to Chapter 5.

We are currently witnessing a major development in the use of physical methods for purifying large molecules and macromolecules, especially of biological origin. Considerable developments in molecular biology are apparent in techniques for the isolation and purification of key biochemicals and substances of high molecular weight. In many cases something approaching homogeneity has been achieved, as evidenced by electrophoresis, immunological and other independent criteria. We have consequently included a new section, Chapter 6, where we list upwards of 100 biological substances to illustrate their current methods of purification. In this chapter the details have been kept to a minimum, but the relevant references have been included.

The lists of individual entries in Chapters 3 and 4 range in length from single line entries to *ca* one page or more for solvents such as acetonitrile, benzene, ethanol and methanol. Some entries include information such as likely contaminants and storage conditions. More data referring to physical properties have been inserted for most entries [i.e. melting and boiling points, refractive indexes, densities, specific optical rotations (where applicable) and UV absorption data]. Inclusion of molecular weights should be useful when deciding on the quantities of reagents needed to carry out relevant synthetic reactions, or preparing analytical solutions. The Chemical Abstracts registry numbers have also been inserted for almost all entries, and should assist in the precise identification of the substances.

In the past ten years laboratory workers have become increasingly conscious of safety in the laboratory environment. We have therefore in three places in Chapter 1 (pp. 3 and 33, and bibliography p. 52) stressed more strongly the importance of safety in the laboratory. Also, where possible, in Chapters 3 and 4 we draw attention to the dangers involved with the manipulation of some hazardous substances.

The world wide facilities for retrieving chemical information provided by the Chemical Abstract Service (CAS on-line) have made it a relatively easy matter to obtain CAS registry numbers of substances, and most of the numbers in this monograph were obtained *via* CAS on-line. We should point out that two other available useful files are CSCHEM and CSCORP which provide, respectively, information on chemicals (and chemical products) and addresses and telephone numbers of the main branch offices of chemical suppliers.

The present edition has been produced on an IBM PC and a Laser Jet printer using the **Microsoft Word (4.0)** word-processing program with a set stylesheet. This has allowed the use of a variety of fonts and font sizes which has made the presentation more attractive than in the previous edition. Also, by altering the format and increasing slightly the sizes of the pages, the length of the monograph has been reduced from 568 to 391 pages. The reduction in the number of pages has been achieved in spite of the increase of *ca* 15% of total text.

We extend our gratitude to the readers whose suggestions have helped to improve the monograph, and to those who have told us of their experiences with some of the purifications stated in the previous editions, and in particular with the hazards that they have encountered. We are deeply indebted to Dr M.D. Fenn for the several hours that he has spent on the terminal to provide us with a large number of CAS registry numbers.

This monograph could not have been produced without the expert assistance of Mr David Clarke who has spent many hours to load the necessary fonts in the computer, and for advising one of the authors (W.L.F.A.) on how to use them together with the idiosyncrasies of Microsoft Word.

D.D.P. & W.L.F.A.
1988

Preface to the Fourth Edition

THE AIMS of the first three editions, to provide purification procedures of commercially available chemicals and biochemicals from published literature data, are continued in this fourth edition. Since the third edition in 1988 the number of new chemicals and biochemicals which have been added to most chemical and biochemical catalogues have increased enormously. Accordingly there is a need to increase the number of entries with more recent useful reagents and chemical and biochemical intermediates. With this in mind, together with the need to reorganise and update general purification procedures, particularly in the area of biological macromolecules, as well as the time lapse since the previous publication, this fourth edition of **Purification of Laboratory Chemicals** has been produced. Chapter 1 has been reorganised with some updating, and by using a smaller font it was kept to a reasonable number of pages. Chapters 2 and 5 were similarly altered and have been combined into one chapter. Eight hundred and three hundred and fifty entries have been added to Chapters 3 (25% increase) and 4 (44% increase) respectively, and four hundred entries (310% increase) were added to Chapter 5 (Chapter 6 in the Third Edition), making a total of 5700 entries; all resulting in an increase from 391 to 529 pages, i.e. by *ca* 35%.

Many references to the original literature have been included remembering that some of the best references happened to be in the older literature. Every effort has been made to provide the best references but this may not have been achieved in all cases. Standard abbreviations, listed on page 1, have been used throughout this edition to optimise space, except where no space advantage was achieved, in which cases the complete words have been written down to improve the flow of the sentences.

With the increasing facilities for information exchange, chemical, biochemical and equipment suppliers are making their catalogue information available on the Internet, e.g. Aldrich-Fluka-Sigma catalogue information is available on the World Wide Web by using the address <http://www.sigma.sial.com>, and GIBCO BRL catalogue information from <http://www.lifetech.com>, as well as on CD-ROMS which are regularly updated. Facility for enquiring about, ordering and paying for items is available *via* the Internet. CAS on-line can be accessed on the Internet, and CAS data is available now on CD-ROM. Also biosafety bill boards can similarly be obtained by sending SUBSCRIBE SAFETY John Doe at the address "listserv@uvmvm.uvm.edu", SUBSCRIBE BIOSAFETY at the address "listserv@mitvma.mit.edu", and SUBSCRIBE RADSAF at the address "listserv@romulus.ehs.uiuc.edu"; and the Occupational, Health and Safety information (Australia) is available at the address "<http://www.worksafe.gov.au/~wsa1>". Sigma-Aldrich provide Material Safety data sheets on CD-ROMs.

It is with much sadness that Dr Douglas D. Perrin was unable to participate in the preparation of the present edition due to illness. His contributions towards the previous editions have been substantial, and his drive and tenacity have been greatly missed.

The Third Edition was prepared on an IBM-PC and the previous IBM files were converted into Macintosh files. These have now been reformatted on a Macintosh LC575 computer and all further data to complete the Fourth Edition were added to these files. The text was printed with a Hewlett-Packard 4MV -600dpi Laser Jet printer which gives a clearer resolution.

I thank my wife Dr Pauline M. Armarego, also an organic chemist, for the arduous and painstaking task of entering the new data into the respective files, and for the numerous hours of proofreading as well as the corrections of typographic errors in the files. I should be grateful to my readers for any comments, suggestions, amendments and criticisms which could, perhaps, be inserted in the second printing of this edition.

W.L.F. Armarego
30 June 1996

Contents

<i>Preface to the Fifth Edition</i>	xi
<i>Preface to the First Edition</i>	xiii
<i>Preface to the Second Edition</i>	xiii
<i>Preface to the Third Edition</i>	xiv
<i>Preface to the Fourth Edition</i>	xv
1. Common Physical Techniques Used in Purification	1
Introduction	1
The Question of Purity	1
Sources of Impurities	2
Practices to Avoid Impurities	3
Cleaning Practices	3
Silylation of Glassware and Plasticware	3
Safety Precautions Associated with the Purification of Laboratory Chemicals	4
Some Hazards of Chemical Manipulation in Purification and Recovery of Residues	4
Perchlorates and Perchloric Acid	5
Peroxides	5
Heavy-Metal-Containing-Explosives	5
Strong Acids	5
Reactive Halides and Anhydrides	5
Solvents	5
Salts	6
Safety Disclaimer	6
Methods of Purification of Reagents and Solvents	6
Solvent Extraction and Distribution	6
Ionization Constants and pK	7
pK and Temperature	8
pK and Solvent	8
Distillation	8
Techniques	9

Distillation of Liquid Mixtures	9
Types of Distillation	10
The Distilling Flask	10
Vacuum Distillation	11
Spinning-Band Distillation	12
Steam Distillation	12
Azeotropic Distillation	13
Kügelrohr Distillation	13
Isopiestic or Isothermal Distillation	13
Recrystallisation	14
Techniques	14
Filtration	14
Choice of Solvents	15
Petroleum Ethers	15
Mixed Solvents	16
Recrystallisation from the Melt	16
Zone Refining	16
Sublimation	17
Chromatography	17
Vapour Phase Chromatography (GC or Gas-Liquid Chromatography)	17
Liquid Chromatography	18
Adsorption Chromatography	18
Graded Adsorbents and Solvents	19
Preparation and Standardization of Alumina	19
Preparation of Other Adsorbents	20
Flash Chromatography	21
Paired-Ion Chromatography	21
Ion-Exchange Chromatography	21
Ion-Exchange Resins	21
Ion-Exchange Celluloses and Sephadex	22
Gel Filtration	24
High Performance Liquid Chromatography (HPLC)	24
Other Types of Liquid Chromatography	25
Drying	25
Removal of Solvents	25
Removal of Water	26
Intensity and Capacity of Common Desiccants	26
Suitability of Individual Desiccants	27
Molecular Sieves	28

Miscellaneous Techniques	29
Freeze-Pump-Thaw and Purging	29
Vacuum-Lines, Schlenk and Glovebox Techniques	30
Abbreviations	30
Tables	30
Table 1. Some Common Immiscible or Slightly Miscible Pairs of Solvents	30
Table 2. Aqueous Buffers	31
Table 3A. Predicted Effect of Pressure on Boiling Point	32
Table 3B. Predicted Effect of Pressure on Boiling Point	33
Figure 1. Nomogram	34
Table 4. Heating Baths	35
Table 5. Whatman Filter Papers	35
Table 6. Micro Filters	36
Table 7. Common Solvents Used in Recrystallisation	37
Table 8. Pairs of Miscible Solvents	37
Table 9. Materials for Cooling Baths	38
Table 10. Liquids for Stationary Phases in Gas Chromatography	39
Table 11. Methods of Visualization of TLC Spots	39
Table 12. Graded Adsorbents and Solvents	40
Table 13. Representative Ion-Exchange Resins	40
Table 14. Modified Fibrous Celluloses for Ion-Exchange	40
Table 15. Bead Form Ion-Exchange Packagings	41
Table 16. Liquids for Drying Pistols	41
Table 17. Vapour Pressures (mm Hg) of Saturated Aqueous Solutions in Equilibrium with Solid Salts	42
Table 18. Drying Agents for Classes of Compounds	43
Table 19. Static Drying for Selected Liquids (25°C)	43
Table 20. Boiling Points of Some Useful Gases at 760 mm	44
Table 21. Solubilities of HCl and NH ₃ at 760 mm (g/100g of Solution)	44
Table 22. Prefixes for Quantities	44
Bibliography	45
2. Chemical Methods Used in Purification	53
General Remarks	53
Removal of Traces of Metals from Reagents	53
Metal Impurities	53
Distillation	53
Use of Ion Exchange Resins	54

Precipitation	54
Removal of Lead Contaminants	54
Removal of Iron Contaminants	54
Removal of Other Metal Contaminants	54
Extraction	54
Complexation	55
Use of Metal Hydrides	55
Lithium Aluminium Hydride	55
Calcium Hydride	55
Sodium Borohydride	55
Potassium Borohydride	56
Purification via Derivatives	56
Alcohols	56
Aldehydes	57
Amines	57
Picrates	57
Salts	57
Double Salts	58
<i>N</i> -Acetyl Derivatives	58
<i>N</i> -Tosyl Derivatives	58
Aromatic Hydrocarbons	58
Adducts	58
Sulfonation	58
Carboxylic Acids	58
4-Bromophenacyl Esters	58
Alkyl Esters	58
Salts	59
Hydroperoxides	59
Ketones	59
Bisulfite Adduct	59
Semicarbazones	59
Phenols	59
Benzoates	59
Acetates	59
Phosphate and Phosphonate Esters	60
Miscellaneous	60
General Methods for the Purification of Classes of Compounds	60
Procedures	60
Criteria of Purity	61

General Procedures for the Purification of Some Classes of Organic Compounds	61
Acetals	61
Acids	61
Carboxylic Acids	61
Sulfonic Acids	62
Sulfinic Acids	62
Acid Chlorides	62
Alcohols	62
Monohydric Alcohols	62
Polyhydric Alcohols	63
Aldehydes	63
Amides	63
Amines	63
Amino Acids	64
Anhydrides	64
Carotenoids	64
Esters	64
Ethers	65
Halides	65
Hydrocarbons	66
Imides	67
Imino Compounds	67
Ketones	67
Macromolecules	67
Nitriles	67
Nitro Compounds	67
Nucleic Acids	68
Phenols	68
Polypeptides and Proteins	68
Quinones	68
Salts (Organic)	68
With Metal Ions	68
With Organic Cations	68
With Sodium Alkane Sulfonates	68
Sulfur Compounds	68
Disulfides	68
Sulfones	68
Sulfoxides	69
Thioethers	69

Thiols	69
Thiolsulfonates (disulfoxides)	69
Bibliography	70
3. The Future of Purification	72
Introduction	72
Solid Phase Synthesis	72
Solid Phase Peptide Synthesis (SPPS)	73
Solid Phase Deoxyribonucleotide Synthesis	73
Solid Phase Oligosaccharide Synthesis	73
Solid Phase Organic Synthesis (SPOS)	73
Polymer Supported React Ants	74
Scavenger Resins	74
Resin Support	74
Choice of Resin for SPOS	74
Combinatorial Chemistry	75
Monitoring Solid Phase Reactions	75
Infrared Analysis of Resin	75
Qualitative and Quantitative Analyses	75
Detection of Reactive Groups on Resins	76
Detection of Hydroxy Groups on Resin	76
Detection of Aldehyde Groups on Resin	76
Detection of Carboxy Groups on Resin	76
Detection of Amino Groups on Resin	76
Detection of Thiol Groups on Resin	76
Fmoc Assay	76
Ionic Liquids	77
Fluorous Chemistry	77
Bibliography	78
4. Purification of Organic Chemicals	80
Abietic acid ... Allantoin	81
Allene ... Azure C	100
B.A.L. ... Bis-(β -chloroethyl)amine hydrochloride	118
Bis-(β -chloroethyl) ether ... Butyryl chloride	134
Cacotheline ... 3-(4-Chlorophenyl)-1,1-dimethylurea	152
4-Chloro-1,2-phenylenediamine ... Cytosine	167
<i>cis</i> -Decahydroisoquinoline ... Diethyl ketone (3-pentanone)	184

Diethyl malonate ... Duroquinone	205
α -Ecdyson ... Furoin	229
Galactaric acid ... Itaconic anhydride	250
Janus green B ... β -D-Lyxose	275
Malachite green ... Myristic acid	280
Naphthacene ... Oxine blue	304
Palmitic acid anhydride ... Phenyltoloxamine	319
Phenyl 4-toluenesulfonate ... Pyruvic acid	333
<i>p</i> -Quaterphenyl ... Syringic acid	346
D(-)-Tagatose ... Thioguanosine	355
Thioindigo ... L-Tyrosine	368
Umbelliferone ... Zeaxanthin	383
5. Purification of Inorganic and Metalorganic Chemicals (Including Organic Compounds of B, Bi, P, Se, Si, and Ammonium and Metal Salts of Organic Acids)	389
Acetarsol ... <i>n</i> -Butylstannoic acid	389
Cacodylic acid ... 1,3-Divinyl-1,1,3,3-tetramethyldisiloxane	405
Eosin B ... Monoperoxyphthalic acid magnesium salt	422
Naphthalene Scarlet Red 4R ... Ruthenocene	443
Samarium (II) iodide ... Sulfuryl chloride	461
Tantalium (V) chloride ... Zirconyl chloride	479
6. Purification of Biochemicals and Related Products	500
Abrin A and Abrin B ... S-Butyryl thiocholine iodide	505
L-Canavanine sulfate ... Fuschin	518
D-Galactal ... Oxitocin	536
Palmitoyl coenzyme A ... Rifamycin SV sodium salt	555
Saccharides ... Zeatin	566
General Subject Index	578
CAS Registry Numbers Index	585
50-00-0 ... 392-56-3	585
404-86-4 ... 3022-16-0	593
3024-83-7 ... 336096-71-0	601

CHAPTER 1

COMMON PHYSICAL TECHNIQUES USED IN PURIFICATION

INTRODUCTION

Purity is a matter of degree. Other than adventitious contaminants such as dust, paper fibres, wax, cork, etc., that may have been incorporated into the sample during manufacture, all commercially available chemical substances are in some measure impure. Any amounts of unreacted starting material, intermediates, by-products, isomers and related compounds may be present depending on the synthetic or isolation procedures used for preparing the substances. Inorganic reagents may deteriorate because of defective packaging (glued liners affected by sulfuric acid, zinc extracted from white rubber stoppers by ammonia), corrosion or prolonged storage. Organic molecules may undergo changes on storage. In extreme cases the container may be incorrectly labelled or, where compositions are given, they may be misleading or inaccurate for the proposed use. Where any doubt exists it is usual to check for impurities by appropriate spot tests, or by recourse to tables of physical or spectral properties such as the extensive infrared and NMR libraries published by the Sigma Aldrich Chemical Co.

The important question, then, is not whether a substance is pure but whether a given sample is sufficiently pure for some intended purpose. That is, are the contaminants likely to interfere in the process or measurement that is to be studied. By suitable manipulation it is often possible to reduce levels of impurities to acceptable limits, but absolute purity is an ideal which, no matter how closely approached, can never be attained. A *negative* physical or chemical test indicates only that the amount of an impurity in a substance lies below a certain sensitivity level; no test can demonstrate that a specified impurity is entirely absent.

When setting out to purify a laboratory chemical, it is desirable that the starting material is of the best grade commercially available. Particularly among organic solvents there is a range of qualities varying from *laboratory chemical* to *spectroscopic* and *chromatographic* grades. Many of these are suitable for use as received. With many of the more common reagents it is possible to obtain from the current literature some indications of likely impurities, their probable concentrations and methods for detecting them. However, in many cases complete analyses are not given so that significant concentrations of unspecified impurities may be present.

THE QUESTION OF PURITY

Solvents and substances that are specified as *pure* for a particular purpose may, in fact, be quite impure for other uses. Absolute ethanol may contain traces of benzene, which makes it unsuitable for ultraviolet spectroscopy, or plasticizers which make it unsuitable for use in solvent extraction.

Irrespective of the grade of material to be purified, it is essential that some criteria exist for assessing the degree of purity of the final product. The more common of these include:

1. Examination of physical properties such as:
 - (a) Melting point, freezing point, boiling point, and the freezing curve (i.e. the variation, with time, in the freezing point of a substance that is being slowly and continuously frozen).
 - (b) Density.
 - (c) Refractive index at a specified temperature and wavelength. The sodium D line at 589.26 nm (weighted mean of D₁ and D₂ lines) is the usual standard of wavelength but results from other wavelengths can often be interpolated from a plot of refractive index versus $1/(\text{wavelength})^2$.

- (d) Specific conductivity. (This can be used to detect, for example, water, salts, inorganic and organic acids and bases, in non-electrolytes).
 - (e) Optical rotation, optical rotatory dispersion and circular dichroism.
2. Empirical analysis, for C, H, N, ash, etc.
 3. Chemical tests for particular types of impurities, e.g. for peroxides in aliphatic ethers (with acidified KI), or for water in solvents (quantitatively by the Karl Fischer method, see Fieser and Fieser, *Reagents for Organic Synthesis* J. Wiley & Sons, NY, Vol 1 pp. 353, **528**, 1967, Library of Congress Catalog Card No 66-27894).
 4. Physical tests for particular types of impurities:
 - (a) Emission and atomic absorption spectroscopy for detecting organic impurities and determining metal ions.
 - (b) Chromatography, including paper, thin layer, liquid (high, medium and normal pressure) and vapour phase.
 - (c) Electron spin resonance for detecting free radicals.
 - (d) X-ray spectroscopy.
 - (e) Mass spectroscopy.
 - (f) Fluorimetry.
 5. Examination of spectroscopic properties
 - (a) Nuclear Magnetic Resonance (^1H , ^{13}C , ^{31}P , ^{19}F NMR etc)
 - (b) Infrared spectroscopy (IR)
 - (c) Ultraviolet spectroscopy (UV)
 - (d) Mass spectroscopy [electron ionisation (EI), electron ionisation (CI), electrospray ionisation (ESI), fast atom bombardment (FAB), matrix-associated laser desorption ionisation (MALDI), etc]
 6. Electrochemical methods (see Chapter 6 for macromolecules).
 7. Nuclear methods which include a variety of radioactive elements as in organic reagents, complexes or salts.

A substance is usually taken to be of an acceptable purity when the measured property is unchanged by further treatment (especially if it agrees with a recorded value). In general, at least two different methods, such as recrystallisation and distillation, should be used in order to ensure maximum purity. Crystallisation may be repeated (from the same solvent or better from different solvents) until the substance has a constant melting point or absorption spectrum, and until it distils repeatedly within a narrow, specified temperature range.

With liquids, the refractive index at a specified temperature and wavelength is a sensitive test of purity. Note however that this is sensitive to dissolved gases such as O_2 , N_2 or CO_2 . Under favourable conditions, freezing curve studies are sensitive to impurity levels of as little as 0.001 moles per cent. Analogous fusion curves or heat capacity measurements can be up to ten times as sensitive as this. With these exceptions, most of the above methods are rather insensitive, especially if the impurities and the substances in which they occur are chemically similar. In some cases, even an impurity comprising many parts per million of a sample may escape detection.

The common methods of purification, discussed below, comprise distillation (including fractional distillation, distillation under reduced pressure, sublimation and steam distillation), crystallisation, extraction, chromatographic and other methods. In some cases, volatile and other impurities can be removed simply by heating. Impurities can also sometimes be eliminated by the formation of derivatives from which the purified material is regenerated (see Chapter 2).

SOURCES OF IMPURITIES

Some of the more obvious sources of contamination of solvents arise from storage in metal drums and plastic containers, and from contact with grease and screw caps. Many solvents contain water. Others have traces of acidic materials such as hydrochloric acid in chloroform. In both cases this leads to corrosion of the drum and contamination of the solvent by traces of metal ions, especially Fe^{3+} . Grease, for example on stopcocks of separating funnels and other apparatus, e.g. greased ground joints, is also likely to contaminate solvents during extractions and chemical manipulation.

A much more general source of contamination that has not received the consideration it merits comes from the use of plastics for tubing and containers. Plasticisers can readily be extracted by organic solvents from PVC and other plastics, so that most solvents, irrespective of their grade (including spectrograde and ultrapure) have been reported to contain 0.1 to 5ppm of plasticiser [de Zeeuw, Jonkman and van Mansvelt *Anal Biochem* **67** 339 1975]. Where large quantities of solvent are used for extraction (particularly of small amounts of compounds), followed by evaporation, this can introduce significant amounts of impurity, even exceeding the weight of the genuine extract and giving rise to spurious peaks in gas chromatography (for example of fatty acid methyl esters [Pascaud, *Anal Biochem* **18** 570 1967]). Likely contaminants are di(2-ethylhexyl)phthalate and dibutyl phthalate, but upwards of 20 different phthalate esters are listed as plasticisers as well as adipates, azelates, phosphates, epoxides, polyesters and various heterocyclic compounds. These plasticisers would enter the solvent during passage through plastic tubing or from storage in containers or from plastic coatings used in cap liners for bottles. Such contamination could arise at any point in the manufacture or distribution of a solvent. The problem with cap liners is avoidable by using corks wrapped in aluminium foil, although even in this case care should be taken because aluminium foil can dissolve in some liquids e.g. benzylamine and propionic acid.

Solutions in contact with polyvinyl chloride can become contaminated with trace amounts of lead, titanium, tin, zinc, iron, magnesium or cadmium from additives used in the manufacture and moulding of PVC.

N-Phenyl-2-naphthylamine is a contaminant of solvents and biological materials that have been in contact with black rubber or neoprene (in which it is used as an antioxidant). Although it was only an artefact of the separation procedure it has been isolated as an apparent component of vitamin K preparations, extracts of plant lipids, algae, livers, butter, eye tissue and kidney tissue [Brown *Chem Br* **3** 524 1967].

Most of the above impurities can be removed by prior distillation of the solvent, but care should be taken to avoid plastic or black rubber as much as possible.

PRACTICES TO AVOID IMPURITIES

Cleaning practices

Laboratory glassware and Teflon equipment can be cleaned satisfactorily for most purposes by careful immersion into a solution of sodium dichromate in concentrated sulfuric acid, followed by draining, and rinsing copiously with distilled water. This is an exothermic reaction and should be carried out **very** cautiously in an efficient fume cupboard. [To prepare the chromic acid bath, dissolve 5 g of sodium dichromate (CARE: cancer suspect agent) in 5 mL of water. The dichromate solution is then cooled and stirred while 100 mL of concentrated sulfuric acid is added slowly. Store in a glass bottle.] Where traces of chromium (adsorbed on the glass) must be avoided, a 1:1 mixture of concentrated sulfuric and nitric acid is a useful alternative. (*Use in a fumehood to remove vapour and with adequate face protection.*) Acid washing is also suitable for polyethylene ware but prolonged contact (some weeks) leads to severe deterioration of the plastic. Alternatively an alcoholic solution of sodium hydroxide (alkaline base bath) can be used. This strongly corrosive solution (CAUTION: Alkali causes serious burns) can be made by dissolving 120g of NaOH in 120 mL water, followed by dilution to 1 L with 95% ethanol. This solution is conveniently stored in suitable alkali resistant containers (e.g. Nalgene heavy duty rectangular tanks) with lids. Glassware can be soaked overnight in the base bath and rinsed thoroughly after soaking. For much glassware, washing with hot detergent solution, using tap water, followed by rinsing with distilled water and acetone, and heating to 200-300° overnight, is adequate. (Volumetric apparatus should not be heated: after washing it is rinsed with acetone, then hexane, and air-dried. Prior to use, equipment can be rinsed with acetone, then with petroleum ether or hexane, to remove the last traces of contaminants.) Teflon equipment should be soaked, first in acetone, then in petroleum ether or hexane for ten minutes prior to use.

For trace metal analyses, prolonged soaking of equipment in 1M nitric acid may be needed to remove adsorbed metal ions.

Soxhlet thimbles and filter papers may contain traces of lipid-like materials. For manipulations with highly pure materials, as in trace-pesticide analysis, thimbles and filter papers should be thoroughly extracted with hexane before use.

Trace impurities in silica gel for TLC can be removed by heating at 300° for 16h or by Soxhlet extraction for 3h with distilled chloroform, followed by 4h extraction with distilled hexane.

Silylation of glassware and plasticware

Silylation of apparatus makes it repellent to water and hydrophilic materials. It minimises loss of solute by adsorption onto the walls of the container. The glassware is placed in a desiccator containing dichloromethyl silane (1mL) in a small beaker and evacuated for 5min. The vacuum is turned off and air is introduced into the desiccator which allows the silylating agent to coat the glassware uniformly. The desiccator is then evacuated, closed and set aside for 2h. The glassware is removed from the desiccator and baked at 180° for 2h before use.

Plasticware is treated similarly except that it is rinsed well with water before use instead of baking. Note that dichloromethyl silane is highly **TOXIC** and **VOLATILE**, and the whole operation should be carried out in an efficient fume cupboard.

An alternative procedure used for large apparatus is to rinse the apparatus with a 5% solution of dichloromethyl silane in chloroform, followed by several rinses with water before baking the apparatus at 180°/2h (for glass) or drying in air (for plasticware).

Plus One REPEL-SILANE ES (a solution of 2% w/v of dichloromethyl silane in octamethyl cyclooctasilane) is used to inhibit the sticking of polyacrylamide gels, agarose gels and nucleic acids to glass surfaces and is available commercially (Amersham Biosciences).

SAFETY PRECAUTIONS ASSOCIATED WITH THE PURIFICATION OF LABORATORY CHEMICALS

Although most of the manipulations involved in purifying laboratory chemicals are inherently safe, care is necessary if hazards are to be avoided in the chemical laboratory. In particular there are dangers inherent in the inhalation of vapours and absorption of liquids and low melting solids through the skin. In addition to the toxicity of solvents there is also the risk of their flammability and the possibility of eye damage. Chemicals, particularly in admixture, may be explosive. Compounds may be carcinogenic or otherwise deleterious to health. Present day chemical catalogues specifically indicate the particular dangerous properties of the individual chemicals they list and these should be consulted whenever the use of commercially available chemicals is contemplated. Radioisotopic labelled compounds pose special problems of human exposure and of disposal of laboratory waste. Hazardous purchased chemicals are accompanied by detailed MSDS (Material Safety Data Sheets), which contain information regarding their toxicity, safety handling procedures and the necessary precautions to be taken. These should be read carefully and filed for future reference. In addition, chemical management systems such as ChemWatch which include information on hazards, handling and storage are commercially available. There are a number of websites which provide selected safety information: they include the Sigma-Aldrich website (www.sigmaaldrich.com) and other chemical websites e.g. www.ilpi.com/msds).

The most common hazards are:

- (1) Explosions due to the presence of peroxides formed by aerial oxidation of ethers and tetrahydrofuran, decahydronaphthalene, acrylonitrile, styrene and related compounds.
- (2) Compounds with low flash points (below room temperature). Examples are acetaldehyde, acetone, acetonitrile, benzene, carbon disulfide, cyclohexane, diethyl ether, ethyl acetate and *n*-hexane.
- (3) Contact of oxidising agents (KMnO₄, HClO₄, chromic acid) with organic liquids.
- (4) Toxic reactions with tissues.

The laboratory should at least be well ventilated and safety glasses should be worn, particularly during distillation and manipulations carried out under reduced pressure or elevated temperatures. With this in mind we have endeavoured to warn users of this book whenever greater than usual care is needed in handling chemicals. As a general rule, however, **all chemicals which users are unfamiliar with should be treated with extreme care and assumed to be highly flammable and toxic.** The safety of others in a laboratory should always be foremost in mind, with ample warning whenever a potentially hazardous operation is in progress. Also, unwanted solutions or solvents should never be disposed of *via* the laboratory sink. The operator should be aware of the usual means for disposal of chemicals in her/his laboratories and she/he should remove unwanted chemicals accordingly. **Organic liquids for disposal should be temporarily stored, as is practically possible, in respective containers. Avoid placing all organic liquids in the same container particularly if they contain small amounts of reagents which could react with each other. Halogenated waste solvents should be kept separate from other organic liquids.**

SOME HAZARDS OF CHEMICAL MANIPULATION IN PURIFICATION AND RECOVERY OF RESIDUES

Performing chemical manipulations calls for some practical knowledge if danger is to be avoided. However, with care, hazards can be kept to an acceptable minimum. A good general approach is to consider every operation as potentially perilous and then to adjust one's attitude as the operation proceeds. A few of the most common dangers are set out below. For a larger coverage of the following sections, and of the literature, the bibliography at the end of this chapter should be consulted.

Perchlorates and perchloric acid. At 160° perchloric acid is an exceedingly strong oxidising acid and a strong dehydrating agent. Organic perchlorates, such as methyl and ethyl perchlorates, are unstable and are violently explosive compounds. A number of heavy-metal perchlorates are extremely prone to explode. The use of anhydrous magnesium perchlorate, *Anhydrone*, *Dehydrite*, as a drying agent for organic vapours is **not** recommended. Desiccators which contain this drying agent should be adequately shielded at all times and kept in a cool place, i.e. **never** on a window sill where sunlight can fall on it.

No attempt should be made to purify perchlorates, except for ammonium, alkali metal and alkaline earth salts which, in water or aqueous alcoholic solutions are insensitive to heat or shock. Note that perchlorates react relatively slowly in aqueous organic solvents, but as the water is removed there is an increased possibility of an explosion. Perchlorates, often used in non-aqueous solvents, are explosive in the presence of even small amounts of organic compounds when heated. Hence stringent care should be taken when purifying perchlorates, and direct flame and infrared lamps should be avoided. Tetra-alkylammonium perchlorates should be dried below 50° under vacuum (and protection). Only very small amounts of such materials should be prepared, and stored, at any one time.

Peroxides. These are formed by aerial oxidation or by autoxidation of a wide range of organic compounds, including diethyl ether, allyl ethyl ether, allyl phenyl ether, dibenzyl ether, benzyl butyl ether, *n*-butyl ether, *iso*-butyl ether, *t*-butyl ether, dioxane, tetrahydrofuran, olefins, and aromatic and saturated aliphatic hydrocarbons. They accumulate during distillation and can detonate violently on evaporation or distillation when their concentration becomes high. If peroxides are likely to be present materials should be tested for peroxides before distillation (for tests see entry under "Ethers", in Chapter 2). Also, distillation should be discontinued when at least one quarter of the residue is left in the distilling flask.

Heavy-metal-containing-explosives. Ammoniacal silver nitrate, on storage or treatment, will eventually deposit the highly explosive silver nitride "*fulminating silver*". Silver nitrate and ethanol may give silver fulminate (see Chapter 5), and in contact with azides or hydrazine and hydrazides may form silver azide. Mercury can also form such compounds. Similarly, ammonia or ammonium ions can react with gold salts to form "*fulminating gold*". Metal fulminates of cadmium, copper, mercury and thallium are powerfully explosive, and some are detonators [Luchs, *Photog Sci Eng* 10 334 1966]. Heavy metal containing solutions, particularly when organic material is present should be treated with great respect and precautions towards possible explosion should be taken.

Strong acids. In addition to perchloric acid (see above), extra care should be taken when using strong mineral acids. Although the effects of concentrated sulfuric acid are well known these cannot be stressed strongly enough. Contact with tissues will leave irreparable damage. **Always dilute the concentrated acid by carefully adding the acid down the side of the flask which contains water, and the process should be carried out under cooling. This solution is not safe to handle until the acid has been thoroughly mixed with the water. Protective face, and body coverage should be used at all times.** Fuming sulfuric acid and chlorosulfonic acid are even more dangerous than concentrated sulfuric acid and adequate precautions should be taken. Chromic acid cleaning mixture contains strong sulfuric acid and should be treated in the same way; and in addition the mixture is potentially *carcinogenic*.

Concentrated and fuming nitric acids are also dangerous because of their severe deleterious effects on tissues.

Reactive halides and anhydrides. Substances like acid chlorides, low molecular weight anhydrides and some inorganic halides (e.g. PCl_3) can be **highly toxic and lachrymatory affecting mucous membranes and lung tissues. Utmost care should be taken when working with these materials. Work should be carried out in a very efficient fume cupboard.**

Solvents. The flammability of low-boiling organic liquids cannot be emphasised strongly enough. These invariably have very low flash points and can ignite spontaneously. Special precautions against explosive flammability should be taken when recovering such liquids. Care should be taken with small volumes (*ca* 250mL) as well as large volumes (> 1L), and the location of all the fire extinguishers, and fire blankets, in the immediate vicinity of the apparatus should be checked. The fire extinguisher should be operational. The following flammable liquids (in alphabetical order) are common fire hazards in the laboratory: acetaldehyde, acetone, acrylonitrile, acetonitrile, benzene, carbon disulfide, cyclohexane, diethyl ether, ethyl acetate, hexane, low-boiling petroleum ether, tetrahydrofuran and toluene. Toluene should always be used in place of benzene wherever possible due to the potential *carcinogenic* effects of the liquid and vapour of the latter.

The drying of flammable solvents with sodium or potassium metal and metal hydrides poses serious potential fire hazards and adequate precautions should be stressed.

Salts. In addition to the dangers of perchlorate salts, other salts such as nitrates, azides and diazo salts can be hazardous and due care should be taken when these are dried. Large quantities should never be prepared or stored for long periods.

SAFETY DISCLAIMER

Experimental chemistry is a very dangerous occupation and extreme care and adequate safety precautions should be taken at all times. Although we have stated the safety measures that have to be taken under specific entries these are by no means exhaustive and some may have been unknowingly or accidentally omitted. The experimenter without prior knowledge or experience must seek further safety advice on reagents and procedures from experts in the field before undertaking the purification of any material. We take no responsibility whatsoever if any mishaps occur when using any of the procedures described in this book.

METHODS OF PURIFICATION OF REAGENTS AND SOLVENTS

Many methods exist for the purification of reagents and solvents. A number of these methods are routinely used in synthetic as well as analytical chemistry and biochemistry. These techniques, outlined below, will be discussed in greater detail in the respective sections in this Chapter. It is important to note that more than one method of purification may need to be implemented in order to obtain compounds of highest purity.

Common methods of purification are:

- (a) Solvent Extraction and Distribution
- (b) Distillation
- (c) Recrystallisation
- (d) Sublimation
- (e) Chromatography

For substances contaminated with water or solvents, drying with appropriate absorbents and desiccants may be sufficient.

SOLVENT EXTRACTION AND DISTRIBUTION

Extraction of a substance from suspension or solution into another solvent can sometimes be used as a purification process. Thus, organic substances can often be separated from inorganic impurities by shaking an aqueous solution or suspension with suitable immiscible solvents such as benzene, carbon tetrachloride, chloroform, diethyl ether, diisopropyl ether or petroleum ether. After several such extractions the combined organic phase is dried and the solvent is evaporated. Grease from the glass taps of conventional separating funnels is invariably soluble in the solvents used. Contamination with grease can be very troublesome particularly when the amounts of material to be extracted are very small. Instead, the glass taps should be lubricated with the extraction solvent; or better, the taps of the extraction funnels should be made of the more expensive material *Teflon*. Immiscible solvents suitable for extractions are given in Table 1. Addition of electrolytes (such as ammonium sulfate, calcium chloride or sodium chloride) to the aqueous phase helps to ensure that the organic layer separates cleanly and also decreases the extent of extraction into the latter. Emulsions can also be broken up by filtration (with suction) through Celite, or by adding a little octyl alcohol or some other paraffinic alcohol. The main factor in selecting a suitable immiscible solvent is to find one in which the material to be extracted is readily soluble, whereas the substance from which it is being extracted is not. The same considerations apply irrespective of whether it is the substance being purified, or one of its contaminants, that is taken into the new phase. (The second of these processes is described as washing.)

Common examples of washing with aqueous solutions include the following:

Removal of acids from water-immiscible solvents by washing with aqueous alkali, sodium carbonate or sodium bicarbonate.

Removal of phenols from similar solutions by washing with aqueous alkali.

Removal of organic bases by washing with dilute hydrochloric or sulfuric acids.

Removal of unsaturated hydrocarbons, of alcohols and of ethers from saturated hydrocarbons or alkyl halides by washing with cold concentrated sulfuric acid.

This process can also be applied to purification of the substance if it is an acid, a phenol or a base, by extracting into the appropriate aqueous solution to form the salt which, after washing with pure solvent, is again converted to

the free species and re-extracted. Paraffin hydrocarbons can be purified by extracting them with phenol (in which aromatic hydrocarbons are highly soluble) prior to fractional distillation.

For extraction of solid materials with a solvent, a *Soxhlet* extractor is commonly used. This technique is applied, for example, in the alcohol extraction of dyes to free them from insoluble contaminants such as sodium chloride or sodium sulfate.

Acids, bases and amphoteric substances can be purified by taking advantage of their ionisation constants.

Ionisation constants and pK.

When substances ionise their neutral species produce positive and negative species. The ionisation constants are those constant values (equilibrium constants) for the equilibria between the charged species and the neutral species, or species with a larger number of charges (e.g. between mono and dications). These ionisation constants are given as **pK** values where $\text{pK} = -\log K$ and K is the dissociation constant for the equilibrium between the species [Albert and Serjeant *The Determination of Ionisation Constants, A Laboratory Manual*, 3rd Edition, Chapman & Hall, New York, London, 1984, ISBN 0412242907].

The advantage of using pK values (instead of K values) is that theory (and practice) states that the pK values of ionisable substances are numerically equal to the pH of the solution at which the concentrations of ionised and neutral species are equal. For example acetic acid has a pK^{25} value of 4.76 at 25° in H₂O, then at pH 4.76 the aqueous solution contains equal amounts of acetic acid [AcOH] and acetate anion [AcO⁻], i.e. [AcOH]/[AcO⁻] of 50/50. At pH 5.76 (pK + 1) the solution contains [AcOH]/[AcO⁻] of 10/90, at pH 6.76 (pK + 2) the solution contains [AcOH]/[AcO⁻] of 1/99 etc; conversely at pH 3.76 (pK - 1) the solution contains [AcOH]/[AcO⁻] of 90/10, and at pH 2.76 (pK - 2) the solution contains [AcOH]/[AcO⁻] of 99/1.

One can readily appreciate the usefulness of pK value in purification procedures, e.g. as when purifying acetic acid. If acetic acid is placed in aqueous solution and the pH adjusted to 7.76 {[AcOH]/[AcO⁻] with a ratio of 0.1/99.9}, and extracted with say diethyl ether, neutral impurities will be extracted into diethyl ether leaving almost all the acetic acid in the form of AcO⁻ in the aqueous solution. If then the pH of the solution is adjusted to 1.67 where the acid is almost all in the form AcOH, almost all of it will be extracted into diethyl ether.

Aniline will be used as a second example. It has a pK^{25} of 4.60 at 25° in H₂O. If it is placed in aqueous solution at pH 1.60 it will exist almost completely (99.9%) as the anilinium cation. This solution can then be extracted with solvents e.g. diethyl ether to remove neutral impurities. The pH of the solution is then adjusted to 7.60 whereby aniline will exist as the free base (99.9%) and can be extracted into diethyl ether in order to give purer aniline.

See Table 2 for the pH values of selected buffers.

A knowledge of the pK allows the adjustment of the pH without the need of large excesses of acids or base. In the case of inorganic compounds a knowledge of the pK is useful for adjusting the ionic species for making metal complexes which could be masked or extracted into organic solvents [Perrin and Dempsey *Buffers for pH and Metal Ion Control*, Chapman & Hall, New York, London, 1974, ISBN 0412117002], or for obtaining specific anionic species in solution e.g. H₂PO₄⁻, HPO₄²⁻ or PO₄³⁻.

The **pK** values that have been entered in Chapters 4, 5 and 6 have been collected directly from the literature or from compilations of literature values for organic bases [Perrin *Dissociation Constants of Organic Bases in Aqueous Solution*, Butterworths, London, 1965, Supplement 1972, ISBN 040870408X; Albert and Serjeant *The Determination of Ionisation Constants, A Laboratory Manual*, 3rd Edition, Chapman & Hall, London, New York, 1984, ISBN 0412242907]; organic acids [Kortum, Vogel and Andrussov, *Dissociation Constants of Organic Acids in Aqueous Solution*, Butterworth, London, 1961; Serjeant and Dempsey, *Dissociation Constants of Organic Acids in Aqueous Solution*, Pergamon Press, Oxford, New York, 1979, ISBN 0080223397; and inorganic acids and bases [Perrin, *Ionisation Constants of Inorganic Acids and Bases in Aqueous Solution*, Second Edition, Pergamon Press, Oxford, New York, 1982, ISBN 0080292143]. Where literature values were not available, values have been predicted and assigned $\text{pK}_{\text{Est}} \sim$. Most predictions should be so close to true values as to make very small difference for the purposes intended in this book. The success of the predictions, i.e. how close to the true value, depends on the availability of pK values for closely related compounds because the effect of substituents or changes in structures are generally additive [Perrin, Dempsey and Serjeant, *pKa Prediction for Organic Acids and Bases*, Chapman & Hall, London, New York, 1981, ISBN 04122190X].

All the pK values in this book are pKa values, the acidic pK, i.e. dissociation of H⁺ from an acid (AH) or from a conjugate base (BH⁺). Occasionally pKb values are reported in the literature but these can be converted using the equation $pK_a + pK_b = 14$. For strong acids e.g. sulfuric acid, and strong bases, e.g. sodium hydroxide, the pK values lie beyond the 1 to 11 scale and have to be measured in strong acidic and basic media. In these cases appropriate scales e.g. the H₀ (for acids) and H_L (for bases) have been used [see Katritzky and Waring *J Chem Soc* 1540 1962]. These values will be less than 1 (and negative) for acids and >11 for bases. They are a rough guides to the strengths of acids and bases. Errors in the stated pK and pK_{Est} ~ values can be judged from the numerical values given. Thus pK values of 4.55, 4.5 and 4 mean that the respective errors are better than ± 0.05, ± 0.3 and ± 0.5. Values taken from the literature are written as pK, and all the values that were estimated because they were not found in the literature are written as pK_{Est}.

pK and Temperature.

The temperatures at which the literature measurements were made are given as superscripts, e.g. pK²⁵. Where no temperature is given, it is assumed that the measurements were carried out at room temperature, e.g. 15—25°. No temperature is given for estimated values (pK_{Est} ~) and these have been calculated from data at room temperature. The variation of pK with temperature is given by the equation:

$$-d(pK)/dT = (pK + 0.052\Delta S^{\circ})/T$$

where T is in degrees Kelvin and ΔS° is in Joules deg⁻¹ mol⁻¹. The $-d(pK)/dT$ in the range of temperatures between 5 to 70° is generally small (e.g. between ~0.0024 and ~0.04), and for chemical purification purposes is not a seriously deterring factor. It does however, vary with the compound under study because ΔS° varies from compound to compound. The following are examples of the effect of temperature on pK values: for imidazole the pK values are 7.57 (0°), 7.33 (10°), 7.10 (20°), 6.99 (25°), 6.89 (30°), 6.58 (40°) and 6.49 (50°), and for 3,5-dinitrobenzoic acid they are 2.60 (10°), 2.73 (20°), 2.85 (30°), 2.96 (40°) and 3.07 (40°), and for *N*-acetyl- β -alanine they are 4.4788 (5°), 4.4652 (10°), 4.4564 (15°), 4.4488 (20°), 4.4452 (25°), 4.4444 (30°), 4.4434 (35°) and 4.4412 (40°).

pK and solvent.

All stated pK values in this book are for data in dilute aqueous solutions unless otherwise stated, although the dielectric constants, ionic strengths of the solutions and the method of measurement, e.g. potentiometric, spectrophotometric etc, are not given. Estimated values are also for dilute aqueous solutions whether or not the material is soluble enough in water. Generally the more dilute the solution the closer is the pK to the real thermodynamic value. The pK in mixed aqueous solvents can vary considerably with the relative concentrations and with the nature of the solvents. For example the pK²⁵ values for *N*-benzylpenicillin are 2.76 and 4.84 in H₂O and H₂O/EtOH (20:80) respectively; the pK²⁵ values for (-)-ephedrine are 9.58 and 8.84 in H₂O and H₂O/MeOCH₂CH₂OH (20:80) respectively; and for cyclopentylamine the pK²⁵ values are 10.65 and 4.05 in H₂O and H₂O/EtOH (50:50) respectively. pK values in acetic acid or aqueous acetic acid are generally lower than in H₂O.

The dielectric constant of the medium affects the equilibria where charges are generated in the dissociations e.g. $AH \rightleftharpoons A^{-} + H^{+}$ and therefore affects the pK values. However, its effect on dissociations where there are no changes in total charge such as $BH^{+} \rightleftharpoons B + H^{+}$ is considerably less, with a slight decrease in pK with decreasing dielectric constant.

DISTILLATION

One of the most widely applicable and most commonly used methods of purification of liquids or low melting solids (especially of organic chemicals) is fractional distillation at atmospheric, or some lower, pressure. Almost without exception, this method can be assumed to be suitable for all organic liquids and most of the low-melting organic solids. For this reason it has been possible in Chapter 4 to omit many procedures for purification of organic chemicals when only a simple fractional distillation is involved - the suitability of such a procedure is implied from the boiling point.

The boiling point of a liquid varies with the 'atmospheric' pressure to which it is exposed. A liquid boils when its vapour pressure is the same as the external pressure on its surface, its normal boiling point being the temperature at which its vapour pressure is equal to that of a standard atmosphere (760mm Hg). Lowering the external pressure lowers the boiling point. For most substances, boiling point and vapour pressure are related by an equation of the form,

$$\log p = A + B/(t + 273),$$

where p is the pressure, t is in °C, and A and B are constants. Hence, if the boiling points at two different pressures are known the boiling point at another pressure can be calculated from a simple plot of $\log p$ versus $1/(t + 273)$. For organic molecules that are not strongly associated, this equation can be written in the form,

$$\log p = 8.586 - 5.703 (T + 273)/(t + 273)$$

where T is the boiling point in °C at 760mm Hg. Tables 3A and 3B give computed boiling points over a range of pressures. Some examples illustrate its application. Ethyl acetoacetate, b 180° (with decomposition) at 760mm Hg has a predicted b of 79° at 16mm; the experimental value is 78°. Similarly 2,4-diaminotoluene, b 292° at 760mm, has a predicted b of 147° at 8mm; the experimental value is 148-150°. For self-associated molecules the predicted b are lower than the experimental values. Thus, glycerol, b 290° at 760mm, has a predicted b of 146° at 8mm: the experimental value is 182°.

Similarly an estimate of the boiling points of liquids at reduced pressure can be obtained using a nomogram (see Figure 1).

For pressures near 760mm, the change in boiling point is given approximately by,

$$\hat{t} = a(760 - p)(t + 273)$$

where $a = 0.00012$ for most substances, but $a = 0.00010$ for water, alcohols, carboxylic acids and other associated liquids, and $a = 0.00014$ for very low-boiling substances such as nitrogen or ammonia [Crafts *Chem Ber* 20 709 1887]. When all the impurities are non-volatile, simple distillation is adequate purification. The observed boiling point remains almost constant and approximately equal to that of the pure material. Usually, however, some of the impurities are appreciably volatile, so that the boiling point progressively rises during the distillation because of the progressive enrichment of the higher-boiling components in the distillation flask. In such cases, separation is effected by fractional distillation using an efficient column.

Techniques.

The distillation apparatus consists basically of a distillation flask, usually fitted with a vertical fractionating column (which may be empty or packed with suitable materials such as glass helices or stainless-steel wool) to which is attached a condenser leading to a receiving flask. The bulb of a thermometer projects into the vapour phase just below the region where the condenser joins the column. The distilling flask is heated so that its contents are steadily vaporised by boiling. The vapour passes up into the column where, initially, it condenses and runs back into the flask. The resulting heat transfer gradually warms the column so that there is a progressive movement of the vapour phase-liquid boundary up the column, with increasing enrichment of the more volatile component. Because of this fractionation, the vapour finally passing into the condenser (where it condenses and flows into the receiver) is commonly that of the lowest-boiling components in the system. The conditions apply until all of the low-boiling material has been distilled, whereupon distillation ceases until the column temperature is high enough to permit the next component to distil. This usually results in a temporary fall in the temperature indicated by the thermometer.

Distillation of liquid mixtures.

The principles involved in fractional distillation of liquid mixtures are complex but can be seen by considering a system which approximately obeys *Raoult's law*. (This law states that the vapour pressure of a solution at any given temperature is the sum of the vapour pressures of each component multiplied by its mole fraction in the solution.) If two substances, A and B, having vapour pressures of 600mm Hg and 360mm Hg, respectively, were mixed in a molar ratio of 2:1 (i.e. 0.666:0.333 mole ratio), the mixture would have (ideally) a vapour pressure of 520mm Hg (i.e. $600 \times 0.666 + 360 \times 0.333$, or $399.6 + 119.88$ mm Hg) and the vapour phase would contain 77% ($399.6 \times 100/520$) of A and 23% ($119.88 \times 100/520$) of B. If this phase was now condensed, the new liquid phase would, therefore, be richer in the volatile component A. Similarly, the vapour in equilibrium with this phase is still further enriched in A. Each such liquid-vapour equilibrium constitutes a "theoretical plate". The efficiency of a fractionating column is commonly expressed as the number of such plates to which it corresponds in operation. Alternatively, this information may be given in the form of the height equivalent to a theoretical plate, or HETP. The number of theoretical plates and equilibria between liquids and vapours are affected by the factors listed to achieve maximum separation by fractional distillation in the section below on techniques. In most cases, systems deviate to a greater or lesser extent from Raoult's law, and vapour pressures may be greater or less than the values calculated. In extreme cases (e.g. azeotropes), vapour pressure-composition curves pass through maxima or minima, so that attempts at fractional distillation lead finally to the separation of a constant-boiling (azeotropic) mixture and one (but not both) of the pure species if either of the latter is present in excess.

Elevation of the boiling point by dissolved solids. Organic substances dissolved in organic solvents cause a rise in boiling point which is proportional to the concentration of the substance, and the extent of rise in temperature is characteristic of the solvent. The following equation applies for dilute solutions and non-associating substances:

$$\frac{M Dt}{c} = K$$

Where M is the molecular weight of the solute, Dt is the elevation of boiling point in °C, c is the concentration of solute in grams for 1000gm of solvent, and K is the *Ebullioscopic Constant* (molecular elevation of the boiling point) for the solvent. K is a fixed property (constant) for the particular solvent. This has been very useful for the determination of the molecular weights of organic substances in solution.

The efficiency of a distillation apparatus used for purification of liquids depends on the difference in boiling points of the pure material and its impurities. For example, if two components of an ideal mixture have vapour pressures in the ratio 2:1, it would be necessary to have a still with an efficiency of at least seven plates (giving an enrichment of $2^7 = 128$) if the concentration of the higher-boiling component in the distillate was to be reduced to less than 1% of its initial value. For a vapour pressure ratio of 5:1, three plates would achieve as much separation.

In a fractional distillation, it is usual to reject the initial and final fractions, which are likely to be richer in the lower-boiling and higher-boiling impurities respectively. The centre fraction can be further purified by repeated fractional distillation.

To achieve maximum separation by fractional distillation:

1. The column must be flooded initially to wet the packing. For this reason it is customary to operate a still at reflux for some time before beginning the distillation.
2. The reflux ratio should be high (i.e. the ratio of drops of liquid which return to the distilling flask and the drops which distil over), so that the distillation proceeds slowly and with minimum disturbance of the equilibria in the column.
3. The hold-up of the column should not exceed one-tenth of the volume of any one component to be separated.
4. Heat loss from the column should be prevented but, if the column is heated to offset this, its temperature must not exceed that of the distillate in the column.
5. Heat input to the still-pot should remain constant.
6. For distillation under reduced pressure there must be careful control of the pressure to avoid flooding or cessation of reflux.

Types of distillation

The distilling flask. To minimise superheating of the liquid (due to the absence of minute air bubbles or other suitable nuclei for forming bubbles of vapour), and to prevent bumping, one or more of the following precautions should be taken:

(a) The flask is heated uniformly over a large part of its surface, either by using an electrical heating mantle or, by partial immersion in a bath above the boiling point of the liquid to be distilled.

(b) Before heating begins, small pieces of unglazed fireclay or porcelain (porous pot, boiling chips), pumice, diatomaceous earth, or platinum wire are added to the flask. These act as sources of air bubbles.

(c) The flask may contain glass siphons or boiling tubes. The former are inverted J-shaped tubes, the end of the shorter arm being just above the surface of the liquid. The latter comprise long capillary tubes sealed above the lower end.

(d) A steady slow stream of inert gas (e.g. N_2 , Ar or He) is passed through the liquid.

(e) The liquid in the flask is stirred mechanically. This is especially necessary when suspended insoluble material is present.

For simple distillations a Claisen flask is often used. This flask is, essentially, a round-bottomed flask to the neck of which is joined another neck carrying a side arm. This second neck is sometimes extended so as to form a

Vigreux column [a glass tube in which have been made a number of pairs of indentations which almost touch each other and which slope slightly downwards. The pairs of indentations are arranged to form a spiral of glass inside the tube].

For heating baths, see Table 4. For distillation apparatus on a micro or semi-micro scale see Aldrich and other glassware catalogues. Alternatively, some useful websites for suppliers of laboratory glassware are www.wheatonsci.com, www.sigmaaldrich.com and www.kimble-kontes.com.

Types of columns and packings. A slow distillation rate is necessary to ensure that equilibrium conditions operate and also that the vapour does not become superheated so that the temperature rises above the boiling point. Efficiency is improved if the column is heat insulated (either by vacuum jacketing or by lagging) and, if necessary, heated to just below the boiling point of the most volatile component. Efficiency of separation also improves with increase in the heat of vaporisation of the liquids concerned (because fractionation depends on heat equilibration at multiple liquid-gas boundaries). Water and alcohols are more easily purified by distillation for this reason.

Columns used in distillation vary in their shapes and types of packing. Packed columns are intended to give efficient separation by maintaining a large surface of contact between liquid and vapour. Efficiency of separation is further increased by operation under conditions approaching total reflux, i.e. under a high reflux ratio. However, great care must be taken to avoid flooding of the column during distillation. The minimum number of theoretical plates for satisfactory separation of two liquids differing in boiling point by t is approximately $(273 + t)/3t$, where t is the average boiling point in $^{\circ}\text{C}$.

The packing of a column greatly increases the surface of liquid films in contact with the vapour phase, thereby increasing the efficiency of the column, but reducing its capacity (the quantities of vapour and liquid able to flow in opposite directions in a column without causing flooding). Material for packing should be of uniform size, symmetrical shape, and have a unit diameter less than one eighth that of the column. (Rectification efficiency increases sharply as the size of the packing is reduced but so, also, does the hold-up in the column.) It should also be capable of uniform, reproducible packing.

The usual *packings* are:

(a) Rings. These may be hollow glass or porcelain (Raschig rings), of stainless steel gauze (Dixon rings), or hollow rings with a central partition (Lessing rings) which may be of porcelain, aluminium, copper or nickel.

(b) Helices. These may be of metal or glass (Fenske rings), the latter being used where resistance to chemical attack is important (e.g. in distilling acids, organic halides, some sulfur compounds, and phenols). Metal single-turn helices are available in aluminium, nickel or stainless steel. Glass helices are less efficient, because they cannot be tamped to ensure uniform packing.

(c) Balls or beads. These are usually made of glass.

Condensers. Some of the more commonly used condensers are:

Air condenser. A glass tube such as the inner part of a Liebig condenser (see below). Used for liquids with boiling points above 90° . Can be of any length.

Coil condenser. An open tube, into which is sealed a glass coil or spiral through which water circulates. The tube is sometimes also surrounded by an outer cooling jacket. A double coil condenser has two inner coils with circulating water.

Double surface condenser. A tube in which the vapour is condensed between an outer and inner water-cooled jacket after impinging on the latter. Very useful for liquids boiling below 40° .

Friedrichs condenser. A "cold-finger" type of condenser sealed into a glass jacket open at the bottom and near the top. The cold finger is formed into glass screw threads.

Liebig condenser. An inner glass tube surrounded by a glass jacket through which water is circulated.

Vacuum distillation. This expression is commonly used to denote a distillation under reduced pressure lower than that of the normal atmosphere. Because the boiling point of a substance depends on the pressure, it is often possible by sufficiently lowering the pressure to distil materials at a temperature low enough to avoid partial or complete decomposition, even if they are unstable when boiled at atmospheric pressure.

Sensitive or high-boiling liquids should invariably be distilled or fractionally distilled under reduced pressure. The apparatus is essentially as described for distillation except that ground joints connecting the different parts of the apparatus should be air tight by using grease, or better Teflon sleeves. For low, moderately high, and very high temperatures Apiezon L, M and T greases respectively, are very satisfactory. Alternatively, it is often preferable to avoid grease and to use thin Teflon sleeves in the joints. The distilling flask, must be supplied with a capillary

bleed (which allows a fine stream of air, nitrogen or argon into the flask), and the receiver should be of the fraction collector type. When distilling under vacuum it is very important to place a loose packing of glass wool above the liquid to buffer sudden boiling of the liquid. The flask should be not more than two-thirds full of liquid. The vacuum must have attained a steady state, i.e. the liquid has been completely degassed, before the heat source is applied, and the temperature of the heat source must be raised *very slowly* until boiling is achieved.

If the pump is a filter pump off a high-pressure water supply, its performance will be limited by the temperature of the water because the vapour pressure of water at 10°, 15°, 20° and 25° is 9.2, 12.8, 17.5 and 23.8 mm Hg respectively. The pressure can be measured with an ordinary manometer. For vacuums in the range 10⁻² mm Hg to 10 mm Hg, rotary mechanical pumps (oil pumps) are used and the pressure can be measured with a Vacostat McLeod type gauge. If still higher vacuums are required, for example for high vacuum sublimations, a mercury diffusion pump is suitable. Such a pump can provide a vacuum up to 10⁻⁶ mm Hg. For better efficiencies, the pump can be backed up by a mechanical pump. In all cases, the mercury pump is connected to the distillation apparatus through several traps to remove mercury vapours. These traps may operate by chemical action, for example the use of sodium hydroxide pellets to react with acids, or by condensation, in which case empty tubes cooled in solid carbon dioxide-ethanol or liquid nitrogen (contained in wide-mouthed Dewar flasks) are used.

Special oil or mercury traps are available commercially and a liquid-nitrogen (b -209.9°C) trap is the most satisfactory one to use between these and the apparatus. It has an advantage over liquid air or oxygen in that it is non-explosive if it becomes contaminated with organic matter. Air should not be sucked through the apparatus before starting a distillation because this will cause liquid oxygen (b -183°C) to condense in the liquid nitrogen trap and this is potentially explosive (especially in mixtures with organic materials). Due to the potential lethal consequences of liquid oxygen/organic material mixtures, care must be exercised when handling liquid nitrogen. Hence, it is advisable to degas the system for a short period before the trap is immersed into the liquid nitrogen (which is kept in a Dewar flask).

Spinning-band distillation. Factors which limit the performance of distillation columns include the tendency to flood (which occurs when the returning liquid blocks the pathway taken by the vapour through the column) and the increased hold-up (which decreases the attainable efficiency) in the column that should, theoretically, be highly efficient. To overcome these difficulties, especially for distillation under high vacuum of heat sensitive or high-boiling highly viscous fluids, spinning band columns are commercially available. In such units, the distillation columns contain a rapidly rotating, motor-driven, spiral band, which may be of polymer-coated metal, stainless steel or platinum. The rapid rotation of the band in contact with the walls of the still gives intimate mixing of descending liquid and ascending vapour while the screw-like motion of the band drives the liquid towards the still-pot, helping to reduce hold-up. There is very little pressure drop in such a system, and very high throughputs are possible, with high efficiency. For example, a 765-mm long 10-mm diameter commercial spinning-band column is reported to have an efficiency of 28 plates and a pressure drop of 0.2 mm Hg for a throughput of 330mL/h. The columns may be either vacuum jacketed or heated externally. The stills can be operated down to 10⁻⁵ mm Hg. The principle, which was first used commercially in the Podbielniak Centrifugal Superfractionator, has also been embodied in descending-film molecular distillation apparatus.

Steam distillation. When two immiscible liquids distil, the sum of their (independent) partial pressures is equal to the atmospheric pressure. Hence in steam distillation, the distillate has the composition

$$\frac{\text{Moles of substance}}{\text{Moles of water}} = \frac{P_{\text{substance}}}{P_{\text{water}}} = \frac{760 - P_{\text{water}}}{P_{\text{water}}}$$

where the *P*'s are vapour pressures (in mm Hg) in the boiling mixture.

The customary technique consists of heating the substance and water in a flask (to boiling), usually with the passage of steam, followed by condensation and separation of the aqueous and non-aqueous phases in the distillate. Its advantages are those of selectivity (because only some water-insoluble substances, such as naphthalene, nitrobenzene, phenol and aniline are volatile in steam) and of ability to distil certain high-boiling substances well below their boiling point. It also facilitates the recovery of a non-steam-volatile solid at a relatively low temperature from a high-boiling solvent such as nitrobenzene. The efficiency of steam distillation is increased if superheated steam is used (because the vapour pressure of the organic component is increased relative to water). In this case the flask containing the material is heated (without water) in an oil bath and the steam passing through it is superheated by prior passage through a suitable heating device (such as a copper coil heated electrically or an oil bath).

Azeotropic distillation. In some cases two or more liquids form constant-boiling mixtures, or azeotropes. Azeotropic mixtures are most likely to be found with components which readily form hydrogen bonds or are otherwise highly associated, especially when the components are dissimilar, for example an alcohol and an aromatic hydrocarbon, but have similar boiling points.

Examples where the boiling point of the distillate is a minimum (less than either pure component) include:

Water with ethanol, *n*-propanol and isopropanol, *tert*-butanol, propionic acid, butyric acid, pyridine,

methanol with methyl iodide, methyl acetate, chloroform,

ethanol with ethyl iodide, ethyl acetate, chloroform, benzene, toluene, methyl ethyl ketone,

benzene with cyclohexane,

acetic acid with toluene.

Although less common, azeotropic mixtures are known which have higher boiling points than their components. These include water with most of the mineral acids (hydrofluoric, hydrochloric, hydrobromic, perchloric, nitric and sulfuric) and formic acid. Other examples are acetic acid-pyridine, acetone-chloroform, aniline-phenol, and chloroform-methyl acetate.

The following azeotropes are important commercially for drying ethanol:

ethanol 95.5% (by weight) - water 4.5%	b 78.1°
ethanol 32.4% - benzene 67.6%	b 68.2°
ethanol 18.5% - benzene 74.1% - water 7.4%	b 64.9°

Materials are sometimes added to form an azeotropic mixture with the substance to be purified. Because the azeotrope boils at a different temperature, this facilitates separation from substances distilling in the same range as the pure material. (Conversely, the impurity might form the azeotrope and be removed in this way). This method is often convenient, especially where the impurities are isomers or are otherwise closely related to the desired substance. Formation of low-boiling azeotropes also facilitates distillation.

One or more of the following methods can generally be used for separating the components of an azeotropic mixture:

1. By using a chemical method to remove most of one species prior to distillation. (For example, water can be removed by suitable drying agents; aromatic and unsaturated hydrocarbons can be removed by sulfonation).
2. By redistillation with an additional substance which can form a ternary azeotropic mixture (as in ethanol-water-benzene example given above).
3. By selective adsorption of one of the components. (For example, of water on to silica gel or molecular sieves, or of unsaturated hydrocarbons onto alumina).
4. By fractional crystallisation of the mixture, either by direct freezing or by dissolving in a suitable solvent.

Kügelrohr distillation. The apparatus (Büchi, see www.buchi.com) is made up of small glass bulbs (*ca* 4-5cm diameter) which are joined together *via* Quickfit joints at each pole of the bulbs. The liquid (or low melting solid) to be purified is placed in the first bulb of a series of bulbs joined end to end, and the system can be evacuated. The first bulb is heated in a furnace at a high temperature whereby most of the material distils into the second bulb (which is outside of the furnace). The second bulb is then moved into the furnace and the furnace temperature is reduced by *ca* 5° whereby the liquid in the second bulb distils into the third bulb (at this stage the first bulb is now out at the back of the furnace and the third and subsequent bulbs are outside the front of the furnace). The furnace temperature is lowered by a further *ca* 5° and the third bulb is moved into the furnace. The lower boiling material will distil into the fourth bulb. The process is continued until no more material distils into the subsequent bulb. The vacuum (if applied) and the furnace are removed, the bulbs are separated and the various fractions of distillates are collected from the individual bulbs. For volatile liquids, it may be necessary to cool the receiving bulb with solid CO₂ held in a suitable container (Kügelrohr distillation apparatus with an integrated cooling system is available). This procedure is used for preliminary purification and the distillates are then redistilled or recrystallised.

Isopiestic or isothermal distillation. This technique can be useful for the preparation of metal-free solutions of volatile acids and bases for use in trace metal studies. The procedure involves placing two beakers, one of distilled water and the other of a solution of the material to be purified, in a desiccator. The desiccator is sealed and left to stand at room temperature for several days. The volatile components distribute themselves between the two beakers whereas the non-volatile contaminants remain in the original beaker. This technique has afforded metal-free pure solutions of ammonia, hydrochloric acid and hydrogen fluoride.

RECRYSTALLISATION

Techniques

The most commonly used procedure for the purification of a solid material by recrystallisation from a solution involves the following steps:

- (a) The impure material is dissolved in a suitable solvent, by shaking or vigorous stirring, at or near the boiling point, to form a near-saturated solution.
- (b) The hot solution is filtered to remove any insoluble particles. To prevent crystallisation during this filtration, a heated filter funnel can be used or the solution can be diluted with more of the solvent.
- (c) The solution is then allowed to cool so that the dissolved substance crystallises out.
- (d) The crystals are separated from the mother liquor, either by centrifuging or by filtering, under suction, through a sintered glass, a Hirsch or a Büchner, funnel. Usually, centrifugation is preferred because of the greater ease and efficiency of separating crystals and mother liquor, and also because of the saving of time and effort, particularly when very small crystals are formed or when there is entrainment of solvent.
- (e) The crystals are washed free from mother liquor with a little fresh cold solvent, then dried.

If the solution contains extraneous coloured material likely to contaminate the crystals, this can often be removed by adding some activated charcoal (decolorising carbon) to the hot, but not boiling, solution which is then shaken frequently for several minutes before being filtered. (The large active surface of the carbon makes it a good adsorbent for this purpose.) In general, the cooling and crystallisation steps should be rapid so as to give small crystals which occlude less of the mother liquor. This is usually satisfactory with inorganic material, so that commonly the filtrate is cooled in an ice-water bath while being vigorously stirred. In many cases, however, organic molecules crystallise much more slowly, so that the filtrate must be set aside to cool to room temperature or left in the refrigerator. It is often desirable to subject material that is very impure to preliminary purification, such as steam distillation, Soxhlet extraction, or sublimation, before recrystallising it. A greater degree of purity is also to be expected if the crystallisation process is repeated several times, especially if different solvents are used. The advantage of several crystallisations from different solvents lies in the fact that the material sought, and its impurities, are unlikely to have similar solubilities as solvents and temperatures are varied.

For the final separation of solid material, sintered-glass discs are preferable to filter paper. Sintered glass is unaffected by strongly acid solutions or by oxidising agents. Also, with filter paper, cellulose fibres are likely to become included in the sample. The sintered-glass discs or funnels can be readily cleaned by washing in freshly prepared *chromic acid cleaning mixture*. This mixture is made by adding 100mL of concentrated sulfuric acid slowly with stirring to a solution of 5g of sodium dichromate (CARE: cancer suspect) in 5mL of water. (The mixture warms to about 70°, see p 3).

For materials with very low melting points it is sometimes convenient to use dilute solutions in acetone, methanol, pentane, diethyl ether or $\text{CHCl}_3\text{-CCl}_4$. The solutions are cooled to -78° in a dry-ice/acetone bath, to give a slurry which is filtered off through a precooled Büchner funnel. Experimental details, as applied to the purification of nitromethane, are given by Parrett and Sun [*J Chem Educ* 54 448 1977].

Where substances vary little in solubility with temperature, *isothermal crystallisation* may sometimes be employed. This usually takes the form of a partial evaporation of a saturated solution at room temperature by leaving it under reduced pressure in a desiccator.

However, in rare cases, crystallisation is not a satisfactory method of purification, especially if the impurity forms crystals that are isomorphous with the material being purified. In fact, the impurity content may even be greater in such recrystallised material. For this reason, it still remains necessary to test for impurities and to remove or adequately lower their concentrations by suitable chemical manipulation prior to recrystallisation.

Filtration. Filtration removes particulate impurities rapidly from liquids and is also used to collect insoluble or crystalline solids which separate or crystallise from solution. The usual technique is to pass the solution, cold or hot, through a fluted filter paper in a conical glass funnel.

If a solution is hot and needs to be filtered rapidly a Büchner funnel and flask are used and filtration is performed under a slight vacuum (water pump), the filter medium being a circular cellulose filter paper wet with solvent. If filtration is slow, even under high vacuum, a pile of about twenty filter papers, wet as before, are placed in the Büchner funnel and, as the flow of solution slows down, the upper layers of the filter paper are progressively removed. Alternatively, a filter aid, e.g. Celite, Florisil or Hyflo-supercel, is placed on top of a filter paper in the funnel. When the flow of the solution (under suction) slows down, the upper surface of the filter aid is scratched gently. Filter papers with various pore sizes are available covering a range of filtration rates. Hardened filter papers are slow filtering but they can withstand acidic and alkaline solutions without appreciable hydrolysis of the

cellulose (see Table 5). When using strong acids it is preferable to use glass micro fibre filters which are commercially available (see Table 5 and 6).

Freeing a solution from extremely small particles [e.g. for optical rotatory dispersion (ORD) or circular dichroism (CD) measurements] requires filters with very small pore size. Commercially available (Millipore, Gelman, Nucleopore) filters other than cellulose or glass include nylon, Teflon, and polyvinyl chloride, and the pore diameter may be as small as 0.01micron (see Table 6). Special containers are used to hold the filters, through which the solution is pressed by applying pressure, e.g. from a syringe. Some of these filters can be used to clear strong sulfuric acid solutions.

As an alternative to the Büchner funnel for collecting crystalline solids, a funnel with a sintered glass-plate under suction may be used. Sintered-glass funnels with various porosities are commercially available and can be easily cleaned with warm chromic or nitric acid (see above).

When the solid particles are too fine to be collected on a filter funnel because filtration is extremely slow, separation by **centrifugation** should be used. Bench type centrifuges are most convenient for this purpose. The solid is placed in the centrifuge tube, the tubes containing the solutions on opposite sides of the rotor should be balanced accurately (at least within 0.05 to 0.1g), and the solutions are spun at maximum speed for as long as it takes to settle the solid (usually *ca* 3-5 minutes). The solid is washed with cold solvent by centrifugation, and finally twice with a pure volatile solvent in which the solid is insoluble, also by centrifugation. After decanting the supernatant, the residue is dried in a vacuum, at elevated temperatures if necessary. In order to avoid "spitting" and contamination with dust while the solid in the centrifuge tube is dried, the mouth of the tube is covered with aluminium foil and held fast with a tight rubber band near the lip. The flat surface of the aluminium foil is then perforated in several places with a pin and the tube and contents are dried in a vacuum desiccator over a desiccant.

Choice of solvents. The best solvents for recrystallisation have the following properties:

- (a) The material is much more soluble at higher temperatures than it is at room temperature or below.
- (b) Well-formed (but not large) crystals are produced.
- (c) Impurities are either very soluble or only sparingly soluble.
- (d) The solvent must be readily removed from the purified material.
- (e) There must be no reaction between the solvent and the substance being purified.
- (f) The solvent must not be inconveniently volatile or too highly flammable. (These are reasons why diethyl ether and carbon disulfide are not commonly used in this way.)

The following generalisations provide a rough guide to the selection of a suitable solvent:

- (a) Substances usually dissolve best in solvents to which they are most closely related in chemical and physical characteristics. Thus, hydroxylic compounds are likely to be most soluble in water, methanol, ethanol, acetic acid or acetone. Similarly, petroleum ether might be used with water-insoluble substances. However, if the resemblance is too close, solubilities may become excessive.
- (b) Higher members of homologous series approximate more and more closely to their parent hydrocarbon.
- (c) Polar substances are more soluble in polar, than in non-polar, solvents.

Although Chapters 4, 5 and 6 provide details of the solvents used for recrystallising a large portion of commercially available laboratory chemicals, they cannot hope to be exhaustive, nor need they necessarily be the best choice. In other cases where it is desirable to use this process, it is necessary to establish whether a given solvent is suitable. This is usually done by taking only a small amount of material in a small test-tube and adding enough solvent to cover it. If it dissolves readily in the cold or on gentle warming, the solvent is unsuitable. Conversely, if it remains insoluble when the solvent is heated to boiling (adding more solvent if necessary), the solvent is again unsuitable. If the material dissolves in the hot solvent but does not crystallise readily within several minutes of cooling in an ice-salt mixture, another solvent should be tried.

Petroleum ethers are commercially available fractions of refined petroleum and are sold in fractions with about 20° boiling ranges. This ensures that little of the hydrocarbon ingredients boiling below the range is lost during standing or boiling when recrystallising a substance. Petroleum ethers with boiling ranges (at 760mm pressure) of 35—60°, 40—60°, 60—80°, 80—100°, and 100—120° are generally free from unsaturated and aromatic hydrocarbons. The lowest boiling petroleum ether commercially available has **b** 30-40°/760mm and is mostly *n*-pentane. The purer spectroscopic grades are almost completely free from olefinic and aromatic hydrocarbons. **Petroleum spirit** (which is sometimes used synonymously with petroleum ether or light

petroleum) is usually less refined petroleum, and *ligroin* is used for fractions boiling above 100°. The lower boiling fractions consist of mixtures of *n*-pentane (b 36°), *n*-hexane (b 68.5°) and *n*-heptane (b 98°), and some of their isomers in varying proportions. For purification of petroleum ether b 35-60° see p. 324.

Solvents commonly used for recrystallisation, and their boiling points, are given in Table 7. For comments on the toxicity and use of **benzene** see the first pages of Chapters 4, 5 and 6.

Mixed Solvents. Where a substance is too soluble in one solvent and too insoluble in another, for either to be used for recrystallisation, it is often possible (provided they are miscible) to use them as a mixed solvent. (In general, however, it is preferable to use a single solvent if this is practicable.) Table 8 contains many of the common pairs of miscible solvents.

The technique of recrystallisation from mixed solvents is as follows:

The material is dissolved in the solvent in which it is the more soluble, then the other solvent (heated to near boiling) is added cautiously to the hot solution until a slight turbidity persists or crystallisation begins. This is cleared by adding several drops of the first solvent, and the solution is allowed to cool and crystallise in the usual way.

A variation of this procedure is simply to precipitate the material in a microcrystalline form from solution in one solvent at room temperature, by adding a little more of the second solvent, filtering off the crystals, adding a little more of the second solvent and repeating the process. This ensures, at least in the first or last precipitation, a material which contains as little as possible of the impurities, which may also be precipitated in this way. With salts, the first solvent is commonly water, and the second solvent is alcohol or acetone.

Recrystallisation from the melt. A crystalline solid melts when its temperature is raised sufficiently for the thermal agitation of its molecules or ions to overcome the restraints imposed by the crystal lattice. Usually, impurities weaken crystal structures, and hence lower the melting points of solids (or the freezing points of liquids). If an impure material is melted and cooled slowly (with the addition, if necessary, of a trace of solid material near the freezing point to avoid supercooling), the first crystals that form will usually contain less of the impurity, so that fractional solidification by partial freezing can be used as a purification process for solids with melting points lying in a convenient temperature range (or for more readily frozen liquids). Some examples of cooling baths that are useful in recrystallisation are summarised in Table 9. In some cases, impurities form higher melting eutectics with substances to be purified, so that the first material to solidify is less pure than the melt. For this reason, it is often desirable to discard the first crystals and also the final portions of the melt. Substances having similar boiling points often differ much more in melting points, so that fractional solidification can offer real advantages, especially where ultrapurity is sought. For further information on this method of recrystallisation, consult the earlier editions of this book as well as references by Schwab and Wichers (*J Res Nat Bur Stand* 25 747 1940). This method works best if the material is already nearly pure, and hence tends to be a final purification step.

Zone refining. Zone refining (or zone melting) is a particular development for fractional solidification and is applicable to all crystalline substances that show differences in the concentrations of impurities in liquid and solid states at solidification. The apparatus used in this technique consists essentially of a device in which the crystalline solid to be purified is placed in a glass tube (set vertically) which is made to move slowly upwards while it passes through a fixed coil (one or two turns) of heated wire. A narrow zone of molten crystals is formed when the tube is close to the heated coil. As the zone moves away from the coil the liquid crystallises, and a fresh molten zone is formed below it at the coil position. The machine can be set to recycle repeatedly. At its advancing side, the zone has a melting interface with the impure material whereas on the upper surface of the zone there is a constantly growing face of higher-melting, resolidified material. This leads to a progressive increase in impurity in the liquid phase which, at the end of the run, is discarded from the bottom of the tube. Also, because of the progressive increase in impurity in the liquid phase, the resolidified material contains correspondingly more of the impurities. For this reason, it is usually necessary to make several zone-melting runs before a sample is satisfactorily purified. This is also why the method works most successfully if the material is already fairly pure. In all these operations the zone must travel slowly enough to enable impurities to diffuse or be convected away from the area where resolidification is occurring.

The technique finds commercial application in the production of metals of extremely high purity (impurities down to 10⁻⁹ ppm), in purifying refractory oxides, and in purifying organic compounds, using commercially available equipment. Criteria for indicating that definite purification is achieved include elevation of melting point, removal of colour, fluorescence or smell, and a lowering of electrical conductivity. Difficulties likely to be found with organic compounds, especially those of low melting points and low rates of crystallisation, are supercooling and, because of surface tension and contraction, the tendency of the molten zone to seep back into the recrystallised areas. The method is likely to be useful in cases where fractional distillation is not practicable, either because of

unfavourable vapour pressures or ease of decomposition, or where super-pure materials are required. It has been used for the latter purpose for purifying anthracene, benzoic acid, chrysene, morphine, 1,8-naphthyridine and pyrene to name a few. [See E.F.G.Herington, *Zone Melting of Organic Compounds*, Wiley & Sons, NY, 1963; W.Pfann, *Zone Melting*, 2nd edn, Wiley, NY, 1966; H.Schildknecht, *Zonenschmelzen*, Verlag Chemie, Weinheim, 1964; W.R.Wilcox, R.Friedenberg et al. *Chem Rev* **64** 187 1964; M.Zief and W.R.Wilcox (Eds), *Fractional Solidification*, Vol I, M Dekker Inc. NY, 1967.]

SUBLIMATION

Sublimation differs from ordinary distillation because the vapour condenses to a solid instead of a liquid. Usually, the pressure in the heated system is diminished by pumping, and the vapour is condensed (after travelling a relatively short distance) onto a cold finger or some other cooled surface. This technique, which is applicable to many organic solids, can also be used with inorganic solids such as aluminium chloride, ammonium chloride, arsenious oxide and iodine. In some cases, passage of a stream of inert gas over the heated substance secures adequate vaporisation. This procedure has the added advantage of removing occluded solvent used in recrystallising the solid.

CHROMATOGRAPHY

Chromatography is often used with advantage for the purification of small amounts of complex organic mixtures. Chromatography techniques all rely on the differential distribution of the various components in a mixture between the mobile phase and the stationary phase. The mobile phase can either be a gas or a liquid whereas the stationary phase can either be a solid or a liquid.

The major chromatographic techniques can also be categorised according to the nature of the mobile phase used - vapour phase chromatography for when a gas is the mobile phase and liquid chromatography for when a liquid is the mobile phase.

A very useful catalog for chromatographic products and information relating to chromatography (from gas chromatography to biochromatography) is that produced by Merck, called the ChromBook and the associated compact disk, ChromCircle.

Vapour phase chromatography (GC or gas-liquid chromatography)

The mobile phase in vapour phase chromatography is a gas (e.g. hydrogen, helium, nitrogen or argon) and the stationary phase is a non-volatile liquid impregnated onto a porous material. The mixture to be purified is injected into a heated inlet whereby it is vaporised and taken into the column by the carrier gas. It is separated into its components by partition between the liquid on the porous support and the gas. For this reason vapour-phase chromatography is sometimes referred to as gas-liquid chromatography (g.l.c). Vapour phase chromatography is very useful in the resolution of a mixture of volatile compounds. This type of chromatography uses either packed or capillary columns. Packed columns have internal diameters of 3-5 mm with lengths of 2-6 m. These columns can be packed with a range of materials including firebrick derived materials (chromasorb P, for separation of non polar hydrocarbons) or diatomaceous earth (chromasorb W, for separation of more polar molecules such as acids, amines). Capillary columns have stationary phase bonded to the walls of long capillary tubes. The diameters in capillary columns are less than 0.5 mm and the lengths of these columns can go up to 50 m! These columns have much superior separating powers than the packed columns. Elution times for equivalent resolutions with packed columns can be up to ten times shorter. It is believed that almost any mixture of compounds can be separated using one of the four stationary phases, OV-101, SE-30, OV-17 and Carbowax-20M. The use of capillary columns in gas chromatography for analysis is now routinely carried out. An extensive range of packed and capillary columns is available from chromatographic specialists such as Supelco, Alltech, Hewlett-Packard, Phenomenex etc.

Table 10 shows some typical liquids used for stationary phases in gas chromatography.

Although vapour gas chromatography is routinely used for the analysis of mixtures, this form of chromatography can also be used for separation/purification of substances. This is known as preparative GC. In preparative GC, suitable packed columns are used and as substances emerge from the column, they are collected by condensing the vapour of these separated substances in suitable traps. The carrier gas blows the vapour through these traps hence these traps have to be very efficient. Improved collection of the effluent vaporised fractions in preparative work is attained by strong cooling, increasing the surface of the traps by packing them with glass wool, and by applying an electrical potential which neutralises the charged vapour and causes it to condense.

When the gas chromatograph is attached to a mass spectrometer, a very powerful analytical tool (*gas chromatography-mass spectrometry*; GC-MS) is produced. Vapour gas chromatography allows the analyses of mixtures but does not allow the definitive identification of unknown substances whereas mass spectrometry is good for the identification of a single compound but is less than ideal for the identification of mixtures of

[Previous Page](#)

unfavourable vapour pressures or ease of decomposition, or where super-pure materials are required. It has been used for the latter purpose for purifying anthracene, benzoic acid, chrysene, morphine, 1,8-naphthyridine and pyrene to name a few. [See E.F.G.Herington, *Zone Melting of Organic Compounds*, Wiley & Sons, NY, 1963; W.Pfann, *Zone Melting*, 2nd edn, Wiley, NY, 1966; H.Schildknecht, *Zonenschmelzen*, Verlag Chemie, Weinheim, 1964; W.R.Wilcox, R.Friedenberg et al. *Chem Rev* **64** 187 1964; M.Zief and W.R.Wilcox (Eds), *Fractional Solidification*, Vol I, M Dekker Inc. NY, 1967.]

SUBLIMATION

Sublimation differs from ordinary distillation because the vapour condenses to a solid instead of a liquid. Usually, the pressure in the heated system is diminished by pumping, and the vapour is condensed (after travelling a relatively short distance) onto a cold finger or some other cooled surface. This technique, which is applicable to many organic solids, can also be used with inorganic solids such as aluminium chloride, ammonium chloride, arsenious oxide and iodine. In some cases, passage of a stream of inert gas over the heated substance secures adequate vapourisation. This procedure has the added advantage of removing occluded solvent used in recrystallising the solid.

CHROMATOGRAPHY

Chromatography is often used with advantage for the purification of small amounts of complex organic mixtures. Chromatography techniques all rely on the differential distribution of the various components in a mixture between the mobile phase and the stationary phase. The mobile phase can either be a gas or a liquid whereas the stationary phase can either be a solid or a liquid.

The major chromatographic techniques can also be categorised according to the nature of the mobile phase used - vapour phase chromatography for when a gas is the mobile phase and liquid chromatography for when a liquid is the mobile phase.

A very useful catalog for chromatographic products and information relating to chromatography (from gas chromatography to biochromatography) is that produced by Merck, called the ChromBook and the associated compact disk, ChromCircle.

Vapour phase chromatography (GC or gas-liquid chromatography)

The mobile phase in vapour phase chromatography is a gas (e.g. hydrogen, helium, nitrogen or argon) and the stationary phase is a non-volatile liquid impregnated onto a porous material. The mixture to be purified is injected into a heated inlet whereby it is vaporised and taken into the column by the carrier gas. It is separated into its components by partition between the liquid on the porous support and the gas. For this reason vapour-phase chromatography is sometimes referred to as gas-liquid chromatography (g.l.c). Vapour phase chromatography is very useful in the resolution of a mixture of volatile compounds. This type of chromatography uses either packed or capillary columns. Packed columns have internal diameters of 3-5 mm with lengths of 2-6 m. These columns can be packed with a range of materials including firebrick derived materials (chromasorb P, for separation of non polar hydrocarbons) or diatomaceous earth (chromasorb W, for separation of more polar molecules such as acids, amines). Capillary columns have stationary phase bonded to the walls of long capillary tubes. The diameters in capillary columns are less than 0.5 mm and the lengths of these columns can go up to 50 m! These columns have much superior separating powers than the packed columns. Elution times for equivalent resolutions with packed columns can be up to ten times shorter. It is believed that almost any mixture of compounds can be separated using one of the four stationary phases, OV-101, SE-30, OV-17 and Carbowax-20M. The use of capillary columns in gas chromatography for analysis is now routinely carried out. An extensive range of packed and capillary columns is available from chromatographic specialists such as Supelco, Alltech, Hewlett-Packard, Phenomenex etc.

Table 10 shows some typical liquids used for stationary phases in gas chromatography.

Although vapour gas chromatography is routinely used for the analysis of mixtures, this form of chromatography can also be used for separation/purification of substances. This is known as preparative GC. In preparative GC, suitable packed columns are used and as substances emerge from the column, they are collected by condensing the vapour of these separated substances in suitable traps. The carrier gas blows the vapour through these traps hence these traps have to be very efficient. Improved collection of the effluent vaporised fractions in preparative work is attained by strong cooling, increasing the surface of the traps by packing them with glass wool, and by applying an electrical potential which neutralises the charged vapour and causes it to condense.

When the gas chromatograph is attached to a mass spectrometer, a very powerful analytical tool (*gas chromatography-mass spectrometry*; GC-MS) is produced. Vapour gas chromatography allows the analyses of mixtures but does not allow the definitive identification of unknown substances whereas mass spectrometry is good for the identification of a single compound but is less than ideal for the identification of mixtures of

compounds. This means that with GC-MS, both separation *and* identification of substances in mixtures can be achieved. Because of the relatively small amounts of material required for mass spectrometry, a splitting system is inserted between the column and the mass spectrometer. This enables only a small fraction of the effluent to enter the spectrometer, the rest of the effluent is usually collected or vented to the air.

Liquid chromatography

In contrast to vapour phase chromatography, the mobile phase in liquid chromatography is a liquid. In general, there are four main types of liquid chromatography: *adsorption, partition, ion-chromatography, and gel filtration.*

Adsorption chromatography is based on the difference in the extent to which substances in solution are adsorbed onto a suitable surface. The main techniques in adsorption chromatography are TLC (Thin Layer Chromatography), paper and column chromatography.

Thin layer chromatography (TLC). In thin layer chromatography, the mobile phase i.e. the solvent, creeps up the stationary phase (the adsorbent) by capillary action. The adsorbent (e.g. silica, alumina, cellulose) is spread on a rectangular glass plate (or solid inert plastic sheet or aluminium foil). Some adsorbents (e.g. silica) are mixed with a setting material (e.g. CaSO_4) by the manufacturers which causes the film to set hard on drying. The adsorbent can be activated by heating at $100\text{--}110^\circ$ for a few hours. Other adsorbents (e.g. celluloses) adhere on glass plates without a setting agent. Thus some grades of adsorbents have prefixes e.g. prefix G means that the adsorbent can cling to a glass plate and is used for TLC (e.g. silica gel GF₂₅₄ is for TLC plates which have a dye that fluoresces under 254nm UV light). Those lacking this binder have the letter H after any coding and is suitable for column chromatography e.g. silica gel 60H. The materials to be purified or separated are spotted in a solvent close to the lower end of the plate and allowed to dry. The spots will need to be placed at such a distance so as to ensure that when the lower end of the plate is immersed in the solvent, the spots are a few mm above the eluting solvent. The plate is placed upright in a tank containing the eluting solvent. Elution is carried out in a closed tank to ensure equilibrium. Good separations can be achieved with square plates if a second elution is performed at right angles to the first using a second solvent system. For rapid work, plates of the size of microscopic slides or even smaller are used which can decrease the elution time and cost without loss of resolution. The advantage of plastic backed and aluminium foil backed plates is that the size of the plate can be made as required by cutting the sheet with scissors or a sharp guillotine. Visualisation of substances on TLC can be carried out using UV light if they are UV absorbing or fluorescing substances or by spraying or dipping the plate with a reagent that gives coloured products with the substance (e.g. iodine solution or vapour gives brown colours with amines), or with dilute sulfuric acid (organic compounds become coloured or black when the plates are heated at 100° if the plates are of alumina or silica, but not cellulose). (see Table 11 for some methods of visualisation.) Some alumina and silica powders are available with fluorescent materials in them, in which case the whole plate fluoresces under UV light. Non-fluorescing spots are thus clearly visible, and fluorescent spots invariably fluoresce with a different colour. The colour of the spots can be different under UV light at 254nm and at 365nm. Another useful way of showing up non-UV absorbing spots is to spray the plate with a 1-2% solution of Rhodamine 6G in acetone. Under UV light the dye fluoresces and reveals the non-fluorescing spots. For preparative work, if the material in the spot or fraction is soluble in ether or petroleum ether, the desired substance can be extracted from the adsorbent with these solvents which leave the water soluble dye behind.

TLC can be used as an analytical technique, or as a guide to establishing conditions for column chromatography or as a preparative technique in its own right.

The thickness of the adsorbent on the TLC plates could be between 0.2mm to 2mm or more. In preparative work, the thicker plates are used and hundreds of milligrams of mixtures can be purified conveniently and quickly. The spots or areas are easily scraped off the plates and the desired substances extracted from the adsorbent with the required solvent. For preparative TLC, non destructive methods for visualising spots and fractions are required. As such, the use of UV light is very useful. If substances are not UV active, then a small section of the plate (usually the right or left edge of the plate) is sprayed with a visualising agent while the remainder of the plate is kept covered.

Thin layer chromatography has been used successfully with ion-exchange celluloses as stationary phases and various aqueous buffers as mobile phases. Also, gels (e.g. Sephadex G-50 to G-200 superfine) have been adsorbed on glass plates and are good for fractionating substances of high molecular weights (1500 to 250,000). With this technique, which is called *thin layer gel filtration (TLG)*, molecular weights of proteins can be determined when suitable markers of known molecular weights are run alongside (see Chapter 6).

Commercially available pre-coated plates with a variety of adsorbents are generally very good for quantitative work because they are of a standard quality. Plates of a standardised silica gel 60 (as medium porosity silica gel with a mean porosity of 6mm) released by Merck have a specific surface of $500\text{ m}^2/\text{g}$ and a specific pore volume of 0.75 mL/g . They are so efficient that they have been called *high performance thin layer chromatography (HPTLC)* plates (Ropphahn and Halpap *J Chromatogr* **112** 81 1975). In another variant of thin layer chromatography the

adsorbent is coated with an oil as in gas chromatography thus producing *reverse-phase thin layer chromatography*. Reversed-phase TLC plates e.g. silica gel RP-18 are available from Fluka and Merck.

A very efficient form of chromatography makes use of a circular glass plate (rotor) coated with an adsorbent (silica, alumina or cellulose). As binding to a rotor is needed, the sorbents used may be of a special quality and/or binders are added to the sorbent mixtures. For example when silica gel is required as the adsorbent, silica gel 60 PF-254 with calcium sulfate (Merck catalog 7749) is used. The thickness of the adsorbent (1, 2 or 4 mm) can vary depending on the amount of material to be separated. The apparatus is called a **Chromatotron** (available from Harrison Research, USA). The glass plate is rotated by a motor, and the sample followed by the eluting solvent is allowed to drip onto a central position on the plate. As the plate rotates the solvent elutes the mixture, centrifugally, while separating the components in the form of circular bands radiating from the central point. The separated bands are usually visualised conveniently by UV and as the bands approach the edge of the plate, the eluent is collected. The plate with the adsorbent can be re-used many times if care is employed in the usage, and hence this form of chromatography utilises less adsorbents as well as solvents.

Recipes and instructions for coating the rotors are available from the Harrison website (<http://pw1.netcom.com/~ithres/harrisonresearch.html>). In addition, information on how to regenerate the sorbents and binders are also included.

Paper chromatography. This is the technique from which thin layer chromatography developed. It uses cellulose paper (filter paper) instead of the TLC adsorbent and does not require a backing like the plastic sheet in TLC. It is used in the **ascending procedure** (like in TLC) whereby a sheet of paper is hung in a jar, the materials to be separated are spotted (after dissolving in a suitable solvent and drying) near the bottom of the sheet which dips into the eluting solvent just below the spot. As the solvent rises up the paper the spots are separated according to their adsorption properties. A variety of solvents can be used, the sheet is then dried in air (fume cupboard), and can then be run again with the solvent running at right angles to the first run to give a two dimensional separation. The spots can then be visualised as in TLC or can be cut out and analysed as required. A **descending procedure** had also been developed where the material to be separated is spotted near the top of the paper and the top end is made to dip into a tray containing the eluting solvent. The whole paper is placed in a glass jar and the solvent then runs down the paper causing the materials in the spots to separate also according to their adsorption properties and to the eluting ability of the solvent. This technique is much cheaper than TLC and is still used (albeit with thicker cellulose paper) with considerable success for the separation of protein hydrolysates for sequencing analysis and/or protein identification.

Column Chromatography. The substances to be purified are usually placed on the top of the column and the solvent is run down the column. Fractions are collected and checked for compounds using TLC (UV and/or other means of visualisation). The adsorbent for chromatography can be packed dry and solvents to be used for chromatography are used to equilibrate the adsorbent by flushing the column several times until equilibration is achieved. Alternatively, the column containing the adsorbent is packed wet (slurry method) and pressure is applied at the top of the column until the column is well packed (i.e. the adsorbent is settled).

Graded Adsorbents and Solvents. Materials used in columns for adsorption chromatography are grouped in Table 12 in an approximate order of effectiveness. Other adsorbents sometimes used include barium carbonate, calcium sulfate, calcium phosphate, charcoal (usually mixed with Kieselguhr or other form of diatomaceous earth, for example, the filter aid Celite) and cellulose. The alumina can be prepared in several grades of activity (see below).

In most cases, adsorption takes place most readily from non-polar solvents, such as petroleum ether and least readily from polar solvents such as alcohols, esters, and acetic acid. Common solvents, arranged in approximate order of increasing eluting ability are also given in Table 12. Eluting power roughly parallels the dielectric constants of solvents. The series also reflects the extent to which the solvent binds to the column material, thereby displacing the substances that are already adsorbed. This preference of alumina and silica gel for polar molecules explains, for example, the use of percolation through a column of silica gel for the following purposes-drying of ethylbenzene, removal of aromatics from 2,4-dimethylpentane and of ultraviolet absorbing substances from cyclohexane.

Mixed solvents are intermediate in strength, and so provide a finely graded series. In choosing a solvent for use as an eluent it is necessary to consider the solubility of the substance in it, and the ease with which it can subsequently be removed.

Preparation and Standardisation of Alumina. The activity of alumina depends inversely on its water content, and a sample of poorly active material can be rendered more active by leaving for some time in a round bottomed flask heated up to about 200° in an oil bath or a heating mantle while a slow stream of a dry inert gas is passed through it. Alternatively, it is heated to red heat (380-400°) in an open vessel for 4-6h with

occasional stirring and then cooled in a vacuum desiccator: this material is then of grade I activity. Conversely, alumina can be rendered less active by adding small amounts of water and thoroughly mixing for several hours. Addition of about 3% (w/w) of water converts grade I alumina to grade II.

Used alumina can be regenerated by repeated extraction, first with boiling methanol, then with boiling water, followed by drying and heating. The degree of activity of the material can be expressed conveniently in terms of the scale due to Brockmann and Schodder (*Chem Ber B 74 73 1941*).

Alumina is normally slightly alkaline. A (less strongly adsorbing) neutral alumina can be prepared by making a slurry in water and adding 2M hydrochloric acid until the solution is acid to Congo red. The alumina is then filtered off, washed with distilled water until the wash water gives only a weak violet colour with Congo red paper, and dried.

Alumina used in TLC can be recovered by washing in ethanol for 48h with occasional stirring, to remove binder material and then washed with successive portions of ethyl acetate, acetone and finally with distilled water. Fine particles are removed by siphoning. The alumina is first suspended in 0.04M acetic acid, then in distilled water, siphoning off 30 minutes after each wash. The process is repeated 7-8 times. It is then dried and activated at 200° [Vogh and Thomson *Anal Chem 53 1365 1981*].

Preparation of other adsorbents

Silica gel can be prepared from commercial water-glass by diluting it with water to a density of 1.19 and, while keeping it cooled to 5°, adding concentrated hydrochloric acid with stirring until the solution is acid to thymol blue. After standing for 3h, the precipitate is filtered off, washed on a Büchner funnel with distilled water, then suspended in 0.2M hydrochloric acid. The suspension is set aside for 2-3 days, with occasional stirring, then filtered, washed well with water and dried at 110°. It can be activated by heating up to about 200° as described for alumina.

Powdered commercial silica gel can be purified by suspending and standing overnight in concentrated hydrochloric acid (6mL/g), decanting the supernatant and repeating with fresh acid until the latter remains colourless. After filtering with suction on a sintered-glass funnel, the residue is suspended in water and washed by decantation until free of chloride ions. It is then filtered, suspended in 95% ethanol, filtered again and washed on the filter with 95% ethanol. The process is repeated with anhydrous diethyl ether before the gel is heated for 24h at 100° and stored for another 24h in a vacuum desiccator over phosphorus pentoxide.

To buffer silica gel for flash chromatography (see later), 200g of silica is stirred in 1L of 0.2M NaH₂PO₄ for 30 minutes. The slurry is then filtered with suction using a sintered glass funnel. The silica gel is then activated at 110°C for 16 hours. The pH of the resulting silica gel is ~4. Similar procedures can be utilized to buffer the pH of the silica gel at various pHs (up to pH ~8: pH higher than this causes degradation of silica) using appropriate phosphate buffers.

Commercial silica gel has also been purified by suspension of 200g in 2L of 0.04M ammonia, allowed to stand for 5min before siphoning off the supernatant. The procedure was repeated 3-4 times, before rinsing with distilled water and drying, and activating the silica gel in an oven at 110° [Vogh and Thomson, *Anal Chem 53 1345 1981*].

Although silica gel is not routinely recycled after use (due to fear of contamination as well as the possibility of reduced activity), the costs of using new silica gel for purification may be prohibitive. In these cases, recycling may be achieved by stirring the used silica gel (1 kg) in a mixture of methanol and water (2L MeOH/4L water) for 30-40 mins. The silica gel is filtered (as described above) and reactivated at 110°C for 16 hours.

Diatomaceous earth (Celite 535 or 545, Hyflo Super-cel, Dicalite, Kieselguhr) is purified before use by washing with 3M hydrochloric acid, then water, or it is made into a slurry with hot water, filtered at the pump and washed with water at 50° until the filtrate is no longer alkaline to litmus. Organic materials can be removed by repeated extraction at 50° with methanol or chloroform, followed by washing with methanol, filtering and drying at 90-100°.

Charcoal is generally satisfactorily activated by heating gently to red heat in a crucible or quartz beaker in a muffle furnace, finally allowing to cool under an inert atmosphere in a desiccator. Good commercial activated charcoal is made from wood, e.g. *Norit* (from Birch wood), *Darco* and *Nuchar*. If the cost is important then the cheaper *animal charcoal* (bone charcoal) can be used. However, this charcoal contains calcium phosphate and other calcium salts and cannot be used with acidic materials. In this case the charcoal is boiled with dilute hydrochloric acid (1:1 by volume) for 2-3h, diluted with distilled water and filtered through a fine grade paper on a Büchner flask, washed with distilled water until the filtrate is almost neutral, and dried first in air then in a vacuum, and activated as above. To improve the porosity, charcoal columns are usually prepared in admixture with diatomaceous earth.

Cellulose for chromatography is purified by sequential washing with chloroform, ethanol, water, ethanol, chloroform and acetone. More extensive purification uses aqueous ammonia, water, hydrochloric acid, water, acetone and diethyl ether, followed by drying in a vacuum. Trace metals can be removed from filter paper by washing for several hours with 0.1M oxalic or citric acid, followed by repeated washing with distilled water.

Flash Chromatography

A faster method of separating components of a mixture is *flash chromatography* (see Still et al. *J Org Chem* **43** 2923 1978). In flash chromatography the eluent flows through the column under a pressure of *ca* 1 to 4 atmospheres. The lower end of the chromatographic column has a relatively long taper closed with a tap. The upper end of the column is connected through a ball joint to a tap. Alternatively a specially designed chromatographic column with a solvent reservoir can also be used (for an example, see the Aldrich Chemical Catalog-glassware section). The tapered portion is plugged with cotton, or quartz, wool and *ca* 1 cm of fine washed sand (the latter is optional). The adsorbent is then placed in the column as a dry powder or as a slurry in a solvent and allowed to fill to about one third of the column. A fine grade of adsorbent is required in order to slow the flow rate at the higher pressure, e.g. Silica 60, 230 to 400 mesh with particle size 0.040-0.063mm (from Merck). The top of the adsorbent is layered with *ca* 1 cm of fine washed sand. The mixture in the smallest volume of solvent is applied at the top of the column and allowed to flow into the adsorbent under gravity by opening the lower tap momentarily. The top of the column is filled with eluent, the upper tap is connected by a tube to a nitrogen supply from a cylinder, or to compressed air, and turned on to the desired pressure (monitor with a gauge). The lower tap is turned on and fractions are collected rapidly until the level of eluent has reached the top of the adsorbent (do not allow the column to run dry). If further elution is desired then both taps are turned off, the column is filled with more eluting solvent and the process repeated. The top of the column can be modified so that gradient elution can be performed. Alternatively, an apparatus for producing the gradient is connected to the upper tap by a long tube and placed high above the column in order to produce the required hydrostatic pressure. Flash chromatography is more efficient and gives higher resolution than conventional chromatography at atmospheric pressure and is completed in a relatively shorter time. A successful separation of components of a mixture by TLC using the same adsorbent is a good indication that flash chromatography will give the desired separation on a larger scale.

Paired-ion Chromatography (PIC)

Mixtures containing ionic compounds (e.g. acids and/or bases), non-ionisable compounds, and zwitterions, can be separated successfully by paired-ion chromatography (PIC). It utilises the 'reverse-phase' technique (Eksberg and Schill *Anal Chem* **45** 2092 1973). The stationary phase is lipophilic, such as μ -BONDAPAK C₁₈ or any other adsorbent that is compatible with water. The mobile phase is water or aqueous methanol containing the acidic or basic counter ion. Thus the mobile phase consists of dilute solutions of strong acids (e.g. 5mM 1-heptanesulfonic acid) or strong bases (e.g. 5 mM tetrabutylammonium phosphate) that are completely ionised at the operating pH values which are usually between 2 and 8. An equilibrium is set up between the neutral species of a mixture in the stationary phase and the respective ionised (anion or cation) species which dissolve in the mobile phase containing the counter ions. The extent of the equilibrium will depend on the ionisation constants of the respective components of the mixture, and the solubility of the unionised species in the stationary phase. Since the ionisation constants and the solubility in the stationary phase will vary with the water-methanol ratio of the mobile phase, the separation may be improved by altering this ratio gradually (gradient elution) or stepwise. If the compounds are eluted too rapidly the water content of the mobile phase should be increased, e.g. by steps of 10%. Conversely, if components do not move, or move slowly, the methanol content of the mobile phase should be increased by steps of 10%.

The application of pressure to the liquid phase in liquid chromatography generally increases the separation (see HPLC). Also in PIC improved efficiency of the column is observed if pressure is applied to the mobile phase (Wittmer, Nuessle and Haney *Anal Chem* **47** 1422 1975).

Ion-exchange Chromatography

Ion-exchange chromatography involves an electrostatic process which depends on the relative affinities of various types of ions for an immobilised assembly of ions of opposite charge. The stationary phase is an aqueous buffer with a fixed pH or an aqueous mixture of buffers in which the pH is continuously increased or decreased as the separation may require. This form of liquid chromatography can also be performed at high inlet pressures of liquid with increased column performances.

Ion-exchange Resins. An ion-exchange resin is made up of particles of an insoluble elastic hydrocarbon network to which is attached a large number of ionisable groups. Materials commonly used comprise synthetic ion-exchange resins made, for example, by crosslinking polystyrene to which has been attached non-

diffusible ionised or ionisable groups. Resins with relatively high crosslinkage (8-12%) are suitable for the chromatography of small ions, whereas those with low cross linkage (2-4%) are suitable for larger molecules. Applications to hydrophobic systems are possible using aqueous gels with phenyl groups bound to the rigid matrix (Phenyl-Superose/Sepharose, Pharmacia-Amersham Biosciences) or neopentyl chains (Alkyl-Superose, Pharmacia-Amersham Biosciences). (Superose is a cross-linked agarose-based medium with an almost uniform bead size.) These groups are further distinguishable as strong $[-\text{SO}_2\text{OH}, -\text{NR}_3^+]$ or weak $[-\text{OH}, -\text{CO}_2\text{H}, -\text{PO}(\text{OH})_2, -\text{NH}_2]$. Their charges are counterbalanced by diffusible ions, and the operation of a column depends on its ability and selectivity to replace these ions. The exchange that takes place is primarily an electrostatic process but adsorptive forces and hydrogen bonding can also be important. A typical sequence for the relative affinities of some common anions (and hence the inverse order in which they pass through such a column), is the following, obtained using a quaternary ammonium (strong base) anion-exchange column:

Fluoride < acetate < bicarbonate < hydroxide < formate < chloride < bromate < nitrite < cyanide < bromide < chromate < nitrate < iodide < thiocyanate < oxalate < sulfate < citrate.

For an amine (weak base) anion-exchange column in its chloride form, the following order has been observed:

Fluoride < chloride < bromide = iodide = acetate < molybdate < phosphate < arsenate < nitrate < tartrate < citrate < chromate < sulfate < hydroxide.

With strong cation-exchangers (e.g. with SO_3H groups), the usual sequence is that polyvalent ions bind more firmly than mono- or di- valent ones, a typical series being as follows:

$\text{Th}^{4+} > \text{Fe}^{3+} > \text{Al}^{3+} > \text{Ba}^{2+} > \text{Pb}^{2+} > \text{Sr}^{2+} > \text{Ca}^{2+} > \text{Co}^{2+} > \text{Ni}^{2+} = \text{Cu}^{2+} > \text{Zn}^{2+} = \text{Mg}^{2+} > \text{UO}_2^+ = \text{Mn}^{2+} > \text{Ag}^+ > \text{Tl}^+ > \text{Cs}^+ > \text{Rb}^+ > \text{NH}_4^+ = \text{K}^+ > \text{Na}^+ > \text{H}^+ > \text{Li}^+.$

Thus, if an aqueous solution of a sodium salt contaminated with heavy metals is passed through the sodium form of such a column, the heavy metal ions will be removed from the solution and will be replaced by sodium ions from the column. This effect is greatest in dilute solution. Passage of sufficiently strong solutions of alkali metal salts or mineral acids readily displaces all other cations from ion-exchange columns. (The regeneration of columns depends on this property.) However, when the cations lie well to the left in the above series it is often advantageous to use a complex-forming species to facilitate removal. For example, iron can be displaced from ion-exchange columns by passage of sodium citrate or sodium ethylenediaminetetraacetate.

Some of the more common commercially available resins are listed in Table 13.

Ion-exchange resins swell in water to an extent which depends on the amount of crosslinking in the polymer, so that columns should be prepared from the wet material by adding it as a suspension in water to a tube already partially filled with water. (This also avoids trapping air bubbles.) The exchange capacity of a resin is commonly expressed as mg equiv./mL of wet resin. This quantity is pH-dependent for weak-acid or weak-base resins but is constant at about 0.6-2 for most strong-acid or strong-base types.

Apart from their obvious applications to inorganic species, sulfonic acid resins have been used in purifying amino acids, aminosugars, organic acids, peptides, purines, pyrimidines, nucleosides, nucleotides and polynucleotides. Thus, organic bases can be applied to the H^+ form of such resins by adsorbing them from neutral solution and, after washing with water, they are eluted sequentially with suitable buffer solutions or dilute acids. Alternatively, by passing alkali solution through the column, the bases will be displaced in an order that is governed by their pK values. Similarly, strong-base anion exchangers have been used for aldehydes and ketones (as bisulfite addition compounds), carbohydrates (as their borate complexes), nucleosides, nucleotides, organic acids, phosphate esters and uronic acids. Weakly acidic and weakly basic exchange resins have also found extensive applications, mainly in resolving weakly basic and acidic species. For demineralisation of solutions without large changes in pH, mixed-bed resins can be prepared by mixing a cation-exchange resin in its H^+ form with an anion-exchange resin in its OH^- form. Commercial examples include Amberlite MB-1 (IR-120 + IRA-400) and Bio-Deminrolit (Zeo-Karb 225 and Zerolit FF). The latter is also available in a self-indicating form.

Ion-exchange Celluloses and Sephadex. A different type of ion-exchange column that finds extensive application in biochemistry for the purification of proteins, nucleic acids and acidic polysaccharides derives from cellulose by incorporating acidic and basic groups to give ion-exchangers of controlled acid and basic strengths. Commercially available cellulose-type resins are given in Tables 14 and 15. AG 501 x 8 (Bio-Rad) is a mixed-bed resin containing equivalents of AG 50W-x8 H^+ form and AG 1-x8 HO^- form, and Bio-Rex MSZ 501 resin. A dye marker indicates when the resin is exhausted. Removal of unwanted cations, particularly of the transition metals, from amino acids and buffer can be achieved by passage of the solution through a column of Chelex 20 or Chelex 100. The metal-chelating abilities of the resin reside in the bonded iminodiacetate groups.

Chelex can be regenerated by washing in two bed volumes of 1M HCl, two bed volumes of 1M NaOH and five bed volumes of water.

Ion-exchange celluloses are available in different particle sizes. It is important that the amounts of 'fines' are kept to a minimum otherwise the flow of liquid through the column can be extremely slow to the point of no liquid flow. Celluloses with a large range of particle sizes should be freed from 'fines' before use. This is done by suspending the powder in the required buffer and allowing it to settle for one hour and then decanting the 'fines'. This separation appears to be wasteful but it is necessary for reasonable flow rates without applying high pressures at the top of the column. Good flow rates can be obtained if the cellulose column is packed dry whereby the 'fines' are evenly distributed throughout the column. Wet packing causes the 'fines' to rise to the top of the column, which thus becomes clogged.

Several ion-exchange celluloses require recycling before use, a process which must be applied for recovered celluloses. Recycling is done by stirring the cellulose with 0.1M aqueous sodium hydroxide, washing with water until neutral, then suspending in 0.1M hydrochloric acid and finally washing with water until neutral. When regenerating a column it is advisable to wash with a salt solution (containing the required counter ions) of increasing ionic strength up to 2M. The cellulose is then washed with water and recycled if necessary. Recycling can be carried out more than once if there are doubts about the purity of the cellulose and when the cellulose had been used previously for a different purification procedure than the one to be used. The basic matrix of these ion-exchangers is cellulose and it is important not to subject them to strong acid (> 1M) and strongly basic (> 1M) solutions.

When storing ion-exchange celluloses, or during prolonged usage, it is important to avoid growth of microorganisms or moulds which slowly destroy the cellulose. Good inhibitors of microorganisms are phenyl mercuric salts (0.001%, effective in weakly alkaline solutions), chlorohexidine (Hibitane at 0.002% for anion exchangers), 0.02% aqueous sodium azide or 0.005% of ethyl mercuric thiosalicylate (Merthiolate) are most effective in weakly acidic solutions for cation exchangers. Trichlorobutanol (Chloretone, at 0.05% is only effective in weakly acidic solutions) can be used for both anion and cation exchangers. Most organic solvents (e.g. methanol) are effective antimicrobial agents but only at high concentrations. These inhibitors must be removed by washing the columns thoroughly before use because they may have adverse effects on the material to be purified (e.g. inactivation of enzymes or other active preparations).

Sephadex. Other carbohydrate matrices such as *Sephadex* (based on dextran) have more uniform particle sizes. Their advantages over the celluloses include faster and more reproducible flow rates and they can be used directly without removal of 'fines'. *Sephadex*, which can also be obtained in a variety of ion-exchange forms (see Table 15) consists of beads of a cross-linked dextran gel which swells in water and aqueous salt solutions. The smaller the bead size, the higher the resolution that is possible but the slower the flow rate. Typical applications of *Sephadex* gels are the fractionation of mixtures of polypeptides, proteins, nucleic acids, polysaccharides and for desalting solutions.

Sephadex is a bead form of cross-linked dextran gel. *Sepharose CL* and *Bio-Gel A* are derived from agarose (see below). *Sephadex* ion-exchangers, unlike celluloses, are available in narrow ranges of particle sizes. These are of two medium types, the G-25 and G-50, and their dry bead diameter sizes are *ca* 50 to 150 microns. They are available as cation and anion exchange *Sephadex*. One of the disadvantages of using *Sephadex* ion-exchangers is that the bed volume can change considerably with alteration of pH. *Ultragels* also suffer from this disadvantage to a varying extent, but ion-exchangers of the bead type have been developed e.g. *Fractogels*, *Toyopearl*, which do not suffer from this disadvantage.

Sepharose (e.g. *Sepharose CL* and *Bio-Gel A*) is a bead form of agarose gel which is useful for the fractionation of high molecular weight substances, for molecular weight determinations of large molecules (molecular weight > 5000), and for the immobilisation of enzymes, antibodies, hormones and receptors usually for affinity chromatography applications.

In preparing any of the above for use in columns, the dry powder is evacuated, then mixed under reduced pressure with water or the appropriate buffer solution. Alternatively it is stirred gently with the solution until all air bubbles are removed. Because some of the wet powders change volumes reversibly with alteration of pH or ionic strength (see above), it is imperative to make allowances when packing columns (see above) in order to avoid overflowing of packing when the pH or salt concentrations are altered.

Cellex CM ion-exchange cellulose can be purified by treatment of 30-40g (dry weight) with 500mL of 1mM cysteine hydrochloride. It is then filtered through a Büchner funnel and the filter cake is suspended in 500mL of 0.05M NaCl/0.5M NaOH. This is filtered and the filter cake is resuspended in 500ml of distilled water and filtered again. The process is repeated until the washings are free from chloride ions. The filter cake is again suspended in 500mL of 0.01M buffer at the desired pH for chromatography, filtered, and the last step repeated several times.

Cellex D and other anionic celluloses are washed with 0.25M NaCl/0.25M NaOH solution, then twice with deionised water. This is followed with 0.25M NaCl and then washed with water until chloride-free. The Cellex is then equilibrated with the desired buffer as above.

Crystalline Hydroxylapatite is a structurally organised, highly polar material which, in aqueous solution (in buffers) strongly adsorbs macromolecules such as proteins and nucleic acids, permitting their separation by virtue of the interaction with charged phosphate groups and calcium ions, as well by physical adsorption. The procedure therefore is not entirely ion-exchange in nature. Chromatographic separations of singly and doubly stranded DNA are readily achievable whereas there is negligible adsorption of low molecular weight species.

Gel Filtration

The gel-like, bead nature of wet Sephadex enables small molecules such as inorganic salts to diffuse freely into it while, at the same time, protein molecules are unable to do so. Hence, passage through a Sephadex column can be used for complete removal of salts from protein solutions. Polysaccharides can be freed from monosaccharides and other small molecules because of their differential retardation. Similarly, amino acids can be separated from proteins and large peptides.

Gel filtration using Sephadex G-types (50 to 200) is essentially useful for fractionation of large molecules with molecular weights above 1000. For Superose, the range is given as 5000 to 5×10^6 . Fractionation of lower molecular weight solutes (e.g. ethylene glycols, benzyl alcohols) can now be achieved with Sephadex G-10 (up to Mol.Wt 700) and G-25 (up to Mol.Wt 1500). These dextrans are used only in aqueous solutions. In contrast, Sephadex LH-20 and LH-60 (prepared by hydroxypropylation of Sephadex) are used for the separation of small molecules (Mol.Wt less than 500) using most of the common organic solvents as well as water.

Sephasorb HP (ultrafine, prepared by hydroxypropylation of crossed-linked dextran) can also be used for the separation of small molecules in organic solvents and water, and in addition it can withstand pressures up to 1400 psi making it useful in HPLC. These gels are best operated at pH values between 2 and 12, because solutions with high and low pH values slowly decompose them (see further in Chapter 6).

High Performance Liquid Chromatography (HPLC)

When pressure is applied at the inlet of a liquid chromatographic column the performance of the column can be increased by several orders of magnitude. This is partly because of the increased speed at which the liquid flows through the column and partly because fine column packings which have larger surface areas can be used. Because of the improved efficiency of the columns, this technique has been referred to as high performance, high pressure, or high speed liquid chromatography and has found great importance in chemistry and biochemistry.

The equipment consists of a hydraulic system to provide the pressure at the inlet of the column, a column, a detector, data storage and output, usually in the form of a computer. The pressures used in HPLC vary from a few psi to 4000-5000 psi. The most convenient pressures are, however, between 500 and 1800psi. The plumbing is made of stainless steel or non-corrosive metal tubing to withstand high pressures. Plastic tubing and connectors are used for low pressures, e.g. up to ~500psi. Increase of temperature has a very small effect on the performance of a column in liquid chromatography. Small variations in temperatures, however, do upset the equilibrium of the column, hence it is advisable to place the column in an oven at ambient temperature in order to achieve reproducibility. The packing (stationary phase) is specially prepared for withstanding high pressures. It may be an adsorbent (for adsorption or solid-liquid HPLC), a material impregnated with a high boiling liquid (e.g. octadecyl sulfate, in *reverse-phase* or *liquid-liquid* or *paired-ion* HPLC), an ion-exchange material (in *ion-exchange* HPLC), or a highly porous non-ionic gel (for high performance *gel filtration*). The mobile phase is water, aqueous buffers, salt solutions, organic solvents or mixtures of these. The more commonly used detectors have UV, visible, diode array or fluorescence monitoring for light absorbing substances, and refractive index monitoring and evaporative light scattering for transparent compounds. UV detection is not useful when molecules do not have UV absorbing chromophores and solvents for elution should be carefully selected when UV monitoring is used so as to ensure the lack of interference in detection. The sensitivity of the refractive index monitoring is usually lower than the light absorbing monitoring by a factor of ten or more. It is also difficult to use a refractive index monitoring system with gradient elution of solvents. When substances have readily oxidised and reduced forms, e.g. phenols, nitro compounds, heterocyclic compounds etc, then electrochemical detectors are useful. These detectors oxidise and reduce these substances and make use of this process to provide a peak on the recorder.

The cells of the monitoring devices are very small (*ca* 5 μ l) and the detection is very good. The volumes of the analytical columns are quite small (*ca* 2mL for a 1 metre column) hence the result of an analysis is achieved very quickly. Larger columns have been used for preparative work and can be used with the same equipment. Most

machines have solvent mixing chambers for solvent gradient or ion gradient elution. The solvent gradient (for two solvents) or pH or ion gradient can be adjusted in a linear, increasing or decreasing exponential manner.

In general two different types of HPLC columns are available. Prepacked columns are those with metal casings with threads at both ends onto which capillary connections are attached. The cartridge HPLC columns are cheaper and are used with cartridge holders. As the cartridge is fitted with a groove for the holding device, no threads are necessary and the connection pieces can be reused. A large range of HPLC columns (including guard columns, i.e. small pre-columns) are available from Alltech, Supelco (see www.sigmaldrich.com), Waters (www.waters.com), Agilent Technologies (www.chem.agilent.com), Phenomenex (www.phenomenex.com), YMC (www.ymc.co.jp/en/), Merck (www.merck.de), SGE (www.sge.com) and other leading companies. Included in this range of columns are also columns with chiral bonded phases capable of separating enantiomeric mixtures, such as Chiralpak AS and Chirex™ columns (e.g. from Restek-www.restekcorp.com, Daicel-www.daicel.co.jp/indexe.html).

HPLC systems coupled to mass spectrometers (LC-MS) are extremely important methods for the separation and identification of substances. If not for the costs involved in LC-MS, these systems would be more commonly found in research laboratories.

Other Types of Liquid Chromatography

New stationary phases for specific purposes in chromatographic separation are being continually proposed. *Charge transfer adsorption chromatography* makes use of a stationary phase which contains immobilised aromatic compounds and permits the separation of aromatic compounds by virtue of the ability to form charge transfer complexes (sometimes coloured) with the stationary phase. The separation is caused by the differences in stability of these complexes (Porath and Dahlgren-Caldwell *J Chromatogr* **133** 180 1977).

In *metal chelate adsorption chromatography* a metal is immobilised by partial chelation on a column which contains bi- or tri- dentate ligands. Its application is in the separation of substances which can complex with the bound metals and depends on the stability constants of the various ligands (Porath, Carlsson, Olsson and Belfrage *Nature* **258** 598 1975; Loennerdal, Carlsson and Porath *FEBS Lett* **75** 89 1977).

An application of chromatography which has found extensive use in biochemistry and has brought a new dimension in the purification of enzymes is *affinity chromatography*. A specific enzyme inhibitor is attached by covalent bonding to a stationary phase (e.g. AH-Sepharose 4B for acidic inhibitors and CH-Sepharose 4B for basic inhibitors), and will strongly bind only the specific enzyme which is inhibited, allowing all other proteins to flow through the column. The enzyme is then eluted with a solution of high ionic strength (e.g. 1M sodium chloride) or a solution containing a substrate or reversible inhibitor of the specific enzyme. (The ionic medium can be removed by gel filtration using a mixed-bed gel.) Similarly, an immobilised lectin may interact with the carbohydrate moiety of a glycoprotein. The most frequently used matrixes are cross-linked (4-6%) agarose and polyacrylamide gel. Many adsorbents are commercially available for nucleotides, coenzymes and vitamins, amino acids, peptides, lectins and related macromolecules and immunoglobulins. Considerable purification can be achieved by one passage through the column and the column can be reused several times.

The affinity method may be *biospecific*, for example as an antibody-antigen interaction, or chemical as in the chelation of boronate by *cis*-diols, or of unknown origin as in the binding of certain dyes to albumin and other proteins.

Hydrophobic adsorption chromatography takes advantage of the hydrophobic properties of substances to be separated and has also found use in biochemistry (Hoftsee *Biochem Biophys Res Commun* **50** 751 1973; Jennissen and Heilmayer Jr *Biochemistry* **14** 754 1975). Specific covalent binding with the stationary phase, a procedure that was called *covalent chromatography*, has been used for separation of compounds and for immobilising enzymes on a support: the column was then used to carry out specific bioorganic reactions (Mosbach *Method Enzymol* **44** 1976; A.Rosevear, J.F.Kennedy and J.M.S.Cabral, *Immobilised Enzymes and Cells: A Laboratory Manual*, Adam Hilger, Bristol, 1987, ISBN 085274515X).

DRYING

Removal of Solvents

Where substances are sufficiently stable, removal of solvent from recrystallised materials presents no problems. The crystals, after filtering at the pump (and perhaps air-drying by suction), are heated in an oven above the boiling point of the solvent (but below this melting point of the crystals), followed by cooling in a desiccator. Where this treatment is inadvisable, it is still often possible to heat to a lower temperature under reduced pressure, for example in an Abderhalden pistol. This device consists of a small chamber which is heated externally by the vapour of a boiling solvent. Inside this chamber, which can be evacuated by a water pump or some other vacuum pump, is

placed a small boat containing the sample to be dried and also a receptacle with a suitable drying agent. Convenient liquids for use as boiling liquids in an Abderhalden pistol, and their temperatures, are given in Table 16. Alternatively an electrically heated drying pistol can also be used. In cases where heating above room temperature cannot be used, drying must be carried out in a vacuum desiccator containing suitable absorbents. For example, hydrocarbons, such as cyclohexane and petroleum ether, can be removed by using shredded paraffin wax, and acetic acid and other acids can be absorbed by pellets of sodium or potassium hydroxide. However, in general, solvent removal is less of a problem than ensuring that the water content of solids and liquids is reduced below an acceptable level.

Removal of Water

Methods for removing water from solids depends on the thermal stability of the solids or the time available. The safest way is to dry in a vacuum desiccator over concentrated sulfuric acid, phosphorus pentoxide, silica gel, calcium chloride, or some other desiccant. Where substances are stable in air and melt above 100°, drying in an air oven may be adequate. In other cases, use of an Abderhalden pistol may be satisfactory.

Often, in drying inorganic salts, the final material that is required is a hydrate. In such cases, the purified substance is left in a desiccator to equilibrate above an aqueous solution having a suitable water-vapour pressure. A convenient range of solutions used in this way is given in Table 17.

The choice of desiccants for drying liquids is more restricted because of the need to avoid all substances likely to react with the liquids themselves. In some cases, direct distillation of an organic liquid is a suitable method for drying both solids and liquids, especially if low-boiling azeotropes are formed. Examples include acetone, aniline, benzene, chloroform, carbon tetrachloride, heptane, hexane, methanol, nitrobenzene, petroleum ether, toluene and xylene. Addition of benzene can be used for drying ethanol by distillation. In carrying out distillations intended to yield anhydrous products, the apparatus should be fitted with guard-tubes containing calcium chloride or silica gel to prevent entry of moist air into the system. (Many anhydrous organic liquids are appreciably hygroscopic).

Traces of water can be removed from solvents such as benzene, 1,2-dimethoxyethane, diethyl ether, pentane, toluene and tetrahydrofuran by refluxing under nitrogen a solution containing sodium wire and benzophenone, and fractionally distilling. Drying with, and distilling from CaH_2 is applicable to a number of solvents including aniline, benzene, *tert*-butylamine, *tert*-butanol, 2,4,6-collidine, diisopropylamine, dimethylformamide, hexamethylphosphoramide, dichloromethane, ethyl acetate, pyridine, tetramethylethylenediamine, toluene, triethylamine.


Removal of water from gases may be by physical or chemical means, and is commonly by adsorption on to a drying agent in a low-temperature trap. The effectiveness of drying agents depends on the vapour pressure of the hydrated compound - the lower the vapour pressure the less the remaining moisture in the gas.

The most usually applicable of the specific methods for detecting and determining water in organic liquids is due to Karl Fischer. (See J.Mitchell and D.M.Smith, *Aquametry*, 2nd Ed, J Wiley & Sons, New York, 1977-1984, ISBN 0471022640; Fieser and Fieser *Reagents for Organic Synthesis*, J.Wiley & Sons, NY, Vol 1, 528 1967, ISBN 0271616X). Other techniques include electrical conductivity measurements and observation of the temperature at which the first cloudiness appears as the liquid is cooled (applicable to liquids in which water is only slightly soluble). Addition of anhydrous cobalt (II) iodide (blue) provides a convenient method (colour change to pink on hydration) for detecting water in alcohols, ketones, nitriles and some esters. Infrared absorption measurements of the broad band for water near 3500 cm^{-1} can also sometimes be used for detecting water in non-hydroxylic substances.

For further useful information on mineral adsorbents and drying agents, go to the SigmaAldrich website, under technical library (Aldrich) for technical bulletin AL-143.

Intensity and Capacity of Common Desiccants

Drying agents are conveniently grouped into three classes, depending on whether they combine with water reversibly, they react chemically (irreversibly) with water, or they are molecular sieves. The first group vary in their drying intensity with the temperature at which they are used, depending on the vapour pressure of the hydrate that is formed. This is why, for example, drying agents such as anhydrous sodium sulfate, magnesium sulfate or calcium chloride should be filtered off from the liquids before the latter are heated. The intensities of drying agents belonging to this group fall in the sequence:



Where large amounts of water are to be removed, a preliminary drying of liquids is often possible by shaking with concentrated solutions of sodium sulfate or potassium carbonate, or by adding sodium chloride to salt out the organic phase (for example, in the drying of lower alcohols), as long as the drying agent does not react (e.g. CaCl_2 with alcohols and amines, see below).

Drying agents that combine irreversibly with water include the alkali metals, the metal hydrides (discussed in Chapter 2), and calcium carbide.

Suitability of Individual Desiccants

Alumina. (Preheated to 175° for about 7h). Mainly as a drying agent in a desiccator or as a column through which liquid is percolated.

Aluminium amalgam. Mainly used for removing traces of water from alcohols *via* refluxing followed by distillation.

Barium oxide. Suitable for drying organic bases.

Barium perchlorate. Expensive. Used in desiccators (*covered with a metal guard*).

Unsuitable for drying solvents or organic material where contact is necessary, because of the danger of **EXPLOSION**

Boric anhydride. (Prepared by melting boric acid in an air oven at a high temperature, cooling in a desiccator, and powdering.) Mainly used for drying formic acid.

Calcium chloride (anhydrous). Cheap. Large capacity for absorption of water, giving the hexahydrate below 30°, but is fairly slow in action and not very efficient. Its main use is for preliminary drying of alkyl and aryl halides, most esters, saturated and aromatic hydrocarbons and ethers. Unsuitable for drying alcohols and amines (which form addition compounds), fatty acids, amides, amino acids, ketones, phenols, or some aldehydes and esters. Calcium chloride is suitable for drying the following gases: hydrogen, hydrogen chloride, carbon monoxide, carbon dioxide, sulfur dioxide, nitrogen, methane, oxygen, also paraffins, ethers, olefins and alkyl chlorides.

Calcium hydride. See Chapter 2.

Calcium oxide. (Preheated to 700-900° before use.) Suitable for alcohols and amines (but does not dry them completely). Need not be removed before distillation, but in that case the head of the distillation column should be packed with glass wool to trap any calcium oxide powder that might be carried over. Unsuitable for acidic compounds and esters. Suitable for drying gaseous amines and ammonia.

Calcium sulfate (anhydrous). (Prepared by heating the dihydrate or the hemihydrate in an oven at 235° for 2-3h; it can be regenerated.) Available commercially as Drierite. It forms the hemihydrate, $2\text{CaSO}_4 \cdot \text{H}_2\text{O}$, so that its capacity is fairly low (6.6% of its weight of water), and hence is best used on partially dried substances. It is very efficient (being comparable with phosphorus pentoxide and concentrated sulfuric acid). Suitable for most organic compounds. Solvents boiling below 100° can be dried by direct distillation from calcium sulfate.

Copper (II) sulfate (anhydrous). Suitable for esters and alcohols. Preferable to sodium sulfate in cases where solvents are sparingly soluble in water (for example, benzene or toluene).

Lithium aluminium hydride. See Chapter 2.

Magnesium amalgam. Mainly used for removing traces of water from alcohols by refluxing the alcohol in the presence of the Mg amalgam followed by distillation.

Magnesium perchlorate (anhydrous). (Available commercially as Dehydrite. Expensive.) Used in desiccators. Unsuitable for drying solvents or any organic material where contact is necessary, because of the danger of **EXPLOSION**.

Magnesium sulfate (anhydrous). (Prepared from the heptahydrate by drying at 300° under reduced pressure.) More rapid and effective than sodium sulfate but is slightly acidic. It has a large capacity, forming $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ below 48°. Suitable for the preliminary drying of most organic compounds.

Molecular sieves. See later.

Phosphorus pentoxide. Very rapid and efficient, but difficult to handle and should only be used after the organic material has been partially dried, for example with magnesium sulfate. Suitable for anhydrides, alkyl and aryl halides, ethers, esters, hydrocarbons and nitriles, and for use in desiccators. Not suitable with acids, alcohols, amines or ketones, or with organic molecules from which a molecule of water can be eliminated. Suitable for drying the following gases: hydrogen, oxygen, carbon dioxide, carbon monoxide, sulfur dioxide, nitrogen, methane, ethene and paraffins. It is available on a solid support with an indicator under the name *Sicapent* (from Merck). The colour changes in *Sicapent* depend on the percentage of water present (e.g. in the absence of water, *Sicapent* is colorless but becomes green with 20% water and blue with 33% w/w water). When the quantity of water in the desiccator is high a crust of phosphoric acid forms a layer over the phosphorus pentoxide powder and decreases its efficiency. The crust can be removed with a spatula to expose the dry powder and restore the desiccant property.

Potassium (metal). Properties and applications are similar to those for sodium but as the reactivity is greater than that of sodium, the hazards are greater than that of sodium. **Handle with extreme care.**

Potassium carbonate (anhydrous). Has a moderate efficiency and capacity, forming the dihydrate. Suitable for an initial drying of alcohols, bases, esters, ketones and nitriles by shaking with them, then filtering off. Also suitable for salting out water-soluble alcohols, amines and ketones. Unsuitable for acids, phenols, thiols and other acidic substances.

Potassium carbonate. Solid potassium hydroxide is very rapid and efficient. Its use is limited almost entirely to the initial drying of organic bases. Alternatively, sometimes the base is shaken first with a concentrated solution of potassium hydroxide to remove most of the water present. Unsuitable for acids, aldehydes, ketones, phenols, thiols, amides and esters. Also used for drying gaseous amines and ammonia.

Silica gel. Granulated silica gel is a commercially available drying agent for use with gases, in desiccators, and (because of its chemical inertness) in physical instruments (pH meters, spectrometers, balances). Its drying action depends on physical adsorption, so that silica gel must be used at room temperature or below. By incorporating cobalt chloride into the material it can be made self indicating (blue when dry, pink when wet), re-drying in an oven at 110° being necessary when the colour changes from blue to pink.

Sodium (metal). Used as a fine wire or as chips, for more completely drying ethers, saturated hydrocarbons and aromatic hydrocarbons which have been partially dried (for example with calcium chloride or magnesium sulfate). Unsuitable for acids, alcohols, alkyl halides, aldehydes, ketones, amines and esters. Reacts violently if water is present and can cause a fire with highly flammable liquids.

Sodium hydroxide. Properties and applications are similar to those for potassium hydroxide.

Sodium-potassium alloy. Used as lumps. Lower melting than sodium, so that its surface is readily renewed by shaking. Properties and applications are similar to those for sodium.

Sodium sulfate (anhydrous). Has a large capacity for absorption of water, forming the decahydrate below 33°, but drying is slow and inefficient, especially for solvents that are sparingly soluble in water. It is suitable for the preliminary drying of most types of organic compounds.

Sulfuric acid (concentrated). Widely used in desiccators. Suitable for drying bromine, saturated hydrocarbons, alkyl and aryl halides. Also suitable for drying the following gases: hydrogen, nitrogen, carbon dioxide, carbon monoxide, chlorine, methane and paraffins. Unsuitable for alcohols, bases, ketones or phenols. Also available on a solid support with an indicator under the name *Sicacide* (from Merck) for desiccators. The colour changes in *Sicacide* depends on the percentage of water present (e.g. when dry *Sicacide* is red-violet but becomes pale violet with 27% water and pale yellow to colorless with 33% w/w water).

For convenience, many of the above drying agents are listed in Table 18 under the classes of organic compounds for which they are commonly used.

Molecular sieves

Molecular sieves are types of adsorbents composed of crystalline zeolites (sodium and calcium aluminosilicates). By heating them, water of hydration is removed, leaving holes of molecular dimensions in the crystal lattices. These holes are of uniform size and allow the passage into the crystals of small molecules, but not of large ones. This *sieving* action explains their use as very efficient drying agents for gases and liquids. The pore size of these sieves can be modified (within limits) by varying the cations built into the lattices. The four types of molecular sieves currently available are:

Type 3A sieves. A crystalline potassium aluminosilicate with a pore size of about 3 Angstroms. This type of molecular sieves is suitable for drying liquids such as acetone, acetonitrile, methanol, ethanol and 2-propanol, and drying gases such as acetylene, carbon dioxide, ammonia, propylene and butadiene. The material is supplied as beads or pellets.

Type 4A sieves. A crystalline sodium aluminosilicate with a pore size of about 4 Angstroms, so that, besides water, ethane molecules (but not butane) can be adsorbed. This type of molecular sieves is suitable for drying chloroform, dichloromethane, diethyl ether, dimethylformamide, ethyl acetate, cyclohexane, benzene, toluene, xylene, pyridine and diisopropyl ether. It is also useful for low pressure air drying. The material is supplied as beads, pellets or powder.

Type 5A sieves. A crystalline calcium aluminosilicate with a pore size of about 5 Angstroms, these sieves adsorb larger molecules than type 4A. For example, as well as the substances listed above, propane, butane, hexane, butene, higher *n*-olefins, *n*-butyl alcohol and higher *n*-alcohols, and cyclopropane can be adsorbed, but not branched-chain C₆ hydrocarbons, cyclic hydrocarbons such as benzene and cyclohexane, or secondary and tertiary alcohols, carbon tetrachloride or boron trifluoride. This is the type generally used for drying gases, though organic liquids such as THF and dioxane can be dried with this type of molecular sieves.

Type 13X sieves. A crystalline sodium aluminosilicate with a pore size of about 10 Angstroms which enables many branched-chain and cyclic compounds to be adsorbed, in addition to all the substances removed by type 5A sieves.

They are unsuitable for use with strong acids but are stable over the pH range 5-11.

Because of their selectivity, molecular sieves offer advantages over silica gel, alumina or activated charcoal, especially in their very high affinity for water, polar molecules and unsaturated organic compounds. Their relative efficiency is greatest when the impurity to be removed is present at low concentrations. Thus, at 25° and a relative humidity of 2%, type 5A molecular sieves adsorb 18% by weight of water, whereas for silica gel and alumina the figures are 3.5 and 2.5% respectively. Even at 100° and a relative humidity of 1.3% molecular sieves adsorb about 15% by weight of water.

The greater preference of molecular sieves for combining with water molecules explains why this material can be used for drying ethanol and why molecular sieves are probably the most universally useful and efficient drying agents. Percolation of ethanol with an initial water content of 0.5% through a 144 cm long column of type 4A molecular sieves reduced the water content to 10ppm. Similar results have been obtained with pyridine.

The main applications of molecular sieves to purification comprise:

1. Drying of gases and liquids containing traces of water.
2. Drying of gases at elevated temperatures.
3. Selective removal of impurities (including water) from gas streams.

(For example, carbon dioxide from air or ethene; nitrogen oxides from nitrogen; methanol from diethyl ether. In general, carbon dioxide, carbon monoxide, ammonia, hydrogen sulfide, mercaptans, ethane, ethene, acetylene (ethyne), propane and propylene are readily removed at 25°. In mixtures of gases, the more polar ones are preferentially adsorbed).

The following applications include the removal of straight-chain from branched-chain or cyclic molecules. For example, type 5A sieves will adsorb *n*-butyl alcohol but not its branched-chain isomers. Similarly, it separates *n*-tetradecane from benzene, or *n*-heptane from methylcyclohexane.

The following liquids have been dried with molecular sieves: acetone, acetonitrile, acrylonitrile, allyl chloride, amyl acetate, benzene, butadiene, *n*-butane, butene, butyl acetate, *n*-butylamine, *n*-butyl chloride, carbon tetrachloride, chloroethane, 1-chloro-2-ethylhexane, cyclohexane, dichloromethane, dichloroethane, 1,2-dichloropropane, 1,1-dimethoxyethane, dimethyl ether, 2-ethylhexanol, 2-ethylhexylamine, *n*-heptane, *n*-hexane, isoprene, isopropyl alcohol, diisopropyl ether, methanol, methyl ethyl ketone, oxygen, *n*-pentane, phenol, propane, *n*-propyl alcohol, propylene, pyridine, styrene, tetrachloroethylene, toluene, trichloroethylene and xylene. In addition, the following gases have been dried: acetylene, air, argon, carbon dioxide, chlorine, ethene, helium, hydrogen, hydrogen chloride, hydrogen sulfide, nitrogen, oxygen and sulfur hexafluoride.

After use, molecular sieves can be regenerated by heating at between 300°–350° for several hours, preferably in a stream of dry inert gas such as nitrogen or preferably under vacuum, then cooling in a desiccator. Special precautions must be taken before regeneration of molecular sieves used in the drying of flammable solvents.

However, care must be exercised in using molecular sieves for drying organic liquids. Appreciable amounts of impurities were *formed* when samples of acetone, 1,1,1-trichloroethane and methyl-*t*-butyl ether were dried in the liquid phase by contact with molecular sieves 4A (Connett *Lab Pract* 21 545 1972). Other, less reactive types of sieves may be more suitable but, in general, it seems desirable to make a preliminary test to establish that no unwanted reaction takes place. Useful comparative data for Type 4A and 5A sieves are in Table 19.

MISCELLANEOUS TECHNIQUES

Freeze-pump-thaw and purging

Volatile contaminants, e.g. traces of low boiling solvent residue or oxygen, in liquid samples or solutions can be very deleterious to the samples on storage. These contaminants can be removed by repeated freeze-pump-thaw cycles. This involves freezing the liquid material under high vacuum in an appropriate vessel (which should be large enough to avoid contaminating the vacuum line with liquid that has bumped) connected to the vacuum line *via* efficient liquid nitrogen traps. The frozen sample is then thawed until it liquefies, kept in this form for some time (*ca* 10-15min), refreezing the sample and the cycle repeated several times without interrupting the vacuum. This procedure applies equally well to solutions, as well as purified liquids, e.g. as a means of removing oxygen from solutions for NMR and other measurements. If the presence of nitrogen, helium or argon, is not a serious contaminant then solutions can be freed from gases, e.g. oxygen, carbon dioxide, and volatile impurities by purging with N₂, He or Ar at room, or slightly elevated, temperature. The gases used for purging are then removed by freeze-pump-thaw cycles or simply by keeping in a vacuum for several hours. Special NMR tubes

with a screw cap thread and a PTFE valve (Wilmad) are convenient for freeze thawing of NMR samples. Alternatively NMR tubes with "J Young" valves (Wilmad) can also be used.

Vacuum-lines, Schlenk and glovebox techniques

Manipulations involving materials sensitive to air or water vapour can be carried out by these procedures. Vacuum-line methods make use of quantitative transfers, and **P**(pressure)-**V**(volume)-**T**(temperature) measurements, of gases, and trap-to-trap separations of volatile substances.

It is usually more convenient to work under an inert-gas atmosphere using **Schlenk** type apparatus. The *principle* of Schlenk methods involve a flask/vessel which has a standard ground-glass joint and a sidearm with a tap. The system can be purged by evacuating and flushing with an inert gas (usually nitrogen, or in some cases, argon), repeating the process until the contaminants in the vapour phases have been diminished to acceptable limits. A large range of Schlenk glassware is commercially available (e.g. see Aldrich Chemical Catalog and the associated technical bulletin AL-166). With these, and tailor-made pieces of glassware, inert atmospheres can be maintained during crystallisation, filtration, sublimation and transfer.

Syringe techniques have been developed for small volumes, while for large volumes or where much manipulation is required, dryboxes (*glove boxes*) or dry chambers should be used.

ABBREVIATIONS

Titles of periodicals are defined as in the Chemical Abstracts Service Source Index (CASSI), except that full stops have been omitted after each abbreviated word. Abbreviations of words in the texts of Chapters 4, 5 and 6 are those in common use and are self evident, e.g. *distn*, *filtd*, *conc* and *vac* are used for distillation, filtered, concentrated and vacuum.

TABLES

TABLE 1. SOME COMMON IMMISCIBLE OR SLIGHTLY MISCIBLE PAIRS OF SOLVENTS

Carbon tetrachloride	with ethanolamine, ethylene glycol, formamide or water.
Dimethyl formamide	with cyclohexane or petroleum ether.
Dimethyl sulfoxide	with cyclohexane or petroleum ether.
Ethyl ether	with ethanolamine, ethylene glycol or water.
Methanol	with carbon disulfide, cyclohexane or petroleum ether.
Petroleum ether	with aniline, benzyl alcohol, dimethyl formamide, dimethyl sulfoxide, formamide, furfuryl alcohol, phenol or water.
Water	with aniline, benzene, benzyl alcohol, carbon disulfide, carbon tetrachloride, chloroform, cyclohexane, cyclohexanol, cyclohexanone, diethyl ether, ethyl acetate, isoamyl alcohol, methyl ethyl ketone, nitromethane, tributyl phosphate or toluene.

[Previous Page](#)

with a screw cap thread and a PTFE valve (Wilmad) are convenient for freeze thawing of NMR samples. Alternatively NMR tubes with "J Young" valves (Wilmad) can also be used.

Vacuum-lines, Schlenk and glovebox techniques

Manipulations involving materials sensitive to air or water vapour can be carried out by these procedures. Vacuum-line methods make use of quantitative transfers, and P (pressure)-V (volume)-T (temperature) measurements, of gases, and trap-to-trap separations of volatile substances.

It is usually more convenient to work under an inert-gas atmosphere using **Schlenk** type apparatus. The *principle* of Schlenk methods involve a flask/vessel which has a standard ground-glass joint and a sidearm with a tap. The system can be purged by evacuating and flushing with an inert gas (usually nitrogen, or in some cases, argon), repeating the process until the contaminants in the vapour phases have been diminished to acceptable limits. A large range of Schlenk glassware is commercially available (e.g. see Aldrich Chemical Catalog and the associated technical bulletin AL-166). With these, and tailor-made pieces of glassware, inert atmospheres can be maintained during crystallisation, filtration, sublimation and transfer.

Syringe techniques have been developed for small volumes, while for large volumes or where much manipulation is required, dryboxes (*glove boxes*) or dry chambers should be used.

ABBREVIATIONS

Titles of periodicals are defined as in the Chemical Abstracts Service Source Index (CASSI), except that full stops have been omitted after each abbreviated word. Abbreviations of words in the texts of Chapters 4, 5 and 6 are those in common use and are self evident, e.g. distn, filtd, conc and vac are used for distillation, filtered, concentrated and vacuum.

TABLES

TABLE 1. SOME COMMON IMMISCIBLE OR SLIGHTLY MISCIBLE PAIRS OF SOLVENTS

Carbon tetrachloride	with ethanolamine, ethylene glycol, formamide or water.
Dimethyl formamide	with cyclohexane or petroleum ether.
Dimethyl sulfoxide	with cyclohexane or petroleum ether.
Ethyl ether	with ethanolamine, ethylene glycol or water.
Methanol	with carbon disulfide, cyclohexane or petroleum ether.
Petroleum ether	with aniline, benzyl alcohol, dimethyl formamide, dimethyl sulfoxide, formamide, furfuryl alcohol, phenol or water.
Water	with aniline, benzene, benzyl alcohol, carbon disulfide, carbon tetrachloride, chloroform, cyclohexane, cyclohexanol, cyclohexanone, diethyl ether, ethyl acetate, isoamyl alcohol, methyl ethyl ketone, nitromethane, tributyl phosphate or toluene.

TABLE 2. AQUEOUS BUFFERS

Approx. pH	Composition
0	2N sulfuric acid or N hydrochloric acid
1	0.1N hydrochloric acid or 0.18N sulfuric acid
2	Either 0.01N hydrochloric acid or 0.013N sulfuric acid Or 50 mL of 0.1M glycine (also 0.1M NaCl) + 50 mL of 0.1N hydrochloric acid
3	Either 20 mL of the 0.2M Na ₂ HPO ₄ + 80 mL of 0.1M citric acid Or 50 mL of 0.1M glycine + 22.8 mL of 0.1N hydrochloric acid in 100 mL
4	Either 38.5 mL of 0.2M Na ₂ HPO ₄ + 61.5 mL of 0.1M citric acid Or 18 mL of 0.2M NaOAc + 82 mL of 0.2M acetic acid
5	Either 70 mL of 0.2M NaOAc + 30 mL of 0.2M acetic acid Or 51.5 mL of 0.2M Na ₂ HPO ₄ + 48.5 mL of 0.1M citric acid
6	63 mL of 0.2M Na ₂ HPO ₄ + 37 mL of 0.1M citric acid
7	82 mL of M Na ₂ HPO ₄ + 18 mL of 0.1M citric acid
8	Either 50 mL of 0.1M Tris buffer + 29 mL of 0.1N hydrochloric acid, in 100 mL Or 30 mL of 0.05M borax + 70 mL of 0.2M boric acid
9	80 mL of 0.05M borax + 20 mL of 0.2M boric acid
10	Either 25 mL of 0.05M borax + 43 mL of 0.1N NaOH, in 100 mL Or 50 mL of 0.1M glycine + 32 mL of 0.1N NaOH, in 100 mL
11	50 mL of 0.15M Na ₂ HPO ₄ + 15 mL of 0.1N NaOH
12	50 mL of 0.15M Na ₂ HPO ₄ + 75 mL of 0.1N NaOH
13	0.1N NaOH or KOH
14	N NaOH or KOH

These buffers are suitable for use in obtaining ultraviolet spectra. Alternatively, for a set of accurate buffers of low, but constant, ionic strength ($I = 0.01$) covering a pH range 2.2 to 11.6 at 20°, see Perrin Aust J Chem 16 572 1963. "In 100 mL" means that the solution is made up to 100 mL with pure water.

TABLE 3A. PREDICTED EFFECT OF PRESSURE ON BOILING POINT*

Temperature in degrees Centigrade

760 mmHg	0	20	40	60	80	100	120	140	160	180
0.1	-111	-99	-87	-75	-63	-51	-39	-27	-15	-4
0.2	-105	-93	-81	-69	-56	-44	-32	-19	-7	5
0.4	-100	-87	-74	-62	-49	-36	-24	-11	2	15
0.6	-96	-83	-70	-57	-44	-32	-19	-6	7	20
0.8	-94	-81	-67	-54	-41	-28	-15	-2	11	24
1.0	-92	-78	-65	-52	-39	-25	-12	1	15	28
2.0	-85	-71	-58	-44	-30	-16	-3	11	25	39
4.0	-78	-64	-49	-35	-21	-7	8	22	36	51
6.0	-74	-59	-44	-30	-15	-1	14	29	43	58
8.0	-70	-56	-41	-26	-11	4	19	34	48	63
10.0	-68	-53	-38	-23	-8	7	22	37	53	68
14.0	-64	-48	-33	-23	-2	13	28	44	59	74
16.0	-61	-45	-29	-14	2	17	33	48	64	79
20.0	-59	-44	-28	-12	3	19	35	50	66	82
30.0	-54	-38	-22	-6	10	26	42	58	74	90
40.0	-50	-34	-17	-1	15	32	48	64	81	97
50.0	-47	-30	-14	3	19	36	52	69	86	102
60.0	-44	-28	-11	6	23	40	56	73	86	107
80.0	-40	-23	-6	11	28	45	62	79	97	114
100.0	-37	-19	-2	15	33	50	67	85	102	119
150.0	-30	-12	6	23	41	59	77	95	112	130
200.0	-25	-7	11	29	47	66	84	102	120	138
300.0	-18	1	19	38	57	75	94	113	131	150
400.0	-13	6	25	44	64	83	102	121	140	159
500.0	-8	11	30	50	69	88	108	127	147	166
600.0	-5	15	34	54	74	93	113	133	152	172
700.0	-2	18	38	58	78	98	118	137	157	177
750.0	0	20	40	60	80	100	120	140	160	180
770.0	0	20	40	60	80	100	120	140	160	180
800.0	1	21	41	61	81	101	122	142	162	182


* *How to use the Table:* Take as an example a liquid with a boiling point of 80°C at 760mm Hg. The Table gives values of the boiling points of this liquid at pressures from 0.1 to 800mm Hg. Thus at 50mm Hg this liquid has a boiling point of 19°C, and at 2mm Hg its boiling point would be -30°C.

TABLE 3B. PREDICTED EFFECT OF PRESSURE ON BOILING POINT*

Temperature in degrees Centigrade											
760mmHg	200	220	240	260	280	300	320	340	360	380	400
0.1	8	20	32	44	56	68	80	92	104	115	127
0.2	17	30	42	54	67	79	91	103	116	128	140
0.4	27	40	53	65	78	91	103	116	129	141	154
0.6	33	40	59	72	85	98	111	124	137	150	163
0.8	38	51	64	77	90	103	116	130	143	156	169
1.0	41	54	68	81	94	108	121	134	147	161	174
2.0	53	66	80	94	108	121	135	149	163	176	190
4.0	65	79	93	108	122	136	151	156	179	193	208
6.0	72	87	102	116	131	146	160	175	189	204	219
8.0	78	93	108	123	137	152	167	182	197	212	227
10.0	83	98	113	128	143	158	173	188	203	218	233
14.0	90	105	120	136	151	166	182	197	212	228	243
18.0	95	111	126	142	157	173	188	204	219	235	251
20.0	97	113	129	144	160	176	191	207	223	238	254
30.0	106	123	139	155	171	187	203	219	235	251	267
40.0	113	130	146	162	179	195	211	228	244	260	277
50.0	119	135	152	168	185	202	218	235	251	268	284
60.0	123	140	157	174	190	207	224	241	257	274	291
80.0	131	148	165	182	199	216	233	250	267	284	301
100.0	137	154	171	189	206	223	241	258	275	293	310
150.0	148	166	184	201	219	237	255	273	290	308	326
200.0	156	174	193	211	229	247	265	283	302	320	338
300.0	169	187	206	225	243	262	281	299	318	337	355
400.0	178	197	216	235	254	273	292	311	330	350	369
500.0	185	205	224	244	263	282	302	321	340	360	379
600.0	192	211	231	251	270	290	310	329	349	368	388
700.0	197	217	237	257	277	296	316	336	356	376	396
750.0	200	220	239	259	279	299	319	339	359	379	399
770.0	200	220	241	261	281	301	321	341	361	381	401
800.0	202	222	242	262	282	302	322	342	362	382	403

* *How to use the Table:* Taking as an example a liquid with a boiling point of 340°C at 760mm Hg, the column headed 340°C gives values of the boiling points of this liquid at each value of pressures from 0.1 to 800mm Hg. Thus, at 100mm Hg its boiling point is 258°C, and at 0.8mm Hg its boiling point will be 130°C.

FIGURE 1: NOMOGRAM



How to use Figure 1:

You can use a nomogram to estimate the boiling points of a substance at a particular pressure. For example, the boiling point of 4-methoxybenzenesulfonyl chloride is 173°C/14mm. Thus to find out what the boiling point of this compound will be at 760mm (atmospheric), draw a point on curve A (pressure) at 14mm (this is shown in (i)). Then draw a point on curve C (observed boiling point) corresponding to 173° (or as close as possible). This is shown in (ii). Using a ruler, find the point of intersection on curve B, drawing a line between points (i) and (ii). This is the point (iii) and is the boiling point of 4-methoxybenzenesulfonyl chloride (i.e. approx. 310°C) at atmospheric pressure. If you want to distil 4-methoxybenzenesulfonyl chloride at 20mm, then you will need to draw a point on curve A (at 20mm). Using a ruler, find the point of intersection on curve C drawing through the line intersecting (iii, curve B, i.e. 310°C) and the point in curve A corresponding to 20mm. You should have a value of 185°C, that is, the boiling point of 4-methoxybenzenesulfonyl chloride is estimated to be at 185°C at 20mm.

TABLE 4. HEATING BATHS

Up to 100°	Water baths
-20 to 200°	Glycerol or di- <i>n</i> -butyl phthalate
Up to about 200°	Medicinal paraffin
Up to about 250°	Hard hydrogenated cotton-seed oil (m 40-60°) or a 1:1 mixture of cotton-seed oil and castor oil containing about 1% of hydroquinone.
-40 to 250° (to 400° under N ₂)	D.C. 550 silicone fluid
Up to about 260°	A mixture of 85% orthophosphoric acid (4 parts) and metaphosphoric acid (1 part)
Up to 340°	A mixture of 85% orthophosphoric acid (2 parts) and metaphosphoric acid (1 part)
60 to 500°	Fisher bath wax (highly unsaturated)
73 to 350°	Wood's Metal*
250 to 800°	Solder*
350 to 800°	Lead*

* In using metal baths, the container (usually a metal crucible) should be removed while the metal is still molten.

TABLE 5. WHATMAN FILTER PAPERS

Grade No.	1	2	3	4	5	6	113
Particle size retained (in microns)	11	8	5	12	2.4	2.8	28
Filtration speed*(sec/100mL)	40	55	155	20	<300	125	9

Routine ashless filters

Grade No.	40	41	42	43	44
Particle size retained (in microns)	7.5	12	3	12	4
Filtration speed* (sec/100mL)	68	19	200	38	125

Hardened**Hardened ashless**

Grade No.	50	52	54	540	541	542
Particle size retained(in microns)	3	8	20	9	20	3
Filtration speed* (sec/100mL)	250	55	10	55	12	250

Glass microfilters

Grade No	GF/A	GF/B	GF/C	GF/D	GF/F
Particle size retained (in microns)	1.6	1.0	1.1	2.2	0.8
Filtration speed (sec/100mL)*	8.3	20.0	8.7	5.5	17.2

*Filtration speeds are rough estimates of initial flow rates and should be considered on a relative basis.

TABLE 6. MICRO FILTERS*

Nucleopore (polycarbonate) Filters						
Mean Pore Size (microns)	8.0	2.0	1.0	0.1	0.03	0.015
Av. pores/cm ²	10 ⁵	2x10 ⁶	2x10 ⁷	3x10 ⁸	6x10 ⁸	1-6x10 ⁹
Water flow rate(mL/min/cm ²)	2000	2000	300	8	0.03	0.1-0.5

Millipore Filters						
Type	—Cellulose ester—		—Teflon—		—Microweb [#] —	
	MF/SC	MF/VF	LC	LS	WS	WH
Mean Pore Size (microns)	8	0.01	10	5	3	0.45
Water flow rate (mL/min/cm ²)	850	0.2	170	70	155	55

Gelman Membranes						
Type	—Cellulose ester—				—Copolymer—	
	GA-1	TCM-450	VM-1	DM-800	AN-200	Tuffryn-450
Mean Pore Size (microns)	5	0.45	5	0.8	0.2	0.45
Water flow rate (mL/min/cm ²)	320	50	700	200	17	50

Sartorius Membrane Filters (SM)					
Application	Gravi-metric	Biological clarificatn	Sterili-sation	Particle count in H ₂ O	For acids & bases
Type No.	11003	11004	11006	11011	12801
Mean PoreSize (microns)	1.2	0.6	0.45	0.01	8
Water flow rate (mL/min/cm ²)	300	150	65	0.6	1100

* Only a few representative filters are tabulated (available ranges are more extensive). # Reinforced nylon.

TABLE 7. COMMON SOLVENTS USED IN RECRYSTALLISATION

Acetic acid (118°)	*Cyclohexane (81°)	*Methanol (64.5°)
*Acetone (56°)	Dichloromethane (41°)	*Methyl ethyl ketone (80°)
Acetylacetone (139°)	*Diethyl ether (34.5°)	Methyl isobutyl ketone (116°)
Acetonitrile (82°C)	Dimethyl formamide (76°/39mm)	Nitrobenzene (210°)
*Benzene (80°)	*Dioxane (101°)	Nitromethane (101°)
Benzyl alcohol (93°/10mm)	*Ethanol (78°)	*Petroleum ether (various)
<i>n</i> -Butanol (118°)	2-Ethoxyethanol (cellosolve 135°)	Pyridine (115.5°)
Butyl acetate (126.5°)	*Ethyl acetate (78°)	Pyridine trihydrate (93°)
<i>n</i> -Butyl ether (142°)	Ethyl benzoate (98°/19mm)	*Tetrahydrofuran (64-66°)
γ -Butyrolactone (206°)	Ethylene glycol (68°/4mm)	Toluene (110°)
Carbon tetrachloride (77°)	Formamide (110°/10mm)	Trimethylene glycol (59°/11mm)
Chlorobenzene (132°)	Glycerol (126°/11mm)	Water (100°)
Chloroform (61°)	Isoamyl alcohol (131°)	Xylenes (<i>o</i> 143-145°, <i>m</i> 138-139°, <i>p</i> 138°)

*Highly flammable, should be heated or evaporated on steam or electrically heated water baths only (preferably under nitrogen). None of these solvents should be heated over a naked flame.

TABLE 8. PAIRS OF MISCIBLE SOLVENTS

- Acetic acid:** with chloroform, ethanol, ethyl acetate, acetonitrile, petroleum ether, or water.
- Acetone:** with benzene, butyl acetate, butyl alcohol, carbon tetrachloride, chloroform, cyclohexane, ethanol, ethyl acetate, methyl acetate, acetonitrile, petroleum ether or water.
- Ammonia:** with ethanol, methanol, pyridine.
- Aniline:** with acetone, benzene, carbon tetrachloride, ethyl ether, *n*-heptane, methanol, acetonitrile or nitrobenzene.
- Benzene:** with acetone, butyl alcohol, carbon tetrachloride, chloroform, cyclohexane, ethanol, acetonitrile, petroleum ether or pyridine.
- Butyl alcohol:** with acetone or ethyl acetate.
- Carbon disulfide:** with petroleum ether.
- Carbon tetrachloride:** with cyclohexane.
- Chloroform:** with acetic acid, acetone, benzene, ethanol, ethyl acetate, hexane, methanol or pyridine.
- Cyclohexane:** with acetone, benzene, carbon tetrachloride, ethanol or diethyl ether.
- Diethyl ether:** with acetone, cyclohexane, ethanol, methanol, methylal (dimethoxymethane), acetonitrile, pentane or petroleum ether.
- Dimethyl formamide:** with benzene, ethanol or ether.
- Dimethyl sulfoxide:** with acetone, benzene, chloroform, ethanol, diethyl ether or water.
- Dioxane:** with benzene, carbon tetrachloride, chloroform, ethanol, diethyl ether, petroleum ether, pyridine or water.
- Ethanol:** with acetic acid, acetone, benzene, chloroform, cyclohexane, dioxane, ethyl ether, pentane, toluene, water or xylene.
- Ethyl acetate:** with acetic acid, acetone, butyl alcohol, chloroform, or methanol.
- Glycerol:** with ethanol, methanol or water.
- Hexane:** with benzene, chloroform or ethanol.
- Methanol:** with chloroform, diethyl ether, glycerol or water.
- Methylal:** with diethyl ether.
- Methyl ethyl ketone:** with acetic acid, benzene, ethanol or methanol.
- Nitrobenzene:** with aniline, methanol or acetonitrile.
- Pentane:** with ethanol or diethyl ether.
- Petroleum ether:** with acetic acid, acetone, benzene, carbon disulfide or diethyl ether.
- Phenol:** with carbon tetrachloride, ethanol, diethyl ether or xylene.
- Pyridine:** with acetone, ammonia, benzene, chloroform, dioxane, petroleum ether, toluene or water.
- Toluene:** with ethanol, diethyl ether or pyridine.
- Water:** with acetic acid, acetone, ethanol, methanol, or pyridine.
- Xylene:** with ethanol or phenol.

TABLE 9. MATERIALS FOR COOLING BATHS

Temperature	Composition	Temperature	Composition
0°	Crushed ice		
-5° to -20°	Ice-salt mixtures	-77°	Solid CO ₂ with chloroform or acetone
Up to -20°	Ice-MeOH mixtures	-78°	Solid CO ₂ (powdered; CO ₂ snow)
-33°	Liquid ammonia		
-40° to -50°	Ice (3.5-4 parts) - CaCl ₂ 6H ₂ O (5 parts)	-100°	Solid CO ₂ with diethyl ether
-72°	Solid CO ₂ with ethanol	-196°	liquid nitrogen (see footnote*)

Alternatively, the following liquids can be used, partially frozen, as cryostats, by adding solid CO₂ from time to time to the material in a Dewar-type container and stirring to make a slush:

13°	<i>p</i> -Xylene	-55°	Diacetone
12°	Dioxane	-56°	<i>n</i> -Octane
6°	Cyclohexane	-60°	Di-isopropyl ether
5°	Benzene	-73°	Trichloroethylene or isopropyl acetate
2°	Formamide	-74°	<i>o</i> -Cymene or <i>p</i> -cymene
-8.6°	Methyl salicylate	-77°	Butyl acetate
-9°	Hexane-2,5-dione	-79°	Isoamyl acetate
-10.5°	Ethylene glycol	-83°	Propylamine
-11.9°	<i>tert</i> -Amyl alcohol	-83.6°	Ethyl acetate
-12°	Cycloheptane or methyl benzoate	-86°	Methyl ethyl ketone
-15°	Benzyl alcohol	-89°	<i>n</i> -Butanol
-16.3°	<i>n</i> -Octanol	-90°	Nitroethane
-18°	1,2-Dichlorobenzene	-91°	Heptane
-22°	Tetrachloroethylene	-92°	<i>n</i> -Propyl acetate
-22.4°	Butyl benzoate	-93°	2-Nitropropane or cyclopentane
-22.8°	Carbon tetrachloride	-94°	Ethyl benzene or hexane
-24.5°	Diethyl sulfate	-94.6°	Acetone
-25°	1,3-Dichlorobenzene	-95.1°	Toluene
-29°	<i>o</i> -Xylene or pentachloroethane	-97°	Cumene
-30°	Bromobenzene	-98°	Methanol or methyl acetate
-32°	<i>m</i> -Toluidine	-99°	Isobutyl acetate
-32.6°	Dipropyl ketone	-104°	Cyclohexene
-38°	Thiophene	-107°	Isooctane
-41°	Acetonitrile	-108°	1-Nitropropane
-42°	Pyridine or diethyl ketone	-116°	Ethanol or diethyl ether
-44°	Cyclohexyl chloride	-117°	Isoamyl alcohol
-45°	Chlorobenzene	-126°	Methylcyclohexane
-47°	<i>m</i> -Xylene	-131°	<i>n</i> -Pentane
-50°	Ethyl malonate or <i>n</i> -butylamine	-160°	Isopentane
-52°	Benzyl acetate or diethylcarbitol		

For other organic materials used in low temperature slush-baths with liquid nitrogen see R.E.Rondeau [*J Chem Eng Data* 11 124 1966]. *NOTE: Use high quality pure nitrogen, do not use liquid air or liquid nitrogen that has been in contact with air for a long period (due to the dissolution of oxygen in it) as this could EXPLODE in contact with organic matter.

TABLE 10. LIQUIDS FOR STATIONARY PHASES IN GAS CHROMATOGRAPHY

Material	Temp.	Retards
Dimethylsulfolane	0-40°	Olefins and aromatic hydrocarbons
Di- <i>n</i> -butyl phthalate	0-40°	General purposes
Squalane	0-150°	Volatile hydrocarbons and polar molecules
Silicone oil or grease	0-250°	General purposes
Diglycerol	20-120°	Water, alcohols, amines, esters, and aromatic hydrocarbons
Dinonyl phthalate	20-130°	General purposes
Polydiethylene glycol succinate	50-200°	Aromatic hydrocarbons, alcohols, ketones, esters.
Polyethylene glycol	50-200°	Water, alcohols, amines, esters and aromatic hydrocarbons
Apiezon grease	50-200°	Volatile hydrocarbons and polar molecules
Tricresyl phosphate	50-250°	General purposes

TABLE 11. METHODS OF VISUALISATION OF TLC SPOTS

Reagent	Compound	Preparation	Observations
Iodine	General	Iodine crystals in a closed chamber or spray 1% methanol solution of Iodine	Brown spots which may disappear upon standing. Limited sensitivity.
H ₂ SO ₄	General	50% solution, followed by heating to 150°C	Black or coloured spots
Molybdate	General	5% (NH ₄) ₆ Mo ₇ O ₂₄ + 0.2% Ce(SO ₄) ₂ in 5% H ₂ SO ₄ , followed by heating to 150°C.	Deep blue spots
Vanillin	General	0.5g vanillin, 0.5 mL H ₂ SO ₄ , 9 mL ethanol	various coloured spots
Ammonia	phenols	Ammonia vapour in a closed chamber	various coloured spots
FeCl ₃	phenols, enolic compounds	1% aqueous FeCl ₃	various coloured spots
2,4-DNP	aldehydes, ketones	0.5% 2,4-dinitrophenylhydrazine/2M HCl	red to yellow spots
HCl	aromatic acids and amines	HCl vapour in a closed chamber	various coloured spots
Ninhydrin	amino acids, and amines	0.3% ninhydrin in <i>n</i> -BuOH with 3% AcOH, followed by heating to 125°C/10 min	blue spots
PdCl ₂	S and Se compds	0.5% aq. PdCl ₂ + few drops of conc. HCl	red and yellow spots
Anisaldehyde	carbohydrates	0.5 mL anisaldehyde in 0.5 mL conc H ₂ SO ₄ + 95% EtOH + a few drops of AcOH Heat at 100-110°C for 20-30 minutes	various blue spots

TABLE 12. GRADED ADSORBENTS AND SOLVENTS

Adsorbents (decreasing effectiveness)	Solvents (increasing eluting ability)
Fuller's earth (hydrated aluminosilicate)	Petroleum ether, b. 40-60°.
Magnesium oxide	Petroleum ether, b. 60-80°.
Charcoal	Carbon tetrachloride.
Alumina	Cyclohexane.
Magnesium trisilicate	Benzene.
Silica gel	Diethyl ether.
Calcium hydroxide	Chloroform.
Magnesium carbonate	Ethyl acetate.
Calcium phosphate	Acetone.
Calcium carbonate	Ethanol.
Sodium carbonate	Methanol.
Talc	Pyridine.
Inulin	Acetic acid.
Sucrose = starch	

TABLE 13. REPRESENTATIVE ION-EXCHANGE RESINS

Sulfonated polystyrene	Aliphatic amine-type
Strong-acid cation exchanger	weak base anion exchangers
AG 50W-x8	Amberlites IR-45 and IRA-67
Amberlite IR-120	Dowex 3-x4A
Dowex 50W-x8	Permutit E
Duolite 225	Permutit A 240A
Permutit RS	
Permutite C50D	
Carboxylic acid-type	Strong Base, anion exchangers
Weak acid cation exchangers	AG 2x8
Amberlite IRC-50	Amberlite IRA-400
Bio-Rex 70	Dowex 2-x8
Chelex 100	Duolite 113
Duolite 436	Permutit ESB
Permutit C	Permutite 330D
Permutits H and H-70	

TABLE 14. MODIFIED FIBROUS CELLULOSES FOR ION-EXCHANGE

Cation exchange	Anion exchange
CM cellulose (carboxymethyl)	DEAE cellulose (diethylaminoethyl)
CM 22, 23 cellulose	DE 22, 23 cellulose
P cellulose (phosphate)	PAB cellulose (<i>p</i> -aminobenzyl)
SE cellulose (sulfoethyl)	TEAE cellulose (triethylaminoethyl)
SM cellulose (sulfomethyl)	ECTEOLA cellulose

SE and SM are much stronger acids than CM, whereas P has two ionisable groups (pK 2-3, 6-7), one of which is stronger, the other weaker, than for CM (3.5-4.5). For basic strengths, the sequence is: TEAE » DEAE (pK 8-9.5) > ECTEOLA (pK 5.5-7) > PAB. Their exchange capacities lie in the range 0.3 to 1.0 mg equiv/g.

TABLE 15. BEAD FORM ION-EXCHANGE PACKAGINGS¹

Cation exchange	Capacity (meq/g)	Anion exchange	Capacity (meq/g)
CM-Sephadex C-25, C-50. ² (weak acid)	4.5±0.5	DEAE-Sephadex A-25, A-50. ⁷ (weak base)	3.5±0.5
SP-Sephadex C-25, C-50. ³ (strong acid)	2.3±0.3	QAE-Sephadex A-25, A-50. ⁸ (strong base)	3.0±0.4
CM-Sepharose CL-6B. ⁴	0.12±0.02	DEAE-Sepharose CL-6B. ⁴	0.13±0.02
		DEAE-Sephacel. ⁹	1.4±0.1
Fractogel EMD, CO ₂ ⁻ (pK ~4.5) , SO ₃ ²⁻ (pK ~<1) . ⁵		Fractogel EMD, DMAE (pK ~9), DEAE (pK ~10.8), TMAE (pK >13). ⁵	
CM-32 Cellulose.		DE-32 Cellulose.	
CM-52 Cellulose. ⁶		DE-52 Cellulose	

¹ May be sterilised by autoclaving at pH 7 and below 120°. ² Carboxymethyl. ³ Sulfopropyl. ⁴ Crosslinked agarose gel, no pre-cycling required, pH range 3-10. ⁵ Hydrophilic methacrylate polymer with very little volume change on change of pH (equivalent to *Toyopearl*, Sigma), available in superfine 650S, and medium 650M particle sizes. ⁶ Microgranular, pre-swollen, no pre-cycling required. ⁷ Diethylaminoethyl. ⁸ Diethyl(2-hydroxypropyl)aminoethyl. ⁹ Bead form cellulose, pH range 2-12, no pre-cycling required. Sephadex and Sepharose from Pharmacia-Amersham Biosciences, Fractogel from Merck, Cellulose from Whatman.

TABLE 16. LIQUIDS FOR DRYING PISTOLS

Boiling points (760mm)		Boiling points (760mm)	
Ethyl chloride	12.2°	Toluene	110.5°
Dichloromethane	39.8°	Tetrachloroethylene	121.2°
Acetone	56.1°	Chlorobenzene	132.0°
Chloroform	62.0°	<i>m</i> -Xylene	139.3°
Methanol	64.5°	Isoamyl acetate	142.5°
Carbon tetrachloride	76.5°	Tetrachloroethane	146.3°
Ethanol	78.3°	Bromobenzene	155.0°
Benzene	79.8°	<i>p</i> -Cymene	176.0°
Trichloroethylene	86.0°	Tetralin	207.0°
Water	100.0°		

TABLE 17. VAPOUR PRESSURES (mm Hg) OF SATURATED AQUEOUS SOLUTIONS IN EQUILIBRIUM WITH SOLID SALTS

Salt	Temperature					% Humidity at 20°
	10°	15°	20°	25°	30°	
LiCl.H ₂ O			2.6			15
CaBr ₂ .6H ₂ O	2.1	2.7	3.3	4.0	4.8	19
KOAc			3.5			20
CaCl ₂ .6H ₂ O	3.5	4.5	5.6	6.9	8.3	20
CrO ₃			6.1			32
Zn(NO ₃) ₂ .6H ₂ O			7.4			42
K ₂ CO ₃ .2H ₂ O			7.7	10.7		44
KCNS			8.2			47
Na ₂ Cr ₂ O ₇ .2H ₂ O			9.1			52
Ca(NO ₃) ₂ .4H ₂ O	6.0	7.7	9.6	11.9	14.2	55
Mg(NO ₃) ₂ .6H ₂ O			9.8			56
NaBr.2H ₂ O	5.8	7.8	10.3	13.5	17.5	58
NaNO ₂			11.6			66
NaClO ₃			13.1			75
NaCl	6.9	9.6	13.2	17.8	21.4	75
NaOAc			13.3			76
NH ₄ Cl			13.8			79
(NH ₄) ₂ SO ₄			14.2			81
KBr			14.7			84
KHSO ₄			15.1			86
KCl			15.1	20.2	27.0	86
K ₂ CrO ₄			15.4			88
ZnSO ₄ .7H ₂ O			15.8			90
NH ₄ .H ₂ PO ₄			16.3			93
KNO ₃			16.7	22.3	29.8	95
Pb(NO ₃) ₂			17.2			98
H ₂ O	9.21	12.79	17.53	23.76	31.82	100

TABLE 18. DRYING AGENTS FOR CLASSES OF COMPOUNDS
Class Dried with

Acetals	Potassium carbonate.
Acids (organic)	Calcium sulfate, magnesium sulfate, sodium sulfate.
Acyl halides	Magnesium sulfate, sodium sulfate.
Alcohols	Calcium oxide, calcium sulfate, magnesium sulfate, potassium carbonate, followed by magnesium and iodine.
Aldehydes	Calcium sulfate, magnesium sulfate, sodium sulfate.
Alkyl halides	Calcium chloride, calcium sulfate, magnesium sulfate, phosphorus pentoxide, sodium sulfate.
Amines	Barium oxide, calcium oxide, potassium hydroxide, sodium carbonate, sodium hydroxide.
Aryl halides	Calcium chloride, calcium sulfate, magnesium sulfate, phosphorus pentoxide, sodium sulfate.
Esters	Magnesium sulfate, potassium carbonate, sodium sulfate.
Ethers	Calcium chloride, calcium sulfate, magnesium sulfate, sodium, lithium aluminium hydride.
Heterocyclic bases	Magnesium sulfate, potassium carbonate, sodium hydroxide.
Hydrocarbons	Calcium chloride, calcium sulfate, magnesium sulfate, phosphorus pentoxide, sodium (not for olefins).
Ketones	Calcium sulfate, magnesium sulfate, potassium carbonate, sodium sulfate.
Mercaptans	Magnesium sulfate, sodium sulfate.
Nitro compounds and Nitriles	Calcium chloride, magnesium sulfate, sodium sulfate.
Sulfides	Calcium chloride, calcium sulfate.

TABLE 19.		STATIC DRYING FOR SELECTED LIQUIDS (25°C)			
Liquid	Water	Linde Type 4 A	Linde Type 5 A	Activated Alumina	Silicic Acid Gel
MeOH	Residual H ₂ O %	0.54	0.55	—	0.60
	Wt % absorbed	2.50	1.50	—	—
EtOH	Residual H ₂ O %	0.25	0.25	0.45	0.68
	Wt % absorbed	7.00	6.80	1.50	—
1-Butylamine	Residual H ₂ O %	1.65	1.31	1.93	2.07
	Wt % absorbed	10.40	18.20	3.40	—
2-Ethyl- hexylamine	Residual H ₂ O %	0.25	0.08	0.43	0.53
	Wt % absorbed	15.10	21.10	6.10	1.70
Diethyl ether	Residual H ₂ O %	0.001	0.013	0.16	0.27
	Wt % absorbed	9.50	9.20	6.20	4.30
Amyl acetate	Residual H ₂ O %	0.002	—	0.33	0.38
	Wt % absorbed	9.30	—	7.40	1.80

TABLE 20. BOILING POINTS OF SOME USEFUL GASES AT 760 mm

Argon	-185.6°	Krypton	-152.3°
Carbon dioxide (sublimes)	-78.5°	Methane	-164.0°
Carbon monoxide	-191.3°	Neon	-246.0°
Ethane	-88.6°	Nitrogen	-209.9°
Helium	-268.6°	Nitrous oxide	-88.5°
Hydrogen	-252.6°	Nitric oxide	-195.8°
		Oxygen	-182.96°

TABLE 21. SOLUBILITIES OF HCl AND NH₃ AT 760mm (g/100g OF SOLUTION)

Gas	Temperature °C	MeOH	EtOH	Et ₂ O
Hydrogen Chloride*	-10	54.6	—	37.5 (-9.2°)
	0	51.3	45.4	35.6
	20	47.0 (18°)	41.0	24.9
	30	43.0	38.1	19.47
Ammonia	15	21.6 (27.6g/100g MeOH)	13.2 (9.2g/100mL soln)	—
	25	16.5 (19.8g/100g MeOH)	10.0 (6.0g/100mL soln)	—

* Saturated EtOH with HCl is ~ 5.7M at 25°C, i.e. 21.5g/100mL of solution.

TABLE 22. PREFIXES FOR QUANTITIES

Fractional	deci (d)	centi (c)	milli (m)	micro (μ)	nano (n)	pico (p)	femto (f)	atto (atto)
	= 10 ⁻¹	= 10 ⁻²	= 10 ⁻³	= 10 ⁻⁶	= 10 ⁻⁹	= 10 ⁻¹²	= 10 ⁻¹⁵	= 10 ⁻¹⁸
Multiple	deca (d)	hecto (h)	kilo (k)	mega (M)	giga (G)	tera (T)	penta (P)	eka (E)
	= 10 ¹	= 10 ²	= 10 ³	= 10 ⁶	= 10 ⁹	= 10 ¹²	= 10 ¹⁵	= 10 ¹⁸

BIBLIOGRAPHY

The following books and reviews provide fuller details of the topics indicated in this chapter. The authors' recommendations for excellent introductory and/or reference texts to the topics are indicated with.* [For earlier bibliographies see *Purification of Laboratory Chemicals*, 4th Edn, ISBN 0750628391 (1996, hardback) and 0750637617 (1997, paperback).

AFFINITY CHROMATOGRAPHY

P. Bailon, G.K. Ehrlich, W. Fung and W. Berthold (Eds), *Affinity Chromatography: Methods and Protocols*, Humana Press, Totowa, 2000. ISBN 0896036944.

I. A. Chaiken, *Analytical Affinity Chromatography*, CRC Press Inc, Florida, 1987. ISBN 084935658X.

* P.D.G. Dean, W.S. Johnson and F. Middle (Eds), *Affinity Chromatography : A Practical Approach*, IRL Press, Oxford, 1985. ISBN 0896036944.

P. Matejtschuk (Ed.), *Affinity Separations: A Practical Approach*, Oxford University Press, Oxford, 1997. ISBN 019963551X, 0199635501.

T.M. Phillips and B.F. Dickens, *Affinity and Immunoaffinity Purification Techniques*, Eaton Publishing, Natick MA, 2000. ISBN 1881299228.

CHIRAL CHROMATOGRAPHY

S. Ahuja (Ed.), *Chiral Separations: Applications and Technology*, ACS, Washington DC, 1997. ISBN 0841234078.

* S. Allenmark, *Chromatographic Enantioseparations (Methods and Applications)*, 2nd Edn, Eliss Horwood Publ. New York, 1991. ISBN 0131329782.

T.E. Beesley and R.P.W. Scott, *Chiral Chromatography*, J.Wiley & Son, Chichester, 1999. ISBN 0471974277.

* W.J. Lough, *Chiral Liquid Chromatography*, Blackie Academic & Professional, Glasgow, 1989. ISBN 0412017415.

D. Stevenson and I.D. Wilson (Eds), *Recent Advances in Chiral Separations* (Chromatographic Society Symposium Series), Plenum Pub Corp., New York, 1991. ISBN 0306438364.

D. Stevenson and I.D. Wilson (Eds), *Chiral Separations*, Plenum Press, New York, 1988. ISBN 0306432528.

G. Subramanian (Ed.), *Chiral Separation Techniques : A Practical Approach*, VCH Verlagsgesellschaft, Weinheim, 2000. ISBN 3527298754.

M. Zief and L.J. Crane (Eds), *Chromatographic Chiral Separations*, Marcel Dekker, New York, 1988. ISBN 0824777867.

CHROMATOGRAPHY

S. Ahuja, *Chromatography and Separation Chemistry: Advances and Developments*, ACS, Washington DC, 1986. ISBN 0841209537.

* K. Blau and J.M. Halket (Eds), *Handbook of Derivatives for Chromatography*, J. Wiley & Sons, New York, 1994. ISBN 047192699X.

P.R. Brown and A. Grushka (Eds), *Advances in Chromatography*, Marcel Dekker, New York, Vol 34 1994, vol 35 1995.

* J. Cazes, *Encyclopedia of Chromatography*, Marcel Dekker, New York, 2001. ISBN 0824705114.

J.C. Giddings, E. Grushka and P.R. Brown (Eds), *Advances in Chromatography*, Marcel Dekker, New York, Vols 1-36, 1955-1995.

E. Heftmann (Ed.), *Chromatography*, 5th Edn, Elsevier Science, Amsterdam, 1992. ISBN 0444882375.

E. Heftmann (Ed.), *Chromatography : Fundamentals and Applications of Chromatographic and Electrophoretic Methods*, 2 volume set, Elsevier Science, New York, 1983. ISBN 0444420452 (set).

K. Hostettmann, A. Marston and M. Hostettmann, *Preparative Chromatography Techniques: Applications in Natural Product Isolation*, 2nd Edn, Springer-Verlag, Berlin, 1998. ISBN 3540624597.

P. Miller (Ed.), *High Resolution Chromatography: A Practical Approach*, Oxford University Press, Oxford, 1999. ISBN 0199636486.

C.F. Poole and S.A. Schuette, *Contemporary Practice of Chromatography*, Elsevier, Amsterdam, 1984. ISBN 0444424105.

C.F. Poole and S.K. Poole, *Chromatography Today*, Elsevier Science, Amsterdam, 4th reprint 1997. ISBN 0444891617.

* G. Zweig and J. Sherma (Eds), *CRC Handbook of Chromatography*, 2 Volumes, CRC Press, Cleveland, 1972. ISBN 0878195602.

CRYSTALLISATION

T.M. Bergfors (Ed.), *Protein Crystallisation*, International University Line, 1999. ISBN 0963681753.

A. Ducruix and R. Giege (Eds), *Crystallisation of Nucleic Acids and Proteins*, IRL Press, Oxford, 1992. ISBN 0199632456. 2nd Edn, Oxford University Press, Oxford, 1999. ISBN 0199636788.

A. Mersmann (Ed.), *Crystallisation Technology Handbook*, Marcel Dekker, New York, 1994. ISBN 0824792335.

H. Michel (Ed.), *Crystallisation of Membrane Proteins*, CRC Press, Boca Raton, 1991. ISBN 0849348161.

J.M. Mullin, *Crystallisation*, 4th Edn, Butterworth-Heinemann, Oxford, 2001. ISBN 0750648333.

A.S. Myerson and K. Toyokura, *Crystallisation as a Separation Process*, ACS, Washington DC, 1990. ISBN 0841218641.

J. Nyvit and J. Ulrich, *Admixtures in Crystallisation*, Wiley-VCH, New York 1995. ISBN 3527287396.

R.S. Tipson (Chapter 3), in A. Weissberger (Ed.), *Techniques of Organic Chemistry*, Vol III, pt I, 2nd Edn, Interscience, New York, 1956.

DISTILLATION

E.S. Perry and A. Weissberger (Eds), *Techniques of Organic Chemistry*, Vol IV, *Distillation*, Interscience, New York, 1965.

J. Stichlmair and J.R. Fair, *Distillation: Principles and Practices*, J. Wiley and Sons, New York, 1998. ISBN 0471252417.

DRYING

G. Broughton (Chapter VI), in A. Weissberger's (Ed.) *Techniques of Organic Chemistry*, Vol III, pt I, 2nd Edn, Interscience, New York, 1956.

J.A. Riddick and W.B. Bunger, *Organic Solvents: Physical Properties and Methods of Purification, Techniques of Chemistry*, Vol II, Wiley-Interscience, New York, 1970.

J.F. Coetzee, *Recommended Methods for the Purification of Solvents and Tests for Impurities*, Pergamon Press, Oxford, 1982. ISBN 0080223702.

GAS CHROMATOGRAPHY

* P.J. Baugh (Ed.), *Gas Chromatography: A Practical Approach*, IRL Press, Oxford, 1994. ISBN 0199632715.

R.E. Clement (Ed.), *Gas Chromatography: Biochemical, Biomedical and Clinical Applications*, J. Wiley & Sons, New York, 1990. ISBN 0471010480.

* I.A. Fowles, *Gas Chromatography: Analytical Chemistry by Open Learning*, 2nd Edn, J. Wiley & Sons, Chichester, 1995. ISBN 0471954683.

W. Jennings, E. Mittlefehldt and P. Stremple, *Analytical Gas Chromatography*, 2nd Edn, Academic Press, San Diego, 1997. ISBN 012384357X.

G. Guichon and C.L. Guillemin, *Quantitative Gas Chromatography for Laboratory and On-Line Process Control*, Elsevier, Amsterdam, 1988. ISBN 0444428577.

H.H. Hill and D.G. McMinn (Eds), *Detectors for Capillary Chromatography*, J. Wiley & Sons, New York, 1992. ISBN 0471506451.

H.-J. Hubschmann, *Handbook of GC/MS: Fundamentals and Applications*, VCH Verlagsgesellschaft MbH, Weinheim, 2001. ISBN 3527301704.

F.W. Karasek and R.E. Clement, *Basic Gas-Chromatography-mass spectrometry: Principles and Techniques*, Elsevier, Amsterdam, 1988. ISBN 0444427600.

* M. McMaster, C. McMaster, *GC/MS: A Practical User's Guide*, Wiley-VCH, New York, 1998. ISBN 0471248266.

* H.M. McNair and J.M. Miller, *Basic Gas Chromatography: Techniques in Analytical Chemistry*, J. Wiley & Sons, New York, 1997. ISBN 047117260X.

* D. Rood, *A Practical Guide to the Care, Maintenance, and Troubleshooting of Capillary Gas Chromatographic Systems*, 3rd Edn, Wiley-VCH, New York; 1999. ISBN 3527297502.

R.D. Sacks, J-M.D. Dimandja and DG. Patterson, *High Speed Gas Chromatography*, J. Wiley & Sons, New York, 2001. ISBN 0471415561.

P. Sandra (Ed.), *Sample Introduction in Capillary Gas Chromatography*, Vol 1, Chromatographic Method Series, Huethig, Basel, 1985.

GEL FILTRATION

P.L. Dubin, *Aqueous Size-Exclusion Chromatography*, Elsevier, Amsterdam, 1988. ISBN 0444429573.

L. Fischer, *Gel Filtration*, 2nd Edn, Elsevier, North-Holland, Amsterdam, 1980. ISBN 0444802231.

M. Potschka and P.L. Dubin (Eds), *Strategies in Size-exclusion Chromatography*, ACS, Washington DC, 1996. ISBN 0841234140.

W.W. Yau, D.D. By and J.J. Kirkland, *Modern Size Exclusion Liquid Chromatography: Practice of Gel Permeation and Gel Filtration Chromatography*, J. Wiley and Sons, New York, 1979. ISBN 0471033871.

C-S. Wu (Ed.), *Handbook of Size Exclusion Chromatography*, Marcel Dekker, NY, 1995. ISBN 0824792882

C-S. Wu (Ed.), *Column Handbook for Size Exclusion Chromatography*, Academic Press, NY, 1998. ISBN 0127655557

HIGH PERFORMANCE LIQUID CHROMATOGRAPHY

T.J. Baker, Chemists, *HPLC Solvents Reference Manual*, T.J.Baker Chemical Co., 1985.

* K.D. Bartle and P. Meyers, *Capillary Electrochromatography*, Royal Society of Chemistry Chromatography Monographs, London, 2000. ISBN 0854045309.

A. Berthold (Ed.), *Countercurrent Chromatography (CCC). The Support-Free Liquid Stationary Phase*, Elsevier, Amsterdam, 2002. ISBN 044450737X.

B.A. Bidlingmeyer, *Preparative Liquid Chromatography*, 2nd Reprint, Elsevier, Amsterdam, 1991. ISBN 0444428321.

R. Freitag, *Modern Advances in Chromatography*, Springer-Verlag, NY, 2002. ISBN 3540430423. [Electronic version available at: <http://link.springer.de/series/abe/>].

W.S. Hancock, J.T. Sparrow, *HPLC Analysis of Biological Compounds: A Laboratory Guide*, (Chromatographic Science Series, Vol 26), Marcel Dekker, New York, 1984. ISBN 0824771400.

T. Hanai and R.M. Smith (Eds), *HPLC: A Practical Guide*, Royal Society of Chemistry, Thomas Graham House, Cambridge, U.K, 1999. ISBN 0854045155.

E.D. Katz (Ed.), *High Performance Liquid Chromatography: Principles and Methods in Biotechnology*, J. Wiley & Sons, Chichester, 1996. ISBN 0471934445.

A.M. Krstulovic and P.R. Brown, *Reversed Phase HPLC. Theory, Practical and Biomedical Applications*, J. Wiley & Sons, New York, 1982. ISBN 0471053694.

C.K. Lim (Ed.), *HPLC of Small Molecules*, IRL Press, Oxford, 1986. ISBN 0947946780, 0947946772.

* W.J. Lough and I.W. Wainer (Eds), *High Performance Liquid Chromatography: Fundamental Principles and Practice*, Blackie Academic and Professional, London, 1996. ISBN 0751400769.

* M.C. McMaster, *HPLC, A Practical User's Guide*, J. Wiley and Sons, New York, 1994. ISBN 0471185868.

* V. Meyer, *Practical High-Performance Liquid Chromatography*, 2nd Edn., J. Wiley & Sons, New York, 1994. ISBN 0471941328, 0471941298.

U.D. Neue, *HPLC Columns: Theory, Technology, and Practice*, Wiley-VCH, New York, 1997. ISBN 0471190373.

R.W.A. Oliver (Ed.), *HPLC of Macromolecules*, 2nd Edn, IRL Press, Oxford, 1998. ISBN 0199635714, 0199635706.

D. Parriott, *A Practical Guide to HPLC Detection*, Academic Press, San Diego, 1997. ISBN 0125456808.

* P.C. Sadek, *The HPLC Solvent Guide*, J. Wiley & Sons, New York, 1996. ISBN 0471118559.

* P.C. Sadek, *Troubleshooting HPLC Systems: A Bench Manual*, Wiley-Interscience, New York, 1999. ISBN 0471178349.

* L.R. Snyder, J. Kirkland and J. Glajch, *Practical HPLC Method Development*, 2nd Edn., Wiley-Interscience, New York, 1997. ISBN 047100703X.

J. Swadesh (Ed.), *HPLC: Practical and Industrial Applications*, CRC Press, Boca Raton, 1997. ISBN 0849326826.

G. Szepesi, *How to Use Reverse Phase HPLC*, VCH, New York, 1992. ISBN 0895737663.

IONIC EQUILIBRIA

G. Kortüm, W. Vogel and K. Andrussow, *Dissociation Constants of Organic Acids in Aqueous Solution*, Butterworths, London, 1961.

D.D. Perrin, *Dissociation Constants of Organic Bases in Aqueous Solution*, Butterworths, London, 1965; and Supplement 1972.

D.D. Perrin, *Dissociation Constants of Inorganic Acids and Bases in Aqueous Solution*, Butterworths, London, 1969.

D.D. Perrin and B. Dempsey, *Buffers for pH and Metal Ion Control*, Chapman and Hall, London, 1974.

E.P. Serjeant and B. Dempsey, *Ionisation Constants of Organic Acids in Aqueous Solution*, Pergamon Press, Oxford, 1979.

ION EXCHANGE

A. Dyer, M.J. Hudson and P.A. Williams (Eds), *Ion Exchange Processes: Advances and Applications*, Royal Society of Chemistry, London, 1993. ISBN 0851864457.

J.S. Fritz and D.T. Gjerde, *Ion Chromatography*, 3rd Edn, VCH Verlagsgesellschaft Mbh, Weinheim, 2000. ISBN 3527299149.

D.T. Gjerde and J.S. Fritz, *Ion Chromatography*, 2nd Edn, J. Wiley and Sons, New York, 1998. ISBN 3527296964.

P.R. Haddad and P.E. Jackson, *Ion Chromatography: Principles and Applications*, Elsevier, New York, 1990. ISBN 0444882324.

F.G. Helfferich, *Ion Exchange*, Dover Publications, New York, 1989. ISBN 0486687848.

J. Korkisch, *Handbook of Ion Exchange Resins: Their Applications to Inorganic Analytical Chemistry*, 6 Vol. Set, CRC Press, Boca Raton, 1989. ISBN 0849331943.

R.E. Smith (Ed.), *Ion Chromatography Applications*, CRC Press, Boca Raton, 1988. ISBN 0849349672.

E.M. Thurman and M.S. Mills, *Solid-phase Extraction: Principles and Practice*, Chemical Analysis Series Vol. 147, J. Wiley & Sons, New York, 1998. ISBN 047161422X.

J. Weiss, *Ion Chromatography*, 2nd Edn, Wiley-VCH, New York, 1994. ISBN 3527286985.

S. Yamamoto, K. Nakanishi and R. Matsuno, *Ion-exchange Chromatography of Proteins*, Chromatographic Science Series Vol. 43, Marcel Dekker, New York, 1988. ISBN 0824779037.

LABORATORY TECHNIQUE AND THEORETICAL DISCUSSION

M. Casey, *Advanced Practical Organic Chemistry*, Chapman and Hall, New York, 1990. ISBN 0412024616, 021692796.

G.S. Coyne, *The Laboratory Companion: A Practical Guide to Materials, Equipment and Technique*, Wiley-Interscience, New York, 1997. ISBN 0471184225.

* L.M. Harwood and C.J. Moody, *Experimental Organic Chemistry: Principles and Practice*, Blackwell Scientific Publications, Oxford, 1989. ISBN 0632020164, 0632020172.

H.J.E. Loewenthal and E. Zass, *A Guide for the Perplexed Organic Experimentalist*, 2nd Edn, J. Wiley & Sons, Chichester, 1992. ISBN 0471935336.

D.W. Mayo, S.S. Butcher, P.K. Trumper, R.M. Pike and S.S. Butcher, *Microscale Techniques for the Organic Laboratory*, J. Wiley & Sons, New York, 1991. ISBN 0471621927.

C.E. Meloan, *Chemical Separations, Principles, Techniques and Experiments*, J. Wiley & Sons, New York, 2000, ISBN 0471351970.

J. Mendham, R.C. Denney, J.D. Barnes and M.J.K. Thomas, *Vogel's Quantitative Chemical Analysis*, 6th Edn, Prentice Hall, Harlow, 2000. ISBN 0582226287.

A.M. Schoffstall, B.A. Gaddis, M.L. Druelinger and M. Druelinger, *Organic Microscale and Miniscale Laboratory Experiments*, McGraw Hill, Boston, 2000. ISBN 0072375493.

J.T. Sharp, I. Gosney and A.G. Rowley, *Practical Organic Chemistry: A Student Handbook of Techniques*, Chapman and Hall, London, 1989. ISBN 0412282305.

* A.I. Vogel and B.S. Furniss, *Vogel's Textbook of Practical Organic Chemistry*, 5th Edn, J. Wiley & Sons, New York, 1989. ISBN 0582462363.

J.W. Zubrick, *The Organic Chemistry Lab Survival Guide*, 5th Edn, J. Wiley & Sons, New York, 2000. ISBN 0471387320.

MOLECULAR SIEVES

A. Dyer, *An Introduction to Zeolite Molecular Sieves*, J. Wiley & Sons, New York, 1988. ISBN 0471919810

R. Szostak, *Molecular Sieves: Principles of Synthesis and Identification*, Van Nostrand Reinhold, New York, 1989. ISBN 0442280238.

REAGENTS AND PURITY

D.J. Bucknell, *Analar Standards for Laboratory Chemicals*, 8th Edn, Analar Standards Ltd., Dorset, 1984. ISBN 095004394X.

K.L. Cheng, K. Ueno, T. Imamura (Eds), *CRC Handbook of Organic Analytical Reagents*, CRC Press, Boca Raton, 1982.

J.A. Dean (Ed.), *Lange's Handbook of Chemistry*, McGraw-Hill, New York, 1999. ISBN 0070163847.

*L.A. Paquette (Ed.), *The Encyclopedia of Reagents for Organic Synthesis*, 8 Volume set, J Wiley & Sons, New York, 1995. ISBN 0471936235.

*A.J. Pearson and W.R. Rouse, *Handbook of Reagents for Organic Synthesis*, 4 Vol. Set, J. Wiley & Sons, Chichester, 2001. ISBN 04719878911.

Reagents Chemicals: American Chemical Society Specifications, Official from January 1 2000 (Reagent Chemicals American Chemical Society Specification) 9th Edn, Washington DC, (Sept. 1999). ISBN 0841236712.

* L.E. Smart, *Separation, Purification and Identification*, The Royal Society of Chemistry, London, 2002 (in paperback and CD-ROM). ISBN 0854046852.

United States Pharmacopeia (USP # 24 NF19) 24-19 edition (January 1, 2000).

G. Wypych (Ed.), *Solvents Database on CD Rom*, William Andrew Publishing, LLC 2001. ISBN 0815514638.

SAFETY IN THE CHEMICAL LABORATORY

* L. Bretherick, P.G. Urben and M.J. Pitt, *Bretherick's Handbook of Reactive Chemical Hazards: An Indexed Guide to Published Data*, 6th Edn, Butterworth-Heinemann, Oxford, 1999. ISBN 075063605X.

N.P. Cheremisinoff, *Handbook of Hazardous Chemical Properties*, Butterworth-Heinemann, Oxford, 2000. ISBN 0750672099.

P. Patniak, *A Comprehensive Guide to the Hazardous Properties of Chemical Substances*, 2nd Edn, J. Wiley & Sons, 1999. ISBN 0471291757.

G.J. Hathaway, *Chemical Hazards of the Workplace*, 4th Edn, J. Wiley & Sons, 1996. ISBN 0471287024.

M.-A. Armour, *Hazardous Laboratory Chemicals: Disposal Guide*, 2nd Edn, CRC Press, Boca Raton, 1996. ISBN 1566701082.

S.G. Luxon (Ed.), *Hazards in the Chemical Laboratory*, 5th Edn, Royal Society of Chemistry, 1992. ISBN 0851862292.

R.E. Lenga (Ed.), *The Sigma-Aldrich Library of Chemical Safety Data*, 3 Volume set, Aldrich Chemical Co., Milwaukee, 1988. ISBN 0941633160.

R.E. Lenga, K.L. Votoupal (Eds), *The Sigma-Aldrich Library of Regulatory and Safety Data*, 3 Volume set, Aldrich Chemical Co., Milwaukee, 1993. ISBN 0941633357.

R.J. Lewis Sr., *Sax's Dangerous Properties of Industrial Materials*, 10th Edn, 3 Vol. Set., J. Wiley & Sons, New York, 1999. ISBN 0471354074.

R.J. Lewis Sr., *Hazardous Chemicals Desk Reference*, 5th Edn, J. Wiley & Sons, 2002. ISBN 0471441651.

SOLVENTS, SOLVENT EXTRACTION AND DISTRIBUTION

L.C. Craig, D. Craig and E.G. Scheibel (Chapter 2), in A. Weissberger's (Ed.) *Techniques of Organic Chemistry*, vol III, pt I, 2nd Edn, Interscience, New York, 1956.

D.R. Lide, *Handbook of Organic Solvents*, CRC Press, Florida, 1995. ISBN 0849389305.

R.P. Pohanish and S.A. Greene, *Rapid Guide to Chemical Incompatibilities*, J. Wiley & Sons, New York, 1997. ISBN 0471288020.

SPECTROSCOPY

E. Breitmaier, *Structure Elucidation by NMR in Organic Chemistry: A Practical Guide*, J. Wiley & Sons, New York, 1993. ISBN 0471937452, 0471933813.

K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 2 Vol. Set, 5th Edn, J. Wiley & Sons, New York, 1997. ISBN 0471194069.

* C.J. Pouchert and J. Behnke, *The Aldrich Library of ^{13}C and ^1H FT NMR Spectra*, 3 Volume set, Aldrich Chemical Co., Milwaukee, 1993. ISBN 0941633349.

* C.J. Pouchert, *The Aldrich Library of Infrared Spectra*, 3rd Edn, Aldrich Chemical Co., Milwaukee, 1981.

THIN LAYER CHROMATOGRAPHY

B. Fried and J. Sherma, *Thin Layer Chromatography*, Marcel Dekker, New York, 1999. ISBN 0824702220.

B. Fried, J. Sherma (Eds), *Practical Thin-Layer Chromatography: A MultiDisciplinary Approach*, CRC Press, Boca Raton, 1996. ISBN 0849326605.

F. Geiss, *Fundamentals of Thin Layer Chromatography*, Huthig, Heidelberg, 1987. ISBN 3778508547.

E. Hahn-Dienstrop, *Applied Thin Layer Chromatography: Best Practice and Avoidance of Mistakes*, VCH Verlagsgesellschaft MbH, Weinheim, 2000. ISBN 3527298398.

* H. Jork (Ed.), *Thin Layer Chromatography: Reagents and Detection Methods*, Vol. 1a, 1b, J. Wiley & Sons, 1989 (Vol. 1a) and 1994 (Vol. 1b). ISBN 3527278346 (Vol. 1a) and 352728205X (Vol. 1b).

CHAPTER 2

CHEMICAL METHODS

USED

IN PURIFICATION

GENERAL REMARKS

Greater selectivity in purification can often be achieved by making use of differences in chemical properties between the substance to be purified and the contaminants. Unwanted metal ions may be removed by precipitation in the presence of a *collector* (see p. 54). Sodium borohydride and other metal hydrides transform organic peroxides and carbonyl-containing impurities such as aldehydes and ketones in alcohols and ethers. Many classes of organic chemicals can be purified by conversion into suitable derivatives, followed by regeneration. This chapter describes relevant procedures.

REMOVAL OF TRACES OF METALS FROM REAGENTS

METAL IMPURITIES

The presence of metal contaminants in reagents may sometimes affect the chemical or biochemical outcomes of an experiment. In these cases, it is necessary to purify the reagents used.

Metal impurities can be determined qualitatively and quantitatively by atomic absorption spectroscopy and the required purification procedures can be formulated. Metal impurities in organic compounds are usually in the form of ionic salts or complexes with organic compounds and very rarely in the form of free metal. If they are present in the latter form then they can be removed by crystallising the organic compound (whereby the insoluble metal can be removed by filtration), or by distillation in which case the metal remains behind with the residue in the distilling flask. If the impurities are in the ionic or complex forms, then extraction of the organic compound in a suitable organic solvent with aqueous acidic or alkaline solutions will reduce their concentration to acceptable levels.

When the metal impurities are present in inorganic compounds as in metals or metal salts, then advantage of the differences in chemical properties should be taken. Properties of the impurities like the solubility, the solubility product (product of the metal ion and the counter-ion concentrations), the stability constants of the metal complexes with organic complexing agents and their solubilities in organic solvents should be considered. Alternatively the impurities can be masked by the addition of complexing agents which could lower the concentration of the metal ion impurities to such low levels that they would not interfere with the main compound (see **complexation** below). Specific procedures and examples are provided below.

DISTILLATION

Reagents such as water, ammonia, hydrochloric acid, nitric acid, perchloric acid, and sulfuric acid can be purified *via* distillation (preferably under reduced pressure and particularly with perchloric acid) using an all-glass still. Isothermal distillation is convenient for ammonia: a beaker containing concentrated ammonia is placed alongside a beaker of distilled water for several days in an empty desiccator so that some of the ammonia distils over into the water. The redistilled ammonia should be kept in polyethylene or paraffin-waxed bottles. Hydrochloric acid can be purified in the same way. To ensure the absence of metal contaminants from some salts (e.g. ammonium acetate), it may be more expedient to synthesise the salts using distilled components rather than to attempt to purify the salts themselves.

USE OF ION EXCHANGE RESINS

Application of ion-exchange columns has greatly facilitated the removal of heavy metal ions such as Cu^{2+} , Zn^{2+} and Pb^{2+} from aqueous solutions of many reagents. Thus, sodium salts and sodium hydroxide can be purified by passage through a column of a cation-exchange resin in its sodium form, prepared by washing the resin with 0.1M aqueous NaOH then washing with water until the pH of the effluent is ~ 7 . Similarly, for acids, a resin in its H^+ form [prepared by washing the column with 0.1M aqueous mineral acid (HCl , H_2SO_4) followed by thorough washing with water until the effluent has pH ~ 7 is used. In some cases, where metals form anionic complexes, they can be removed by passage through an anion-exchange resin. Iron in hydrochloric acid solution can be removed in this way.

Ion exchange resins are also useful for demineralising biochemical preparations such as proteins. Removal of metal ions from protein solutions using polystyrene-based resins, however, may lead to protein denaturation. This difficulty may be avoided by using a weakly acidic cation exchanger such as Bio-Rex 70.

Heavy metal contamination of pH buffers can be removed by passage of the solutions through a Chelex X-100 column. For example when a solution of 0.02M HEPES [4-(2-HydroxyEthyl)Piperazine-1-Ethanesulfonic acid] containing 0.2M KCl (1L, pH 7.5) alone or with calmodulin, is passed through a column of Chelex X-100 (60g) in the K^+ form, the level of Ca^{2+} ions falls to less than 2×10^{-7} M as shown by atomic absorption spectroscopy. Such solutions should be stored in polyethylene containers that have been washed with boiling deionised water (5min) and rinsed several times with deionised water. TES [*N,N,N',N'*-Tetraethylsulfamide] and TRIS [Tris-(hydroxymethyl)aminomethane] have been similarly decontaminated from metal ions.

Water, with very low concentrations of ionic impurities (and approaching conductivity standards), is very readily obtained by percolation through alternate columns of cation- and anion-exchange resins, or through a mixed-bed resin, and many commercial devices are available for this purpose. For some applications, this method is unsatisfactory because the final deionised water may contain traces of organic material after passage through the columns. However, organic matter can also be removed by using yet another special column in series for this purpose (see Milli Q water preparation, Millipore Corp., www.millipore.com).

PRECIPITATION

In removing traces of impurities by precipitation it is necessary to include a material to act as a *collector* of the precipitated substance so as to facilitate its removal by filtration or decantation. The following are a few examples:

Removal of lead contaminants

Aqueous hydrofluoric acid can be freed from lead by adding 1mL of 10% strontium chloride per 100mL of acid, lead being co-precipitated as lead fluoride with the strontium fluoride. If the hydrofluoric acid is decanted from the precipitate and the process repeated, the final lead content in the acid is less than 0.003 ppm. Similarly, lead can be precipitated from a nearly saturated sodium carbonate solution by adding 10% strontium chloride dropwise (1-2mL per 100mL) followed by filtration. (If the sodium carbonate is required as a solid, the solution can be evaporated to dryness in a platinum dish.) Removal of lead from potassium chloride uses precipitation as lead sulfide by bubbling H_2S , followed, after filtration, by evaporation and recrystallisation of the potassium chloride.

Removal of iron contaminants

Iron contaminants have been removed from potassium thiocyanate solutions by adding a slight excess of an aluminium salt, then precipitating aluminum and iron as their hydroxides by adding a few drops of ammonia. Iron is also carried down on the hydrated manganese dioxide precipitate formed in cadmium chloride or cadmium sulfate solutions by adding 0.5% aqueous potassium permanganate (0.5mL per 100mL of solution), sufficient ammonia to give a slight precipitate, and 1mL of ethanol. The solution is heated to boiling to coagulate the precipitate, then filtered. Ferrous ion can be removed from copper solutions by adding some hydrogen peroxide to the solution to oxidise the iron, followed by precipitation of ferric hydroxide by adding a small amount of sodium hydroxide.

Removal of other metal contaminants

Traces of calcium can be removed from solutions of sodium salts by precipitation at pH 9.5-10 as the 8-hydroxyquinolate. The excess 8-hydroxyquinoline acts as a *collector*.

EXTRACTION

In some cases, a simple solvent extraction is sufficient to remove a particular impurity. For example, traces of gallium can be removed from titanous chloride in hydrochloric acid by extraction with diisopropyl ether.

Similarly, ferric chloride can be removed from aluminium chloride solutions containing hydrochloric acid by extraction with diethyl ether. Usually, however, it is necessary to extract an undesired metal with an organic solvent in the presence of a suitable complexing agent such as dithizone (diphenylthiocarbazone) or sodium diethyl dithiocarbamate. When the former is used, weakly alkaline solutions of the substance containing the metal impurity are extracted with dithizone in chloroform (at about 25mg/L of chloroform) or carbon tetrachloride until the colour of some fresh dithizone solution remains unchanged after shaking. Dithizone complexes metals more strongly in weakly alkaline solutions. Excess dithizone in the aqueous medium is removed by extracting with the pure solvent (chloroform or carbon tetrachloride), the last traces of which, in turn, are removed by aeration. This method has been used to remove metal impurities from aqueous solutions of ammonium hydrogen citrate, potassium bromide, potassium cyanide, sodium acetate and sodium citrate. The advantage of dithizone for such a purpose lies in the wide range of metals with which it combines under these conditions. 8-Hydroxyquinoline (oxine) can also be used in this way. Sodium diethyl dithiocarbamate has been used to remove metals from aqueous hydroxylamine hydrochloride (made just alkaline to thymol blue by adding ammonia) from copper and other heavy metals by repeated extraction with chloroform until no more diethyl dithiocarbamate remained in the solution (which was then acidified to thymol blue by adding hydrochloric acid).

COMPLEXATION

Although not strictly a removal of an impurity, addition of a suitable complexing agent such as ethylenediaminetetraacetic acid often overcomes the undesirable effects of contaminating metal ions by reducing the concentrations of the free metal species to very low levels, i.e. sequestering metal ions by complexation. For a detailed discussion of this *masking*, see *Masking and Demasking of Chemical Reactions*, D.D.Perrin, Wiley-Interscience, New York, 1970.

USE OF METAL HYDRIDES

This group of reagents is commercially available in large quantities; some of its members - notably lithium aluminium hydride (LiAlH_4), calcium hydride (CaH_2), sodium borohydride (NaBH_4) and potassium borohydride (KBH_4) - have found widespread use in the purification of chemicals.

LITHIUM ALUMINIUM HYDRIDE

This solid is stable at room temperature, and is soluble in ether-type solvents. It reacts violently with water, liberating hydrogen, and is a powerful drying and reducing agent for organic compounds. It reduces aldehydes, ketones, esters, carboxylic acids, peroxides, acid anhydrides and acid chlorides to the corresponding alcohols. Similarly, amides, nitriles, aldimines and aliphatic nitro compounds yield amines, while aromatic nitro compounds are converted to azo compounds. For this reason it finds extensive application in purifying organic chemical substances by the removal of water and carbonyl containing impurities as well as peroxides formed by autoxidation. Reactions can generally be carried out at room temperature, or in refluxing diethyl ether, at atmospheric pressure. *When drying organic liquids with this reagent it is important that the concentration of water in the liquid is below 0.1% - otherwise a violent reaction or EXPLOSION may occur. LiAlH_4 should be added cautiously to a cooled solution of organic liquid in a flask equipped with a reflux condenser.*

CALCIUM HYDRIDE

This powerful drying agent is suitable for use with hydrogen, argon, helium, nitrogen, hydrocarbons, chlorinated hydrocarbons, esters and higher alcohols.

SODIUM BOROHYDRIDE

This solid which is stable in dry air up to 300° , is a less powerful reducing agent than lithium aluminium hydride, from which it differs also by being soluble in hydroxylic solvents and to a lesser extent in ether-type solvents. Sodium borohydride forms a dihydrate melting at $36-37^\circ$, and its aqueous solutions decompose slowly unless stabilised to above pH 9 by alkali. (For example, a useful sodium borohydride solution is one that is nearly saturated at $30-40^\circ$ and containing 0.2% sodium hydroxide.) Its solubility in water is 25, 55 and 88g per 100mL of water at 0° , 25° and 60° , respectively. Boiling or acidification rapidly decomposes aqueous sodium borohydride solutions. The reagent, available either as a hygroscopic solid or as an aqueous sodium hydroxide solution, is useful as a water soluble reducing agent for aldehydes, ketones and organic peroxides. This explains its use for the removal of carbonyl-containing impurities and peroxides from alcohols, polyols, esters, polyesters, amino alcohols, olefins, chlorinated hydrocarbons, ethers, polyethers, amines (including aniline), polyamines and aliphatic sulfonates.

Purifications using sodium borohydride can be carried out conveniently using alkaline aqueous or methanolic solutions of sodium borohydride, allowing the reaction mixture to stand at room temperature for several hours. Other solvents that can be used with this reagent include isopropyl alcohol (without alkali), amines (including liquid ammonia, in which its solubility is 104g per 100g of ammonia at 25°, and ethylenediamine), diglyme, formamide, dimethylformamide and tetrahydrofurfuryl alcohol. Alternatively, the material to be purified can be percolated through a column of the borohydride. In the absence of water, sodium borohydride solutions in organic solvents such as dioxane or amines decompose only very slowly at room temperature. Treatment of ethers with sodium borohydride appears to inhibit peroxide formation.

POTASSIUM BOROHYDRIDE

Potassium borohydride is similar in properties and reactions to sodium borohydride, and can similarly be used as a reducing agent for removing aldehydes, ketones and organic peroxides. It is non-hygroscopic and can be used in water, ethanol, methanol or water-alcohol mixtures, provided some alkali is added to minimise decomposition, but it is somewhat less soluble than sodium borohydride in most solvents. For example, the solubility of potassium borohydride in water at 25° is 19g per 100mL of water (as compared to sodium borohydride, 55g).

PURIFICATION *via* DERIVATIVES

Relatively few derivatives of organic substances are suitable for use as aids to purification. This is because of the difficulty in regenerating the starting material. For this reason, we list below, the common methods of preparation of derivatives that can be used in this way.

Whether or not any of these derivatives is likely to be satisfactory for the use of any particular case will depend on the degree of difference in properties, such as solubility, volatility or melting point, between the starting material, its derivative and likely impurities, as well as on the ease with which the substance can be recovered. Purification *via* a derivative is likely to be of most use when the quantity of pure material that is required is not too large. Where large quantities (for example, more than 50g) are available, it is usually more economical to purify the material directly (for example, in distillations and recrystallisations).

The most generally useful purifications *via* derivatives are as follows:

ALCOHOLS

Aliphatic or aromatic alcohols are converted to solid esters. *p*-Nitrobenzoates are examples of convenient esters to form because of their sharp melting points, and the ease with which they can be recrystallised as well as the ease with which the parent alcohol can be recovered. The *p*-nitrobenzoyl chloride used in the esterification is prepared by refluxing dry *p*-nitrobenzoic acid with a 3 molar excess of thionyl chloride for 30min on a steam bath (*in a fume cupboard*). The solution is cooled slightly and the excess thionyl chloride is distilled off under vacuum, keeping the temperature below 40°. Dry toluene is added to the residue in the flask, then distilled off under vacuum, the process being repeated two or three times to ensure complete removal of thionyl chloride, hydrogen chloride and sulfur dioxide. (This freshly prepared *p*-nitrobenzoyl chloride cannot be stored without decomposition; it should be used directly.) A solution of the acid chloride (1mol) in dry toluene or alcohol-free chloroform (distilled from P₂O₅ or by passage through an activated Al₂O₃ column) under a reflux condenser is cooled in an ice bath while the alcohol (1mol), with or without a solvent (preferably miscible with toluene or alcohol-free chloroform), is added dropwise to it. When addition is over and the reaction subsides, the mixture is refluxed for 30min and the solvent is removed under reduced pressure. The solid ester is then recrystallised to constant melting point from toluene, acetone, low boiling point petroleum ether or mixtures of these, but not from alcohols.

Hydrolysis of the ester is achieved by refluxing in aqueous N or 2N NaOH solution until the insoluble ester dissolves. The solution is then cooled, and the alcohol is extracted into a suitable solvent, e.g. ether, toluene or alcohol-free chloroform. The extract is dried (CaSO₄, MgSO₄) and distilled, then fractionally distilled if liquid or recrystallised if solid. (The *p*-nitrobenzoic acid can be recovered by acidification of the aqueous layer.) In most cases where the alcohol to be purified can be readily extracted from ethanol, the hydrolysis of the ester is best achieved with N or 2N ethanolic NaOH or 85% aqueous ethanolic N NaOH. The former is prepared by dissolving the necessary alkali in a minimum volume of water and diluting with absolute alcohol. The ethanolic solution is refluxed for one to two hours and hydrolysis is complete when an aliquot gives a clear solution on dilution with four or five times its volume of water. The bulk of the ethanol is distilled off and the residue is

extracted as above. Alternatively, use can be made of ester formation with benzoic acid, toluic acid or 3,5-dinitrobenzoic acid, by the above method.

Other derivatives can be prepared by reaction of the alcohol with an acid anhydride. For example, phthalic or 3-nitrophthalic anhydride (1 mol) and the alcohol (1mol) are refluxed for half to one hour in a non-hydroxylic solvent, e.g. toluene or alcohol-free chloroform, and then cooled. The phthalate ester crystallises out, is precipitated by the addition of low boiling petroleum ether or is isolated by evaporation of the solvent. It is recrystallised from water, 50% aqueous ethanol, toluene or low boiling petroleum ether. Such an ester has a characteristic melting point and the alcohol can be recovered by acid or alkaline hydrolysis.

ALDEHYDES

The best derivative from which an aldehyde can be recovered readily is its bisulfite addition compound, the main disadvantage being the lack of a sharp melting point. The aldehyde (sometimes in ethanol) is shaken with a cold saturated solution of sodium bisulfite until no more solid adduct separates. The adduct is filtered off, washed with a little water, followed by alcohol. A better reagent to use is a freshly prepared saturated aqueous sodium bisulfite solution to which 75% ethanol is added to near-saturation. (Water may have to be added dropwise to render this solution clear.) With this reagent the aldehyde need not be dissolved separately in alcohol and the adduct is finally washed with alcohol. The aldehyde is recovered by dissolving the adduct in the least volume of water and adding an equivalent quantity of sodium carbonate (not sodium hydroxide) or concentrated hydrochloric acid to react with the bisulfite, followed by steam distillation or solvent extraction.

Other derivatives that can be prepared are the Schiff bases and semicarbazones. Condensation of the aldehyde with an equivalent of primary aromatic amine yields the Schiff base, for example aniline at 100° for 10-30min. Semicarbazones are prepared by dissolving semicarbazide hydrochloride (*ca* 1g) and sodium acetate (*ca* 1.5g) in water (8-10mL) and adding the aldehyde or ketone (0.5-1g) with stirring. The semicarbazone crystallises out and is recrystallised from ethanol or aqueous ethanol. These are hydrolysed by steam distillation in the presence of oxalic acid or better by exchange with pyruvic acid (Hershberg *J Org Chem* 13 542 1948) [see entry under Ketones].

AMINES

Picrates

The most versatile derivative from which the free base can be readily recovered is the picrate. This is very satisfactory for primary and secondary aliphatic amines and aromatic amines and is particularly so for heterocyclic bases. The amine, dissolved in water or alcohol, is treated with excess of a saturated solution of picric acid in water or alcohol, respectively, until separation of the picrate is complete. If separation does not occur, the solution is stirred vigorously and warmed for a few minutes, or diluted with a solvent in which the picrate is insoluble. Thus, a solution of the amine and picric acid in ethanol can be treated with petroleum ether to precipitate the picrate. Alternatively, the amine can be dissolved in alcohol and aqueous picric acid added. The picrate is filtered off, washed with water or ethanol and recrystallised from boiling water, ethanol, methanol, aqueous ethanol, methanol or chloroform. The solubility of picric acid in water and ethanol is 1.4 and 6.23 % respectively at 20°.

It is not advisable to store large quantities of picrates for long periods, *particularly when they are dry due to their potential EXPLOSIVE nature*. The free base should be recovered as soon as possible. The picrate is suspended in an excess of 2N aqueous NaOH and warmed a little. Because of the limited solubility of sodium picrate, excess hot water must be added. Alternatively, because of the greater solubility of lithium picrate, aqueous 10% lithium hydroxide solution can be used. The solution is cooled, the amine is extracted with a suitable solvent such as diethyl ether or toluene, washed with 5N NaOH until the alkaline solution remains colourless, then with water, and the extract is dried with anhydrous sodium carbonate. The solvent is distilled off and the amine is fractionally distilled (under reduced pressure if necessary) or recrystallised.

If the amines are required as their hydrochlorides, picrates can often be decomposed by suspending them in acetone and adding two equivalents of 10N HCl. The hydrochloride of the base is filtered off, leaving the picric acid in the acetone. Dowex No 1 anion-exchange resin in the chloride form is useful for changing solutions of the more soluble picrates (for example, of adenosine) into solutions of their hydrochlorides, from which sodium hydroxide precipitates the free base.

Salts

Amines can also be purified *via* their salts, e.g. hydrochlorides. A solution of the amine in dry toluene, diethyl ether, dichloromethane or chloroform is saturated with dry hydrogen chloride (generated by addition of concentrated sulfuric acid to dry sodium chloride, or to concentrated HCl followed by drying the gas through sulfuric acid, or from a hydrogen chloride cylinder) and the insoluble hydrochloride is filtered off and dissolved in water. The solution is made alkaline and the amine is extracted, as above. Hydrochlorides can also be prepared by dissolving the amine in ethanolic HCl and adding diethyl ether or petroleum ether. Where

hydrochlorides are too hygroscopic or too soluble for satisfactory isolation, other salts, e.g. nitrate, sulfate, bisulfate or oxalate, can be used.

Double salts

The amine (1mol) is added to a solution of anhydrous zinc chloride (1mol) in concentrated hydrochloric acid (42mL) in ethanol (200mL, or less depending on the solubility of the double salt). The solution is stirred for 1h and the precipitated salt is filtered off and recrystallised from ethanol. The free base is recovered by adding excess of 5-10N NaOH (to dissolve the zinc hydroxide that separates) and is steam distilled. Mercuric chloride in hot water can be used instead of zinc chloride and the salt is crystallised from 1% hydrochloric acid. Other double salts have been used, e.g. cuprous salts, but are not as convenient as the above salts.

N-Acetyl derivatives

Purification as their *N*-acetyl derivatives is satisfactory for primary, and to a limited extent secondary, amines. The base is refluxed with slightly more than one equivalent of acetic anhydride for half to one hour, cooled and poured into ice-cold water. The insoluble derivative is filtered off, dried, and recrystallised from water, ethanol, aqueous ethanol or benzene (CAUTION toxic!). The derivative can be hydrolysed to the parent amine by refluxing with 70% sulfuric acid for a half to one hour. The solution is cooled, poured onto ice, and made alkaline. The amine is steam distilled or extracted as above. Alkaline hydrolysis is very slow.

N-Tosyl derivatives

Primary and secondary amines are converted into their tosyl derivatives by mixing equimolar amounts of amine and *p*-toluenesulfonyl chloride in dry pyridine (ca 5-10mols) and allowing to stand at room temperature overnight. The solution is poured into ice-water and the pH adjusted to 2 with HCl. The solid derivative is filtered off, washed with water, dried (vac. desiccator) and recrystallised from an alcohol or aqueous alcohol solution to a sharp melting point. The derivative is decomposed by dissolving in liquid ammonia (*fume cupboard*) and adding sodium metal (in small pieces with stirring) until the blue colour persists for 10-15min. Ammonia is allowed to evaporate (*fume cupboard*), the residue treated with water and the solution checked that the pH is above 10. If the pH is below 10 then the solution has to be basified with 2N NaOH. The mixture is extracted with diethyl ether or toluene, the extract is dried (K₂CO₃), evaporated and the residual amine recrystallised if solid or distilled if liquid.

AROMATIC HYDROCARBONS

Adducts

Aromatic hydrocarbons can be purified as their picrates using the procedures described for amines. Instead of picric acid, 1,3,5-trinitrobenzene or 2,4,7-trinitrofluorenone can also be used. In all these cases, following recrystallisation, the hydrocarbon can be isolated either as described for amines or by passing a solution of the adduct through an activated alumina column and eluting with toluene or petroleum ether. The picric acid and nitro compounds are more strongly adsorbed on the column.

Sulfonation

Naphthalene, xylenes and alkyl benzenes can be purified by sulfonation with concentrated sulfuric acid and crystallisation of the sodium sulfonates. The hydrocarbon is distilled out of the mixture with superheated steam.

CARBOXYLIC ACIDS

4-Bromophenacyl esters

A solution of the sodium salt of the acid is prepared. If the salt is not available, the acid is dissolved in an equivalent of aqueous NaOH and the pH adjusted to 8-9 with this base. A solution of one equivalent of 4-bromophenacyl bromide (for a monobasic acid, two equivalents for a dibasic acid, etc) in ten times its volume of ethanol is then added. The mixture is heated to boiling, and, if necessary, enough ethanol is added to clarify the solution which is then refluxed for half an hour to three hours depending on the number of carboxylic groups that have to be esterified. (One hour is generally sufficient for monocarboxylic acids.) On cooling, the ester should crystallise out. If it does not, then the solution is heated to boiling, and enough water is added to produce a slight turbidity. The solution is again cooled. The ester is collected, and recrystallised or fractionally distilled.

The ester is hydrolysed by refluxing for 1-2h with 1-5% of barium carbonate suspended in water or with aqueous sodium carbonate solution. The solution is cooled and extracted with diethyl ether, toluene or chloroform. It is then acidified and the acid is collected by filtration or extraction, and recrystallised or fractionally distilled.

p-Nitrobenzyl esters can be prepared in an analogous manner using the sodium salt of the acid and *p*-nitrobenzyl bromide. They are readily hydrolysed.

Alkyl esters

Of the alkyl esters, methyl esters are the most useful because of their rapid hydrolysis. The acid is refluxed with one or two equivalents of methanol in excess alcohol-free chloroform (or dichloromethane) containing about 0.1g of *p*-toluenesulfonic acid (as catalyst), using a Dean-Stark apparatus. (The water formed by the

esterification is carried away into the trap.) When the theoretical amount of water is collected in the trap, esterification is complete. The chloroform solution in the flask is washed with 5% aqueous sodium carbonate solution, then water, and dried over anhydrous sodium sulfate or magnesium sulfate. The chloroform is distilled off and the ester is fractionally distilled through an efficient column. The ester is hydrolysed by refluxing with 5-10% aqueous NaOH solution until the insoluble ester has completely dissolved. The aqueous solution is concentrated a little by distillation to remove all of the methanol. It is then cooled and acidified. The acid is either extracted with diethyl ether, toluene or chloroform, or filtered off and isolated as above. Other methods for preparing esters are available.

Salts

The most useful salt derivatives for carboxylic acids are the isothiuronium salts. These are prepared by mixing almost saturated solutions containing the acid (carefully neutralised with N NaOH using phenolphthalein indicator) then adding two drops of N HCl and an equimolar amount of *S*-benzylisothiuronium chloride in ethanol and filtering off the salt that crystallises out. After recrystallisation from water, alcohol or aqueous alcohol the salt is decomposed by suspending or dissolving in 2N HCl and extracting the carboxylic acid from aqueous solution into diethyl ether, chloroform or toluene.

HYDROPEROXIDES

These can be converted to their sodium salts by precipitation below 30° with aqueous 25% NaOH. The salt is then decomposed by addition of solid (powdered) carbon dioxide and extracted with low-boiling petroleum ether. The solvent should be removed under reduced pressure below 20°. **The manipulation should be adequately shielded at all times to guard against EXPLOSIONS for the safety of the operator.**

KETONES

Bisulfite adduct

The adduct can be prepared and decomposed as described for aldehydes. Alternatively, because no Cannizzaro reaction is possible, it can also be decomposed with 0.5N NaOH.

Semicarbazones

A powdered mixture of semicarbazide hydrochloride (1mol) and anhydrous sodium acetate (1.3mol) is dissolved in water by gentle warming. A solution of the ketone (1mol) in the minimum volume of ethanol needed to dissolve it is then added. The mixture is warmed on a water bath until separation of the semicarbazone is complete. The solution is cooled, and the solid is filtered off. After washing with a little ethanol followed by water, it is recrystallised from ethanol or dilute aqueous ethanol. The derivative should have a characteristic melting point. The semicarbazone is decomposed by refluxing with excess of oxalic acid or with aqueous sodium carbonate solution. The ketone (which steam distils) is distilled off. It is extracted or separated from the distillate (after saturating with NaCl), dried with CaSO₄ or MgSO₄ and fractionally distilled using an efficient column (under vacuum if necessary). [See entry under Aldehydes.]

PHENOLS

The most satisfactory derivatives for phenols that are of low molecular weight or monohydric are the benzoate esters. (Their acetate esters are generally liquids or low-melting solids.) Acetates are more useful for high molecular weight and polyhydric phenols.

Benzoates

The phenol (1mol) in 5% aqueous NaOH is treated (while cooling) with benzoyl chloride (1mol) and the mixture is stirred in an ice bath until separation of the solid benzoyl derivative is complete. The derivative is filtered off, washed with alkali, then water, and dried (in a vacuum desiccator over NaOH). It is recrystallised from ethanol or dilute aqueous ethanol. The benzylation can also be carried out in dry pyridine at low temperature (*ca* 0°) instead of in NaOH solution, finally pouring the mixture into water and collecting the solid as above. The ester is hydrolysed by refluxing in an alcohol (for example, ethanol, *n*-butanol) containing two or three equivalents of the alkoxide of the corresponding alcohol (for example sodium ethoxide or sodium *n*-butoxide) and a few (*ca* 5-10) millilitres of water, for half an hour to three hours. When hydrolysis is complete, an aliquot will remain clear on dilution with four to five times its volume of water. Most of the solvent is distilled off. The residue is diluted with cold water and acidified, and the phenol is steam distilled. The latter is collected from the distillate, dried and either fractionally distilled or recrystallised.

Acetates

These can be prepared as for the benzoates using either acetic anhydride with 3N NaOH or acetyl chloride in pyridine. They are hydrolysed as described for the benzoates. This hydrolysis can also be carried out with aqueous 10% NaOH solution, completion of hydrolysis being indicated by the complete dissolution of the acetate in the aqueous alkaline solution. On steam distillation, acetic acid also distils off but in these cases the phenols (see above) are invariably solids which can be filtered off and recrystallised.

PHOSPHATE AND PHOSPHONATE ESTERS

These can be converted to their uranyl nitrate addition compounds. The crude or partially purified ester is saturated with uranyl nitrate solution and the adduct filtered off. It is recrystallised from *n*-hexane, toluene or ethanol. For the more soluble members crystallisation from hexane using low temperatures (-40°) has been successful. The adduct is decomposed by shaking with sodium carbonate solution and water, the solvent is steam distilled (if hexane or toluene is used) and the ester is collected by filtration. Alternatively, after decomposition, the organic layer is separated, dried with CaCl₂ or BaO, filtered, and fractionally distilled under high vacuum.

MISCELLANEOUS

Impurities can sometimes be removed by conversion to derivatives under conditions where the major component does not react or reacts much more slowly. For example, normal (straight-chain) paraffins can be freed from unsaturated and branched-chain components by taking advantage of the greater reactivity of the latter with chlorosulfonic acid or bromine. Similarly, the preferential nitration of aromatic hydrocarbons can be used to remove e.g. benzene or toluene from cyclohexane by shaking for several hours with a mixture of concentrated nitric acid (25%), sulfuric acid (58%), and water (17%).

GENERAL METHODS FOR THE PURIFICATION OF CLASSES OF COMPOUNDS

Chapters 4, 5 and 6 list a large number of individual compounds, with a brief statement of how each one may be purified. For substances that are not included in these chapters the following procedures may prove helpful.

PROCEDURES

If the laboratory worker does not know of a reference to the preparation of a commercially available substance, he may be able to make a reasonable guess at the synthetic method used from published laboratory syntheses. This information, in turn, can simplify the necessary purification steps by suggesting probable contaminants. However, for other than macromolecules it is important that *at least* the ¹H NMR spectrum and/or the mass spectrum of the substance should be measured. These measurements require no more than two to three milligrams of material and provide a considerable amount of information about the substance. From the bibliography at the end of this chapter, references to NMR, IR and mass spectral data for a large number of the compounds in the Aldrich catalogue are available and are extremely useful for identifying compounds and impurities. If the material appears to have several impurities these spectra should be followed by examination of their chromatographic properties and spot tests. Purification methods can then be devised to remove these impurities, and a monitoring method will have already been established.

Physical methods of purification depend largely on the melting and boiling points of the materials. For gases and low-boiling liquids use is commonly made of the *freeze-pump-thaw* procedure (see Chapter 1). Gas chromatography is also useful, especially for low-boiling point liquids. Liquids are usually purified by refluxing with drying agents, acids or bases, reducing agents, charcoal, etc., followed by fractional distillation under reduced pressure. For solids, general methods include fractional freezing of the melted material, taking the middle fraction. Another procedure is sublimation of the solid under reduced pressure. The other commonly used method for purifying solids is by recrystallisation from a solution in a suitable solvent, by cooling with or without the prior addition of a solvent in which the solute is not very soluble.

The nature of the procedure will depend to a large extent on the quantity of purified material that is required. For example, for small quantities (50-250mg) of a pure volatile liquid, preparative gas chromatography is probably the best method. Two passes through a suitable column may well be sufficient. Similarly, for small amounts (100-500mg) of an organic solid, column chromatography is likely to be very satisfactory, the eluate being collected as a number of separate fractions (*ca* 5-10mL) which are examined by FT-IR, NMR or UV spectroscopy, TLC or by some other appropriate analytical technique. (For information on suitable adsorbents and eluents the texts referred to in the bibliography at the end of Chapters 1 and 2 should be consulted.) Preparative thin layer chromatography or HPLC can also be used successfully for purifying up to 500mg of solid. HPLC is the more and more commonly used procedure for the purification of small molecules as well as large molecules such as polypeptides and DNA.

Where larger quantities (upwards of 1g) are required, most of the impurities should be removed by preliminary treatments, such as solvent extraction, liquid-liquid partition, or conversion to a derivative (*vide supra*) which can be purified by crystallisation or fractional distillation before being reconverted to the starting material. The substance is then crystallised or distilled. If the final amounts must be in excess of 25g, preparation of a derivative is sometimes omitted because of the cost involved. In all of the above cases, purification is likely to be more laborious if the impurity is an isomer or a derivative with closely similar physical properties.

CRITERIA OF PURITY

Purification becomes meaningful only insofar as adequate tests of purity are applied: the higher the degree of purity that is sought, the more stringent must these tests be. If the material is an organic solid, its melting point should first be taken and compared with the recorded value. *Note that the melting points of most salts, organic or inorganic, are generally decomposition points and are not reliable criteria of purity.* As part of this preliminary examination, the sample might be examined by thin layer chromatography in several different solvent systems and in high enough concentrations to facilitate the detection of minor components. On the other hand, if the substance is a liquid, its boiling point should be measured. If, further, it is a high boiling liquid, its chromatographic behaviour should be examined. Liquids, especially volatile ones, can be studied very satisfactorily by gas chromatography, preferably using at least two different stationary and/or mobile phases. Atomic absorption spectroscopy is a useful and sensitive method for detecting metal impurities and the concentrations of metals and metal salts or complexes.

Application of these tests at successive steps will give a good indication of whether or not the purification is satisfactory and will also show when adequate purification has been achieved. Finally elemental analyses, e.g. of carbon, hydrogen, nitrogen, sulfur, metals etc. are very sensitive to impurities (other than with isomers), and are good criteria of purity.

GENERAL PROCEDURES FOR THE PURIFICATION OF SOME CLASSES OF ORGANIC COMPOUNDS

In the general methods of purification described below, it is assumed that the impurities belong essentially to a class of compounds different from the one being purified. They are suggested for use in cases where substances are not listed in Chapters 4, 5 and the low molecular weight compounds in Chapter 6. In such cases, the experimenter is advised to employ them in conjunction with information given in these chapters for the purification of suitable analogues. Also, for a wider range of drying agents, solvents for extraction and solvents for recrystallisation, the reader is referred to Chapter 1. See Chapter 6 for general purification procedures used for macromolecules.

ACETALS

These are generally diethyl or dimethyl acetal derivatives of aldehydes. They are more stable to alkali than to acids. Their common impurities are the corresponding alcohol, aldehyde and water. Drying with sodium wire removes alcohols and water, and polymerizes aldehydes so that, after decantation, the acetal can be fractionally distilled. In cases where the use of sodium is too drastic, aldehydes can be removed by shaking with alkaline hydrogen peroxide solution and the acetal is dried with sodium carbonate or potassium carbonate. Residual water and alcohols (up to *n*-propyl) can be removed with Linde type 4A molecular sieves. The acetal is then filtered and fractionally distilled. Solid acetals (i.e. acetals of high molecular weight aldehydes) are generally low-melting and can be recrystallised from low-boiling petroleum ether, toluene or a mixture of both.

ACIDS

Carboxylic acids

Liquid carboxylic acids are first freed from neutral and basic impurities by dissolving them in aqueous alkali and extracting with diethyl ether. (The pH of the solution should be at least three units above the pK_a of the acid, see pK in Chapter 1). The aqueous phase is then acidified to a pH at least three units below the pK_a of the acid and again extracted with ether. The extract is dried with magnesium sulfate or sodium sulfate and the ether is distilled off. The acid is fractionally distilled through an efficient column. It can be further purified by

conversion to its methyl or ethyl ester (*vide supra*) which is then fractionally distilled. Hydrolysis yields the original acid which is again purified as above.

Acids that are solids can be purified in this way, except that distillation is replaced by repeated crystallisation (preferable from at least two different solvents such as water, alcohol or aqueous alcohol, toluene, toluene/petroleum ether or acetic acid.) Water-insoluble acids can be partially purified by dissolution in N sodium hydroxide solution and precipitation with dilute mineral acid. If the acid is required to be free from sodium ions, then it is better to dissolve the acid in hot N ammonia, heat to *ca* 80°, adding slightly more than an equal volume of N formic acid and allowing to cool slowly for crystallisation. Any ammonia, formic acid or ammonium formate that adhere to the acid are removed when the acid is dried in a vacuum — they are volatile. The separation and purification of naturally occurring fatty acids, based on distillation, salt solubility and low temperature crystallisation, are described by K.S.Markley (Ed.), *Fatty Acids*, 2nd Edn, part 3, Chap. 20, Interscience, New York, 1964.

Aromatic carboxylic acids can be purified by conversion to their sodium salts, recrystallisation from hot water, and reconversion to the free acids.

Sulfonic acids

The low solubility of sulfonic acids in organic solvents and their high solubility in water makes necessary a treatment different from that for carboxylic acids. Sulfonic acids are strong acids, they have the tendency to hydrate, and many of them contain water of crystallisation. The lower-melting and liquid acids can generally be purified with only slight decomposition by fractional distillation, preferably under reduced pressure. A common impurity is sulfuric acid, but this can be removed by recrystallisation from concentrated aqueous solutions. The wet acid can be dried by azeotropic removal of water with toluene, followed by distillation. The higher-melting acids, or acids that melt with decomposition, can be recrystallised from water or, occasionally, from ethanol. For a typical purification of aromatic sulfonic acids using their barium salts refer to benzenesulfonic acid in Chapter 4.

Sulfinic acids

These acids are less stable, less soluble and less acidic than the corresponding sulfonic acids. The common impurities are the respective sulfonyl chlorides from which they have been prepared, and the thioisulfonates (neutral) and sulfonic acids into which they decompose. The first two of these can be removed by solvent extraction from an alkaline solution of the acid. On acidification of an alkaline solution, the sulfinic acid crystallises out leaving the sulfonic acid behind. The lower molecular weight members are isolated as their metal (e.g. ferric) salts, but the higher members can be crystallised from water (made slightly acidic), or alcohol.

ACID CHLORIDES

The corresponding acid and hydrogen chloride are the most likely impurities. Usually these can be removed by efficient fractional distillation. Where acid chlorides are not readily hydrolysed (e.g. aryl sulfonyl chlorides) the compound can be freed from contaminants by dissolving in a suitable solvent such as alcohol-free chloroform, dry toluene or petroleum ether and shaking with dilute sodium bicarbonate solution. The organic phase is then washed with water, dried with anhydrous sodium sulfate or magnesium sulfate, and distilled or recrystallised. This procedure is *hazardous* with readily hydrolysable acid chlorides such as acetyl chloride and benzoyl chloride. Solid acid chlorides should be thoroughly dried *in vacuo* over strong drying agents and are satisfactorily recrystallised from toluene, toluene-petroleum ether, petroleum ethers, alcohol-free chloroform/toluene, and, occasionally, from dry diethyl ether. Hydroxylic or basic solvents should be strictly avoided. *All operations should be carried out in a fume cupboard because of the irritant nature of these compounds which also attack the skin.*

ALCOHOLS

Monohydric alcohols

The common impurities in alcohols are aldehydes or ketones, and water. [*Ethanol* in Chapter 4 is typical.] Aldehydes and ketones can be removed by adding a small amount of sodium metal and refluxing for 2 hours, followed by distillation. Water can be removed in a similar way but it is preferable to use magnesium metal instead of sodium because it forms a more insoluble hydroxide, thereby shifting the equilibrium more completely from metal alkoxide to metal hydroxide. The magnesium should be activated with iodine (or a small amount of methyl iodide), and the water content should be low, otherwise the magnesium will be deactivated. If the amount of water is large it should be removed by azeotropic distillation (see below), or by drying over anhydrous $MgSO_4$ (not $CaCl_2$ which combines with alcohols). Acidic materials can be removed by treatment

with anhydrous Na_2CO_3 , followed by a suitable drying agent, such as calcium hydride, and fractional distillation, using gas chromatography to establish the purity of the product [Ballinger and Long, *J Am Chem Soc* **82** 795 1960]. Alternatively, the alcohol can be refluxed with freshly ignited CaO for 4 hours and then fractionally distilled [McCurdy and Laidler, *Can J Chem* **41** 1867 1963].

With higher-boiling alcohols it is advantageous to add some freshly prepared magnesium ethoxide solution (only slightly more than required to remove the water), followed by fractional distillation. Alternatively, in such cases, water can be removed by azeotropic distillation with toluene. Higher-melting alcohols can be purified by crystallisation from methanol or ethanol, toluene/petroleum ether or petroleum ether. Sublimation in vacuum, molecular distillation and gas chromatography are also useful means of purification. For purification *via* derivatives, *vide supra*.

Polyhydric alcohols

These alcohols are more soluble in water than are monohydric alcohols. Liquids can be freed from water by shaking with type 4A Linde molecular sieves and can safely be distilled only under high vacuum. Carbohydrate alcohols can be crystallised from strong aqueous solution or, preferably, from mixed solvents such as ethanol/petroleum ether or dimethyl formamide/toluene. Crystallisation usually requires seeding and is extremely slow. Further purification can be effected by conversion to the acetyl or benzoyl derivatives which are much less soluble in water and which can readily be recrystallised, e.g. from ethanol. Hydrolysis of the acetyl derivatives, followed by removal of acetate or benzoate and metal ions by ion-exchange chromatography, gives the purified material. On no account should solutions of carbohydrates be concentrated above 40° because of darkening and formation of *caramel*. Ion exchange, charcoal or cellulose column chromatography has been used for the purification and separation of carbohydrates.

ALDEHYDES

Common impurities found in aldehydes are the corresponding alcohols, aldols and water from self-condensation, and the corresponding acids formed by autoxidation. Acids can be removed by shaking with aqueous 10% sodium bicarbonate solution. The organic liquid is then washed with water. It is dried with anhydrous sodium sulfate or magnesium sulfate and then fractionally distilled. Water soluble aldehydes must be dissolved in a suitable solvent such as diethyl ether before being washed in this way. Further purification can be effected *via* the bisulfite derivative (see pp. 57 and 59) or the Schiff base formed with aniline or benzidine. Solid aldehydes can be dissolved in diethyl ether and purified as above. Alternatively, they can be steam distilled, then sublimed and crystallised from toluene or petroleum ether.

AMIDES

Amides are stable compounds. The lower-melting members (such as acetamide) can be readily purified by fractional distillation. Most amides are solids which have low solubilities in water. They can be recrystallised from large quantities of water, ethanol, ethanol/ether, aqueous ethanol, chloroform/toluene, chloroform or acetic acid. The likely impurities are the parent acids or the alkyl esters from which they have been made. The former can be removed by thorough washing with aqueous ammonia followed by recrystallisation, whereas elimination of the latter is by trituration or recrystallisation from an organic solvent. Amides can be freed from solvent or water by drying below their melting points. These purifications can also be used for sulfonamides and acid hydrazides.

AMINES

The common impurities found in amines are nitro compounds (if prepared by reduction), the corresponding halides (if prepared from them) and the corresponding carbamate salts. Amines are dissolved in aqueous acid, the pH of the solution being at least three units below the pK_a value of the base to ensure almost complete formation of the cation. They are extracted with diethyl ether to remove neutral impurities and to decompose the carbamate salts. The solution is then made strongly alkaline and the amines that separate are extracted into a suitable solvent (ether or toluene) or steam distilled. The latter process removes coloured impurities. Note that chloroform cannot be used as a solvent for primary amines because, in the presence of alkali, poisonous carbylamines (isocyanides) are formed. However, chloroform is a useful solvent for the extraction of heterocyclic bases. In this case it has the added advantage that while the extract is being freed from the chloroform most of the moisture is removed with the solvent.

Alternatively, the amine may be dissolved in a suitable solvent (e.g. toluene) and dry HCl gas is passed through the solution to precipitate the amine hydrochloride. This is purified by recrystallisation from a suitable solvent mixture (e.g. ethanol/diethyl ether). The free amine can be regenerated by adding sodium hydroxide and isolated as above.

Liquid amines can be further purified *via* their acetyl or benzoyl derivatives (*vide supra*). Solid amines can be recrystallised from water, alcohol, toluene or toluene-petroleum ether. *Care should be taken in handling large quantities of amines because their vapours are harmful (possibly carcinogenic) and they are readily absorbed through the skin.*

AMINO ACIDS

Because of their zwitterionic nature, amino acids are generally soluble in water. Their solubility in organic solvents rises as the fat-soluble portion of the molecule increases. The likeliest impurities are traces of salts, heavy metal ions, proteins and other amino acids. Purification of these is usually easy, by recrystallisation from water or ethanol/water mixtures. The amino acid is dissolved in the boiling solvent, decolorised if necessary by boiling with 1g of acid-washed charcoal/100g amino acid, then filtered hot, chilled, and set aside for several hours to crystallise. The crystals are filtered off, washed with ethanol, then ether, and dried.

Amino acids have high melting or decomposition points and are best examined for purity by paper or thin layer chromatography. The spots are developed with ninhydrin. Customary methods for the purification of small quantities of amino acids obtained from natural sources (i.e. 1-5g) are ion-exchange chromatography (see Chapter 1). For general treatment of amino acids see Greenstein and Winitz [*The Amino Acids*, Vols 1-3, J. Wiley & Sons, New York 1961] and individual amino acids in Chapters 4 and 6.

A useful source of details such as likely impurities, stability and tests for homogeneity of amino acids is *Specifications and Criteria for Biochemical Compounds*, 3rd edn, National Academy of Sciences, USA, 1972.

ANHYDRIDES

The corresponding acids, resulting from hydrolysis, are the most likely impurities. Distillation from phosphorus pentoxide, followed by fractional distillation, is usually satisfactory. With high boiling or solid anhydrides, another method involves boiling under reflux for 0.5-1h with acetic anhydride, followed by fractional distillation. Acetic acid distils first, then acetic anhydride and finally the desired anhydride. Where the anhydride is a solid, removal of acetic acid and acetic anhydride at atmospheric pressure is followed by heating under vacuum. The solid anhydride is then either crystallised as for acid chlorides or (in some cases) sublimed in a vacuum. A preliminary purification when large quantities of acid are present in a solid anhydride (such as phthalic anhydride) is by preferential solvent extraction of the (usually) more soluble anhydride from the acid (e.g. with CHCl_3 in the case of phthalic anhydride). *All operations with liquid anhydrides should be carried out in a fume cupboard because of their LACHRYMATORY properties. Almost all anhydrides attack skin.*

CAROTENOIDS

These usually are decomposed by light, air and solvents, so that degradation products are probable impurities. Chromatography and adsorption spectra permit the ready detection of coloured impurities, and separations are possible using solvent distribution, chromatography or crystallisation. Thus, in partition between immiscible solvents, xanthophyll remains in 90% methanol while carotenes pass into the petroleum ether phase. For small amounts of material, thin-layer or paper chromatography may be used, while column chromatography is suitable for larger amounts. Colorless impurities may be detected by IR, NMR or mass spectrometry. The more common separation procedures are described by P. Karrer and E. Jucker in *Carotenoids*, E.A. Braude (translator), Elsevier, NY, 1950.

Purity can be checked by chromatography (on thin-layer plates, Kieselguhr, paper or columns), by UV or NMR procedures.

ESTERS

The most common impurities are the corresponding acid and hydroxy compound (i.e. alcohol or phenol), and water. A liquid ester from a carboxylic acid is washed with 2N sodium carbonate or sodium hydroxide to remove acid material, then shaken with calcium chloride to remove ethyl or methyl alcohols (if it is a methyl or ethyl ester). It is dried with potassium carbonate or magnesium sulfate, and distilled. Fractional distillation then removes residual traces of hydroxy compounds. This method does not apply to esters of inorganic acids (e.g. dimethyl sulfate) which are more readily hydrolysed in aqueous solution when heat is generated in the neutralisation of the excess acid. In such cases, several fractional distillations, preferably under vacuum, are usually sufficient.

Solid esters are easily crystallisable materials. It is important to note that esters of alcohols must be recrystallised either from non-hydroxylic solvents (e.g. toluene) or from the alcohol from which the ester is derived. Thus methyl esters should be crystallised from methanol or methanol/toluene, but not from ethanol, *n*-butanol or other alcohols, in order to avoid alcohol exchange and contamination of the ester with a second ester. Useful solvents for crystallisation are the corresponding alcohols or aqueous alcohols, toluene, toluene/petroleum ether, and chloroform (ethanol-free)/toluene. Esters of carboxylic acid derived from phenols

are more difficult to hydrolyse and exchange, hence any alcoholic solvent can be used freely. Sulfonic acid esters of phenols are even more resistant to hydrolysis: they can safely be crystallised not only from the above solvents but also from acetic acid, aqueous acetic acid or boiling *n*-butanol.

Fully esterified phosphoric acid and phosphonic acids differ only in detail from the above mentioned esters. Their major contaminants are alcohols or phenols, phosphoric or phosphonic acids (from hydrolysis), and (occasionally) basic material, such as pyridine, which is used in their preparation. Water-insoluble esters are washed thoroughly and successively with dilute acid (e.g. 0.2N sulfuric acid), water, 0.2N sodium hydroxide and water. After drying with calcium chloride they are fractionally distilled. Water-soluble esters should first be dissolved in a suitable organic solvent and, in the washing process, water should be replaced by saturated aqueous sodium chloride. Some esters (e.g. phosphate and phosphonate esters) can be further purified through their uranyl adducts (*vide supra*). Traces of water or hydroxy compounds can be removed by percolation through, or shaking with, activated alumina (about 100g/L of liquid solution), followed by filtration and fractional distillation in a vacuum. For high molecular weight esters (which cannot be distilled without some decomposition) it is advisable to carry out distillation at as low a pressure as possible. Solid esters can be crystallised from toluene or petroleum ether. Alcohols can be used for recrystallising phosphoric or phosphonic esters of phenols.

ETHERS

The purification of diethyl ether (see Chapter 4) is typical of liquid ethers. The most common contaminants are the alcohols or hydroxy compounds from which the ethers are prepared, their oxidation products (e.g. aldehydes), peroxides and water. Peroxides, aldehydes and alcohols can be removed by shaking with alkaline potassium permanganate solution for several hours, followed by washing with water, concentrated sulfuric acid [CARE], then water. After drying with calcium chloride, the ether is distilled. It is then dried with sodium or with lithium aluminium hydride, redistilled and given a final fractional distillation. The drying process should be repeated if necessary.

Alternatively, methods for removing peroxides include leaving the ether to stand in contact with iron filings or copper powder, shaking with a solution of ferrous sulfate acidified with N sulfuric acid, shaking with a copper-zinc couple, passage through a column of activated alumina, and refluxing with phenothiazine. Cerium(III) hydroxide has also been used.

A simple test for ether peroxides is to add 10mL of the ether to a stoppered cylinder containing 1mL of freshly prepared 10% solution of potassium iodide containing a drop of starch indicator. No colour should develop during one minute if free from peroxides. Alternatively, a 1% solution of ferrous ammonium sulfate, 0.1M in sulfuric acid and 0.01M in potassium thiocyanate should not increase appreciably in red colour when shaken with two volumes of the ether.

As a safety precaution against **EXPLOSION** (in case the purification has been insufficiently thorough) at least a quarter of the total volume of ether should remain in the distilling flask when the distillation is discontinued as peroxides are generally higher boiling. To minimize peroxide formation, ethers should be stored in dark bottles and, if they are liquids, they should be left in contact with type 4A Linde molecular sieves, in a cold place, over sodium amalgam. The rate of formation of peroxides depends on storage conditions and is accelerated by heat, light, air and moisture. The formation of peroxides is inhibited in the presence of diphenylamine, di-*tert*-butylphenol, or other antioxidants as stabiliser.

Ethers that are solids (e.g. phenyl ethers) can be steam distilled from an alkaline solution which will hold back any phenolic impurity. After the distillate is made alkaline with sodium carbonate, the insoluble ether is collected either by extraction (e.g. with chloroform, diethyl ether or toluene) or by filtration. It is then crystallised from alcohols, alcohol/petroleum ether, petroleum ether, toluene or mixtures of these solvents, sublimed in a vacuum and recrystallised if necessary.

HALIDES

Aliphatic halides are likely to be contaminated with halogen acids and the alcohols from which they have been prepared, whereas in aromatic halides the impurities are usually aromatic hydrocarbons, amines or phenols. In both groups the halogen is less reactive than it is in acid halides. Purification is by shaking with concentrated hydrochloric acid, followed by washing successively with water, 5% sodium carbonate or bicarbonate, and water. After drying with calcium chloride, the halide is distilled and then fractionally distilled using an efficient column. For a solid halide the above purification is carried out by dissolving it in a suitable solvent such as toluene. Solid halides can also be purified by chromatography using an alumina column and eluting with toluene or petroleum ether. They can be crystallised from toluene, petroleum ethers, toluene/petroleum ether or toluene/chloroform/petroleum ether. Care should be taken when handling organic halogen compounds because of their **TOXICITY**. It should be noted that methyl iodide is a cancer suspect.

Liquid aliphatic halides are obtained alcohol-free by distillation from phosphorus pentoxide. They are stored in dark bottles to prevent oxidation and, in some cases, the formation of phosgene.

A general method for purifying *chlorohydrocarbons* uses repeated shaking with concentrated sulfuric acid [CARE] until no further colour develops in the acid, then washing with water then a solution of sodium bicarbonate, followed by water again. After drying with calcium chloride, the chlorohydrocarbon is fractionally redistilled to constant boiling point.

HYDROCARBONS

Gaseous hydrocarbons are best freed from water and gaseous impurities by passage through suitable adsorbents and (if olefinic material is to be removed) oxidants such as alkaline potassium permanganate solution, followed by fractional cooling (see Chapter 1 for cooling baths) and fractional distillation at low temperature. To effect these purifications and also to store the gaseous sample, a vacuum line is necessary.

Impurities in hydrocarbons can be characterised and evaluated by gas chromatography and mass spectrometry. The total amount of impurities present can be estimated from the thermometric freezing curve.

Liquid aliphatic hydrocarbons are freed from aromatic impurities by shaking with concentrated sulfuric acid [CARE] whereby the aromatic compounds are sulfonated. Shaking is carried out until the sulfuric acid layer remains colourless for several hours. The hydrocarbon is then freed from the sulfuric acid and the sulfonic acids by separating the two phases and washing the organic layer successively with water, 2N sodium hydroxide, and water. It is dried with CaCl_2 or Na_2SO_4 , and then distilled. The distillate is dried with sodium wire, P_2O_5 , or metallic hydrides, or passage through a dry silica gel column, or preferably, and more safely, with molecular sieves (see Chapter 1) before being finally fractionally distilled through an efficient column. If the hydrocarbon is contaminated with olefinic impurities, shaking with aqueous alkaline permanganate is necessary prior to the above purification. Alicyclic and paraffinic hydrocarbons can be freed from water, non-hydrocarbon and aromatic impurities by passage through a silica gel column before the final fractional distillation. This may also remove isomers. (For the use of chromatographic methods to separate mixtures of aromatic, paraffinic and alicyclic hydrocarbons see references in the bibliography in Chapter 1 under *Chromatography, Gas Chromatography and High Performance Liquid Chromatography*). Another method of removing branched-chain and unsaturated hydrocarbons from straight-chain hydrocarbons depends on the much faster reaction of the former with chlorosulfonic acid.

Isomeric materials which have closely similar physical properties can be serious contaminants in hydrocarbons. With aromatic hydrocarbons, e.g. xylenes and alkyl benzenes, advantage is taken of differences in ease of sulfonation. If the required compound is sulfonated more readily, the sulfonic acid is isolated, crystallised (e.g. from water), and decomposed by passing superheated steam through the flask containing the acid. The sulfonic acid undergoes hydrolysis and the liberated hydrocarbon distils with the steam. It is separated from the distillate, dried, distilled and then fractionally distilled. For small quantities (10-100mg), vapour phase chromatography is the most satisfactory method for obtaining a pure sample (for column materials for packings see Chapter 1).

Azeotropic distillation with methanol or 2-ethoxyethanol (cellosolve) has been used to obtain highly purified saturated hydrocarbons and aromatic hydrocarbons such as xylenes and isopropylbenzenes.

Carbonyl-containing impurities can be removed from hydrocarbons (and other oxygen-lacking solvents such as CHCl_3 and CCl_4) by passage through a column of Celite 545 (100g) mixed with concentrated sulfuric acid (60mL). After first adding some solvent and about 10g of granular Na_2SO_4 , the column is packed with the mixture and a final 7-8cm of Na_2SO_4 is added at the top [Hornstein and Crowe, *Anal Chem* **34** 1037 1962]. Alternatively, Celite impregnated with 2,4-dinitrophenylhydrazine can be used.

With solid hydrocarbons such as naphthalene and polycyclic hydrocarbons, preliminary purification by sublimation in vacuum (or high vacuum if the substance is high melting), is followed by zone refining and finally by chromatography (e.g. on alumina) using low-boiling liquid hydrocarbon eluents. These solids can be recrystallised from alcohols, alcohol/petroleum ether or from liquid hydrocarbons (e.g. toluene) and dried below their melting points. Aromatic hydrocarbons that have been purified by zone melting include anthracene, biphenyl, fluoranthrene, naphthalene, perylene, phenanthrene, pyrene and terphenyl, among others. Some polycyclic hydrocarbons, e.g. benzpyrene, are CARCINOGENIC.

Olefinic hydrocarbons have a very strong tendency to polymerise and commercially available materials are generally stabilised, e.g. with hydroquinone. When distilling compounds such as vinylpyridine or styrene, the stabiliser remains behind and the purified olefinic material is more prone to polymerisation. The most common impurities are higher-boiling dimeric or polymeric compounds. Vacuum distillation in a nitrogen atmosphere not only separates monomeric from polymeric materials but in some cases also depolymerises the impurities. The distillation flask should be charged with a polymerisation inhibitor and the purified material should be used immediately or stored in the dark and mixed with a small amount of stabiliser (e.g. 0.1% of hydroquinone or di-*tert*-butylcatechol). It is also advisable to add to the flask a small amount (ca 5-10% by volume of liquid in the flask) of a ground mixture of Kieselguhr and NaCl which will provide nuclei for facilitating boiling and finally for cleaning the flask from insoluble polymeric residue (due to the presence of the water soluble NaCl).

IMIDES

Imides (e.g. phthalimide) can be purified by conversion to their potassium salts by reaction in ethanol with ethanolic potassium hydroxide. The imides are regenerated when the salts are hydrolysed with dilute acid. Like amides, imides readily crystallise from alcohols and, in some cases (e.g. quinolinic imide), from glacial acetic acid.

IMINO COMPOUNDS

These substances contain the $-C=NH$ group and, because they are strong, unstable bases, they are kept as their more stable salts, such as the hydrochlorides. (The free base usually hydrolyses to the corresponding oxo compound and ammonia.) Like amine hydrochlorides, the salts are purified by solution in alcohol containing a few drops of hydrochloric acid. After treatment with charcoal, and filtering, dry diethyl ether (or petroleum ether if ethanol is used) is added until crystallisation sets in. The salts are dried and kept in a vacuum desiccator.

KETONES

Ketones are more stable to oxidation than aldehydes and can be purified from oxidisable impurities by refluxing with potassium permanganate until the colour persists, followed by shaking with sodium carbonate (to remove acidic impurities) and distilling. Traces of water can be removed with type 4A Linde molecular sieves. Ketones which are solids can be purified by crystallisation from alcohol, toluene, or petroleum ether, and are usually sufficiently volatile for sublimation in vacuum. Ketones can be further purified *via* their bisulfite, semicarbazone or oxime derivatives (*vide supra*). The bisulfite addition compounds are formed only by aldehydes and methyl ketones but they are readily hydrolysed in dilute acid or alkali.

MACROMOLECULES See Chapter 6.

NITRILES

All purifications should be carried out in an efficient fume cupboard because of the TOXIC nature of these compounds.

Nitriles are usually prepared either by reacting the corresponding halide or diazonium salts with a cyanide salt or by dehydrating an amide. Hence, possible contaminants are the respective halide or alcohol (from hydrolysis), phenolic compounds, amines or amides. Small quantities of phenols can be removed by chromatography on alumina. More commonly, purification of liquid nitriles or solutions of solid nitriles in a solvent such as diethyl ether is by shaking with dilute aqueous sodium hydroxide, followed by washing successively with water, dilute acid and water. After drying with sodium sulfate, the solvent is distilled off. Liquid nitriles are best distilled from a small amount of P_2O_5 which, besides removing water, dehydrates any amide to the nitrile. About one fifth of the nitrile should remain in the distilling flask at the end of the distillation (*the residue may contain some inorganic cyanide*). This purification also removes alcohols and phenols. Solid nitriles can be recrystallised from ethanol, toluene or petroleum ether, or a mixture of these solvents. They can also be sublimed under vacuum. Preliminary purification by steam distillation is usually possible.

Strong alkali or heating with dilute acids may lead to hydrolysis of the nitrile, and should be avoided.

NITRO COMPOUNDS

Aliphatic nitro compounds are generally acidic. They are freed from alcohols or alkyl halides by standing for a day with concentrated sulfuric acid, then washed with water, dried with magnesium sulfate followed by calcium sulfate and distilled. The principal impurities are isomeric or homologous nitro compounds. In cases where the nitro compound was originally prepared by vapour phase nitration of the aliphatic hydrocarbon, fractional distillation should separate the nitro compound from the corresponding hydrocarbon. Fractional crystallisation is more effective than fractional distillation if the melting point of the compound is not too low.

The impurities present in aromatic nitro compounds depend on the aromatic portion of the molecule. Thus, benzene, phenols or anilines are probable impurities in nitrobenzene, nitrophenols and nitroanilines, respectively. Purification should be carried out accordingly. Isomeric compounds are likely to remain as impurities after the preliminary purifications to remove basic and acidic contaminants. For example, *o*-nitrophenol may be found in samples of *p*-nitrophenol. Usually, the *o*-nitro compounds are more steam volatile than the *p*-nitro isomers, and can be separated in this way. Polynitro impurities in mononitro compounds can be readily removed because of their relatively lower solubilities in solvents. With acidic or basic nitro compounds which cannot be separated in the above manner, advantage may be taken of their differences in pK values (see Chapter 1). The compounds can thus be purified by preliminary extractions with several sets of aqueous buffers

of known pH (see for example Table 19, Chapter 1) from a solution of the substance in a suitable solvent such as diethyl ether. This method is more satisfactory and less laborious the larger the difference between the pK value of the impurity and the desired compound. Heterocyclic nitro compounds require similar treatment to the nitroanilines. Neutral nitro compounds can be steam distilled.

NUCLEIC ACIDS See Chapter 6.

PHENOLS

Because phenols are weak acids, they can be freed from neutral impurities by dissolution in aqueous N sodium hydroxide and extraction with a solvent such as diethyl ether, or by steam distillation to remove the non-acidic material. The phenol is recovered by acidification of the aqueous phase with 2N sulfuric acid, and either extracted with ether or steam distilled. In the second case the phenol is extracted from the steam distillate after saturating it with sodium chloride (salting out). A solvent is necessary when large quantities of liquid phenols are purified. The phenol is fractionated by distillation under reduced pressure, preferably in an atmosphere of nitrogen to minimise oxidation. Solid phenols can be crystallised from toluene, petroleum ether or a mixture of these solvents, and can be sublimed under vacuum. Purification can also be effected by fractional crystallisation or zone refining. For further purification of phenols *via* their acetyl or benzoyl derivatives (*vide supra*).

POLYPEPTIDES AND PROTEINS See Chapter 6.

QUINONES

These are neutral compounds which are usually coloured. They can be separated from acidic or basic impurities by extraction of their solutions in organic solvents with aqueous basic or acidic solutions, respectively. Their colour is a useful property in their purification by chromatography through an alumina column with, e.g. toluene, as eluent. They are volatile enough for vacuum sublimation, although with high-melting quinones a very high vacuum is necessary. *p*-Quinones are stable compounds and can be recrystallised from water, ethanol, aqueous ethanol, toluene, petroleum ether or glacial acetic acid. *o*-Quinones, on the other hand, are readily oxidised. They should be handled in an inert atmosphere, preferably in the absence of light.

SALTS (ORGANIC)

With metal ions

Water-soluble salts are best purified by preparing a concentrated aqueous solution to which, after decolorising with charcoal and filtering, ethanol or acetone is added so that the salts crystallise. They are collected, washed with aqueous ethanol or aqueous acetone, and dried. In some cases, water-soluble salts can be recrystallised satisfactorily from alcohols. Water-insoluble salts are purified by Soxhlet extraction, first with organic solvents and then with water, to remove soluble contaminants. The purified salt is recovered from the thimble.

With organic cations

Organic salts (e.g. trimethylammonium benzoate) are usually purified by recrystallisation from polar solvents (e.g. water, ethanol or dimethyl formamide). If the salt is too soluble in a polar solvent, its concentrated solution should be treated dropwise with a miscible nonpolar, or less polar, solvent (see Table 8, Chapter 1) until crystallisation begins.

With sodium alkane sulfonates

Purified from sulfites by boiling with aqueous HBr. Purified from sulfates by adding BaBr₂. Sodium alkane disulfonates are finally pptd by addition of MeOH. [Pethybridge and Taba *J Chem Soc Faraday Trans 1* **78** 1331 1982].

SULFUR COMPOUNDS

Disulfides

These can be purified by extracting acidic and basic impurities with aqueous base or acid, respectively. However, they are somewhat sensitive to strong alkali which slowly cleaves the disulfide bond. The lower-melting members can be fractionally distilled under vacuum. The high members can be recrystallised from alcohol, toluene or glacial acetic acid.

Sulfones

Sulfones are neutral and very stable compounds that can be distilled without decomposition. They are freed from acidic and basic impurities in the same way as disulfides. The low molecular weight members are quite

soluble in water but the higher members can be recrystallised from water, ethanol, aqueous ethanol or glacial acetic acid.

Sulfoxides

These compounds are odourless, rather unstable compounds, and should be distilled under vacuum in an inert atmosphere. They are water-soluble but can be extracted from aqueous solution with a solvent such as diethyl ether.

Thioethers

Thioethers are neutral stable compounds that can be freed from acidic and basic impurities as described for disulfides. They can be recrystallised from organic solvents and distilled without decomposition. They have sulfurous odours.

Thiols

Thiols, or mercaptans, are stronger acids than the corresponding aliphatic hydroxy or phenolic compounds, but can be purified in a similar manner. However, care must be exercised in handling thiols to avoid their oxidation to disulfides. For this reason, purification is best carried out in an inert atmosphere in the absence of oxidising agents. Similarly, thiols should be stored out of contact with air. They can be distilled without change, and the higher-melting thiols (which are usually more stable) can be crystallised, e.g. from water or dilute alcohol. They oxidise readily in alkaline solution but can be separated from the disulfide which is insoluble in this medium. They should be stored in the dark below 0°. *All operations with thiols should be carried out in an efficient fume cupboard because of their very unpleasant odour and their TOXICITY.*

Thiolsulfonates (disulfoxides)

Thiolsulfonates are neutral and are somewhat light-sensitive compounds. Their most common impurities are sulfonyl chlorides (neutral) or the sulfinic acid or disulfide from which they are usually derived. The first can be removed by partial freezing or crystallisation, the second by shaking with dilute alkali, and the third by recrystallisation because of the higher solubility of the disulfide in solvents. Thiolsulfonates decompose slowly in dilute, or rapidly in strong, alkali to form disulfides and sulfonic acids. Thiolsulfonates also decompose on distillation but they can be steam distilled. The solid members can be recrystallised from water, alcohols or glacial acetic acid.

BIBLIOGRAPHY

[For earlier bibliographies see *Purification of Laboratory Chemicals*, 4th Edn, ISBN 0750628391 (1996, hardback) and 0750637617 (1997, paperback)].

CHARACTERISATION OF ORGANIC AND INORGANIC COMPOUNDS

M. Fieser and L. Fieser, *Reagents for Organic Synthesis*, J. Wiley & Sons, Inc., New York, Vol 1 1967 to Vol 15, 1990. ISSN 0271 616X.

F. Fiegl, *Spot Tests in Organic Analysis*, 7th Edn, Elsevier Science, New York, 1989. ISBN 0444402098.

A.J. Gordon and R.A. Ford, *The Chemist's Companion: A Handbook of Practical Data, Techniques, and References*, J. Wiley & Sons, New York, 1973. ISBN 0471315907.

W.L. Jolly, *The Synthesis and Characterisation of Inorganic Compounds*, Waveland Press, 1991. ISBN 0881335789.

R.B. King (Ed.), *Encyclopedia of Inorganic Chemistry*, (8 volumes), J. Wiley & Sons, New York, 1994. ISBN 0471936200.

D.R. Lide, *CRC Handbook of Chemistry and Physics*, 82nd Edn, CRC Press, Boca Raton, 2001. ISBN 0849304822.

D.R. Lide, *CRC Handbook of Chemistry and Physics on CD-ROM 2002 Version*, CRC Press, Boca Raton, 2002. ISBN 0849308798. CD-ROM (Windows PCs only) edition.

* J. Mendham, R.C. Denney, J.D. Barnes and M.J.K. Thomas, *Vogel's Quantitative Chemical Analysis*, 6th Edn, Prentice Hall, Harlow, 2000. ISBN 0582226287.

* L.A. Paquette (Ed.), *The Encyclopedia of Reagents for Organic Synthesis*, 8 Volume set, J. Wiley & Sons, New York, 1995. ISBN 0471936235.

R.M. Silverstein, F.X. Webster, and R. M. Silverstein, *Spectrometric Identification of Organic Compounds*, 6th Edn, J. Wiley & Sons, New York, 1998. ISBN 0471134570.

R.L. Shriner, C.K.F. Hermann, T.C. Morrill, D.Y. Curtin, and R.C. Fuson, *The Systematic Identification of Organic Compounds*, 7th Edn, J. Wiley & Sons, New York, 1997. ISBN 0471 59748.

METAL HYDRIDES

A. Dedieu, (Ed.), *Transition Metal Hydrides*, J. Wiley & Sons, New York, 1992. ISBN 0471187682

M.V.C. Sastri, B. Viswanathan, and S.S. Murthy, *Metal Hydrides*, Narosa Publishing House, New Delhi, 1998. ISBN 8173192294.

J. Seyden-Penne, *Reductions by the Alumino- and Borohydrides in Organic Synthesis*, 2nd Edn, J. Wiley & Sons, New York, 1997. ISBN 1560819391.

SPECTROSCOPY

F.W. McLafferty and D.B. Stauffer, *The Wiley/NBS Registry of Mass Spectral Data*, (7 volumes), J. Wiley & Sons, New York, 1989. ISBN 0471628867.

* C.J. Pouchert and J. Behnke, *The Aldrich Library of ^{13}C and ^1H FT NMR Spectra*, 3 Volume set, Aldrich Chemical Co., Milwaukee, 1993. ISBN 0941633349.

* C.J. Pouchert, *The Aldrich Library of FT-IR Spectra*, Aldrich Chemical Co., Milwaukee, 3 Volume set, 2nd Edn, 1997. ISBN 0941633349X.

H. Gunzler and M.H. Heise, *IR Spectroscopy*, Wiley-VCH, New York, 2000. ISBN 3527288961.

TRACE METAL ANALYSIS

I.S. Krull (Ed.), *Trace Metal Analysis and Speciation*, Elsevier Science, New York, 1991. ISBN 044488209X.

A.Varma, *Handbook of Atomic Absorption Analysis*, (2 volumes), CRC Press, Boca Raton, Florida, 1984. ISBN 084932985X, 0849329868.

B. Welz (translated by C. Skegg), *Atomic Absorption Spectrometry*, VCH, Weinheim, 1985. ISBN 0895734184.

CHAPTER 3

THE FUTURE OF PURIFICATION

INTRODUCTION

The essence of research is to seek answers wherever there are questions. Regardless of what the answers are the experiments to be conducted must be carried out with utmost care. For this, one must ensure that the quality of the reactants used and the products obtained are of the highest possible purity. In general terms, one can broadly categorise experimental chemistry and biological chemistry into the following areas:

Isolation and identification of substances (natural products from nature, protein purification and characterisation, etc).

Synthesis of substances (organic, or inorganic in nature; these substances may be known substances or new compounds).

Analysis of substances (this is a key process in the identification of new or known chemical and biological substances. Methods of analysis include spectroscopic methods, derivatisation and sequencing methods).

Measurements of particular properties of a compound or substance (enzyme kinetics, reaction kinetics, FACS, fluorescence-activated cell sorting, assay).

Impressive and sophisticated strategies, in the form of new reagents, catalysts and chemical transformations, are currently available for the syntheses of molecules. In recent years there is a deviation in focus from developing new synthetic routes and reactions to improving methods for carrying out reactions. In particular, traditional reactions can be carried out in new ways such that those efficiencies of reactions are greatly improved. The efficiencies of reactions can be measured in terms of the yields of the desired product(s), or in terms of the time taken to obtain the desired product(s). Some of the 'new' lateral ways of thinking to improve efficiencies of reactions recognise the importance of purification of products in the planning of a synthetic sequence. Thus methods such as solid phase synthesis, fluororous chemistry as well as the use of ionic liquids minimise purification procedures and thus improve the ability to rapidly access pure compounds. These techniques also contribute to the efficiencies of reactions in terms of yields. In looking ahead to synthesis in the 21st century, a brief outline of the key aspects of these techniques are presented. In time many commercially available chemicals will be prepared using methods described in this chapter, and knowledge of these now should be useful to the experimenter. Some of these compounds (e.g. peptides) have already been synthesised by such methods (e.g. SPPS, see below).

SOLID PHASE SYNTHESIS

Solid phase synthesis (SPS) has emerged as an important methodology for the rapid and efficient synthesis of molecules. The ease of work-up and purification procedures in solid phase as compared to solution phase chemistry, as well as the scope for combinatorial chemistry provides impetus for further development in this field. The earliest studies on solid phase chemistry were focused on solid phase peptide synthesis (SPPS). The concept of carrying out reactions on a polymer support as distinct to reactants in solution, was conceived by R.B. Merrifield who received the Nobel Prize in Chemistry in 1984 for his pioneering work. However since the mid 1990's, advances in solid phase chemistry have moved beyond the routine (often robotic) synthesis of small to medium peptides and oligonucleotides. SPOS (solid phase organic synthesis) has gained much prominence due to the wealth of compounds (combinatorial libraries) that can be synthesised rapidly. This is especially important for pharmaceutical companies, screening for compounds with certain biological profiles or for chemical companies, screening for new catalysts or reagents. In SPOS, it is envisaged that difficult reactions can be driven to completion by using a large excess of reagents, which are easily removed by filtration. Furthermore, expensive reagents in the form of catalysts or chiral auxiliaries may be recycled easily if supported on a polymer and hence solid phase reactions provide economy in terms of costs and labour. Another strength of SPS is the ease in purification procedures which generally involves filtration of polymer supported products (solid) from soluble

reaction components (liquid) in what is effectively a solid-liquid extraction. In the final step of the synthetic sequence, the desired product is then cleaved from the polymer support.

Despite the relative infancy in the development of solid phase reactions, a wide range of functionalised resins are commercially available. The main uses of these functionalised resins can be roughly classified as follows:

SOLID PHASE PEPTIDE SYNTHESIS (SPPS)

Extensive studies on the synthesis of peptides on solid phase have been carried out, so much so that the technique of SPPS can be reliably and routinely used for the synthesis of short peptides by novices in the field. A large number of resins and reagents have been developed specifically for this purpose, and much is known on problems and avoidance of racemisation, difficult couplings, compatibility of reagents and solvents. Methods for monitoring the success of coupling reactions are available. Automated synthesisers are available commercially (e.g. from Protein Technologies, Rainin Inst Inc, Tuscon AZ; protan@dakotacom.net) which can carry out as many as a dozen polypeptide syntheses simultaneously. The most satisfactory chemistry currently used is Fmoc (9-fluorenylmethoxycarbonyl) chemistry whereby the amino group of the individual amino acid residues is protected as the Fmoc. A large number of Fmoc-amino acids are commercially available as well as polymer resins to which the specific Fmoc-amino acid (which will eventually become the carboxy terminal residue of the peptide) is attached. With automated synthesisers, the solvent used is *N*-methylpyrrolidone and washings are carried out with dimethylformamide. Deprotection of the polypeptide is carried out with anhydrous trifluoroacetic acid (TFA). A cycle for one residue varies with the residue but can take an hour or more. This means that 70-80 mer polypeptides could take more than a week to prepare. This is not a serious drawback because several different polypeptides can be synthesised simultaneously. The success of the synthesis is dependent on the amino acid sequence since there are some twenty or more different amino acids and the facility of forming a peptide bond varies with the pair of residues involved. However, generally 70 to 80 mers are routinely prepared, and if the sequence is favourable, up to 120 mer polypeptides can be synthesised. After deprotection with TFA the polypeptide is usually purified by HPLC using a C18 column with reverse phase chromatography. There are many commercial firms that will supply custom made polypeptides at a price depending on the degree of purity required.

SOLID PHASE DEOXYRIBONUCLEOTIDE SYNTHESIS

The need for oligodeoxyribonucleotides mainly as primers for the preparation of deoxyribonucleic acids (DNA) and for DNA sequencing has resulted in considerable developments in oligodeoxyribonucleotide synthesis. The solid phase procedure is the method commonly used. Automated synthesisers are commercially available, but with the increase in the number of firms which will provide custom made oligodeoxyribonucleotides, it is often not economical to purchase a synthesiser to make one's own oligodeoxyribonucleotides. Unlike in polypeptide synthesis where there are some twenty different residues to "string" together, in DNA synthesis there are only four deoxyribonucleotides, consequently there is usually little difficulty in synthesising 100 mers in quantities from 10 µg to 10 milligrams of material. The deprotected deoxyribonucleic acid which is separated from the solid support is purified on an anion exchange column followed by reverse phase HPLC using C8 to C18 columns for desalting. As for the polypeptides, the cost of DNA will depend on the purification level required.

SOLID PHASE OLIGOSACCHARIDE SYNTHESIS

Although automated solid phase peptide and oligonucleotide synthetic procedures are well established, automated solid phase oligosaccharide synthesis is considerably more difficult. The current awareness of the importance of polysaccharides as surface recognition molecules and in glycoproteins and glycolipids has prompted much interest in oligosaccharide synthesis and some progress has been made (see Kochetkov *Russ Chem Rev* **69** 795 2000; Ito and Manabe *Curr Opin Chem Biol* **2** 701 1998; Seeberger and Danishefsky *Acc Chem Res* **31** 685 1998). A general method for automated oligosaccharide synthesis is not as yet available. An example of an automated synthesis of *specific* glycosides has been reported by Seeberger (*Science* **291** 1523 2001; see also Houlton *Chem Br* **38** (4) 46 2002).

SOLID PHASE ORGANIC SYNTHESIS (SPOS)

At the time of writing this book, SPOS is in an area of relative infancy but has considerable potential. One of the main difficulties in SPOS lies in the lack of techniques available to monitor reactions carried out on polymer supports. Unlike reactions in solution phase, reactions on solid support cannot be monitored with relative ease and this has hindered the progress as well as the efficacy of solid supported synthesis of small non-peptidic molecules. Despite these difficulties, a large body of studies is available for SPOS. Recent reviews incorporate

information on the types of reactions that can be carried out, as well as outline the difficulties and differences with SP (solid phase) reactions as compared with their solution phase counterparts (see bibliography). An interesting application of such procedures is the synthesis of polymeric esters (e.g. polycaprolactones, polyhydroxybutyrate, polylactates) and starch- and cellulose- like polymers using a plasticised starch support. These have been useful for making biodegradable trays and containers for foodstuffs (BenBrahim *Chem Br* 38(4) 40 2002).

POLYMER SUPPORTED REACTANTS

These have become of increasing importance in synthesis and a broad classification of polymer supported reactants are as follows: Polymer bound bases (e.g. dimethylaminopyridine, morpholine, piperidine); Polymer supported catalysts (e.g. Grubbs catalyst for metathesis reactions, palladium for hydrogenation reactions, tributylmethylammonium chloride for phase transfer reactions); Polymer supported condensation reagents (e.g. DEAD (diethyl azodicarboxylate) for Mitsunobu reactions, DEC [1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride] (or EDCI [1-ethyl-3-(3-di-methylaminopropyl) carbodiimide HCl]) for peptide synthesis, HOBt (1-hydroxybenzotriazole) for peptide synthesis; Polymer supported oxidizing agents (e.g. osmium tetroxide, perruthenate, pyridinium chlorochromate); Polymer supported reducing agents (e.g. borohydride, tributyltin fluoride); Polymer supported phosphines (for miscellaneous applications depending on the structure) and so on. Commercially available polymer supported reactants are identified in Chapters 4 and 5 of this book.

SCAVENGER RESINS

Though not as extensively utilised as polymer supported reactants, the use of resins to clean up reactions is gaining favour. The type of commercially available scavenger resins are electrophilic scavenger resins (e.g. benzaldehyde derivatised resins to scavenge amines; isocyanate resins to scavenge amines, anilines and hydrazines; tosyl chloride resins to scavenge nucleophiles) and nucleophilic scavenger resins (e.g. diethylenetriamine resins to scavenge acids, acid chlorides, anhydrides; sulfonyl amide resins to scavenge acids, acid chlorides, aldehydes, isocyanates and chloroformates).

RESIN SUPPORT

The common resin matrixes comprise of polystyrene crosslinked with divinylbenzene, graft polymers of polystyrene-polyethylene glycol (PS-PEG) and polyethyleneglycol acrylamide (PEGA) composite resins. For each type of resin matrixes, a range of functionalised polymer supports are available. In addition, a number of these resins are available with different percentage of crosslinking as well as a range of loadings of the reactive functionality. Polystyrene based resins are the most extensively used. Unfortunately these resins do not swell, i.e. do not imbibe water, in polar solvents such as water and methanol and thus cannot be used for carrying out reactions in these solvents. In contrast grafted PS-PEG resins swell in a range of solvents from toluene to water. Examples of grafted PS-PEG resins are NovaSyn® TG and NovaGel® resins. As the success of transformations to be carried out on SPOS depends in part on the swelling properties as well as the robustness of the resin, the choice of resin matrix to be used must be carefully considered. The swelling properties of a number of resin types in a variety of solvents have been documented (see NovaBiochem catalog and also Santini, Griffith and Qi *Tetrahedron Lett* 39 8951 1998). For example, the swelling of a polystyrene resin in DMF is 3 mL/g of resin as compared to that in dichloromethane which is 7 mL/g of resin. It is thought that swelling of resins in the order of greater than 4 mL/g constitutes a good solvent, between 2-4 mL/g a moderate solvent and that less than 2 mL/g a poor solvent choice for carrying out solid phase reactions.

Lightly crosslinked resins are less robust but have greater ability to swell in appropriate solvents. Typically a 1-2% crosslinked divinyl benzene polystyrene resin is employed in organic synthesis.

An extensive list of the commercially available resins is available from Sigma-Aldrich (www.sigmaaldrich.com), Novabiochem (www.novabiochem.com), Fluka and other chemical companies. Sigma-Aldrich and Novabiochem have excellent catalogs. In addition, the Novabiochem catalog and website are a rich source of useful technical information.

CHOICE OF RESIN FOR SPOS

There is a large range of resins available for SPOS. These resins are derivatised polymer supports with a range of linkers. The roles of linkers are (i) to provide point(s) of attachment for the tethered molecule, akin to a solid supported protecting group(s), (ii) to provide distance from the polymeric backbone in order to minimise interactions with the backbone, (iii) to enable cleavage of product molecules under conditions compatible with the stability of the molecules and the reaction conditions employed for chemical transformations. Hence in order to

choose an appropriate resin for use in SPOS, one would need to consider the nature of the attachment of the reactant molecule onto the solid support (e.g. in order to tether the carboxy group in a reactant as an ester linkage on a solid support, one may choose to use a hydroxy functionalised linker), the stability of the resin under conditions employed in the chemical transformations (e.g. issues of orthogonality - will the conditions utilised cause premature cleavage of the linker or premature cleavage of the products?), the solvents and reactants needed in the transformations (e.g. will the solvents swell the resin?), conditions of cleavage of products (e.g. will this cause racemisation or rearrangement of the product?), the functionality of the resultant product after cleavage (e.g. will cleavage of the product result in a residual functionality in the molecule?) and so on. Linkers which leave no residual functionalities in the products upon cleavage are known as *traceless* linkers and those which need to be activated in order to be cleaved are known as *safety catch* linkers. A fascinating array of linkers (commercial or otherwise) is available and some excellent reviews are cited in the bibliography at the end of this chapter.

COMBINATORIAL CHEMISTRY

The major impetus for the development of solid phase synthesis centers around applications in combinatorial chemistry. The notion that new drug leads and catalysts can be discovered in a high throughput fashion has been demonstrated many times over as is evidenced from the number of publications that have arisen (see references at the end of this chapter). A number of approaches to combinatorial chemistry exist. These include the split-mix method, serial techniques and parallel methods to generate libraries of compounds. The advances in combinatorial chemistry are also accompanied by sophisticated methods in deconvolution and identification of compounds from libraries. In a number of cases, innovative hardware and software has been developed for these purposes.

Depending on the size of the combinatorial library to be generated as well as the scale of the reactions to be carried out, a wide range of specialised glassware and equipment are commercially available. For example, in order to carry out parallel combinatorial synthesis, reaction stations equipped with temperature and stirring control are available from a number of sources (e.g. www.fisher.co.uk; www.radleys.com; www.sigmaaldrich.com). These reaction stations are readily adapted, using appropriate modules, for conditions under reflux or under inert atmosphere. For automated synthesis of large libraries of compounds, reactions can be carried out using reaction blocks on microtiter plates.

Ready to use CombiKits™ which contain a variety of pre-weighed building blocks are available from Aldrich Chemical Company.

MONITORING SOLID PHASE REACTIONS

This remains the bane of solid phase reactions. Unlike solution phase reactions, where the progress of reactions can be monitored rapidly *via* TLC, GC or HPLC methods, procedures for the rapid monitoring of progress in solid phase reactions are limited. Although a number of spectroscopic methods have been developed for direct monitoring of reactions on solid supports, these methods usually require specialised equipment, not routinely available in chemical laboratories. These methods include on-bead IR analysis (e.g. Huber et al. *Anal Chim Acta* **393** 213 1999; Yan and Gremlich *J Chromatogr. B.* **725** 91 1999; Yan et al. *J Org Chem* **60** 5736 1995) and solid state magic angle spinning NMR techniques (e.g. Warrass and Lippens *J Org Chem* **65** 2946 2000; Rousselot-Pailley, Ede and Lippens *J. Comb. Chem.* **3** 559 2001).

The most common methods for monitoring solid phase reactions utilized in normal research laboratories are:

Infrared analysis of resin

This is a destructive method in which the resin is ground and pelleted as a KBr disc and analysed by FT-IR analysis. This method works best for systems where distinct functional group transformations (C=O, C-OH, C=C, etc) are expected. No special equipment is needed.

Qualitative and quantitative analyses

There are a number of colour or UV tests which are available for monitoring the presence or absence of certain functional groups. Although some of these tests are routinely used for the quantitative analysis of functional groups in solution phase, the quantification on solid phase is less than reliable. An exception to this is the Fmoc (9-fluorenylmethoxycarbonyl) assay, which is routinely used for quantification of coupling in SPPS using Fmoc amino acids. It should also be noted that the generality of some of these colour tests on a variety of solid phase resins is not known and hence these tests serve only as a *guide* to functional group identification. Some (not an exhaustive list) of the reported methods of analyses are outlined below.

DETECTION OF REACTIVE GROUPS ON RESINS

Detection of hydroxy groups on resin

A method in which the resin is treated with cyanuric chloride (trichlorotriazine, TCT) in DMF followed by a nucleophilic dye (AliR or Mordant Orange 1, beads appear red in the presence of hydroxyl groups, or with fuschin, beads appear fuschia, or with fluorescein, they become fluorescent) has been reported (Attardi and Taddei *Tetrahedron Lett* **42** 2927 2001; Attardi, Falchi and Taddei *Tetrahedron Lett* **41** 7395 2001). Another colorimetric test for the detection of polymer supported tertiary alcohols utilizes the conversion of the alcohols to the polymer supported diphenylsilylchloride ether, followed by subsequent treatment with methyl red. The beads form a readily distinguishable orange/red colour. The test is also positive for the hydroxy Wang resin and the aminomethylpolystyrene resin [Burkett, Brown and Meloni *Tetrahedron Lett* **42** 5773 2001]. Alternatively the conversion of polymer supported alcohols to the tosylate followed by displacement by *p*-nitrobenzylpyridine (PNBP) gives a strongly coloured salt upon treatment with bases such as piperidine, followed by gentle heating [Kuisle et al. *Tetrahedron* **55** 14807 1999].

Detection of aldehyde groups on resin

The use of an acidic solution of *p*-anisaldehyde in ethanol to detect aldehyde functionalities on polystyrene polymer supports has been reported (beads are treated with a freshly made solution of *p*-anisaldehyde (2.55 mL), ethanol (88 mL), sulfuric acid (9 mL), acetic acid (1 mL) and heated at 110°C for 4 min). The colour of the beads depends on the percentage of CHO content such that at 0% of CHO groups, the beads are colourless, ~50% CHO content, the beads appear red and at 98% CHO the beads appear burgundy [Vázquez and Albericio *Tetrahedron Lett* **42** 6691 2001]. A different approach utilises 4-amino-3-hydrazino-5-mercapto-1,2,4-triazole (Purpald) as the visualizing agent for CHO groups. Resins containing aldehyde functionalities turn dark brown to purple after a 5 min reaction followed by a 10 minute air oxidation [Cournoyer et al. *J Comb Chem* **4** 120 2002].

Detection of carboxy groups on resin

The presence of a COOH functionality on a polystyrene resin can be detected using a 0.25% solution of malachite green-oxalate in ethanol in the presence of a drop of triethylamine. Beads with COOH functionalities are coloured dark green or appear as clear gel beads [Attardi, Porcu and Taddei *Tetrahedron Lett* **41** 7391 2000].

Detection of amino groups on resin

The methods for the detection of amine functional groups are well established. For example the Kaiser test can be used to detect the presence of amine groups on resins (blue colour is observed). In the Kaiser test, two reagents are prepared. Reagent 1 comprises of a mixture of two solutions: A and B. A is a solution of phenol in absolute ethanol (40g of phenol in 10 mL of absolute ethanol, followed by treatment of this clear solution with 4g of Amberlite mixed bed resin MB-3 for 45 mins. The solution is then filtered.). Solution B is made up of 65 mg of KCN in 100 mL water; 2 mL of this solution is diluted to 100 mL of freshly distilled pyridine. The solution is then stirred with 4 g of Amberlite mixed-bed resin MB-3 and filtered. Solutions A and B are then mixed. Reagent 2 is a solution of ninhydrin (2.5g) in absolute ethanol (50 mL). For a qualitative Kaiser test, 6 drops of reagent 1 and 2 drops of reagent 2 are added to the well washed dried resin (2-5 mg) and mixed, followed by heating to 100°C for 4-6 min. A method for the quantitative determination of amino groups using this test has also been reported [Sarin et al. *Anal Biochem* **117** 147 1981]. It is however known that the Kaiser test does not give a positive test with a secondary amino acid such as proline or some 'unnatural' amino acids. In addition some deprotected amino acids (Ser, Asn, Asp) do not show the expected intense blue colour typical of free primary amino groups.

A test for secondary amines (e.g. proline) is the Chloranil test (1 drop of a 2% acetaldehyde solution in DMF, followed by one drop of a 2% solution of *p*-chloranil in DMF, leave for 5 mins). A positive test gives blue stained beads.

Other tests for the detection of amino functionalities on solid supports include the TNBS (2,4,6-trinitrobenzenesulfonic acid, picrylsulfonic acid) [Hancock and Battersby *Anal Biochem* **71** 260 1976], the DABITC [Shah et al. *Anal. Commun.* **34** 325 1997] and the NF31 [Madder et al. *Eur J Org Chem* 2787 1999] tests.

Detection of thiol groups on resin

For quantitative analysis of solid supported thiol residues on free macroporous or PEG grafts, Ellman's reagent has been used [5,5'-dithio-bis-(2-nitrobenzoic acid)]. However only qualitative information can be gained using lightly crosslinked polystyrene resins [Badyal et al. *Tetrahedron Lett* **42** 8531 2001].

Fmoc assay

This is a very important and well tested method for the quantitative determination of loading of Fmoc protected compounds particularly that of Fmoc (fluorenylmethoxycarbonyl) amino acids on solid support. Fmoc groups can

be readily deprotected in the presence of base. Generally, in the deprotection and quantification procedures, a known amount of resin is treated with 20% piperidine in DMF at room temperature for 30 mins. The resin is washed with more DMF and the pooled filtrates are combined in a volumetric flask and made up to an accurate volume (e.g. to 10 mL) with more DMF. The UV absorbance at 301 nm of the piperidine-dibenzofulvene adduct which is formed can then be measured against a blank solution of piperidine in DMF [Meienhofer et al. *Int J Pept Protein Res* 13 35 1979]. The loading **L** is then determined using the equation :

$$L = \frac{(\text{Absorbance value}) \times (\text{Solution volume in litres})}{7.8 \times (\text{Weight of resin in mg})}$$

IONIC LIQUIDS

Ionic liquids are organic or inorganic salts that are liquid at room or reaction temperatures. Although ionic liquids are themselves not new discoveries (e.g. the ionic liquid [EtNH₃][NO₃] was described in 1914), the use of ionic liquids in synthesis is only recent. In particular, the potential applications of ionic liquids as solvents in synthesis and in catalysis have recently been realised. The physical properties of ionic liquids make them unique solvents for synthesis. For example, ionic liquids are good solvents for both organic and inorganic substances and hence can be used to bring reagents into the same phase for reaction. Ionic liquids are also immiscible with a number of organic solvents and thus provide a non-aqueous, polar alternative for two-phase extraction systems. As ionic liquids are non-volatile, they can be used in high vacuum systems without the possibility of loss or contaminants. In addition, this also facilitates the isolation of products as the products can be distilled from the ionic liquid or alternatively extracted with an organic solvent that is immiscible with the ionic liquid. Although ionic liquids are frequently composed of poorly coordinating ions, they are highly polar which are important characteristics in the activation of catalysts.

Commonly used ionic liquids are *N*-alkylpyridinium, *N,N'*-dialkylimidazolium, alkylammonium and alkylphosphonium salts.

To date a number of reactions have been carried out in ionic liquids [for examples, see Dell'Anna et al. *J Chem Soc, Chem Commun* 434 2002; Nara, Harjani and Salunkhe *Tetrahedron Lett* 43 1127 2002; Semeril et al. *J Chem Soc Chem Commun* 146 2002; Buijsman, van Vuuren and Sterrenburg *Org Lett* 3 3785 2001]. These include Diels-Alder reactions, transition-metal mediated catalysis, e.g. Heck and Suzuki coupling reactions, and olefin metathesis reactions. An example of ionic liquid acceleration of reactions carried out on solid phase is given by Revell and Ganesan [*Org Lett* 4 3071 2002].

FLUOROUS CHEMISTRY

This new approach to synthesis was introduced by Curran early in 1997 and involves the attachment of fluororous phase labels to substrates such that the subsequent fluorinated products can be extracted into the fluororous phase. For example in liquid-liquid extractions (typical work-up procedures), a three-phase extraction is now possible (organic, fluororous and aqueous phases). As organic and inorganic compounds have little or no tendency to dissolve in highly fluorinated solvents and compounds, phase labeling a compound as fluororous will enable successful extraction into the fluororous phase. However in order to carry out homogenous reactions with these fluorinated compounds, organic solvents with a good dissolving power for fluororous compounds or miscible organic and fluororous solvents can be used. Alternatively organic solvents with a few fluorine atoms e.g. trifluoroethanol, benzotrifluoride ('hybrid solvents') will dissolve both organic and fluororous compounds. A number of synthetic applications utilising fluororous chemistry have been reported in the literature. [For examples, see Schneider and Bannwarth *Helv Chim Acta* 84 735 2001; Galante, Lhoste and Sinou *Tetrahedron Lett* 42 5425 2001; Studer and Curran *Tetrahedron* 53 6681 1997; Studer et al. *J Org Chem* 62 2917 1997; Crich and Neelamkavil *Tetrahedron* 58 3865 2002].

BIBLIOGRAPHY

A SAMPLE OF REVIEWS ON SOLID PHASE SYNTHESIS

General

- C. Blackburn, Polymer supports for solid-phase organic synthesis, *Biopolymers* **47** 311-351 1998.
- R.J. Booth and J.C. Hodges, Solid-supported reagent strategies for rapid purification of combinatorial synthesis products, *Acc Chem Res* **32** 18-26 1999.
- J.A. Ellman, Design, Synthesis, and Evaluation of Small-Molecule Libraries, *Acc Chem. Res* **29** 132-143 1996.
- J.S. Fruchtel and G. Jung, Organic Chemistry On Solid Supports, *Angew Chem, Int Ed Engl* **35** 17-42 1996.
- K. Gordon and S. Balasubramanian, Solid phase synthesis - designer linkers for combinatorial chemistry: A review *J Chem Technol Biotechnol* **74** 835-851 1999.
- F. Guillier, D. Orain and M. Bradley, Linkers and Cleavage Strategies in Solid-phase Organic Synthesis and Combinatorial Chemistry, *Chem Rev* **100** 2091-2157 2000.
- P. Hodge, Polymer-supported Organic Reactions: What Takes Place In The Beads? *Chem Soc Rev* **26** 417-424 1997.
- I.W. James, Linkers for solid phase organic synthesis, *Tetrahedron* **55** 4855-4946 1999.
- J.W. Labadie, Polymeric Supports for Solid Phase Synthesis, *Curr Opin Chem Biol* **2** 346-352 1998.
- S.V. Ley, I.R. Baxendale, G. Brusotti, M. Caldarelli, A. Massi and M. Nesi, Solid-supported Reagents for Multi-step Organic Synthesis: Preparation and Application, *Il Farmaco* **57** 321-330 2002.
- P. Seneci, *Solid Phase and Combinatorial Technologies*, J. Wiley & Sons, New York, 2000, ISBN 0471331953.
- A.R. Vaino and K.D. Janda, Solid-Phase Organic Synthesis: A Critical Understanding of the Resin, *J Comb Chem* **2** 579-596 2000.

Applications to SPOS, SPPS

- P. Blaney, R. Grigg and V. Sridharan, Traceless Solid-Phase Organic Synthesis, *Chem Rev* **102** 2607-2624 2002.
- B. Clapham, T.S. Reger and K.D. Janda, Polymer-supported Catalysis in Synthetic Organic Chemistry, *Tetrahedron* **57** 4637-4662 2001.
- D.H. Drewry, D.M. Coe and S. Poon, Solid Supported Reagents in Organic Synthesis, *Med Res Rev* **19** 97-148 1999 .
- S.J. Shuttleworth, S.M. Allin and P.K. Sharma, Functionalised Polymers - Recent Developments and New Applications in Synthetic Organic Chemistry, *Synthesis* 1217-1239 1997.
- S.J. Shuttleworth, S.M. Allin, R.D. Wilson and D. Nasturica, Functionalised polymers in organic chemistry; Part 2, *Synthesis* **8** 1035-1074 2000.

Monitoring SP reactions

- Y. Bing, Monitoring the Progress and the Yield of Solid Phase Organic Reactions Directly On Resin Supports, *Acc Chem Res* **31** 621-630 1998.

M. Dal Cin, S. Davalli, C. Marchioro, M. Passarini, O. Perini, S. Provera and A. Zaramella, Analytical Methods for Monitoring of Solid Phase Organic Synthesis, *Il Farmaco* **57** 497-510 2002.

S.W. Gerritz, Quantitative Techniques for the Comparison of Solid Supports, *Curr Opin Chem Biol*, **5** 264-268 2001.

M. Irving, J. Cournoyer, R.S. Li, C. Santos and B. Yan, Qualitative and quantitative analyses of resin-bound organic compounds, *Comb Chem High Throughput Screening* **4** 353-362 2001.

COMBINATORIAL CHEMISTRY

F. Balkenhohl, C. Vondembusschehunnefeld, A. Lansky and C. Zechel, Combinatorial Synthesis of Small Organic Molecules, *Angew Chem, Int Ed Engl* **35** 2289-2337 1996.

S. Bhattacharyya, Polymer-supported reagents and catalysts: Recent advances in synthetic applications, *Comb Chem High Throughput Screening* **3** 65-92 2000.

A. Ganesan, Recent developments in combinatorial organic synthesis, *Drug Discovery Today* **7** 47-55 2002.

G. Lowe, Combinatorial Chemistry, *Chem Soc Rev* **24** 309-317 1995.

H. Wennemers, Combinatorial chemistry: A tool for the discovery of new catalysts, *Comb Chem High Throughput Screening* **4** 273-285 2001.

IONIC LIQUIDS

T. Welton, Room temperature ionic liquids. Solvents for synthesis and catalysis, *Chem Rev* **99** 2071-2083 1999.

C.M. Gordon, New developments in catalysis using ionic liquids, *Appl. Catal A: General* **222** 101-117 2001.

R. Sheldon, Catalytic reactions in ionic liquids, *J Chem Soc, Chem Commun* 2399-2407 2001.

FLUOROUS CHEMISTRY

D.P. Curran, Strategy-level separations in organic synthesis: from planning to practice, *Angew Chem, Int Ed Engl* **37** 1174-1196 1998.

D.P. Curran, Combinatorial Organic Synthesis and Phase Separation: Back to the Future, *Chemtracts: Org Chem* **9** 75-87 1996.

D.L. Flynn, Phase-trafficking reagents and phase-switching strategies for parallel synthesis, *Med Res Rev* **19** 408-431 1999.

A. Studer, S. Hadida, R. Ferritto, S.Y. Kim, P. Jeyer, P. Wipf, D.P. Curran, Fluorous synthesis — A fluorous-phase strategy for improving separation efficiency in organic synthesis, *Science* **275** 823-826 1997.

Tetrahedron Symposium **58** Issue 20, pp 3823-4131 2002.

CHAPTER 4

PURIFICATION OF ORGANIC CHEMICALS

The general principles, techniques and methods of purification in Chapters 1 and 2 are applicable to this chapter. Most organic liquids and a number of solids can readily be purified by fractional distillation, usually at atmospheric pressure. Sometimes, particularly with high boiling or sensitive liquids, or when in doubt about stability, distillation or fractionation under reduced pressure should be carried out. To save space, the present chapter omits many substances for which the published purification methods involve simple distillation. Where boiling points are given, purification by distillation is another means of removing impurities. Literature references are omitted for methods which require simple recrystallisation from solution if the correct solvent can be guessed readily, and where no further information is given, e.g. spectra. Substances are listed alphabetically, usually with some criteria of purity, giving brief details of how they can be purified. Also noted are the molecular weights (to the first decimal place), melting points and/or boiling points together with the respective densities and refractive indexes for liquids, and optical rotations when the compounds are chiral. When the temperatures and/or the wavelengths are not given for the last three named properties then they should be assumed to be 20°C and the average of the wavelengths of the sodium D lines respectively; and densities are relative to water at 4°.

The present chapter includes commercially available organic chemicals. Most of the inorganic, metalorganic, organo- bismuth, boron, phosphorus, selenium, silicon and alkali metal compounds and metal ion salts of organic acids are in Chapter 5. Naturally occurring commercially available organic compounds for use in biochemistry, molecular biology and biology are in Chapter 6. Commercially available polymer supported reagents are indicated with § under the appropriate reagent.

Rapid purification procedures are noted for commonly used solvents and reagents which make them suitable for general use in synthetic chemistry.

Abbreviations of titles of periodicals are defined as in the Chemical Abstracts Service Source Index (CASSI). Other abbreviations are self evident (see Chapter 1, p. 30).

Ionisation constants of ionisable compounds are given as **pK** values (published from the literature) and refer to the **pKa** values at room temperature (~ 15°C to 25°C). The values at other temperatures are given as superscripts, e.g. **pK²⁵** for 25°C. Estimated values are entered as **pK_{Est.}** (see Chapter 1, p. 7 for further information).

As a good general rule, all low boiling (<100°) organic liquids should be treated as highly flammable and toxic (because they can be inhaled in large quantities) and the necessary precautions should be taken.

Benzene, which has been used as a solvent successfully and extensively in the past for reactions and purification by chromatography and crystallisation is now considered a **very dangerous substance** so it has to be used with extreme care. We emphasise that an alternative solvent system to benzene (e.g. toluene, toluene-petroleum ether, or a petroleum ether to name a few) should be used first. However, if no other solvent system can be found then all operations involving benzene have to be performed in an efficient fumehood and precautions must be taken to avoid inhalation and contact with skin and eyes. Whenever benzene is mentioned in the text an asterisk e.g. *C₆H₆ or *benzene, is inserted to remind the user that special precaution should be adopted.

Abietic acid [514-10-3] M 302.5, m 172-175°, $[\alpha]_D^{25}$ -116° (-106°)(c 1, EtOH), pK 5.27. Crystd by dissolving 100g of acid in 95% EtOH (700mL), adding to H₂O (600mL) and cooling. Filter, dry in a vacuum (over KOH or CaSO₄) store in an O₂-free atmosphere. λ in EtOH nm(log ϵ): 2343(4.3), 241(4.4), 2505(4.2), 235(4.34) and 240(4.36). [*Org Synth* 23 1 1952 ; *J Am Chem Soc* 35 3736 1949; *Monatsh Chem* 116 1345 1985.]

S-Abscisic acid [21293-29-8] M 264.3, m 160-161°, 161-163° (sublimation), $[\alpha]_{287} + 24,000^\circ$, $[\alpha]_{245} -69,000^\circ$ (c 1-50 μ g/mL in acidified MeOH or EtOH), pK_{Est} ~3.9. Crystd from CCl₄-pet.ether, EtOH + hexane and sublimes at 120°.

Acenaphthalene [208-96-8] M 152.2, m 92-93°. Dissolved in warm redistd MeOH, filtered through a sintered glass funnel and cooled to -78° to ppt the material as yellow plates [Dainton, Ivin and Walmsley *Trans Faraday Soc* 56 1784 1960]. Alternatively can be sublimed *in vacuo*.

Acenaphthaquinone [82-86-0] M 182.2, m 260-261°. Extracted with, then recrystd twice from *C₆H₆. [LeFevre, Sundaram and Sundaram *J Chem Soc* 974 1963].

Acenaphthene [83-32-9] M 154.2, m 94.0°. Crystd from EtOH. Purified by chromatography from CCl₄ on alumina with *benzene as eluent [McLaughlin and Zainal *J Chem Soc* 2485 1960].

RS-Acenaphthenol [6306-07-6] M 170.2, m 144.5-145.5°, 146°, 148°. If highly coloured (yellow), dissolve in boiling *benzene (14g in 200mL), add charcoal (0.5g), filter through a heated funnel, concentrate to 100mL and cool to give almost colourless needles. *Benzene vapour is TOXIC use an efficient fume cupboard. The acetate has b 166-168°/5mm (bath temp 180-185°). [*Org Synth* Col.Vol. III 3 1955.] It can also be recrystd from *C₆H₆ or EtOH [Fieser and Cason *J Am Chem Soc* 62 432 1940]. It forms a brick-red crystalline complex with 2,4,5,7-tetranitrofluoren-9-one which is recrystd from AcOH and dried in a vacuum over KOH and P₂O₅ at room temp, m 170-172° [Newman and Lutz *J Am Chem Soc* 78 2469 1956].

Acetal (acetaldehyde diethylacetal) [105-57-7] M 118.2, b 103.7-104°, d 0.831, n 1.38054, n²⁵ 1.3682. Dried over Na to remove alcohols and water, and to polymerise aldehydes, then fractionally distd. Or, treat with alkaline H₂O₂ soln at 40-45° to remove aldehydes, then the soln is saturated with NaCl, separated, dried with K₂CO₃ and distd from Na [Vogel *J Chem Soc* 616 1948].

Acetaldehyde [75-07-0] M 44.1, b 20.2°, d 0.788, n 1.33113, pK²⁵ 13.57 (hydrate). Usually purified by fractional distn in a glass helices-packed column under dry N₂, discarding the first portion of distillate. Or, shaken for 30min with NaHCO₃, dried with CaSO₄ and fractionally distd at 760mm through a 70cm Vigreux column. The middle fraction was taken and further purified by standing for 2h at 0° with a small amount of hydroquinone, followed by distn [Longfield and Walters *J Am Chem Soc* 77 810 1955].

Acetaldehyde ammonia trimer (hexahydro-2,4,6-trimethyl-1,3,5-triazine trihydrate) [76231-37-3] M 183.3, m 94-96°, 95-97°, 97°, b 110°(partly dec). Crystd from EtOH-Et₂O. When prepared it separates as the trihydrate which can be dried in a vacuum over CaCl₂ at room temp to give the anhydrous compound with the same melting point. The dihydrate melts at 25-28° then resolidifies and melts again at 94-95°. IRRITATES THE EYES AND MUCOUS MEMBRANES. [*J Org Chem* 38 3288 1973.]

Acetaldehyde dimethyl acetal [534-15-6] M 90.1, b 63-65°, d₄²⁰ 0.852, n_D²⁵ 1.36678. Distd through a fractionating column and fraction boiling at 63.8°/751mm is collected. It forms an azeotrope with MeOH. It has been purified by GLC.

Acetamide [60-35-5] M 59.1, m 81°, pK₁²⁵ -1.4, pK₂²⁵ +0.37. Crystd by soln in hot MeOH (0.8mL/g), dilted with Et₂O and allowed to stand [Wagner *J Chem Educ* 7 1135 1930]. Alternate crystns are from acetone, *benzene, chloroform, dioxane, methyl acetate or from *benzene-ethyl acetate mixture (3:1 and 1:1). It has also been recrystd from hot water after treating with HCl-washed activated charcoal (which had been

repeatedly washed with water until free from chloride ions), then crystd again from hot 50% aq. EtOH and finally twice from hot 95% EtOH [Christoffers and Kegeles *J Am Chem Soc* **85** 2562 1963]. Final drying is in a vacuum desiccator over P₂O₅. Acetamide is also purified by distn (b 221-223°) or by sublimation *in vacuo*. Also purified by recrystn twice from cyclohexane containing 5% (v/v) of *benzene. Needle-like crystals separated by filt'n, washed with a small volume of distd H₂O and dried with a flow of dry N₂. [Slebocka-Tilk et al. *J Am Chem Soc* **109** 4620 1987.]

Acetamidine hydrochloride [124-42-5] M 94.5, m 164-166°, 165-170° (dec), 174°, pK²⁵ 12.40. It can be recrystd from small volumes of EtOH. Alternatively dissolve in EtOH, filter, add Et₂O, filter the crystalline salt off under N₂, dry in a vacuum desiccator over H₂SO₄. The salt is deliquescent and should be stored in a tightly stoppered container. Solubility in H₂O is 10% at room temperature, soluble in Me₂CO. The *free base* reacts strongly alkaline in H₂O. It has λ_{max} 224nm (ε 4000) in H₂O. The *picrate* has m 252° (sintering at ~245°). [Dox *Org Synth Coll Vol I* 5 1941; Davies and Parsons *Chem Ind (London)* 628 1958; Barnes et al. *J Am Chem Soc* **62** 1286 1940 give m 177-178°.]

N-(2-Acetamido)-2-aminoethanesulfonic acid (ACES) [7365-82-4] M 182.2, m > 220°(dec), pK_{Est} ~1.5, pK₂ 6.9. Recrystd from hot aqueous EtOH.

4-Acetamidobenzaldehyde [122-85-0] M 163.2, m 156°. Recrystd from water.

p-Acetamidobenzenesulfonyl chloride (N-acetylsulfanilyl chloride) [121-60-8] M 233.7, m 149°(dec). Crystd from toluene, CHCl₃, or ethylene dichloride.

α-Acetamidocinnamic acid [5469-45-4] M 205.2, m 185-186° (2H₂O), 190-191°(anhydr), 193-195°, pK_{Est} ~3.2. Recrystd from H₂O as the dihydrate and on drying at 100° it forms the anhydrous compound which is *hygroscopic*. Alkaline hydrolysis yields NH₃ and phenylpyruvic acid. [Erlenmeyer and Früstück *Justus Liebigs Ann Chem* **284** 47 1895.]

Z-O-(2-Acetamido-2-deoxy-D-glycopyranosylideneamino)N-phenylcarbamate (PUGNAC) [132063-05-9] M 335.3, m 171-174°(dec), 174-180°(dec), [α]_D²⁰ +67.5° (c 0.2, MeOH). Purified by flash chromatography (silica gel and eluted with AcOEt-hexane 3:2) evaporated, and the foam recrystallised from AcOEt-MeOH. TLC on Merck SiO₂ gel 60 F₂₅₄ and detected by spraying with 0.025M I₂ in 10% aqueous H₂SO₄ and heat at 200° gave R_F 0.21. The acetate is hydrolysed with NH₃-MeOH. [*Helv Chim Acta* **68** 2254 1985; **73** 1918 1990.]

2-Acetamidofluorene [53-96-3] M 223.3, m 194°, 196-198°. Recrystd from toluene (1.3mg in 100mL). Solubility in H₂O is 1.3mg/L; UV λ_{max} nm(log ε) : 288(4.43), 313(4.13). [*J Org Chem* **21** 271 1956.] It can also be recrystd from 50% AcOH and sol in H₂O is 1.3mg/100mL at 25° [*Chem Ber* **35** 3285 1902]. 9-¹⁴C and ω-¹⁴C 2-acetamidofluorene were recrystd from aqueous EtOH and had m 194-195° and 194° respectively. **Potent CARCINOGEN**. [*Cancer Res* **10** 616 1950; Sadin et al. *J Am Chem Soc* **74** 5073 1952.]

N-(2-Acetamido)iminodiacetic acid (ADA) [26239-55-4] M 190.2, m 219° (dec), pK₁ ~2.3, pK₂ 6.6. Dissolved in water by adding one equivalent of NaOH soln (to final pH of 8-9), then acidified with HCl to ppt the free acid. Filtered and washed with water.

Acetamidomethanol [625-51-4] M 89.1, m 47-50°, 54-56°, 55°. Recryst from freshly distd Me₂CO, wash the crystals with dry Et₂O and dry in a vacuum desiccator over P₂O₅. R_F 0.4 on paper chromatography with CHCl₃/EtOH (2:8) as solvent and developed with ammoniacal AgNO₃. Also crystallises in needles from EtOAc containing a few drops of Me₂CO. It is *hygroscopic* and should be stored under dry conditions. [*J Am Chem Soc* **73** 2775 1951; *Chem Ber* **99** 3204 1966; *Justus Liebigs Ann Chem* **343** 265 1905.]

2-Acetamido-5-nitrothiazole [140-40-9] M 187.2, m 264-265°. Recrystd from EtOH or glacial acetic acid.

2-Acetamidophenol [614-80-2] M 151.2, m. 209°, pK_{Est} ~9.4. Recrystd from water or aqueous EtOH.

3-Acetamidophenol [621-42-1] M 151.2, m 148-149°, pK²⁵ ~9.59. Recrystd from water.

4-Acetamidophenol [103-90-2] M 151.2, m 169-170.5°, pK_{Est} ~10.0. Recrystd from water or EtOH.

4-Acetamido-2,2,6,6-tetramethylpiperidine-1-oxyl (acetamidoTEMPO) [14691-89-5] M 213.3, m 144-146°, 146-147°. Dissolve in CH₂Cl₂, wash with saturated K₂CO₃, then saturated aqueous NaCl, dry (Na₂SO₄), filter and evaporate. The red solid is recrystd from aqueous MeOH, m 147.5°. [*J Org Chem* 56 6110 1991; *Bull Acad Sci USSR, Div Chem Sci* 15 1422 1966.]

5-Acetamido-1,3,4-thiadiazole-2-sulfonamide [59-66-5] M 222.3, m 256-259° (dec). Recrystd from water.

Acetanilide [103-84-4] M 135.2, m 114°, pK²⁵ 0.5. Recrystd from water, aqueous EtOH, *benzene or toluene.

Acetic acid (glacial) [64-19-7] M 60.1, m 16.6°, b 118°, d 1.049, n 1.37171, n²⁵ 1.36995, pK²⁵ 4.76. Usual impurities are traces of acetaldehyde and other oxidisable substances and water. (Glacial acetic acid is very *hygroscopic*. The presence of 0.1% water lowers its m by 0.2°.) Purified by adding some acetic anhydride to react with water present, heating for 1h to just below boiling in the presence of 2g CrO₃ per 100mL and then fractionally distilling [Orton and Bradfield *J Chem Soc* 960 1924, 983 1927]. Instead of CrO₃, 2-5% (w/w) of KMnO₄, with boiling under reflux for 2-6h, has been used.

Traces of water have been removed by refluxing with tetraacetyl diborate (prepared by warming 1 part of boric acid with 5 parts (w/w) of acetic anhydride at 60°, cooling, and filtering off), followed by distn [Eichelberger and La Mer *J Am Chem Soc* 55 3633 1933].

Refluxing with acetic anhydride in the presence of 0.2g % of 2-naphthalenesulfonic acid as catalyst has also been used [Orton and Bradfield *J Chem Soc* 983 1927]. Other suitable drying agents include CuSO₄ and chromium triacetate: P₂O₅ converts some acetic acid to the anhydride. Azeotropic removal of water by distn with thiophene-free *benzene or with butyl acetate has been used [Birdwhistell and Griswold *J Am Chem Soc* 77 873 1955]. An alternative purification uses fractional freezing.

Rapid procedure: Add 5% acetic anhydride, and 2% of CrO₃. Reflux and fractionally distil.

Acetic anhydride [108-24-7] M 102.1, b 138°, d 1.082, n 1.3904. Adequate purification can usually be obtained by fractional distn through an efficient column. Acetic acid can be removed by prior refluxing with CaC₂ or with coarse Mg filings at 80-90° for 5days, or by distn from a large excess of quinoline (1% AcOH in quinoline) at 75mm pressure. Acetic anhydride can also be dried by standing with Na wire for up to a week, removing the Na and distilling from it under vacuum. (Na reacts vigorously with acetic anhydride at 65-70°). Dippy and Evans [*J Org Chem* 15 451 1950] let the anhydride (500g) stand over P₂O₅ (50g) for 3h, then decanted it and stood it with ignited K₂CO₃ for a further 3h. The supernatant liquid was distd and the fraction b 136-138°, was further dried with P₂O₅ for 12h, followed by shaking with ignited K₂CO₃, before two further distns through a five-section Young and Thomas fractionating column. The final material distd at 137.8-138.0°. Can also be purified by azeotropic distn with toluene: the azeotrope boils at 100.6°. After removal of the remaining toluene, the anhydride is distd [sample had a specific conductivity of 5 x 10⁻⁹ ohm⁻¹cm⁻¹].

Rapid procedure: Shake with P₂O₅, separate, shake with dry K₂CO₃ and fractionally distil.

Acetic hydrazide [1068-57-1] M 74.1, m 67°, b 127°/18mm. Cryst as needles from EtOH. Reduces NH₃/AgNO₃.

Acetoacetamide [5977-14-0] M 101.1, m 54-55°, 54-56°. Recrystallise from CHCl₃, or Me₂CO/pet ether. Crystallises from pyridine with 4mol of solvent. Slightly soluble in H₂O, EtOH and AcOH but

insoluble in Et₂O. *Phenylhydrazone* has *m* 128°. [*Beilstein* 3, 4th Suppl, p 1545; *Kato Chem Pharm Bull Jpn* 15 921,923 1967; *Chem Ber* 35 583 1902.]

Acetoacetanilide [102-01-2] *M* 177.2, *m* 86°, *pK* 10.68. Crystd from H₂O, aqueous EtOH or pet ether (b 60-80°).

Acetoacetylpiiperidide [1128-87-6] *M* 169.2, *b* 88.9°/0.1mm, *n*⁵² 1.4983. Dissolved in *benzene, extracted with 0.5M HCl to remove basic impurities, washed with water, dried, and distd at 0.1mm [*Wilson J Org Chem* 28 314 1963].

α-Acetobromoglucose (2,3,4,6-tetraacetyl-α-D-glucopyranosyl bromide) [572-09-8] *M* 411.2, *m* 88-89°, [α]_D²⁵ +199.3° (c 3, CHCl₃). Crystd from isopropyl ether or pet ether (b 40-60°) [*Org Synth* 65 236 1897].

Acetone [67-64-1] *M* 58.1, *b* 56.2°, *d* 0.791, *n* 1.35880, *pK*₁²⁵ -6.1 (**basic, mono-protonated**), *pK*₂²⁵ 20.0 (**acidic**) The commercial preparation of acetone by catalytic dehydrogenation of isopropyl alcohol gives relatively pure material. Analytical reagent quality generally contains less than 1% organic impurities but may have up to about 1% H₂O. Dry acetone is appreciably *hygroscopic*. The main organic impurity in acetone is mesityl oxide, formed by the aldol condensation. It can be dried with anhydrous CaSO₄, K₂CO₃ or type 4A Linde molecular sieves, and then distd. Silica gel and alumina, or mildly acidic or basic desiccants cause acetone to undergo the aldol condensation, so that its water content is increased by passage through these reagents. This also occurs to some extent when P₂O₅ or sodium amalgam is used. Anhydrous MgSO₄ is an inefficient drying agent, and CaCl₂ forms an addition compound. Drierite (anhydrous CaSO₄) offers the minimum acid and base catalysis of aldol formation and is the recommended drying agent for this solvent [Coetzee and Siao *Inorg Chem* 14v 2 1987; Riddick and Bunger *Organic Solvents* Wiley-Interscience, N.Y., 3rd edn, 1970]. Acetone was shaken with Drierite (25g/L) for several hours before it was decanted and distd from fresh Drierite (10g/L) through an efficient column, maintaining atmospheric contact through a Drierite drying tube. The equilibrium water content is about 10⁻²M. Anhydrous Mg(ClO₄)₂ **should not be used as drying agent because of the risk of EXPLOSION with acetone vapour.**

Organic impurities have been removed from acetone by adding 4g of AgNO₃ in 30mL of water to 1L of acetone, followed by 10mL of M NaOH, shaking for 10min, filtering, drying with anhydrous CaSO₄ and distilling [Werner *Analyst (London)* 58 335 1933]. Alternatively, successive small portions of KMnO₄ have been added to acetone at reflux, until the violet colour persists, followed by drying and distn. Refluxing with chromium trioxide (CrO₃) has also been used. Methanol has been removed from acetone by azeotropic distn (at 35°) with methyl bromide, and treatment with acetyl chloride.

Small amounts of acetone can be purified as the NaI addition compound, by dissolving 100g of finely powdered NaI in 400g of boiling acetone, then cooling in ice and salt to -8°. Crystals of NaI.3Me₂CO are filtered off and, on warming in a flask, acetone distils off readily. [This method is more convenient than the one using the bisulfite addition compound.] Also purified by gas chromatography on a 20% free fatty acid phthalate (on Chromosorb P) column at 100°.

For efficiency of desiccants in drying acetone see Burfield and Smithers [*J Org Chem* 43 3966 1978]. The water content of acetone can be determined by a modified Karl Fischer titration [Koupparis and Malmstadt *Anal Chem* 54 1914 1982].

Rapid procedure: Dry over anhydrous CaSO₄ and distil.

Acetone cyanohydrin [75-86-5] *M* 85.1, *b* 48°/2.5mm, 68-70°/11mm, 78-82°/15mm, *d*₄²⁰ 0.93. Dry with Na₂SO₄, and distil as rapidly as possible under vacuum to avoid decomposition. Discard fractions boiling below 78-82°/15mm. Store in the dark. **USE AN EFFICIENT FUME HOOD as HCN (POISONOUS) is always present.** [*Org Synth* Col.Vol. II 7 1940.]

Acetonedicarboxylic acid [542-05-2] *M* 146.1, *m* 138° (dec), *pK*²⁵ 3.10. Crystd from ethyl acetate and stored over P₂O₅. Decarboxylates in hot water.

Acetone semicarbazone [110-20-3] *M* 115.1, *m* 187°, *pK*²⁵ 1.33. Crystd from water or from aqueous EtOH.

Acetonitrile (methyl cyanide) [75-05-8] **M 41.1, b 81.6°, d²⁵ 0.77683, n 1.3441, n²⁵ 1.34163.** Commercial acetonitrile is a byproduct of the reaction of propylene and ammonia to acrylonitrile. The following procedure that significantly reduces the levels of acrylonitrile, allyl alcohol, acetone and benzene was used by Kiesel [*Anal Chem* **52** 2230 1988]. Methanol (300mL) is added to 3L of acetonitrile fractionated at high reflux ratio until the boiling temperature rises from 64° to 80°, and the distillate becomes optically clear down to $\lambda = 240\text{nm}$. Add sodium hydride (1g) free from paraffin, to the liquid, reflux for 10min, and then distil rapidly until about 100mL of residue remains. Immediately pass the distillate through a column of acidic alumina, discarding the first 150mL of percolate. Add 5g of CaH₂ and distil the first 50mL at a high reflux ratio. Discard this fraction, and collect the following main fraction. The best way of detecting impurities is by gas chromatography.

Usual contaminants in commercial acetonitrile include H₂O, acetamide, NH₄OAc and NH₃. Anhydrous CaSO₄ and CaCl₂ are inefficient drying agents. Preliminary treatment of acetonitrile with cold, satd aq KOH is undesirable because of base-catalysed hydrolysis and the introduction of water. Drying by shaking with silica gel or Linde 4A molecular sieves removes most of the water in acetonitrile. Subsequent stirring with CaH₂ until no further hydrogen is evolved leaves only traces of water and removes acetic acid. The acetonitrile is then fractionally distd at high reflux, taking precaution to exclude moisture by refluxing over CaH₂ [Coetzee *Pure Appl Chem* **13** 429 1966]. Alternatively, 0.5-1% (w/v) P₂O₅ is often added to the distilling flask to remove most of the remaining water. Excess P₂O₅ should be avoided because it leads to the formation of an orange polymer. Traces of P₂O₅ can be removed by distilling from anhydrous K₂CO₃.

Kolthoff, Bruckenstein and Chantooni [*J Am Chem Soc* **83** 3297 1961] removed acetic acid from 3L of acetonitrile by shaking for 24h with 200g of freshly activated alumina (which had been reactivated by heating at 250° for 4h). The decanted solvent was again shaken with activated alumina, followed by five batches of 100-150g of anhydrous CaCl₂. (Water content of the solvent was then less than 0.2%). It was shaken for 1h with 10g of P₂O₅, twice, and distd in a 1m x 2cm column, packed with stainless steel wool and protected from atmospheric moisture by CaCl₂ tubes. The middle fraction had a water content of 0.7 to 2mM.

Traces of unsaturated nitriles can be removed by an initial refluxing with a small amount of aq KOH (1mL of 1% solution per L). Acetonitrile can be dried by azeotropic distn with dichloromethane, benzene or trichloroethylene. Isonitrile impurities can be removed by treatment with conc HCl until the odour of isonitrile has gone, followed by drying with K₂CO₃ and distn.

Acetonitrile was refluxed with, and distd from alkaline KMnO₄ and KHSO₄, followed by fractional distn from CaH₂. (This was better than fractionation from molecular sieves or passage through a type H activated alumina column, or refluxing with KBH₄ for 24h and fractional distn)[Bell, Rodgers and Burrows *J Chem Soc, Faraday Trans 1* **73** 315 1977; Moore et al. *J Am Chem Soc* **108** 2257 1986].

Material suitable for polarography was obtained by refluxing over anhydrous AlCl₃ (15g/L) for 1h, distilling, refluxing over Li₂CO₃ (10g/L) for 1h and redistg. It was then refluxed over CaH₂ (2g/L) for 1h and fractionally distd, retaining the middle portion. The product was not suitable for UV spectroscopy use. A better purification procedure used refluxing over anhydrous AlCl₃ (15g/L) for 1h, distg, refluxing over alkaline KMnO₄ (10g KMnO₄, 10g Li₂CO₃/L) for 15min, and distg. A further reflux for 1h over KHSO₄ (15g/L), then distn, was followed by refluxing over CaH₂ (2g/L) for 1h, and fractional distn. The product was protected from atmospheric moisture and stored under nitrogen [Walter and Ramalay *Anal Chem* **45** 165 1973]. Purification of "General Purity Reagent" for this purpose is not usually satisfactory because very large losses occur at the KMnO₄, LiCO₃ step. For electrochemical work involving high oxidation fluorides, further reflux over P₂O₅ (1g/mL for 0.5h) and distilling (discarding 3% of first and last fractions) and repeating this step is necessary. The distillate is kept over molecular sieves in vac after degassing, for 24h and vac distd onto freshly activated 3A molecular sieves. The MeCN should have absorption at 200nm of <0.05 (H₂O reference) and UV cutoff at ca 175nm. Also the working potential range of purified Et₄N⁺ BF₄⁻ (0.1mol.dcm⁻³ in the MeCN) should be +3.0 to -2.7V vs Ag⁺/Ag⁰. If these criteria are not realised then further impurities can be removed by treatment with activated neutral alumina (60 mesh) in vacuo before final molecular sieves treatment [Winfield *J Fluorine Chem* **25** 91 1984].

Acetonitrile has been distd from AgNO₃, collecting the middle fraction over freshly activated Al₂O₃. After standing for two days, the liquid was distd from the activated Al₂O₃. Specific conductivity 0.8-1.0 x 10⁻⁸ mhos [Harkness and Daggett *Can J Chem* **43** 1215 1965]. Acetonitrile ¹⁴C was purified by gas chromatography and is water free and distd at 81°. [*J Mol Biol* **87** 541 1974.]

Rapid procedure: Dry over anhydrous K_2CO_3 for 24h, followed by further drying for 24h over 3A molecular sieves or boric anhydride, followed by distn. Alternatively, stir over P_2O_5 (5% w/v) for 24h then distil. However this last method is not suitable for use in reactions with very acid sensitive compounds.

Acetylacetone (2,5-hexanedione) [110-13-4] M 114.2, m -9° , b 76-78°/13 mm, 88°/25mm, 137°/150mm, 188°/atm, d_4^{20} 0.9440, n_D^{20} 1.423, pK 18.7. Purified by dissolving in Et_2O , stirred with K_2CO_3 (a quarter of the wt of dione), filtered, dried over anhydrous Na_2SO_4 (not $CaCl_2$), filtered, evapd and distd in a vacuum. It is then redistd through a 30cm Vigreux column (oil bath temp 150°). It is miscible with H_2O and $EtOH$. The *dioxime* has m 137° (plates from $*C_6H_6$), *mono-oxime* has b 130°/11mm, and the *2,4-dinitrophenylhydrazone* has m 210-212° (red needles from $EtOH$). [Chem Ber 22 2100 1989; for enol content see J Org Chem 19 1960 1954.]

4-Acetophenetidine (phenacetin) [62-44-2] M 179.2, m 136°. Crystd from H_2O or purified by soln in cold dilute alkali and reprecipitated by addn of acid to neutralisation point. Air-dried.

Acetophenone [98-86-2] M 120.2, m 19.6°, b 54°/2.5mm, 202°/760mm, d^{25} 1.0238, n^{25} 1.5322, pK 19.2. Dried by fractional distn or by standing with anhydrous $CaSO_4$ or $CaCl_2$ for several days, followed by fractional distn under reduced pressure (from P_2O_5 , optional), and careful, slow and repeated partial crystallisations from the liquid at 0° excluding light and moisture. It can also be crystallised at low temperatures from isopentane. Distn can be followed by purification using gas-liquid chromatography [Earls and Jones J Chem Soc, Faraday Trans 1 71 2186 1975.]

§ A commercial polystyrene supported version is available — scavenger resin (for diol substrate).

Aceto-*o*-toluidide [120-66-1] M 149.2, m 110°, b 296°/760mm. Crystd from H_2O , $EtOH$ or aqueous $EtOH$.

Aceto-*m*-toluidide [537-92-8] M 149.2, m 65.5°, b 182-183°/14mm, 303°/760mm. Crystd from H_2O , $EtOH$ or aqueous $EtOH$.

Aceto-*p*-toluidide [103-89-9] M 149.2, m 146°, b 307°/760mm. Crystd from aqueous $EtOH$.

Acetoxime (acetone oxime) [127-06-0] M 73.1, m 63°, b 135°/760mm, pK⁴⁰ 0.99. Crystd from pet ether (b 40-60°). Can be sublimed.

Acetoxyacetone (acetol acetone) [592-20-1] M 116.1, b 65°/11mm, 73-75°/17mm, 174-176°/atm, d_4^{20} 1.0757, n_D^{20} 1.4141. Distil under reduced pressure, then redistil at atm pressure. It is miscible with H_2O but is slowly decomposed by it. Store in dry atmosphere. The *2,4-dinitrophenylhydrazone* has m 115-115.5° (from $CHCl_3$ /hexane). [J Chem Soc 59 789 1891; J Org Chem 21 68 1956; Justus Liebigs Ann Chem 335 260 1904.]

4-Acetoxy-2-azetidinone [28562-53-0] M 129.1, m 38-41°. Dissolve in $CHCl_3$, dry ($MgSO_4$) concentrate at 40°/70mm, or better at room temperature to avoid decomposition. Wash and stir the residual oil with hexane by decantation and discard wash. Dry the oil at high vacuum when it should solidify, m 34°. It can be distilled at high vacuum, 80-82°/10⁻³mm, but this results in extensive losses. The purity can be checked by TLC using Merck Silica Gel F₂₅₄ and eluting with $EtOAc$. The azetidinone has R_F 0.38 (typical impurities have R_F 0.67). The spots can be detected by the TDM spray. This is prepared from (A) 2.5g 4,4'-tetramethyldiaminodiphenylmethane (TDM) in 10mL $AcOH$ and diluted with 50mL of H_2O , (B) 5g KI in 100mL of H_2O and (C) 0.3g ninhydrin in 10mL of $AcOH$ and 90mL of H_2O . The spray is prepared by mixing (A) and (B) with 1.5mL of (C) and stored in a brown bottle. [Justus Liebigs Ann Chem 539 1974; Org Synth 65 135 1987.]

1-Acetoxy-1,3-butadiene (1,3-butadienyl acetate) *cis-trans* mixture [1515-76-0] M 112.1, b 42-43°/16mm, 51-52°/20mm, 60-61°/40mm, d_4^{20} 0.9466, n_D^{20} 1.4622. The commercial sample is stabilised with 0.1% of *p-tert*-butylcatechol. If the material contains crotonaldehyde (by IR, used in its synthesis) it should be dissolved in Et_2O , shaken with 40% aqueous sodium bisulfite, then 5% aqueous

Na_2CO_3 , water, dried (Na_2SO_4) and distilled several times in a vac through a Widmer (*Helv Chim Acta* 7 59 1924) or Vigreux column [Wicterle and Hudlicky *Collect Czech Chem Commun* 12 564 1947; Hagemeyer and Hull *Ind Eng Chem* 41 2920 1949].

1-Acetoxy-2-butoxyethane [112-07-2] M 160.2, b 61-62°/0.2mm, 75-76°/12mm, 185.5°/740mm, 188-192°/atm, d_4^{20} 0.9425, n_D^{20} 1.4121. Shake with anhydrous Na_2CO_3 , filter and distil in a vacuum. Redistn can be then be carried out at atmospheric pressure. [*J Org Chem* 21 1041 1956.]

3R,4R,1'R-4-Acetoxy-3-[1-(tert-butylmethylsilyloxy)ethyl]-2-azetinone see Chapter 5.

2-Acetoxyethanol [542-59-6] M 104.1, b 187°/761mm, 187-189°/atm, d_4^{20} 1.108, n_D^{20} 1.42. Dry over K_2CO_3 (not CaCl_2), and distil. [*J Chem Soc* 3061 1950; rate of hydrolysis: *J Chem Soc* 2706 1951.]

1-Acetoxy-2-ethoxyethane [111-15-9] M 132.2, b 156-159°, d_4^{20} 0.97, n_D^{20} 1.406. Shake with anhydr Na_2CO_3 , filter and distil in vac. Redistn can then be carried out at atm pressure. [*J Org Chem* 21 1041 1956.]

1-Acetoxy-2-methoxyethane [110-49-6] M 118.1, b 141°/732mm, 140144°/atm, d_4^{20} 1.009, n_D^{20} 1.4011. Shake with anhydrous Na_2CO_3 , filter and distil in a vacuum. Redistn can be then be carried out at atmospheric pressure. [*J Org Chem* 21 1041 1956.]

R(-)- α -Acetoxyphenylacetic (acetyl mandelic) acid [51019-43-3] M 194.2, m 96-98°, $[\alpha]_D^{20}$ -153.7° (c 2.06, Me_2CO), $[\alpha]_{546}^{20}$ -194° (c 2.4, Me_2CO), $\text{pK}_{\text{Est}} \sim 2.9$ Recryst from H_2O with 1mol of solvent which is removed on drying, or from solvents as for the S-isomer. [*J Chem Soc* 227 1943.]

S-(+)- α -Acetoxyphenylacetic (acetyl mandelic) acid [7322-88-5] M 194.2, m 80-81°, 95-97.5°, $[\alpha]_D^{27}$ +158° (c 1.78, Me_2CO), $[\alpha]_{546}^{20}$ +186° (c 2, Me_2CO). Recryst from *benzene-hexane or toluene and has characteristic NMR and IR spectra. [*Justus Liebigs Ann Chem* 622 10 1959; *J Org Chem* 39 1311 1974.]

21-Acetoxypregnenolone [566-78-9] M 374.5, m 184-185°. Crystd from Me_2CO .

S(-)-2-Acetoxypropionyl chloride [36394-75-9] M 150.6, b 51-53°/11mm, d_4^{20} 1.19, n_D^{20} 1.423, $[\alpha]_D^{27}$ -33°, (c 4, CHCl_3), $[\alpha]_{546}^{20}$ -38° (c 4, CHCl_3). It is moisture sensitive and is hydrolysed to the corresponding acid. Check the IR spectrum. If the OH band above 3000cm^{-1} is too large and broad then the mixture should be refluxed with pure acetyl chloride for 1h, evapd and distd under reduced pressure.

S-Acetoxy succinic anhydride [59025-03-5] M 158.1, m 58° (RS 81.5-82.5°, 86-87°), $[\alpha]_D^{20}$ -26.0° (c 19, Me_2CO), $[\alpha]_D^{20}$ -28.4° (c 13, Ac_2O). Recrystd from Ac_2O and dry in a vacuum over KOH, or by washing with dry Et_2O due to its deliquescent nature. [*J Chem Soc* 788 1933; *Synth Commun* 16 183 1986; *J Org Chem* 52 1040 1988; RS : *J Am Chem Soc* 88 5306 1966.]

Acetylacetone (2,4-pentanedione) [123-54-6] M 100.1, b 45°/30mm, $d^{30.2}$ 0.9630, $n^{18.5}$ 1.45178, pK_1^{25} -5.0 (enol), -6.6 (keto), pK_2^{25} 8.95 Small amounts of acetic acid were removed by shaking with small portions of 2M NaOH until the aqueous phase remained faintly alkaline. The sample, after washing with water, was dried with anhydrous Na_2SO_4 , and distd through a modified Vigreux column [*Cartledge J Am Chem Soc* 73 4416 1951]. An additional purification step is fractional crystn from the liquid. Alternatively, there is less loss of acetylacetone if it is dissolved in four volumes of *benzene and the soln is shaken three times with an equal volume of distd water (to extract acetic acid): the *benzene is then removed by distn at 43-53° and 20-30mm through a helices-packed column. It is then refluxed over P_2O_5 (10g/L) and fractionally distd under reduced pressure. The distillate (sp conductivity $4 \times 10^{-8} \text{ohm}^{-1}\text{cm}^{-1}$) was suitable for polarography [Fujinaga and Lee *Talanta* 24 395 1977]. To recover used acetylacetone, metal ions were stripped from the soln at pH 1 (using 100mL 0.1M H_2SO_4 /L of acetylacetone). The acetylacetone was washed with

(1:10) ammonia soln (100mL/L) and with distd water (100mL/L, twice), then treated as above. It complexes with Al, Be, Ca, Cd, Ce, Cu, Fe²⁺, Fe³⁺, Mn, Mg, Ni, Pb and Zn.

***N*-Acetyl-L-alaninamide** [15962-47-7] M 130.2, m 162°. Crystd repeatedly from EtOH-diethyl ether.

***N*-Acetyl-β-alanine** [3025-95-4] M 127.2, m 78.3-80.3°, pK²⁵ 4.45. Crystd from acetone.

***N*-Acetyl-L-alanyl-L-alaninamide** [30802-37-0] M 201.2, m 250-251°. Crystd repeatedly from EtOH/diethyl ether.

***N*-Acetyl-L-alanyl-L-alanyl-L-alaninamide** [29428-34-0] M 272.3, m 295-300°. Crystd from MeOH/diethyl ether.

***N*-Acetyl-L-alanylglycinamide** [76571-64-7] M 187.2, m 148-149°. Crystd repeatedly from EtOH/diethyl ether.

Acetyl-α-amino-*n*-butyric acid [34271-24-4] M 145.2, pK²⁵ 3.72. Crystd twice from water (charcoal) and air dried [King and King *J Am Chem Soc* 78 1089 1956].

9-Acetylanthracene [784-04-3] M 220.3, m 75-76°. Crystd from EtOH. [Masnori et al. *J Am Chem Soc* 108 1126 1986.]

***N*-Acetylanthranilic acid** [89-52-1] M 179.1, m 182-184°, 185-186°, 190°(dec), pK²⁰ 3.61. Wash with distilled H₂O and recrystallise from aqueous AcOH, dry and recrystallise again from EtOAc. Also recryst from water or EtOH. [*J Chem Soc* 2495 1931; *J Am Chem Soc* 77 6698 1955.]

2-Acetylbenzoic acid [577-56-0] M 164.2, m 115-116°, 116-118°, pK²⁵ 4.10. Recrystallises from *C₆H₆ and H₂O (15g/100mL). The *oxime* has m 156-157°, and the *2,4-dinitrophenylhydrazone* has m 185-186°(needles from EtOH). [*J Am Chem Soc* 69 1547 1947.]

4-Acetylbenzoic acid [586-89-0] M 164.2, m 207.5-209.5°, 208.6-209.4°, pK²⁵ 3.70, 5.10 (EtOH). Dissolve in 5% aqueous NaOH, extract with Et₂O, and acidify the aqueous soln. Collect the ppte, and recrystallise from boiling H₂O (100 parts) using decolorising charcoal [*J Org Chem* 24 504 1959; *J Chem Soc* 265 1957; *J Am Chem Soc* 72 2882 1050, 74 1058 1952].

Acetylbenzoxonitrile [1443-80-7] M 145.2, m 57-58°. Recrystd from EtOH [Wagner et al. *J Am Chem Soc* 108 7727 1986].

4-Acetylbiphenyl [92-91-1] M 196.3, m 120-121°, b 325-327°/760mm. See 4'-phenylacetophenone on p. 327.

Acetyl-5-bromosalicylic acid [1503-53-3] M 259.1, m 168-169°, pK_{Est} ~3.0. Crystd from EtOH.

2-Acetylbutyrolactone [517-23-7] M 128.1, b 105°/5mm, 120-123°/11mm, 142-143°/30mm, d₄²⁰ 1.1846, n_D²⁰ 1.459. Purified by distillation, which will convert any free acid to the lactone, alternatively dissolve in Et₂O, wash well with 0.5N HCl, dry the organic layer and distil. The solubility in H₂O is 20% v/v. The *2,4-dinitrophenylhydrazone* forms orange needles from MeOH, m 146°. The lactone hydrolyses in mineral acid to 2-acetyl-4-hydroxybutyric acid which can be converted to the *di-n-propylamine salt* with m 68-70°. The lactone is a SKIN IRRITANT. [*Yakugaku Zasshi (J Pharm Soc Japan)* 62 417(439) 1942; *Helv Chim Acta* 35 2401 1952.]

Acetyl chloride [75-36-5] M 78.5, b 52°, d 1.1051, n 1.38976. Refluxed with PCl₅ for several hours to remove traces of acetic acid, then distd. Redistd from one-tenth volume of dimethylaniline or quinoline to remove free HCl. A.R. quality is freed from HCl by pumping it for 1h at -78° and distg into a trap at -196°.

Acetyl bromide [506-96-7] **M 123.0, b 76-77°, d 1.65.** Boiled with $\text{PBr}_3/\text{Ac}_2\text{O}$ for 1h then distd off and redistd. Store dry. [Burton and Degering *J Am Chem Soc* **62** 227 1940.] **LACHRYMATORY.**

Acetylcyclohexane (cyclohexyl methylketone) [823-76-7] **M 126.2, b 64°/11mm, 76.2-77°/25mm, d_4^{20} 0.9178, n_D^{20} 1.4519.** Dissolve in Et_2O , shake with H_2O , dry, evaporate and fractionate under reduced pressure. [UV: *J Am Chem Soc* **74** 518 1952; enol content: *J Org Chem* **19** 1960 1954.] The semicarbazone has **m 174°**; the 2,4-dinitrophenylhydrazone has **m 139-140°** [*Helv Chim Acta* **39** 1290 1956].

2-Acetylcyclohexanone [874-23-7] **M 140.2, m -11°, b 62-64°/2.5mm, 95-98°/10mm, 111-112°/18mm, d_4^{20} 1.08, n_D^{20} 1.51.** Dissolve in ligroin (b 30-60°), wash with saturated aqueous NaHCO_3 dry over Drierite and fractionate in a vacuum. [*J Am Chem Soc* **75** 626, 5030 1953; *Chem Ber* **87** 108 1954.] It forms a *Cu salt* which crystallises in green leaflets from EtOH , **m 162-163°** [UV: *J Chem Soc* 4419 1957].

2-Acetylcyclopentanone [1670-46-8] **M 126.2, b. 72-75°/8mm, 82-86°/12mm, 88°/18mm, d_4^{20} 1.043, n_D^{20} 1.490.** Dissolve in pet ether (b 30-60°), wash with satd aq NaHCO_3 , dry over Drierite and fractionate in a vacuum. It gives a violet colour with ethanolic FeCl_3 and is only slowly hydrolysed by 10% aq KOH but rapidly on boiling to yield 6-oxoheptanoic acid. [*J Am Chem Soc* **75** 5030 1953; *J Chem Soc* 4232 1956; UV: *J Am Chem Soc* **81** 2342 1959 .] It gives a gray green *Cu salt* from Et_2O -pentane, **m 237-238°** [*J Am Chem Soc* **79** 1488 1957].

Acetyldigitoxin- α [25395-32-8] **M 807.0, m 217-221°, $[\alpha]_D^{20} +5.0$ (c 0.7, pyridine).** Crystd from MeOH as plates.

Acetylene [74-86-2] **M 26.0, m -80.8°, b -84°, pK ~25.** If very impure it should be purified by successive passage through spiral wash bottles containing, in this order, satd aq NaHSO_4 , H_2O , 0.2M iodine in aq KI (two bottles), sodium thiosulfate soln (two bottles), alkaline sodium hydrosulfite with sodium anthraquinone-2-sulfonate as indicator (two bottles), and 10% aqueous KOH soln (two bottles). The gas was then passed through a Dry-ice trap and two drying tubes, the first containing CaCl_2 , and the second, Dehydrite [$\text{Mg}(\text{ClO}_4)_2$] [Conn, Kistiakowsky and Smith *J Am Chem Soc* **61** 1868 1939]. Acetone vapour can be removed from acetylene by passage through H_2O , then concd H_2SO_4 , or by passage through two gas traps at -65° and -80°, concd H_2SO_4 and a soda lime tower, a tower of 1-mesh Al_2O_3 then into H_2SO_4 [*Org Synth Coll Vol 1* 229 1941, 3 853 1955; 4 793 1963]. Sometimes it contains acetone and air. These can be removed by a series of bulb-to-bulb distns, e.g. a train consisting of a conc H_2SO_4 trap and a cold EtOH trap (-73°), or passage through H_2O and H_2SO_4 , then over KOH and CaCl_2 . [See Brandsma *Preparative Acetylenic Chemistry*, 1st Edn Elsevier 1971, for pK p15, ISBN 0444409475; 2nd Edn Elsevier 1988, ISBN 0444429603, and Chapter 5 for sodium acetylide.] It is also available commercially as 10ppm in helium, and several concentrations in N_2 for instrumental calibration.

Sodium acetylide [1066-26-8] **M 48.0,** was prepd by dissolving Na (23g) in liquid NH_3 (1L) and bubbling acetylene until the blue color was discharged (ca 30min) and evapd to dryness [Saunders *Org Synth Coll Vol III* 416 1955]; and is available commercially as a suspension in xylene/light mineral oil. [See entry in Chapter 5.]

Acetylenedicarboxamide [543-21-5] **M 112.1, m 294°(dec).** Crystd from MeOH .

Acetylenedicarboxylic acid [142-45-0] **M 114.1, m 179°(anhydrous), pK_1^{19} 1.04, pK_2^{19} 2.50.** Crystd from aqueous ether as dipicrate. For mono K salt see entry in Chapter 5.

***N*-Acetylenethylenediamine** [1001-53-2] **M 102.1, m 50-51°, 51°, b 128°/3mm, 125-130°/5mm, 133-139°/27mm, pK^{25} 9.28.** It has been fractionated under reduced pressure and fraction **b** 125-130°/5mm was refractionated; fraction **b** 132-135°/4mm was collected and solidified. It is a low melting *hygroscopic* solid which can be recrystd from dioxane- Et_2O . It is soluble in H_2O , Et_2O and $^*\text{C}_6\text{H}_6$. The *p-toluenesulfonate salt* can be recrystd from EtOH - EtOAc 1:8, has **m 125-126°** but the free base cannot be recovered from it by basifying and extracting with CH_2Cl_2 . The *picrate* has **m 175°** (from EtOH) [*J Am Chem Soc* **63** 853 1941, **78** 2570 1956].

2-Acetylfluorene [781-73-7] **M 208.3, m 132°.** Crystd from EtOH .

Acetyl fluoride [557-99-3] M 62.0, b 20.5°/760mm, d 1.032. Purified by fractional distn.

N-Acetyl-D-galactosamine [14215-68-0] M 221.2, m 160-161°, $[\alpha]_{546} +102^\circ$ (c 1, H₂O). Crystd from MeOH/Et₂O.

N-Acetyl-D-glucosamine [7512-17-6] M 221.2, m ca 215°, $[\alpha]_{546} +49^\circ$ after 2h (c 2, H₂O). Crystd from MeOH/Et₂O.

N-Acetylglutamic acid [1188-37-0] M 189.2, m 185° (RS); 201° (S), $[\alpha]^{25} -16.6^\circ$ (in H₂O), pK_{Est}(1) ~3.4, pK_{Est}(2) ~4.3. Likely impurity is glutamic acid. Crystd from boiling water.

N-Acetylglucinamide [2620-63-5] M 116.1, m 139-139.5°. Repeated crystn from EtOH/Et₂O. Dried in a vacuum desiccator over KOH.

N-Acetyl glycine [543-24-8] M 117.1, m 206-208°, pK₁²⁵ -1.92, pK₂²⁵ 3.69. Treated with acid-washed charcoal and recrystd three times from water or EtOH/Et₂O and dried *in vacuo* over KOH [King and King *J Am Chem Soc* 78 1089 1956].

N-Acetyl glycy-L-alaninamide [34017-20-4] M 175.2. Repeated crystn from EtOH/Et₂O. Dried in a vacuum desiccator over KOH.

N-Acetyl glycyglycinamide [27440-00-2] M 173.2, m 207-208°. Repeated crystn from EtOH/Et₂O. Dried in a vacuum desiccator over KOH.

N-Acetyl glycyglycyglycinamide [35455-24-4] M 230.2, m 253-255°. Repeated crystn from EtOH/Et₂O. Dried in a vacuum desiccator over KOH.

N-Acetylhistidine (H₂O) [39145-52-3] M 171.2, m 148° (RS); 169° (S) $[\alpha]^{25} +46.2^\circ$ (H₂O). Likely impurity is histidine. Crystd from water, then 4:1 acetone:water.

N-Acetyl-RS-homocysteine thiolactone (Citolone) [1195-16-0] [17896-21-8 for ±] M 159.2, m 110°, 109-111°, 111.5-112.5°. Dry in a vacuum desiccator and recrystallise from toluene as needles. It is a ninhydrin -ve substance which gives a "slow" nitroprusside test. λ_{\max} 238nm (ϵ 4,400 M⁻¹cm⁻¹); ν (nujol) 1789s and 851ms cm⁻¹. [*J Am Chem Soc* 78 1597 1956; *J Chem Soc* 2758 1963.]

N-Acetylimidazole [2466-76-4] M 110.1, m 101.5-102.5°, pK²⁵ 3.6. Crystd from isopropenyl acetate. Dried in a vacuum over P₂O₅.

3-Acetylindole [703-80-0] M 159.2, m 188-190°, 191-193°, 194°, pK²⁵ 12.99 (acidic). Recrystd from MeOH or *C₆H₆ containing a little EtOH. The *phenylureido* derivative has m 154°. [*J Chem Soc* 461 1946.]

Acetyl iodide [507-02-8] M 170.0, b 108°/760mm. Purified by fractional distn.

N-Acetyl-L-leucinamide [28529-34-2] M 177.2, m 133-134°. Recrystd from CHCl₃ and pet ether (b 40-60°).

3-(S-Acetylmercapto)isobutyric acid [RS 33325-40-5] M 162.2, m 40-40.5°, b ca 120°/1.25mm, pK_{Est} ~4.0. Distil under vacuum and recrystd from *C₆H₆. [*Chem Abstr* 38 3616 1944.]

Acetyl methanesulfonate [5539-53-7] M 170.2, b <120°/<0.01mm. The main impurity is methanesulfonic acid. Reflux with redistd acetyl chloride for 6-10h, i.e. until no further HCl is absorbed in a trap, and exclude moisture. Dist off excess of AcCl and carefully dist below 0.001mm with the bath temp below 120° to give the anhydride as a pale yellow oil which solidifies below 0°. Below ~130° it decomps to the

disulfonic anhydride and above $\sim 130^\circ$ polymers are formed. It is used for cleaving ethers [Prep, IR, NMR: Karger and Mazur *J Org Chem* **36** 528, 532 1971].

***N*-Acetyl-L-methionine** [65-82-7] M 191.3, m 104°, $[\alpha]_{546} -24.5^\circ$ (c 1, in H₂O), pK_{Est} ~ 3.4 . Crystd from water or ethyl acetate. Dried in a vacuum over P₂O₅.

Acetylmethionine nitrile [538-14-7] M 172.3, m 44-46°. Crystd from diethyl ether.

5-Acetyl-2-methoxybenzaldehyde [531-99-7] M 166.2, m 144°. Crystd from EtOH or Et₂O.

***N*-Acetyl-*N'*-methyl-L-alanimide** [19701-83-8] M 144.2. Crystd from EtOAc/Et₂O, then from EtOH and Et₂O.

4-Acetyl-1-methyl-1-cyclohexene [6090-09-1] M 138.2, b 73-75°/7.5mm, 85-86°/13mm, 94-94.7°/20mm, 204.5-206°/747mm, d_4^{20} 1.0238, n_D^{20} 1.469. Purified by fractionation under reduced pressure *in vacuo*, and when almost pure it can be fractionated at atmospheric pressure, preferably in an inert atm. Forms two *semicarbazones* one of which is more soluble in *C₆H₆, and both can be recryst from EtOH, more soluble has m 149°(151°), and the less soluble has m 172-175°(191°). *4-Nitrophenylhydrazone* has m 166-167° and the *2,4-dinitrophenylhydrazone* has m 114-115°. [*Helv Chim Acta* **17** 129, 140 1934; *Justus Liebigs Ann Chem* **564** 109 1949.]

***N*-Acetyl-6*N'*-methylglycinamide** [7606-79-3] M 130.2. Recrystd from EtOH/Et₂O mixture.

***N*-Acetyl-6*N'*-methyl-L-leucine amide** [32483-15-1] M 186.3. Recrystd from EtOH/hexane mixture.

4-Acetylmorpholine [1696-20-4] M 129.2, m 13.8-14°, 14°, 14.5°, b 96-97°/6mm, 113-128°/22mm, 242-247°/760mm, d_4^{20} 1.0963, n_D^{20} 1.4830. Distd through an 8inch Fenske (glass helices packing) column with a manual take-off head. Purified by fractional distn. The *hydrobromide* has m 172-175°. [*J Am Chem Soc* **75** 357 1953, *J Org Chem* **21** 1072 1956.]

1-Acetylnaphthalene [941-98-0] M 170.1, m 10.5°, b 93-95°/0.1mm, 167°/12mm, 302°/atm, d_4^{20} 1.12, pK -6.22 (H₀ scale, aq H₂SO₄). If the NMR spectrum indicates the presence of impurities, probably 2-acetylnaphthalene, convert the substance to its picrate by dissolving in *benzene or EtOH and adding excess of satd picric acid in these solvents until separation of picrates is complete. Recryst the picrate till m is 118°. Decompose the picrate with dil NaOH and extract with Et₂O. Dry the extract (Na₂SO₄), filter, evap and dist. The *2,4-dinitrophenylhydrazone* crystals from EtOH and has m 259°. [*Justus Liebigs Ann Chem* **380** 95 1911; *J Am Chem Soc* **61** 3438 1939.]

2-Acetylnaphthalene (2-acetonaphthenone, β -Acetonaphthone, 2-acetonaphthalene, methyl-2-naphthylketone) [93-08-3] M 170.2, m 52-53°, 55°, 55.8°, b 164-166°/8mm, 171-173°/17mm, 301-303°/atm, pK -6.16 (H₀ scale, aq H₂SO₄). Separated from the 1-isomer by fractional crystn of the picrate in EtOH (see entry for the 1-isomer above) m 82°. Decomposition of the picrate with dil NaOH and extraction with Et₂O then evaporation gives purer 2-acetylnaphthalene. If this residue solidifies it can be recrystd from pet ether, EtOH or acetic acid; otherwise it should be distilled in a vac and the solid distillate is recrystd [Gorman and Rodgers *J Am Chem Soc* **108** 5074 1986; Levanon et al. *J Phys Chem* **91** 14 1987]. Purity should be checked by high field NMR spectroscopy. *Oxime* has m 145° decomp, and the *semicarbazone* has m 235°. [*Justus Liebigs Ann Chem* **380** 95 1911; *J Am Chem Soc* **72** 753 and 5626 1950, *J Org Chem* **5** 512 1940.]

***N*-Acetyl-D-penicillamine** [15537-71-0] M 191.3, m 189-190° (dec), $[\alpha]_D +18^\circ$ (c 1, in 50% EtOH). See *N*-acetyl penicillamine on p. 507 in Chapter 6.

***N*-Acetyl-L-phenylalanine** [2018-61-3] M 207.2, m 170-171°, $[\alpha]_D +41^\circ$ (c 1, EtOH), (DL) m 152.5-153°, pK_{Est} 3.5. Crystd from CHCl₃ and stored at 4°. (DL)-isomer crystd from water or acetone.

***N*-Acetyl-L-phenylalanine ethyl ester** [2361-96-8] M 235.3, m 93-94°. Crystd from aq EtOH or H₂O. [Izumiya and Fruton *J Biol Chem* 218 59 1956.]

1-Acetyl-2-phenylhydrazine [114-83-0] M 150.2, m 128.5°, pK²⁵ 1.3. Crystd from aq EtOH.

1-Acetylpiperazine [13889-98-0] M 128.2, m 32-34°, 52°, pK²⁵ 7.94. Purified by recrystn from 40% aqueous EtOH or from EtOH-Et₂O. It is an irritant, and is *hygroscopic*. The *hydrochloride* has m 191° (from EtOH), and the *tosylate* has m 148-149° (from EtOH-EtOAc, 1:16). The free base, however, cannot be isolated by basifying the tosylate salt and extractn with CH₂Cl₂. [*Chem Ber* 66 113 1933; *J Am Chem Soc* 75 4949 1953, 2570 78 1956.]

1-Acetyl-4-piperidone [32161-06-1] M 141.2, b 124-128°/0.2mm, 218°/760mm, d₄²⁵ 1.1444, n_D²⁵ 1.5023. Purified by fractional distn through a short Vigreux column (15mm). The 2,4-dinitrophenylhydrazone has m 212-213° (from EtOH). It is freely soluble in H₂O but insoluble in Et₂O. [*J Am Chem Soc* 901 71 1949.]

3-Acetylpyridine [350-03-8] M 121.1, m 13-14°, b 65-66°/1mm, 92-95°/8-9mm, 105°(113°)/16mm, 219-221°/760mm, d₄²⁰ 1.1065, n_D²⁰ 1.1065, pK²⁵ 3.18. It is purified by dissolving in HCl, extracting with Et₂O to remove the possible impurity of nicotinic acid, basified with NaOH and extracted with Et₂O. The dried extract is filtered, evaporated and the residual oil distd. If the NMR spectrum indicates further impurities then convert to the *phenylhydrazone* (m 137°, yellow needles from EtOH). This is hydrolysed with HCl [*Chem Ber* 22 597 1889], the phenylhydrazine HCl is removed by filtration, NaNO₂ is added, the soln is basified with aq NaOH and extracted with Et₂O as before and distd at atmospheric pressure to give 3-acetylpyridine as a colourless oil. Purification can be achieved by shaking with 50% aq KOH, extracting with Et₂O, drying the extract and distilling at atmospheric pressure or in a vacuum. [*J Am Chem Soc* 79 4226 1957]. The *hydrochloride* has m 180-181° (from MeOH-EtOH), the *picrate* has m 133.8-134.8° (from H₂O), and the *phenylhydrazone* has m 137° (129-130°) (from EtOH) [*J Am Chem Soc* 71 2285 1949]. The *ketoxime* has m 112° (from EtOH or *C₆H₆). [*J Am Chem Soc* 55 816 1933, 63 490 1941, 67 1468 1945, 79 4226 1957.]

Acetylsalicylic acid (Aspirin) [50-78-2] M 180.2, m 133.5-135°, pK²⁵ 3.38, (pK¹⁷ 4.56). Crystd twice from toluene, washed with cyclohexane and dried at 60° under vacuum for several hours [Davis and Hetzer *J Res Nat Bur Stand* 60 569 1958]. Has also been recrystd from isopropanol and from diethyl ether/pet ether (b 40-60°).

***O*-Acetylsalicyloyl chloride** [5538-51-2] M 198.6, m 45°, 46-49°, 48-52°, b 107-110°/0.1mm, 135°/12mm, n_D²⁰ 1.536. Check first the IR to see if an OH frequency is present. If so then some free acid is present. Then reflux with acetyl chloride for 2-3h and fractionate at high vac. The distillate should crystallise. It can be recryst from hexane. [*J Chem Soc* 89 1318 1906.]

***O*-Acetylsalicylsalicylic acid** [530-75-6] M 300.3, m 159°. Crystd from dilute acetic acid.

***N*-(4)-Acetylsulfanilamide** [144-80-9] M 214.2, m 216°. Crystd from aqueous EtOH.

2-Acetylthiazole [24295-03-2] M 127.2, b 89-91° (90-95°)/12mm, 95-105°/15mm, d₄²⁰ 1.23, n_D²⁰ 1.55. Check NMR spectrum, if not too bad, distil through an efficient column in a vacuum. The *oxime* sublimes at 140-145°, m 159° (cryst from H₂O) has m 163-165.5° [*Helv Chim Acta* 31 1142 1948; *J Am Chem Soc* 79 4524 1957; *Helv Chim Acta* 40 554 1957].

2-Acetylthiophene (methyl 2-thienyl ketone) [88-15-3] M 126.2, m 9.2-10.5°, 10.45°, 10-11°, b 77°/4mm, 89-91°/9mm, 94.5-96.5°/13mm, 213-214°/atm, d₄²⁰ 1.17, n_D²⁰ 1.5666. Fractionally distd through a 12 plate column and fraction b 77°/4mm was collected. Also wet the acetylthiophene in order to remove and free thiophene which forms an azeotrope with H₂O, b 68°. Store in a brown bottle and the clear colourless liquid remains thus for extended periods. [*Org Synth* 28 1 1948; *J Am Chem Soc* 69 3093 1947.] The red 4-nitrophenylhydrazone crystals from EtOH, m 181-182°.

3-Acetylthiophene (methyl 3-thienyl ketone) [1468-83-3] M 126.2, m 57°, 60-63°, b 106-107°/25 mm, 208-210°/748mm. Recrystd from pet ether (b 30-60°) or EtOH. 2,4-dinitrophenylhydrazone crystallises from CHCl₃, m 265°, and the semicarbazone crystallises from EtOH, m 174-175°. [*J Am Chem Soc* 70 1555 1948.]

N-Acetylthiourea [591-08-2] M 118.2, m 164-165°, 165-168°. Recrystd from AcOH, the solid is washed with Et₂O and dried in air then at 100°. [*Collect Czech Chem Commun* 24 3678 1959.]

Acetyl p-toluenesulfonate [26908-82-7] M 214.2, m 54-56°. The most likely impurity is p-toluenesulfonic acid (could be up to 10%). This can be removed by dissolving in dry Et₂O and cooling until the anhydride crystallises out. It decomp on heating; below ~130° it gives the disulfonic anhydride and above ~130° polymers are formed. It is used for cleaving ethers [Prep, IR, NMR: Karger and Mazur *J Org Chem* 36 528, 532 1971].

1-O-Acetyl-2,3,5-tri-O-benzoyl-β-D-ribofuranose [6974-32-9] M 504.5, m 128-130°, 130-131°, 131-132°, [α]_D²⁰ +44.2° (c 1, CHCl₃). Recrystd from EtOH or isoPrOH. [*Helv Chim Acta* 42 1171 1959; NMR: *J Org Chem* 33 1799 1968; IR: *Chem Pharm Bull Jpn* 11 188 1963.]

N-Acetyltryptophan M 246.3, [87-32-1] m 206° (RS), pK_{Est} ~3.8; [1218-34-4] m 188° (S), [α]²⁵ +30.1° (aq NaOH). Likely impurity is tryptophan. Crystd from EtOH by adding water.

N-Acetyl-L-valine amide [37933-88-3] M 158.2, m 275°. Recrystd from CH₃OH/Et₂O.

cis-Aconitic acid [585-84-2] M 174.1, m 126-129°(dec). Crystd from water by cooling (sol: 1g in 2mL of water at 25°). Dried in a vacuum desiccator.

trans-Aconitic acid (1,2,3-propenetrisarcoxylic acid) [4023-65-8] M 174.1, m 195°(dec), m 198-199°(dec), 204-205°(dec), pK₁²⁵ 2.81, pK₂²⁵ 4.46. Purified by dissolving in AcOH (77g/150mL), filtering and cooling. The acid separates (55g) as colourless needles. A further quantity (10g) can be obtained by reducing the vol of the filtrate. The acid is dried in air then in a vacuum desiccator over NaOH. The acid can be recrystd from Me₂CO-CHCl₃. The highest m is obtained with the very dry acid. The m (209°) is obtained on a Dennis bar [*J Am Chem Soc* 52 3128 1930, *Org Synth Coll Vol II* 12 1943].

cis-Aconitic anhydride [6318-55-4] M 156.1, m 75°, 76-78°, 78-78.5°. Reflux in xylene (7.5 parts) for 1h, then evaporate and recrystallise the residue from *C₆H₆. Alternatively, reflux in Ac₂O, evaporate and recrystallise from *C₆H₆. It is sensitive to moisture. [IR: *Acta Chem Scand* 21 291 1967, *Chem Ber* 61 2523 1928; NMR: *Biochemistry* 5 2335 1966.]

Aconitine [302-27-2] M 645.8, m 204°, [α]₅₄₆ +20° (c 1, CHCl₃), pK₁₅ 8.35. Crystd from EtOH, CHCl₃ or toluene.

Aconitine hydrobromide [6034-57-7] M 726.7, m 207°. Crystd from water or EtOH/ether.

Acridine (2,3-benzoquinoline) [260-94-6] M 179.2, m 111° (sublimes), b 346°, pK 5.58 (pK²⁵ of excited state 10.65). Crystd twice from *benzene/cyclohexane, or from aqueous EtOH, then sublimed, removing and discarding the first 25% of the sublimate. The remainder was again crystd and sublimed, discarding the first 10-15% [Wolf and Anderson *J Am Chem Soc* 77 1608 1955].

Acridine can also be purified by crystn from n-heptane and then from ethanol/water after pre-treatment with activated charcoal, or by chromatography on alumina with pet ether in a darkened room. Alternatively, acridine can be ppted as the hydrochloride from *benzene soln by adding HCl, after which the base is regenerated, dried at 110°/50mm, and crystd to constant melting point from pet ether [Cumper, Ginman and Vogel *J Chem Soc* 4518 1962]. The regenerated free base may be recrystd, chromatographed on basic alumina, then vac-sublimed and zone-refined. [Williams and Clarke, *J Chem Soc, Faraday Trans 1* 73 514 1977; Albert, *The Acridines*

Arnold Press 1966]. It can exist in five crystalline forms and is steam volatile. It is a strong IRRITANT to skin and mucous membranes and can become a chronic irritant— handle with CARE.

Acridine Orange [494-38-2] **M 349.94, m 181-182° (free base)**. The double salt with $ZnCl_2$ (6g) was dissolved in water (200mL) and stirred with four successive portions (12g each) of Dowex-50 ion-exchange resin (K^+ form) to remove the zinc. The soln was then concentrated in vacuum to 20mL, and 100mL of ethanol was added to ppt KCl which was removed. Ether (160mL) was added to the soln from which, on chilling, the dye crystallises as its chloride. It was separated by centrifuging, washed with chilled ethanol and ether, and dried under vac, before being recryst from ethanol (100mL) by adding ether (50mL), and chilling. Yield 1g. [Pal and Schubert *J Am Chem Soc* **84** 4384 1962].

It was recrystd twice as the free base from ethanol or methanol/water by dropwise addition of NaOH (less than 0.1M). The ppt was washed with water and dried under vacuum. It was dissolved in $CHCl_3$ and chromatographed on alumina: the main sharp band was collected, concentrated and cooled to -20° . The ppt was filtered, dried in air, then dried for 2h under vacuum at 70° . [Stone and Bradley *J Am Chem Soc* **83** 3627 1961; Blauer and Linschitz *J Phys Chem* **66** 453 1962.]

Acridine Yellow G [135-49-9] **M 273.8, m 325°, CI 46025**. Crystd from 1:1 *benzene/methanol.

Acridone [578-95-0] **M 195.2, m >300°, pK₁ -0.32 (basic), pK₂ 14 (acidic)**. Dissolve ~1g in ca 1% NaOH (100mL), add 3M HCl to pH 4 when acridone separates as a pale yellow solid with **m** just above 350° (sharp). It can be recrystd from large vols of H_2O to give a few mg. It is soluble in 160 parts of boiling EtOH (540 parts at 22°) [*J Chem Soc* 1294 1956]. A few decigrams are best crystallised as the *hydrochloride* from 400 parts of 10N HCl (90% recovery) from which the free base is obtained by washing the salt with H_2O . A small quantity can be recrystd (as the neutral species) from boiling AcOH. Larger quantities are best recrystallised from a mixture of 5 parts of freshly distd aniline and 12.5 parts of glacial acetic acid. Acridone distils unchanged at atmospheric pressure, but the boiling point was not recorded, and some sublimation occurs below 350° . It has UV: λ_{max} 399nm. [see Albert, *The Acridines* Arnold Press pp. 201, 372 1966.]

N-(9-Acridinyl)maleimide (NAM) [49759-20-8] **M 274.3, m 248°, 255-258°**. Purified by chromatography on silica gel using CH_2Cl_2 as eluant. Evaporation of pooled fractions that gave the correct NMR spectra gave a solid which was recrystd from Me_2CO as pale yellow prisms. IR ν (nujol): 1710 (imide); UV (MeOH): λ_{max} (nm), (ϵ $M^{-1}cm^{-1}$): 251 (159 500), 343 shoulder (7 700), 360 (12 400) and 382shoulder (47 000). [*Chem Pharm Bull Jpn* **26** 596 1978; *Eur J Biochem* **25** 64 1972.]

Acriflavine [8048-52-0] **M 196.2, pK >12**. Treated twice with freshly ppted AgOH to remove proflavine, then recrystd from absolute methanol [Wen and Hsu *J Phys Chem* **66** 1353 1962].

Acriflavin Mixture (Euflavin, 3,6-diamino-10-methylacridinium chloride) [8063-24-9] **M 259.7, m 179-181°**. Purified by dissolving in 50 parts of H_2O , shake with a small excess of freshly ppted and washed Ag_2O . The mixture is set aside overnight at 0° and filtered. The cake is not washed. The pH of the filtrate is adjusted to 7.0 with HCl and evaporated to dryness. The residue is then crystd twice from MeOH, twice from H_2O and dried at 120° . λ_{max} at 452nm has a log ϵ value of 4.67. It is a red powder which readily absorbs H_2O . The solubility is increased in the presence of proflavin. The *dihydrochloride* is a deep red crystn powder. It is available as a mixture of 3,6-diaminoacridinium chloride (35%) and its 10-metho-chloride (65%). [see Albert, *The Acridines* Arnold Press p. 346 1966; *Chem Ber* **45** 1787 1912].

Acrolein (acraldehyde) [107-02-8] **M 56.1, b 52.1°, n 1.3992, d 0.839**. Purified by fractional distn. under nitrogen, drying with anhydrous $CaSO_4$ and then distilling under vac. Blacet, Young and Roof [*J Am Chem Soc* **59** 608 1937] distd under nitrogen through a 90cm column packed with glass rings. To avoid formation of diacryl, the vapour was passed through an ice-cooled condenser into a receiver cooled in an ice-salt mixture and containing 0.5g catechol. The acrolein was then distd twice from anhydrous $CuSO_4$ at low pressure, catechol being placed in the distilling flask and the receiver to avoid polymerization. [Alternatively, hydroquinone (1% of the final soln) can be used.]

Acrolein diacetyl acetal (1,1-diacetoxy-2-propene). [869-29-4] M 158.2, b 75°/10 mm, 184°/atm, d_4^{20} 1.08, n_D^{20} 1.4203. Check the NMR spectrum. If it is not satisfactory then add Ac₂O and a drop of conc H₂SO₄ and heat at 50° for 10min. Then add anhydrous NaOAc (ca 3g/100g of liquid) and fractionate. Note that it forms an azeotrope with H₂O, so do not add H₂O at any time. It is a **highly flammable and TOXIC** liquid, keep away from the skin. [*J Am Chem Soc* 73 5282 1951.]

Acrolein diethyl acetal [3054-95-3] M 130.2, b 120-125°/atm, n_4^{20} 1.398-1.407. Add Na₂CO₃ (ca 3.5%) and distil using an efficient column, or better a spinning band column. [*Org Synth* 25 1 1945.]

Acrolein dimethyl acetal (1,1-dimethoxy-2-propene) [6044-68-4] M 102.1, b 87.5-88°/750mm, 89-90°/760mm, d_4^{20} 0.86, n_D^{20} 1.3962. Fractionally distil (after adding 0.5g of hydroquinone) under reduced press through an all glass column (40cm x 2.5 cm) packed with glass helices and provided with a heated jacket and a total reflux variable take-off head. Stainless steel Lessing rings (1/8 x 1/8 in) or gauze have been used as packing. It is a **highly flammable and TOXIC** liquid, keep away from the skin. [*J Chem Soc* 2657 1955.]

Acrolein semicarbazone [6055-71-6] M 113.1, m 171°. Crystd from water.

Acrylamide [79-06-1] M 71.1, m 84°, b 125°/25mm. Crystd from acetone, chloroform, ethyl acetate, methanol or *benzene/chloroform mixture, then vac dried and kept in the dark under vac. Recryst from CHCl₃ (200g dissolved in 1L heated to boiling and filtered without suction in a warmed funnel through Whatman 541 filter paper. Allowed to cool to room temp and kept at -15° overnight). Crystals were collected with suction in a cooled funnel and washed with 300mL of cold MeOH. Crystals were air-dried in a warm oven. [Dawson et al. *Data for Biochemical Research*, Oxford Press 1986 p. 449.]

CAUTION: Acrylamide is extremely **TOXIC** and precautions must be taken to avoid skin contact or inhalation. Use gloves and handle in a well ventilated fume cupboard.

Acrylic acid [79-10-7] M 72.1, m 13°, b 30°/3mm, d 1.051, pK^{25} 4.25. Can be purified by steam distn, or vacuum distn through a column packed with copper gauze to inhibit polymerisation. (This treatment also removes inhibitors such as methylene blue that may be present.) Azeotropic distn of the water with *benzene converts aqueous acrylic acid to the anhydrous material.

Acrylonitrile [107-13-1] M 53.1, b 78°, d 0.806, n^{25} 1.3886. Washed with dilute H₂SO₄ or dilute H₃PO₄, then with dilute Na₂CO₃ and water. Dried with Na₂SO₄, CaCl₂ or (better) by shaking with molecular sieves. Fractionally distd under nitrogen. Can be stabilised by adding 10ppm *tert*-butyl catechol. Immediately before use, the stabilizer can be removed by passage through a column of activated alumina (or by washing with 1% NaOH soln if traces of water are permissible in the final material), followed by distn. Alternatively, shaken with 10% (w/v) NaOH to extract inhibitor, and then washed in turn with 10% H₂SO₄, 20% Na₂CO₃ and distd water. Dried for 24h over CaCl₂ and fractionally distd under N₂ taking the fraction boiling at 75.0 to 75.5°C (at 734mm Hg). Stored with 10ppm *tert*-butyl catechol. Acrylonitrile is distilled off as required. [Burton et al, *J Chem Soc, Faraday Trans 1* 75 1050 1979.]

Acryloyl chloride [814-68-6] M 90.5, b 72-74°/740mm, 74°/760mm, d_4^{20} 1.1127, n_D^{20} 1.4337. Distil rapidly through an efficient 25cm column after adding 0.5g of hydroquinone/200g of chloride, and then redistil carefully at atmospheric pressure preferably in a stream of dry N₂. [*J Am Chem Soc* 72 72, 2299 1950.] **The liquid is an irritant and is TOXIC.**

Actarit (p-acetamidophenylacetic acid) [18699-02-0] M 193.2, m 174-175°. Crystd from MeOH + Me₂CO or aq EtOH.

Adamantane [281-23-2] M 136.2, m 269.6-270.8° (sublimes). Crystd from acetone or cyclohexane, sublimed in a vacuum below its melting point. [Butler et al. *J Chem Soc, Faraday Trans 1* 82 535 1986.] Adamantane was also purified by dissolving in *n*-heptane (ca 10mL/g of adamantane) on a hot plate, adding activated charcoal (2g/100g of adamantane), and boiling for 30min, filtering the hot soln through a filter paper, concentrating the filtrate until crystn just starts, adding one quarter of the original volume *n*-heptane and

allowing to cool slowly over a period of hours. The supernatant was decanted off and the crystals were dried on a vacuum line at room temperature. [Walter et al. *J Am Chem Soc* **107** 793 1985.]

1-Adamantane acetic acid [4942-47-6] **M 194.3, m 136°**, **pK_{Est} ~4.8**. Dissolve in hot N NaOH, treat with charcoal, filter and acidify. Collect solid, wash with H₂O, dry and recryst from MeOH. [*Chem Ber* **92** 1629 1959.]

1-Adamantane carboxylic acid [828-51-3] **M 180.3, m 175-176.5°, 177°, pK_{Est} ~4.9**. Possible impurities are trimethylacetic acid and C9 and C13 acids. Dissolve 15g of acid in CCl₄ (300mL) and shake with 110mL of 15N aqueous NH₃ and the ammonium salt separates and is collected. Acid impurities form soluble ammonium salts. The salt is washed with cold Me₂CO (20mL) and suspended in H₂O (250mL). This is treated with 12N HCl and extracted with CHCl₃ (100mL). The dried (Na₂SO₄) is evaporated and the residue recrystd from a mixture of MeOH (30mL) and H₂O (ca 10mL) to give the pure acid (10-11g). [*Org Synth Coll Vol.V* **20** 1973.] Also recrystd from absolute EtOH and dried under vacuum at 100°.

Alternatively, the acid (5g) is refluxed for 2h with 15mL of MeOH and 2mL of 98% H₂SO₄ (cool when mixing this soln). Pour into 10 volumes of H₂O and extract with the minimum volume of CHCl₃ to give clear separation of phases. The extract is washed with H₂O and dried (CaCl₂) and distd. The methyl ester is collected at 77-79°/1mm, **m 38-39°**. The ester is hydrolysed with the calculated amount of N KOH and refluxed until clear. Acidification with HCl provides the pure acid with 90% recovery. [*Org Synth* **4** 1 1964.] The *amide* crystals from cyclohexane, **m 189°**. [*Chem Ber* **62** 1629 1959.]

1,3-Adamantane diamine dihydrochloride [26562-81-2] **M 239.2, m >310°, pK_{Est(1)} ~8.1, pK_{Est(2)} ~10.1**. Dissolve in boiling conc HCl (400mg in 15mL) and evaporate to dryness. Dissolve in absolute EtOH and add dry Et₂O to crystallise the *dihydrochloride*. [*Chem Ber* **93** 1366 1960.]

1,3-Adamantane dicarboxylic acid [39269-10-8] **M 224.3, m 276°, 276-278°, 279°, pK_{Est(1)} ~4.9, pK_{Est(2)} 5.9**. Dissolve in aq NaOH, treat with charcoal, filter and acidify with dilute HCl. Recryst from MeOH. [*Chem Ber* **93** 1366 1960.]

1-Adamantane methylamine [17768-41-1] **M 165.3, b 83-85°/0.3mm, d₄²⁰ 0.93, pK_{Est} ~10.2**. Dissolve in Et₂O, dry over KOH and distil. The *N-Tosyl* derivative has **m 134-135°** (from EtOH). [*Chem Ber* **96** 550 1963.]

1-Adamantanol (1-hydroxyadamantane) [768-95-6] **M 152.4, m 288.5-290°**. If 2-adamantanol is a suspected impurity then dissolve substance (10g) in acetone (100mL) and Jones's reagent {CrO₃ (10.3g) in H₂O (30mL)} and conc H₂SO₄ (8.7mL) is added dropwise (turns green in colour) until excess reagent is present (slight red colour). Allow to stir overnight, decant the acetone soln from the Cr salts and adamantan-2-one, and dry (Na₂SO₄) and evaporate to dryness. The residue (ca 7g) is chromatographed through Al₂O₃ (250g) and washed with 50% *benzene-pet ether (b 40-60°), then 100% Et₂O (to remove any adamantan-2-one present) and the 1-adamantanol is then eluted with 5% MeOH in Et₂O. The eluate is evaporated, and the residue is recrystd from pet ether (b 30-60°) at -70°, **m 287.2-288.5°**. It has characteristic IR, ν 3640, 1114, 1086, 982 and 930cm⁻¹. [*J Am Chem Soc* **83** 182 1961.]

Alternatively, if free from the 2-isomer, dissolve in tetrahydrofuran, dilute with H₂O to ppt the alcohol. Collect, dry and sublime in a vacuum at 130°. [*Chem Ber* **92** 1629 1959.]

2-Adamantanol (2-hydroxyadamantane) [700-57-2] **M 152.4, m 296.2-297.7°**. Can be purified by chromatography as for the 1-isomer. It crystallises from cyclohexane and has characteristic IR, ν 3600, 1053, 1029 and 992cm⁻¹ [*J Am Chem Soc* **8** 182 1961].

2-Adamantanone [700-58-3] **M 150.2, m 256-258°(sublimes)**. Purified by repeated sublimation *in vacuo*. [Butler et al. *J Chem Soc, Faraday Trans 1* **82** 535 1986.]

N-(1-Adamantyl)acetamide [880-52-4] **M 193.3, m 149°**. Wash well with H₂O, dry and recrystallise from cyclohexane. *It is an irritant*. [*Chem Ber* **92** 1629 1959.]

1-Adamantylamine [768-94-5] M 151.2, m 160-190°, 208-210°, pK²⁵ 10.58. Dissolve in Et₂O, dry over KOH, evaporate and sublime in a vacuum. [*Chem Ber* 93 226 1960.]

1-Adamantylamine hydrochloride [665-66-7] M 187.7, m 360° (dec). Dissolve in dry EtOH, add a few drops of dry EtOH saturated with HCl gas, followed by dry Et₂O to crystallise the hydrochloride. Dry the salt in vacuum. [*Chem Ber* 93 226 1960.]

2-Adamantylamine hydrochloride [10523-68-9] M 187.7, m >300°, pK_{Est} ~10.4. The free amine in Et₂O, liberated by the action of alkali in H₂O, is dried over KOH, filtered, evap and sublimed at 110°/12Torr, m 230-236°. The base is dissolved in EtOH and crystd by the addition of Et₂O, and dried in vac. [*Justus Liebigs Ann Chem* 658 151 1962].

1-Adamantyl bromide [768-90-1] M 215.1, m 117-119°, 118°, 119.5-120°. If coloured, dissolve in CCl₄, wash with H₂O, treat with charcoal, dry (CaCl₂), filter, evap to dryness. Dissolve in a small volume of MeOH and cool in a CO₂/trichloroethylene bath and collect the crystals. Sublime at 90-100°/water pump vacuum. [*Chem Ber* 92 1629 1959; *J Am Chem Soc* 83 2700 1961.]

1-Adamantyl bromomethylketone [5122-82-7] M 257.2, m 76-79°, 78-79°. Dissolve in Et₂O, wash with H₂O, dry (MgSO₄), evaporate and crystallise residue from small volumes of MeOH. **LACHRYMATORY.** [*Chem Ber* 93 2054 1960.]

1-Adamantyl chloride [935-56-8] M 170.7, m 164.3-165.6°. Crystd from aqueous MeOH and sublimed at 100°/12Torr. Also crystd from MeOH at -70°. [*Chem Ber* 92 1629 1959; *J Am Chem Soc* 83 2700 1961.]

1-Adamantyl fluoride (1-fluoroadamantane) [768-92-3] M 154.2, m 210-212° (dec), 259-260° (dec). Dissolve in Et₂O, dry over Na₂SO₄, evaporate to dryness and sublime the residue at 90-100°/12mm. Recryst sublimate from MeOH, m 259-260°. [*Zh Org Khim* 30 1609 1965.] To remove 1-hydroxyadamantane impurity, dissolve in cyclohexane cool for many hours, filter off the hydroxyadamantane, and evaporate to dryness. Recrystallise the residue from pet ether at -77° and sublime in vacuum, m 210-212° dec (sealed tube). [*J Org Chem* 30 789 1965.]

1-Adamantyl fluoroformate [62087-82-5] M 198.2, m 31-32°. Dissolve in *n*-hexane (*ca* 10g in 150 mL) and keep at 0° for 24h. Any 1-adamantanol present will separate. Filter and evaporate to dryness. Crystalline residue has m 31-32° (ν 1242, 1824 and 2340 cm⁻¹). There should be no OH str band above 2500 cm⁻¹. [*Z Phys Chem* 357 1647 1976; Haas et al. *J Am Chem Soc* 88 1988 1966.]

1-Adamantyl iodide (1-iodoadamantane) [768-93-4] M 262.1, m 75.3-76.4°. Dissolve in Et₂O, shake with aqueous NaHSO₃, aqueous K₂CO₃, and H₂O, dry (Na₂SO₄), evaporate and recrystallise from MeOH at -70° (to avoid alcoholysis) giving white crystals. [*J Am Chem Soc* 83 2700 1961; lit m of 151-152.5° is incorrect.] Also purified by recrystn from pet ether (40-60°C) followed by rigorous drying and repeated sublimation.

1-Adamantyl isocyanate [4411-25-0] M 177.3, m 144-145°. Recryst from *n*-hexane and sublime. **Irritant.** [*Chem Ber* 95 2302 1962.]

1-Adamantyl isothiocyanate [4411-26-1] M 193.3, m 168-169°. Dissolve in Et₂O, wash with H₂O, dry (Na₂SO₄), evaporate and sublime the residue in a vacuum at 140°, and recryst from MeOH. **Irritant.** [*Chem Ber* 95 2302 1962.]

***N*-(1-Adamantyl)urea** [13072-69-0] M 194.2, m >250° (dec), 268-272° (dec). Wash with H₂O and dioxane and recryst from EtOH. [*Chem Ber* 95 2302 1962.]

Adenine [73-24-5] M 135.1, m 360-365° (dec rapid heating), pK₁²⁵ 4.12, pK₂²⁵ 9.83. Crystd from distd water.

Adenosine [58-61-7] M 267.3, m 234-236°, $[\alpha]_{546} -85^\circ$ (c 2, 5% NaOH), pK_1^{25} 3.48, pK_2^{25} 12.5. Crystd from distilled water.

Adipic acid [124-04-9] M 146.1, m 154°, pK_1^{25} 4.44, pK_2^{25} 5.45. For use as a volumetric standard, adipic acid was crystd once from hot water with the addition of a little animal charcoal, dried at 120° for 2h, then recrystd from acetone and again dried at 120° for 2h. Other purification procedures include crystn from ethyl acetate and from acetone/petroleum ether, fusion followed by filtration and crystn from the melt, and preliminary distn under vac.

Adiponitrile (1,4-dicyanobutane) [111-69-3] M 108.14, m 2.4°, b 123°/0.5mm, 153°/6mm, 175°/26mm, 184°/30mm, 295°/atm, d_4^{20} 0.9396, n_D^{20} 1.4371. Reflux over P₂O₅ and POCl₃, and fractionally distil, then fractionate through an efficient column. **The liquid is TOXIC and is an IRRITANT.** [Chem Ber 67 1770 1934; Justus Liebigs Ann Chem 596 127 1955; Can J Chem 34 1662 1956; J Am Chem Soc 62 228 1940.]

Adonitol (Ribitol) [488-81-3] M 152.2, m 102°. Crystallise from EtOH by addition of diethyl ether.

Adrenalin see epinephrine.

Adrenochrome [54-06-8] M 179.2, m 125-130°. Crystd from MeOH/formic acid, as hemihydrate, and stored in a vacuum desiccator.

Adrenosterone (Reichstein's G) [382-45-6] M 300.4, m 220-224°. Crystd from EtOH. Can be sublimed under high vacuum.

Agaricic acid [1-(n-hexadecyl)citric acid] [666-99-9] M 416.6, m 142°(dec), $[\alpha]_D -9.8^\circ$ (in NaOH), $pK_{Est(1)}$ ~2.7, $pK_{Est(2)}$ ~4.2, $pK_{Est(3)}$ ~5.5. Crystd from EtOH.

Agmatine sulfate [5-guanidinopent-1-ylamine sulfate] [2482-00-0] M 228.3, m 231°, $pK_{Est(1)}$ ~9.1, $pK_{Est(2)}$ ~13.0. Crystd from aqueous MeOH.

Agroclavin [548-42-5] M 238.3, m 198-203°(dec), 205-206°, $[\alpha]_D^{20} -155^\circ$ (c 1, CHCl₃), pK_{Est} ~8.0. Crystd from diethyl ether.

Ajmalicine [483-04-5] M 352.4, m 250-252°(dec), $[\alpha]_{546} -76^\circ$ (c 0.5, CHCl₃), pK_{Est} ~7.4. Crystd from MeOH.

Ajmalicine hydrochloride [4373-34-6] M 388.9, m 290°(dec), $[\alpha]_D -17^\circ$ (c 0.5, MeOH). Crystd from EtOH.

Ajmaline [γ -yohimbine] [4360-12-7] M 326.4, m 160° (MeOH), 205-206° (anhyd), $[\alpha]_D^{20} +144^\circ$ (c 0.8, CHCl₃), pK_{Est} ~7.5. Crystd from MeOH.

Ajmaline hydrochloride [4410-48-4] M 388.9, m 140°. Crystd from water.

Alanine (RS) [302-72-7] M 89.1, m 295-296°, (S) [56-41-7] m 297°(dec), $[\alpha]_D^{15} +14.7^\circ$ (in 1M HCl), pK_1^{25} 2.34, pK_2^{25} 9.87. Crystd from water or aqueous EtOH, e.g. crystd from 25% EtOH in water, recrystd from 62.5% EtOH, washed with EtOH and dried to constant weight in a vacuum desiccator over P₂O₅. [Gutter and Kegeles J Am Chem Soc 75 3893 1953.] 2,2'-Iminodipropionic acid is a likely impurity.

β -Alanine [107-95-9] M 89.1, m 205°(dec), pK_1^{25} 3.55, pK_2^{25} 10.24. Crystd from filtered hot saturated aqueous soln by adding four volumes of absolute EtOH and cooling in an ice-bath. Recrystd in the same way and then finally, crystd from a warm saturated soln in 50% EtOH by adding four volumes of absolute

EtOH cooled in an ice bath. Crystals were dried in a vacuum desiccator over P_2O_5 . [Donovan and Kegeles *J Am Chem Soc* **83** 255 1961.]

S-Alaninol [*S*-2-Aminopropan-1-ol] [2749-11-3] **M 75.1, b 167-169°/760mm, d_4^{20} 0.961, n_D^{20} 1.456, $[\alpha]_{546} +26.0^\circ$ (c 2, EtOH), pK^{25} 9.43.** Purification as for *S*-2-amino-3-methylbutan-1-ol

Aldol (3-hydroxybutanal) [107-89-1] **M 88.1, b 80-81°/20mm.** An ethereal soln was washed with a saturated aqueous soln of $NaHCO_3$, then with water. The non-aqueous layer was dried with anhydrous $CaCl_2$ and distd immediately before use. The fraction, **b 80-81°/20mm**, was collected, [Mason, Wade and Pouncy *J Am Chem Soc* **76** 2255 1954].

Aldosterone [52-39-1] **M 360.5, m 108-112°(hydrate), 164°(anhydr), $[\alpha]_D^{25} +161^\circ$ (c 1, $CHCl_3$).** Crystd from aqueous acetone. *Acetate*, cryst from $Me_2CO + Et_2O$, has **m 198-199°**, $[\alpha]_D^{24} +121.7^\circ$ (c 0.7, $CHCl_3$)

Aldrin [309-00-2] **M 354.9, m 103-104.5°.** Crystd from MeOH. **POISONOUS**

Aleuritic acid [*RS-erythro-9,10,16-trihydroxyhexadecanoic acid*] [533-87-9] **M 304.4, m 100-101°.** Crystd from aqueous EtOH. *Hydrazide* cryst from EtOH has **m 139-140°.**

Alginic acid [9005-32-7] **M 48,000-186000.** To 5g in 550mL water containing 2.8g $KHCO_3$, were added 0.3mL acetic acid and 5g potassium acetate. EtOH to make the soln 25% (v/v) in EtOH was added and any insoluble material was discarded. Further addition of EtOH, to 37% (v/v), ppted alginic acid. [Pal and Schubert *J Am Chem Soc* **84** 4384 1962.]

Aliquat 336 (methyltricaprylammonium chloride, tri-*n*-octylmethylammonium chloride) [5137-55-3] **M 404.2, d 0.884.** A 30% (v/v) soln in *benzene was washed twice with an equal volume of 1.5M HBr. [Petrow and Allen, *Anal Chem* **33** 1303 1961.] Purified by dissolving 50g in $CHCl_3$ (100mL) and shaking with 20% NaOH soln (200mL) for 10min, and then with 20% NaCl (200mL) for 10min. Washed with small amount of H_2O and filtered through a dry filter paper [Adam and Pribil *Talanta* **18** 733 1971].

Alizarin (1,2-dihydroxyanthraquinone) [72-48-0] **M 240.2, m 290°, d 0.884, pK_1^{25} 7.45, pK_2^{25} 11.80.** Crystd from glacial acetic acid or 95% EtOH. Can also be sublimed at 110°/2mm.

Alizarin-3-methyliminodiacetic acid (Alizarin Complexone) (2 H_2O) [3952-78-1] **M 421.4, m 189°(dec), $pK_{Est(1)} \sim 4.9$, $pK_{Est(2)} \sim 7.5$.** Purified by suspending in 0.1M NaOH (1g in 50mL), filtering the solution and extracting alizarin with 5 successive portions of CH_2Cl_2 . Then add HCl dropwise to precipitate the reagent, stirring the solution in a bath. Filter ppte on glass filter, wash with cold water and dry in a vacuum desiccator over KOH [Ingman *Talanta* **20** 135 1973].

Alizarin Yellow R [5-(4-nitrophenylazosalicylic acid), Mordant Orange I] [2243-76-7] **M 287.2, m 253-254°(dec), $>300^\circ$, pK^{25} 11.17.** The free acid is ppted by adding HCl to an aq soln of the Na salt. After 2 recrystns from aq AcOH, it has **m 255°(dec)**; [m 253-254° dec was reported *J Chem Soc* **79** 49 1901]. The free acid can be recrystd from dilute AcOH as orange brown needles. The Na salt changes colour from yellow to red when the pH is increased from 10.2 to 12.0. [*J Am Chem Soc* **75** 5838 1953.]

***n*-Alkylammonium chloride $n=2,4,6$.** Recrystd from EtOH or an EtOH/ Et_2O mixture. [Hashimoto and Thomas *J Am Chem Soc* **107** 4655 1985; Chu and Thomas *J Am Chem Soc* **108** 6270 1986.]

***n*-Alkyltrimethylammonium bromide $n=10,12,16$.** Recrystd from an EtOH/ Et_2O mixture. [Hashimoto and Thomas *J Am Chem Soc* **107** 4655 1985.]

Allantoin [97-59-6] **M 158.1, m 238°(dec).** Crystd from water or EtOH.

Allene (prodiene) [463-49-0] M 40.1, m -146°, b -32°. Frozen in liquid nitrogen, evacuated, then thawed out. This cycle was repeated several times, then the allene was frozen in a methyl cyclohexane-liquid nitrogen bath and pumped for some time. Also purified by HPLC. [Cripps and Kiefer *Org Synth* 42 12 1962.]

(-)-Alloaromadendrene [25246-27-9] M 204.4, b 96°/2mm, 265-267°/atm, $[\alpha]_D^{25}$ -22° (neat), d_4^{20} 0.923, n_D^{23} 1.501. Fractionally distd from Na. IR has bands at 6.06 and 11.27 μ due to C=CH₂. [*J Chem Soc* 715 1953; cf *J Am Chem Soc* 91 6473 1969.]

neo-Allocimene (tc-2,6-dimethyl-2,4,6-octatriene) [7216-56-0] M 136.2, b 80°/13mm, 196-198°/atm, d_4^{20} 0.8161, n_D^{20} 1.5437. Fractionally distd through an efficient column and stabilised with ca 0.1% of hydroquinone. UV: λ_{max} nm(ϵ M⁻¹cm⁻¹) 290 (32 500), 279 (41 900) and 270 (32 600). [*Justus Liebigs Ann Chem* 609 1 1957; *Anal Chem* 26 1726 1954.]

5 α -Allopregnane-3 α ,20 α -diol [566-58-5] M 320.5, m 248-248.5°, $[\alpha]_D$ +17° (c 0.15, EtOH). Crystd from EtOH.

D-Allothreonine [2R,3R(-)-isomer] [24830-94-2] M 119.1, m 272-273°(dec), 276°(dec), $[\alpha]_D^{25}$ -9.1° (c 3.9, H₂O), pK_1^{25} 2.11, pK_2^{25} 9.10. Recrystd from aqueous EtOH or 50% EtOH. [*J Chem Soc* 62 1950; *J Am Chem Soc* 194 455 1952; IR: Greenstein & Winitz *The Chemistry of the Amino Acids* J. Wiley, Vol 3 1961.]

Alloxan [2,4,5,6(1H,3H)pyrimidine, tetrone] [50-71-5] M 142.0, m -170°(dec), pK^{25} 6.64. Crystn from water gives the tetrahydrate. Anhydrous crystals are obtained by crystn from acetone, glacial acetic acid or by sublimation *in vacuo*.

Alloxan monohydrate [2244-11-3] M 160.1, m 255°(dec), pK 6.64. Recryst from H₂O as the *tetrahydrate* in large prisms or rhombs. On heating at 100°, or on exposure to air, this is converted to the *monohydrate*. Dissolve it in its own weight of boiling H₂O and cool for several days below 0° [the *tetrahydrate* crystallises from soln much more slowly when free from HNO₃. It is less sol in HCO₃ solns than in H₂O]. Drying the solid over H₂SO₄ yields the *monohydrate*. The *anhydrous* crystals can be obtained by recrystn from dry Me₂CO or AcOH followed by washing with dry Et₂O or by sublimation in a vacuum. On heating it turns pink at 230° and decomposes at ca 256°. It is acidic to litmus. [*Org Synth Coll Vol III* 37 1955.] It forms a compound with urea which crystallises from H₂O in yellow needles that become red at 170° and dec at 185-186°.

Alloxantin [76-24-4] M 286.2, m 253-255°(dec) (yellow at 225°). Crystd from water or EtOH and kept under nitrogen. Turns red in air.

Allyl acetate [591-87-7] M 100.1, b 103°, d 0.928, n_4 1.40488, n_D^{27} 1.4004. Freed from peroxides by standing with crystalline ferrous ammonium sulfate, then washed with 5% NaHCO₃, followed by saturated CaCl₂ soln. Dried with Na₂SO₄ and fractionally distd in an all-glass apparatus.

Allylacetic acid (pent-4-enoic acid) [591-80-0] M 100.1, m -22.5°, b 83-84°/12mm, 90°/15mm, d_4^{20} 0.9877, n_D^{20} 1.4280, pK^{25} 4.68. Distil through an efficient column (allyl alcohol has b 95-97°). It is characterised as the *S-benzyl isothiuronium salt* m 155-158° (96% EtOH, aq EtOH) [*Acta Chem Scand* 9 1425 1955], *4-bromophenacyl ester* m 59.5-60.5° (from 90% EtOH). Solubility at 18°: in pyridine (57%), AcOH (7.3%), MeOH (5.4%), Me₂CO (3.2%), MeOAc (2.8%), EtOH (5.4%), H₂O (1.8%), PrOH (1.6%), isoPrOH (0.27%). [*J Am Chem Soc* 74 1894 1952.]

Allyl alcohol [107-18-6] M 58.1, b 98°, d_4 0.857, n_D 1.4134. Can be dried with K₂CO₃ or CaSO₄, or by azeotropic distn with *benzene followed by distn under nitrogen. It is difficult to obtain peroxide free. Also reflux with magnesium and fractionally distd [Hands and Norman *Ind Chem* 21 307 1945].

Allylamine [107-11-9] M 57.1, b 52.9°, d 0.761, n 1.42051, pK²⁵ 9.49. Purified by fractional distn from calcium chloride. Causes sneezing and tears.

1-Allyl-6-amino-3-ethyluracil [642-44-4] M 195.2, m 143-144° (anhydr). Crystd from water (as monohydrate).

Allyl bromide [106-95-6] M 121, b 70°, d 1.398, n 1.46924. Washed with NaHCO₃ soln then distd water, dried (CaCl₂ or MgSO₄), and fractionally distd. Protect from strong light. **LACHRYMATOR, HIGHLY TOXIC and FLAMMABLE.**

Allyl butyl ether [3739-64-8] M 114.2, b 64-65°/120mm, 117.8-118°/763mm, d₄²⁰ 1.4057, n_D²⁰ 0.7829. Check the IR for the presence of OH str vibrations, if so then wash well with H₂O, dry with CaCl₂ and distil through a good fractionating column. **The liquid is an irritant.** [*J Org Chem* 23 1666 1958; *J Am Chem Soc* 73 3528 1951.]

Allyl chloride [107-05-1] M 76.5, b 45.1°, d 0.939, n 1.4130. Likely impurities include 2-chloropropene, propyl chloride, *i*-propyl chloride, 3,3-dichloropropane, 1,2-dichloropropane and 1,3-dichloropropane. Purified by washing with conc HCl, then with Na₂CO₃ soln, drying with CaCl₂, and distn through an efficient column [Oae and Vanderwerf *J Am Chem Soc* 75 2724 1953]. **LACHRYMATOR, TOXIC.**

Allyl chloroformate [2937-50-0] M 120.5, b 56°/97mm, 109-110°/atm, d₄²⁰ 1.14, n_D²⁰ 1.4223. Wash several times with cold H₂O to remove alcohol and HCl and dry over CaCl₂. It is **important** to dry well before distilling *in vacuo*. Note that the receiver should be cooled in ice to avoid loss of distillate into the trap and vacuum pump. The liquid is **highly TOXIC and flammable.** [*J Am Chem Soc* 72 1254 1950.]

Allyl cyanide (3-butene nitrile) [109-75-1] M 67.1, b -19.6°/1.0mm, 2.9°/5 mm, 14.1°/5mm, 26.6°/20mm, 48.8°/60mm, 60.2°/100mm, 98°/400mm, 119°/760mm, d₄²⁰ 0.8341, n_D²⁰ 1.406. It should be redistd at atmospheric pressure then distilled under a vacuum to remove final traces of HCN from the residue. Note that the residue from the first distiln may be difficult to remove from the flask and should be treated with conc HNO₃ then H₂O and finally hot EtOH (**CARE**). Allyl cyanide has an onion-like odour and is stable to heat. It forms a complex with AlCl₃ (2:2) m 41°, and (3:2) m 120°. **All operations should be done in an efficient fume hood as the liquid is flammable and HIGHLY TOXIC.** [*Org Synth Coll Vol I* 46 1941.]

Allyl disulfide (diallyl disulfide) [2179-57-9] M 146.3, b 58-59°/5mm, 79-81°/20mm, 138-139°/atm, d₄²⁰ 1.01, n_D²⁰ 1.541. Purified by fractional distn until their molar refractivities are in uniformly good agreement with the calculated values [*J Am Chem Soc* 69 1710 1947]. Also purified by gas chromatography [retention times: *J Org Chem* 24 175 1959; UV: *J Chem Soc* 395 1949].

RS- α -Allylglycine (2-aminopent-4-enoic acid). [7685-44-1] M 115.1, m 250-255°(dec), pK_{Est(1)} ~2.3, pK_{Est(2)} ~9.6. Dissolve in absolute EtOH and ppte with pyridine, then recrystallise from aqueous EtOH [R_F in BuOH:EtOH:NH₃:H₂O (4:4:1:1:) 0.37]. The *hydrobromide* has m 136-140° (from EtOAc) and the *phenylureido* derivative has m 159-161°. [*Monatsh Chem* 89 377 1958.]

1-N-Allyl-3-hydroxymorphinan [152-02-3] M 283.4, m 180-182°. Crystd from aqueous EtOH.

Allyl iodide (3-iodopropene) [556-56-9] M 167.7, b 103°, d¹² 1.848. Purified in a dark room by washing with aq Na₂SO₃ to remove free iodine, then drying with MgSO₄ and distilling at 21mm pressure, to give a very pale yellow liquid. (This material, dissolved in hexane, was stored in a light-tight container at -5° for up to three months before free iodine could be detected, by its colour in the soln) [*Sibbett and Noyes J Am Chem Soc* 75 761 1953].

5-Allyl-5-isobutylbarbituric acid [77-26-9] M 224.3, m 139°, 139-140°, 140-142°, pK³⁸ 12.36. It can be recrystallised from H₂O or dilute EtOH, and sublimes at 100-120°/8-12mm. It is soluble in *C₆H₆, cyclohexane, tetralin and pet ether at 20°. [J Am Chem Soc 77 1486 1955.]

Allylisocyanate [1476-23-9] M 83.1, b 84°/atm, 87-89°/atm, d₄²⁰ 0.94, n_D²⁰ 1.417. Purify as for allylthiocyanate.

Allylthiocyanate [57-06-7] M 99.2, m -80°, b 84-85°/80mm, 150°/760mm, 151°/atm, d₄²⁰ 1.017, n_D²⁰ 1.5268. Fractionate using an efficient column, preferably in a vacuum. It is a yellow pungent irritating and TOXIC (suspected CARCINOGEN) liquid. Store in a sealed tube under N₂. The *N'*-benzylthiourea derivative has m 94.5° (from aq EtOH) [J Am Chem Soc 74 1104 1952].

Allyl Phenyl sulfide [5296-64-0] M 150.2, b 59-60°/1.5mm, 79-80°/3mm, 114-114.3°/23.5mm, 225-226°/740mm, 215-218°/750mm, d₄²⁰ 1.0275, n_D²⁰ 1.5760. Dissolve in Et₂O, wash with alkali, H₂O, dry over CaCl₂, evaporate and fractionally distil, preferably under vacuum. It should not give a ppte with an alcoholic soln of Pb(OAc)₂. [J Am Chem Soc 52 3356 1930, 74 48 1952.]

***N*-Allylthiourea (thiosinamine)** [109-57-9] M 116.2, m 70-73°, 78°. Recrystd from H₂O. Soluble in 30 parts of cold H₂O, soluble in EtOH but insoluble in *C₆H₆. Also recrystd from acetone, EtOH or ethyl acetate, after decolorising with charcoal. The white crystals have a bitter taste with a slight garlic odour and are TOXIC. [Anal Chem 21 421 1949.]

***N*-Allylurea** [557-11-9] M 100.1, m 85°. Crystd from EtOH, EtOH/ether, EtOH/chloroform or EtOH/toluene.

Aloin [10-glucopyranosyl-1,8-dihydroxy-3-(hydroxymethyl)-9(10*H*)anthracenone, Barbaloin] [8015-61-0] M 418.4, m 148-148.5°, 148-150°. Lemon yellow crystals from H₂O (450g/1.5L) as the *monohydrate* which has a lower m (70-80°). [J Chem Soc 2573 1932, 3141 1956.]

D-Altrose [1990-29-0] M 180.2, m 103-105°, [α]₅₄₆ +35° (c 7.6, H₂O). Crystd fom aq EtOH.

Amberlite IRA-904 Anion exchange resin (Rohm and Haas) [9050-98-0]. Washed with 1M HCl, CH₃OH (1:10) and then rinsed with distilled water until the washings were neutral to litmus paper. Finally extracted successively for 24h in a Soxhlet apparatus with MeOH, *benzene and cyclohexane [Shue and Yan Anal Chem 53 2081 1981]. Strongly basic resin also used for base catalysis [Fieser & Fieser Reagents for Org Synth 1 511, Wiley 1967].

Aminoacetaldehyde dimethyl acetal (2,2-dimethoxyethylamine) [22483-09-6] M 105.1, m <-78°, b 139.5°/768mm, 137-139°/atm, d₄²⁰ 0.9676 n_D²⁰ 1.4144. Dry over KOH pellets and distil through a 30cm vac jacketed Vigreux column. [J Am Chem Soc 75 3398 1953, 77 6640 1955.]

***p*-Aminoacetanilide** [122-80-5] M 150.2, m 162-163°. Crystd from water.

Aminoacetic acid (Glycine) [56-40-6] M 75.1, m 262° (dec, goes brown at 226°, sublimes at 200°/0.1mm), pK₁²⁵ 2.35, pK₂²⁵ 9.78. Crystd from distilled water by dissolving at 90-95°, filtering, cooling to about -5°, and draining the crystals centrifugally. Alternatively, crystd from distilled water by addition of MeOH or EtOH (e.g. 50g dissolved in 100mL of warm water, and 400mL of MeOH added). The crystals can be washed with MeOH or EtOH, then with diethyl ether. Likely impurities are ammonium glycinate, iminodiacetic acid, nitrilotriacetic acid, ammonium chloride.

Aminoacetonitrile bisulfate [151-63-3] M 154.1, m 188°(dec) Crystd from aqueous EtOH.

Aminoacetonitrile hydrochloride [6011-14-9] M 92.5, m 166-167°, 172-174°, pK²⁵ 5.34. Recrystd from dil EtOH *hygroscopic* leaflets. Best to crystallise from absolute EtOH-Et₂O (1:1) and then recryst from absolute EtOH. The m recorded range from 144° to 174°. The free base has b 58°/15mm with

partial decomposition. [*J Prakt Chem* [2] **65** 189 1902; *J Am Chem Soc* **56** 2197 1934; *J Chem Soc* 1371 1947.]

2-Aminoacetophenone hydrochloride [5468-37-1] **M 171.6, m 188°(dec), 194°(dec), pK²⁵ 5.34.** Crystd from acetone/EtOH or 2-propanol [Castro *J Am Chem Soc* **108** 4179 1986].

***m*-Aminoacetophenone** [99-03-6] **M 135.2, m 98-99°, pK²⁵ 3.56.** Recrystd from EtOH.

***p*-Aminoacetophenone** [99-92-3] **M 135.2, m 104-106°, 105-107°, b 293°/atm, pK²⁵ 2.19** Recryst from CHCl₃, *C₆H₆ or H₂O. Soluble in hot H₂O. UV (EtOH) has λ_{max} 403nm (log ε 4.42) [*J Am Chem Soc* **75** 2720 1953]. [*Anal Chem* **26** 726 1954.] The *2,4-dinitrophenylhydrazone* has **m 266-267°** (from CHCl₃ or EtOH), and the *semicarbazone* has **m 193-194°(dec)**(from MeOH) and the *hydrochloride* has **m 98°(dec)**(from H₂O).

α-Amino acids see Chapter 6.

9-Aminoacridine [**9-acridineamine**] [90-45-9] **M 194.2, m 241°, pK²⁰ 9.95.** Crystd from EtOH or acetone and sublimes at 170-180°/0.04mm [Albert and Ritchie *Org Synth Coll Vol III* 53 1955; for hydrochloride see Chapter 6.]

***dl*-α-Aminoadipic acid (hydrate)** [542-32-5] **M 161.2, m 196-198°, pK_{Est(1)} ~2.0, pK_{Est(2)} ~4.5, pK_{Est(3)} ~9.8.** Crystd from water.

2-Amino-4-anilino-*s*-triazine [537-17-7] **M 168.2, m 235-236°, pK_{Est} ~5.5.** Crystd from dioxane or 50% aqueous EtOH.

1-Aminoanthraquinone-2-carboxylic acid [82-24-6] **M 276.2, m 295-296°.** Crystd from nitrobenzene.

4-Aminoantipyrine [83-07-8] **M 203.3, m 109°.** Crystd from EtOH or EtOH/ether.

***p*-Aminoazobenzene (*p*-phenylazoaniline)** [60-09-3] **M 197.2, m 126°, pK²⁵ ~2.82.** Crystd from EtOH, CCl₄, pet ether/*benzene, or a MeOH/water mixture.

***o*-Aminoazotoluene (Fast Garnet GBC base, 4'-amino-2,3'-dimethylazobenzene)** [97-56-3] **M 225.3, m 101.4-102.6°, CI 11160, pK_{Est} ~2.8.** Crystd twice from EtOH, once from *benzene, then dried in an Abderhalden drying apparatus [Cilento *J Am Chem Soc* **74** 968 1952]. **CARCINOGENIC.**

2-Aminobenzaldehyde [529-23-7] **M 121.1, m 39-40°, pK²⁰ 1.36.** Distd in steam and crystd from water or EtOH/ether.

2-Aminobenzaldehyde phenylhydrazone (Nitrin) [63363-93-9] **M 211.3, m 227-229°.** Crystd from acetone. [Knöpfer *Monatsh Chem* **31** 97 1910.]

3-Aminobenzaldehyde [29159-23-7] **M 121.1, m 28-30°, pK_{Est} ~2.0.** Crystd from ethyl acetate.

4-Aminobenzamide hydrochloride [59855-11-7] **M 199.6, m 284-285°, pK_{Est} ~1.7.** Recrystd from EtOH.

***p*-Aminobenzeneazodimethylaniline** [539-17-3] **M 240.3, m 182-183°.** Crystd from aqueous EtOH.

***o*-Aminobenzoic acid (anthranilic acid)** [118-92-3] **M 137.1, m 145°, pK₁²⁵ 2.94, pK₂²⁵ 4.72.** Crystd from water (charcoal). Has also been crystd from 50% aqueous acetic acid. Can be vacuum sublimed.

***m*-Aminobenzoic acid** [99-05-8] M 137.1, m 174°, pK₁²⁵ 3.29, pK₂²⁵ 5.10. Crystd from water.

***p*-Aminobenzoic acid** [150-13-0] M 137.1, m 187-188°, pK₁²⁵ 2.45, pK₂²⁵ 4.85. Purified by dissolving in 4-5% aqueous HCl at 50-60°, decolorising with charcoal and carefully precipitating with 30% Na₂CO₃ to pH 3.5-4 in the presence of ascorbic acid. It can be crystd from water, EtOH or EtOH/water mixtures.

***p*-Aminobenzonitrile** [873-74-5] M 118.1, m 86-86.5°, 85-87°, pK²⁵ 1.74. Crystd from water, 5% aqueous EtOH or EtOH and dried over P₂O₅ or dried *in vacuo* for 6h at 40°. [Moore et al. *J Am Chem Soc* 108 2257 1986; Edidin et al. *J Am Chem Soc* 109 3945 1987.]

4-Aminobenzophenone [1137-41-3] M 197.2, m 123-124°, pK²⁵ 2.17. Dissolved in aq acetic acid, filtered and ppted with ammonia. Process repeated several times, then recrystd from aqueous EtOH.

2-Aminobenzothiazole [136-95-8] M 150.2, m 132°, pK²⁰ 4.48. Crystd from aqueous EtOH.

6-Aminobenzothiazole [533-30-2] M 150.2, m 87°, pK_{Est} ~3. Crystd from aqueous EtOH.

***N*-(*p*-Aminobenzoyl)-L-glutamic acid** [4271-30-1] M 266.3, m 173° (L-form), [α]₅₄₆ -17.5° (c 2, 0.1m HCl); 197° (DL), pK_{Est(1)} ~1.7, pK_{Est(2)} ~3.4, pK_{Est(3)} ~4.3. Crystd from H₂O.

3-*o*-Aminobenzyl-4-methylthiazolium chloride hydrochloride [534-94-1] M 277.4, m 213°(dec). Crystd from aqueous EtOH.

4-Amino-1-benzylpiperidine [50541-93-0] M 190.3, b ~180°/20mm, d 0.933, n 1.543, pK_{Est(1)} ~ 8.3 pK_{Est(2)} ~ 10.4. Purified by distn *in vacuo*, and stored under N₂, because it absorbs CO₂. The *dihydrochloride salt* [1205-72-7] has m 270-273° (255°) after recrystn from MeOH + EtOAc or EtOH. [*J Chem Soc* 3165, 3172 1957.] The *4-methylamino-1-benzylpiperidine* derivative has b 168-172°/17mm, n 1.5367 [*J Am Chem Soc* 70 4009 1948]. The *1-(1-benzyl-4-piperidinyl)-3-cyano-2-methylisothiourea* derivative has m 160° from CHCl₃/Et₂O [Prepn, IR, NMR: Ried et al. *Chem Ber* 116 1547 1983].

***o*-Aminobiphenyl** [90-41-5] M 169.2, m 49.0°, pK¹⁸ 3.83. Crystd from aqueous EtOH (charcoal).

***p*-Aminobiphenyl** [92-67-1] M 169.2, m 53°, b 191°/16mm, pK¹⁸ 4.38. Crystd from water or EtOH. CARCINOGENIC.

2-Amino-5-bromotoluene [583-75-5] M 186.1, m 59°, pK²⁵ 3.58. Steam distd, and crystd from EtOH.

***RS*-2-Aminobutyric acid** [2835-81-6] M 103.1, m 303°(dec), pK₁²⁵ 2.29, pK₂²⁵ 9.83. Crystd from water.

***S*-2-Aminobutyric acid** [1492-24-6] M 103.1, m 292°(dec), [α]_D + 20.4° (c 2, 2.5N HCl). Crystd from aqueous EtOH.

3-Aminobutyric acid [2835-82-7] M 103.1, m 193-194°, pK_{Est(1)} ~3.5, pK_{Est(2)} ~10.3. Crystd form aqueous EtOH or MeOH + Et₂O.

4-Aminobutyric acid (GABA) [56-12-2] M 103.1, m 202°(dec), pK₁²⁵ 4.14, pK₂²⁵ 10.55. Crystd form aqueous EtOH or MeOH + Et₂O.

2-Amino-5-chlorobenzoic acid [635-21-1] M 171.6, m 100°, pK₁²⁵ 1.69, pK₂²⁵ 4.35. Crystd from water, EtOH or chloroform.

3-Amino-4-chlorobenzoic acid [2840-28-0] M 171.6, m 216-217°, pK_{Est(1)} ~2.7, pK_{Est(2)} ~2.9. Crystd from water.

4-Amino-4'-chlorobiphenyl [135-68-2] M 203.5, m 134°, pK_{Est} ~4.0. Crystd from pet ether.

2-Amino-4-chloro-6-methylpyrimidine [5600-21-5] M 143.6, m 184-186°, pK_{Est} ~1.0. Crystd from EtOH.

2-Amino-5-chloropyridine [1072-98-6] M 128.6, m 135-136°, pK 4.38. Crystd from pet ether, sublimes at 50°/0.5mm.

1-Amino-1-cyclopentanecarboxylic acid (cycloleucine) [52-52-8] M 129.2, m 330°(dec), pK_{Est(1)} ~2.5 pK_{Est(2)} ~10.3. Crystd from aq EtOH.

2-Amino-3,5-dibromopyridine [35486-42-1] M 251.9, m 103-104°, pK_{Est} ~2.4. Steam distd and crystd from aqueous EtOH or pet ether.

2-Amino-4,6-dichlorophenol [527-62-8] M 175.0, m 95-96°, pK_{Est(1)} ~3.1, pK_{Est(2)} ~6.8. Crystd from CS₂ or *benzene.

3-Amino-2,6-dichloropyridine [62476-56-6] M 164.0, m 119°, b 110°/0.3mm, pK_{Est} ~2.0. Crystd from water.

4-Amino-N,N-diethylaniline hydrochloride [16713-15-8] M 200.7, m 233.5°, pK²² 6.61. Crystd from EtOH.

4-Amino-3,5-diiodobenzoic acid [2122-61-4] M 388.9, m >350°, pK_{Est(1)} 0.4, pK_{Est(2)} ~1.6. Purified by soln in dilute NaOH and pptn with dilute HCl. Air dried.

2-Amino-4,6-dimethylpyridine [5407-87-4] M 122.2, m 69-70.5°, pK 7.84. Crystd from hexane, ether/pet ether or *benzene. Residual *benzene was removed over paraffin-wax chips in an evacuated desiccator.

2-Amino-4,6-dimethylpyrimidine [767-15-7] M 123.2, m 152-153°, pK²⁵ 4.95. Crystn from water gives m 197°, and crystn from acetone gives m 153°.

2-Aminodiphenylamine [534-85-0] M 184.2, m 79-80°, pK_{Est(1)} ~3.8 (NH₂), pK_{Est(2)} <~0. Crystd from H₂O.

4-Aminodiphenylamine [101-54-2] M 184.2, b 155°/0.026mm, pK²⁵ 5.20. Crystn from EtOH gives m 66°, and crystn from ligroin gives m 75°.

2-Amino-1,2-diphenylethanol [530-36-9] M 213.3, m 165°, pK_{Est(1)} ~7.5. Crystd from EtOH.

2-Aminodiphenylmethane [28059-64-5] M 183.3, m 52°, b 172°/12mm and 190°/22mm, pK_{Est(1)} ~4.2. Crystd from ether.

2-Aminoethanethiol see cysteamine in Chapter 6.

2-Aminoethanol (ethanolamine) [141-43-5] M 61.1, f 10.5°, b 72-73°/12mm, 171.1°/760mm, d 1.012, n 1.14539, pK²⁵ 9.51. Decomposes slightly when distd at atmospheric pressure, with the formation of conducting impurities. Fractional distn at about 12mm pressure is satisfactory. After distn, 2-aminoethanol was further purified by repeated washing with ether and crystn from EtOH (at low temperature). After fractional distn in the absence of CO₂, it was twice crystd by cooling, followed by distn.

Hygroscopic. [Reitmeier, Silvertz and Tartar *J Am Chem Soc* **62** 1943 1940.] It can be dried by azeotropic distn with dry *benzene.

2-Aminoethanol hydrochloride [2002-24-6] M 97.6, m 75-77°. Crystd from EtOH. It is deliquescent.

2-Aminoethyl hydrogen sulfate (sulfuric acid mono-2-aminoethyl ester) [926-39-6] M 141.1, m 285-287° (chars at 275°). Crystd from water or dissolved in water and EtOH added.

S-(2-Aminoethyl)isothiuronium bromide hydrobromide [56-10-0] M 281.0, m 194-195°. Crystd from absolute EtOH/ethyl acetate. It is *hygroscopic*.

(2-Aminoethyl)trimethylammonium chloride hydrochloride (chloramine chloride hydrochloride) [3399-67-5] M 175.1, m 260°(dec). Crystd from EtOH. (Material is very soluble in H₂O).

2-Aminofluorene [153-78-6] M 181.2, m 127.8-128.8°, 132-133°, pK²⁵ 4.64. Wash well with H₂O and recrystd from Et₂O or 50% aqueous EtOH (25g with 400mL), and dry in a vacuum. Store in the dark. [*Org Synth Coll Vol II* 447 1943; *Coll Vol V* 30 1973].

4-Amino hippuric acid [61-78-9] M 194.2, m 198-199°, pK_{Est(1)}~1.7(NH₂), pK_{Est(2)}~3.4 (CO₂H). Crystd from H₂O.

4-Amino-3-hydrazino-5-mercapto-1,2,4-triazole (Purpald) [1750-12-5] M 146.2, m 228-230°(dec), 234-235°(dec), pK_{Est(1)}~2, pK_{Est(2)}~3 (NH₂), pK_{Est(3)}~8 (SH). Crystd from H₂O (0.6g in 300-400mL). The *benzylidene deriv* has m 245-246°(dec) from *i*-PrOH [Hogarth *J Chem Soc* 4817 1952].

1-Amino-4-hydroxyanthraquinone [116-85-8] M 293.2, m 207-208°, pK_{Est(1)}~2.6 (NH₂), pK_{Est(2)}~9.0 (OH). Purified by TLC on SiO₂ gel plates using toluene/acetone (9:1) as eluent. The main band was scraped off and extracted with MeOH. The solvent was evaporated and the dye was dried in a drying pistol [Land, McAlpine, Sinclair and Truscott *J Chem Soc, Faraday Trans 1* 72 2091 1976]. Crystd from aq EtOH.

dl-4-Amino-3-hydroxybutyric acid [924-49-2] M 119.1, m 225°(dec), pK₁²⁵~3.80 (CO₂H), pK_{Est(2)}~9.3. Crystd from H₂O or aqueous EtOH.

5-Amino-8-hydroxyquinoline hydrochloride [3881-33-2] M 196.7, pK₁²⁰ 5.67, pK₂²⁰ 11.24. Dissolved in minimum of MeOH, then Et₂O was added to initiate pptn. Ppte was filtered off and dried [Lovell et al. *J Phys Chem* **88** 1885 1984].

3-Amino-4-hydroxytoluene [95-84-1] M 123.2, m 137-138°, pK_{Est(1)}~4.7(NH₂), pK_{Est(2)}~9.6 (OH). Crystd from H₂O or toluene.

4-Amino-5-hydroxytoluene [2835-98-5] M 123.2, m 159°, pK_{Est(1)}~5.4 (NH₂), pK_{Est(2)}~10.2 (OH). Crystd from H₂O, 50% EtOH, or toluene.

6-Amino-3-hydroxytoluene [2835-99-6] M 123.2, m 162°(dec), pK_{Est(1)}~5.4 (NH₂), pK_{Est(2)}~10.4 (OH). Crystd from 50% EtOH.

4-Aminoimidazole-5-carboxamide hydrochloride (AICAR HCl) [72-40-2] M 162.6, m 255-256°(dec), pK_{Est(1)}~3.5, pK_{Est(2)}~9.4. Recrystd from EtOH.

5-Aminoindane [24425-40-9] M 133.2, m 37-38°, b 131°/15mm, 146-147°/25mm, 247-249°/745mm, pK¹⁶ 5.31. Distd and then crystd from pet ether.

6-Aminoindazole [6967-12-0] M 133.2, m 210°, pK²⁵ 3.99. Crystd from H₂O or EtOH and sublimed in a vacuum.

2-Amino-5-iodotoluene [13194-68-8] M 233.0, m 87°, pK_{Est} ~3.6. Crystd from 50% EtOH.

α-Aminoisobutyric acid [62-57-7] M 103.1, m sublimes at 280°, pK₁²⁵ 2.36, pK₂²⁵ 10.21. Crystd from aqueous EtOH and dried at 110°.

RS- β-Aminoisobutyric acid (α-methyl-β-alanine) [10569-72-9] M 103.1, m 176-178°, 178-180°, 181-182°, R (-)- isomer [144-90-1] m 183°, [α]_D²⁵ -21° (c 0.43, H₂O), pK_{Est(1)} ~ 3.7, pK_{Est(2)} ~ 10.2. Colorless prisms from hot H₂O, were powdered and dried *in vacuo*. The purity is checked by paper chromatography (Whatman 1) using ninhydrin spray to visualise the amino acid; R_F values in 95% MeOH and *n*-PrOH/5N HCOOH (8:2) are 0.36 and 0.50 respectively. [Kupiecki and Coon *Biochem Prep* 7 20 1960; Pollack *J Am Chem Soc* 65 1335 1943.] The *R*-enantiomer, isolated from iris bulbs or human urine was crystd from H₂O and sublimed *in vacuo* [Asen et al. *J Biol Chem* 234 343 1959]. The *RS*-hydrochloride was recrystd from EtOH/Et₂O m 128-129°, 130° [Böhme et al. *Chem Ber* 92 1258, 1260, 1261 1959].

5-Aminolaevulinic acid hydrochloride [5451-09-2] M 167.6, m 156-158°(dec), pK₁²² 4.05, pK₂²² 8.90. Dried in a vacuum desiccator over P₂O₅ overnight then crystd by dissolving in cold EtOH and adding dry Et₂O.

Aminomalonitrile toluene-4-sulfonate [5098-14-6] M 253.4, m 168-170°, 172°(dec), pK_{Est} ~ 1.3. Colourless crystals on recrystn from MeCN (1.8g in 100mL) using activated charcoal. Wash the crystals with dry Et₂O and dry at 25°/1mm. Recovery of ~80%. [Ferris et al. *Org Synth Coll Vol V* 32 1973.]

3-Amino-5-mercapto-1,2,4-triazole [16691-43-3] M 116.1, m 298°, pK_{Est(1)} ~ 3.0, pK_{Est(2)} ~ 9. Recrystd from H₂O and dried *in vacuo*. The *acetyl derivative* has m 325°(dec) after recrystn from H₂O. [Beilstein 26, 3rd/4th Suppl p. 1351.] Also recrystd from EtOH/H₂O (3:1, 1g in 50 mL, 50% recovery), m 300-302° dec subject to heating rate (λ_{max} 263nm, log ε 4.12), and *S-Benzyl derivative* when crystd from *C₆H₆/EtOH (20:1), or CHCl₃/Et₂O has m 109-111° [Godfrey and Kruzer *J Chem Soc* 3437 1960].

2-Amino-4-methoxy-6-methylpyrimidine [7749-47-5] M 139.2, m 157-159°, 158-158.5°, 158-160°, pK_{Est} ~ 6.0. Recrystd from H₂O. The *picrate* has m 220-221°(dec). [Baker et al. *J Am Chem Soc* 69 3072, 3075 1947; Sirakawa et al. *Yakugaku Zasshi* 73 598 1953; Backer and Grevenstuck *Recl Trav Chim, Pays-Bas* 61 291 1942.]

8-Amino-6-methoxyquinoline [90-52-8] M 174.2, m 41-42°, 51°, b 137-138°/1mm, pK^{70.1} 3.38. Distd under N₂ and high vac, then recrystd several times from MeOH (0.4mL/g). It remains colourless for several months when purified in this way [Elderfield and Rubin *J Am Chem Soc* 75 2963 1953].

1-Amino-4-methylaminoanthraquinone [1220-94-6] M 252.3, pK_{Est(1)} ~1, pK_{Est(2)} <~4. Purified by TLC on silica gel plates using toluene/acetone (3:1) as eluent. The main band was scraped off and extracted with MeOH. The solvent was evaporated and the residue dried in a drying pistol [Land, McAlpine, Sinclair and Truscott *J Chem Soc, Faraday Trans 1* 72 2091 1976].

4-Aminomethylbenzenesulfonamide hydrochloride [138-37-4] M 222.3, m 265-267°, pK₁²⁰ 8.18 (NH₂), pK₂²⁰ 10.23 (SONH₂). Crystd from dilute HCl and dried in a vacuum at 100°.

S-2-Amino-3-methyl-1-butanol (S-valinol) [2026-48-4] M 103.2, m 31-32°, b 88°/11mm, d 0.92, [α]₅₄₆ + 16.5° (c 6.32, l = 2 H₂O), [α]_D + 15.6° (EtOH), pK_{Est} ~10.4. Purified by vacuum distn using short Vigreux column. Alternatively it is purified by steam distn. The steam distillate is acidified with HCl, the aq layer is collected and evapd. The residue is dissolved in butan-1-ol, filtered and dry Et₂O added to cryst the hydrochloride salt (*hygroscopic*), m 113°. The free base can be obtained by suspending the salt in Et₂O adding small vols of satd K₂CO₃ until effervescence is complete and the mixture is distinctly alkaline. At this stage the aqueous layer should appear as a white sludge. The mixture is heated to boiling and refluxed for 30 min (more Et₂O is added if necessary). The Et₂O is decanted off from the white sludge, the

sludge is extracted twice with Et₂O (by boiling for a few minutes), the combined organic layers are dried (KOH pellets), evapd and the residue distd in a vacuum.

7-Amino-4-methylcoumarin [26093-31-2] M 175.2, m 221-442°(dec), pK_{Est} ~3.2. Dissolved in 5% HCl, filtered and basified with 2M ammonia. The ppt is dried in a vacuum, and crystd from dilute EtOH. It yields a blue soln and is light sensitive.

4-Amino-2-methyl-1-naphthol hydrochloride [130-24-5] M 209.6, m 283°(dec), pK_{Est(1)}~5.6 (NH₂), pK_{Est(2)} ~10.4 (OH). Crystd from dilute HCl.

2-Amino-2-methyl-1,3-propanediol [115-69-5] M 105.1, m 111°, b 151-152°/10mm, pK²⁵ 8.80. Crystd three times from MeOH, dried in a stream of dry N₂ at room temp, then in a vacuum oven at 55°. Stored over CaCl₂ [Hetzer and Bates *J Phys Chem* 66 308 1962].

2-Amino-2-methyl-1-propanol (β-aminoisobutanol) [124-68-5] M 89.4, m 24°, 31°, b 67°/10mm, 164-166°/760mm, d 0.935, n 1.45, pK²⁵ 9.71. Purified by distn and fractional freezing. The *hydrochloride* has m 204°-206°.

2-Amino-3-methylpyridine [1603-40-3] M 108.1, m 33.2°, b 221-222°, pK²⁵ 7.24. Crystd three times from *benzene, most of the residual *benzene being removed from the crystals over paraffin wax chips in an evacuated desiccator. The amine, transferred to a separating funnel under N₂, was left in contact with NaOH pellets for 3h with occasional shaking. It was then placed in a vacuum distilling flask where it was refluxed gently in a stream of dry N₂ before being fractionally distd [Mod, Magne and Skau *J Phys Chem* 60 1651 1956].

2-Amino-4-methylpyridine [695-34-1] M 108.1, m 99.2°, b 230°, pK²⁵ 7.48. Crystd from EtOH or a 2:1 *benzene/acetone mixture, and dried under vacuum.

2-Amino-5-methylpyridine [1603-41-4] M 108.1, m 76.5°, b 227°, pK²⁵ 7.22. Crystd from acetone.

2-Amino-6-methylpyridine [1824-81-3] M 108.1, m 44.2°, b 208-209°, pK²⁵ 7.41. Crystd three times from acetone, dried under vacuum at ca 45°. After leaving in contact with NaOH pellets for 3h, with occasional shaking, it was decanted and fractionally distd [Mod, Magne and Skau *J Phys Chem* 60 1651 1956]. Also recrystd from CH₂Cl₂ by addition of pet ether. [Marzilli et al. *J Am Chem Soc* 108 4830 1986.]

2-Amino-5-methylpyrimidine [50840-23-8] M 109.1, m 193.5°, pK_{Est} ~4.0. Crystd from water and *benzene. Sublimes at 50°/0.5mm.

4-Amino-2-methylquinoline [6628-04-2] M 158.2, m 168°, b 333°/760mm, pK²⁰ 9.42. Crystd from *benzene/pet ether.

2-Aminonaphthalene (β-naphthylamine) [91-59-8] M 143.2, m 111-113°, pK²⁵ 4.20. See entry on p. 306.

3-Amino-2-naphthoic acid [5959-52-4] M 187.2, m 214°(dec), pK_{Est(1)}~1.5 pK_{Est(2)} ~4.0. Crystd from aqueous EtOH.

4-Amino-5-naphthol-2,7-disulfonic acid [90-20-0] M 320.3, pK₁²⁵ 3.63, pK₂²⁵ 8.83. Sufficient Na₂CO₃ (ca 22g) to make the soln slightly alkaline to litmus was added to a soln of 100g of the dry acid in 750mL of hot distd water, followed by 5g of activated charcoal and 5g of Celite. The suspension was stirred for 10min and filtered by suction. The acid was pptd by adding ca 40mL of conc HCl (soln blue to Congo Red), then filtered by suction through sharkskin filter circular sheet (or hardened filter paper) and washed with 100mL of distd water. The purification process was repeated. The acid was dried overnight in an oven at 60° and stored in a dark bottle [Post and Moore *Anal Chem* 31 1872 1959].

1-Amino-2-naphthol hydrochloride [1198-27-2] M 195.7, m 250°(dec), $pK_{Est(1)} \sim 3.7$ (NH₂), $pK_{Est(2)} \sim 9.9$ (OH). Crystd from the minimum volume of hot water containing a few drops of stannous chloride in an equal weight of hydrochloric acid (to reduce atmospheric oxidation).

1-Amino-2-naphthol-4-sulfonic acid [116-63-2] M 239.3, m 295°(dec), $pK_{Est(1)} < 0$, $pK_{Est(2)} \sim 2.8$ (NH₂), $pK_{Est(2)} \sim 8.8$. Purified by warming 15g of the acid, 150g of NaHSO₃ and 5g of Na₂SO₃ (anhydrous) with 1L of water to ca 90°, shaking until most of the solid had dissolved, then filtering hot. The precipitate obtained by adding 10mL of conc HCl to the cooled filtrate was collected, washed with 95% EtOH until the washings were colourless, and dried under vacuum over CaCl₂. It was stored in a dark coloured bottle, in the cold [Chanley, Gindler and Sobotka *J Am Chem Soc* 74 4347 1952].

6-Aminonicotinic acid [3167-49-5] M 138.1, m 312°(dec), $pK_{Est(1)} \sim 2.2$ (CO₂H), $pK_{Est(2)} \sim 6.5$. Crystd from aq acetic acid.

2-Amino-4-nitrobenzoic acid [619-17-0] M 182.1, m 269°(dec), $pK_1^{25} 0.65$, $pK_2^{25} 3.70$. Crystd from water or aq EtOH.

5-Amino-2-nitrobenzoic acid [13280-60-9] M 182.1, m 235°(dec), $pK_{Est(1)} \sim 1.1$, $pK_{Est(2)} \sim 1.2$. Crystd from water.

1-Amino-4-nitronaphthalene [776-34-1] M 188.2, m 195°, $pK^{20} 0.54$. Crystd from EtOH or ethyl acetate.

2-Amino-4-nitrophenol [99-57-0] M 154.1, m 80-90° (hydrate), 142-143° (anhydr), $pK_{Est(1)} \sim 3.9$ (NH₂), $pK_{Est(2)} \sim 9.2$. Crystd from water.

2-Amino-5-nitrophenol [121-88-0] M 154.1, m 207-208°, $pK_{Est(1)} \sim 3.8$, $pK_{Est(2)} \sim 9.3$. Crystd from water.

6-Aminopenicillanic acid [551-16-6] M 216.2, m 208-209°, $[\alpha]_{546} +327^\circ$ (in 0.1M HCl), $pK_1^{25} 2.30$, $pK_2^{25} 4.90$. Crystd from water.

2-Aminoperimidine hydrobromide [40835-96-9] M 264.1, m 299°, $pK_{Est} \sim 7.9$ (free base). Purified by boiling a saturated aqueous soln with charcoal, filtering and leaving the salt to crystallise. Stored in a cool, dark place.

2-Aminophenol [95-55-6] M 109.1, m 175-176°, $pK_1^{25} 4.65$, $pK_2^{25} 9.75$. Purified by soln in hot water, decolorised with activated charcoal, filtered and cooled to induce crystn. Maintain an atmosphere of N₂ over the hot phenol soln to prevent its oxidation [Charles and Freiser *J Am Chem Soc* 74 1385 1952]. Can also be crystd from EtOH.

3-Aminophenol [591-27-5] M 109.1, m 122-123°, $pK_1^{25} 4.25$, $pK_2^{25} 9.90$. Crystd from hot water or toluene.

4-Aminophenol [123-30-8] M 109.1, m 190° (under N₂), $pK_1^{25} 5.38$, $pK_2^{25} 10.4$. Crystd from EtOH, then water, excluding oxygen. Can be sublimed at 110°/0.3mm. Has been purified by chromatography on alumina with a 1:4 (v/v) mixture of absolute EtOH/*benzene as eluent.

4-Aminophenol hydrochloride [51-78-5] M 145.6, m 306°(dec). Purified by treating an aqueous soln with saturated Na₂S₂O₃, filtering under an inert atmosphere, then recrystd from 50% EtOH twice and once from absolute EtOH [Livingston and Ke *J Am Chem Soc* 72 909 1950].

4-Aminophenylacetic acid [1197-55-3] M 151.2, m 199-200°(dec), $pK_1^{20} 3.60$, $pK_2^{20} 5.26$. Crystd from hot water (60-70mL/g).

2-Amino-1-phenylbutan-1-ol [α -(α -aminopropyl)benzyl alcohol] [(\pm)-*threo* 5897-76-7] M 165.1, m 79-80°, pK_{Est} ~9.7. Crystd from *benzene/pet ether.

S-(-)-2-Amino-3-phenyl-1-propanol (L-phenylalaninol) [3182-95-4] M 151.2, m 95°. See phenylalaninol on p. 327.

N-Aminophthalimide [1875-48-5] M 162.2, m 200-202°, pK_{Est} ~0. It has been recrystd from 96% EtOH (sol ~2% at boiling temperature) to form a yellow solution. It sublimes *in vacuo* at ca 150°. Resolidifies after melting, and remelts at 338-341°.

4-Aminopropiophenone [70-69-9] M 163.1, m 140°, pK_{Est} ~2.2. Crystd from water or EtOH.

4-(2-Aminopropyl)phenol [103-86-6] M 151.2, m 125-126°, pK_{Est(1)} ~9.4 (OH), pK_{Est(2)} ~9.7(NH₂). Crystd from *benzene.

1-Aminopyrene [1606-67-3] M 217.3, m 117-118°, pK₁²⁵ 2.91 (50% aq EtOH), pK₂²⁵ 2.77 (50% aq EtOH). Crystd from hexane.

2-Aminopyridine [504-29-0] M 94.1, m 58°, b 204-210°, pK₁²⁵ -7.6, pK₂²⁵ 6.71. Crystd from *benzene/pet ether (b 40-60°) or CHCl₃/pet ether.

3-Aminopyridine [462-08-8] M 94.1, m 64°, b 248°, pK₁²⁵ -1.5, pK₂²⁵ 6.03. Crystd from *benzene, CHCl₃/pet ether (b 60-70°), or *benzene/pet ether (4:1).

4-Aminopyridine [504-24-5] M 94.1, m 160°, b 180°/12-13mm, pK₁²⁵ -6.55, pK₂²⁵ 9.11 (9.18). Crystd from *benzene/EtOH, then recrystd twice from water, crushed and dried for 4h at 105° [Bates and Hetzer *J Res Nat Bur Stand* 64A 427 1960]. Has also been crystd from EtOH, *benzene, *benzene/pet ether, toluene and sublimes in vacuum.

2-Aminopyrimidine [109-12-6] M 95.1, m 126-127.5°, pK²⁰ 3.45. Crystd from *C₆H₆, EtOH or H₂O.

4-Aminopyrimidine [591-54-8] M 95.1, m 149-151°, 154-156°, pK²⁵ 5.69. Recryst 10.6g from hot EtOAc (200mL), 7.4g colorless needles, first crop, evap to 25mL gave 1.7g of second crop. The *Hydriodide* has m 180°. *Picrate* has m 225°. [Brown *J Soc Chem Ind* (London) 69 353 1950.]

5-Aminopyrimidine [591-55-9] M 95.1, m 171-172° (with sublimation), pK²⁵ 2.52. Purified by conversion to the MgCl₂ complex in a small vol of H₂O. The complex (~ 5g) is dissolved in the minimum vol of hot H₂O, passed through a column of activated Al₂O₃ (200g) and the column washed with EtOH. Evapn of the EtOH gives a colorless residue of the aminopyrimidine which is recrystd from *C₆H₆ (toluene could also be used) which forms needles at first then prisms. It melts with sublimation. Acetylation yields 5-*acetamidopyrimidine* which crystals from *C₆H₆, m 148-149°. [Whittaker *J Chem Soc* 1565 1951.]

Aminopyrene (4-dimethylaminoantipyrene) [58-15-1] M 231.3, m 107-109°, pK₁²⁵ -2.22, pK₂²⁵ 4.94. Crystd from pet ether.

3-Aminoquinoline [580-17-6] M 144.2, m 93.5°, pK²⁰ 4.94. Crystd from *C₆H₆.

4-Aminoquinoline [578-68-7] M 144.2, m 158°, pK₁²⁰ -7.11, pK₂²⁰ 9.13. Purified by zone refining.

5-Aminoquinoline [611-34-7] M 144.2, m 110°, b 184°/10mm, 310°/760mm, pK₁²⁰ 0.97, pK₂²⁰ 5.42. Crystd from pentane, then from *benzene or EtOH.

6-Aminoquinoline [580-15-4] M 144.2, m 117-119°, pK₁²⁰ 1.63, pK₂²⁰ 5.59. Purified by column chromatography on a SiO₂ column using CHCl₃/MeOH (4:1) as eluent. It is an irritant.

8-Aminoquinoline [578-66-5] M 144.2, m 70°, pK²⁰ 3.95. Crystd from EtOH or ligroin.

4-Aminosalicylic acid [65-49-6] M 153.1, m 150-151°(dec), pK₁²⁵ 1.78 (CO₂H), pK₂²⁵ 3.63 (NH₂), pK²⁰ 13.74 (OH). Cryst from EtOH.

5-Aminosalicylic acid (5-amino-2-hydroxybenzoic acid) [89-57-6] M 153.1, m 276-280°, 283°(dec), pK₁²⁵ 2.74 (CO₂H), pK₂²⁵ 5.84 (NH₂). Cryst as needles from H₂O containing a little NaHSO₃ to avoid aerial oxidation to the quinone-imine. The *Me ester* gives needles from *C₆H₆, m 96°, and the *hydrazide* has m 180-182° (From H₂O). [Fallab et al. *Helv Chim Acta* 34 26 1951, Shavel *J Amer Pharm Assoc* 42 402 1953.]

2-Amino-5-sulfanylthiazole [473-30-3] M 238.3, m 219-221°(dec), pK_{Est} ~4.5 (OH). Crystd from EtOH.

4-Amino-2-sulfobenzoic acid [527-76-4] M 217.1. Crystd from water.

2-Aminothiazole [96-50-4] M 108.1, m 93°, b 140°/11mm, pK²⁰ 5.36. Crystd from pet ether (b 100-120°), or EtOH.

1-Amino-1,2,4-triazole [24994-60-3] M 84.1, m 91-93°, pK_{Est} ~2. Crystd from water. [Barszez et al. *J Chem Soc, Dalton Trans* 2025 1986.]

3-Amino-1,2,4-triazole [61-82-5] M 84.1, m 159°, pK₁²⁰ 4.04, pK₂²⁰ 11.08. Crystd from EtOH (charcoal), then three times from dioxane [Williams, McEwan and Henry *J Phys Chem* 61 261 1957].

4-Amino-1,2,4-triazole [584-13-4] M 84.1, m 80-81°, pK 3.23. Crystd from water. [Barszez et al. *J Chem Soc, Dalton Trans* 2025 1986.]

7-Amino-4-(trifluoromethyl)coumarin, [53518-15-3] M 229.1, m 222°, pK_{Est} ~3.1. Purified by column chromatography on a C18 column, eluted with acetonitrile/0.01M aq HCl (1:1), and crystd from isopropanol. Alternatively, it is eluted from a silica gel column with CH₂Cl₂, or by extracting a CH₂Cl₂ solution (4g/L) with 1M aq NaOH (3 x 0.1L), followed by drying (MgSO₄), filtration and evaporation. [Bissell *J Org Chem* 45 2283 1980.]

9-Aminotryptcene [793-41-9] M 269.3, m 223.5-224.5°. Recrystd from ligroin [Imashiro et al. *J Am Chem Soc* 109 729 1987].

5-Amino-*n*-valeric acid (5-aminopentanoic acid) [660-88-8] M 117.2, m 157-158°, pK₁²⁵ 4.25, pK₂²⁵ 10.66. Crystd by dissolving in H₂O and adding EtOH.

5-Amino-*n*-valeric acid hydrochloride [627-95-2] M 153.6, m 103-104°. Crystd from CHCl₃.

Amodiaquin [4-(3-dimethylaminomethyl-4-hydroxyanilino)-7-chloroquinoline] [86-42-0] M 287.5, m 208°. Crystd from 2-ethoxyethanol.

D-Amygdalin [29883-15-6] M 457.4, m 214-216°, [α]_D²² -38° (c 1.2, H₂O). Crystd from water.

***n*-Amyl acetate** [628-63-7] M 130.2, b 149.2°, d 0.876, n 1.40228. Shaken with saturated NaHCO₃ soln until neutral, washed with water, dried with MgSO₄ and distd.

***n*-Amyl alcohol (1-pentanol)** [71-41-0] M 88.2, b 138.1°, d¹⁵ 0.818, n 1.4100. Dried with anhydrous K₂CO₃ or CaSO₄, filtered and fractionally distd. Has also been treated with 1-2% of sodium and

heated at reflux for 15h to remove water and chlorides. Traces of water can be removed from the near-dry alcohol by refluxing with a small amount of sodium in the presence of 2-3% *n*-amyl phthalate or succinate followed by distn (see *ethanol*).

Small amounts of amyl alcohol have been purified by esterifying with *p*-hydroxybenzoic acid, recrystallising the ester from CS₂, saponifying with ethanolic-KOH, drying with CaSO₄ and fractionally distilling [Olivier *Recl Trav Chim Pays-Bas* 55 1027 1936].

***tert*-Amyl alcohol** [75-85-4] M 88.2, b 102.3°, d¹⁵ 0.8135, n 1.4058. Refluxed with anhydrous K₂CO₃, CaH₂, CaO or sodium, then fractionally distd. Near-dry alcohol can be further dried by refluxing with magnesium activated with iodine, as described for *ethanol*. Further purification is possible using fractional crystn, zone refining or preparative gas chromatography.

***n*-Amylamine [1-aminopentane]** [110-58-7] M 87.2, b 105°, d 0.752, pK²⁵ 10.63. Dried by prolonged shaking with NaOH pellets, then distd.

***n*-Amyl bromide (*n*-pentylbromide)** [110-53-2] M 151.1, b 129.7°, d 1.218, n 1.445. Washed with conc H₂SO₄, then water, 10% Na₂CO₃ soln, again with water, dried with CaCl₂ or K₂CO₃, and fractionally distd just before use.

***n*-Amyl chloride** [543-59-9] M 106.6, b 107.8°, d 0.882, n 1.41177. Same as *sec*-amyl chloride.

***sec*-Amyl chloride (1-chloro-2-methylbutane)** [616-13-7] M 106.6, b 96-97°. Purified by stirring vigorously with 95% H₂SO₄, replacing the acid when it became coloured, until the layer remained colourless after 12h stirring. The amyl chloride was then washed with satd Na₂CO₃ soln, then distd water, and dried with anhydrous MgSO₄, followed by filtration, and distn through a 10-in Vigreux column. Alternatively a stream of oxygen containing 5% ozone was passed through the amyl chloride for three times as long as it took to cause the first coloration of starch iodide paper by the exit gas. Washing the liquid with NaHCO₃ soln hydrolyzed ozonides and removed organic acids prior to drying and fractional distn [Chien and Willard *J Am Chem Soc* 75 6160 1953].

***tert*-Amyl chloride** [594-36-5] M 106.6, b 86°, d 0.866. Methods of purification commonly used for other alkyl chlorides lead to decomposition. Unsatd materials were removed by chlorination with a small amount of chlorine in bright light, followed by distn [Chien and Willard *J Am Chem Soc* 75 6160 1953].

Amyl ether [693-65-2] M 158.3, b 186.8°, d 0.785, n 1.41195. Repeatedly refluxed over sodium and distd.

***p*-*tert*-Amylphenol** [80-46-6] M 146.3, m 93.5-94.2°, pK_{Est} ~10.2. Purified *via* its benzoate, as for phenol. After evaporating the solvent from its soln in ether, the material was crystd (from the melt) to constant melting point [Berliner, Berliner and Nelidow *J Am Chem Soc* 76 507 1954].

2-*n*-Amylpyridine [2294-76-0] M 149.2, b 63.0°/2mm, n²⁶ 1.4861, pK²⁵ 6.00. Dried with NaOH for several days, then distd from CaO under reduced pressure, taking the middle fraction and redistilling it.

4-*n*-Amylpyridine [2961-50-4] M 149.2, b 78.0°/2.5mm, n 1.4908, pK_{Est} ~6.1. Dried with NaOH for several days, then distd from CaO under reduced pressure, taking the middle fraction and redistilling it.

α-Amyrin [638-95-9] M 426.7, m 186°. Crystd from EtOH.

β-Amyrin [508-04-3] M 426.7, m 197-197.5°. Crystd from pet ether or EtOH.

Androstane [24887-75-0] M 260.5, m 50-50.5°. Crystd from acetone/MeOH.

***epi*-Androsterone** [481-29-8] M 290.4, m 172-173°, [α]₅₄₆ +115° (c 1, MeOH). Crystd from aq EtOH.

cis-Androsterone [53-41-8] M 290.4, m 185-185.5°. Crystd from acetone/Et₂O.

Angelic acid [565-63-9] M 100.1, m 45°, pK¹⁸ 4.29. Steam distd, then crystd from H₂O.

Aniline [62-53-3] M 93.1, f -6.0°, b 68.3/10mm, 184.4°/760mm, d 1.0220, n 1.585, n²⁵ 1.5832, pK²⁵ 4.60. Aniline is *hygroscopic*. It can be dried with KOH or CaH₂, and distd at reduced pressure. Treatment with stannous chloride removes sulfur-containing impurities, reducing the tendency to become coloured by aerial oxidn. Can be crystd from Et₂O at low temps. More extensive purifications involve preparation of derivatives, such as the double salt of aniline hydrochloride and cuprous chloride or zinc chloride, or *N*-acetylaniline (m 114°) which can be recrystd from water.

Recrystd aniline was dropped slowly into an aqueous soln of recrystd oxalic acid. Aniline oxalate was filtered off, washed several times with water and recrystd three times from 95% EtOH. Treatment with satd Na₂CO₃ soln, regenerated aniline which was distd from the soln, dried and redistd under reduced pressure [Knowles *Ind Eng Chem* 12 881 1920].

After refluxing with 10% acetone for 10h, aniline was acidified with HCl (Congo Red as indicator) and extracted with Et₂O until colourless. The hydrochloride was purified by repeated crystn before aniline was liberated by addition of alkali, then dried with solid KOH, and distd. The product was sulfur-free and remained colourless in air [Hantzsch and Freese *Chem Ber* 27 2529, 2966 1894].

Non-basic materials, including nitro compounds were removed from aniline in 40% H₂SO₄ by passing steam through the soln for 1h. Pellets of KOH were added to liberate the aniline which was steam distd, dried with KOH, distd twice from zinc dust at 20mm, dried with freshly prepared BaO, and finally distd from BaO in an all-glass apparatus [Few and Smith *J Chem Soc* 753 1949]. Aniline is absorbed by skin and is **TOXIC**

Aniline hydrobromide [542-11-0] M 174.0, m 286°. Crystd from water or EtOH and dried at 5mm over P₂O₅. Crystd four times from MeOH containing a few drops of conc HCl by addition of pet ether (b 60-70°), then dried to constant weight over paraffin chips, under vacuum [Gutbezahl and Grunwald *J Am Chem Soc* 75 559 1953]. It was pptd from EtOH soln by addition of Et₂O, and the filtered solid was recrystd from EtOH and dried *in vacuo*. [Buchanan et al. *J Am Chem Soc* 108 1537 1986.]

Aniline hydrochloride [142-04-1] M 129.6, m 200.5-201°. Same as aniline HBr above.

Aniline hydroiodide [45497-73-2] M 220.0, m dec on heating. Same as aniline HBr, store in the dark.

m-Anisaldehyde [591-31-1] M 136.2, b 143°/50mm, d 1.119. Washed with saturated aq NaHCO₃, then H₂O, dried with anhydrous MgSO₄ and distd at reduced pressure under N₂. Stored under N₂ in glass sealed ampoules.

p-Anisaldehyde (*p*-methoxybenzaldehyde) [123-11-5] M 136.2, m -1°, b 249°/atm, 89-90°/2mm, d 1.119, n 1.576. Washed with saturated aq NaHCO₃, then H₂O, steam distd, extracted distillate with Et₂O, dried (MgSO₄) and distd under vac and N₂. Store in glass ampoules under N₂ in the dark.

o-Anisidine [2-methoxyaniline] [90-04-0] M 123.2, m ~5°, b 109°/17mm, 119°/21mm, 225°/atm, d 1.096, n 1.575, pK²⁵ 4.52. It is separated from the *m*- and *p*- isomers by steam distn. It is also separated from its usual synthetic precursor *o*-nitroanisole by dissolving in dil HCl (pH <2.0) extracting the nitro impurity with Et₂O, adjusting the pH to ~8.0 with NaOH extracting the amine in Et₂O or steam distg. Extract the distillate with Et₂O, dry extract (Na₂SO₄), evaporate and fractionate the residual oil. Protect the almost colorless oil from light which turns it yellow in color. [Biggs and Robinson *J Chem Soc* 3881961; Nodzu et al. *Yakugaku Zasshi (J Pharm Soc Japan)* 71 713, 715 1951.]

m-Anisidine [3-methoxyaniline] [536-90-3] M 123.2, m ~5°, b 79°/1mm, 128°/17mm, 251°/atm, d 1.101, n 1.583, pK²⁵ 4.20. *o*-Isomer impurity can be removed by steam distn. Possible impurity is the precursor 3-nitroanisole which can be removed as for the preceding *o*-isomer and fractionated using an efficient column. Yellow liquid. [Gilman and Kyle *J Am Chem Soc* 74 3027 1952; Bryson *J Am Chem Soc* 82 4858 1960; Kadaba and Massie *J Org Chem* 22 333 1957.]

***p*-Anisidine [4-methoxyaniline]** [104-94-9] M 123.2, m 57°, pK²⁵ 5.31. Crystd from H₂O or aqueous EtOH. Dried in a vacuum oven at 35° for 6h and stored in a dry box. [More et al. *J Am Chem Soc* 108 2257 1986.] Purified by vacuum sublimation [Guarr et al. *J Am Chem Soc* 107 5104 1985].

Anisole [100-66-3] M 108.1, f -37.5°, b 43°/11mm, 153.8°/760mm, d¹⁵ 0.9988, n²⁵ 1.5143, pK⁰ -6.61 (aq H₂SO₄). Shaken with half volume of 2M NaOH, and emulsion allowed to separate. Repeated 3 times, then washed twice with water, dried over CaCl₂, filtered, dried over sodium wire and finally distd from fresh sodium under N₂, using a Dean-Stark trap, samples in the trap being rejected until free from turbidity [Caldin, Parbov, Walker and Wilson *J Chem Soc, Faraday Trans 1* 72 1856 1976].

Dried with CaSO₄ or CaCl₂, or by refluxing with sodium or BaO with crystalline FeSO₄ or by passage through an alumina column. Traces of phenols have been removed by prior shaking with 2M NaOH, followed by washing with water. Can be purified by zone refining.

2-*p*-Anisyl-1,3-indanone [117-37-3] M 252.3, m 156-157°, pK²⁰ 4.09. Crystd from acetic acid or EtOH.

Anserine [N,β-alanyl-1-methylhistidine] [584-85-0] M 240.3, m 238-239°, [α]_D +11.3° (H₂O), pK₁²⁵ 2.64, pK₂²⁵ 7.04, pK₃²⁵ 9.49. Crystd from aqueous EtOH. It is *hygroscopic*.

***S*-Anserine nitrate** [5937-77-9] M 303.3, m 225°(dec), [α]_D³⁰ +12.2°. Likely impurities: 1-methylimidazole-5-alanine, histidine. Crystd from aqueous MeOH.

Antheraxanthin [68831-78-7] M 584.8, m 205°, λ_{max} 460.5, 490.5nm, in CHCl₃. Likely impurities: violaxanthin and mutatoxanthin. Purified by chromatography on columns of Ca(OH)₂ and of ZnCO₃. Crystd from *C₆H₆/MeOH as needles or thin plates. Stored in the dark, in an inert atmosphere, at -20°.

Anthracene [120-12-7] M 178.2, m 218°, pK -7.4 (aq H₂SO₄). Likely impurities are anthraquinone, anthrone, carbazole, fluorene, 9,10-dihydroanthracene, tetracene and bianthryl. Carbazole is removed by continuous-adsorption chromatography [see Sangster and Irvine *J Phys Chem* 24 670 1956] using a neutral alumina column and passing *n*-hexane. [Sherwood in *Purification of Inorganic and Organic Materials*, Zief (ed), Marcel Dekker, New York, 1969.] The solvent is evaporated and anthracene is sublimed under vacuum, then purified by zone refining, under N₂ in darkness or non-actinic light.

Has been purified by co-distillation with ethylene glycol (boils at 197.5°), from which it can be recovered by additn of water, followed by crystn from 95% EtOH, *benzene, toluene, a mixture of *benzene/xylene (4:1), or Et₂O. It has also been chromatographed on alumina with pet ether in a dark room (to avoid photo-oxidation of adsorbed anthracene to anthraquinone). Other purification methods include sublimation in a N₂ atmosphere (in some cases after refluxing with sodium), and recrystd from toluene [Gorman et al. *J Am Chem Soc* 107 4404 1985].

Anthracene has also been crystd from EtOH, chromatographed through alumina in hot *benzene (*fume hood*) and then vac sublimed in a pyrex tube that has been cleaned and baked at 100°. (For further details see Craig and Rajikan *J Chem Soc, Faraday Trans 1* 74 292 1978; and Williams and Zboinski *J Chem Soc, Faraday Trans 1* 74 611 1978.) It has been chromatographed on alumina, recrystd from *n*-hexane and sublimed under reduced pressure. [Saltiel *J Am Chem Soc* 108 2674 1986; Masnori et al. *J Am Chem Soc* 108 1126 1986.] Alternatively, it was recrystd from cyclohexane, chromatographed on alumina with *n*-hexane as eluent, and recrystd two more times [Saltiel et al. *J Am Chem Soc* 109 1209 1987].

Anthracene-9-carboxylic acid [723-62-6] M 222.2, m 214°(dec), pK²⁰ 3.65. Crystd from EtOH.

9-Anthraldehyde [642-31-9] M 206.2, m 104-105°. Crystd from acetic acid or EtOH. [Masnori et al. *J Am Chem Soc* 108 1126 1986.]

Anthranol [529-86-2] M 196.2, m 160-170°(dec). Crystd from glacial acetic acid or aqueous EtOH.

Anthranthrone [641-13-4] M 306.3, m 300°, pK -7.9 (aq H₂SO₄). Crystd from chlorobenzene or nitrobenzene.

Anthraquinone [84-65-1] M 208.2, m 286°, pK²⁵ -8.27 (aq H₂SO₄). Crystd from CHCl₃ (38mL/g), *benzene, or boiling acetic acid, washing with a little EtOH and drying under vacuum over P₂O₅.

Anthrarufin [1,5-dihydroxy-9,10-anthraquinone] [117-12-4] M 240.1, m 280°(dec), pK₁²⁵ 9.90, pK₂²⁵ 11.05. Purified by column chromatography on silica gel with CHCl₃/Et₂O as eluent, followed by recrystn from acetone. Alternatively recrystd from glacial acetic acid [Flom and Barbara *J Phys Chem* 89 4489 1985].

1,8,9-Anthratrionol [480-22-8] M 226.2, m 176-181°, pK_{Est} ~9.5. Crystd from pet ether.

Anthrimide [1,1'-imino-bis-anthraquinone] [82-22-4] M 429.4, m >250°(dec). Crystd from chlorobenzene (red needles) or nitrobenzene (red rhombs)

Anthrone [90-44-8] M 194.2, m 155°, pK -5.5 (aq H₂SO₄). Crystd from a 3:1 mixture of *benzene/pet ether (b 60-80°) (10-12mL/g), or successively from *benzene then EtOH. Dried under vacuum.

Antipyrine [2,3-dihydro-1,5-dimethyl-3-oxo-2-phenylpyrazole] [60-80-0] M 188.2, m 114°, b 319°, pK²⁵ 1.45. Crystd from EtOH/water mixture, *benzene, *benzene/pet ether or hot water (charcoal), and dried under vacuum.

β-Apo-4'-carotenal, β-Apo-8'-carotenal, β-Apo-8'-carotenoic acid ethyl ester, β-Apo-8'-carotenoic acid methyl ester, Apocodeine, Apomorphine see entries in Chapter 6.

β-L-Arabinose (natural) [87-72-9] M 150.1, m 158°, [α]_D +104° (c 4, H₂O after 24h). Crystd slowly twice from 80% aq EtOH, then dried under vacuum over P₂O₅.

D-Arabinose [10323-20-3, 28697-53-2 (pyranoside)] M 150.1, m 164°, [α]₅₄₆ -123° (c 10, H₂O after 24h), pK²⁵ 12.54. Crystd three times from EtOH, vacuum dried at 60° for 24h and stored in a vacuum desiccator.

L-Arabitol [7643-75-6] M 152.2, m 102°, [α]₅₄₆ -16° (c 5, 8% borax soln). Crystd from 90% EtOH.

DL-Arabitol [2152-56-9] M 152.2, m 105-106°. Crystd from 90% EtOH.

Arachidic (eicosanoic C₂₀) acid [506-30-9] M 312.5, m 77°, pK_{Est} ~5.0. Crystd from abs EtOH.

Arachidic alcohol (1-eicosanol) [629-96-9] M 298.6, m 65.5° (71°), b 200°/3mm. Crystd from *benzene or *benzene/pet ether.

p-Arbutin [497-76-7] M 272.3, m 163-164°. Crystd from water.

S-Arginine [74-79-3] M 174.2, m 207°(dec), [α]_D +26.5° (c 5, in 5M HCl), [α]₅₄₆ +32° (c 5, in 5M HCl), pK₁²⁵ 2.18, pK₂²⁵ 9.36, pK₃²⁵ 11.5. Crystd from 66% EtOH.

S-Arginine hydrochloride [1119-34-2] M 210.7, m 217°(dec), [α]_D²⁰ +26.9° (c 6, M HCl). Likely impurity is ornithine. Crystd from water at pH 5-7, by adding EtOH to 80% (v/v).

S-Argininosuccinic acid [2387-71-5] M 290.3, [α]_D²⁴ +16.4° (H₂O). Likely impurity is fumaric acid. In neutral or alkaline soln it readily undergoes ring closure to the 'anhydride'. Crystd from water by adding 1.5 vols of EtOH. Barium salt is stable at 0-5° if dry. [Westfall *Biochem J* 77 135 1960.]

S-Argininosuccinic anhydride [28643-94-9] M 272.3, $[\alpha]_D^{23} -10^\circ$ (H₂O for anhydride formed at neutral pH). Crystd from water by adding two volumes of EtOH. An isomeric anhydride is formed if the free acid is allowed to stand at acid pH. In soln, the mixture of anhydrides and free acid is formed [see above entry].

L(+)-Ascorbic acid [50-81-7] M 176.1, m 193°(dec), $[\alpha]_{546} +23^\circ$ (c 10, H₂O), pK₁²⁵ 4.04, pK₂²⁵ 11.34. Crystd from MeOH/Et₂O/pet ether [Herbert et al. *J Chem Soc* 1270 1933].

S-Asparagine [70-47-3] M 150.1, m 234-235°, (monohydrate) [5794-13-8] $[\alpha]_D +32.6^\circ$ (0.1M HCl), pK₁²⁵ 1.98, pK₂²⁵ 8.84. Likely impurities are aspartic acid and tyrosine. Crystd from H₂O or aqueous EtOH. Slowly effloresces in dry air.

Aspartic acid M 133.1, m 338-339° (RS, [617-45-8]); m 271° (S, requires heating in a sealed tube [56-84-8]), $[\alpha]_D^{25} +25.4^\circ$ (3M HCl), pK₁²⁵ 1.99, pK₂²⁵ 3.90. Likely impurities are glutamic acid, cystine and asparagine. Crystd from water by adding 4 volumes of EtOH and dried at 110°.

L-Aspartic acid β-methyl ester hydrochloride [16856-13-6] M 183.6, m 194°, pK²⁵ 8.62. Recrystd from MeOH by using anhydrous diethyl ether [Bach et al. *Biochem Prep* 13 20 1971].

DL-Aspartic acid dimethyl ester hydrochloride [14358-33-9] M 197.7, 116-117°. Crystd from absolute MeOH. [Kovach et al. *J Am Chem Soc* 107 7360 1985.] Diethyl ester has pK²⁵ 6.4.

Aspergillic acid [490-02-8] M 224.3, m 97-99°, pK 5.5. Sublimed at 80°/10⁻³mm. Crystd from MeOH.

Astacin (β,β-carotene-3,3',4,4'-tetraone) [514-76-1] M 592.8, m 228°, 240-243°(evacuated tube), $\epsilon_{1\text{cm}}^{1\%} 550,000$ at 498mm (pyridine). Probable impurity is astaxanthin. Purified by chromatography on alumina/fibrous clay (1:4) or sucrose, or by partition between pet ether and MeOH (alkaline). Crystd from pyridine/water. Stored in the dark under N₂ at -20°. [Davis and Weedon *J Chem Soc* 182 1960.]

Atrolactic acid (0.5H₂O) (2-hydroxy-2-phenylpropionic acid) [515-30-0] M 166.2, m 94.5° (anhydr), 88-91° (0.5H₂O), pK¹⁸ 3.53. Crystd from water and dried at 55°/0.5mm.

Atropine [51-55-8] M 289.4, m 114-116°, pK¹⁸ 9.85. Crystd from acetone or hot water.

Auramine O (4,4'-bis-dimethylaminobenzophenone imine hydrochloride) [2465-27-2] M 321.9, pK²⁵ 10.71 (free base), 9.78 (carbinolamine). Crystd from EtOH as hydrochloride, very slightly soluble in CHCl₃, UV: λ_{max} 434 (370) nm. The free base has m 136° after crystn from *benzene. [*J Chem Soc* 1724 1949; *Biochemistry* 9 1540 1970].

Aurin tricarboxylic acid [4431-00-9] M 422.4, m 300°. The acid is dissolved in aqueous NaOH, NaHSO₃ solution is added until the colour is discharged and then the tricarboxylic acid is ppted with HCl [*Org Synth Coll Vol I* 54 1947]. Do not extract the acid with hot water because it softens forming a viscous mass. Make a solution by dissolving in aqueous NH₃. See **Aluminon** for the ammonium salt.

8-Azaadenine [1123-54-2] M 136.1, m 345°(dec), pK₁²⁰ 2.65, pK₂²⁰ 6.29. Crystd from H₂O.

2-Azacyclotridecanone (lauro lactam) [947-04-6] M 197.3, m 152°. Crystd from CHCl₃, stored over P₂O₅ in a vacuum desiccator.

8-Azaguanine [134-58-7] M 152.1, m >300°, pK₁²⁰ 1.04, pK₂²⁰ 6.29. Dissolved in hot M NH₄OH, filtered, and cooled; recrystd, and washed with water.

7-Azaindole [271-63-6] M 118.1, m 105-106°, pK²⁰ 4.57. Repeatedly recrystd from EtOH, then vacuum sublimed [Tokumura et al. *J Am Chem Soc* 109 1346 1987].

1-Azaindolizine [274-76-0] **M 118.1, b 72-73°/1mm, pK²⁰ 1.43.** Purified by distn or gas chromatography.

Azaserine [115-02-6] **M 173.1, m 146-162°(dec), [α]_D^{27.5} -0.5° (c 8.5, H₂O, pH 5.2), pK_{Est(1)}~4.53, pK_{Est(2)} ~5.40.** Crystd from 90% EtOH.

8-Azapurine (1H-1,2,3-triazolo[4,5-d]pyrimidine, 1,2,3,4,6[3H]penta-azaindene) [273-40-5] **M 121.1, m 174-175° (effervescence, m depends on heating rate), pK₁²⁰ 2.05 (equilib with covalent hydrate), pK₂²⁰ 4.84.** Sublimed at 120-130°/0.01mm and recryst from 3 parts of EtOH. [Albert *J Chem Soc(B)* 427 1966.]

Azelaic acid [123-99-9] **M 188.2, m 105-106°.** Crystd from H₂O (charcoal) or thiophene-free *benzene. The material cryst from H₂O was dried by azeotropic distn in toluene, the residual toluene soln was cooled and filtered, the ppte being dried in a vacuum oven. Also purified by zone refining or by sublimation onto a cold finger at 10⁻³ torr.

Azetidine (trimethyleneimine) [503-29-7] **M 57.1, b 61-62°, d 0.846, n 1.432, pK²⁵ 11.29.** It is a flammable, hygroscopic liquid smelling of ammonia, which absorbs CO₂ from air and should be kept under Argon. Purified by drying over solid KOH and distd through a short Vigreux column at atm pressure (under Argon) and keeping the pot temp below 210°. [Searles et al. *J Am Chem Soc* 78 4917 1956.]

Aziridine (ethyleneimine) [151-56-4] **M 43.1, b 55-56°/756mm, 56°/760mm, d²⁴ 0.8321, pK²⁵ 8.00.** Redistd in an Ar or N₂ atmosphere in a fume hood, and stored over KOH in sealed bottles in a refrigerator. Commercial aziridine has been dried over sodium and distd from the metal through an efficient column before use [Jackson and Edwards *J Am Chem Soc* 83 355 1961; Wenker *J Am Chem Soc* 57 2328 1935]. It is a weaker base than Me₂NH (pK 10.87) but is caustic to the skin. It should not be inhaled, causes inflammation of the eyes, nose and throat and one may become sensitised. It is sol in H₂O and has an ammoniacal smell and reacts with CO₂. Pure aziridine is comparatively stable but polymerises in the presence of traces of H₂O and is occasionally explosive in the presence of acids. CO₂ is sufficiently acidic to cause polymerisation (forms linear polymers) which is not free radical promoted. It is stable in the presence of bases. The violet 2:1 Cu complex crystd from EtOH containing a few drops of Aziridine and adding Et₂O has m 142°(decomp). The picrate has m 142°. [O'Rourke et al. *J Am Chem Soc* 78 2159 1956.] It has also been dried with BaO, and distd from sodium under nitrogen. **TOXIC.**

Azobenzene [103-33-3] **M 182.2, m 68°, pK²⁵ 2.48.** Ordinary azobenzene is nearly all in the *trans*-form. It is partly converted into the *cis*-form on exposure to light [for isolation see Hartley *J Chem Soc* 633 1938, and for spectra of *cis*- and *trans*-azobenzenes, see Winkel and Siebert *Chem Ber* 74B 6701941]. *trans*-Azobenzene is obtained by chromatography on alumina using 1:4 *benzene/heptane or pet ether, and crystd from EtOH (after refluxing for several hours) or hexane. All operations should be carried out in diffuse red light or in the dark.

1,1'-Azobis(cyclohexane carbonitrile) [2094-98-6] **M 244.3, m 114-114.5°, ϵ_{350nm} 16.0.** Crystd from EtOH.

α,α' -Azobis(isobutyronitrile) (AIBN) [78-61-1] **M 164.2, m 103°(dec).** Crystd from acetone, Et₂O, CHCl₃, aq EtOH or MeOH. Has also been crystd from abs EtOH below 40° in subdued light. Dried under vacuum at room temp over P₂O₅ and stored under vacuum in the dark at <-10° until used. Also crystd from CHCl₃ soln by addn of pet ether (b <40°). [Askham et al. *J Am Chem Soc* 107 7423 1985; Ennis et al. *J Chem Soc, Dalton Trans* 2485 1986; Inoue and Anson *J Phys Chem* 91 1519 1987; Tanner *J Org Chem* 52 2142 1987].

Azolitmin B [1395-18-2] **M ~3300, m >250°(dec).** Crystd from water as dark violet scales, or ppted from H₂O by addtn of EtOH as a red powder. It is an indicator which is red at pH 4.5 and blue at pH 8.3.

Azomethane [503-28-6] M 58.1, m -78° , b 1.5° . Purified by vacuum distn and stored in the dark at -80° . Can be **EXPLOSIVE**.

***p,p'*-Azoxyanisole (4,4'-dimethoxyazoxybenzene)** [1562-94-3] M 258.3, transition temps: 118.1-118.8 $^{\circ}$, 135.6-136.0 $^{\circ}$, pK^{25} -5.23 (20% aq EtOH + 80% aq H₂SO₄). Crystd from absolute or 95% EtOH, or acetone, and dried by heating under vacuum or sublimed in a vac onto a cold finger.

Azoxybenzene [495-48-7] M 198.2, m 36° , pK^{25} -6.16 (20% aq EtOH + 80% aq H₂SO₄). Crystd from EtOH or MeOH, and dried for 4h at 25° and 10^{-3} mm. Sublimed before use.

***p,p*-Azoxyphenetole** [4792-83-0] M 286.3, m 137-138 $^{\circ}$ (turbid liquid clarifies at 167 $^{\circ}$). Crystd from toluene or EtOH.

Azulene [275-51-4] M 128.2, m 98.5-99 $^{\circ}$, pK^{25} -1.65 (aq H₂SO₄). Crystd from EtOH.

Azuleno(1,2-b)thiophene [25043-00-9] M 184.2. Crystd from cyclohexane, then sublimed *in vacuo*.

Azuleno(2,1-b)thiophene [248-13-5] M 184.2. Crystd from cyclohexane, then sublimed *in vacuo*.

Azure A (3-amino-7-dimethylaminophenazin-5-ium chloride) [531-53-3] M 291.8, CI 52005, m > 290 $^{\circ}$ (dec), λ_{max} 633nm, pK 7.2. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

Azure B (3-dimethylamino-7-methylaminophenazin-5-ium chloride) [531-55-5] M 305.8, CI 52010, m > 201 $^{\circ}$ (dec), λ_{max} 648nm, pK 7.4. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

Azure C (3-amino-7-methylaminophenazin-5-ium chloride) [531-57-7] M 277.8, λ_{max} 616nm, pK 7.0. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

B.A.L. (British Anti-Lewesite) see 1,2-dimercapto-3-propanol.

Barbituric acid [6-hydroxypyrimidin-2,4-dione] [67-52-7] M 128.1, m 250 $^{\circ}$ (dec), pK_1^{25} 3.99, pK_2^{25} 12.5. Crystd twice from H₂O, then dried for 2 days at 100 $^{\circ}$.

Bathophenanthroline (4,7-diphenyl-1,10-phenanthroline) [1662-01-7] M 332.4, m 215-216 $^{\circ}$, 218-220 $^{\circ}$, pK^{25} 4.67. Best purified by recrystn from *C₆H₆ or toluene. Its solubility (per L): H₂O (1mg), M HCl (20mg), heptane (110mg), Et₂O (530mg), Me₂CO (2.3g), dioxane (3.4g), MeOH (6.0g), EtOH (10.5g), isoPrOH (10.0g), *n*-pentanol (18.7g), *C₆H₆ (12.2g), pyridine (33g), nitrobenzene (44.7g), CHCl₃ (78g) and AcOH (450.4g). [UV: *Bull Soc Chim Fr* 371 1972.] For di-Na salt 3H₂O see entry in Chapter 5.

Batyl alcohol [544-62-7] M 344.6, m 70.5-71 $^{\circ}$. Crystd from aq Me₂CO, EtOH or pet ether (b 40-60 $^{\circ}$).

Behenoyl chloride (docosanoyl chloride) [21132-76-3] M 359.0, m 40 $^{\circ}$. If the IR shows OH bands then it should be dissolved in oxalyl chloride in *C₆H₆ soln and warmed at 35 $^{\circ}$ for 24h in the absence of moisture, evaporated and distd in a vacuum of 10^{-5} mm. It is sol in *C₆H₆ and Et₂O. It is moisture sensitive and is **LACHRYMATORY**. [*J Chem Soc* 1001 1937; *J Biol Chem* 59 905 1924.]

Benzalacetone (trans-4-phenyl-3-buten-2-one) [122-57-6] M 146.2, m 42 $^{\circ}$. Crystd from pet ether (b 40-60 $^{\circ}$), or distd (b 137-142 $^{\circ}$ /16mm).

[Previous Page](#)

Azomethane [503-28-6] M 58.1, m -78° , b 1.5° . Purified by vacuum distn and stored in the dark at -80° . Can be **EXPLOSIVE**.

***p,p'*-Azoxyanisole (4,4'-dimethoxyazoxybenzene)** [1562-94-3] M 258.3, transition temps: 118.1-118.8 $^{\circ}$, 135.6-136.0 $^{\circ}$, pK^{25} -5.23 (20% aq EtOH + 80% aq H₂SO₄). Crystd from absolute or 95% EtOH, or acetone, and dried by heating under vacuum or sublimed in a vac onto a cold finger.

Azoxybenzene [495-48-7] M 198.2, m 36° , pK^{25} -6.16 (20% aq EtOH + 80% aq H₂SO₄). Crystd from EtOH or MeOH, and dried for 4h at 25° and 10^{-3} mm. Sublimed before use.

***p,p*-Azoxyphenetole** [4792-83-0] M 286.3, m 137-138 $^{\circ}$ (turbid liquid clarifies at 167 $^{\circ}$). Crystd from toluene or EtOH.

Azulene [275-51-4] M 128.2, m 98.5-99 $^{\circ}$, pK^{25} -1.65 (aq H₂SO₄). Crystd from EtOH.

Azuleno(1,2-b)thiophene [25043-00-9] M 184.2. Crystd from cyclohexane, then sublimed *in vacuo*.

Azuleno(2,1-b)thiophene [248-13-5] M 184.2. Crystd from cyclohexane, then sublimed *in vacuo*.

Azure A (3-amino-7-dimethylaminophenazin-5-ium chloride) [531-53-3] M 291.8, CI 52005, m $> 290^{\circ}$ (dec), λ_{max} 633nm, pK 7.2. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

Azure B (3-dimethylamino-7-methylaminophenazin-5-ium chloride) [531-55-5] M 305.8, CI 52010, m $> 201^{\circ}$ (dec), λ_{max} 648nm, pK 7.4. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

Azure C (3-amino-7-methylaminophenazin-5-ium chloride) [531-57-7] M 277.8, λ_{max} 616nm, pK 7.0. Twice recrystd from H₂O, and dried at 100 $^{\circ}$ /1h in an oven.

B.A.L. (British Anti-Lewesite) see 1,2-dimercapto-3-propanol.

Barbituric acid [6-hydroxypyrimidin-2,4-dione] [67-52-7] M 128.1, m 250° (dec), pK_1^{25} 3.99, pK_2^{25} 12.5. Crystd twice from H₂O, then dried for 2 days at 100 $^{\circ}$.

Bathophenanthroline (4,7-diphenyl-1,10-phenanthroline) [1662-01-7] M 332.4, m 215-216 $^{\circ}$, 218-220 $^{\circ}$, pK^{25} 4.67. Best purified by recrystn from *C₆H₆ or toluene. Its solubility (per L): H₂O (1mg), M HCl (20mg), heptane (110mg), Et₂O (530mg), Me₂CO (2.3g), dioxane (3.4g), MeOH (6.0g), EtOH (10.5g), isoPrOH (10.0g), *n*-pentanol (18.7g), *C₆H₆ (12.2g), pyridine (33g), nitrobenzene (44.7g), CHCl₃ (78g) and AcOH (450.4g). [UV: *Bull Soc Chim Fr* 371 1972.] For di-Na salt 3H₂O see entry in Chapter 5.

Batyl alcohol [544-62-7] M 344.6, m 70.5-71 $^{\circ}$. Crystd from aq Me₂CO, EtOH or pet ether (b 40-60 $^{\circ}$).

Behenoyl chloride (docosanoyl chloride) [21132-76-3] M 359.0, m 40 $^{\circ}$. If the IR shows OH bands then it should be dissolved in oxalyl chloride in *C₆H₆ soln and warmed at 35 $^{\circ}$ for 24h in the absence of moisture, evaporated and distd in a vacuum of 10^{-5} mm. It is sol in *C₆H₆ and Et₂O. It is moisture sensitive and is **LACHRYMATORY**. [*J Chem Soc* 1001 1937; *J Biol Chem* 59 905 1924.]

Benzalacetone (trans-4-phenyl-3-buten-2-one) [122-57-6] M 146.2, m 42 $^{\circ}$. Crystd from pet ether (b 40-60 $^{\circ}$), or distd (b 137-142 $^{\circ}$ /16mm).

Benzalacetophenone (Chalcone) [94-41-7] M 208.3, m 56-58°, b 208°/25mm, pK²⁵ -5.73 (aq H₂SO₄). Crystd from EtOH warmed to 50° (about 5mL/g), iso-octane, or toluene/pet ether, or recrystd from MeOH, and then twice from hexane. SKIN IRRITANT.

Benzaldehyde [100-52-7] M 106.1, f -26°, b 62° (58°)/10mm, 179.0°/760mm, d 1.044, n 1.5455, pK²⁵ -7.1 (aq H₂SO₄). To diminish its rate of oxidation, benzaldehyde usually contains additives such as hydroquinone or catechol. It can be purified *via* its bisulfite addition compound but usually distn (under nitrogen at reduced pressure) is sufficient. Prior to distn it is washed with NaOH or 10% Na₂CO₃ (until no more CO₂ is evolved), then with satd Na₂SO₃ and H₂O, followed by drying with CaSO₄, MgSO₄ or CaCl₂.

anti-Benzaldoxime [932-90-1] M 121.1, m 33-34°. Crystd from diethyl ether by adding pet ether (b 60-80°). The *syn*-isomer [622-32-2] has b 121-124°/12mm, m 34-36°.

Benzamide [55-21-0] M 121.1, m 129.5°, pK²⁵ -2.16 (aq H₂SO₄). Crystd from hot water (about 5mL/g), EtOH or 1,2-dichloroethane, and air dried. Crystd from dilute aqueous ammonia, water, acetone and then *benzene (using a Soxhlet extractor). Dried in an oven at 110° for 8h and stored in a desiccator over 99% H₂SO₄. [Bates and Hobbs *J Am Chem Soc* 73 2151 1951.]

Benzamidine [618-39-3] M 120.2, m 64-66°, pK²⁰ 11.6. Liberated from chloride by treatment with 5M NaOH. Extracted into diethyl ether. Sublimed *in vacuo*.

Benzanilide [93-98-1] M 197.2, m 164°, pK⁵⁵ 1.26. Crystd from pet ether (b 70-90°) using a Soxhlet extractor, and dried overnight at 120°. Also crystd from EtOH.

Benz[a]anthracene [56-55-3] M 228.3, m 159-160°. Crystd from MeOH, EtOH or *benzene (charcoal), then chromatographed on alumina from sodium-dried *benzene (twice), using vacuum distn to remove *benzene. Final purification was by vacuum sublimation.

Benz[a]anthracene-7,12-dione [2498-66-0] M 258.3, m 169.5-170.5°. Crystd from MeOH (charcoal).

Benzanthrone [82-05-3] M 230.3, m 170°, pK -3.2 (aq H₂SO₄). Crystd from EtOH or xylene.

***Benzene** [71-43-2] M 78.1, f 5.5°, b 80.1°, d 0.874, n 1.50110, n²⁵ 1.49790. For most purposes, *benzene can be purified sufficiently by shaking with conc H₂SO₄ until free from thiophene, then with H₂O, dilute NaOH and water, followed by drying (with P₂O₅, sodium, LiAlH₄, CaH₂, 4X Linde molecular sieve, or CaSO₄, or by passage through a column of silica gel, for a preliminary drying, CaCl₂ is suitable), and distn. A further purification step to remove thiophene, acetic acid and propionic acid, is crystn by partial freezing. The usual contaminants in dry thiophene-free *benzene are non-benzenoid hydrocarbons such as cyclohexane, methylcyclohexane, and heptanes, together with naphthenic hydrocarbons and traces of toluene. Carbonyl-containing impurities can be removed by percolation through a Celite column impregnated with 2,4-dinitrophenylhydrazine, phosphoric acid and H₂O. (Prepared by dissolving 0.5g DNPH in 6mL of 85% H₃PO₄ by grinding together, then adding and mixing 4mL of distd H₂O and 10g Celite.) [Schwartz and Parker *Anal Chem* 33 1396 1961.] *Benzene has been freed from thiophene by refluxing with 10% (w/v) of Raney nickel for 15min, after which the nickel was removed by filtration or centrifugation.

Dry *benzene was obtained by doubly distilling high purity *benzene from a soln containing the blue ketyl formed by the reaction of sodium-potassium alloy with a small amount of benzophenone.

Thiophene has been removed from *benzene (absence of bluish-green coloration when 3mL of *benzene is shaken with a soln of 10mg of isatin in 10mL of conc H₂SO₄) by refluxing the *benzene (1Kg) for several hours with 40g HgO (freshly pptd) dissolved in 40mL glacial acetic acid and 300mL of water. The ppte was filtered off, the aq phase was removed and the *benzene was washed twice with H₂O, dried and distd. Alternatively, *benzene dried with CaCl₂ has been shaken vigorously for half an hour with anhydrous AlCl₃ (12g/L) at 25-35°, then decanted, washed with 10% NaOH, and water, dried and distd. The process was repeated, giving thiophene-free *benzene. [Holmes and Beeman *Ind Eng Chem* 26 172 1934.]

After shaking successively for about an hour with conc H_2SO_4 , distd water (twice), 6M NaOH, and distd water (twice), *benzene was distd through a 3-ft glass column to remove most of the water. Abs EtOH was added and the *benzene-alcohol azeotrope was distd. (This low-boiling distn leaves any non-azeotrope-forming impurities behind.) The middle fraction was shaken with distd water to remove EtOH, and again redistd. Final slow and very careful fractional distn from sodium, then LiAlH_4 under N_2 , removed traces of water and peroxides. [Peebles, Clarke and Stockmayer *J Am Chem Soc* **82** 2780 1960.] *Benzene liquid and vapour are very **TOXIC and HIGHLY FLAMMABLE**, and all operations should be carried out in an efficient fume cupboard and in the absence of naked flames in the vicinity.

Rapid purification: To dry benzene, alumina, CaH_2 or 4A molecular sieves (3% w/v) may be used (dry for 6h). Then benzene is distd, discarding the first 5% of distillate, and stored over molecular sieves (3A, 4A) or Na wire.

[$^2\text{H}_6$]*Benzene (*benzene- d_6) [1076-43-3] M 84.2, b $80^\circ/773.6\text{mm}$, $70^\circ/562\text{mm}$, $60^\circ/399\text{mm}$, $40^\circ/186.3\text{mm}$, $20^\circ/77.1\text{mm}$, $10^\circ/49.9\text{mm}$, $0^\circ/27.5\text{mm}$, d 0.9488, d^{40} 0.9257, n 1.4991, n^{40} 1.4865. Hexadeuteriobenzene of 99.5% purity is refluxed over and distd from CaH_2 onto Linde type 5A sieves under N_2 .

Benzeneazodiphenylamine (4-phenylazodiphenylamine) [28110-26-1] M 273.3, m 82° , pK^{22} 1.52. Purified by chromatography on neutral alumina using anhydrous $^*\text{C}_6\text{H}_6$ with 1% anhydrous MeOH. The major component, which gave a stationary band, was cut out and eluted with EtOH or MeOH. [Högfeldt and Bigeleisen *J Am Chem Soc* **82** 15 1960.] Crystd from pet ether or EtOH. See Sudan I.

1-Benzeneazo-2-naphthol [842-07-9] M 248.3, m 134° , pK_{Est} ~9.5 (OH). Crystd from EtOH.

1-Benzeneazo-2-naphthylamine (Yellow AB) [85-84-7] M 247.3, m $102-104^\circ$, pK_{Est} ~4.1. Crystd from glacial acetic acid, acetic acid/water or ethanol.

1,2-Benzenedimethanol (1,2-bishydroxymethylbenzene) [612-14-6] M 138.2, m $61-64^\circ$, $63-64^\circ$, $64-65^\circ$, $65-66.5^\circ$, b $145^\circ/3\text{mm}$. Recrystd from $^*\text{C}_6\text{H}_6$, H_2O , pet ether or pentane. It has been extracted in a Soxhlet with Et_2O , evaporated and recrystd from hot pet ether. Also dissolve in Et_2O , allow to evaporate till crystals are formed, filter off and wash the colourless crystals with warm pet ether or pentane. The diacetate has m 35° , $35-36^\circ$. [*J Am Chem Soc* **69** 1197 1947, IR and UV: *J Am Chem Soc* **74** 441 1952.]

m-Benzenedisulfonic acid [98-48-6] M 238.2, pK_{Est} <0. Freed from H_2SO_4 by conversion to the calcium or barium salts (using $\text{Ca}(\text{OH})_2$ or $\text{Ba}(\text{OH})_2$, and filtering). The calcium salt was then converted to the potassium salt, using K_2CO_3 . Both the potassium and the barium salts were recrystd from H_2O , and the acid was regenerated by passing through the H^+ form of a strong cation exchange resin. The acid was recrystd twice from conductivity water and dried over CaCl_2 at 25° . [Atkinson, Yokoi and Hallada *J Am Chem Soc* **83** 1570 1961.] It has also been crystd from Et_2O and dried in a vacuum oven.

m-Benzenedisulfonyl chloride [585-47-7] M 275.1, m 63° . Crystd from CHCl_3 (EtOH free, by passing through an alumina column) and dried at 20mm pressure.

Benzene-1,2-dithiol [17534-15-5] M 142.2, m $24-25^\circ$, $27-28^\circ$, b $110-112^\circ$, $\text{pK}_{\text{Est}(1)}$ ~6.0, $\text{pK}_{\text{Est}(2)}$ ~9.4. Likely impurities are the oxidation products, the disulfides which could be polymeric. Dissolve in aq NaOH until the soln is alkaline. Extract with Et_2O and discard the extract. Acidify with cold HCl (diluted 1:1 by vol with H_2O) to Congo Red paper under N_2 and extract three times with Et_2O . Dry the Et_2O with Na_2SO_4 , filter, evaporate and distil residue under reduced press in an atmosphere of N_2 . The distillate solidifies on cooling. [UV: *J Chem Soc* 3076 1958; *J Am Chem Soc* **81** 4939 1951; *Org Synth Coll Vol V* 419 1973.]

Benzenesulfinic acid [618-41-7] M 142.2, m 84° , pK^{25} 2.16 (2.74). The acid is purified by dissolving the Na salt in H_2O , acidifying to Congo Red paper with HCl and adding a concentrated soln of FeCl_3 whereby Fe sulfinate ppts. Collect the salt, wash with a little H_2O , drain, suspend in H_2O and add a slight excess of 1.5M aq NaOH. The $\text{Fe}(\text{OH})_3$ ppts, it is fild off, the sulfinic acid in the aq soln is extracted with

Et_2O , the extract is dried (Na_2SO_4) and evapd to give colorless crystals of benzenesulfinic acid **m** 84° which are stored under N_2 in the dark, as it slowly oxidises in air to the sulfonic acid [see *Org Synth* **42** 62 1966].

Benzenesulfonic acid [98-11-3] **M** 158.2, **m** 43-44°, 50-55° (anhydrous), 65-66°, **pK**²⁵ -2.7, 0.70 (2.53?) Purified by dissolving in a small volume of distd H_2O and stirring with slightly less than the theoretical amount of BaCO_3 . When effervescence is complete and the solution is still acidic, filter off the insoluble barium benzenesulfonate. The salt is collected and dried to constant weight *in vacuo*, then suspended in H_2O and stirred with a little less than the equivalent (half mol.) of sulfuric acid. The insoluble BaSO_4 (containing a little barium benzenesulfonate) is filt'd off and the filtrate containing the free acid is evapd in a high vacuum. The oily residue will eventually crystallise when completely anhydrous. A 32% commercial acid was caused to fractionally cryst at room temp over P_2O_5 in a vac desiccator giving finally colorless deliquescent plates **m** 52.5°. The anhydrous crystn acid is deliquescent and should be stored over anhyd Na_2SO_4 in the dark and should be used in subdued sunlight as it darkens under sunlight. The main impurity is Fe which readily separates as the Fe salt in the early fractions [Taylor and Vincent *J Chem Soc* 3218 1952]. It is an IRRITANT to the skin and eyes. [see *Org Synth Coll Vol I* 84 1948; Michael and Adair *Chem Ber* **10** 585 1877.]

Benzenesulfonic anhydride [512-35-6] **M** 298.3, **m** 88-91°. Crystd from Et_2O .

Benzenesulfonyl chloride [98-09-9] **M** 176.6, **m** 14.5°, **b** 120°/10mm, 251.2°/760mm(dec), **d** 1.384. Distd, then treated with 3mole % each of toluene and AlCl_3 , and allowed to stand overnight. The free benzenesulfonyl chloride was distd off at 1mm pressure, and then carefully fractionally distd at 10mm in an all-glass column. [Jensen and Brown *J Am Chem Soc* **80** 4042 1958.]

Benzene-1,2,4,5-tetracarboxylic (pyromellitic) acid [89-05-4] **M** 254.2, **m** 281-284°, **pK**₁²⁵ 1.87, **pK**₂²⁵ 2.72, **pK**₃²⁵ 4.30, **pK**₄²⁵ 5.52. See entry on p. 345.

Benzene-1,2,3-tricarboxylic (hemimellitic) acid (H_2O) [36362-97-7] **M** 210.1, **m** 190°(dec), **pK**₁²⁵ 2.62, **pK**₂²⁵ 3.82, **pK**₃²⁵ 5.51. Crystd from water.

Benzene-1,3,5-tricarboxylic (trimesic or trimellitic) acid [554-95-0] **M** 210.1, **m** 360°(dec), **pK**₁²⁵ 2.64, **pK**₂²⁵ 3.71, **pK**₃²⁵ 5.01. Crystd from water.

1,2,4-Benzenetriol [533-73-3] **M** 126.1, **m** 141°, **pK**₁²⁰ 9.08, **pK**₂²⁰ 11.82. Crystd from Et_2O .

Benzethonium chloride [121-54-0] **M** 448.1, **m** 164-166°. Crystd from 1:9 MeOH/ Et_2O mixture.

Benzhydrol (diphenylmethanol) [91-01-0] **M** 184.2, **m** 69°, **b** 297°/748mm, 180°/20mm. Crystd from hot H_2O or pet ether (b 60-70°), pet ether containing a little *benzene, from CCl_4 , or EtOH (1mL/g). An additional purification step is passage of a *benzene soln through an activated alumina column. Sublimes in a vacuum. Also crystd three times from MeOH/ H_2O [Naguib *J Am Chem Soc* **108** 128 1986]. § A commercial polystyrene supported version is available.

Benzidine (4,4'-diaminobiphenyl) [92-87-5] **M** 184.2, **m** 128-129°, **pK**₁²⁰ 3.85, **pK**₂²⁰ 4.95. Its soln in *benzene was decolorized by percolation through two 2-cm columns of activated alumina, then concentrated until benzidine crystd on cooling. Recrystd alternatively from EtOH and *benzene to constant absorption spectrum [Carlin, Nelb and Odioso *J Am Chem Soc* **73** 1002 1951]. Has also been crystd from hot water (charcoal) and from diethyl ether. Dried under vac in an Abderhalden pistol. Stored in the dark in a stoppered container. **CARCINOGENIC**.

Benzidine dihydrochloride [531-85-1] **M** 257.2, **m** >250°(dec). Crystd by soln in hot H_2O , with addition of conc HCl to the slightly cooled soln. **CARCINOGENIC**.

Benzil [134-81-6] **M** 210.2, **m** 96-96.5°. Crystd from *benzene after washing with alkali. (Crystn from EtOH did not free benzil from material reacting with alkali.) [Hine and Howarth *J Am Chem Soc* **80** 2274

1958.] Has also been crystd from CCl_4 , diethyl ether or EtOH [Inoue et al. *J Chem Soc, Faraday Trans 1* **82** 523 1986].

Benzilic acid (diphenylglycollic acid) [76-93-7] **M 228.3, m 150°**, **pK¹⁸ 3.06**. Crystd from *benzene (ca 6mL/g), or hot H_2O .

Benzil monohydrazone [5433-88-7] **M 224.3, m 151°**. Crystd from EtOH.

α -Benzil monoxime [14090-77-8], [E, 574-15-2], [Z, 574-16-3] **M 105.1, m 140°**. Crystd from * C_6H_6 (must not use animal charcoal).

Benzimidazole [51-17-2] **M 118.1, m 172-173°**, **pK₁²⁵ 5.53, pK₂²⁵ 11.70**. Crystd from water or aqueous EtOH (charcoal), and dried at 100° for 12h.

2-Benzimidazolylacetonitrile [4414-88-4] **M 157.2, m 200-205° dec, 209.7-210.7°(corrected), 210°**. Recrystd from aqueous EtOH. It has been recrystd from hot H_2O using charcoal, and finally from aqueous EtOH. [*J Am Chem Soc* **65** 1072 1943].

Benzo[b]biphenylene [259-56-3] **M 202.2, m dec >250°**. Purified by sublimation under reduced pressure.

Benzo-15-crown-5 [14098-44-3] **M 268.3, m 78-80°**. Recrystd from *n*-heptane. **IRRITANT**.

Benzo-18-crown-6 [14098-24-9] **M 312.2, m 42-45°, 43-43.5°**. Purified by passage through a DEAE cellulose column in cyclohexane. Recryst from *n*-hexane. Its complex with thiourea has **m 127°** [5-6 mol of urea to ether, *J Org Chem* **36** 1690 1971]. The stability constants of Na^+ , K^+ , Rb^+ , Cs^+ , Tl^+ and Ba^{2+} are in *Inorg Chim Acta* **28** 73 1978] [NMR: *J Am Chem Soc* **98** 3769 1976]. **IRRITANT**.

Benzo[3,4]cyclobuta[1,2-*b*]quinoxaline [259-57-4] **M 204.2, m dec >250°**. Purified by sublimation under reduced pressure.

Benzofuran (coumarone) [271-89-6] **M 118.1, b 62-63°/15mm, 97.5-99.0°/80mm, 170-173°/atm, 173-175°(169)/760mm, d₄²⁰ 1.0945, n_D²⁰ 1.565**. Steam distil, dissolve in Et_2O , wash with 5% aqueous NaOH, saturated NaCl, dry (Na_2SO_4), evaporate and distil. UV: λ_{max} 245, 275, 282nm (log ϵ 4.08, 3.45, 3.48). The *picrate* has **m 102-103°**. [*Org Synth Coll Vol V* 251 1973; NMR: Black and Heffernan *Aust J Chem* **18** 353 1965.]

2-Benzofurancarboxylic acid [496-41-3] **M 162.1, m 192-193°**, **pK_{Est} ~3.2**. Crystd from water.

Benzofurazan [273-09-6] **M 120.1, m 55°**. Purified by crystn from EtOH and sublimed.

Benzoic acid [65-85-0] **M 122.1, m 122.6-123.1°**, **pK²⁵ 4.12**. For use as a volumetric standard, analytical reagent grade benzoic acid should be carefully fused to ca 130° (to dry it) in a platinum crucible, and then powdered in an agate mortar. Benzoic acid has been crystd from boiling water (charcoal), aq acetic acid, glacial acetic acid, * C_6H_6 , aq EtOH, pet ether (b 60-80°), and from EtOH soln by adding water. It is readily purified by fractional crystn from its melt and by sublimation in a vacuum at 80°.

***o*-Benzoic acid sulfimide (saccharin, 1,2-benzisothiazol-3(2*H*)-one 1,1-dioxide)** [81-07-2] **M 183.2, m 227-229°, 229°, 228.8-229.7°**, **pK²⁵ 1.31, pK²⁵ 12.8**. Purified by recrystn from Me_2CO [solubility 7.14% at 0°, 14.4% at 50°], or aqueous isoPrOH to give a fluorescent soln. [*Am J Pharm* **41** 17 1952.]

Benzoic anhydride [93-97-0] **M 226.2, m 42°**. Freed from benzoic acid by washing with NaHCO_3 , then water, and drying. Crystd from *benzene (0.5mL/g) by adding just enough pet ether (b 40-60°), to cause cloudiness, then cooling in ice. Can be distd at 210-220°/20mm.

(±)-**Benzoin (2-hydroxy-2-phenylacetophenone)** [119-53-9] **M 212.3, m 137°**. Crystd from CCl₄, hot EtOH (8mL/g), or 50% acetic acid. Crystd from high purity *benzene, then twice from high purity MeOH, to remove fluorescent impurities [Elliott and Radley *Anal Chem* 33 1623 1961]. Sublimes.

(±)-**α-Benzoinoxime** [441-38-3] **M 227.3, m 151°**. Crystd from diethyl ether.

Benzonitrile [100-47-0] **M 103.1, f -12.9°, b 191.1°, d 1.010, n 1.528**. Dried with CaSO₄, CaCl₂, MgSO₄ or K₂CO₃, and distd from P₂O₅ in an all-glass apparatus, under reduced pressure (**b 69°/10mm**), collecting the middle fraction. Distn from CaH₂ causes some decomposition of solvent. Isonitriles can be removed by preliminary treatment with conc HCl until the smell of isonitrile has gone, followed by preliminary drying with K₂CO₃. (This treatment also removes amines).

Steam distd (to remove small quantities of carbylamine). The distillate was extracted into ether, washed with dil Na₂CO₃, dried overnight with CaCl₂, and the ether removed by evaporation. The residue was distd at 40mm (**b 96°**) [Kice, Perham and Simons *J Am Chem Soc* 82 834 1960].

Conductivity grade benzonitrile (specific conductance 2×10^{-8} mho) was obtained by treatment with anhydrous AlCl₃, followed by rapid distn at 40-50° under vacuum. After washing with alkali and drying with CaCl₂, the distillate was vac distd several times at 35° before being fractionally crystd several times by partial freezing. It was dried over finely divided activated alumina from which it was withdrawn as required [Van Dyke and Harrison *J Am Chem Soc* 73 402 1951].

Benzo[ghi]perylene (1,12-benzoperylene) [191-24-2] **M 276.3, m 273°, 277-278.5°, 278-280°**. Purified as light green crystals by recrystn from *C₆H₆ or xylene and sublimes at 320-340° and 0.05mm [UV *Helv Chim Acta* 42 2315 1959; *Chem Ber* 65 846 1932; Fluoresc. Spectrum: *J Chem Soc* 3875 1954]. *1,3,5-Trinitrobenzene complex* **m 310-313°** (deep red crystals from *C₆H₆); *picrate* **m 267-270°** (dark red crystals from *C₆H₆); *styphnate (2,4,6-trinitroresorcinol complex)* **m 234°** (wine red crystals from *C₆H₆). It recrystallises from propan-1-ol [*J Chem Soc* 466 1959].

3,4-Benzophenanthrene [195-19-7] **M 228.3, m 68°**. Crystd from EtOH, pet ether, or EtOH/Me₂CO.

Benzophenone [119-61-9] **M 182.2, m 48.5-49°, pK -6.0 (aq H₂SO₄)**. Crystd from MeOH, EtOH, cyclohexane, *benzene or pet ether, then dried in a current of warm air and stored over BaO or P₂O₅. Also purified by zone melting and by sublimation [Itoh *J Phys Chem* 89 3949 1985; Naguib et al. *J Am Chem Soc* 108 128 1986; Gorman and Rodgers *J Am Chem Soc* 108 5074 1986; Ohamoto and Teranishi *J Am Chem Soc* 108 6378 1986; Naguib et al. *J Phys Chem* 91 3033 1987].

Benzophenone oxime [574-66-3] **M 197.2, m 142°, pK 11.18**. Crystd from MeOH (4mL/g).

Benzopinacol [464-72-2] **M 366.5, m 170-180° (depends on heating rate)**. Crystd from EtOH.

Benzo[a]pyrene (3,4 benzpyrene) [50-32-8] **M 252.3, m 177.5-178°, 179.0-179.5°**. A soln of 250mg in 100mL of *benzene was diluted with an equal volume of hexane, then passed through a column of alumina, Ca(OH)₂ and Celite (3:1:1). The adsorbed material was developed with a 2:3 *benzene/hexane mixture. (It showed as an intensely fluorescent zone.) The main zone was eluted with 3:1 acetone/EtOH, and was transferred into 1:1 *benzene-hexane by adding H₂O. The soln was washed, dried with Na₂SO₄, evaporated and crystd from *benzene by the addition of MeOH [Lijinsky and Zechmeister *J Am Chem Soc* 75 5495 1953]. Alternatively it can be chromatographed on activated alumina, eluted with a cyclohexane-*benzene mixture containing up to 8% *benzene, and the solvent evapd under reduced pressure [Cahnmann *Anal Chem* 27 1235 1955], and recrystd from EtOH [Nithipatikom and McGown *Anal Chem* 58 3145 1986]. **CARCINOGENIC.**

Benzo[e]pyrene (1,2-benzpyrene) [192-97-2] **M 252.3, m 178-179°, 178-180°**. Purified by passage through an Al₂O₃ column (Woelm, basic, activity I) and eluted with *C₆H₆ and recrystd from 2 volumes of EtOH-*C₆H₆ (4:1). Forms colourless or light yellow prisms or needles. [*J Chem Soc* 3659 1954; *Justus Liebigs Ann Chem* 705 190 1967.] *1,3,5-Trinitrobenzene complex* **m 253-254°** (orange needles from

EtOH); the *picrate* prepared by mixing 20mg in 1mL of *C_6H_6 with 20mg of picric acid in 2mL *C_6H_6 , collecting the deep red crystals, and recrystallising from *C_6H_6 m 228-229° [Synth J Chem Soc 398 1967; NMR: J Chem Phys 47 2020 1967]. **CARCINOGEN.**

3,4-Benzoquinoline (phenanthridine) [229-87-8] M 179.2, m 108-109°, b 350°, pK^{20} 4.61. Chromatographed on activated alumina from * benzene soln, with diethyl ether as eluent. Evapn of ether gave crystalline material which was freed from residual solvent under vacuum, then further purified by fractional crystn under N_2 , from its melt. Sublimes in vacuo. See also p. 324.

5,6-Benzoquinoline [85-02-9] M 179.0, m 93°, b 350°, pK^{20} 5.11. As 3,4-benzoquinoline above.

7,8-Benzoquinoline [230-27-3] M 179.0, m 52.0-52.5°, pK^{20} 4.21. As 3,4-benzoquinoline above.

p-Benzoquinone [106-51-4] M 108.1, m 115.7°. Usually purified in one or more of the following ways: steam distn, followed by filtration and drying (e.g. in a desiccator over $CaCl_2$); crystn from pet ether (b 80-100°), * benzene (with, then without, charcoal), water or 95% EtOH; sublimation under vacuum (e.g. from room temperature to liquid N_2). It slowly decomposes, and should be stored, refrigerated, in an evacuated or sealed glass vessel in the dark. It should be resublimed before use. [Wolfenden et al. J Am Chem Soc 109 463 1987.]

1-Benzosuberone (6,7,8,9-tetrahydrobenzocyclohepten-5-one) [826-73-3] M 160.2, b 80-85°/0.5mm, 90-93°/1mm, 138-139°/12mm, 154°/15mm, 175-175°/40mm, d_4^{20} 1.086, n_D^{20} 1.5638. Purified by dissolving in toluene, washing with aqueous 5% NaOH, then brine, dried ($MgSO_4$), and distd. *2,4-Dinitrophenylhydrazone* has m 210.5°, 207-208° (from $CHCl_3$ + MeOH). *Z-O-Picryloxime* has m 156-157° (from Me_2CO +MeOH); the *E-O-picryloxime* has m 107°. The *oxime* has m 106.5-107.5°. [UV J Am Chem Soc 73 1411 1951, 75 3744 1953; Chem Ber 90 1844 1957.]

1,2,3-Benzothiadiazole [273-77-8] M 136.2, m 35°, pK_{Est} \sim <0. Crystd from pet ether.

2,1,3-Benzothiadiazole [272-13-2] M 136.2, m 44°, b 206°/760mm, pK_{Est} <0. Crystd from pet ether.

1-Benzothiophene (benzo[b]thiophene, thianaphthene) [95-15-8] M 134.2, m 29-32°, 30°, 31-32°, 32°, b 100°/16mm, 103-105°/20mm, 221-222°/760mm, $d_4^{32.2}$ 1.1484, n_D^{39} 1.6306. It has the odour of naphthalene. If the IR spectrum is not very good then suspend in a faintly alkaline aqueous soln and steam distil. Extract the distillate with Et_2O , dry the extract with $CaCl_2$, filter, evaporate the solvent and fractionate the residue. Distillate sets solid. The *sulfoxide* has m 142°, the *picrate* has m 148-149° (yellow crystals from EtOH) and the *stypnate* has m 136-137°. [J Org Chem 10, 381 1945; Chem Ber 52B 1249 1919, 53 1551 1920; The Chemistry of Heterocyclic Compounds Hartough and Weisel eds, Interscience Publ, NY, p23, 28, 1954.]

1,2,3-Benzotriazole [95-14-7] M 119.1, m 96-97°, 98.5°, 100°, b 159°/0.2mm, 204°/15mm, pK_1^{20} 1.6, pK_2^{20} 8.64. Crystd from toluene, $CHCl_3$, Me_2NCHO or satd aq soln, and dried at room temperature or in a vacuum oven at 65°. Losses are less if material is distd in a vacuum. **CAUTION: may EXPLODE during vac distn, necessary precautions must be taken.** [Org Synth Coll Vol III 106 1955.]

O-Benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) [94790-37-1] M 379.2, m 200° (dec), 250°, 254°(dec). Wash with H_2O (3 x), CH_2Cl_2 (3 x), dry and recryst from MeCN. Dry in a vacuum and store cold in the dark [Dourtoglou et al. Tetrahedron Lett 1269 1978, NMR: Synthesis 572 1984].

Benzoylacetone (1-phenyl-1,3-butanedione) [93-91-4] M 162.2, m 58.5-59.0°. Crystd from Et_2O or MeOH and dried under vacuum at 40°.

2-Benzoylbenzoic acid [85-52-9] M 226.2, m 126-129°, 129.2, 130°, pK²⁵ 3.54. Recrystd from *C₆H₆ or cyclohexane, but is best recrystallised by dissolving in a small volume of hot toluene and then adding just enough pet ether to cause pptn and cool. Dry in a low vacuum at 80°. It can be sublimed at 230-240°/0.3mm [J Chem Soc 265 1957]. The *S*-benzylthiuronium salt has m 177-178° (from EtOH). [J Am Chem Soc 75 4087 1953; Chem Ber 90 1208 1957.]

3-Benzoylbenzoic acid [579-18-0] M 226.2, m 164-166°, pK_{Est}~3.5. Cryst from EtOH; vac subl.

4-Benzoylbenzoic acid [611-95-0] M 226.2, m 196.5-198°, 197-200°, pK_{Est} ~3.7. Dissolve in hot H₂O by adding enough aqueous KOH soln till distinctly alkaline, filter and then acidify with drops of conc HCl. Filter off, wash solid with cold H₂O, dry at 100°, and recrystallise from EtOH. [J Am Chem Soc 55 2540 1933.]

(S +) and (R -) 1-Benzoyl-2-tert-butyl-3-methyl-4-imidazolinone [R- 101055-57-6] [S-101055-56-5] M 260.3, m 142-143°, 145.6-146.6°, 145-147°, [α]₅₄₆²⁰ (+) and (-) 155°, [α]_D²⁰ (+) and (-) 133° (c 1, CHCl₃). Recrystd from boiling EtOH (sol 1.43g/mL) or better by dissolving in CH₂Cl₂ and adding pentane, filter and dry for at least 12h at 60°/0.1mm and sublimed at 135°/0.01mm. It has also been purified by flash column chromatography with Merck silica gel at 0.04-0.063mm and using Et₂O/pet ether/MeOH (60:35:5) as eluent. It is then recrystd from EtOH/pet ether. [IR, NMR: *Helv Chim Acta* 70 237 1987; *Angew Chem, Int Ed Engl* 25 345 1986.] The *racemate* is purified in a similar manner and has m 104-105° [NMR: *Helv Chim Acta* 68 949 1985].

Benzoyl chloride [98-88-4] M 140.6, b 56°/4mm, 196.8°/745mm, d 1.2120, n¹⁰ 1.5537.

A soln of benzoyl chloride (300mL) in *C₆H₆ (200mL) was washed with two 100mL portions of cold 5% NaHCO₃ soln, separated, dried with CaCl₂ and distd [Oakwood and Weisgerber *Org Synth* III 113 1955]. Repeated fractional distn at 4mm Hg through a glass helices-packed column (avoiding porous porcelain or silicon-carbide boiling chips, and hydrocarbon or silicon greases on the ground joints) gave benzoyl chloride that did not darken on addition of AlCl₃. Further purification was achieved by adding 3 mole% each of AlCl₃ and toluene, standing overnight, and distilling off the benzoyl chloride at 1-2mm [Brown and Jenzen *J Am Chem Soc* 80 2291 1958]. Refluxing for 2h with an equal weight of thionyl chloride before distn, has also been used. **Strong IRRITANT.** Use in a fume cupboard.

Benzoylformic acid (phenylglyoxylic acid) [611-73-4] M 150.14, m 62-65°, 64.5-65.5°, 67°, b 84°/0.1mm, 163-167°/15mm, pK²⁵ 1.39 (1.79). If the sample is oily then it may contain H₂O. In this case dry in a vacuum desiccator over P₂O₅ or KOH until crisp. For further purification dissolve 5.5g in hot CCl₄ (750mL), add charcoal (2g, this is necessary otherwise the acid may separate as an oil), filter, cool in ice-water until crystallisation is complete. Filter the acid, and the solvent on the crystals is removed by keeping the acid (4.5g) in a vacuum desiccator for 2 days. Slightly yellow crystals are obtained. It can be recrystd also from *C₆H₆/pet ether, and can be distilled in vacuum. The acid is estimated by titration with standard NaOH. The *phenylhydrazone* is recrystallised from EtOH, m 163-164°; the *semicarbazone acid* has m 259°(dec) (from EtOH). The *methyl ester* distils at 137°/14mm, 110-111°/2mm, n_D²⁰ 1.5850. [J Am Chem Soc 67 1482 1945; J Org Chem 24 1825 1959.]

Benzoyl glycine (hippuric acid) [495-69-2] M 179.2, m 188°, pK⁴⁰ 3.59. Crystd from boiling H₂O. Dried over P₂O₅.

Benzoyl isothiocyanate [532-55-8] M 163.2, m 25.5-26°, b 72.5-73°/6mm, 88-91°/20mm, 94-96°/21mm, 202.5-204°/724mm, 250-255°/atm, d₄²⁰ 1.213, n_D²⁰ 1.637. Distil over a small amount of P₂O₅, whereby the distillate crystallises in prisms. It is readily hydrolysed by H₂O to give benzamide and benzoylurea, but with NH₃ it gives *benzoylurea* m 210° which can be recrystd from EtOH. [J Am Chem Soc 62 1595 1940, 76 580 1954; *Org Synth Coll Vol* III 735 1955.]

Benzoyl peroxide [94-36-0] M 242.2, m 95°(dec). Dissolved in CHCl₃ at room temperature and ppted by adding an equal volume of MeOH or pet ether. Similarly ppted from acetone by adding two volumes of distilled water. Has also been crystd from 50% MeOH, and from diethyl ether. Dried under vacuum at room

temperature for 24h. Stored in a desiccator in the dark at 0°. When purifying in the absence of water it can be **EXPLOSIVE** and it should be done on a very small scale with adequate protection. Large amounts should be kept moist with water and stored in a refrigerator. [Kim et al. *J Org Chem* **52** 3691 1987.]

***p*-Benzoylphenol (4-hydroxybenzophenone)** [1137-42-4] **M 198.2, m 133.4-134.8°, pK²⁵ 7.95.** Dissolved in hot EtOH (charcoal), crystd once from EtOH/H₂O and twice from *benzene [Grunwald *J Am Chem Soc* **73** 4934 1951; Dryland and Sheppard *J Chem Soc Perkin Trans 1* 125 1986].

***N*-Benzoyl-*N*-phenylhydroxylamine** [304-88-1] **M 213.2, m 121-122°.** Recrystd from hot water, *benzene or acetic acid.

2-Benzoylpyridine [91-02-1] **M 183.2, m 41-43°, 48-50°, 72°/0.02mm, 104-105°/0.01, n_D²⁴ 1.6032, pK_{Est} ~2.4.** Dissolve in Et₂O, shake with aqueous NaHCO₃, H₂O, dry over MgSO₄, it solidifies on cooling. The solid can be recrystd from pet ether. Its *hydrochloride* crystallises from Me₂CO, **m 126-127°**, and the *2,4-dinitrophenylhydrazone* has **m 193-195°.** [*J Organomet Chem* **24** 623 1970.]

Benzoyl sulfide [644-32-6] **M 174.4, m 131.2-132.3°.** About 300mL of solvent was blown off from a filtered soln of benzoyl disulfide (25g) in acetone (350mL). The remaining acetone was decanted from the solid which was recrystd first from 300mL of 1:1 (v/v) EtOH/ethyl acetate, then from 300mL of EtOH, and finally from 240mL of 1:1 (v/v) EtOH/ethyl acetate. Yield about 40% [Pryor and Pickering *J Am Chem Soc* **84** 2705 1962]. *Handle in a fume cupboard because of TOXICITY and obnoxious odour.*

2,1-Benzoxathiol-3-one-1,1-dioxide (sulfobenzoic acid anhydride) [81-08-3] **M 184.2, m 116-124°, 126-127°, 128°, b 184-186°/18mm.** Purified by distn in a vacuum and readily solidifies to a crystalline mass on cooling. [*J Am Chem Soc* **34** 1594 1912.] Alternatively purified by dissolving in the minimum vol of toluene and reflux for 2h using a Dean-Stark trap. Evaporate under reduced pressure and distil the anhydride at 18mm. It can then be recrystd three times from its own weight of dry *C₆H₆. It is sensitive to moisture and should be stored in the dark in a dry atmosphere. The *O-methyl oxime* has **m 110-112°** [*Tetrahedron Lett* 3289 1972]. [*Org Synth Coll Vol I* 495 1941.] (See also p. 568 in Chapter 6.)

Benzoxazolinone [59-49-4] **M 135.1, m 137-139°, 142-143°(corrected), b 121-213°/17mm, 335-337°/760mm.** It can be purified by recrystn from aqueous Me₂CO then by distn at atm pressure then in a vacuum. The *methyl mercury salt* recryst from aq EtOH has **m 156-158°.** [*J Am Chem Soc* **67** 905 1945.]

***N*-Benzoyl-*o*-tolylhydroxylamine** [1143-74-4] **M 227.3, m 104°.** Recrystd from aqueous EtOH.

Benzyl-2-acetamido-4,6-*O*-benzylidene-2-deoxy- α -D-glucopyranoside [13343-63-0] **M 399.4, m 256-261°, 263-264°, [α]_D²⁶ +120° (c 1, pyridine).** Wash with cold isoPrOH and crystallise from dioxane/isoPrOH. [*J Org Chem* **32** 2759 1967.]

Benzyl acetate [140-11-4] **M 150.2, m -51°, b 92-93°/10mm, 134°/102mm, 214.9°/760mm, d₄²⁰ 1.0562, n_D²⁵ 1.4994.** Purified by fractional distn, preferably in a good vacuum. Values of n²⁵ of 1.5232-1.5242 seem too high and should be 1.4994. [*J Org Chem* **26** 5180 1961.]

Benzyl acetoacetate [5396-89-4] **M 192.2, b 130°/2mm, 156-157°/10mm, 162-167°/15mm, 275-277°/atm, d₄²⁰ 1.114, n_D²⁰ 1.514.** Fractionate and collect fractions of expected physical properties. Otherwise add *ca* 10% by weight of benzyl alcohol and heat in an oil bath (160-170°, open vessel) for 30min during which time excess of benzyl alcohol will have distd off, then fractionate. [*J Org Chem* **17** 77 1952.]

4'-Benzylacetophenone [782-92-3] **M 210.3, m 73°.** Crystd from EtOH (*ca* 1mL/g).

Benzyl alcohol [100-51-6] **M 107.2, f -15.3°, b 205.5°, 93°/10mm, d 0.981, n 1.54033, pK²⁵ 15.4.** Usually purified by careful fractional distn at reduced pressure in the absence of air. Benzaldehyde, if present, can be detected by UV absorption at 283nm. Also purified by shaking with aq KOH and extracting with peroxide-free diethyl ether. After washing with water, the extract was treated with satd NaHS sol, filtered,

washed and dried with CaO and distd under reduced pressure [Mathews *J Am Chem Soc* **48** 562 1926]. Peroxy compounds can be removed by shaking with a soln of Fe(II) followed by washing the alcohol layer with distd water and fractionally distd.

Benzylamine [100-46-9] M 107.2, b 178°/742mm, 185°/768mm, d 0.981, n 1.5392, pK²⁵ 9.33. Dried with NaOH or KOH, then distd under N₂, through a column packed with glass helices, taking the middle fraction. Has also been distd from zinc dust under reduced pressure.

Benzylamine hydrochloride [3287-99-8] M 143.6, m 248° (rapid heating). Crystd from water.

N-Benzylaniline (N-phenylbenzylamine) [103-32-2] M 183.4, m 36°, b 306-307°, d 1.061, pK²⁵ 4.04. Crystd from pet ether (b 60-80°) (ca 0.5mL/g).

1-Benzyl-1-aza-12-crown-4 (10-benzyl-1,4,7-trioxa-10-azacyclododecane) [84227-47-4] M 265.4, 122-125°/0.03mm, 140-143°/0.05mm, d₄²⁰ 1.09, n_D²⁰ 1.52, pK_{Est} ~ 7.7. Dissolve in CH₂Cl₂ or CCl₄ (1g in 30mL) wash with H₂O (30mL), brine (30mL), H₂O (30 mL) again, dry over MgSO₄ or Na₂SO₄ and evaporate. The residue in CH₂Cl₂ is chromatographed through Al₂O₃ (eluting with 10% EtOAc in hexane), evaporate, collect the correct fractions and distil (Kügelrohr). Log K_{Na} in dry MeOH at 25° for Na⁺ complex is 2.08. [*Tetrahedron Lett* **26** 151 1985; *J Org Chem* **53** 5652 1988.]

Benzyl bromide [100-39-0] M 171.0, m -4°, b 85°/12mm, 192°/760mm, d 1.438, n 1.575. Washed with conc H₂SO₄ (CARE), water, 10% Na₂CO₃ or NaHCO₃ soln, and again with water. Dried with CaCl₂, Na₂CO₃ or MgSO₄ and fractionally distd in the dark, under reduced pressure. It has also been thoroughly degassed at 10⁻⁶ mm and redistd in the dark. This gave material with λ_{max} (MeCN): 226nm (ε 8200) [Mohammed and Kosower *J Am Chem Soc* **93** 2709 1971]. Handle in a fume cupboard, extremely LACHRYMATORY.

Benzyl bromoacetate [5437-45-6] M 229.1, b 96-98°/0.1mm, 146°/12mm, 166-170°/22mm, d₄²⁰ 1.444, n_D²⁵ 1.5412. Dilute with Et₂O, wash with 10% aqueous NaHCO₃, H₂O, dry (MgSO₄) and fractionate using a Fenske (glass helices packing) column. [*J Chem Soc* 1521 1956.] LACHRYMATORY

N-Benzyl-tert-butylamine (N-tert-butylbenzylamine) [3378-72-1] M 163.3, b 91°/12mm, 109-110°/25mm, 218-220°/atm, d₄²⁰ 0.899, n_D²⁵ 1.4942., pK²⁵ 10.19. Dissolve in Et₂O, dry over KOH pellets, filter and fractionate in a N₂ atmosphere to avoid reaction with CO₂ from the air. The hydrochloride has m 245-246°(dec) (from MeOH + Me₂CO) and the perchlorate has m 200-201°. [*J Am Chem Soc* **80** 4320 1958.]

Benzyl carbamate [621-84-1] M 151.2, m 86°, 86-88°, 90-91°. If it smells of NH₃ then dry in a vac desiccator and recryst from 2 vols of toluene and dry in a vac desiccator again. It forms glistening plates from toluene, and can be recrystd from H₂O [*J Org Chem* **6** 878 1941; *Org Synth Coll Vol III* 168 1955].

Benzyl chloride [100-44-7] M 126.6, m 139°, b 63°/8mm, d 1.100, n 1.538. Dried with MgSO₄ or CaSO₄, or refluxed with fresh Ca turnings, then fractionally distd under reduced pressure, collecting the middle fraction and storing with CaH₂ or P₂O₅. Has also been purified by passage through a column of alumina. Alternatively it is dried over MgSO₄ and distd in a vacuum. The middle fraction is degassed by several freeze-thaw cycles and then fractionated in an 'isolated fractionating column' (which has been evacuated and sealed off at ~10⁻⁶ mm) over a steam bath. The middle fraction is retained. The final samples were vacuum distd from this sample and again retaining the middle fraction. The purity is >99.9% (no other peaks are visible on GLC and the NMR spectrum is consistent with the structure. [Mohammed and Kosower *J Am Chem Soc* **93** 1709 1971.] IRRITANT and strongly LACHRYMATORY.

N-Benzyl-β-chloropropionamide [24752-66-7] M 197.7, m 94°. Crystd from MeOH.

Benzyl cinnamate [103-41-3] M 238.3, m 34-35°, 39°, b 154-157°/0.5mm, 228-230°/22mm. Recrystd to constant melting point from 95% EtOH and has the odour of balsam. Alternatively dissolve in

Et₂O, wash with 10% aqueous Na₂CO₃, H₂O, dry (Na₂SO₄), evaporate and fractionate under reduced press using a short Vigreux column. It decomposes when boiled at atm press. [*J Am Chem Soc* **74** 547 1952; **84** 2550 1962.]

Benzyl cyanide [140-29-4] **M 117.1, b 100°/8mm, 233.5°/760mm, d 1.015, n 1.523.** Benzyl isocyanide can be removed by shaking vigorously with an equal volume of 50% H₂SO₄ at 60°, washing with satd aq NaHCO₃, then half-saturated NaCl soln, drying and fractionally distilling under reduced pressure. Distn from CaH₂ causes some decomposition of this compound: it is better to use P₂O₅. Other purification procedures include passage through a column of highly activated alumina, and distn from Raney nickel. *Precautions should be taken because of possible formation of free TOXIC cyanide; use an efficient fume cupboard.*

N-Benzyl dimethylamine [103-83-3] **M 135.2, b 66-67°/15mm, 83-84°/30mm, 98-99°/24mm, d₄²⁰ 0.898, n_D²⁰ 1.516, pK²⁵ 8.91.** Dry over KOH pellets and fractionate over Zn dust in a CO₂-free atmosphere. It has a pK_a²⁵ of 8.25 in 45% aq EtOH. Store under N₂ or in a vacuum. The *picrate* has **m 94-95°**, and the *picrolonate* has **m 151°** (from EtOH). [*Chem Ber* **63** 34 1930; *J Am Chem Soc* **55** 3001 1933; *J Chem Soc* 2845 1957.] The *tetraphenyl borate salt* has **m 182-185°**. [*Anal Chem* **28** 1794 1956.]

Benzyl dimethyloctadecylammonium chloride [122-19-0] **M 442.2, m 63°.** Crystd from acetone.

2-Benzyl-1,3-dioxolane [101-49-5] **M 164.2, b 98-99°/1mm, 110°/5mm, 137-138°/34mm, 240-242°/atm, d₄²⁰ 1.087, n_D²⁰ 1.532.** Dissolve in CH₂Cl₂, wash well with 1M NaOH, dry over K₂CO₃, filter, evaporate and distil through a short path still (Kügelrohr). It has also been purified by preparative gas chromatography. [*Synthesis* 808 1974; *J Org Chem* **34** 3949 1969.]

Benzyl ether [103-50-4] **M 198.3, b 298°, 158-160°/0.1mm, d 1.043, n 1.54057.** Refluxed over sodium, then distd under reduced pressure. Also purified by fractional freezing.

N-Benzyl-N-ethylaniline [92-59-1] **M 221.3, b 212-222°/54mm, 285-286°/710mm, 312-313°/atm (dec), d₄²⁰ 1.029, n_D²⁰ 1.595, pK_{Est} ~4.6.** Dry over KOH pellets and fractionate. The *picrate* crystallises from *C₆H₆ as yellow lemon crystals **m 126-128°** (softening at 120°). [*J Chem Soc* 303 1951; IR: *J Chem Soc* 760 1958.]

Benzyl ethyl ether [539-30-0] **M 136.2, b 186°, 65°/10mm, d 0.949, n 1.4955.** Dried with CaCl₂ or NaOH, then fractionally distd. [*J Am Chem Soc* **78** 6079 1956.]

Benzyl ethyl ketone (1-phenylbutan-2-one) [1007-32-5] **M 148.2, b 49-49.5°/0.01mm, 66-69°/1mm, 83-85°/5mm, 101-102°/10mm, 229-233°/atm, d₄²⁰ 0.989, n_D²⁵ 1.5015.** Purified by fractionation using an efficient column. It can be converted into the *oxime* and distd, **b 117-118°/2mm, 145-146°/15mm, d₂₅²⁵ 1.036, n_D²⁵ 1.5363,** decompose oxime and the ketone is redistilled. It can also be purified via the *semicarbazone* which has **m 154 155°**. [*J Am Chem Soc* **77** 5655 1955; *J Org Chem* **15** 8 1950.]

S-(+)- and R-(-)- Benzyl glycidyl ether (1-benzyloxyoxirane) [*S*:14618-80-5] [*R*:16495-13-9] **M 164.2, b 68°/10⁻⁴ mm, 105°/0.4mm, d₄²⁰ 1.072, n_D²⁰ 1.517, [α]₅₄₆²⁰ (+) and (-) 5.5°, [α]_D²⁰ (+) and (-) 5.1° (c 5, toluene), [α]_D²⁰ (+) and (-) 1.79° (c 5.02, CHCl₃), [α]_D²¹ (+) and (-) 15.3° (neat).** The ether in EtOAc is dried (Na₂SO₄) then purified by flash chromatography using pet ether/EtOAc (5:1) as eluent. The ether is then distd through a short path dist apparatus (Kugelrohr) as a colourless liquid. Alternatively, dissolve in CHCl₃, wash with H₂O, dry (Na₂SO₄), evaporate and purify through silica gel chromatography. [*J Chem Soc* 1021 1967; *Heterocycles* **16** 381 1981; *Org Synth* **69** 82 1990; *Synthesis* 539 1989; *Chem Pharm Bull Jpn* **39** 1385 1991.]

3-Benzyl-5-(2-hydroxyethyl)-4-methylthiazolinium chloride [4568-71-2] **M 269.8, m 142-144°, 145-147°.** Purified by recrystn from EtOH or H₂O. If placed in a bath at 125° and heated at 2°/min the **m** is 140.5-141.4°. [*J Biol Chem* **167** 699 1947, *J Am Chem Soc* **79** 4386 1957.]

O-Benzylhydroxylamine hydrochloride [2687-43-6] M 159.6, m 234-238°(sublimes), pK_{Est} ~5.9. Recrystd from H₂O or EtOH.

N-Benzylideneaniline [538-51-2] M 181.2, m 48° (54°), b 300°/760mm. Steam volatile and crystd from *benzene or 85% EtOH.

Benzyl isocyanate [3173-56-6] M 133.2, b 82-84°/10mm, 87°/14mm, 95°/17mm, 101-104°/33mm, d_4^{20} 1.08, n_D^{20} 1.524. Purified by fractionation through a two-plate column. It is a viscous liquid and is TOXIC. [J Chem Soc 182 1947; J Am Chem Soc 81 4838 1959; IR: *Monatsh Chem* 88 35 1957.]

Benzyl isothiocyanate [622-78-6] M 149.2, b 123-124°/1mm, 138-140°/20mm, 255-260°/atm, d_4^{20} 1.1234, n_D^{20} 1.6039. Dissolve in Et₂O, filter, if there is any solid, and distil through an efficient column at 1mm with bath temperature at ca 150°. Characterise by reacting (0.5mL) in EtOH (1mL) with 50% NH₂NH₂.H₂O (2 mL) to give 4-benzylthiosemicarbazide as colourless needles which are recrystallised from EtOH, m 130°. [J Chem Soc 1582 1950; *Justus Liebigs Ann Chem* 612 11 1958; IR and UV: *Acta Chem Scand* 13 442 1959.]

S-Benzyl-isothiuronium chloride [538-28-3] M 202.7, two forms, m 150° and 175°, pK_{Est} ~9.8 (free base). Crystd from 0.2M HCl (2mL/g) or EtOH and dried in air.

Benzylmalonic acid [616-75-1] M 194.2, m 121°, pK_1^{25} 2.91, pK_2^{25} 5.87. Crystd from *C₆H₆.

Benzylidene malononitrile [2700-22-3] M 154.2, m 83-84°. Recrystd from EtOH [Bernasconi et al. *J Am Chem Soc* 107 3612 1985].

Benzyl mercaptan [100-53-8] M 124.2, b 70.5-70.7°/9.5mm, d 1.058, n 1.5761, pK^{25} 9.43. Purified via the mercury salt [see Kern *J Am Chem Soc* 75 1865 1953], which was crystd from *benzene as needles (m 121°), and then dissolved in CHCl₃. Passage of H₂S gas regenerated the mercaptan. The HgS ppte was filtered off, and washed thoroughly with CHCl₃. The filtrate and washings were evaporated to remove CHCl₃, then residue was fractionally distd under reduced pressure [Mackle and McClean, *Trans Faraday Soc* 58 895 1962].

(-)-N-Benzyl-N-methylephedrinium bromide [benzyl(2-hydroxy-1-methyl-2-phenethyl) dimethylammonium bromide] [58648-09-2] M 350.3, m 209-211°, 212-214°, $[\alpha]_D^{25}$ -3.8° (c 1.45, MeOH), $[\alpha]_D^{20}$ -5.3° (c 1.45, MeOH). Recrystd from MeOH/Et₂O. [*Justus Liebigs Ann Chem* 710 1978.] The chloride is recrystd from EtOAc/n-hexane, m 198-199° $[\alpha]_D^{25}$ -8.67° (c 1.45, MeOH). [*J Chem Soc, Perkin Trans 1* 574 1981.]

Benzyl 4-nitrophenyl carbonate [13795-24-9] M 273.2, m 78-80°. Dissolve in Et₂O, wash with H₂O (3x) and satd aq NaCl, dry (MgSO₄), evap in vac and recryst residue from a small vol of MeOH, m 78-79°. Alternatively dissolve in Et₂O, wash with N HCl (2x), 0.5N NaHCO₃ (4x) then H₂O, dry (Na₂SO₄), evap Et₂O and recryst residue from *C₆H₆-pet ether, m 79-80°. [Khosla et al. *Indian J Chem* 5 279 1967; Wolman et al. *J Chem Soc (C)* 596 1976.]

Benzylloxyacetyl chloride [19810-31-2] M 184.6, b 81°/0.2mm, 84-87°/0.4mm, 105-107°/5mm, d_4^{20} 1.19, n_D^{20} 1.523. Check IR to see if there are OH bands. If so then it may be contaminated with free acid formed by hydrolysis. Add oxalyl chloride (amount depends on contamination and needs to be judged, ca 3mols) heat at 50° in the absence of moisture for 1h and fractionate twice, b 81°/0.2mm (with bath temp at 81°). Excessive heating results in decomposition to give benzyl chloride. The anilide is formed by adding aniline in CHCl₃ soln, m 49°. [*Helv Chim Acta* 16 1130 1933.]

Benzylloxybutan-2-one [6278-91-7] M 178.2, b 90-92°/0.1mm, 88-91°/0.5mm, 121-126°/5mm, d_4^{20} 1.0275, n_D^{20} 1.5040. Dissolve in CHCl₃, wash with H₂O, aqueous saturated NaHCO₃, H₂O, dry (MgSO₄), evaporate the CHCl₃, and fractionate. [*J Am Chem Soc* 79 2316 1957.]

Benzyloxycarbonyl chloride (Cbz-Cl, benzyl chloroformate) [501-53-1] M 170.6, b 103°/20mm, d 1.195, n 1.5190. Commercial material is better than 95% pure and may contain some toluene, benzyl alcohol, benzyl chloride and HCl. After long storage (e.g. two years at 4°, Greenstein and Winitz [*The Chemistry of the Amino Acids Vol 2* p. 890, J Wiley and Sons NY, 1961] recommended that the liquid should be flushed with a stream of dry air, filtered and stored over sodium sulfate to remove CO₂ and HCl which are formed by decomposition. It may further be distilled from an oil bath at a temperature below 85° because Thiel and Dent [*Annalen* 301 257 1898] stated that benzyloxycarbonyl chloride decarboxylates to benzyl chloride slowly at 100° and vigorously at 155°. Redistillation at higher vac below 85° yields material which shows no other peaks than those of benzyloxycarbonyl chloride by NMR spectroscopy. **LACHRYMATORY and TOXIC.**

N-Benzyloxycarbonylglycyl-L-alaninamide [17331-79-2] M 279.3, m dec >200°. Recrystd from EtOH/Et₂O.

N-Benzyloxycarbonyl-N'-methyl-L-alaninamide [33628-84-1] M 236.3, m dec >200°. Recrystd from EtOAc.

5-Benzyloxyindole [1215-59-4] M 223.3, m 96-97°; 100-103°, 104-106°, pK <0. Recrystd from *C₆H₆-pet ether or pet ether. The *picrate*, red crystals from *C₆H₆, has m 142-143°. [*Chem Ind (London)* 1035 1953; *J Am Chem Soc* 76 5579 1954; fluorescence: *Biochem J* 107 225 1968.]

p-(Benzyloxy)phenol [103-16-2] M 200.2, m 122.5°, pK_{Est} ~10.1. Crystd from EtOH or water, and dried over P₂O₅ under vacuum. [Walter et al. *J Am Chem Soc* 108 5210 1986.]

S-(-)-3-Benzyloxypropan-1,2-diol [17325-85-8] M 182.2, m 24-26°, b 117-118°/10⁻⁴mm, 115-116°/0.02mm, 121-123°/0.2mm, d₄²⁰ 1.1437, n_D²² 1.5295, [α]_D²⁵ -5.9° (neat). Purified by repeated fractional distn. [*J Biol Chem* 193 835 1951, 230 447 1958.]

2-Benzylphenol [28994-41-4] M 184.2, m 54.5°, b 312°/760mm, 175°/18mm, pK_{Est} ~10.0 Crystd from EtOH, stable form has m 52° and unstable form has m 21°.

4-Benzylphenol (α-Phenyl-p-cresol) [101-53-1] M 184.2, m 84°, pK_{Est} ~10.2. Crystd from water.

1-Benzyl-4-piperidone [3612-20-2] M 189.3, b 107-108°/0.2mm, 114-116°/0.3mm, 143-146°/5mm, 157-158°/11mm, d 1.059, n 1.538. If physical properties show contamination then dissolve in the minimum volume of H₂O, made strongly alkaline with aqueous KOH, extract with toluene several times, dry the extract with K₂CO₃, filter, evaporate and distil the residue at high vacuum using a bath temp of 160-190°, and redistil. [*J Chem Soc* 3173 1957, *J Am Chem Soc* 53 1030 1930.] The *hydrochloride* has m 159-161° (from Me₂CO + Et₂O), and the *picrate* has m 174-182° (from Me₂CO + Et₂O). [*Helv Chim Acta* 41 1184 1958.]

2-Benzylpyridine [101-82-6] M 169.2, b 98.5°/4mm, d 1.054, n²⁶ 1.5771, pK²⁵ 5.13. Dried with NaOH for several days, then distd from CaO under reduced pressure, redistilling the middle fraction.

4-Benzylpyridine [2116-65-6] M 169.2, b 110.0°/6mm, d 1.065, n²⁶ 1.5814, pK²⁵ 5.59. Dried with NaOH for several days, then distd from CaO under reduced pressure, redistilling the middle fraction.

4-N-Benzylsulfanilamide [1709-54-2] M 262.3, m 175°. Crystd from dioxane/H₂O.

Benzyl sulfide [538-74-9] M 214.3, m 50°. See dibenzylsulfide on p. 192.

Benzylthiocyanate [3012-37-1] M 149.2, m 43°, b 256°(dec). Crystd from EtOH or aqueous EtOH.

Benzyl toluene-*p*-sulfonate [1024-41-5] M 162.3, m 58°. Crystd from pet ether (b 40-60°).

Benzyltributylammonium bromide [25316-59-0] M 356.4, m 169-171°, 174-175°. Recrystd from EtOAc/EtOH and EtOH/Et₂O. [*J Am Chem Soc* 73 4122 1951, 81 3264 1959.]

Benzyl 2,2,2-trichloroacetimidate [81927-55-1] M 252.5, b 106°/0.5mm, m 3°, d 1.349, n 1.545. Purify by distn to remove up to 1% of PhCH₂OH as stabiliser. A soln in hexane can be stored for up to 2 months without decompn. It is hygroscopic and has to be stored dry. [Wessel et al. *J Chem Soc, Perkin Trans I* 2247 1985.]

Benzyltrimethylammonium chloride [56-93-9] M 185.7, m 238-239°(dec). A 60% aq soln was evapd to dryness under vac on a steam bath, and then left in a vac desiccator containing a suitable dehydrating agent. The solid residue was dissolved in a small amount of boiling absolute EtOH and pptd by adding an equal volume of diethyl ether and cooling. After washing, the ppte was dried under vac [Karusch *J Am Chem Soc* 73 1246 1951].

Benzyltrimethylammonium hydroxide (Triton B) [100-85-6] M 167.3, d 0.91. A 38% soln (as supplied) was decolorized (charcoal), then evaporated under reduced pressure to a syrup, with final drying at 75° and 1mm pressure. Prepared anhydrous by prolonged drying over P₂O₅ in a vacuum desiccator.

Berbamine [478-61-5] M 608.7, m 197-210°, [α]_D²⁰ +115° (CHCl₃), pK²⁰ 7.33. Crystd from pet ether.

Berberine [2086-83-1] M 336.4, m 145°, pK₁²⁰ 2.47, pK₂²⁰ 11.73 (pseudobase?). Crystd from pet ether or ether as yellow needles.

Berberine hydrochloride (2H₂O) [633-65-8] M 407.9, m 204-206°(dec), pK 2.47. Crystn from water gives the dihydrate. The anhydrous salt may be obtained by recrystn from EtOH/Et₂O, wash with Et₂O and dry in a vacuum. The *iodide* has m 250°(dec) (from EtOH). [*J Chem Soc* 113 503 1918; *J Chem Soc* 2036 1969.]

Betaine [107-43-7] M 117.1, m 301-305°(dec) (anhydrous), pK²⁵ 1.83. Crystd from aq EtOH.

Betamethasone (9α-fluoro-11β,17α,21-trihydroxy-16β-methylpregna-1,4-diene-3,20-dione) [378-44-9] M 392.5, m 231-136°(dec), 235-237°(dec), [α]_D²⁰ +108° (c 1, Me₂CO). Crystd from ethyl acetate, and has λ_{max} 238nm (log ε 4.18) in MeOH.

Biacetyl (butan-2,3-dione) [431-03-8] M 86.1, b 88°, d 0.981, n^{18.5} 1.3933. Dried with anhydrous CaSO₄, CaCl₂ or MgSO₄, then vacuum distd under nitrogen, taking the middle fraction and storing it at Dry-ice temperature in the dark (to prevent polymerization).

Bibenzyl [103-29-7] M 182.3, m 52.5-53.5°. Crystd from hexane, MeOH, or 95% EtOH. It has also been sublimed under vacuum, and further purified by percolation through columns of silica gel and activated alumina.

Bicuculline [485-49-4] M 367.4, m 215° (196°, 177°), [α]₅₄₆²⁰ +159° (c 1, CHCl₃), pK 4.84. See bicuculline entry on p. 515 in Chapter 6.

Bicyclohexyl [92-51-3] M 166.3, b 238° (*cis-cis*), 217-219° (*trans-trans*). Shaken repeatedly with aqueous KMnO₄ and with conc H₂SO₄, washed with water, dried, first from CaCl₂ then from sodium, and distd. [Mackenzie *J Am Chem Soc* 77 2214 1955.]

Bicyclo[3.2.1]octane [6221-55-2] M 110.2, m 141°. Purified by zone melting.

Biguanide [56-03-1] M 101.1, m 130° pK₁²⁵ 3.1, pK₂²⁵ 12.8. Crystd from EtOH.

Bilirubin [635-65-4] M 584.7, ε_{450nm} 55,600 in CHCl₃, pK_{Est} ~3.0. An acyclic tetrapyrrole bile pigment with impurities which can be eliminated by successive Soxhlet extraction with diethyl ether and MeOH. It crystallises from CHCl₃ as deep red-brown rhombs, plates or prisms, and is dried to constant weight at 80° under vacuum. [Gray et al. *J Chem Soc* 2264, 2276 1961.]

Biliverdine [114-25-0] M 582.6, m >300°, pK 3.0. The precursor of bilirubin (above) and forms dark green plates or prisms, with a violet reflection, from MeOH. [Gray et al. *J Chem Soc* 2264 1961; Sheldrick *J Chem Soc, Perkin Trans 2* 1457 1976.]

(±)-1,1'-Bi-(2-naphthol) [1,1'-di-(2-naphthol)] [602-09-5; 41024-90-2] M 286.3, m 215-217°, 218°, pK_{Est(1)}~7.1, pK_{Est(2)}~11.2. Crystd from toluene or *benzene (10mL/g). When crystd from chlorobenzene it had m 238°. Sol in dioxane is 5%.

1,1'-Bi-(2-naphthol) [1,1'-di-(2-naphthol)] [*R*(+)- 18531-94-7], [*S*(-)- 18531-99-2] M 286.3, m 207.5-208.5°, 209-211°, [α]_D²⁰ (+) and (-) 37.4.0° (c 0.5, THF), [α]₅₄₆²⁵ (+) and (-) 51° (c 0.1, THF), pK as above. Dissolve in cold 2.5N NaOH, extract with CH₂Cl₂, and acidify with 5% HCl. Collect the white ppte and recryst from aq EtOH and dry in a vacuum [*Tetrahedron* 27 5999 1971]. Optically stable in dioxane-water (100°/24h). *Racemisation*: 72% in 1.2N HCl at 100°/24h and 68% in 0.67M KOH in BuOH at 118°/23h [*J Am Chem Soc* 95 2693 1973]. Cryst from *C₆H₆ (sol 1%) using Norite or aq EtOH after chromatography through silica gel, eluting with Me₂CO-*C₆H₆. [Kyba et al. *J Org Chem* 42 4173 1977; see also Brussee and Jansen *Tetrahedron Lett* 24 3261 1983; Akimoto and Yamada *Tetrahedron* 27 5999 1971.]

1,1'-Binaphthyl [± 32507-32-7 and 604-53-5; *R*(-)- 24161-30-6; *S*(+)- 734-77-0] M 254.3, m 145°, 159° (±, 2 forms), 153-154° (+ and -), [α]_D²⁰ (-) and (+) ~220° (*C₆H₆). Purified through a silica gel column with Me₂CO-*C₆H₆ [or Al₂O₃ with 10% *C₆H₆/pet ether (b 30-60°)] and recrystd from EtOH, pentane, or slow evap of *C₆H₆, Me₂CO or Et₂O solns. Half life ~10h at 25° in various solvents. [Wilson and Pincock *J Am Chem Soc* 97 1474 1975; Akimoto and Yamada *Tetrahedron* 27 5999 1971.]

2,2'-Binaphthyl (β, β'-binaphthyl) [61-78-2] M 254.3, m 188°. Crystd from *benzene.

Biphenyl [92-52-4] M 154.2, m 70-71°, b 255°, d 0.992. Crystd from EtOH, MeOH, aq MeOH, pet ether (b 40-60°) or glacial acetic acid. Freed from polar impurities by passage through an alumina column in *benzene, followed by evapn. A in CCl₄ has been purified by vac distn and by zone refining. Treatment with maleic anhydride removed anthracene-like impurities. Recrystd from EtOH followed by repeated vacuum sublimation and passage through a zone refiner. [Taliani and Bree *J Phys Chem* 88 2351 1984.]

p-Biphenylamine [92-67-1] M 169.2, m 53°, b 191°/15mm, pK¹⁸ 4.38. See *p*-aminobiphenyl entry on p. 104.

4-Biphenylcarbonyl chloride [14002-51-8] M 216.7, m 114-115°. Dissolve in a large volume of pet ether (10 x, b 50-70°), filter through a short column of neutral alumina, evaporate to dryness *in vacuo* and recryst from pet ether (b 60-80°). **LACHRYMATORY.**

Biphenyl-2-carboxylic (2-phenylbenzoic) acid [947-84-2] M 198.2, m 114°, b 343-344°, pK 3.46. Crystd from *C₆H₆-pet ether or aq EtOH.

Biphenyl-4-carboxylic (4-phenylbenzoic) acid [92-92-2] M 198.2, m 228°, pK²⁵ 5.66 (in 50% 2-butoxyethanol). Crystd from *C₆H₆-pet ether or aq EtOH.

2,4'-Biphenyldiamine [492-17-1] M 184.2, m 45°, b 363°/760mm, pK_{Est(1)}~4.8, pK_{Est(2)}~3.9. Crystd from aqueous EtOH.

Biphenylene [259-79-0] M 152.2, m 152°. Recrystd from cyclohexane then sublimed in vacuum.

α -(4-Biphenyl)butyric acid [959-10-4] M 240.3, m 175-177°, pK_{Est} ~4.5. Crystd from MeOH.

γ -(4-Biphenyl)butyric acid [6057-60-9] M 240.3, m 118°, pK_{Est} ~4.8. Crystd from MeOH.

2,2'-Bipyridyl [366-18-7] M 156.2, m 70.5°, b 273°, pK₁²⁵ -0.52, pK₂²⁵ 4.44. Crystd from hexane, or EtOH, or (after charcoal treatment of a CHCl₃ soln) from pet ether. Also pptd from a conc soln in EtOH by addition of H₂O. Dried in a vacuum over P₂O₅. Further purification by chromatography on Al₂O₃ or by sublimation. [Airoldi et al. *J Chem Soc, Dalton Trans* 1913 1986.]

4,4'-Bipyridyl [553-26-4] M 156.2, m 73°(hydrate), 114° (171-171°)(anhydrous), b 305°/760mm, 293°/743mm, pK₁²⁰ 3.17, pK₂²⁰ 4.82. Crystd from water, *benzene/pet ether, ethyl acetate and sublimed *in vacuo* at 70°. Also purified by dissolving in 0.1M H₂SO₄ and twice pptd by addition of 1M NaOH to pH 8. Recrystd from EtOH. [Man et al. *J Chem Soc, Faraday Trans 1* 82 869 1986; Collman et al. *J Am Chem Soc* 109 4606 1987.]

2,2'-Bipyridylamine [1202-34-2] M 171.2, m 95.1°, pK_{Est} ~5.0. Crystd from Me₂CO.

2,2'-Biquinolin-4,4'-dicarboxylic (2,2'-bicinchoninic) acid [1245-13-2] M 344.3, m 367°, pK_{Est(1)} ~1.5, pK_{Est(2)} ~4.0. Dissolve in dilute NaOH and ppte with acetic acid, filter, wash well with H₂O and dry at 100° in a vacuum oven. Attempts to form a picrate failed. The *methyl ester* (SOCl₂-MeOH) has m 165.6-166°. [*J Am Chem Soc* 64 1897 1942; 68 2705 1946.] For di-K salt see entry in Chapter 5.

2,2'-Biquinolyl (α, α' -diquinolyl) [119-91-5] M 256.3, m 196°, pK_{Est} ~4.2. Decolorised in CHCl₃ soln (charcoal), then crystd to constant melting point from EtOH or pet ether [Cumper, Ginman and Vogel *J Chem Soc* 1188 1962].

Bis-acrylamide (*N,N'*-methylene bisacrylamide) [110-26-9] M 154.2, m >300°. Recrystd from MeOH (100g dissolved in 500mL boiling MeOH) and filtered without suction in a warmed funnel. Allowed to stand at room temperature and then at -15°C overnight. Crystals collected with suction in a cooled funnel and washed with cold MeOH). Crystals air-dried in a warm oven. TOXIC.

Bis-(4-aminophenyl)methane [101-77-9] M 198.3, m 92-93°, b 232°/9mm, pK_{Est} ~4.9. See 4,4'-diaminodiphenylmethane on p. 189.

2,5-Bis-(4-aminophenyl)-1,3,4-oxadiazole (BAO) [2425-95-8] M 252.3, m 252-255°, 254-255°. Recrystd from EtOH using charcoal and under N₂ to avoid oxidation.

2,5-Bis(2-benzothiazolyl)hydroquinone [33450-09-8] M 440.3, m dec >200°. Purified by repeated crystn from dimethylformamide followed by sublimation in vacuum [Erusting et al. *J Phys Chem* 91 1404 1987].

Bis-(*p*-bromophenyl)ether [53563-56-7] M 328.0, m 60.1-61.7°. Crystd twice from EtOH, once from *benzene and dried under vac [Purcell and Smith *J Am Chem Soc* 83 1063 1961].

Bis-*N*-tert-butylloxycarbonyl-L-cystine, [10389-65-8] M 440.5, m 144.5-145°, [α]_D²⁰ -133.2° (c 1, MeOH), pK_{Est} ~2.9. Crystd from in EtOAc by adding hexane [Ferraro *Biochem Prep* 13 39 1971].

2*R*,3*R*-(+)-1,4-Bis-(4-chlorobenzoyl)-2,3-butanediol [85362-86-3] and 2*S*,3*S*-(-)-1,4-Bis-(4-chlorobenzoyl)-2,3-butanediol [85362-85-2] M 371.3, m 76-77°, [α]_D²⁰ (+) and (-) 6.4° (c 3.11 CHCl₃). Recrystd from toluene-hexane. [*Tetrahedron* 40 4617 1984.]

Bis-(β -chloroethyl)amine hydrochloride [821-48-7] M 178.5, m 214-215°, pK_{Est} ~5.8 (free base). Crystd from Me₂CO.

[Previous Page](#)

Bis-(β -chloroethyl) ether [111-44-4] M 143.0, b 94°/33mm, 178.8°, d 1.220, n 1.45750. Wash with conc H₂SO₄, then Na₂CO₃ soln, dry with anhydrous Na₂CO₃, and finally pass through a 50cm column of activated alumina before distn. Alternatively, wash with 10% ferrous sulfate soln to remove peroxides, then H₂O, dry with CaSO₄, and dist in vac. Add 0.2% of catechol to stabilise it. **VERY TOXIC.**

***N,N*-Bis-(2-chloroethyl)2-naphthylamine (chlornaphthazine)** [494-03-1] M 268.3, m 54-56°, b 210°/5mm, pK_{Est} ~5.3. Crystd from pet ether. **CARCINOGENIC.**

Bis-(chloromethyl)durene [3022-16-0] M 231.2, m 197-198°. Crystd three times from *benzene, then dried under vacuum in an Abderhalden pistol.

3,3'-Bis-(chloromethyl)oxacyclobutane [78-71-7] M 155.0, m 18.9°. Shaken with aqueous NaHCO₃ or FeSO₄ to remove peroxides. Separated, dried with anhydrous Na₂SO₄, then distd under reduced pressure from a little CaH₂ [Dainton, Ivin and Walmsley *Trans Faraday Soc* 65 17884 1960].

2,2-Bis-(*p*-chlorophenyl)-1,1-dichloroethane (*p,p'*-DDD) [72-54-8] M 320.1, m 109-111°, 111-112°. Crystd from EtOH and dried in a vac. Purity checked by TLC. **TOXIC INSECTICIDE.**

2,2-Bis-(*p*-chlorophenyl)-1,1-dichloroethylene (*p,p'*-DDE) [72-55-9] M 318.0, m 89-91°. Crystd from EtOH and dried in a vac. Purity checked by TLC. **POSSIBLE CARCINOGEN.**

2,2-Bis-(4-chlorophenyl)-1,1,1-trichloroethane (*p,p'*-DDT, 1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)ethane) [50-29-3] M 354.5, m 108.5-109°, 108°. Crystd from *n*-propyl alcohol (5mL/g), then dried in air or an air oven at 50-60°. Alternatively crystd from 95% EtOH, and checked by TLC.

2,2'-Bis-[di-(carboxymethyl)-amino]diethyl ether, (HOOCCH₂)₂NCH₂CH₂OCH₂CH₂N-(CH₂COOH)₂ [923-73-9] M 336.3, pK₁²⁰ 1.8, pK₂²⁰ 2.76, pK₃²⁰ 8.84, K₄²⁰ 9.47. Crystd from EtOH.

4,4'-Bis-(diethylamino)benzophenone [90-93-7] M 324.5, m 95-96°, pK_{Est(1)} ~1.8, pK_{Est(2)} ~3.3. Crystd from EtOH (25mL/g) and dried under vacuum.

Bis-(4-dimethylaminobenzylidene)benzidine [6001-51-0] M 454.5, m 318°, pK_{Est} ~0. Crystd from nitrobenzene.

1,8-Bis-(dimethylamino)naphthalene (Proton sponge) [20734-58-1] M 214.3, m 47-48°, pK 12.34 (pK₂ -10.5 from half protonation in 86% H₂SO₄). Crystd from EtOH and dried in a vacuum oven. Stored in the dark. Also see *N,N,N',N'*-Tetramethyl-1,8-naphthalenediamine on p. 364.

Bis-(dimethylthiocarbamyl)disulfide (tetramethylthiuram disulfide, Thiram) [137-26-8] M 240.4, m 155-156°. See tetramethylthiuram disulfide on p. 365.

Bis-(4-fluoro-3-nitrophenyl) sulfone [312-30-1] M 344.3, m 193-194°. Recrystd from Me₂CO and H₂O (5:1). It should give a yellow colour in aqueous base. [*Chem Ber* 86 172 1953.]

***N,N*-Bis-(2-hydroxyethyl)-2-aminoethanesulfonic acid (BES)** [10191-18-1] M 213.3, m 150-155°, pK²⁰ 7.17. Crystd from aqueous EtOH.

Bis-(2-hydroxyethyl)amino-tris-(hydroxymethyl)methane (BIS-TRIS) [6976-37-0] M 209.2, m 89°, pK²⁰ 6.46. Crystd from hot 1-butanol. Dried in a vacuum at 25°.

***N,N*-Bis-(2-hydroxyethyl)glycine** [150-25-4] M 163.2, m 191-194°(dec). See *N,N*-di-(hydroxyethyl)glycine (BICINE) on p. 208.

3,4-Bis-(4-hydroxyphenyl)hexane [5635-50-7] M 270.4, m 187°. Freed from diethylstilboestrol by zone refining.

1,4-Bismethylaminoanthraquinone (Disperse Blue 14) [2475-44-7] M 266.3, λ_{\max} 640 (594)nm. Purified by thin-layer chromatography on silica gel plates, using toluene/acetone (3:1) as eluent. The main band was scraped off and extracted with MeOH. The solvent was evapd and the dye was dried in a drying pistol [Land, McAlpine, Sinclair and Truscott *J Chem Soc, Faraday Trans 1* 72 2091 1976].

Bis-(1-naphthylmethyl)amine [5798-49-2] M 329.4, m 62°, $pK_{\text{Est}} \sim 8.4$. Crystd from pet ether.

N,N'-Bis-(nicotinic acid) hydrazide [840-78-8] M 227-228°, m dec 200°, $pK_{\text{Est}} \sim 3.3$. Crystd from water.

Bis-(4-nitrophenyl) carbonate [5070-13-3] M 304.3, m 142-143°. Dissolve in CHCl_3 , wash with 2N NaOH (3 x) and once with conc HCl, dry (Na_2SO_4), evaporate and crystallise from toluene (authors say 15 vols of *benzene, prisms). [*Helv Chim Acta* 46 795 1963.]

Bis-(2-nitrophenyl) disulfide [1155-00-6] M 308.3, m 192-195°, 195°, 194-197°, 198-199°. Purified by recrystn from glacial AcOH or from * C_6H_6 and the yellow needles are dried in an oven at 100° until the odour of the solvent is absent. It is sparingly soluble in EtOH and Me_2CO . [Bogert and Stull *Org Synth Coll Vol I* 220 1941; Bauer and Cymerman *J Chem Soc* 3434 1949.]

Bis-(4-nitrophenyl) ether [101-63-3] M 260.2, m 142-143°. Crystd twice from * C_6H_6 , and dried under vacuum.

Bis-(4-nitrophenyl) methane [1817-74-9] M 258.2, m 183°. Crystd twice from * C_6H_6 , and dried under vacuum.

Bisnorcholanolic acid (pregnane-20-carboxylic acid) [28393-20-6] M 332.5, m 214° (α -form), 242° (β -form), 210-211° (γ -form), 184° (δ -form), 181° (ϵ -form), $pK_{\text{Est}} \sim 5.0$. Crystd from EtOH (α -form), or acetic acid (all forms).

3,3'-Bis-(phenoxymethyl)oxacyclobutane [1224-69-7] M 270.3, m 67.5-68°. Crystd from MeOH.

1,4-Bis-(2-pyridyl-2-vinyl)benzene [20218-87-5] M 284.3, $pK_{\text{Est}} \sim 5.4$. Recrystd from xylene, then chromatography (in the dark) on basic silica gel (60-80-mesh), using CH_2Cl_2 as eluent. Vacuum sublimed in the dark to a cold surface at 10^{-3} torr.

Bis-(trichloromethyl) carbonate (triphosgene) [32315-10-9] M 296.8, m 79-83°, 81-83°, b 203-206°(slight decomp). A good solid substitute for phosgene (using a third mol per mol). Cryst from pet ether and wash with anhydrous cold Et_2O , degas at 200mm then dry at 0.1mm (over H_2SO_4). It is a lachrymator, is TOXIC and should be handled with gloves and in an efficient fume hood. [Eckert and Forster *Angew Chem, Int Ed Engl* 26 894 1987; *Aldrichimica Acta* 21 47 1988.]

Bistrifluoroacetamide [407-24-9] M 209.1, m 85°, b 135-136°/744mm, 141°/760mm. Major impurity is trifluoroacetamide. Add trifluoroacetic anhydride, reflux for 2h and fractionate using a Vigreux column at atmospheric pressure. [*J Chromatogr* 78 273 1973.]

Bis-(trifluoroacetoxy)iodobenzene [2712-78-9] M 430.0, m 112-114°(dec), 120-121°, 124-126°. Cryst from warm trifluoroacetic acid and dry over NaOH pellets. Recrystd from Me_2CO /pet ether. Melting point depends on heating rate. [*Synthesis* 445 1975.]

Biuret (allophanic acid amide, carbamoylurea) [108-19-0] M 103.1, sinters at 218° and chars at 270°, $pK_1^{25} \sim -0.88$, $pK_2^{25} > 4$. Crystd from EtOH.

Bixin (6,6'-diapo- ψ,ψ - carotenedioic acid monomethyl ester) [6983-79-5] **M 394.5, m 198°**, 217°(dec), λ_{\max} (CHCl₃) 209, 475 and 443nm, $pK_{\text{Est}} \sim 4.3$. Crystd from Me₂CO (violet prisms) [Pattenden et al. *J Chem Soc (C)* 235 1970].

Blue Tetrazolium [1871-22-3] **M 727.7, m 254-255°(dec)**. Crystd from 95% EtOH/anhydrous diethyl ether, to constant absorbance at 254nm.

R-2-endo-Borneol [464-43-7] **M 154.3, m 208°** [α]_D²⁰ +15.8° (in EtOH). Crystd from boiling EtOH (charcoal).

(±)-Borneol [6627-72-1] **M 154.3, m 130°(dec)**. Crystd from pet ether (b 60-80°).

Brazilin [474-07-7] **M 269.3, m 130°(dec)**, $pK_{\text{Est}(1)} \sim 9.3$, $pK_{\text{Est}(2)} \sim 10.0$, $pK_{\text{Est}(3)} \sim 12.5$ (all phenolic). Crystd from EtOH.

Brilliant Cresyl Blue [4712-70-3] **M 332.8, pK^{25} 3.2**. Crystd from pet ether.

Brilliant Green (4-dimethylaminotriphenyl carbinol) [633-03-4] **M 482.7, m 209-211°(dec)**, pK^{25} 4.75. Purified by pptn as the perchlorate from aqueous soln (0.3%) after filtering, heating to 75° and adjustment to pH 1-2. Recrystd from EtOH/water (1:4) [Kerr and Gregory *Analyst (London)* 94 1036 1969].

N-Bromoacetamide [79-15-2] **M 138.0, m 102-105°, 107-109°, 108° (anhyd)**. Possible contaminant is CH₃CONBr₂. Recrystd from CHCl₃/hexane (1:1, seed if necessary) or water and dried over CaCl₂. [Oliveto and Gerold *Org Synth Coll Vol IV* 104 1963].

4-Bromoacetanilide [103-88-8] **M 214.1, m 167°**. Crystd from aq MeOH or EtOH. Purified by zone refining.

Bromoacetic acid [79-08-3] **M 138.9, m 50°, b 118°/15mm, 208°/760mm, pK^{25} 2.92**. Crystd from pet ether (b 40-60°). Diethyl ether soln passed through an alumina column, and the ether evaporated at room temperature under vacuum. **LACHRYMATORY**.

Bromoacetone [598-31-2] **M 137.0, b 31.5°/8mm**. Stood with anhydrous CaCO₃, distd under low vacuum, and stored with CaCO₃ in the dark at 0°. **LACHRYMATORY**.

4-Bromoacetophenone [99-90-1] **M 199.1, m 54°**. Crystd from EtOH, MeOH or from pet ether (b 80-100°). [Tanner *J Org Chem* 52 2142 1987.]

ω -Bromoacetophenone (phenacyl bromide) [70-11-1] **M 199.1, m 57-58°**. Crystd from EtOH, MeOH or from pet ether (b 80-100°). [Tanner *J Org Chem* 52 2142 1987.]

4-Bromoaniline [106-40-1] **M 172.0, m 66°, pK^{25} 3.86**. Crystd (with appreciable loss) from aqueous EtOH.

2-Bromoanisole [578-57-4] **M 187.0, f 2.5°, b 124°/40mm, d 1.513, n^{25} 1.5717**. Crystd by partial freezing (repeatedly), then distd under reduced pressure.

4-Bromoanisole [104-92-7] **M 187.0, f 13.4°, b 124°/40mm, d 1.495, n^{25} 1.5617**. Crystd by partial freezing (repeatedly), then distd under reduced pressure.

9-Bromoanthracene [1564-64-3] **M 257.1, m 98-100°**. Crystd from MeOH or EtOH followed by sublimation *in vacuo*. [Masnori et al. *J Am Chem Soc* 108 126 1986.]

4-Bromobenzal diacetate [55605-27-1] **M 287.1, m 95°**. Crystd from hot EtOH (3mL/g).

Bromobenzene [108-86-1] M 157.0, b 155.9°, d 1.495, n 1.5588, n¹⁵ 1.56252. Washed vigorously with conc H₂SO₄, then 10% NaOH or NaHCO₃ solns, and H₂O. Dried with CaCl₂ or Na₂SO₄, or passed through activated alumina, before refluxing with, and distilling from, CaH₂, using a glass helix-packed column.

4-Bromobenzene diazonium tetrafluoroborate [673-40-5] M 270.8, m 133° (dec), 135-140° (dec), 135° (dec). Wash with Et₂O until the wash is colourless and allow to dry by blowing N₂ over it. Store at 0-4° in the dark. [Chem Ber 64 1340 1931.]

4-Bromobenzenesulfonyl chloride [98-58-8] M 255.5, m 73-75°, 74.3-75.1, 75-76°, 77°, b 153°/15mm, 150.6°/13mm. Wash with cold water, dry and recryst from pet ether, or from ethyl ether cooled in powdered Dry-ice after the ether soln had been washed with 10% NaOH until colourless, then dried with anhydrous Na₂SO₄. Alternatively dissolve in CHCl₃, wash with H₂O, dry (Na₂SO₄), evaporate and crystallise. [J Am Chem Soc 62 511 1940.] Test for the SO₂Cl group by dissolving in EtOH and boiling with NH₄CNS whereby a yellow amorphous ppte forms on cooling [J Am Chem Soc 25 198 1901].

***o*-Bromobenzoic acid** [88-65-3] M 201.0, m 148.9°, pK²⁰ 2.88. Crystd from *C₆H₆ or MeOH.

***m*-Bromobenzoic acid** [585-76-2] M 201.0, m 155°, pK²⁵ 3.81. Crystd from acetone/water, MeOH or acetic acid.

***p*-Bromobenzoic acid** [586-76-5] M 201.0, m 251-252°, 254-256°, 257-258°, pK²⁵ 3.96. Crystd from MeOH, or MeOH/water mixture, 90% EtOH and Et₂O. The *methyl ester* has m 81° from Et₂O or dilute MeOH. [Male and Thorp J Am Chem Soc 35 269 1913; Lamneck J Am Chem Soc 76 406 1954, Vandenbelt et al. Anal Chem 26 926 1954.]

***p*-Bromobenzophenone** [90-90-4] M 261.1, m 81°. Crystd from EtOH.

***p*-Bromobenzoyl chloride** [586-75-4] M 219.5, m 36-39°, 39.8°, 41°, b 62°/0.1mm, 104.5°/6mm, 126.4-127.2°/14mm. Check IR of a film to see if OH bands are present. If absent then recryst from pet ether and dry in a vacuum. If OH bands are weak then distil *in vacuo* and recryst if necessary. If OH bands are very strong then treat with an equal volume of redistilled SOCl₂ reflux for 2h then evaporate excess of SOCl₂ and distil residual oil or low melting solid. Store in the dark away from moisture. LACHRYMATORY. [Martin and Partington J Chem Soc 1175 1936.]

***p*-Bromobenzyl bromide** [589-15-1] M 249.9, m 60-61°. Crystd from EtOH. LACHRYMATORY.

***p*-Bromobenzyl chloride** [589-17-3] M 205.5, m 40-41°, b 105-115°/12mm. Crystd from EtOH. LACHRYMATORY.

***p*-Bromobiphenyl** [92-66-0] M 233.1, m 88.8-89.2°. Crystd from abs EtOH and dried under vacuum.

2-Bromobutane [78-76-2] M 137.0, b 91.2°, d 1.255, n 1.4367, n²⁵ 1.4341. Washed with conc HCl, water, 10% aqueous NaHSO₃, and then water. Dried with CaCl₂, Na₂SO₄ or anhydrous K₂CO₃, and fractionally distd through a 1m column packed with glass helices.

(+)-3-Bromocamphor-8-sulfonic acid [5344-58-1] M 311.2, m 195-196°(anhydrous), [α]_D²⁰ +88.3° (in H₂O), pK ~0. Crystd from water.

IR(endo,anti)-3-Bromocamphor-8-sulfonic acid ammonium salt see entry in Chapter 5.

(+)-3-Bromocamphor-10-sulfonic acid hydrate [67999-30-8] M 329.2, m 119-121°, [α]_D²⁰ +98.3° (in H₂O), pK ~0. Crystd from water.

4-Bromo-4'-chlorobenzophenone [27428-57-5] **M 295.6, m 150°**. Crystd from EtOH or *C₆H₆ and further purified by zone refining (100 passes) [Grove and Turner *J Chem Soc* 509 1929; Lin and Hanson *J Phys Chem* 91 2279 1987].

Bromocresol Green (3',3'',5',5''-tetrabromo-*m*-cresolsulfonephthalein) [76-60-8] **M 698.0, m 218-219°(dec), 225°(dec), pK 4.51**. Crystd from glacial acetic acid or dissolved in aqueous 5% NaHCO₃ soln and ppted from hot soln by dropwise addition of aqueous HCl. Repeated until the extinction did not increase (λ_{\max} 423nm). Indicator at pH 3.81 (yellow) and pH 5.4 (blue-green).

Bromocresol Purple (5',5''-dibromo-*o*-cresolsulfonephthalein) [115-40-2] **M 540.2, m 241-242°(dec), pK₁ -2.15, pK₂ 6.3**. Dissolved in aqueous 5% NaHCO₃ soln and ppted from hot soln by dropwise addition of aqueous HCl. Repeated until the extinction did not increase (λ_{\max} 419nm). Can also be crystd from *benzene. Indicator at pH 5.2 (yellow) and pH 6.8 (purple).

5-Bromocytosine [2240-25-7] **M 190.0, m 245-255°(dec), 250°(dec), pK₁²⁵ 3.04, pK₂²⁵ 10.33**. Recryst from H₂O or 50% aq EtOH. Alternatively, dissolve *ca* 3g in conc HCl (10mL) and evaporate to dryness. Dissolve the residual hydrochloride in the minimum volume of warm H₂O and make faintly alkaline with aq NH₃. Collect the crystals and dry in a vacuum at 100°. [*J Am Chem Soc* 56 134 1934.]

***p*-Bromo-*N,N*-dimethylaniline** [586-77-6] **M 200.1, m 55°, b 264°, pK²⁵ 4.23**. Refluxed for 3h with two equivalents of acetic anhydride, then fractionally distd under reduced pressure

1-Bromo-2,4-dinitrobenzene [584-48-5] **M 247.0, m 75°**. Crystd from ethyl ether, isopropyl ether, 80% EtOH or absolute EtOH.

2-(2-Bromoethyl)-1,3-dioxane [33884-43-4] **M 195.1, b 67-70°/2.8mm, 71-72°/4mm, 95°/15mm, d₄²⁰ 1.44, n_D²⁰ 1.4219**. Purify by vacuum fractionation. Also dissolve in Et₂O, wash with aqueous NaHCO₃, dry extract with Na₂SO₄, filter and fractionate. NMR in CCl₄ has δ 1.3 (m, 1H), 2.1 (m, 3H), 3.36 (t, 2H), 3.90 (m, 4H) and 4.57 (t, H). [*J Org Chem* 41 560 1976; NMR, MS: *Tetrahedron* 35 1969 1979; *J Pharm Sci* 60 1250 1971.]

2-(2-Bromoethyl)-1,3-dioxolane [18742-02-4] **M 181.1, b 68-80°/8mm, 68-73°/10mm, 78-80°/20mm, d₄²⁰ 1.510, n_D²⁰ 1.479**. Dissolve in pentane, wash with 5% aqueous NaHCO₃, dry (Na₂SO₄), and evaporate. Distil the residue. [NMR: *J Org Chem* 34 1122 1969; *J Pharm Sci* 60 1250 1971.]

***N*-(2-Bromoethyl)phthalimide** [574-98-1] **M 254.1, m 81-83°, 82.5-83.5°**. The following is to be carried out in an efficient **FUME HOOD**. Dissolve the compound (180g) in CS₂ (500 mL) by refluxing for 15 min (to cause the separation of the most likely impurity, 1,2-diphthalimidoethane), filter and evaporate under reduced pressure. The product forms light tan crystals. (m 78-80°). Recryst from EtOH (charcoal) [the compound (50g) is dissolved in hot 75% EtOH (200mL), boiled for *ca* 10 min, carbon added (5g, Norite), filtered and cooled to 0°], as white crystals (40g) which can be recrystd (m 80-81°) and further recrystn gave m 82-83°. [*Org Synth Coll Vol I* 119 1932, *Synthesis* 389 1976; NMR: *Bull Soc Chim Fr* II-165 1979.]

Bromoform [75-25-2] **M 252.8, f 8.1°, 55-56°/35mm, 149.6°/760mm, d¹⁵ 2.9038, d³⁰ 2.86460, n¹⁵ 1.60053, n 1.5988**. Storage and stability of bromoform and chloroform are similar. Ethanol, added as a stabilizer, is removed by washing with H₂O or with saturated CaCl₂ soln, and the CHBr₃, after drying with CaCl₂ or K₂CO₃, is fractionally distd. Prior to distn, CHBr₃ has also been washed with conc H₂SO₄ until the acid layer no longer became coloured, then dilute NaOH or NaHCO₃, and H₂O. A further purification step is fractional crystn by partial freezing.

3-Bromofuran [22037-28-1] **M 147.0, b 38.5°/40mm, 50°/110mm, 102.5-103°/atm, d 1.661, n 1.4970**. Purified by two steam distillations and dried over fresh CaO. It can be dried over Na metal (no obvious reaction) and fractionated. It is not very soluble in H₂O but soluble in organic solvents. Freshly distilled, it is a clear oil, but darkens on standing and eventually resinifies. It can be stored for long periods by covering the oil with an alkaline soln of hydroquinone and redistilled when required. It forms a characteristic

maleic anhydride adduct, **m** 131.5-132°. [*J Am Chem Soc* **52** 2083 1930, **53** 737 1931, adduct: **55** 430 1933.]

(±)-2-Bromohexadecanoic acid (2-bromopalmitic acid) [18263-25-7] **M** 335.3, **m** 51-53°, 52.3-52.5°, 53°, **pK_{Est}** ~3.2. Recrystd from pet ether (b 60-80°, charcoal) and finally from EtOH. The *ethyl ester* has **b** 177-178°/2mm, d_{28}^{28} 1.0484, n_D^{20} 1.4560. [IR: *J Org Chem* **21** 1426 1956.]

5-Bromoindole [10075-50-0] **M** 196.1, **m** 90.5-91°, 90-92°, **pK** 16.13 (NH). Purified by steam distn from a faintly alkaline soln. Cool the aqueous distillate, collect the solid, dry in a vacuum desiccator over P₂O₅ and recryst from aqueous EtOH (35% EtOH) or pet ether-Et₂O. λ_{max} in MeOH: 279, 287 and 296 (log ϵ 3.70, 3.69 and 3.53. The *picrate* has **m** 137-138°(dec) (from Et₂O-pet ether). [UV: *Chem Ber* **95** 2205 1962; UV and NMR: *Bull Soc Chim Fr* 4091 1970.]

5-Bromoisatin [87-48-9] **M** 226.0, **m** 245°(dec), 251-153°, 255-256°. Forms red prisms or needles from EtOH. The *N-acetate* crystallises as yellow prisms from *C₆H₆, **m** 170-172°, and the *N-methyl* derivative form orange-red needles from MeOH, **m** 172-173°. [*Chem Ber* **47** 360 1914, **53** 1545 1920; *Recl Trav Chim Pays-Bas* **73** 197 1954; *Tetrahedron Lett* 215 1978.]

6-Bromoisatin [6326-79-0] **M** 226.0, **m** 270°, **pK** 10.35. Recrystd from AcOH (yellow needles). It is a plant growth substance. [Sadler *J Org Chem* **21** 169 1956.]

2-Bromomethylanthraquinone [7598-10-9] **M** 301.1, **m** 200-202°. Recrystd from AcOH, the crystals are washed with a little Et₂O, dried in air and then in vac at 100°. It is prepared by bromination of 2-methylanthraquinone with Br₂/PhNO₂ at 145-150°, or *N*-bromosuccinimide in CCl₄ containing a trace of (PhCOO)₂.

2-(Bromomethyl)benzonitrile [22115-41-9] **M** 195.1, **m** 72-73°, 79°, **b** 152-155°/15mm. Purified by steam distn. Extract the distillate with Et₂O, dry extract (Na₂SO₄), evap and distil residue. The solidified distillate can be recrystd from pet ether or cyclohexane. NMR (CDCl₃) δ : 7.8-7.2 (m 4H), 4.62 (s, 2H); IR ν : 2238 cm⁻¹. **LACHRYMATORY** [*Chem Ber* **24** 2570 1891, **74** 675 1934; *Aust J Chem* **22** 577 1969.]

S-(+)-1-Bromo-2-methylbutane [534-00-9] **M** 151.1, **b** 38.2°/39mm, 49°/62mm, 60.8°(57-58°)/100mm, 65-65.6°/140mm, 116-122°/atm, d_4^{20} 1.2232, n_D^{20} 1.4453, $[\alpha]_D^{20}$ +5.1° (neat, +5.8° (c 5, CHCl₃). Wash with ice-cold H₂O, dried by freezing, shake twice with an equal vol of H₂SO₄ at 0°, and twice with an equal volume of H₂O at 0°. Freeze dried and kept over freshly heated (and then cooled) K₂CO₃, and distd through a vacuum jacketed column of broken glass. Alternatively, dissolve in pet ether (b 40-60°), wash with 5% NaOH, conc H₂SO₄ (at 0°), then H₂O, dry (CaCl₂), evaporate and distil. [*J Am Chem Soc* **74** 4858 1952, **81** 2779 1959; *J Chem Soc* 1413 1959, 2685 1950.]

2-Bromo-3-methylindole (2-bromoskatole) [1484-28-2] **M** 210.1, **m** 102-104°, **pK_{Est}** <0. Purified by chromatography on silica gel in CHCl₃/pet ether (1:2) followed by crystn from aqueous EtOH. [Phillips and Cohen *J Am Chem Soc* **108** 2023 1986.]

4-(Bromomethyl)-7-methoxycoumarin [35231-44-8] **M** 269.1, **m** 208-209°, 213-215°, 216-218°. Cryst from boiling AcOH, crystals are washed with AcOH, EtOH and dried in a vacuum, NMR (TFA) δ 3.97s, 4.57s, 6.62s, 6.92-7.19m and 7.80d. [*Biochem Biophys Res Commun* **45** 1262 1971.]

2-(Bromomethyl)-naphthalene [939-26-4] **M** 221.1, **m** 52-54°, 56°, 56-57°, **b** 133-136°/0.8mm, 214°/100mm. Dissolve in toluene, wash with saturated aqueous NaHCO₃, dry (Mg SO₄), evaporate and fractionally distil the residue and recrystallise the distillate from EtOH. [*J Chem Soc* 5044, 1952; *Bull Soc Chim Fr* 566 1953.]

2-Bromo-2-methylpropane [507-19-7] M 137.0, b 71-73°, d 1.218, n 1.429. Neutralised with K_2CO_3 , distd, and dehydrated using molecular sieves (5A), then vacuum distd and degassed by freeze-pump-thaw technique. Sealed under vacuum.

1-Bromonaphthalene [90-11-9] M 207.1, b 118°/6mm, d 1.489. Purified by passage through activated alumina, and three vacuum distns.

2-Bromonaphthalene [580-13-2] M 207.1, m 59°. Purified by fractional elution from a chromatographic column. Crystd from EtOH.

1-Bromo-2-naphthol [573-97-7] M 223.1, m 76-78°, pK_{Est} ~8.0. Crystd from EtOH.

6-Bromo-2-naphthol [15231-91-1] M 223.1, m 122-126°, pK_{Est} ~9.1. Crystd from EtOH.

5-Bromonicotinic acid [20826-04-4] M 202.0, m 178-182°, 189-190°, pK_{Est} ~4.4, pK^{25} 4.02 (50% aq EtOH). Recryst from H_2O and then from EtOH using charcoal. The *amide* has m 219-219.5° (from aq EtOH) and the *methyl ester* prepared by addition of ethereal diazomethane can be purified by sublimation in a vacuum and has m 98-99°, the *acid chloride* also can be sublimed in *vacuo* and has m 74-75° and gives the methyl ester in MeOH. [J Prakt Chem 138 244 1933; J Am Chem Soc 70 2381 1948; 82 4430 1960; J Chem Soc 35 1978.]

ω -Bromo-4-nitroacetophenone [99-81-0] M 244.1, m 98°. Crystd from $*C_6H_6$ -pet ether.

***o*-Bromonitrobenzene** [577-19-5] M 202.1, m 43°. Crystd twice from pet ether, using charcoal before the first crystn.

***m*-Bromonitrobenzene** [585-79-5] M 202.1, m 55-56°. Crystd twice from pet ether, using charcoal before the first crystn.

***p*-Bromonitrobenzene** [586-78-7] M 202.1, m 127°. Crystd twice from pet ether, using charcoal before the first crystn.

1-Bromooctadecane [112-89-0] M 333.4, m 26°, 27.3°, 28-30°, b 178-179°/2mm, 214-218°/15mm, d_4^{20} 0.976, n_D^{20} 1.461. Twice recrystd from the melt then distilled under vacuum three times and using the middle cut. Alternatively, wash the oil with aqueous Na_2SO_4 , then conc H_2SO_4 (cool) and again with aqueous Na_2SO_4 and then fractionate. [J Am Chem Soc 55 1574 1933, 72 171 1950; IR: Aust J Chem 12 743 1959; IR: Bull Soc Chim Fr 516 1957.]

(\pm)-2-Bromopentane [107-81-3] M 151.1, b 117.2°/753mm, 116-117°/atm, 117.5°/740mm, d_4^{20} 1.2190, n_D^{20} 1.4401. Dry over K_2CO_3 and distil through a short Vigreux column. [IR: J Am Chem Soc 74 4063 1952, 78 2199 1956.]

***p*-Bromophenacyl bromide** [99-73-0] M 277.9, m 110-111°. Crystd from EtOH (*ca* 8mL/g).

***o*-Bromophenol** [95-56-7] M 173.0, b 194°, d 1.490, pK^{25} 8.45. Purified by at least two passes through a chromatographic column.

***p*-Bromophenol** [106-41-2] M 173.0, m 64°, pK^{25} 9.36. Crystd from $CHCl_3$, CCl_4 , pet ether (b 40-60°), or water and dried at 70° under vacuum for 2h.

Bromophenol Blue (3,3',5,5'-tetrabromophenolsulfonephthalein) [115-39-9] M 670.0, m 270-271°(dec), 273°(dec), λ_{max} 422max, pK 3.62. Crystd from $*benzene$ or acetone/glacial acetic acid, and air dried. Indicator at pH 3.0 (yellow) and pH 4.6 (purple).

(4-Bromophenoxy)acetic acid [1878-91-7] M 231.1, m 158°, pK^{25} 3.13. Crystd from EtOH.

3-(4-Bromophenoxy)propionic acid [93670-18-9] M 247.1, m 146°, pK_{Est} ~4.2. Crystd from EtOH.

4-Bromophenylacetic acid [1878-68-8] M 215.1, m 112-113°, 113-115°, 114°, pK 4.19. Recrystd from H₂O as needles. The *acid chloride* has b 238°/atm, m 50°, and the *anilide* has m 174-175°. [*J Chem Soc* 161 1934, 1251 1948; *J Org Chem* 11 798 1946.]

4-Bromophenylhydrazine [589-21-9] M 187.1, m 108-109°, pK²⁰ -5.6 (aq H₂SO₄), pK²⁵ 5.05. Crystd from H₂O.

4-Bromophenyl isocyanate [2492-02-9] M 189.0, m 41-42°. Crystd from pet ether (b 30-40°).

4-Bromophenyl isothiocyanate [1985-12-2] M 214.1, m 56-58°. Recryst from boiling *n*-hexane. Any insoluble material is most probably the corresponding urea. It can be purified by steam distn, cool the receiver, add NaCl and extract in Et₂O, wash extract with N H₂SO₄; dry (MgSO₄), evaporate and recrystallise the residual solid. [*Org Synth Coll Vol IV* 700 1963; *Coll Vol I* 447 1941.]

Bromopicrin (tribromonitromethane) [464-10-8] M 297.8, m 10.2-10.3°, b 85-87°/16mm, d 2.788, n 1.579. Steam distd, dried with anhydrous Na₂SO₄ and vacuum distd. TOXIC.

R-(+)-2-Bromopropionic acid [10009-70-8] M 153.0, b 78°/4mm, [α]_D²⁵ +27.2° (neat), pK²⁵ 4.07. Dissolve in Et₂O, dry (CaCl₂), evap and distil through a short column. Distillation through a Podbielniak column led to decomposition. [Podbielniak column. A plain tube containing "Heli-Grid" Nichrome or Inconel wire packing. This packing provides a number of passage-ways for the reflux liquid, while the capillary spaces ensure very even spreading of the liquid, so that there is a very large area of contact between liquid and vapour while, at the same time, channelling and flooding are minimised. A column 1m high has been stated to have an efficiency of 200-400 theoretical plates (for further details, see Podbielniak *Ind Eng Chem (Anal Ed)* 13 639 1941; Mitchell and O'Gorman *Anal Chem* 20 315 1948)]. Store in the dark under N₂, preferably in sealed ampoules. Even at -10° it slowly decomposes. [*J Am Chem Soc* 76 6054 1954.]

3-Bromopropionic acid [590-92-1] M 153.0, m 62.5°, 62.5-63.5°, 63-64°. Crystallises as plates from CCl₄. It is soluble in organic solvents and H₂O. It has a pK_a²⁵ in H₂O of 4.01, and its *methyl ester* has b 65°/18mm and 80°/27mm. The *S-benzylisothiuronium salt* has m 136°. [*Org Synth Coll Vol I* 134 1948; *Justus Liebigs Ann Chem* 599 140 1956.]

N-(3-Bromopropyl)phthalimide [5460-29-7] M 268.1, m 72-74°, 74°. Place in a Soxhlet and extract with Et₂O, whereby the bis-phthalimido impurity is not extracted. Evaporate the Et₂O and recryst from EtOH or aqueous EtOH or pet ether. [*Chem Ber* 21 2669 1888; *Justus Liebigs Ann Chem* 614 83 1958; *Can J Chem* 31 1060 1953.]

2-Bromopyridine [109-04-6] M 158.0, b 49.0°/2.7mm, d 1.660, n 1.5713, pK²⁵ 0.90. Dried over KOH for several days, then distd from CaO under reduced pressure, taking the middle fraction.

Bromopyrogallol Red (5,5'-dibromopyrogallolsulfonephthalein) [16574-43-9] M 576.2, m 300°, λ_{max} 538nm (ε 54,500 H₂O pH 5.6-7.5), pK₁ 2.9, pK₂ 4.39, pK₃ 9.15, pK₄ 11.72. Crystd from 50% EtOH, or aq alkaline soln by acid [*Suk Collect Czech Chem Commun* 31 3127 1966].

Bromopyruvic acid [1113-59-3] M 167.0, m 79-82°, pK_{Est} ~1.6. Dried by azeotropic distn (toluene), and then recrystd from dry CHCl₃. Dried for 48h at 20° (0.5 Torr) over P₂O₅. Stored at 0°. [Labandiniere et al. *J Org Chem* 52 157 1987].

5-Bromosalicyl hydroxamic acid [5798-94-7] M 210.1, m 232°(dec), pK_{Est(1)} ~ 1.5, pK_{Est(2)} ~ 7.0, pK_{Est(3)} ~ 8.7. Crystd from EtOH.

4-Bromostyrene [2039-82-9] M 183.1, b 49.5-50°/2.5mm, 87-88°/12mm, 102-104°/20mm, d 1.3984, n 1.5925. It polymerises above 75° in the presence of benzoyl peroxide. To purify, if it has not gone to a solid resin, dissolve in Et₂O, dry (MgSO₄), add ca 0.1g of 4-*tert*butylcatechol (polymerisation inhibitor) per 100g of bromostyrene. Filter, evap under reduced press (use as high a vac as possible) and distil. Store in dark bottles in the presence of the inhibitor (concn as above). [*Org Synth Coll Vol III* 204 1955.]

N-Bromosuccinimide [128-08-5] M 178.0, m 183-184°(dec). *N*-Bromosuccinimide (30g) was dissolved rapidly in 300mL of boiling water and filtered through a fluted filter paper into a flask immersed in an ice bath, and left for 2h. The crystals were filtered, washed thoroughly with ca 100mL of ice-cold water and drained on a Büchner funnel before drying under vac over P₂O₅ or CaCl₂ [Dauben and McCoy *J Am Chem Soc* 81 4863 1959]. Has also been crystd from acetic acid or water (10 parts, washed in water and dried *in vacuo*, [Wilcox et al. *J Am Chem Soc* 108 7693 1986; Shell et al. *J Am Chem Soc* 108 121 1986; Phillips and Cohen *J Am Chem Soc* 108 2013 1986.]

Bromotetronic acid [21151-51-9] M 179.0, m 183°(dec), pK²⁵ 2.23. Decolorised, and free bromine was removed by charcoal treatment of an ethyl acetate soln, then recrystd from ethyl acetate [Schuler, Bhatia and Schuler *J Phys Chem* 78 1063 1974].

Bromotheophylline [10381-75-6] M 259.1, m 309°, 315-320° (with browning and dec), pK_{Est(1)} ~5.5, pK_{Est(2)} ~9.2. It is purified by dissolving in the minimum volume of dilute NaOH (charcoal), filter and acidify to pH ca 3.5-4 and the solid that separates is collected, dried *in vacuo* at 100° and stored in a dark container. [*J Prakt Chem* [2] 118 158 1928; *Chem Ber* 28 3142 1895.]

Bromothymol Blue (3',3''-dibromothymolsulfonephthalein) [76-59-5] M 624.4, m 201-203°, pK₁ -0.66, pK₂ 6.99. Dissolved in aq 5% NaHCO₃ soln and ppted from the hot soln by dropwise addn of aq HCl. Repeated until the extinction did not increase (λ_{\max} 420nm). Indicator at pH 6.0 (yellow) and 7.6 (blue).

***p*-Bromotoluene** [106-38-7] M 171.0, m 28°, b 184°, d 1.390. Crystd from EtOH [Taylor and Stewart *J Am Chem Soc* 108 6977 1986].

α -Bromo-4-toluic acid [6232-88-8] M 215.1, m 229-230°, pK_{Est} ~3.2. Crystd from Me₂CO.

Bromotrichloromethane [75-62-7] M 198.5, f -5.6°, m 21°, b 104.1°, d 2.01, n 1.5061. Washed with aq NaOH soln or dilute Na₂CO₃, then with H₂O, and dried with CaCl₂, BaO, MgSO₄ or P₂O₅ before distilling in diffuse light and storing in the dark. Has also been purified by treatment with charcoal and by fractional crystn by partial freezing. Purified also by vigorous stirring with portions of conc H₂SO₄ until the acid did not discolour during several hours stirring. Washed with Na₂CO₃ and water, dried with CaCl₂ and then illuminated with a 1000W projection lamp at 15cm for 10h, after making it 0.01M in bromine. Passed through a 30 x 1.5cm column of activated alumina and fractionally redistilling through a 12-in Vigreux column. [Firestone and Willard *J Am Chem Soc* 83 3511 1961; see also Cadogan and Duell *J Chem Soc* 4154 1962.]

1-Bromo-3,3,3-trifluoroethane [421-06-7] M 163.0, m -94°, b 26-27°, d 1.788, n 1.332. Washed with water, dried (CaCl₂) and distd.

Bromotrifluoromethane (Freon 13B1) [75-63-8] M 148.9, b -59°, d 1.590. Passed through a tube containing P₂O₅ on glass wool into a vac system where it was frozen out in a quartz sample tube and degassed by a series of cycles of freezing, evacuating and thawing.

5-Bromouracil [51-20-7] M 191.0, m 293°, 303-305°, 312°(dec), pK₁²⁵ -7.25, pK₂²⁵ 7.83. Purified by dissolving in 2N NaOH (charcoal), filter and acidify with HCl. The ppte is dried *in vacuo* at 100° and recryst (prisms) twice from H₂O. [*J Am Chem Soc* 56 134 1934, UV: *J Am Chem Soc* 81 3786 1959; *J Org Chem* 23 1377 1958.]

5-Bromovaleric (γ -bromopentanoic) acid [2067-33-6] M 181.0, m 40°, pK_{Est} ~4.6. Crystd from pet ether.

α -Bromo-*p*-xylene [104-81-4] M 185.1, m 35°, b 218-220°/740mm. Crystd from EtOH or pentane.

Bromural [*N*-(aminocarbonyl)-2-bromo-3-methylbutanamide, bromisovalum] [496-67-3] M 223.1, m 154-155°. Crystd from toluene, and air dried.

Bufotenine hydrogen oxalate [2963-79-3] M 294.3, m 96.5°. Crystd from Et₂O.

1,3-Butadiene [106-99-0] M 54.1, b -2.6°. Dried by condensing with a soln of triethylaluminium in decahydronaphthalene; then flash distd. Also dried by passage over anhydrous CaCl₂ or distd from NaBH₄. Also purified by passage through a column packed with molecular sieves (4A), followed by cooling in Dry-ice/MeOH bath overnight, filtering off the ice and drying over CaH₂ at -78° and distd in a vacuum line.

***n*-Butane** [106-97-8] M 58.1, m -135°, b -0.5°. Dried by passage over anhydrous Mg(ClO₄)₂ and molecular sieves type 4A. Air was removed by prolonged and frequent degassing at -107°.

1,4-Butanediol (tetramethylene glycol) [110-63-4] M 90.1, f 20.4°, b 107-108°/4mm, 127°/20mm, d 1.02, n 1.4467. Distd and stored over Linde type 4A molecular sieves, or crystd twice from anhydrous diethyl ether/acetone, and redistd. Also purified by recrystn from the melt and doubly distd *in vacuo* in the presence of Na₂SO₄.

***meso*-2,3-Butanediol** [513-85-9] M 90.1, m 25°. Crystd from isopropyl ether.

***threo*-2,3-butanediol** [*R,R*(-): 24347-58-8] [*S,S*(+):19132-06-0] M 90.1, m 16-19°, 19.7°, b 77,5-78°/10mm, 179-180°/atm, [α]_D²⁰ (-) or (+) 13.1° (neat). Purified by fractional distn. The *bis*-(4-nitrobenzoate) has m 141-142° and [α]_D²⁵ ± 52° (c 4 CHCl₃). [*J Am Chem Soc* 79 734 1957, 74 425 1952, *Can J Res* 27 457 1949.]

1-Butanesulfonyl chloride [2386-60-9] M 156.6, b 75-76°/7mm, 98°/13mm, d₄²⁰ 1.2078, n_D²⁰ 1.4559. It has a pungent odour and is LACHRYMATORY. If IR shows OH bands then dissolve in Et₂O, wash with cold saturated aq NaHCO₃ (care since CO₂ will be generated) then H₂O, dry over solid Na₂SO₄, filter evaporate and distil the residue twice. Characterised by shaking a soln in Et₂O or *C₆H₆ with aq NH₃, collect the solid and recryst from CHCl₃, CCl₄ or Et₂O-pet ether, m 48°. [*J Am Chem Soc* 60 1488 1938; *J Org Chem* 5 83 1940.]

1-Butanethiol [109-79-5] M 90.2, b 98.4°, d²⁵ 0.837, n 1.443, n²⁵ 1.440, pK_{Est} ~11.3. Dried with CaSO₄ or Na₂SO₄, then refluxed from magnesium; or dried with, and distd from CaO, under nitrogen [Roberts and Friend *J Am Chem Soc* 108 7204 1986.] Has been separated from hydrocarbons by extractive distn with aniline.

Dissolved in 20% NaOH, extracted with a small amount of *C₆H₆, then steam distd, until clear. The soln was then cooled and acidified slightly with 15% H₂SO₄. The thiol was distd out, dried with CaSO₄ or CaCl₂, and fractionally distd under N₂ [Mathias and Filho *J Phys Chem* 62 1427 1958]. Also purified by pptn as lead mercaptide from alcoholic soln, with regeneration by adding dilute HCl to the residue after steam distn. *All operations should be carried out in a fume cupboard due to the TOXICITY and obnoxious odour of the thiol.*

2-Butanethiol [513-53-1] M 90.2, b 37.4°/134mm, d²⁵ 0.846, n²⁵ 1.4338, pK_{Est} ~11.4. Purified as for 1-butanethiol.

***n*-Butanol** [71-36-3] M 74.1, b 117.7°, d²⁵ 0.80572, n 1.39922, n¹⁵ 1.40118. Dried with MgSO₄, CaO, K₂CO₃, Ca or solid NaOH, followed by refluxing with, and distn from, calcium, magnesium activated with iodine, aluminium amalgam or sodium. Can also dry with molecular sieves, or by refluxing with *n*-butyl phthalate or succinate. (For method, see *Ethanol*.) *n*-Butanol can also be dried by efficient fractional distn, water passing over in the first fractn as a binary azeotrope (contains about 37% water). An ultraviolet-

transparent distillate has been obtained by drying with magnesium and distilling from sulfanilic acid. To remove bases, aldehydes and ketones, the alcohol has been washed with dil H_2SO_4 , then NaHSO_4 soln; esters were removed by boiling for 1.5h with 10% NaOH .

Also purified by adding 2g NaBH_4 to 1.5L butanol, gently bubbling with argon and refluxing for 1 day at 50° . Then added 2g of freshly cut sodium (washed with butanol) and refluxed for 1 day. Distd and the middle fraction collected [Jou and Freeman *J Phys Chem* **81** 909 1977].

2-Butanone (methyl ethyl ketone, MEK) [78-93-0] **M 72.1, b 79.6°, d 0.853, n 1.37850, n^{25} 1.37612, pK^{25} -7.2 (aq H_2SO_4)**. In general, purification methods are the same as for acetone. Aldehydes can be removed by refluxing with $\text{KMnO}_4 + \text{CaO}$, until the Schiff aldehyde test is negative, prior to distn. Shaking with satd K_2CO_3 , or passage through a small column of activated alumina, removes cyclic impurities. The ketone can be dried by careful distn (an azeotrope containing 11% water boils at 73.4°), or by CaSO_4 , P_2O_5 , Na_2SO_4 , or K_2CO_3 , followed by fractional distn. Purification as the bisulfite addition compound is achieved by shaking with excess satd Na_2SO_3 , cooled to 0° , filtering off the ppt, washing with a little ethyl ether and drying in air; this is followed by decomposition with a slight excess of Na_2CO_3 soln and steam distn, the distillate being satd with K_2CO_3 so that the ketone can be separated, dried with K_2CO_3 , filtered, and distd. Purification as the *NaI addition compound* (**m 73-74°**) is more convenient. (For details, see *Acetone*.) Small quantities of 2-butanone can be purified by conversion to the semicarbazone, recrystn to constant melting point, drying under vac over CaCl_2 and paraffin wax, refluxing for 30min with excess oxalic acid, followed by steam distn, salting out, drying and distilling [Cowan, Jeffery and Vogel *J Chem Soc* 171 1940].

cis-2-Butene [590-18-1] **M 56.1, b 2.95-3.05°/746mm**. The gas is dried with CaH_2 . Purified by gas chromatography. **HIGHLY FLAMMABLE**.

trans-2-Butene [624-64-6] **M 56.1, b 0.3-0.4°/744mm**. The gas is dried with CaH_2 . Purified by gas chromatography. **HIGHLY FLAMMABLE**.

2-Butene-1,4-dicarboxylic acid (trans-3-hexenedioic acid, trans- β -hydromuconic acid) [4436-74-2] **M 144.1, m 194-197°, 195-196°, $pK_{\text{Est}(1)} \sim 4.2$, $pK_{\text{Est}(2)} \sim 5.00$** . Crystd from boiling water, then dried at $50-60^\circ$ in a vacuum oven.

But-3-en-2-one (methyl vinyl ketone) [78-94-4] **M 70.1, b 79-80°/760mm, d 0.842**. See entry on p.302.

2-tert-Butoxycarbonyloxyimino-2-phenylacetonitrile (BOC-ON) [58632-95-4] **M 246.3, m 87-89°**. Triturate solid with 90% aq MeOH , filter, wash with 90% aq MeOH and dry in a vac. Recryst from MeOH (needles or plates), but use warm MeOH and cool to cryst, *do not boil as it decomposes slowly*. IR has ν 1785 ($\text{C}=\text{O}$) cm^{-1} and NMR (CDCl_3) usually shows two *tert*-butyl singlets for *syn* and *anti* isomers. Store in a brown bottle (fridge). It evolves CO_2 at room temp (stoppered bottle can explode!), but can be stored over silica gel which can extend its useful life to more than a year. [Itoh et al. *Org Synth* **59** 95 1980.]

2-Butoxyethanol (butyl cellosolve) [111-76-2] **M 118.2, b 171°/745mm, d 0.903, n 1.4191**. Peroxides can be removed by refluxing with anhydrous SnCl_2 or by passage under slight pressure through a column of activated alumina. Dried with anhydrous K_2CO_3 and CaSO_4 , filtered and distd, or refluxed with, and distd from NaOH .

4-Butoxyphenylacetic acid [4547-57-3] **M 208.3, m 86-87°, 88.5°, $pK_{\text{Est}} \sim 4.4$** . Purified by recrystn from pet ether (b $40-60^\circ$). [*J Am Chem Soc* **68** 2592 1946.]

n-Butyl acetate [123-86-4] **M 116.2, b 126.1°, d 0.882, n 1.394**. Distd, refluxed with successive small portions of KMnO_4 until the colour persisted, dried with anhydrous CaSO_4 , filtered and redistd.

tert-Butyl acetate [540-88-5] **M 116.2, b 97-98°, d 0.72**. Washed with 5% Na_2CO_3 soln, then saturated aqueous CaCl_2 , dried with CaSO_4 and distd.

tert-Butyl acetoacetate [1694-31-1] M 158.2, b 71°/10mm, 85°/20mm, d_4^{20} 0.954, n_D^{20} 1.42. Dist under reduced press through a short column. [Org Synth 42 28 1962.] HARMFUL VAPOUR.

tert-Butylacetylchloride [7065-46-5] M 134.6, b 68-71°/100mm, 81°/180mm, 128-132°/atm, d_4^{20} 0.964, n_D^{20} 1.423. Distil under vacuum. If IR shows OH group then treat with thionyl chloride or oxalyl chloride at ca 50° for 30min, evap and fractionate using a short column. Strongly LACHRYMATORY, use a good fume hood. [J Am Chem Soc 72 222 1950; J Org Chem 22 1551 1957.]

Butyl acrylate [141-32-2] M 128.2, b 59°/25mm, d 0.894, n_D^{12} 1.4254. Washed repeatedly with aqueous NaOH to remove inhibitors such as hydroquinone, then with distilled water. Dried with CaCl₂. Fractionally distd under reduced pressure in an all-glass apparatus. The middle fraction was sealed under nitrogen and stored at 0° in the dark until used [Mallik and Das J Am Chem Soc 82 4269 1960].

(±)-**sec-Butyl alcohol** [15892-23-6] M 74.1, b 99.4°, d 0.808. Purification methods are the same as for *n*-Butanol. These include drying with K₂CO₃ or CaSO₄, followed by filtration and fractional distn, refluxing with CaO, distn, then refluxing with magnesium and redistn; and refluxing with, then distn from CaH₂. Calcium carbide has also been used as a drying agent. Anhydrous alcohol is obtained by refluxing with *sec*-butyl phthalate or succinate. (For method see *Ethanol*.) Small amounts of alcohol can be purified by conversion to the alkyl hydrogen phthalate and recrystn [Hargreaves, J Chem Soc 3679 1956]. For purification of optical isomers, see Timmermans and Martin [J Chem Phys 25 411 1928].

tert-Butyl alcohol [75-65-0] M 74.1, m 23-25°, 25.7°, b 28.3°/60mm, 43.3°/123.8mm, 61.8°/315mm, 72.5°/507mm, 82.45°/760mm, d_4^{20} 0.7858, n_D^{20} 1.3878. Synthesised commercially by the hydration of 2-methylpropene in dilute H₂SO₄. Dried with CaO, K₂CO₃, CaSO₄ or MgSO₄, filtered and fractionally distd. Dried further by refluxing with, and distilling from, either magnesium activated with iodine, or small amounts of calcium, sodium or potassium, under nitrogen. Passage through a column of type 4A molecular sieve is another effective method of drying. So, also, refluxing with *tert*-butyl phthalate or succinate. (For method see *Ethanol*.) Other methods include refluxing with excess aluminium *tert*-butylate, or standing with CaH₂, and distilling as needed. Further purification is achieved by fractional crystn by partial freezing, taking care to exclude moisture. *tert*-Butyl alcohol samples containing much water can be dried by adding *benzene, so that the water distils off as a tertiary azeotrope, b 67.3°. Traces of isobutylene have been removed from dry *tert*-butyl alcohol by bubbling dry pre-purified nitrogen through for several hours at 40-50° before using. It form azeotropic mixtures with a large number of compounds. It has also been purified by distn from CaH₂ into Linde 4A molecular sieves which had been activated at 350° for 24h [Jaeger et al. J Am Chem Soc 101 717 1979].

Rapid purification: Dry *tert*-butanol with CaH₂ (5% w/v), distil and store over 3A molecular sieves.

***n*-Butylamine** [109-73-9] M 73.1, b 77.8°, d 0.740, n 1.4009, n_D^{25} 1.399, pK²⁵ 10.66. Dried with solid KOH, K₂CO₃, LiAlH₄, CaH₂ or MgSO₄, then refluxed with, and fractionally distd from P₂O₅, CaH₂, CaO or BaO. Further purified by pptn as the *hydrochloride*, m 213-213.5°, from ether soln by bubbling HCl gas into it. Re-pptd three times from EtOH by adding ether, followed by liberation of the free amine using excess strong base. The amine was extracted into ether, which was separated, dried with solid KOH, the ether removed by evapn and then the amine was distd. It was stored in a desiccator over solid NaOH [Bunnett and Davis J Am Chem Soc 82 665 1960; Lycan et al. Org Synth Coll Vol II 319 1943]. SKIN IRRITANT.

***R*-(-)-*sec*-Butylamine** [13250-12-9] M 73.1, b 61-63°/atm, 62.5°/atm, d_4^{20} 0.731, n_D^{20} 1.393, $[\alpha]_D^{20} +7.5^\circ$ (neat), pK 10.56. Dry over solid NaOH overnight and fractionate through a short helices packed column. The *L*-hydrogen tartrate salt has m 139-140° (from H₂O), the *1H*₂O has m 96° $[\alpha]_D^{21} +18.1^\circ$ (c 11, H₂O); the *hydrochloride* has m 152° $[\alpha]_D^{21} -1.1^\circ$ (c 13, H₂O) and the *benzoyl derivative* crystallises from EtOH as needles m 97°, $[\alpha]_D^{21} -34.9^\circ$ (c 11, H₂O). [J Chem Soc 921 1956; Acta Chem Scand 11 898 1957.]

***tert*-Butylamine** [75-64-9] M 73.1, b 42°, d 0.696, pK 10.68. Dried with KOH or LiAlH₄. Distd from CaH₂ or BaO.

n-Butyl *p*-aminobenzoate [94-25-7] M 193.2, m 57-59°, pK_{Est} ~2.5. Crystd from EtOH.

tert-Butylammonium bromide [60469-70-7] M 154.1, m >250°(dec). Recrystd several times from absolute EtOH and thoroughly dried at 105°.

4-*tert*-Butylaniline [769-92-6] M 149.2, m 14.5-15°, 15-16°, b 98.5-99°/3mm, 122°/20mm, d₄²⁰ 0.945, n_D²⁰ 1.538, pK²⁵ 4.95. Isolate as sulfate salt then liberate the free base with 10% aqueous NaOH, separate layers, dry over solid KOH and dist twice from Zn dust in a vacuum and store in brown containers. It has pK_a²⁵ (H₂O) 4.95 and (50% aq EtOH) 4.62. [J Am Chem Soc 76 2349 1954.] The *anilide* has m 171.5-172.3°, and the *hydrochloride* has m 270-274°. [J Chem Soc 680 1952; J Am Chem Soc 76 6179 1954.]

2-*tert*-Butylanthracene [13719-97-6] M 234.3, m 148-149°. Recrystd from EtOH and finally purified by TLC.

n-Butylbenzene [104-51-8] M 134.2, b 183.3°, d 0.860, n 1.4897, n²⁵ 1.487. Distd from sodium. Washed with small portions of conc H₂SO₄ until the acid was no longer coloured, then with water and aqueous Na₂CO₃. Dried with anhydrous MgSO₄, and distd twice from Na, collecting the middle fraction [Vogel J Chem Soc 607 1948].

tert-Butylbenzene [98-06-6] M 134.2, b 169.1°, d 0.867, n 1.493, n²⁵ 1.490. Washed with cold conc H₂SO₄ until a fresh portion of acid was no longer coloured, then with 10% aqueous NaOH, followed by distd water until neutral. Dried with CaSO₄ and distd in a glass helices-packed column, taking the middle fraction.

4-*tert*-Butyl benzoyl chloride [1710-98-1] M 196.7, b 135°/10mm, 149.9-150.5°/14mm, 266-268°(dec), d₄²⁰ 1.082, n_D²⁰ 1.536. Distil under vac. If IR shows OH group then treat with thionyl chloride or oxalyl chloride at ca 50° for 30min, evap and fractionate in a vac using a short column. Strongly LACHRYMATORY, use a good fume hood. [Bull Chem Soc Jpn 32 960 1959; J Am Chem Soc 72 5433 1950.]

n-Butyl bromide [109-65-9] M 137.0, b 101-102°, d²⁵ 1.2678, n 1.4399, n²⁵ 1.4374. Washed with conc H₂SO₄, water, 10% Na₂CO₃ and again with H₂O. Dried with CaCl₂, CaSO₄ or K₂CO₃, and distd. Redistd after drying with P₂O₅, or passed through two columns containing 5:1 silica gel/Celite mixture and stored with freshly activated alumina.

tert-Butyl bromoacetate [5292-43-3] M 195.1, b 52°/10mm, 74-76°/25mm, d₄²⁰ 1.324, n_D²⁵ 1.4162. Dissolve in Et₂O, wash well with ice cold 10% aqueous K₂CO₃, dry over CaCl₂, filter and evaporate the Et₂O then fractionate through a Vigreux column in a vacuum. LACHRYMATORY [Org Synth 34 28 1954, Coll Vol III 144 1955; J Am Chem Soc 64 2274 1942, 65 986 1943.]

4-*tert*-Butylcalix[4]arene [60705-62-6] M 648.9, m >300° (dec), 380° (dec), 344-346°. Recrystd from CHCl₃ in large solvated prisms (m 380° dec) effloresces on drying in air; *tetra-acetate* crystals from Ac₂O in colourless prisms m 332-333° dec. Crystals from CCl₄ or chlorobenzene + EtOH (m >300°) and *tetra-acetate* crystal from CHCl₃ + EtOH m >290° dec. Crystals from toluene in white plates with toluene of crystallisation m 344-346° (330-332°); the *tetra-acetate* crystallises with 1AcOH of crystallisation m 383-386° (softening at 330-340°, also m 283-286°), but acetylation with Ac₂O-NaOAc gives *triacetate* which recrystd from AcOH with 1AcOH of crystn m 278-281°. 4-*tert*-Butylcalix[4]arene (100mg) is unchanged after boiling for 4h with 10N KOH (0.04mL) in xylene (4mL). [Br J Pharmacol 10 73 1955; Monatsh Chem 109 767 1978; J Am Chem Soc 103 3782 1981; see also J.Vicens and V.Böhner eds, *Calixarenes*, Kluwer Academic Publ., Boston, 1991.]

4-*tert*-Butylcalix[6]arene [78092-53-2] M 972.3, m >300°, 380-381°. Recryst from CHCl₃ or CHCl₃ - MeOH as a white solid from the mother liquors of the calix[8]arene preparation. The *hexa-acetate* (Ac₂O-H₂SO₄) crystallises from CHCl₃-MeOH m 360-362° dec, and the (SiMe₃)₆ derivative crystallises from

$\text{CHCl}_3\text{-MeOH}$ **m** 410-412°. Stability in KOH-xylene is same as for the 4-*tert*-butylcalix[4]arene. [*J Am Chem Soc* **103** 3782 1981; see also J.Vicens and V.Böhner eds, *Calixarenes*, Kluwer Academic Publ., Boston, 1991.]

4-*tert*-Butylcalix[8]arene [68971-82-4] **M** 1297.8, **m** 411-412°. Recryst from CHCl_3 in fine colourless glistening needles. It melts sharply between 400-401° and 411-412° depending on the sample and is sensitive to traces of metal ions. TLC on silica gel (250 μm thick) and elution with $\text{CHCl}_3\text{-hexane}$ (3:4); it has R_F 0.75. The *octa-acetate* is prepared from 8g in Ac_2O (50mL) and 2 drops of conc H_2SO_4 refluxed for 2h. On cooling a colourless ppte separates and is recrystd from Ac_2O (1.2g 48%) **m** 353-354°. The $(\text{SiMe}_3)_8$ is prepared from 4-*tert*-butylcalix[8]arene (0.65g) in pyridine (4mL) with excess of hexamethyldisilazane (1mL) and trimethylchlorosilane (0.5mL) and refluxed under N_2 for 2h. Cool, evaporate the pyridine, triturate gummy residue with MeOH. Chromatography on silica gel using hexane- CH_2Cl_2 gave 0.5g (61%) with one spot on TLC. Crystallises from hexane- Me_2CO as colourless needles **m** 358-360°. [*J Am Chem Soc* **103** 3782 1981; *J Org Chem* **43** 4905 1978; **44** 3962 1979; *J Chem Soc, Chem Commun* 533 1981; see also J.Vicens and V.Böhner eds, *Calixarenes*, Kluwer Academic Publ., Boston, 1991.]

***tert*-Butyl carbazate** [870-46-2] **M** 132.2, **m** 41-42°, **b** 64°/0.01mm, **55-57°/0.4mm**. Dist in a Claisen flask with a water or oil bath at *ca* 80°. After a couple of drops have distd the carbazate is collected as an oil which solidifies to a snow white solid. It can be crystd with 90% recovery from a 1:1 mixt of pet ether (**b** 30-60°) and pet ether (**b** 60-70°). [*Org Synth* **44** 20 1964.]

4-*tert*-Butylcatchol [98-29-3] **M** 166.22, **m** 47-48°, **55-56°**, **75°**, **b** 265°/atm, **pK_{Est(1)}~9.5**, **pK_{Est(2)}~13.0**. Vacuum distd and recrystd from pentane or pet ether (or $^*\text{C}_6\text{H}_6$).

***n*-Butyl chloride** [109-69-3] **M** 92.6, **b** 78°, **d** 0.886, **n** 1.4021. Shaken repeatedly with conc H_2SO_4 (until no further colour developed in the acid), then washed with water, aq NaHCO_3 or Na_2CO_3 , and more water. Dried with CaCl_2 , or MgSO_4 (then with P_2O_5 if desired), decanted and fractionally distd. Alternatively, a stream of oxygen continuing *ca* three times as long as was necessary to obtain the first coloration of starch iodide paper by the exit gas. After washing with NaHCO_3 soln to hydrolyze ozonides and to remove the resulting organic acid, the liquid was dried and distd [Chien and Willard *J Am Chem Soc* **75** 6160 1953].

***tert*-Butyl chloride** [507-20-0] **M** 92.6, **f** -24.6°, **b** 50.4°, **d** 0.851, **n** 1.38564. Purification methods commonly used for other alkyl halides lead to decomposition. Some impurities can be removed by photochlorination with a small amount of chlorine prior to use. The liquid can be washed with ice water, dried with CaCl_2 or $\text{CaCl}_2 + \text{CaO}$ and fractionally distd. It has been further purified by repeated fractional crystn by partial freezing.

***tert*-Butyl chloroacetate** [107-59-5] **M** 150.6, **b** 48-49°/11mm, **60.2°/15mm**, **155°/atm (dec)**, **d₄²⁵ 1.4204**, **n_D²⁰ 1.4259**. Check the NMR spectrum, if satisfactory then dist in a vac, if not then dissolve in Et_2O , wash with H_2O , 10% H_2SO_4 until the acid extract does not become cloudy when made alkaline with NaOH. Wash the organic layer again with H_2O , then satd aq NaHCO_3 , dry over Na_2SO_4 , evap and fractionate through a carborundum-packed column or a 6-inch Widmer column (see *tert-butyl ethyl malonate for precautions to avoid decomposition during distn*). [*J Chem Soc* 940 1940; *J Am Chem Soc* **75** 4995 1953; *Org Synth Coll Vol* 144 1944.]

6-*tert*-Butyl-1-chloro-2-naphthol [525-27-9] **M** 232.7, **m** 76°, **b** 185°/15mm, **pK_{Est} ~8.0**. Crystd from pet ether.

***tert*-Butyl cyanide (trimethylacetoneitrile)** [630-18-2] **M** 83.1, **m** 16-18°, **d** 0.765, **b** 104-106°. Purified by a two stage vac distn and degassed by freeze-pump-thaw technique. Stored under vac at 0°. **TOXIC**, use efficient fume hood.

***tert*-Butyl cyanoacetate** [1116-98-9] **M** 141.2, **b** 40-42°/0.1mm, **54-56°/0.3mm**, **90°/10mm**, **107-108°/23mm**, **d₄²⁰ 0.989**, **n_D²⁰ 1.4198**. The IR spectrum of a film should have bands at 1742 (ester

CO) and 2273 (C≡N) but not OH band (*ca* 3500 broad) cm^{-1} . If it does not have the last named band then fractionally dist, otherwise dissolve in Et_2O , wash with satd aq NaHCO_3 , dry over K_2CO_3 , evap Et_2O , and dist residue under a vacuum (*see tert-butyl ethyl malonate for precautions to avoid decomposition during distn*). [*J Chem Soc* 423 1955; *Helv Chim Acta* 42 1214 1959].

4-*tert*-Butyl-1-cyclohexanone [98-53-3] M 154.3, m 49-50°. Crystd from pentane.

***n*-Butyl disulfide** [629-45-8] M 178.4, b 110-113°/15mm, d 0.938, n^{22} 1.494. Shaken with lead peroxide, filtered and distd in vacuum under N_2 .

***n*-Butyl ether (di-*n*-butyl ether)** [142-96-1] M 130.2, b 52-53°/26mm, 142.0°/760mm, d 0.764, n 1.39925, n^{25} 1.39685, pK^{25} -5.40 (aq H_2SO_4). Peroxides (detected by the liberation of iodine from weakly acid (HCl) solns of 2% KI) can be removed by shaking 1L of ether with 5-10mL of a soln comprising 6.0g of ferrous sulfate and 6mL conc H_2SO_4 and 110mL of water, with aq Na_2SO_3 , or with acidified NaI, water, then $\text{Na}_2\text{S}_2\text{O}_3$. After washing with dil NaOH, KOH, or Na_2CO_3 , then water, the ether is dried with CaCl_2 and distd. It can be further dried by distn from CaH_2 or Na (after drying with P_2O_5), and stored in the dark with Na or NaH. The ether can also be purified by treating with CS_2 and NaOH, expelling the excess sulfide by heating. The ether is then washed with water, dried with NaOH and distd [Kusama and Koike *J Chem Soc Japan, Pure Chem Sect* 72 229 1951]. Other purification procedures include passage through an activated alumina column to remove peroxides, or through a column of silica gel, and distn after adding about 3% (v/v) of a 1M soln of MeMgI in *n*-butyl ether.

***n*-Butyl ethyl ether** [628-81-9] M 102.2, b 92.7°, d 0.751, n 1.38175, n^{25} 1.3800. Purified by drying with CaSO_4 , by passage through a column of activated alumina (to remove peroxides), followed by prolonged refluxing with Na and then fractional distn.

***tert*-Butyl ethyl ether** [637-92-3] M 102.2, b 71-72°, d 0.741. Dried with CaSO_4 , passed through an alumina column, and fractionally distd.

***tert*-Butyl ethyl malonate** [32864-38-3] M 188.2, b 83-85°/8mm, 93-95°/17mm, 107-109°/24mm, d_4^{25} 0.994, n_D^{24} 1.4150. Likely impurity is monoethyl malonate, check IR for OH bands at 3330 br. To *ca* 50g of ester add ice cold NaOH (50g in 200mL of H_2O and 200g of ice). Swirl a few times (filter off ice if necessary), place in a separating funnel and extract with 2 x 75mL of Et_2O . Dry extract (MgSO_4) (since traces of acid decompose the *t*-Bu group of the ester, the distillation flask has to be washed with aq NaOH, rinsed with H_2O and allowed to dry). Addition of some K_2CO_3 or MgO before distilling is recommended to inhibit decomposition. Distil under reduced press through a 10 cm Vigreux column. *Decomposition is evidenced by severe foaming due to autocatalytic decomposition and cannot be prevented from accelerating except by stopping the distillation and rewashing the distillation flask with alkali again.* [*J Am Chem Soc* 66 1287 1944, 64 2714 1942; *Org Synth Coll Vol IV* 417 1963; *Org Synth* 37 35 1957.]

***n*-Butyl formate** [592-84-7] M 102.1, b 106.6°, d 0.891, n 1.3890. Washed with satd NaHCO_3 soln in the presence of satd NaCl, until no further reaction occurred, then with saturated NaCl soln, dried (MgSO_4) and fractionally distd.

Butyl glycolate [7397-62-8] M 132.2, b 191-192°/755mm, 187-190°/atm, d_4^{20} 1.019, n_D^{20} 1.4263. Dissolve in CHCl_3 (EtOH-free), wash with 5% KHCO_3 until effervescence ceases (if free acid is present), dry over CaCl_2 , filter, evaporate and distil through a short column. [Bøhme and Opfer *Z Anal Chem* 139 255 1953; *cf J Am Chem Soc* 73 5265 1951.]

***tert*-Butyl hydroperoxide (TBHP)** [75-91-2] M 90.1, f 5.4°, m 0.5-2.0°, b 38°/18mm, d 0.900, n 1.4013, pK^{20} 12.8. *Care should be taken when handling this peroxide because of the possibility of EXPLOSION. It explodes when heating over an open flame.* Alcoholic and volatile impurities can be removed by prolonged refluxing at 40° under reduced pressure, or by steam distn. For example, Bartlett, Benzing and Pincock [*J Am Chem Soc* 82 1762 1960] refluxed at 30mm pressure in an azeotropic separation apparatus until two phases no longer separated, and then distilled at 41°/23mm. Pure

material is stored under N_2 , in the dark at 0° . Crude commercial material has been added to 25% NaOH below 30° , and the crystals of the sodium salt have been collected, washed twice with benzene and dissolved in distilled water. After adjusting the pH of the soln to 7.5 by adding solid CO_2 , the peroxide was extracted into pet ether, from which, after drying with K_2CO_3 , it was recovered by distilling off the solvent under reduced pressure at room temperature [O'Brien, Beringer and Mesrobian *J Am Chem Soc* **79** 6238 1957]. **The temperatures should be kept below 75° .** It has also been distilled through a helices packed column (*ca* 15 plates) and material collected had **b** 34-35 $^\circ$ /20 mm. Similarly, a soln in pet ether has been extracted with cold aq NaOH, and the hydroperoxide has been regenerated by adding at 0° , $KHSO_4$ at a pH not higher than 4.5, then extracted into diethyl ether, dried with $MgSO_4$, filtered and the ether evapd in a rotary evaporator under reduced pressure [Milac and Djokic *J Am Chem Soc* **84** 3098 1962].

A 3M soln of TBHP in CH_2Cl_2 is prepared by swirling 85mL (0.61mol) of commercial TBHP (70% TBHP-30% H_2O , **d** 0.935 *ca* 7.2mmol/mL) with 140mL of CH_2Cl_2 in a separating funnel. The milky mixture is allowed to stand until the phases separate (*ca* 30min). The organic (lower) layer (*ca* 200mL) containing 0.60mole of TBHP was separated from the aqueous layer (*ca* 21mL) and used without further drying. TBHP is assayed by iodometric titration. With 90% grade TBHP (w/w, **d** 0.90, *ca* 9.0mmole/mL) no separation of layers occurs; i.e. when TBHP (66.67mL, 0.60mole) is added to CH_2Cl_2 (140mL) the resulting soln (*ca* 200mL) is clear. [*J Am Chem Soc* **77** 60032 1955, **74** 4742 1952; Akashi, Palermo and Sharpless *J Org Chem* **43** 2063 1978 states quality of available grades, handling and compatibility for reactions.]

2-tert-Butyl hydroquinone [1948-33-0] **M 166.2, m 125-127 $^\circ$, 127-128 $^\circ$, 129 $^\circ$, $pK_{Est(1)} \sim 10.5$, $pK_{Est(2)} \sim 11.6$.** Recryst from H_2O or MeOH and dried in a vacuum at 70° . Store in a dark container. [*Angew Chem* **69** 699 1957.]

n-Butyl iodide (1-iodobutane) [542-69-8] **M 184.0, b 130.4 $^\circ$, d 1.616, n_D^{25} 1.44967.** Dried with $MgSO_4$ or P_2O_5 , fractionally distd through a column packed with glass helices, taking the middle fraction and storing with calcium or mercury in the dark. Also purified by prior passage through activated alumina or by shaking with conc H_2SO_4 then washing with Na_2SO_3 soln. It has also been treated carefully with sodium to remove free HI and H_2O , before distilling in a column containing copper turnings at the top. Another purification consisted of treatment with bromine, followed by extraction of free halogen with $Na_2S_2O_3$, washing with H_2O , drying and fractional distn.

tert-Butyl iodide [558-17-8] **M 184.0, b 100 $^\circ$ (dec), d 1.544.** Vacuum distn has been used to obtain a distillate which remained colourless for several weeks at -5° . More extensive treatment has been used by Boggs, Thompson and Crain [*J Phys Chem* **61** 625 1957] who washed with aq $NaHSO_3$ soln to remove free iodine, dried for 1h with Na_2SO_3 at 0° , and purified by four or five successive partial freezings of the liquid to obtain colourless material which was stored at -78° .

tert-Butyl isocyanate [1609-86-5] **M 99.1, m 10.5-11.5 $^\circ$, b 30.5-32 $^\circ$ /10mm, 64 $^\circ$ /52mm, d_{25}^{25} 0.9079, n_D^{25} 1.470.** It is **LACHRYMATORY** and **TOXIC**, and should have IR with 2251 ($C\equiv N$) cm^{-1} and no OH bands. The NMR should have one band at 1.37 ppm from TMS. Purified by fractional distn under reduced pressure. [*J Org Chem* **36** 3056 1971; *J Prakt Chem.* **125** 152 1930.]

tert-Butyl isocyanide [7188-38-7] **M 83.1, b 91-92 $^\circ$ /730mm, 90 $^\circ$ /758mm, d^{20} 0.735.** Dissolve in pet ether (b 40-60 $^\circ$) wash with H_2O , dry (Na_2SO_4), remove pet ether under slight vacuum, dist using a vacuum-jacketed Vigreux column at atmospheric pressure, IR: ν 2134 cm^{-1} . [*Chem Ber* **93** 239 1960.]

tert-butyl isocyanoacetate [2769-72-4] **M 141.2, b 50 $^\circ$ /0.1mm, 49-50 $^\circ$ /10mm, 63-65 $^\circ$ /15mm, d_4^{20} 0.970, n_D^{20} 1.420.** If it contains some free acid (OH bands in IR) then dissolve in Et_2O , shake with 20% Na_2CO_3 , dry over anhydrous K_2CO_3 , evaporate and distil. [*Chem Ber* **94** 2814 1961.]

n-Butyl methacrylate [97-88-1] **M 142.2, b 49-52 $^\circ$ /0.1mm, d 0.896, n 1.424.** Purified as for butyl acrylate.

tert-Butyl methacrylate [585-07-9] M 142.2, f -48°, b 135-136°/760mm, d 0.878, n 1.415. Purified as for butyl acrylate.

2-tert-Butyl-4-methoxyphenol (2-tert-butyl-4-hydroxyanisole) [121-00-6] M 180.3, m 64.1°, pK_{Est} ~10.8. Fractionally distd *in vacuo*, then passed as a soln in CHCl₃ through alumina, and the solvent evaporated from the eluate. Recrystd from pet ether.

n-Butyl methyl ether [628-28-4] M 88.2, b 70°, d 0.744, pK -3.50 (aq H₂SO₄). Dried with CaSO₄, passed through an alumina column to remove peroxides, and fractionally distd.

tert-Butyl methyl ether [1634-04-4] M 88.2, b 56°, n 1.369. Same as for *n*-butyl methyl ether.

tert-Butyl methyl ketone [75-97-8] M 100.2, b 105°/746mm, 106°/760mm, d 0.814, n 1.401. Refluxed with a little KMnO₄. Dried with CaSO₄ and distd.

8-sec-Butylmetrazole [25717-83-3] M 194.3, m 70°. Crystd from pet ether, and dried for 2 days under vacuum over P₂O₅.

tert-Butyl nitrite [540-80-7] M 103.1, b 34°/250mm, 61-63°/atm, d₄²⁰ 0.8671, n_D²⁵ 1.3660. If it is free from OH bands (IR) then distil through a 12inch helices packed column under reduced pressure, otherwise wash with aq 5% NaHCO₃ (effervescence), then H₂O, dry (Na₂SO₄) and fractionate through a 10 theoretical plates column at *ca* 10mm pressure. [*J Chem Soc* 1968 1954, *J Am Chem Soc* 70 1516 1948; UV: *J Org Chem* 21 993 1956; IR: *Bull Soc Chim Belg* 60 240 1951.]

p-tert-Butylnitrobenzene [3282-56-2] M 179.2, m 28.4°. Fractionally crystd three times by partially freezing a mixture of the mono-nitro isomers, then recryst from MeOH twice and dried under vacuum [*Brown J Am Chem Soc* 81 3232 1959].

N-(n-Butyl)-5-nitro-2-furamide [14121-89-2] M 212.2, m 89-90°. Recrystd twice from EtOH/water mixture.

Butyloxirane (1-hexene oxide) [1436-34-6] M 100.2, b 116-117°/atm, 116-119°/atm, d₄²⁰ 0.833, n_D²⁰ 1.44051. Purified by fractional distn through a 2ft helices packed column at atmospheric pressure in a N₂ atm. [*J Org Chem* 30 1271 1965; *J Chem Soc* 2433 1927; ¹³C NMR *J Chem Soc Perkin Trans 2* 861 1975.]

tert-Butyl peracetate [107-71-1] M 132.2, b 23-24°/0.5mm, n²⁵ 1.4030. Washed with NaHCO₃ from a *benzene soln, then redistd to remove *benzene [*Kochi J Am Chem Soc* 84 774 1962]. Handle with adequate protection due to possible **EXPLOSIVE** nature.

tert-Butylperoxy isobutyrate [109-13-7] M 160.2, f -45.6°. After diluting 90mL of the material with 120mL of pet ether, the mixture was cooled to 5° and shaken twice with 90mL portions of 5% NaOH soln (also at 5°). The non-aqueous layer, after washing once with cold water, was dried at 0° with a mixture of anhydrous MgSO₄ and MgCO₃ containing *ca* 40% MgO. After filtering, this material was passed, twice, through a column of silica gel at 0° (to remove *tert*-butyl hydroperoxide). The soln was evapd at 0°/0.5-1mm to remove the solvent, and the residue was recrystd several times from pet ether at -60°, then subjected to high vac to remove traces of solvent [*Milos and Golubovic J Am Chem Soc* 80 5994 1958]. Handle with adequate protection due to possible **EXPLOSIVE** nature.

tert-Butyl perphthalic acid [15042-77-0] M 238.2, pK_{Est} ~6.2. Crystd from Et₂O and dried over H₂SO₄. Possibly **EXPLOSIVE**.

p-tert-Butylphenol [98-54-4] M 150.2, m 99°, pK²⁵ 10.39. Crystd to constant melting point from pet ether (b 60-80°). It sublimes. Also purified *via* its benzoate, as for phenol.

***p*-tert-Butylphenoxyacetic acid** [1798-04-5] M 208.3, m 88-89°, pK_{Est} ~2.9. Crystd from pet ether/*C₆H₆ mixture.

***tert*-Butyl phenyl carbonate** [6627-89-0] M 194.2, b 74-78°/0.5mm, 83°/0.6mm, d₄²⁰ 1.05, n_D²⁰ 1.480. If IR is free from OH then purify by redistillation, otherwise, dissolve in Et₂O, wash with 5% HCl, then H₂O, dry over MgSO₄, evap and distil through a Claisen head under vacuum. Care should be taken in the distillation as distn of large quantities can lead to decomposition with liberation of CO₂ and isobutylene, use the necessary precautions. [J Am Chem Soc 79 98 1957.]

***n*-Butyl phenyl ether** [1126-79-0] M 150.2, b 210.5°, d 0.935. Dissolved in diethyl ether, washed first with 10% aq NaOH to remove traces of phenol, then repeatedly with distilled water, followed by evaporation of the solvent and distn under reduced pressure [Arnett and Wu J Am Chem Soc 82 5660 1960].

***N*-tert-Butyl α -phenyl nitrone** [3376-24-7] M 177.2, m 73-74°. Crystd from hexane.

Butyl phthalate [84-74-2] M 278.4, f -35°, b 340°/760mm, d 1.043. Freed from alcohol by washing with H₂O, or from acids and butyl hydrogen phthalate by washing with dilute NaOH. Distd at 10torr or less. (See also p. 195.)

4-*tert*-Butyl pyridine [3978-81-2] M 135.2, f -44.4°, b 194-197°atm, 197°/765mm, d₄²⁰ 0.923, n_D²⁰ 1.495, pK²⁵ 5.82. It is dried over solid KOH and is purified by fractional distn through an efficient column under dry N₂. Its *picrate* has m 153.9-154°, and the *hydrochloride* has m 151.7-154.8° (from Me₂CO). [J Am Chem Soc 73 3308, 3310 1951, IR: J Am Chem Soc 100 214 1978; J Chem Soc 4454 1960.]

Butyl stearate [123-95-5] M 340.6, m 26.3°, d 0.861. Acidic impurities removed by shaking with 0.05M NaOH or a 2% NaHCO₃ soln, followed by several water washes, then purified by fractional freezing of the melt and fractional crystn from solvents with boiling points below 100°.

***S*-tert-Butyl thioacetate** [999-90-6] M 132.2, b 31-32°/11mm, 38°/14mm, 44-45°/28mm, 67°/54mm, 135.6-135.9°/773mm, d₄²⁵ 0.9207, n_D²⁰ 1.4532. Dissolve in CHCl₃ (EtOH-free), wash with H₂O, 10% H₂SO₄, saturated aqueous NaHCO₃ (care CO₂ liberated), H₂O again, dried over Drierite and anhydrous K₂CO₃, and fractionate under reduced pressure. [J Am Chem Soc 72 3021 1950.]

***p*-tert-Butyltoluene** [98-51-1] M 148.3, f -53.2°, b 91°/28mm, d 0.854, n 1.4920. A sample containing 5% of the *meta*-isomer was purified by selective mercuration. Fractional distn of the solid arylmercuric acetate, after removal from the residual hydrocarbon, gave pure *p*-tert-butyltoluene [Stock and Brown J Am Chem Soc 81 5615 1959].

***tert*-Butyl 2,4,6-trichlorophenyl carbonate** [16965-08-5] M 297.6, m 64-66°. Crystd from a mixture of MeOH (90mL) and water (6mL) using charcoal [Broadbent et al. J Chem Soc (C) 2632 1967].

***N*-tert-Butyl urea** [1118-12-3] M 116.2, m 182°, 185°(dec). Possible impurity is *N,N'*-di-*tert*-butyl urea which is quite insol in H₂O. Recrystd from hot H₂O, filter off insol material, and cool to 0° to -5° with stirring. Dry in vac at room temp over KOH or H₂SO₄. If dried at higher temperatures it sublimes slowly. It can be recrystd from EtOH as long white needles or from 95% aq EtOH as plates. During melting point determination the bath temp has to be raised rapidly as the urea sublimes slowly above 100° at 760mm. [Org Synth Coll Vol III 151 1955.]

***n*-Butyl vinyl ether** [111-34-2] M 100.2, b 93.3°, d 0.775. After five washings with equal volumes of water to remove alcohols (made slightly alkaline with KOH), the ether was dried with sodium and distd under vacuum, taking the middle fraction [Coombes and Eley J Chem Soc 3700 1957]. Stored over KOH.

2-Butyne [503-17-3] M 54.1, b 0°/253mm, d 0.693. Stood with sodium for 24h, then fractionally distd under reduced pressure.

2-Butyne-1,4-diol [110-65-6] M 86.1, m 54-57°, 56-58°, b 238°. Crystd from EtOAc.

***n*-Butyraldehyde** [123-72-8] M 72.1, b 74.8°, d 0.810, n 1.37911, n¹⁵ 1.38164. Dried with CaCl₂ or CaSO₄, then fractionally distd under N₂. Lin and Day [*J Am Chem Soc* 74 5133 1952] shook with batches of CaSO₄ for 10min intervals until a 5mL sample, on mixing with 2.5mL of CCl₄ containing 0.5g of aluminium isopropoxide, gave no ppt and caused the soln to boil within 2min. Water can be removed from *n*-butyraldehyde by careful distn as an azeotrope distilling at 68°. The aldehyde has also been purified through its bisulfite compound which, after decomposing with excess NaHCO₃ soln, was steam distd, extracted under N₂ into ether and, after drying, the extract was fractionally distd [Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960].

Butyramide [514-35-5] M 87.1, m 115°, b 230°. Crystd from acetone, *benzene, CCl₄-pet ether, 20% EtOH or water. Dried under vacuum over P₂O₅, CaCl₂ or 99% H₂SO₄.

***n*-Butyric acid** [107-92-6] M 88.1, f -5.3°, b 163.3°, d 0.961, n²⁵ 1.396, pK²⁵ 2.82. Distd, mixed with KMnO₄ (20g/L), and fractionally redistd, discarding the first third [Vogel *J Chem Soc* 1814 1948].

***n*-Butyric anhydride** [106-31-0] M 158.2, b 198°, d 0.968. Dried by shaking with P₂O₅, then distd.

γ -Butyrolactone [96-48-0] M 86.1, b 83.8°/12mm, d 1.124. Dried with anhydrous CaSO₄, then fractionally distd. *Handle in a fume cupboard due to TOXICITY.*

Butyronitrile [109-74-0] M 69.1, b 117.9°, d 0.793, n 1.3846, n³⁰ 1.37954. Treated with conc HCl until the smell of the isonitrile had gone, then dried with K₂CO₃ and fractionally distd [Turner *J Chem Soc* 1681 1956]. Alternatively it was twice heated at 75° and stirred for several hours with a mixture of 7.7g Na₂CO₃ and 11.5g KMnO₄ per L of butyronitrile. The mixture was cooled, then distd. The middle fraction was dried over activated alumina. [Schoeller and Wiemann *J Am Chem Soc* 108 22 1986.]

Butyryl chloride (butanoyl chloride) [141-75-3] M 106.6, f -89°, b 101-102°/atm, d₄²⁰ 1.026, n_D²⁰ 1.412. Check IR to see if there is a significant peak at 3000-3500 cm⁻¹ (br) for OH. If OH is present then reflux with less than one mol equiv of SOCl₂ for 1h and distil directly. The fraction boiling between 85-100° is then refractionated at atm pressure. Keep all apparatus free from moisture and store the product in sealed glass ampoules under N₂. **LACHRYMATORY** - *handle in a good fume hood.* [*Org Synth Coll Vol I* 147 1941.]

Cacotheline (2,3-dihydro-4-nitro-2,3-dioxo-9,10-*secostrychnidin*-10-oic acid) [561-20-6] M 508.4, pK_{Est(1)} ~4.4 (CO₂H), pK_{Est(2)} ~10.2 (N). Yellow crystals from H₂O. It is then dried over H₂SO₄ which gives the *dihydrate*, and in a vacuum over H₂SO₄ at 105° to give the anhydrous compound. The *hydrochloride* separates as the hydrate (on heating in vacuum at 80°) in orange-yellow prisms or plates, m 250°(dec), and forms a *resorcinol complex* which gives brown crystals from EtOH, m 325°, and a *hydroquinone complex* as dark red crystals from EtOH, m 319°. [*Chem Ber* 43 1042 1910, 86 232, UV: 242 1953; complexes: Gatto *Gazz Chim Ital* 85 1441 1955.] Used in the titrimetric estimation of Sn²⁺ ions [Szrvas and Lantos *Talanta* 10 477 1963].

Caffeic (3,4-dihydroxycinnamic) acid [331-39-5] M 180.2, m 195°, 223-225°, pK₁²⁵ 4.62, pK₂²⁵ 9.07 Crystd from water.

Caffeine [58-08-2] M 194.2, m 237°, pK₁⁴⁰ -0.10, pK₂⁵⁵ 1.22. Crystd from water or absolute EtOH.

(+)-Calarene (+ β -gurjunen, 1,3,3,11-tetramethyltricyclo[5.4.0.0^{2,4}]undecan-7-ane, (1aR)-1,1,7c,7ac-tetramethyl-1a,2,3,5,6,7,7a,7b-octahydro-1H-cyclopropa[α]naphthalene, new name 1(10)aristolene) [17334-55-3] M 204.35, b 45-47°/0.008-0.01mm, 255-258°/atm, d₄²⁰ 0.9340, n_D²⁰ 1.55051, [α]_D²⁰ +58° (EtOH), +81.8° (neat). Purified by gas chromatography (7%

2-Butyne-1,4-diol [110-65-6] M 86.1, m 54-57°, 56-58°, b 238°. Crystd from EtOAc.

***n*-Butyraldehyde** [123-72-8] M 72.1, b 74.8°, d 0.810, n 1.37911, n¹⁵ 1.38164. Dried with CaCl₂ or CaSO₄, then fractionally distd under N₂. Lin and Day [*J Am Chem Soc* 74 5133 1952] shook with batches of CaSO₄ for 10min intervals until a 5mL sample, on mixing with 2.5mL of CCl₄ containing 0.5g of aluminium isopropoxide, gave no ppt and caused the soln to boil within 2min. Water can be removed from *n*-butyraldehyde by careful distn as an azeotrope distilling at 68°. The aldehyde has also been purified through its bisulfite compound which, after decomposing with excess NaHCO₃ soln, was steam distd, extracted under N₂ into ether and, after drying, the extract was fractionally distd [Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960].

Butyramide [514-35-5] M 87.1, m 115°, b 230°. Crystd from acetone, *benzene, CCl₄-pet ether, 20% EtOH or water. Dried under vacuum over P₂O₅, CaCl₂ or 99% H₂SO₄.

***n*-Butyric acid** [107-92-6] M 88.1, f -5.3°, b 163.3°, d 0.961, n²⁵ 1.396, pK²⁵ 2.82. Distd, mixed with KMnO₄ (20g/L), and fractionally redistd, discarding the first third [Vogel *J Chem Soc* 1814 1948].

***n*-Butyric anhydride** [106-31-0] M 158.2, b 198°, d 0.968. Dried by shaking with P₂O₅, then distd.

γ -Butyrolactone [96-48-0] M 86.1, b 83.8°/12mm, d 1.124. Dried with anhydrous CaSO₄, then fractionally distd. *Handle in a fume cupboard due to TOXICITY.*

Butyronitrile [109-74-0] M 69.1, b 117.9°, d 0.793, n 1.3846, n³⁰ 1.37954. Treated with conc HCl until the smell of the isonitrile had gone, then dried with K₂CO₃ and fractionally distd [Turner *J Chem Soc* 1681 1956]. Alternatively it was twice heated at 75° and stirred for several hours with a mixture of 7.7g Na₂CO₃ and 11.5g KMnO₄ per L of butyronitrile. The mixture was cooled, then distd. The middle fraction was dried over activated alumina. [Schoeller and Wiemann *J Am Chem Soc* 108 22 1986.]

Butyryl chloride (butanoyl chloride) [141-75-3] M 106.6, f -89°, b 101-102°/atm, d₄²⁰ 1.026, n_D²⁰ 1.412. Check IR to see if there is a significant peak at 3000-3500 cm⁻¹ (br) for OH. If OH is present then reflux with less than one mol equiv of SOCl₂ for 1h and distil directly. The fraction boiling between 85-100° is then refractionated at atm pressure. Keep all apparatus free from moisture and store the product in sealed glass ampoules under N₂. **LACHRYMATORY - handle in a good fume hood.** [*Org Synth Coll Vol I* 147 1941.]

Cacotheline (2,3-dihydro-4-nitro-2,3-dioxo-9,10-*secostrychnidin*-10-oic acid) [561-20-6] M 508.4, pK_{Est(1)} ~4.4 (CO₂H), pK_{Est(2)} ~10.2 (N). Yellow crystals from H₂O. It is then dried over H₂SO₄ which gives the *dihydrate*, and in a vacuum over H₂SO₄ at 105° to give the anhydrous compound. The *hydrochloride* separates as the hydrate (on heating in vacuum at 80°) in orange-yellow prisms or plates, m 250°(dec), and forms a *resorcinol complex* which gives brown crystals from EtOH, m 325°, and a *hydroquinone complex* as dark red crystals from EtOH, m 319°. [*Chem Ber* 43 1042 1910, 86 232, UV: 242 1953; complexes: Gatto *Gazz Chim Ital* 85 1441 1955.] Used in the titrimetric estimation of Sn²⁺ ions [Szrvas and Lantos *Talanta* 10 477 1963].

Caffeic (3,4-dihydroxycinnamic) acid [331-39-5] M 180.2, m 195°, 223-225°, pK₁²⁵ 4.62, pK₂²⁵ 9.07 Crystd from water.

Caffeine [58-08-2] M 194.2, m 237°, pK₁⁴⁰ -0.10, pK₂⁵⁵ 1.22. Crystd from water or absolute EtOH.

(+)-Calarene (+ β -gurjunen, 1,3,3,11-tetramethyltricyclo[5.4.0.0^{2,4}]undecan-7-ane, (1aR)-1,1,7c,7ac-tetramethyl-1a,2,3,5,6,7,7a,7b-octahydro-1H-cyclopropa[α]naphthalene, new name 1(10)aristolene) [17334-55-3] M 204.35, b 45-47°/0.008-0.01mm, 255-258°/atm, d₄²⁰ 0.9340, n_D²⁰ 1.55051, [α]_D²⁰ +58° (EtOH), +81.8° (neat). Purified by gas chromatography (7%

propylene glycol adipate on unglazed tile particles of size 0.2-0.3mm, 400 cm column length and 0.6 cm diameter, at 184°, with N₂ carrier gas at a flow rate of 0.54 mL/sec using a thermal detector). Also purified by chromatography on alumina (200 times the weight of calarene) and eluted with pet ether. UV: λ_{\max} 200 and 210 nm (ϵ 9560, 5480) in EtOH. [IR: *Sorm Collect Czech Chem Commun* **18** 512 1953, **29** 795 1964; *Tetrahedron Lett* 827 1962, 225 1963.]

Calcon carboxylic acid [3-hydroxy-4-(2-hydroxy-4-sulfo-1-naphthylazo)naphthalene-2-carboxylic acid] [3737-95-9] **M 428.4, m 300°, λ_{\max} 560nm, pK₁ 1.2, pK₂ 3.8, pK₃ 9.26, pK₄ 13.14.** Purified through its *p*-toluidinium salt. The dye was dissolved in warm 20% aq MeOH and treated with *p*-toluidine to ppt the salt after cooling. Finally recrystd from hot water. [Itoh and Ueno *Analyst (London)* **95** 583 1970.] Patton and Reeder (*Anal Chem* **28** 1026 1956) indicator and complexes with Ca in presence of Mg and other metal ions.

Calmagite [3147-14-6] **M 358.4, m 300°, pK₁ 8.1, pK₂ 12.4.** Crude sample was extracted with anhydrous diethyl ether [Lindstrom and Diehl *Anal Chem* **32** 1123 1960]. Complexes with Ca, Mg and Th.

Campesterol (24R-24-methylcholest-5-en-3 β -ol) [474-62-4] **M 400.7, m 156-159°, 157-158°, $[\alpha]_D^{24}$ -35.1° (c 1.2, CHCl₃).** Recryst twice from hexane and once from Me₂CO. The *benzoyl derivative* has **m 158-160° $[\alpha]_D^{23}$ -8.6° (CHCl₃), the acetyl derivative has m 137-138° (from EtOH) and $[\alpha]_D^{23}$ -35.1° (c 2.9, CHCl₃)** [*J Am Chem Soc* **63** 1155 1941].

1R,4S-(-)-Camphanic acid [13429-83-9] **M 198.2, m 190-192°, 198-200°, $[\alpha]_{548}^{20}$ -22.5° (c 1, dioxane), -4.4° (c 8, EtOH), pK_{Est} ~3.8.** Dissolve in CH₂Cl₂, dry (MgSO₄), filter, evaporate and residue is sublimed at 120°/0.5mm or 140°/1mm. [*Helv Chim Acta* **61** 2773 1978.]

1R,4S-(-)-Camphanic acid chloride [39637-74-6] **M 216.7, m 65-66.5°, 70.5-71°, $[\alpha]_{548}$ -23° (c 2, CCl₄), -7.5° (c 0.67, *benzene).** Soluble in toluene (50g/100mL at 0°) and crystals from pet ether (b 40-60°). It sublimes at 70°/5mm. Store dry at 0°, v (CCl₄) 1805s and 1780m cm⁻¹. [*J Chem Soc, Dalton Trans* 2229 1976.]

RS-Camphene [565-00-4] **M 136.2, m 51-52°, b 40-70°/10mm.** Crystd twice from EtOH, then repeatedly melted and frozen at 30mm pressure. [Williams and Smyth *J Am Chem Soc* **84** 1808 1962.] Alternatively it is dissolved in Et₂O, dried over CaCl₂ and Na, evaporated and the residue sublimed in a vacuum [NMR: *Chem Ber* **111** 2527 1978].

(-)-Camphene (1S-2,2-dimethyl-3-methylene norbornane) [5794-04-7] **M 136.2, m 49.2-49.6°, 49-50°, b 79-80°/58mm, 91.5°/100mm, d_4^{54} 0.8412, n_D^{54} 1.4564, $[\alpha]_D^{25}$ -106.2° (c 40, *C₆H₆), -117.5° (c 19, toluene), -113.5° (c 9.7, Et₂O).** Purified by fractionation through a Stedman column (see p. 441) at 100mm in a N₂ atmosphere, crystallised from EtOH and sublimed in a vacuum below its melting point. It is characterised by its *camphenilone semicarbazone*, **m 217-218.5°**, or *camphor semicarbazone*, **m 236-238°**. [NMR: *Chem Ber* **111** 2527 1978; *Justus Liebigs Ann Chem* **623** 217 1959; Bain et al. *J Am Chem Soc* **72** 3124 1950]

Camphor (1R-bornan-2-one) [*R*-(+)- 464-49-3]; [*S*-(-)- 464-48-2] **M 136.2, m 178.8°, 179.97°(open capillary), b 204°/atm, $[\alpha]_{546}^{35}$ (+) and (-) 59.6° (in EtOH), $[\alpha]_D^{20}$ (+) and (-) 44.3° (c 10, EtOH), $[\alpha]_{579}^{179}$ (+) and (-) 70.85° (melt).** Crystd from EtOH, 50% EtOH/water, MeOH, or pet ether or from glacial acetic acid by addition of water. It can be sublimed (50°/14mm) and also fractionally crystd from its own melt. It is steam volatile. It should be stored in tight containers as it is appreciably volatile at room temperature. The solubility is 0.1% (H₂O), 100% (EtOH), 173% (Et₂O) and 300% (CHCl₃). The *R*-oxime (from Et₂O, CHCl₃, or dil EtOH) **m 119° $[\alpha]_D^{20}$ -42.4° (c 3, EtOH);** the \pm oxime has **m 118-119°**. [*Chem Ber* **67** 1432 1934; Allan and Rodgers *J Chem Soc (B)* 632 1971; UV, NMR: Fairley et al. *J Chem Soc, Perkin Trans 1* 2109 1973; *J Am Chem Soc* **62** 8 1940.]

Camphoric acid (1,2,2-trimethyl-cyclopentan-1r,3c-dicarboxylic acid) [*1R,2S*-(+)- 124-83-4; *1S,2R*-(-)- 560-09-8] **M 200.2, m 186-188°, 187°, 186.5-189°, $[\alpha]_{546}^{20}$ (+) and (-) 57° (c 1,**

EtOH), $[\alpha]_D^{20}$ (+) and (-) 47.7° (c 4, EtOH), pK_1^{25} 4.71, pK_2^{25} 5.83 (for + isomer). Purified by reprecipitation from an alkaline solution by HCl, filtered, and recrystallized from water several times, rejecting the first crop. It forms leaflets from EtOH and Me₂CO and H₂O and is insoluble in CHCl₃. Soluble in H₂O is 0.8% at 25° and 10% at 100°; 50% (EtOH) and 5% in ethylene glycol. The (±)-acid has m 202-203°. The (+)-1-methyl ester had m 86° (from petroleum ether) $[\alpha]_D^{20}$ +45° (c 4, EtOH), and the (+)-3-methyl ester has m 77° (from petroleum ether) $[\alpha]_D^{17.5}$ +53.9° (c 3, EtOH). [*J Am Chem Soc* 53 1661 1931; *Helv Chim Acta* 30 933 1947; *Acta Chem Scand* 2 597 1948; *J Am Chem Soc* 80 6316 1958.]

(±)-Camphoric anhydride [595-30-2] M 182.2, transition temp 135°, m 223.5°. Crystallized from EtOH.

Camphorquinone (borna-2,3-dione) [1R(-)- 10334-26-6; 1S(+)- 2767-84-2] M 166.2, m 198.7°, 198-199°, 197-201°, $[\alpha]_D^{25}$ (-) and (+) 101.1° (c 2, EtOH). It can be purified by steam distillation, recrystallized (yellow prisms) from EtOH, C₆H₆ or Et₂O-petroleum ether and can be sublimed in a vacuum. The (±)-quinone forms needles from EtOH, m 197-198°, 203°. [*Helv Chim Acta* 13 1026; *Chem Ber* 67 1432 1934.]

RS-Camphorquinone [10373-78-1] M 166.2, m 199-202°. Purification is same as for above isomers.

(1R)-(-)-Camphor-10-sulfonic acid [35963-20-3] M 232.3, m 197.4-198°(dec), 197-198°, $[\alpha]_D^{20}$ -20.7° (c 5.4, H₂O), pK_{Est} ~-1. Forms prisms from AcOH or EtOAc, and is deliquescent in moist air. Store in tightly stoppered bottles. The NH₄ salt forms needles from H₂O $[\alpha]_D^{16}$ ±20.5° (c 5, H₂O). [*J Chem Soc* 127 279 1925; *J Am Chem Soc* 78 3063 1956.]

(1S)-(+)-Camphor-10-sulfonic acid [3144-16-9] M 232.3, m 193°(dec), 197-198°, $[\alpha]_{546}^{20}$ +27.5° (c 10, H₂O), $[\alpha]_D^{20}$ +43.5° (c 4.3, EtOH), pK_{Est} ~-1. Crystallized from ethyl acetate and dried under vacuum.

Camphor-10-sulfonyl chloride [1S(+)- 21286-54-4; 1R(-)- 39262-22-1] M 250.7, m 67-68°, 70°, $[\alpha]_D^{20}$ (+) and (-) 32.2° (c 3, CHCl₃). If free from OH bands in the IR then recrystallized from Et₂O or petroleum ether, otherwise treat with SOCl₂ at 50° for 30min, evaporate, dry residue over KOH in a vacuum and recrystallize. The (±)-acid chloride has m 85°. Characterized as the amide (prisms from EtOH) m 132°, $[\alpha]_D^{17}$ (+) and (-) 1.5° (EtOH). [Read and Storey *J Chem Soc* 2761 1930; *J Am Chem Soc* 58 62 1936.]

2,10-Camphorsultam [1R(+)- 108448-77-7; 1S(-)- 94594-90-8] M 215.3, m 181-183°, 185-187°, $[\alpha]_D^{20}$ (+) and (-) 32° (c 5, CHCl₃). The (-)-enantiomer is recrystallized from 95% EtOH. It dissolves in dilute aqueous NaOH and can be precipitated without hydrolysis by acidifying. It forms the N-Na salt in EtOH (by addition of Na to the EtOH solution) and the salt can be methylated with MeI to give the (-)-N-Me lactam with m 80° after recrystallization from hot H₂O and has $[\alpha]_D^{25}$ -59.6° (c 5, CHCl₃) [Shriner et al. *J Am Chem Soc* 60 2794 1938].

S-Canavanine [543-38-4] M 176.2, m 184°, $[\alpha]_D^{17}$ +19.4° (c 2, H₂O), pK_1^{25} 2.43, pK_2^{25} 9.41. Crystallized from aqueous EtOH.

S(L)-Canavanine sulfate [2219-31-0] M 274.3, m 172°(dec). See L-canavanine sulfate on p. 518 in Chapter 6.

Cannabinol [521-35-7] M 310.4, m 76-77°, b 185°/0.05mm. Crystallized from petroleum ether. Sublimed.

Canthaxanthin (trans) [514-78-3] M 564.9, m 211-212°, $A_{1cm}^{1\%}$ 2200 (470nm) in cyclohexane. Purified by chromatography on a column of deactivated alumina or magnesium oxide, or on a thin layer of silica gel G (Merck), using dichloromethane/diethyl ether (9:1) to develop the chromatogram. Stored in the dark and in an inert atmosphere at -20°.

Capric acid (decanoic acid) [334-48-5] M 172.3, m 31.5°, b 148°/11mm, d 0.886, n_D^{25} 1.424, pK_{Est} ~4.9. Purified by conversion to its methyl ester, b 114.0°/15mm (using excess MeOH, in the presence of H₂SO₄). After removal of the H₂SO₄ and excess MeOH, the ester was distilled under vacuum through a 3ft

column packed with glass helices. The acid was then obtained from the ester by saponification. [Trachtman and Miller *J Am Chem Soc* **84** 4828 1962].

***n*-Caproamide (*n*-hexanamide)** [628-02-4] M 115.2, m 100°. Crystd from hot water.

Caproic acid (hexanoic acid) [142-62-1] M 116.2, b 205.4°, d 0.925, n 1.417, pK²⁵ 4.85. Dried with MgSO₄ and fractionally distilled from CaSO₄.

ϵ -Caprolactam (azepan-2-one, aza-2-cycloheptanone, 2-oxohexamethyleneimine) [105-60-2] M 113.2, m 70°, 70.5-71.5°, 70-71°, 262.5°/760mm. Distd at reduced pressure, crystd from acetone or pet ether and redistd. Purified by zone melting. Very *hygroscopic*. Discolours in contact with air unless small amounts (0.2g/L) of NaOH, Na₂CO₃ or NaBO₂ are present. Crystd from a mixture of pet ether (185mL of b 70°) and 2-methyl-2-propanol (30mL), from acetone, or pet ether. Distd under reduced pressure and stored under nitrogen. [*Synthesis* 614 1978.]

Capronitrile (hexanenitrile) [124-12-9] M 125.2, b 163.7°, n 1.4069, n²⁵ 1.4048. Washed twice with half-volumes of conc HCl, then with saturated aqueous NaHCO₃, dried with MgSO₄, and distilled.

Caprylolactam (azanon-2-one, azacyclononan-2-one, 8-aminooctanoic acid lactam) [935-30-8] M 141.2, m 72°, 73°, 74-76°, 75°, 76-77°, b 119-122°/0.7mm, 150-151°/7-8mm, 164°/14mm, d₄⁷³ 1.009, n_D⁷³ 1.489, pK²⁵ 0.55 (AcOH). Dissolve in CHCl₃, decolorise with charcoal, evaporate to dryness and recrystallise from CHCl₃-hexane. Sublime at high vacuum. The *oxime* has m 117° (from *C₆H₆ or pet ether). [*J Med Chem* **14** 501 1971; *Justus Liebigs Ann Chem* **607** 67 1957.]

Capsaicin (*E-N*-[(4-hydroxy-3-methoxyphenyl)-methyl]-8-methyl-6-nonenamide) [404-86-4] M 305.4, m 64-66°, 65°, 66.1°, b 210-220°/0.01mm. Recrystd from pet ether (b 40-60°), or pet ether-Et₂O (9:1). Also purified by chromatography on neutral Al₂O₃ (grade V) and eluted successively with *C₆H₆, *C₆H₆-EtOAc (17:3) and *C₆H₆-EtOAc (7:3), and distilled at 120°/10⁻⁵mm, and repeatedly recrystd from isopropanol (charcoal), needles. [*J Chem Soc* 11025 1955, *J Chem Soc(C)* 442 1968.]

Capsorubin (3,3'-dihydroxy- κ,κ -carotene-6,6'dione) [470-38-2] M 604.9, m 218°, λ_{\max} 443, 468, 503 nm, in hexane. Possible impurities: zeaxanthin and capsanthin. Purified by chromatography on a column of CaCO₃ or MgO. Crystd from *benzene/pet ether or CS₂.

Captan (*N*-trichloromethylmercapto-cyclohex-4-ene-1,2-dicarboxamide) [133-06-2] M 300.5, m 172-173°. Crystd from CCl₄. Large quantities internally cause diarrhoea and vomiting.

Captopril (*S*-1-[3-mercapto-2-methyl-1-oxopropyl]-L-proline) [62571-86-2] M 217.3, m 103-104°(polymorphic unstable form m 86°, melts at 87-88° solidifies and then melts again at 104-105°), [α]_D²² -131° (c 1.7, EtOH), pK₁ 3.7, pK₂ 9.8. Purified by recrystn from EtOAc-hexane. Also purified by dissolving in EtOAc and chromatographed on a column of Wakogel C200 using a linear gradient of MeOH in EtOAc (0-100°) and fractions which give a positive nitroprusside test (for SH) are combined, evap and recrystd from EtOAc-hexane (1:1), white crystals [α]_D²⁰ -128.2° (c 2.0, EtOH). [Nam *J Pharm Sci* **73** 1843 1984]. Alternatively, dissolve in H₂O, apply to a column of AG-50Wx2 (BioRad) and eluted with H₂O. The free acid is converted to the dicyclohexylamine salt in MeCN by addition until the pH is 8-9 (moist filter paper). The salt is converted to the free acid by shaking with EtOAc and 10% aq KHSO₄ or passage through an AG50Wx2 column. The EtOAc soln is dried (MgSO₄) and recrystd as above from EtOAc-hexane [*Biochem J* **16** 5484 1977; NMR and IR: Horii and Watanabe *Yakugaku Zasshi (J Pharm Soc Japan)* **81** 1786 1961].

4-(Carbamoylmethoxy)acetanilide [14260-41-4] M 208.2, m 208°. Crystd from water.

3-Carbamoyl-1-methylpyridinium chloride (1-methylnicotinamide chloride, Trigonellamide) [1005-24-9] M 172.6, m 240°(dec). Crystd from MeOH.

Carbanilide (*sym*-diphenylurea) [102-07-8] M 212.3, m 242°. Crystd from EtOH or a large volume (40mL/g) of hot water.

9-Carbazolacetic acid [524-80-1] M 225.2, m 215°, pK_{Est} ~3.5. Crystd from ethyl acetate.

Carbazole [86-74-8] M 167.2, m 240-243°, pK <0. Dissolved (60g) in conc H₂SO₄ (300mL), extracted with three 200mL portions of *benzene, then stirred into 1600mL of an ice-water mixture. The ppt was filtered off, washed with a little water, dried, crystd from *benzene and then from pyridine/*benzene. [Feldman, Pantages and Orchin *J Am Chem Soc* 73 4341 1951]. Has also been crystd from EtOH or toluene, sublimed in vacuum, zone-refined, and purified by TLC.

Carbazole-9-carbonyl chloride [73500-82-0] M 300.0, m 100-103°, 103.5-104.5°. Recrystd from *C₆H₆. If it is not very pure (presence of OH or NH bands in the IR) dissolve in pyridine, shake with phosgene in toluene, evaporate and recrystallise the residue. Carry out this experiment in a good fume cupboard as COCl₂ is very TOXIC, and store the product in the dark. It is moisture sensitive. The *amide* has m 246.5-247°, and the *dimethylaminoethylamide hydrochloride* has m 197-198°. [Weston et al. *J Am Chem Soc* 75 4006 1953.]

4-Carboethoxy-3-methyl-2-cyclohexen-1-one (Hagemann's ester) [487-51-4] M 182, b 79-80°/0.2mm, 121-123°/4mm, 142-144°/15mm, d₄²⁰ 1.038. Dissolve in ether, shake with solid K₂CO₃, aqueous saturated NaHCO₃, dry (MgSO₄) and distil. *Semicarbazone* has m 165-167° (169°). [*J Am Chem Soc* 65, 631, 1943.]

1-Carboethoxy-4-methylpiperazine hydrochloride [532-78-5] M 204.7, m 168.5-169°, pK 7.31. Crystd from absolute EtOH.

N-Carboethoxyphthalimide (*N*-ethoxycarbonylphthalimide) [22509-74-6] M 219.2, m 87-89°, 90-92°. Crystd from toluene-pet ether (or *benzene-pet ether). Partly soluble in Et₂O, *benzene and CHCl₃. [*Chem Ber* 54 1112 1921.]

γ-Carboline {9*H*-pyrido[3,4-*b*]indole} [244-69-9] M 168.2, m 225°, pK~0. Crystd from water.

Carbon Black Leached for 24h with 1:1 HCl to remove oil contamination, then washed repeatedly with distd water. Dried in air, and eluted for one day each with *benzene and acetone. Again dried in air at room temp, then heated in a vacuum for 24h at 600° to remove adsorbed gases. [Tamamushi and Tamaki *Trans Faraday Soc* 55 1007 1959.]

Carbon disulfide [75-15-0] M 76.1, b 46.3°, d 1.264, n 1.627. Shaken for 3h with three portions of KMnO₄ soln (5g/L), twice for 6h with mercury (to remove sulfide impurities) until no further darkening of the interface occurred, and finally with a soln of HgSO₄ (2.5g/L) or cold, satd HgCl₂. Dried with CaCl₂, MgSO₄, or CaH₂ (with further drying by refluxing with P₂O₅), followed by fractional distn in diffuse light. **Alkali metals cannot be used as drying agents.** Has also been purified by standing with bromine (0.5mL/L) for 3-4h, shaking with KOH soln, then copper turnings (to remove unreacted bromine), and drying with CaCl₂. CS₂ is highly TOXIC and highly FLAMMABLE. *Work in a good fumehood.* Small quantities of CS₂ have been purified (including removal of hydrocarbons) by mechanical agitation of a 45-50g sample with a soln of 130g of sodium sulfide in 150mL of H₂O for 24h at 35-40°. The aqueous sodium thiocarbonate soln was separated from unreacted CS₂, then pptd with 140g of copper sulfate in 350g of water, with cooling. After filtering off the copper thiocarbonate, it was decomposed by passing steam into it. The distillate was separated from H₂O and distd from P₂O₅. [Ruff and Golla *Z Anorg Chem* 138 17 1924.]

Carbon tetrabromide [558-13-4] M 331.7, m 92.5°. Reactive bromide was removed by refluxing with dilute aqueous Na₂CO₃, then steam distd, crystd from EtOH, and dried in the dark under vacuum. [Sharpe and Walker *J Chem Soc* 157 1962.] Can be sublimed at 70° at low pressure.

Carbon tetrachloride [56-23-5] M 153.8, b 76.8°, d²⁵ 1.5842. For many purposes, careful fractional distn gives adequate purification. Carbon disulfide can be removed by shaking vigorously for several hours with saturated KOH, separating, and washing with water: this treatment is repeated. The CCl₄ is shaken with conc H₂SO₄ until there is no further coloration, then washed with water, dried with CaCl₂ or MgSO₄ and distd (from P₂O₅ if desired). **It must not be dried with sodium.** An initial refluxing with mercury for 2h removes sulfides. Other purification steps include passage of dry CCl₄ through activated alumina, and distn from KMnO₄. Carbonyl containing impurities can be removed by percolation through a Celite column impregnated with 2,4-dinitrophenylhydrazine (DNPH), H₃PO₄ and water. (Prepared by dissolving 0.5g DNPH in 6mL of 85% H₃PO₄ by grinding together, then mixing with 4mL of distd water and 10g Celite.) [Schwartz and Parks *Anal Chem* 33 1396 1961]. Photochlorination of CCl₄ has also been used: CCl₄ to which a small amount of chlorine has been added is illuminated in a glass bottle (e.g. for 24h with a 200W tungsten lamp near it), and, after washing out the excess chlorine with 0.02M Na₂SO₃, the CCl₄ is washed with distd water and distd from P₂O₅. It can be dried by passing through 4A molecular sieves and distd. Another purification procedure is to wash CCl₄ with aq NaOH, then repeatedly with water and N₂ gas bubbled through the liquid for several hours. After drying over CaCl₂ it is percolated through silica gel and distd under dry N₂ before use [Klassen and Ross *J Phys Chem* 91 3664 1987].

Rapid purification: Distil, discarding the first 10% of distillate or until the distillate is clear. The distilled CCl₄ is then stored over 5A molecular sieves.

Carbon tetrafluoride [75-73-0] M 88.0, b -15°. Purified by repeated passage over activated charcoal at solid-CO₂ temperatures. Traces of air were removed by evacuating while alternately freezing and melting. Alternatively, liquefied by cooling in liquid air and then fractionally distilled under vacuum. (The chief impurity originally present was probably CF₃Cl).

Carbon tetraiodide [507-25-5] M 519.6, m 168°(dec). Sublimed *in vacuo*.

N,N'-Carbonyldiimidazole [530-62-1] M 162.2, m 115.5-116°. Crystd from *benzene or tetrahydrofuran, in a dry-box.

1,1'-Carbonyldi(1,2,4-triazole) [41864-22-6] M 164.1, m 134-136°, 145-150°. Dissolve in tetrahydrofuran and evaporate at 10mm until it crystallises. Wash crystals with cold tetrahydrofuran and dry in a vacuum desiccator over P₂O₅ in which it can be stored for months. [*Recl Trav Chim Pays-Bas* 80 1372 1961; Potts *J Org Chem* 27 2631 1962; Staab *Justus Liebigs Ann Chem* 106 75 1957.]

Carbonyl sulfide [463-58-1] M 60.1. See carbonyl sulfide entry on p. 409 in Chapter 5.

***o*-Carboxyphenylacetonitrile** [6627-91-4] M 161.2, m 114-115°. Crystd (with considerable loss) from *benzene or glacial acetic acid.

(-)-Caryophyllene oxide (1-S-5c-6t-epoxy-6c,10,10-trimethyl-2-methylene-1r,9t-bicyclo-[7.2.0]undecane) [1139-30-6] M 220.4, m 62-63°, 63.5-64°, 64°, b 114-117°/1.8mm, 141-142°/11mm, d₄²⁰ 0.9666, n_D²⁰ 1.49564, [α]_D²⁰ -79° (c 2,CHCl₃), [α]_D²⁰ -68° (supercooled melt). Purified by TLC on silica gel with EtOAc-pet ether (b 60-80°) (15:85), and recrystallised from MeOH or *C₆H₆. [NMR: Warnhoff *Can J Chem* 42 1664 1964, Ramage and Whitehead *J Chem Soc* 4336 1954.]

(±)-Catechin [7295-85-4] M 272.3, m 177° (anhyd). Crystd from hot water. Dried at 100°.

Catechol (1,2-dihydroxybenzene, pyrocatechol) [120-80-9] M 110.1, m 105°, pK₁²⁵ 9.45, pK₂²⁵ 12.8. Crystd from *benzene or toluene. Sublimed under vacuum. [Rozo et al. *Anal Chem* 58 2988 1986.]

Cation exchange resin. Conditioned before use by successive washing with water, EtOH and water, and taken through two H⁺-Na⁺-H⁺ cycles by successive treatment with M NaOH, water and M HCl then washed with water until neutral.

(+)-Cedrol (octahydro-3,6,8,8-tetramethyl-1-3a,7-methanoazulen-6-ol, 8a*S*-6*c*-hydroxy-3*c*,6*t*,8,8-tetramethyl[8*ar*-*H*]-octahydro-3*H*,3*a**t*,7*t*-methanoazulene), [77-53-2] **m 82-86°**, 86-87°, $[\alpha]_D^{28} +10.5^\circ$ (c 5, CHCl₃), $[\alpha]_D^{18} +13.1^\circ$ (c 5.5, EtOH), $[\alpha]_D^{18} +14.3^\circ$ (c 10, dioxane). Purified by recrystn from aqueous MeOH. It is estimated colorimetrically with H₃PO₄ in EtOH followed by vanillin and HCl [Hayward and Seymour *Anal Chem* 20 572 1948]. The 3,5-dinitrobenzoyl derivative has **m 92-93°**. [*J Am Chem Soc* 83 3114 1961.]

β-Cellobiose [528-50-7] **M 342.3**, **m 228-229°(dec)**, $[\alpha]_D^{25} +33.3^\circ$ (c 2, water). Crystd from 75% aqueous EtOH.

Cellulose triacetate [9012-09-3] **M 72,000-74,000**. Extracted with cold EtOH, dried in air, washed with hot distd water, again dried in air, then dried at 50° for 30min. [Madorsky, Hart and Straus *J Res Nat Bur Stand* 60 343 1958.]

Cerulenin (helicocerin, 2*R*,3*S*-2,3-epoxy-4-oxo-7*E*,10*E*-dodecadienamide) [17397-89-6] **M 223.3**, **m 93-94°**, 93-95°, **b 120°/10⁻⁸mm**, $[\alpha]_D^{16} +63^\circ$ (c 2, MeOH). White needles from *C₆H₆. Also purified by repeated chromatography from Florisil and silica gel. It is soluble in EtOH, MeOH, *C₆H₆, slightly soluble in H₂O and pet ether. The dl-form has **m 40-42°** (from *C₆H₆-hexane), and the 2*R*,3*S*-tetrahydrocerulenin has **m 86-87°**, $[\alpha]_D^{20} +44.4$ (c 0.25, MeOH after 24h). [*Tetrahedron Lett* 2095 1978, 2039 1979; *J Am Chem Soc* 99 2805 1977; *J Org Chem* 47 1221 1982.]

Cetyl acetate [629-70-9] **M 284.5**, **m 18.3°**. Vacuum distd twice, then crystd several times from diethyl ether/MeOH.

Cetyl alcohol (1-hexadecanol) [36653-82-4] **M 242.5**, **m 49.3°**. Crystd from aqueous EtOH or from cyclohexane. Purified by zone refining. Purity checked by gas chromatography.

Cetylamine [629-54-9] **M 255.4**, **m 106-107°**, **b 235-236°/12mm**. Crystd from thiophene-free *benzene and dried under vacuum over P₂O₅.

Cetylamine (1-hexadecylamine) [143-27-1] **M 241.5**, **m 48°**, **b 162-165°/5.2mm**, **pK²⁵ 10.60**. Crystd from thiophene-free *benzene and dried under vacuum over P₂O₅.

Cetylammonium chloride [1602-97-7] **M 278.0**, **m 178°**. Crystd from MeOH.

Cetyl bromide (1-bromohexadecane) [112-82-3] **M 305.4**, **m 15°**, **b 193-196°/14mm**. Shaken with H₂SO₄, washed with water, dried with K₂CO₃ and fractionally distd.

Cetyl ether [4113-12-6] **M 466.9**, **m 54°**. Vacuum distd then crystd several times from MeOH/*benzene.

Cetylpyridinium chloride (H₂O) [6004-24-6] **M 358.0**, **m 80-83°**. Crystd from MeOH or EtOH/diethyl ether and dried *in vacuo*. [Moss et al. *J Am Chem Soc* 108 788 1986; Lennox and McClelland *J Am Chem Soc* 108 3771 1986.]

Cetyltrimethylammonium bromide (cetrimonium bromide, CTAB) [57-09-0] **M 364.5**, **m 227-235°(dec)**. Crystd from EtOH, EtOH/*benzene or from wet acetone after extracting twice with pet ether. Shaken with anhydrous diethyl ether, filtered and dissolved in a little hot MeOH. After cooling in the refrigerator, the ppt was filtered at room temperature and redissolved in MeOH. Anhydrous ether was added and, after warming to obtain a clear soln, it was cooled and crystalline material was filtered. [Dearden and Wooley *J Phys Chem* 91 2404 1987; Hakemi et al. *J Am Chem Soc* 91 120 1987.]

Cetyltrimethylammonium chloride [112-02-7] **M 320.0**. Crystd from acetone/ether mixture, EtOH/ether, or from MeOH. [Moss et al. *J Am Chem Soc* 109 4363 1987.]

Charcoal. Charcoal (50g) was added to 1L of 6M HCl and boiled for 45min. The supernatant was discarded, and the charcoal was boiled with two more lots of HCl, then with distilled water until the supernatant no longer gave a test for chloride ion. The charcoal (which was now phosphate-free) was filtered on a sintered-glass funnel and air dried at 120° for 24h. [Lippin, Talbert and Cohn *J Am Chem Soc* **76** 2871 1954.] The purification can be carried out using a Soxhlet extractor (without cartridge), allowing longer extraction times. Treatment with conc H₂SO₄ instead of HCl has been used to remove reducing substances.

Chaulmoogric acid [(13-cyclopent-2-enyl)tridecanoic acid] [29106-32-9] **M 280.4, m 68.5°, b 247-248°/20mm**, $[\alpha]_D^{20} +60^\circ$ (c 4, CHCl₃), **pK_{Est} ~5.0**. Crystd from pet ether or EtOH. The *Me ester* [24828-59-9] has **m 22°, b 227°/20mm** and $[\alpha]_D^{15} +50^\circ$ (c 5, CHCl₃).

Chelerythrine [34316-15-9] **M 389.4, m 207°**. Crystd from CHCl₃ by addition of MeOH.

Chelex 100 [11139-85-8]. Washed successively with 2M ammonia, water, 2M nitric acid and water. Chelex 100 may develop an odour on long standing. This can be removed by heating to 80° for 2h in 3M ammonia, then washing with water. [Ashbrook *J Chromatogr* **105** 151 1975.]

Chelidonic acid (4-oxopyran-2,6-dicarboxylic acid) [99-32-1] **M 184.1, m 262°, pK₂²⁵ 2.36**. Crystd from aqueous EtOH.

Chenodesoxycholic acid [474-25-9] **M 392.6, m 143°**. See 3 α ,7 α -dihydroxycholanolic acid on p. 207.

Chimyl alcohol (1-O-n-hexadecylglycerol) [(±) 506-03-6; 10550-58-0 (*chimyl alcohol*)] **M 316.5, m 64°**. Crystd from hexane.

Chloral [75-87-6; 302-17-0 (*hydrate*)] **M 147.4, b 98°, pK²⁵ 10.04**. Distd, then dried by distilling through a heated column of CaSO₄.

Chloralacetone chloroform [512-47-0] **M 324.9, m 65°**. Crystd from *benzene.

α -Chloralose (*R*-1,2-*O*-[2,2,2-trichloroethylidene]- α -D-glucofuranose) [15879-93-3] **M 309.5, m 180-182°, 187°, 186-188°, $[\alpha]_D^{26} +19.5^\circ$ (c 11, pyridine)**. Recrystd from EtOH, 38% aqueous EtOH, Et₂O, H₂O or CHCl₃. The solubility is 0.44% in H₂O at 15°, 0.83% in H₂O at 37°, 6.7% in EtOH at 25°. [Whiton and Hixon *J Am Chem Soc* **55** 2438 1933; *Helv Chim Acta* **6** 621 1923.] The β -isomer is less soluble in H₂O, EtOH or Et₂O and has **m 237.5-238°** [*J Am Chem Soc* **59** 1955 1937; *Acta Chem Scand* **19** 359 1965].

2-Chloroacetophenone [532-27-4] **M 154.6, m 54-56°**. Crystd from MeOH [Tanner *J Org Chem* **52** 2142 1987].

***p*-Chloranil (2,3,5,6-tetrachloro-1,4-benzoquinone)** [118-75-2] **M 245.9, m 290°, 294.2-294.6° (sealed tube)**. Crystd from acetic acid, acetone, *benzene, EtOH or toluene, drying under vac over P₂O₅, or from acetic acid, drying over NaOH in a vacuum desiccator. It can be sublimed under vacuum at 290°. Sample may contain significant amounts of the *o*-chloranil isomer as impurity. Purified by triple sublimation under vacuum. Recrystd before use. **It is a skin and mucous membrane irritant.** [UV: *Rec Trav Chim Pays Bas* **276** 684 1924; Brook *J Chem Soc* 5040 1952.]

Chloranilic acid (2,5-dichloro-3,6-dihydroxy-1,4-benzoquinone) [87-88-7] **M 209.0, m 283-284° pK₁²⁵ 1.22, pK₂²⁵ 3.01**. A soln of 8g in 1L of boiling water was filtered while hot, then extracted twice at about 50° with 200mL portions of *benzene. The aq phase was cooled in ice-water. The crystals were filtered off, washed with three 10mL portions of water, and dried at 115°. It can be sublimed in vacuum. [*J Phys Chem* **61** 765 1957.] The *diacetate* has **m 182-185°** [*J Am Chem Soc* **46** 1866 1924; Thamer and Voight *J Phys Chem* **56** 225 1952].

Chlorendic anhydride (1,4,5,6,7,7,-hexachloro-5-norbornene-2,3-dicarboxylic anhydride) [115-27-5] **M 370.9, m 234-236°**. 235-237°, 238°. Steam distn or recrystn from H₂O yields the diacid. The purified diacid yields the anhydride with Ac₂O. [Prill *J Am Chem Soc* **69** 62 1947.]

Chloroacetaldehyde dimethyl acetal [97-97-2] **M 124.6, m -34.4°, b 64°/23mm, 71-72°/35mm, d₄²⁰ 1.0172, n_D²⁰ 1.4175**. Purified by fractional distillation. [Melhotra *J Indian Chem Soc* **36** 4405 1959; *Bull Soc Chim Belg* **61** 393 1952.]

α-Chloroacetamide [79-07-2] **M 93.5, m 121°, b 224-225°/743mm**. Crystd from acetone and dried under vacuum over P₂O₅.

p-Chloroacetanilide [539-03-7] **M 169.6, m 179°**. Crystd from EtOH or aqueous EtOH.

Chloroacetic acid [79-11-8] **M 94.5, m 62.8°, b 189°, pK²⁵ 2.87**. Crystd from CHCl₃, CCl₄, *benzene or water. Dried over P₂O₅ or conc H₂SO₄ in a vacuum desiccator. Further purification by distn from MgSO₄, and by fractional crystn from the melt. Stored under vac or under dry N₂. [Bernasconi et al. *J Am Chem Soc* **107** 3621 1985.]

Chloroacetic anhydride [541-88-8] **M 171.0, m 46°, d 1.5494**. Crystd from *benzene.

Chloroacetone [78-95-5] **M 92.5, b 119°/763mm, d 1.15**. Dissolved in water and shaken repeatedly with small amounts of diethyl ether which extracts, preferentially, 1,1-dichloroacetone present as an impurity. The chloroacetone was then extracted from the aqueous phase using a large amount of diethyl ether, and distd at slightly reduced pressure. It was dried with CaCl₂ and stored at Dry-ice temperature. Alternatively, it was stood with CaSO₄, distd and stored over CaSO₄. **LACHRYMATORY**.

Chloroacetonitrile [107-14-2] **M 75.5, b 125°**. Refluxed with P₂O₅ for one day, then distd through a helices-packed column. Also purified by gas chromatography. **LACHRYMATOR, HIGHLY TOXIC**.

o-Chloroaniline [95-51-2] **M 127.6, m -1.9°, b 208.8°, d 1.213, n 1.588, pK²⁵ 2.66**. Freed from small amounts of the *p*-isomer by dissolving in one equivalent of H₂SO₄ and steam distilling. The *p*-isomer remains behind as the sulfate. [Sidgwick and Rubie *J Chem Soc* 1013 1921.] An alternative method is to dissolve in warm 10% HCl (11mL/g of amine) and on cooling, the hydrochloride of *o*-chloroaniline separates out. The latter can be recrystd until the acetyl derivative has a constant melting point. (In this way, yields are better than for the recrystn of the picrate from EtOH or of the acetyl derivative from pet ether.) [King and Orton *J Chem Soc* 1377 1911].

p-Chloroaniline [106-47-8] **M 127.6, m 70-71°, pK²⁵ 3.98**. Crystd from MeOH, pet ether (b 30-60°), or 50% aq EtOH, then *benzene/pet ether (b 60-70°), then dried in a vacuum desiccator. Can be distd under vacuum (b 75-77°/33mm).

p-Chloroanisole [623-12-1] **M 142.6, b 79°/11.5mm, 196.6°/760mm, d 1.164, n^{25.5} 1.5326**. Washed with 10% (vol) aqueous H₂SO₄ (three times), 10% aqueous KOH (three times), and then with water until neutral. Dried with MgSO₄ and fractionally distd from CaH₂ through a glass helices-packed column under reduced pressure.

9-Chloroanthracene [716-53-0] **M 212.9, m 105-107°**. Crystd from EtOH. [Masnori *J Am Chem Soc* **108** 1126 1986.]

10-Chloro-9-anthraldehyde [10527-16-9] **M 240.7, m 217-219°**. Crystd from EtOH.

o-Chlorobenzaldehyde [89-98-5] **M 140.6, m 11°, b 213-214°, d 1.248, n 1.566**. Washed with 10% Na₂CO₃ soln, then fractionally distd in the presence of a small amount of catechol.

3-Chlorobenzaldehyde [587-04-2] M 140.6, m 18°, b 213-214°, d 1.241, n 1.564. Purified by low temperature crystn from pet ether (b 40-60°).

4-Chlorobenzaldehyde [104-88-1] M 140.6, m 47°. Crystd from EtOH/water (3:1), then sublimed twice at 2mm pressure at a temperature slightly above the melting point.

Chlorobenzene [108-90-7] M 112.6, b 131.7°, d 1.107, n 1.52480. The main impurities are likely to be chlorinated impurities originally present in the *benzene used in the synthesis of chlorobenzene, and also unchlorinated hydrocarbons. A common purification procedure is to wash several times with conc H₂SO₄ then with aq NaHCO₃ or Na₂CO₃, and water, followed by drying with CaCl₂, K₂CO₃ or CaSO₄, then with P₂O₅, and distn. It can also be dried with Linde 4A molecular sieve. Passage through, and storage over, activated alumina has been used to obtain low conductance material. [Flaherty and Stern *J Am Chem Soc* 80 1034 1958.]

4-Chlorobenzenesulfonyl chloride [98-60-2] M 211.1, m 53°, b 141°/15mm. Crystd from ether in powdered Dry-ice, after soln had been washed with 10% NaOH until colourless and dried with Na₂SO₄.

4-Chlorobenzhydrazide [536-40-3] M 170.6, m 164°. Crystd from water.

2-Chlorobenzoic acid [118-91-2] M 156.6, m 139-140°, pK²⁵ 2.91. Crystd successively from glacial acetic acid, aq EtOH, and pet ether (b 60-80°). Other solvents include hot water or toluene (ca 4mL/g). Crude material can be given an initial purification by dissolving 30g in 100mL of hot water containing 10g of Na₂CO₃, boiling with 5g of charcoal for 15min, then filtering and adding 31mL of 1:1 aq HCl: the ppte is washed with a little water and dried at 100°.

3-Chlorobenzoic acid [535-80-8] M 156.6, m 154-156°, 158°, d₄²⁵ 1.496, pK²⁵ 3.82 (5.25 in 50% dimethylacetamide). Crystd successively from glacial acetic acid, aqueous EtOH and pet ether (b 60-80°). It also recrystd from *C₆H₆ or Et₂O-hexane, and sublimes at 55° in a vacuum. [*Anal Chem* 26 726 1954] The methyl ester has m 21°, b 231°/atm. The *S*-benzyl thiuronium salt has m 164-165° (from EtOH) [*Acta Chem Scand* 9 1425 1955; *J Chem Soc* 1318 1960].

4-Chlorobenzoic acid [74-11-3] M 156.6, m 238-239°, pK²⁵ 3.99. Same as for *m*-chlorobenzoic acid. Has also been crystd from hot water, and from EtOH.

2-Chlorobenzonitrile [873-32-5] M 137.6, m 45-46°. Crystd to constant melting point from *benzene/pet ether (b 40-60°).

4-Chlorobenzophenone [134-85-0] M 216.7, m 75-76°. Recrystd for EtOH. [Wagner et al. *J Am Chem Soc* 108 7727 1986.]

2-Chlorobenzothiazole [615-20-3] M 169.6, m 21°, 90-91.4°/4mm, 135-136°/28mm, d₄²⁰ 1.303, n_D²⁰ 1.6398. It is purified by fractional distn *in vacuo*. The 2-chloro-3-methylbenzothiazolinium 2,4-dinitrobenzenesulfonate crystallises from Ac₂O, m 162-163° (dec). [*J Am Chem Soc* 73 4773 1951; *J Org Chem* 19 1830 1954; *J Chem Soc* 2190 1930.]

***o*-Chlorobenzotrifluoride** [88-16-4] M 180.6, b 152.3°. Dried with CaSO₄, and distd at high reflux ratio through a silvered vacuum-jacketed glass column packed with one-eighth inch glass helices [Potter and Saylor *J Am Chem Soc* 73 90 1951].

***m*-Chlorobenzotrifluoride** [98-15-7] M 180.6, b 137.6°. Same as for *o*-chlorobenzotrifluoride above.

***p*-Chlorobenzotrifluoride** [98-56-6] M 180.6, b 138.6°. Same as for *o*-chlorobenzotrifluoride above.

2-Chlorobenzoxazole [615-18-9] M 153.6, b 95-96°/20mm, 198-202°/atm, d_4^{20} 1.331, n_D^{20} 1.570. Purified by fractional distn, preferably in a vacuum. [Siedel *J Prakt Chem* (2) 42 456 1890; *J Am Chem Soc* 75 712 1953.]

p-Chlorobenzyl chloride [104-83-6] M 161.0, m 28-29°, b 96°/15mm. Dried with CaSO₄, then fractionally distd under reduced pressure. Crystd from heptane or dry diethyl ether. **LACHRYMATORY.**

p-Chlorobenzylisothiuronium chloride [544-47-8] M 237.1, m 197°, pK_{Est} ~9.6 (free base). Crystd from conc HCl by addition of water.

2-Chlorobutane [78-86-4] M 92.6, b 68.5°, d 0.873, n^{25} 1.3945. Purified in the same way as *n*-butyl chloride.

2-(4-Chlorobutyl)-1,3-dioxolane [118336-86-0] M 164.6, b 56-58°/0.1mm, d_4^{20} 1.106, n_D^{20} 1.457. If the IR has a CHO band then just distil in vacuum. If it is present then dissolve in Et₂O, wash with H₂O, then saturated NaHCO₃, dry over MgSO₄, evaporate and distil. [*J Am Chem Soc* 73 1365 1951.]

N-Chlorocarbonyl isocyanate [27738-96-1] M 105.5, m -68°, b 63.6°/atm, d_4^{20} 1.310. Fractionally distd at atmospheric pressure using a 40cm column. **TOXIC vapour use a good fume hood.** Store dry, ν 2260 (NCO), 1818 (CO) and 1420 (NCO sym) cm⁻¹. [*Chem Ber* 106 1752 1975.]

trans-4-Chlorocinnamic acid [1615-02-7] M 182.6, m 243°, 248-250°, 249-251°, pK^{25} 4.41. Recrystd from EtOH or aq EtOH (charcoal). [*Org Synth Coll Vol IV* 731 1963, Walling and Wolfstin *J Am Chem Soc* 69 852 1947.]

Chlorocyclohexane [542-18-7] M 118.6, b 142.5°, d 1.00, n^{25} 1.46265. Washed several times with dilute NaHCO₃, then repeatedly with distilled water. Dried with CaCl₂ and fractionally distd.

4-Chloro-2,6-diaminopyrimidine (2,4-diamino-6-chloropyrimidine) [156-83-2] M 144.6, m 198°, 199-202°, pK^{25} 3.57. Purified by recrystn from boiling H₂O (charcoal) as needles; also crystallises from Me₂CO. [Büttner *Chem Ber* 36 2232 1903; Roth *J Am Chem Soc* 72 1914 1950; UV: *J Chem Soc* 3172 1962.]

4-Chloro-3,5-dimethylphenol [88-04-0] M 156.6, m 115.5°, pK^{25} 9.70. Crystd from *benzene or toluene.

1-Chloro-2,4-dinitrobenzene [97-00-7] M 202.6, m 48-50°, 51°, 52-54°, 54°, b 315°/atm, d_4^{22} 1.697. Usually crystd from EtOH or MeOH. Has also been crystd from Et₂O, *C₆H₆, *C₆H₆-pet ether or isopropyl alcohol. A preliminary purification step has been to pass its soln in *benzene through an alumina column. Also purified by zone refining. It exists in three forms: one stable and two unstable. The stable form crysts as yellow needles from Et₂O, m 51°, b 315°/atm with some dec, and is sol in EtOH. The labile forms also crystallises from Et₂O, m 43°, and is more soluble in organic solvents. The second labile form has m 27°. [Hoffman and Dame, *J Am Chem Soc* 41 1015 1919, Welsh *J Am Chem Soc* 63 3276 1941; *J Chem Soc* 2476 1957.]

4-Chloro-3,5-dinitrobenzoic acid [118-97-8] M 246.6, m 159-161°, pK_{Est} ~2.5. Crystd from EtOH/water, EtOH or *benzene.

2-Chloro-3,5-dinitropyridine [2578-45-2] M 203.5, m 62-65°, 63-65°, 64°, pK_{Est} <-5. Dissolve in CHCl₃, shake with saturated NaHCO₃, dry (MgSO₄), evaporate and apply to an Al₂O₃ column, elute with pet ether (b 60-80°), evaporate and recryst from *C₆H₆ or pet ether. [*Chem Pharm Bull Jpn* 8 28 1960; *Recl Trav Chim Pays-Bas* 72 573 1953.]

2-Chloroethanol (ethylene chlorohydrin) [107-07-3] M 80.5, b 51.0°/31 mm, 128.6°/760mm, d 1.201, n¹⁵ 1.444. Dried with, then distd from, CaSO₄ in the presence of a little Na₂CO₃ to remove traces of acid.

2-Chloroethyl bromide (1-bromo-2-chloroethane) [107-04-0] M 143.4, b 106-108°. Washed with conc H₂SO₄, water, 10% Na₂CO₃ soln, and again with water, then dried with CaCl₂ and fractionally distd before use.

2-Chloroethyl chloroformate [627-11-2] M 143.0, b 52-54°/12mm, 153°/760mm, d₄¹⁸ 1.3760, n_D²⁰ 1.446. Purified by fractional distn, preferably in a vacuum and stored in dry atmosphere. [*J Chem Soc* 2735 1957.]

1-(2-Chloroethyl)pyrrolidine hydrochloride [7250-67-1] M 170.1, m 167-170°, 173.5-174°, pK_{Est} ~8.5 (free base). Purified by recrystn from isopropanol-di-isopropyl ether (charcoal) and recrystallised twice more. The free base, b 55-56°/11mm, 60-63°/23mm and 90°/56mm, is relatively unstable and should be converted to the hydrochloride immediately, by dissolving in isopropanol and bubbling dry HCl through the soln at 0°, and filtering off the hydrochloride and recrystallising it. The picrate has m 107.3-107.8° (from EtOH) [*Cason J Org Chem* 24 247 1959; *J Am Chem Soc* 70 3098 1948].

2-Chloroethyl vinyl ether [110-75-8] M 106.6, b 109°/760mm, d 1.048, n 1.437. Washed repeatedly with equal volumes of water made slightly alkaline with KOH, dried with sodium, and distd under vacuum. TOXIC.

Chloroform [67-66-3] M 119.4, b 61.2°, d¹⁵ 1.49845, d¹⁰ 1.47060, n¹⁵ 1.44858. Reacts slowly with oxygen or oxidising agents, when exposed to air and light, giving, mainly, phosgene, Cl₂ and HCl. Commercial CHCl₃ is usually stabilized by addn of up to 1% EtOH or of dimethylaminoazobenzene. Simplest purifications involve washing with water to remove the EtOH, drying with K₂CO₃ or CaCl₂, refluxing with P₂O₅, CaCl₂, CaSO₄ or Na₂SO₄, and distilling. It must not be dried with sodium. The distd CHCl₃ should be stored in the dark to avoid photochemical formation of phosgene. As an alternative purification, CHCl₃ can be shaken with several small portions of conc H₂SO₄, washed thoroughly with water, and dried with CaCl₂ or K₂CO₃ before filtering and distilling. EtOH can be removed from CHCl₃ by passage through a column of activated alumina, or through a column of silica gel 4-ft long by 1.75-in diameter at a flow rate of 3mL/min. (The column, which can hold about 8% of its weight of EtOH, is regenerated by air drying and then heating at 600° for 6h. It is pre-purified by washing with CHCl₃, then EtOH, leaving in conc H₂SO₄ for about 8hr, washing with water until the washings are neutral, then air drying, followed by activation at 600° for 6h. Just before use it is reheated for 2h to 154°.) [McLaughlin, Kaniecki and Gray *Anal Chem* 30 1517 1958].

Carbonyl-containing impurities can be removed from CHCl₃ by percolation through a Celite column impregnated with 2,4-dinitrophenylhydrazine, phosphoric acid and water. (Prepared by dissolving 0.5g DNPH in 6mL of 85% H₃PO₄ by grinding together, then mixing with 4mL of distilled water and 10g of Celite.) [Schwartz and Parks *Anal Chem* 33 1396 1961]. Chloroform can be dried by distn from powdered type 4A Linde molecular sieves. For use as a solvent in IR spectroscopy, chloroform is washed with water (to remove EtOH), then dried for several hours over anhydrous CaCl₂ and fractionally distd. This treatment removes material absorbing near 1600 cm⁻¹. (Percolation through activated alumina increases this absorbing impurity). [Goodspeed and Millson *Chem Ind (London)* 1594 1967].

Rapid purification: Pass through a column of basic alumina (Grade I, 10g/mL of CHCl₃), and either dry by standing over 4A molecular sieves, or alternatively, distil from P₂O₅ (3% w/v). Use immediately.

Chlorogenic [3-(3,4-dihydroxycinnamoyl)quinic] acid [327-97-9] M 354.3, m 208°, [α]_D²⁵ -36° (c 1, H₂O), pK₁²⁵ 3.59, pK₂²⁵ 8.59. Crystd from water. Dried at 110°.

Chlorohydroquinone (2-chloro-1,4-dihydroxybenzene) [615-67-8] M 144.6, m 106°, b 263°, pK₁²⁵ 8.81, pK₂²⁵ 10.78. Crystd from CHCl₃ or toluene.

5-Chloro-8-hydroxy-7-iodoquinoline (vioform) [130-26-7] M 305.5, m 178-179°, pK_{Est(1)} ~2.6, pK_{Est(2)} ~7.0. Crystd from abs EtOH.

5-Chloroindole [17422-32-1] M 151.6, m 69-71°, 72-73°, b 120-130°/0.4mm, pK_{Est} <0 It is distd at high vacuum and recrystallises from pet ether (b 40-60°) or (b 80-100°) as glistening plates. The *picrate* has m 147° (146.5-147.5°)(from *C₆H₆). [*J Chem Soc* 3493 1955; *J Org Chem* 44 578 1979].

4-Chloriodobenzene [637-87-6] M 238.5, m 53-54°. Crystd from EtOH.

2,3-Chloromaleic anhydride [1122-17-4] M 167.0, m 121-121.5°. See 2,3-dichloromaleic anhydride on p. 198.

5-Chloro-2-methoxyaniline (2-amino-4-chloroanisole) [95-03-4] M 157.6, m 81-83°, 82-84°, 84°, pK²⁵ 3.56. Purified by steam distn and recrystn from H₂O or 40% aqueous EtOH. The *N*-acetate forms needles from hot H₂O m 104°; the *N*-benzoyl derivative forms needles from aq EtOH m 77-78°; the *picrate* has m 194° dec. [*J Am Chem Soc* 48 2657 1926.]

9-Chloromethyl anthracene [24463-19-2] M 226.7, m 141-142° dec, 141-142.5°. If it is free from OH in the IR then recryst from hexane-*C₆H₆ or *C₆H₆ as needles. If OH is present then some solvolysis has occurred. In this case treat 8.5g with SOCl₂ (4.8g) in dioxane (60mL) and reflux for 5h, then evaporate to dryness and wash the residue with cold *C₆H₆ and recrystallise. With KI/Me₂CO it forms the *iodomethyl* derivative. [Martin et al. *Helv Chim Acta* 38 2009 1955; *J Org Chem* 21 1512 1956.]

2-Chloro-3-methylindole (2-chloroskatole) [51206-73-6] M 165.6, m 114.5-115.5°, pK_{Est} <0. Purified by chromatography on silica gel in CH₂Cl₂/pet ether (1:2), followed by recrystn from aqueous EtOH or aqueous acetic acid. [Phillips and Cohen *J Am Chem Soc* 108 2023 1986.]

Chloromethyl methyl ether (MOMCl) [107-30-2] M 80.5, b 55-57°, d 1.060, n 1.396. If suspect (check IR), shake with satd aq CaCl₂ soln, dry over CaCl₂ and fractionally distil taking middle fraction. [Marvel and Porter *Org Synth Coll Vol I* 377 1941.] **VERY TOXIC and CARCINOGENIC.**

4-Chloro-2-methylphenol [1570-64-5] M 142.6, m 49°, pK²⁵ 9.71. Purified by zone melting.

4-Chloro-3-methylphenol [59-50-7] M 142.6, m 66°, pK²⁵ 9.55. Crystd from pet ether.

4-Chloro-2-methylphenoxyacetic acid (MCPA) [94-74-6] M 200.6, m 113-117°, 120°, 122-123°, pK²⁰ 3.62 (3.05). It is insoluble in H₂O (sol 0.55g/L at 20°), and recrystallises from *C₆H₆ or chlorobenzene as plates [*Acta Chem Scand* 6 993 1952]. The *S*-benzylthiuronium salt has m 164-165°, and the Cu²⁺ salt has m 247-249°dec [Armarego et al. *Nature* 183 1176 1959; UV: Duvaux and Grabe *Acta Chem Scand* 4 806 1950; IR: Jöberg *Acta Chem Scand* 4 798 1950].

Chloromethyl phenyl sulfide [7205-91-6] M 158.7, b 63°/0.1mm, 98°/12mm, 113-115°/20mm. Dissolve in CH₂Cl₂ or CCl₄ and dry over CaCl₂, or pass through a tube of CaCl₂ and fractionally distil using a fractionating column. *Harmful vapours*. It gives the *sulfone* (b 130°/1mm and m 53° from EtOH) on oxidation with permonophthalic acid. [*Justus Liebigs Ann Chem* 563 54 64 1949.]

***N*-(Chloromethyl)phthalimide** [17564-64-6] M 195.6, m 131-135°, 134-135°, 136.5°. Purified by recrystn from EtOAc or CCl₄ [*J Am Chem Soc* 70 2822 1948; Böhme et al. *Chem Ber* 92 1258 1959].

4-(Chloromethyl)pyridine hydrochloride [1822-51-1] M 164.0, m 170-175°, 172-173°, pK_{Est} ~5.6. Purified by recrystn from EtOH or EtOH-dry Et₂O. It melts between 171° and 175° and the clear melt resolidifies on further heating at 190° and turns red to black at 280° but does not melt again. The *picrate-hydrochloride* (prepared in EtOH) has m 146-147°. The free base is an oil, [Mosher and Tessieri *J Am Chem Soc* 73 4925 1951.]

2-Chloro-1-methylpyridinium iodide [14338-32-0] M 255.5, m 203-205°, 205-206°(dec), 207°. Purified by dissolving in EtOH and adding dry Et₂O. The solid is washed with Me₂CO and dried at

20°/0.35mm. Store in the dark. Attempted recrystn from Me₂CO-EtOH-pet ether (b 40-60°) caused some exchange of the Cl substituent by I. The *picrate* has *m* 106-107°, and the *perchlorate* has *m* 212-213°. [UV and solvolysis: Barlin and Benbow *J Chem Soc, Perkin Trans 2* 790 1974.]

1-Chloronaphthalene [90-13-1] *M* 162.6, *f* -2.3°, *b* 136-136.5°/20mm, 259.3°/760mm, *d* 1.194, *n* 1.6326. Washed with dilute NaHCO₃, then dried with Na₂SO₄ and fractionally distd under reduced pressure. Alternatively, before distn, it was passed through a column of activated alumina, or dried with CaCl₂, then distd from sodium. It can be further purified by fractional crystn by partial freezing or by crystn of its picrate to constant melting point (132-133°) from EtOH, and recovering from the picrate.

2-Chloronaphthalene [91-58-7] *M* 162.6, *m* 61°, *b* 264-266°. Crystd from 25% EtOH/water and dried under vacuum.

1-Chloro-2-naphthol [633-99-8] *M* 178.6, *m* 70°, *pK*_{Est} ~8.3. Cryst from pet ether. *Acetate* has *m* 42-43°.

2-Chloro-1-naphthol [606-40-6] *M* 178.6, *m* 64-65°, *pK*_{Est} ~7.9. Crystd from pet ether.

4-Chloro-1-naphthol [604-44-4] *M* 178.6, *m* 116-117°, 120-121°, *pK*²⁵ 8.86. Crystd from EtOH or chloroform.

6-Chloronicotinic acid [5326-23-8] *M* 157.6, *m* 190-193°, 198-199°(dec), *pK*²⁵ 4.22 (50% aq EtOH). Purified by recrystn from hot H₂O and is sublimed in a vacuum. [Pechmann and Welsch *Chem Ber* 17 2384 1884; Herz and Murty *J Org Chem* 26 122 1961.]

4-Chloro-2-nitroaniline [89-63-4] *M* 172.6, *m* 116-116.5°, *pK*²⁵ -0.99. Crystd from hot water or EtOH/water and dried for 10h at 60° under vacuum.

2-Chloro-4-nitrobenzamide [3011-89-0] *M* 200.6, *m* 172°. Crystd from EtOH.

2-Chloro-1-nitrobenzene [88-73-3] *M* 157.6, *m* 32.8-33.2°. Crystd from EtOH, MeOH or pentane (charcoal).

3-Chloro-1-nitrobenzene [121-73-3] *M* 157.6, *m* 45.3-45.8°. Crystd from MeOH or 95% EtOH (charcoal), then pentane.

4-Chloro-1-nitrobenzene [100-00-5] *M* 157.6, *m* 80-83°, 83.5-84°, *b* 113°/8mm, 242°/atm, *d*^{100.5} 1.2914. Crystd from 95% EtOH (charcoal) and sublimes in a vacuum. [Emmons *J Am Chem Soc* 76 3470 1954; Newman and Forres *J Am Chem Soc* 69 1221 1947.]

4-Chloro-7-nitrobenzofurazane (7-chloro-4-nitrobenzoxadiazole) [10199-89-0] *M* 199.6, *m* 96.5-97°, 97°, 99-100°. Wash the solid with H₂O and recrystallise from aqueous EtOH (1:1) as pale yellow needles. It sublimes in a vacuum [UV, NMR: Bolton, Gosh and Katritzky *J Chem Soc* 1004 1966].

1-Chloro-2-nitroethane [625-47-8] *M* 109.5, *b* 37-38°/20mm, *n* 1.4224, *n*²⁵ 1.4235. Dissolved in alkali, extracted with ether (discarded), then the aqueous phase was acidified with hydroxylamine hydrochloride, and the nitro compound fractionally distd under reduced pressure. [Pearson and Dillon *J Am Chem Soc* 75 2439 1953.]

2-Chloro-3-nitropyridine [5470-18-8] *M* 158.5, *m* 100-103°, 101-102°, 103-104° (sublimes), *pK*²⁰ -2.6. Forms needles from H₂O. Purified by continuous sublimation over a period of 2 weeks at 50-60°/0.1mm [Barlin *J Chem Soc* 2150 1964]. The *N-oxide* has *m* 99-100°(from CH₂Cl₂-Et₂O). [Taylor and Driscoll *J Org Chem* 25 1716 1960; Ochiai and Kaneko *Chem Pharm Bull Jpn* 8 28 1960.]

2-Chloro-5-nitropyridine [4548-45-2] M 158.5, m 108°, pK_{Est} ~-2.6. Crystd from *benzene or *benzene/pet ether.

3-Chloroperbenzoic acid [937-14-4] M 172.6, m 92-94°(dec), pK²⁵ 7.57. Recrystd from CH₂Cl₂ [Traylor and Mikzta *J Am Chem Soc* 109 2770 1987]. Peracid of 99+% purity can be obtained by washing commercial 85% material with phosphate buffer pH 7.5 and drying the residue under reduced pressure. Alternatively the peracid can be freed from *m*-chlorobenzoic acid by dissolving 50g/L of *benzene and washing with an aq soln buffered at pH 7.4 (NaH₂PO₄/NaOH) (5 x 100mL). The organic layer was dried over MgSO₄ and carefully evaporated under vacuum. *Necessary care should be taken in case of EXPLOSION.* The solid was recrystd twice from CH₂Cl₂/Et₂O and stored at 0° in a plastic container as glass catalyses the decomposition of the peracid. The acid is assayed iodometrically. [*J Org Chem* 29 1976 1964; Bortolini et al. *J Org Chem* 52 5093 1987.]

2-Chlorophenol [95-57-8] M 128.6, m 8.8°, b 61-62°/10mm, 176°/atm, pK²⁵ 8.34. Passed at least twice through a gas chromatograph column. Also purified by fractional distn.

3-Chlorophenol [108-43-0] M 128.6, m 33°, b 44.2°/1mm, 214°/atm, pK²⁵ 9.13. Could not be obtained solid by crystn from pet ether. Purified by distn under reduced pressure.

4-Chlorophenol [106-48-9] M 128.6, m 43°, 100-101°/10mm, pK²⁵ 9.38. Distd, then crystd from pet ether (b 40-60°) or hexane, and dried under vacuum over P₂O₅ at room temp. It has pK_a 9.38 at 20° in water. [Bernasconi and Paschalis *J Am Chem Soc* 108 2969 1986.]

Chlorophenol Red (3,3'-dichlorophenolsulfonephthalein) [4430-20-0] M 423.3, m dec on heating, λ_{max} 573nm, pK²⁵ 5.96. Crystd from glacial acetic acid. It is yellow at pH 4.8 and violet at pH 6.7.

4-Chlorophenoxyacetic acid [122-88-3] M 186.6, m 157°, pK²⁰ 3.00. Crystd from EtOH.

α-4-Chlorophenoxypropionic acid [3307-39-9] M 200.6, m 116°, pK_{Est} ~3.2. Crystd from EtOH.

β-4-Chlorophenoxypropionic acid [3284-79-5] M 200.6, m 138°, pK_{Est} ~4.2. Crystd from EtOH.

3-Chlorophenylacetic acid [1878-65-5] M 170.6, m 74°, pK²⁵ 4.11. Crystd from EtOH/water, or as needles from *C₆H₆ or H₂O (charcoal). The *acid chloride* (prepared by boiling with SOCl₂) has b 127-129°/15mm. [Dippy and Williams *J Chem Soc* 161 1934; Misra and Shukla *J Indian Chem Soc* 28 480 1951.]

4-Chlorophenylacetic acid [1878-66-6] M 170.6, m 105°, 106°, pK²⁵ 4.12. Same as for 3-chlorophenylacetic acid.

4-Chloro-1-phenylbutan-1-one [939-52-6] M 182.7, m 19-20°, b 134-137°/5mm, d₄²⁰ 1.149, n_D²⁰ 1.55413. Fractionate several times using a short column. It can be recrystd from anhydrous pet ether at -20° as glistening white rosettes and filtered at 0° and dried in a vacuum desiccator over H₂SO₄. The *semicarbazone* has m 136-137°. [*J Am Chem Soc* 46 1882 1924, 51 1174 1929, Hart and Curtis *J Am Chem Soc* 79 931 1957.]

1-(2-Chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane (Mitotane, *o,p'*-DDD) [53-19-0] M 320.1, m 75.8-76.8°, 76-78°. Purified by recrystallisation from pentane and from MeOH or EtOH. It is sol in isoctane and CCl₄. [Haller et al. *J Am Chem Soc* 67 1600 1945.]

3-(4-Chlorophenyl)-1,1-dimethylurea (monuron) [150-68-5] M 198.7, m 171°. Crystd from MeOH.

Previous Page

4-Chloro-1,2-phenylenediamine [95-83-0] M 142.6, m 69-70°, pK_1^{25} -0.27 (aq H_2SO_4), pK_2^{25} 3.35 (3.67). Recrystd from pet. ether.

4-Chlorophenyl isocyanate [104-12-1] M 153.6, m 28-31°, 31-32°, 32°, 32.5°, b 80.6-80.9°/9.5mm, 115-117°/45mm. Purified by recrystn from pet ether (b 30-40°) or better by fractional distn. **TOXIC irritant.**

4-Chlorophenyl isothiocyanate [2131-55-7] M 169.6, m 44°, 43-45°, 45°, 46°, 47°, b 110-115°/4mm, 135-136°/24mm. Check the IR first. Triturate with pet ether (b 30-60°) and decant the solvent. Repeat 5 times. The combined extracts are evap under reduced press to give almost pure compound as a readily crystallisable oil with a pleasant anise odour. It can be recrystd from the minimum vol of EtOH at 50° (do not boil too long in case it reacts). It can be purified by vac distn. **IRRITANT.** [*Org Synth Coll Vol V* 223 1973.]

4-Chlorophenyl 2-nitrobenzyl ether [109669-56-9] M 263.7, m 69°. Crystd from EtOH.

4-Chlorophenyl 4-nitrobenzyl ether [5442-44-4] M 263.7, m 102°. Crystd from EtOH.

9-Chloro-9-phenylxanthene (Pixyl chloride) [42506-03-6] M 292.8, m 105-106°. Possible impurity is 9-hydroxy-9-phenylxanthene. If material contains a lot of the hydroxy product then boil 10g in $CHCl_3$ (50mL) with redistd acetyl chloride (1mL) until liberation of HCl is complete. Evapn leaves the chlorophenylxanthene as the hydrochloride which on heating with *benzene loses HCl; and on adding pet ether prisms of chlorophenylxanthene separate and contain 0.5mol of *benzene. The *benzene-free compound is obtained on drying and melts to a colourless liquid. [*Justus Liebigs Ann Chem* 370 142 1909.] The 9-phenylxanthyl group is called pixyl. [*J Chem Soc, Chem Commun* 639 1978.]

Chlorophyll a [479-61-8] M 983.5, m 117-120°, 150-153°, 178-180° (sinters at ~150°), $[\alpha]_D^{20}$ -262° (Me_2CO). Forms green crystals from Me_2CO , $Et_2O + H_2O$, $Et_2O + hexane + H_2O$ or $Et_2O + pentane + H_2O$. It is sparingly soluble in MeOH and insol in pet ether. In alkaline soln it gives a blue-green colour with deep red fluorescence. A very crude chlorophyll mixture has been purified by chromatography on low melting polyethylene (MI 0.044; 'Dow' melting index MI <2) and developed with 70% aq Me_2CO . The order of effluent from the bottom of the column is: xanthophylls, chlorophyll *b*, chlorophyll *a*, phaeophytins and carotenes. A mixture of chlorophylls *a* and *b* is best separated by chromatography on sugar and the order is chlorophyll *b* elutes first followed by chlorophyll *a*. To an Me_2CO-H_2O soln of chlorophylls 200mL of iso-octane is added and the mixt shaken in a separating funnel and the H_2O is carefully removed. The iso-octane layer is dried (Na_2SO_4) and applied to a glass column (5cm diameter) dry packed with 1000mL of powdered sucrose which has been washed with 250mL of iso-octane. Elution with 0.5% of isopropanol in iso-octane gives chlorophyll *a*. Keeping the eluate overnight at 0° yields micro crystals which are collected by filtration or centrifugation (Yield 40mg). UV_{EtOH} has λ_{max} 660, 613, 577, 531, 498, 429 and 409 nm. [Anderson and Calvin *Nature* 194 285 1962; Stoll and Weidemann *Helv Chim Acta* 16 739 757 1933; NMR: Katz et al. *J Am Chem Soc* 90 6841 1968, 85 3809 1963 for *a* and *b*; ORD: Inhoffen et al. *Justus Liebigs Ann Chem* 704 208 1967; Willstätter and Isler *Justus Liebigs Ann Chem* 390 269, 233 1912.]

Chlorophyll b [519-62-0] M 907.52, sinters at 86-92°, sinters at 170°, dec at 160-170°, m 183-185°, 190-195°, $[\alpha]_D^{20}$ -267° ($Me_2CO + MeOH$), $[\alpha]_{720}^{25}$ -133° ($MeOH + Pyridine$ 95:5). See purification of chlorophyll *a*, and is separated from "a" by chromatography on sucrose [UV, IR: Stoll and Weidemann *Helv Chim Acta* 42 679, 681 1959]. It forms red-black hexagonal bipyramids or four sided plates from dilute EtOH and has been recrystd from $CHCl_3-MeOH$. It is soluble in MeOH, EtOH, EtOAc and insoluble in pet ether. [*J Am Chem Soc* 88 5037 1966.]

Chloropicrin (trichloronitromethane) [76-06-2] M 164.5, b 112°. Dried with $MgSO_4$ and fractionally distd. **EXTREMELY NEUROTOXIC, use appropriate precautions.**

RS-2-Chloropropionic acid [598-78-7] M 108.5, b 98°/3mm, d 1.182, n 1.453 pK^{25} 2.89. Dried with P_2O_5 and fractionally distd under vacuum.

S-(-)-2-Chloropropionic acid [29617-66-1] M 108.5, b 77°/10mm, 80.7°/10mm, 185-188°/atm, d_4^{25} 1.2485, n_D^{25} 1.436, $[\alpha]_D^{25}$ -14.6° (neat). Purified by twice fractionating through a 115cm Podbielniak column (calcd 50 theoretical plates at atm pressure, see p. 141) using a take-off ratio of 1:5. This *acid chloride* is prepared by dissolving the acid in SOCl_2 adding a few drops of PCl_3 , refluxing and then distilling through a 30 cm column, b 53°/100mm, $[\alpha]_D^{25}$ -4.6° (neat), d_4^{25} 1.2689, n_D^{25} 1.4368. [Fu et al. *J Am Chem Soc* 76 6954 1954].

3-Chloropropionic acid [107-94-8] M 108.5, m 41°, pK^{25} 4.08. Crystd from pet ether or *benzene.

3-Chloropropyl bromide (1-bromo-3-chloropropane) [109-70-6] M 157.5, b 142-145°, n_D^{25} 1.4732. Washed with conc H_2SO_4 , water, 10% Na_2CO_3 soln, water again and then dried with CaCl_2 and fractionally distd just before use [Akagi, Oae and Murakami *J Am Chem Soc* 78 4034 1956].

6-Chloropurine [87-42-3] M 154.6, m 179°(dec), pK_1^{20} 0.45, pK_2^{20} 7.88. Crystd from water.

2-Chloropyrazine [14508-49-7] M 114.5, b 62-63°/31mm, 153-154°/atm, d_4^{20} 1.302, n_D^{26} 1.535, $\text{pK}_{\text{Est}} < 0$. Fractionally distil through a short column packed with glass helices. It has a penetrating mildly pungent odour with a high vapour pressure at room temperature. [Erickson and Spoerri *J Am Chem Soc* 68 400 1946; *J Org Chem* 28 1682 1963.]

2-Chloropyridine [109-09-1] M 113.6, b 49.0°/7mm, d 1.20, n 1.532, pK^{20} 0.49 (0.72). Dried with NaOH for several days, then distd from CaO under reduced pressure.

3-Chloropyridine [626-60-8] M 113.6, b 148°, d 1.194, n 1.5304, pK^{25} 2.84. Distd from KOH pellets.

4-Chloropyridine [626-61-9] M 113.6, b 85-86°/100mm, 147-148°/760mm, pK^{20} 3.84. Dissolved in distilled water and excess of 6M NaOH was added to give pH 12. The organic phase was separated and extracted with four volumes of diethyl ether. The combined extracts were filtered through paper to remove water and the solvent evaporated. The dark brown residual liquid was kept under high vacuum [Vaidya and Mathias *J Am Chem Soc* 108 5514 1986]. It can be distd but readily darkens and is best kept as the *hydrochloride* [7379-35-3] M 150.1, m 163-165°(dec).

2-Chloropyrimidine [1722-12-9] M 114.5, m 63-65°, 66°, b 91°/26mm, pK^{20} -1.90. It has been recrystd from * C_6H_6 , pet ether or a mixture of both. It sublimes at 50°/18mm and can be distd in a vacuum. [IR: Short and Thompson *J Chem Soc* 168 1952; Boarland and McOmie *J Chem Soc* 1218 1951.]

2-Chloroquinoline [612-62-4] M 163.6, m 34°, b 147-148°/15mm, d_4^{35} 1.235, n_D^{25} 1.629, $\text{pK}_{\text{Est}} \sim 0.3$. Purified by crystn of its picrate to constant melting point (123-124°) from *benzene, regenerating the base and distilling under vacuum [Cumper, Redford and Vogel *J Chem Soc* 1183 1962]. 2-Chloroquinoline can be crystd from EtOH. Its *picrate* has m 122° (from EtOH).

4-Chloroquinoline [611-35-8] M 163.6, m 29-32°, 31°, b 130°/15mm, 261°/744mm, pK 3.72. Possible impurities include the 2-isomer. Best purified by converting to the *picrate* (m 212-213° dec) in EtOH and recryst from EtOH (where the *picrate* of the 2-chloroquinoline stays in soln) or EtOAc. The *picrate* is decomposed with 5% aqueous NaOH, extracted in CHCl_3 , washed with H_2O , dried (MgSO_4), evapd and distd in a vacuum. It can be steam distd from slightly alkaline aqueous solns, the aqueous distillate is extracted with Et_2O , evaporated and distd. The distillate solidifies on cooling. [Bobránski *Chem Ber* 71 578 1938.]

8-Chloroquinoline [611-33-6] M 163.6, b 171-171.5°/24mm, d 1.278, n 1.644, pK^{25} 3.12. Purified by crystn of its ZnCl_2 complex (m 228°) from aqueous EtOH.

4-Chlororesorcinol [95-88-5] M 144.6, m 105°, $\text{pK}_{\text{Est}(1)} \sim 9.2$, $\text{pK}_{\text{Est}(2)} \sim 10.1$. Crystd from boiling CCl_4 (10g/L, charcoal) and air dried.

5-Chlorosalicylaldehyde [635-93-8] M 156.6, m 98.5-99°. Steam distd, then crystd from aq EtOH.

N-Chlorosuccinimide [128-09-6] M 133.5, m 149-150°. Rapidly crystd from *benzene, or glacial acetic acid and washed well with water then dried *in vacuo*. [Phillips and Cohen *J Am Chem Soc* 108 2023 1986.]

2-Chlorothiophene (2-thienyl chloride) [96-43-5] M 118.6, b 126-128°, d 1.285, n 1.551. Purified by fractional distn at atmospheric pressure or by gas chromatography.

8-Chlorotheophylline [85-18-7] M 214.6, m 311°(dec), $pK_{Est(1)} \sim 5.4$, $pK_{Est(2)} \sim 9.1$. Crystd from H₂O.

4-Chlorothiophenol [106-54-7] M 144.6, m 51-52°, pK^{25} 6.14. Recrystd from aqueous EtOH [D'Sousa et al. *J Org Chem* 52 1720 1987].

2-Chlorotoluene [95-49-8] M 126.6, b 159°, d 1.083, n 1.5255. Dried for several days with CaCl₂, then distd from Na using a glass helices-packed column.

3-Chlorotoluene [108-41-8] M 126.6, m -48°, b 161-163°, d 1.072, n 1.522. Purified as for 2-chlorotoluene above.

4-Chlorotoluene [106-43-4] M 126.6, f 7.2°, b 162.4°, d 1.07, n 1.521. Dried with BaO, fractionally distd, then fractionally crystd by partial freezing.

2-Chlorotriethylamine hydrochloride [869-24-9] M 172.1, m 208-210°, $pK_{Est} \sim 8.6$ (free base). Crystd from absolute MeOH (to remove highly coloured impurities).

Chlorotrifluoroethylene [79-38-9] M 116.5, b -26 to -24°. Scrubbed with 10% KOH soln, then 10% H₂SO₄ soln to remove inhibitors, and dried. Passed through silica gel.

Chlorotrifluoromethane [75-72-9] M 104.5, m -180°, b -81.5°. Main impurities were CO₂, O₂, and N₂. The CO₂ was removed by passage through saturated aqueous KOH, followed by conc H₂SO₄. The O₂ was removed using a tower packed with activated copper on Kieselguhr at 200°, and the gas dried over P₂O₅.

Chlorotriphenylmethane see triphenylmethyl chloride (trityl chloride).

5-Chlorouracil (5-chloro-2,4(6)-dihydroxypyrimidine) [1820-81-1] M 146.5, m 314-418° dec, 324-325° dec, pK_1^{25} 7.95, $pK_2^{25} > 13$. Recrystallised from hot H₂O (4g/500mL) using charcoal. [McOmie et al. *J Chem Soc* 3478 1955; West and Barrett *J Am Chem Soc* 76 3146 1954.]

5 β -Cholanic acid [25312-65-6] M 360.6, m 164-165°, $[\alpha]_D^{14} + 21.7^\circ$ (CHCl₃), $pK_{Est} \sim 4.9$. Crystd from EtOH. The *Ethyl ester* has m 93-94° (from 80% EtOH), b 273°/12mm, $[\alpha]_D^{20} + 21^\circ$ (CHCl₃).

Cholanthrene (1,2-dihydrobenz[j]aceanthrylene) [479-23-2] M 254.3, m 173°. Crystd from *benzene/diethyl ether.

5 α -Cholestane [481-21-0] M 372.7, m 80°, $[\alpha]_{546}^{20} + 29.5^\circ$ (c 2, CHCl₃). Crystd from diethyl ether/EtOH.

5 α -Cholestan-3 β -ol [80-97-7] M 388.7, m 142-143°(monohydrate), $[\alpha]_{546}^{20} + 28^\circ$ (c 1, CHCl₃), $[\alpha]_D + 27.4^\circ$ (in CHCl₃). Crystd from EtOH or slightly aqueous EtOH, or MeOH. [Mizutani and Whitten *J Am Chem Soc* 107 3621 1985.]

Cholest-2-ene [15910-23-3] M 370.6, m 75-76°, $[\alpha]_D^{24} + 64^\circ$. Recrystd from MeOH or diethyl ether/acetone. [Berzbrester and Chandran *J Am Chem Soc* 109 174 1987.]

Cholesterol [57-88-5] M 386.7, m 148.9-149.4°, $[\alpha]_D^{25}$ -35° (hexane). Crystd from ethyl acetate, EtOH or isopropyl ether/MeOH. [Hiromitsu and Kevan *J Am Chem Soc* 109 4501 1987.] For extensive details of purification through the dibromide, see Fieser [*J Am Chem Soc* 75 5421 1953] and Schwenk and Werthessen [*Arch Biochem Biophys* 40 334 1952], and by repeated crystn from acetic acid, see Fieser [*J Am Chem Soc* 75 4395 1953].

Cholesteryl acetate [604-35-3] M 428.7, m 112-115°, $[\alpha]_{546}^{20}$ -51° (c 5, CHCl₃). Crystd from *n*-pentanol.

Cholesteryl myristate [1989-52-2] M 597.0. Crystd from *n*-pentanol. Purified by column chromatography with MeOH and evaporated to dryness. Recrystd and finally, dried in vacuum over P₂O₅. [Malanik and Malat *Anal Chim Acta* 76 464 1975].

Cholesteryl oleate [303-43-5] M 651.1, m 48.8-49.4°. Purified by chromatography on silica gel.

Cholic acid [81-25-4] M 408.6, m 198-200°, $[\alpha]_{546}$ +41° (c 0.6, EtOH), pK²⁰ 4.98. Crystd from EtOH. Dried under vacuum at 94°.

Choline chloride [67-48-1] M 139.6, m 302-305°(dec). *Extremely deliquescent*. Purity checked by AgNO₃ titration or by titration of free base after passage through an anion-exchange column. Crystd from absolute EtOH, or EtOH-diethyl ether, dried under vacuum and stored in a vacuum desiccator over P₂O₅ or Mg(ClO₄)₂.

4-Chromanone [491-37-2] M 148.2, m 35-37°, 39°, 41°, b 92-93°/3mm, 130-132°/15mm, 160°/50mm. It has been recryst from pet ether, or purified by dissolving in *C₆H₆ washing with H₂O, drying (MgSO₄), evaporate and dist in a vacuum, then recryst the residue. The liquid has a pleasant lemon-like odour. The *semicarbazone* has m 227°. [Loudon and Razdan *J Chem Soc* 4299 1954.] The *oxime* is prepared from 3g of chromanone, 3g NH₂OH.HCl in EtOH (50mL), 6g K₂CO₃ and refluxed on a water bath for 6h. The soln is poured into H₂O, the solid is filtered off, dried and dissolved in hot *C₆H₆ which on addition of pet ether yields the oxime as glistening needles m 140°. Decomposition of this gives very pure chromanone. The *benzal derivative* is prepared from 3g of chromanone, 4g PhCHO in 50mL EtOH, heated to boiling, 10mL of conc HCl are added dropwise and set aside for several days. The derivative separates and is recrystd from EtOH to give yellow needles, m 112° [*J Am Chem Soc* 45 2711 1923]. Reaction with Pb(OAc)₄ yields the *3-acetoxy derivative* m 74° (from pet ether + trace of EtOAc) [Cavill et al. *J Chem Soc* 4573 1954].

Chrysene [218-01-9] M 228.3, m 255-256°. Purified by chromatography on alumina from pet ether in a darkened room. Its soln in *C₆H₆ was passed through a column of decolorising charcoal, then crystd by concentration of the eluate. Also purified by crystn from *C₆H₆ or *C₆H₆-pet ether, and by zone refining. [Gorman et al. *J Am Chem Soc* 107 4404 1985]. It was freed from 5*H*-benzo[*b*]carbazole by dissolving in *N,N*-dimethylformamide and successively adding small portions of alkali and iodomethane until the fluorescent colour of the carbazole anion no longer appeared when alkali was added. The chrysene (and alkylated 5*H*-benzo[*b*]carbazole) separated on addition of water. Final purification was by crystn from ethylcyclohexane and from 2-methoxyethanol [Bender, Sawicki and Wilson *Anal Chem* 36 1011 1964]. It can be sublimed in a vacuum.

Chrysoidine G (4-phenylazo-1,3-benzenediamine monohydrochloride) [532-82-1] M 248.7, m 118-118.5°, pK₁ 3.32, pK₂ 5.21. Red-brown powder which is recrystd from H₂O. It gives a yellow soln in conc H₂SO₄ which turns orange on dilution. Its solubility at 15° is 5.5% (H₂O), 4.75% (EtOH), 6.0% (cellosolve), 9.5% (ethylene glycol), 0.005% (xylene) and insol in *C₆H₆. The *hydroiodide* has m 184° (from EtOH) and the *picrate* forms red needles m 196°. [*Bull Chem Soc Jpn* 31 864 1958; *Chem Ber* 10 213 1877.]

1,8-Cineole (1,8-epoxy-*p*-menthane) [470-82-6] M 154.2, f 1.3°, b 176.0°, d 0.9251. See eucaliptol on p. 242.

trans-Cinnamaldehyde [14271-10-9] M 132.2, m -4° , -7.5° , -9° , b $80^{\circ}/0.4\text{ mm}$, $85.8^{\circ}/1.1\text{ mm}$, $125-128^{\circ}/11\text{ mm}$, $152.2^{\circ}/40\text{ mm}$, $163.7^{\circ}/60\text{ mm}$, $199.3^{\circ}/200\text{ mm}$, $246^{\circ}/760\text{ mm}$ dec, d_4^{20} 1.0510, n_D^{20} 1.623. Purified by steam distn (sol 1 in 700 parts H_2O) followed by distn *in vacuo*. The *cis*-isomer has b $67-69^{\circ}/40\text{ mm}$ and d_4^{20} 1.0436 and n_D^{20} 1.5937. The *trans*-semicarbazone has m 210° dec from CHCl_3 -MeOH (*cis*-semicarbazone has m 196°); the *trans*-phenylsemicarbazone has m 177° from CHCl_3 -MeOH (the *cis*-phenylsemicarbazone has m 146°); the *trans*-2,4-dinitrophenylhydrazone has m 250° dec from MeOH as the *cis*-isomer [Gamboni et al. *Helv Chim Acta* **38** 255 1955; Peine *Chem Ber* **17** 2117 1884; *J Org Chem* **26** 4814 1961; *J Am Chem Soc* **86** 198 1964].

cis-Cinnamic acid (Z-3-phenyl-2-propenoic acid) [102-94-3] M 148.2, m 68° (for *allo*-form), pK²⁵ 3.93. The *cis*-acid is prepared by catalytic reduction of phenylpropionic acid and after distn in high vacuum at $\sim 95^{\circ}$ gives the most stable *allo*-isomer m 68° . Recrystn from pet ether yields Liebermann's *iso*-cinnamic acid m 58° . When the *allo*-acid (m 68°) is heated at 20° above its melting point in a sealed capillary for 0.5h and allowed to cool slowly Erlenmyer's *iso*-cinnamic acid m 42° is formed. This form can also be obtained in larger amounts by heating the *allo*-acid at 80° for 3h and on cooling it remains liquid for several weeks but gives the 42° acid on inoculation with the crystals from the capillary tube. This form is unchanged in 6 weeks when kept in a dark cupboard. All three forms have the same pK values and the same rate of bromination. There is also a very labile form with m 32° . [Liebermann, *Chem Ber* **26** 1572 1893; Claisen and Crismer *Justus Liebigs Ann Chem* **218** 135 1883; Robinson and James *J Chem Soc* 1453 1933; Berthoud and Urech *Helv Chim Acta* **13** 437 1930; McCoy and McCoy *J Org Chem* **33** 2354 1968.]

trans-Cinnamic (E-3-phenyl-2-propenoic) acid [140-10-3; 621-82-9 for E-Z mixture] M 148.2, m $134.5-135^{\circ}$, pK²⁵ 4.42 (4.50). Crystd from *benzene, CCl_4 , hot water, water/EtOH (3:1), or 20% aqueous EtOH. Dried at 60° *in vacuo*. Steam volatile.

Cinnamic anhydride [538-56-7] M 278.4, m 136° . Crystd from C_6H_6 or toluene/pet ether (b $60-80^{\circ}$).

N-Cinnamoyl-N-phenylhydroxylamine [7369-44-0] M 239.3, m $158-163^{\circ}$. Recrystd from EtOH.

Cinnamyl alcohol [104-54-1] M 134.2, m 33° , b $143.5^{\circ}/14\text{ mm}$, λ_{max} 251nm (ϵ 18,180M⁻¹cm⁻¹). Crystd from diethyl ether/pentane.

Cinnoline [253-66-7] M 130.2, m 38° , pK²⁰ 2.37. Crystd from pet ether. Kept under N_2 in sealed tubes in the dark at 0° .

Citraconic acid [498-23-7] M 130.1, m 91° , pK₁²⁵ 2.2, pK₂²⁵ 5.60 (*cis*). Steam distd and crystd from EtOH/ligroin.

Citraconic anhydride [616-02-4] M 112.1, m $8-9^{\circ}$, b $47^{\circ}/0.03\text{ mm}$, $213^{\circ}/760\text{ mm}$, d_4^{20} 1.245, n_D^{20} 1.472. Possible contamination is from the acid formed by hydrolysis. If the IR has OH bands then reflux with Ac_2O for 30 min, evaporate then distil the residue in a vacuum; otherwise distil in a vacuum. Store in a dry atmosphere. [*Biochem J* **191** 269 1980.]

Citrazinic acid (2,6-dihydroxyisonicotinic acid) [99-11-6] M 155.1, m $>300^{\circ}$, pK₁ 3.0, pK₂ 4.76. Yellow powder with a greenish shade, but is white when ultra pure and turns blue on long standing. It is insoluble in H_2O but slightly soluble in hot HCl and soluble in alkali or carbonate solutions. Purified by precipitation from alkaline solutions with dilute HCl, and dry in a vacuum over P_2O_5 . The *ethyl ester* has m 232° (evacuated tube) and a pKa of 4.81 in $\text{MeOCH}_2\text{CH}_2\text{OH}$ [IR: Pitha *Coll Czech Chem Comm* **28** 1408 1963].

Citric acid (H_2O) [5949-29-1; 77-92-9 (*anhydr*)] M 210.1, m $156-157^{\circ}$, 153° (*anhyd*), pK₁²⁵ 2.96, pK₂²⁵ 4.38, pK₃²⁵ 5.68. Crystd from water.

Citronellal (3,7-dimethyloctan-6-al) [*R*(+): 2385-77-5; *S*(-): 5949-05-3] **M 154.3, b 67°/4mm, 89°/14mm, 104-105°/21mm, 207°/760mm, $[\alpha]_{546}^{20}$ (+) and (-) 20°, $[\alpha]_{\text{D}}^{20}$ (+) and (-) 16.5° (neat).** Fractionally distd. Alternatively extracted with NaHSO₃ solution, washed with Et₂O then acidified to decompose the bisulfite adduct and extracted with Et₂O, dried (Na₂SO₄), evaporated and distd. Check for purity by hydroxylamine titration. The ORD in MeOH (c 0.167) is: $[\alpha]_{700} +9^\circ$, $[\alpha]_{589} +11^\circ$, $[\alpha]_{275} +12^\circ$ and $[\alpha]_{260} 12^\circ$. The *semicarbazone* has *m* 85°, and the *2,4-dinitrophenylhydrazone* has *m* 79-80°. [IR: *J Chem Soc* 3457 1950; ORD: Djerassi and Krakower *J Am Chem Soc* 81 237 1959.]

β-Citronellene (2,6-dimethylocta-2,7-diene) [*S*(+): 2436-90-0; *R*(-): 10281-56-8] **M 138.3, b 153-154°/730mm, 155°/atm, $d_4^{22} 0.757$, $n_{\text{D}}^{22} 1.431$, $[\alpha]_{546}^{20}$ (+) and (-) 13°, $[\alpha]_{546}^{20}$ (+) and (-) 10° (neat).** Purified by distillation over Na three times and fractionation. [(-) Arigoni and Jeager *Helv Chim Acta* 37 881 12954; (+) Eschenmoser and Schinz *Helv Chim Acta* 33 171 1950.]

β-Citronellol (3,7-dimethyloctan-6-ol) [*R*(+): 11171-61-9; *S*(-): 106-22-9] **M 156.3, b 47°/1mm, 102-104(110°)/10mm, 112-113°/12mm, 221-224°/atm, 225-226°/atm, $d_4^{24} 0.8551$, $n_{\text{D}}^{24} 1.4562$, $[\alpha]_{546}^{20}$ (+) and (-) 6.3°, $[\alpha]_{\text{D}}^{20}$ (+) and (-) 5.4° (neat).** Purified by distn through a cannon packed (Ni) column and the main cut collected at 84°/14mm and redistd. Also purified *via* the benzoate. [IR: Eschenazi *J Org Chem* 26 3072 1961; *Bull Soc Chim Fr* 505 1951.]

S-Citrulline (2-amino-5-ureidopentanoic acid) [372-75-8] **M 175.2, m 222°, $[\alpha]_{\text{D}}^{20} +24.2^\circ$ (in 5M HCl), $\text{pK}_{\text{Est}(1)}^{25} 9.71$.** Likely impurities are arginine, and ornithine. Crystd from water by adding 5 volumes of EtOH. Also crystd from water by addn of MeOH.

Clofazimine [2-(4-chloroanilino)-3-isopropylimino-5-(4-chlorophenyl)-3,4-dihydrophenazine] [2030-63-9] **M 473.5, m 210-212°.** Recrystd from acetone.

Coenzyme Q₀ (2,3-Dimethoxy-5-methyl-1,4-benzoquinone, 3,4-dimethoxy-2,5-toluquinone, fumigatin methyl ether), colchicine and colchicoside see entries in Chapter 6.

Conessine [546-06-5] M 356.6, m 125°, 127-128.5°, $[\alpha]_{\text{D}}^{20} -1.9^\circ$ (in CHCl₃) and +25.3° (in EtOH), $\text{pK}_{\text{Est}(1)} \sim 10.4$, $\text{pK}_{\text{Est}(2)} \sim 10.7$. Crystd from acetone. The *dihydrochloride* has *m* >340° and $[\alpha]_{\text{D}}^{20} +9.3^\circ$ (c 0.9, H₂O).

Coniferyl alcohol [4-hydroxy-3-methoxy-cinnamyl alcohol, 3-(4-hydroxy-3-methoxyphenyl)-2-propen-1-ol] [458-35-5] **M 180.2, m 73-75°, b 163-165°/3mm, $\text{pK}_{\text{Est}}^{25} 9.54$.** It is soluble in EtOH and insoluble in H₂O. It can be recrystd from EtOH and distd in a vacuum. It polymerises in dilute acid. The *benzoyl derivative* has *m* 95-96° (from pet ether), and the *tosylate* has *m* 66°. [Derivatives: Freudenberg and Achtzehn *Chem Ber* 88 10 1955; UV: Herzog and Hillmer *Chem Ber* 64 1288 1931.]

Congo Red [573-58-0] M 696.7, λ_{max} 497nm, $\text{pK}_2^{28} 4.19$. Crystd from aq EtOH (1:3). Dried in air.

(-)-α-Copaene {*1R,2S,6S,7S,8S*-8-isopropyl-1,3-dimethyltricyclo[4.4.0.0^{2,7}]dec-3-ene} [3856-25-5] **M 204.4, b 119-120°/10mm, 246-251°, d 0.908, n 1.489, $[\alpha]_{\text{D}}^{20} -6.3^\circ$ (c 1.2, CHCl₃).** Purified by distn, preferably under vacuum.

4,5-Coprosten-3-ol (cholest-4-ene-3β-ol) [517-10-2] **M 386.7, m 132°.** Crystd from MeOH/diethyl ether.

Coprosterol (5α-cholestan-3β-ol, dihydrocholesterol) [80-97-7] **M 388.7, m 101°, 139-140°, $[\alpha]_{\text{D}}^{20} +24^\circ$ (c 1, CHCl₃).** See entry on p. 169.

Coronene [191-07-1] M 300.4, m 438-440°, λ_{max} 345nm (log ε 4.07). Crystd from *benzene or toluene, then sublimed in vacuum.

Cortisol, corticosterone, cortisone and cortisone-21-acetate see entries in Chapter 6.

Coumalic acid (2-pyrone-5-carboxylic acid) [500-05-0] M 140.1, m 205-210°(dec), pK_{Est} ~0. Crystd from MeOH. *Methyl ester*, from pet ether, has m 74-74° and b 178-180°/60 mm.

Coumarin [91-64-5] M 146.2, m 68-69°, b 298°, pK -4.97 (aq H₂SO₄). Crystd from ethanol or water and sublimed *in vacuo* at 43° [Srinivasan and deLevie *J Phys Chem* 91 2904 1987].

Coumarin-3-carboxylic acid [531-81-7] M 190.2, m 188°(dec), pK_{Est} ~1.5. Crystd from water.

Creatine (H₂O) and **creatinine** see entries in Chapter 6.

***o*-Cresol** [95-48-7] M 108.1, m 30.9°, b 191°, n⁴¹ 1.536, n⁴⁶ 1.534, pK²⁵ 10.22. Can be freed from *m*- and *p*-isomers by repeated fractional distn. Crystd from *benzene by addition of pet ether. Fractional crystd by partial freezing of its melt.

***m*-Cresol** [108-39-4] M 108.1, f 12.0°, b 202.7°, d 1.034, n 1.544, pK²⁵ 10.09. Separation of the *m*- and *p*-cresols requires chemical methods, such as conversion to their sulfonates [Brüchner *Anal Chem* 75 289 1928]. An equal volume of H₂SO₄ is added to *m*-cresol, stirred with a glass rod until soln is complete. Heat for 3h at 103-105°. Dilute carefully with 1-1.5 vols of water, heat to boiling point and steam distil until all unsulfonated cresol has been removed. Cool and extract residue with ether. Evaporate the soln until the boiling point reaches 134° and steam distil off the *m*-cresol. Another purification involves distn, fractional crystn from the melt, then redistn. Freed from *p*-cresol by soln in glacial acetic acid and bromination by about half of an equivalent amount of bromine in glacial acetic acid. The acetic acid was distd off, then fractional distn of the residue under vac gave bromocresols from which 4-bromo-*m*-cresol was obtained by crystn from hexane. Addn of the bromocresol in glacial acetic acid slowly to a reaction mixture of HI and red phosphorus or (more smoothly) of HI and hypophosphorus acid, in glacial acetic acid, at reflux, removed the bromine. After an hour, the soln was distd at atmospheric pressure until layers were formed. Then it was cooled and diluted with water. The cresol was extracted with ether, washed with water, NaHCO₃ soln and again with water, dried with a little CaCl₂ and distd [Baltzly, Ide and Phillips *J Am Chem Soc* 77 2522 1955].

***p*-Cresol** [106-44-5] M 108.1, m 34.8°, b 201.9°, n⁴¹ 1.531, n⁴⁶ 1.529, pK²⁵ 10.27. Can be separated from *m*-cresol by fractional crystn of its melt. Purified by distn, by pptn from *benzene soln with pet ether, and *via* its benzoate, as for phenol. Dried under vacuum over P₂O₅. Has also been crystd from pet ether (b 40-60°) and by conversion to sodium *p*-cresoxyacetate which, after crystn from water was decomposed by heating with HCl in an autoclave [Savard *Ann Chim (Paris)* 11 287 1929].

***o*-Cresolphthalein complexon** [2411-89-4] M 636.6, m 186°(dec), λ_{max} 575nm, pK₁ 2.2, pK₂ 2.9, pK₃ 7.0, pK₄ 7.8, pK₅ 11.4, pK₆ 12.0. *o*-Cresolphthalein (a complexone precursor without the two bis-carboxymethylamino groups) is a contaminant and is one of the starting materials. It can be removed by dissolving the reagent in water and adding a 3-fold excess of sodium acetate and fractionally precipitating it by dropwise addition of HCl to the clear filtrate. Wash the ppte with cold H₂O and dry the monohydrate at 30° in a vacuum. The pure material gives a single spot on paper chromatography (eluting solvent EtOH/water/phenol, 6:3:1; and developing with NaOH). [Anderegg et al. *Helv Chim Acta* 37 113 1954.] Complexes with Ba, Ca, Cd, Mg and Sr.

***o*-Cresol Red** [1733-12-6] M 382.4, m 290°(dec), pK²⁵ 1.26. Crystd from glacial acetic acid. Air dried. Dissolved in aqueous 5% NaHCO₃ soln and pted from hot soln by dropwise addition of aqueous HCl. Repeated until extinction coefficients did not increase.

***o*-Cresotic acid (methyalsalicylic acid)** [83-40-9] M 152.2, m 163-164°, pK₁²⁵ 3.32. Crystd from water.

***m*-Cresotic acid** [50-85-1] M 152.2, m 177°, pK₁²⁵ 3.15, pK₂²⁵ 13.35. Crystd from water.

***p*-Cresotic acid** [89-56-5] M 152.2, m 151°, pK₁²⁵ 3.40, pK₂²⁵ 13.45. Crystd from water.

Crocetin diethyl ester [5056-14-4] M 384.5, m 218-219°, 222.5°, $A_{1\text{cm}}^{1\%}$ (λ_{max}) 2340 (400nm), 3820 (422nm), 3850 (450nm) in pet ether. Purified by chromatography on a column of silica gel G. Crystd from *benzene. Stored in the dark, under an inert atmosphere, at 0°.

Crotonaldehyde (2-butenal) [123-73-9] M 70.1, b 104-105°, d 0.851, n 1.437. Fractionally distd under N₂, through a short Vigreux column. Stored in sealed ampoules. Stabilised with 0.01% of 2,6-di-tert-butyl-*p*-cresol

trans-Crotonic acid [107-93-7] M 86.1, m 72-72.5°, pK_1^{25} -6.17 (aq H₂SO₄), pK_2^{18} 4.71. Distd under reduced pressure. Crystd from pet ether (b 60-80°) or water, or by partial freezing of the melt.

E- and Z-Crotonitrile (mixture) [4786-20-3] M 67.1, b 120-121°, d 1.091, n 1.4595. Separated by preparative GLC on a column using 5% FFAP on Chromosorb G. [Lewis et al. *J Am Chem Soc* 108 2818 1986.]

γ -Crotonolactone [2(5H)-furanone] [497-23-4] M 84.1, m 3-4°, 76-77°/3.5mm, 90.5-91°/11.5mm, 92-93°/14mm, 107-109°/24mm, 212-214°/760mm, d_4^{20} 1.197, n_D^{20} 1.470. Fractionally distd under reduced pressure. IR: (CCl₄) 1784 and 1742 cm⁻¹, UV no max above 205nm (ϵ 1160 cm⁻¹ M⁻¹) and ¹H NMR: (CCl₃) τ : 2.15 (pair of triplets 1H), 3.85 (pair of triplets 1H) and 5.03 (triplet 2H). [Org Synth Coll Vol V 255 1973; Smith and Jones *Can J Chem* 37 2007, 2092 1959].

Crotyl bromide [29576-14-5] M 135.0, b 103-105°/740mm, n_D^{25} 1.4792. Dried with MgSO₄, CaCO₃ mixture. Fractionally distd through an all-glass Todd column. [Todd column. A column (which may be a Dufton type, fitted with a Monel metal rod and spiral, or a Hempel type, fitted with glass helices) which is surrounded by an open heating jacket so that the temperature can be adjusted to be close to the distillation temperature (Todd *Ind Eng Chem (Anal Ed)* 17 175 1945)].

15-Crown-5 [33100-27-5] M 220.3, b 93-96°/0.1mm, d 1.113, n 1.465. Dried over 3A molecular sieves.

18-Crown-6 [17455-13-9] M 264.3, m 37-39°. Recrystd from acetonitrile and vacuum dried. Purified by pptn of 18-crown-6/nitromethane 1:2 complex with Et₂O/nitromethane (10:1 mixture). The complex is decomposed in vacuum and distilled under reduced pressure. Also recrystd from acetonitrile and vacuum dried.

Cryptopine [482-74-6] M 369.4, m 220-221°. Crystd from *benzene.

Cryptoxanthin [472-70-8] M 552.9, $A_{1\text{cm}}^{1\%}$ 2370 (452nm), 2080 (480nm) in pet ether. Purified by chromatography on MgO, CaCO₃ or deactivated alumina, using EtOH or diethyl ether to develop the column. Crystd from CHCl₃/EtOH. Stored in the dark, under inert atmosphere, at -20°.

Crystal Violet Chloride {Gentian violet, N-4[bis[4-(dimethylaminophenyl)methylene]-2,5-cyclohexadien-1-ylidene]-N-methylmethaninium chloride} [548-62-9] M 408.0, pK 9.36. Crystd from water (20mL/g), the crystals being separated from the chilled soln by centrifugation, then washed with chilled EtOH (sol 1g in 10 mL of hot EtOH) and diethyl ether and dried under vac. It is sol in CHCl₃ but insol in Et₂O. The carbinol was ppted from an aqueous soln of the HCl dye, using excess NaOH, then dissolved in HCl and recrystd from water as the chloride [UV and kinetics: Turgeon and La Mer *J Am Chem Soc* 74 5988 1952]. The *carbinol base* has m 195° (needles from EtOH). The *diphthalate* (blue and turns red in H₂O) crystallises from H₂O, m 153-154° (dec 185-187°)[Chamberlain and Dull *J Am Chem Soc* 50 3089 1928].

Cumene (isopropyl benzene) [98-82-8] M 120.2, b 69-70°/41mm, 152.4°/760mm, d 0.864, n 1.49146, n_D^{25} 1.48892. Usual purification is by washing with several small portions of conc H₂SO₄ (until the acid layer is no longer coloured), then with water, 10% aq Na₂CO₃, again with water, and drying with MgSO₄, MgCO₃ or Na₂SO₄, followed by fractional distn. It can then be dried with, and distd from, Na, NaH or CaH₂. Passage through columns of alumina or silica gel removes oxidation products. Has also been steam

distd from 3% NaOH, and azeotropically distd with 2-ethoxyethanol (which was subsequently removed by washing out with water).

Cumene hydroperoxide [80-15-9] M 152.2, b 60°/0.2mm, d 1.028, n²⁴ 1.5232. Purified by adding 100mL of 70% material slowly and with agitation to 300mL of 25% NaOH in water, keeping the temperature below 30°. The resulting crystals of the sodium salt were filtered off, washed twice with 25 mL portions of *benzene, then stirred with 100mL of *benzene for 20min. After filtering off the crystals and repeating the washing, they were suspended in 100mL of distilled water and the pH was adjusted to 7.5 by addn of 4M HCl. The free hydroperoxide was extracted into two 20mL portions of *n*-hexane, and the solvent was evaporated under vacuum at room temperature, the last traces being removed at 40-50° and 1mm [Fordham and Williams *Canad J Res* 27B 943 1949]. Petroleum ether, **but not diethyl ether**, can be used instead of *benzene, and powdered solid CO₂ can replace the 4M HCl. *The material is potentially EXPLOSIVE.*

Cuminaldehyde (4-isopropylbenzaldehyde) [122-03-2] M 148.2, b 82-84°/3.5 mm, 120°/23mm, 131-135°/35mm, 235-236°/760mm, d²⁰ 0.978, n_D²⁰ 1.5301. Likely impurity is the benzoic acid. Check the IR for the presence of OH from CO₂H and the CO frequencies. If acid is present then dissolve in Et₂O, wash with 10% NaHCO₃ until effervescence ceases, then with brine, dry over CaCl₂, evap and distil the residual oil, preferably under vacuum. It is almost insoluble in H₂O, but soluble in EtOH and Et₂O. The *thiosemicarbazone* has m 147° after recrystn from aqueous EtOH, or MeOH or *C₆H₆. [Crouse *J Am Chem Soc* 71 1263 1949; Bernstein et al. *J Am Chem Soc* 73 906 1951; Gensler and Berman *J Am Chem Soc* 80 4949 1958.]

Cuprein (6'-hydroxyquinchonidine) [524-63-0] M 310.4, m 202°, [α]_D⁷ -176° (in MeOH), pK¹⁵ 7.63. Crystd from EtOH.

Curcumin [bis-(4-hydroxy-3-methoxycinnamoyl)methane] [458-37-7] M 368.4, m 183°. Crystd from EtOH or acetic acid.

Cyanamide [420-04-2] M 42.0, m 41°, pK₁²⁵ -0.36, pK₂²⁵ 10.27. See cyanamide on p. 416 in Chapter 5.

Cyanoacetamide [107-91-5] M 84.1, m 119.4°. Crystd from MeOH/dioxane (6:4), then water. Dried over P₂O₅ under vacuum.

Cyanoacetic acid [372-09-8] M 85.1, m 70.9-71.1°, pK²⁵ 2.47. Crystd to constant melting point from *benzene/acetone (2:3), and dried over silica gel.

Cyanoacetic acid hydrazide [140-87-4] M 99.1, m 114.5-115°. Crystd from EtOH.

p-**Cyanoaniline** [873-74-5] M 118.1, m 85-87°. See *p*-aminobenzonitrile on p. 104.

9-Cyanoanthracene (anthracene-9-carbonitrile) [1210-12-4] M 203.2, m 134-137°. Purified by crystn from EtOH or toluene, and vacuum sublimed in the dark and in an inert atmosphere [Ebied et al. *J Chem Soc, Faraday Trans 1* 76 2170 1980; Kikuchi et al. *J Phys Chem* 91 574 1987].

9-Cyanoanthracene photodimer [33998-38-8] M 406.4, dec to monomer above ~147°. Purified by dissolving in the minimum amount of CHCl₃ followed by addition of EtOH at 5° [Ebied et al. *J Chem Soc, Faraday Trans 1* 75 1111 1979; 76 2170 1980].

p-**Cyanobenzoic acid** [619-65-8] M 147.1, m 219°, pK²⁵ 3.55. Crystd from water and dried in vacuum desiccator over Sicapent.

4-Cyanobenzoyl chloride [6068-72-0] M 165.6, m 68-70°, 69-70°, 73-74°, b 132°/8 mm, 150-151°/25mm. If the IR shows presence of OH then treat with SOCl₂ boil for 1h, evaporate and distil in

vacuum. The distillate solidifies and can be recrystallised from pet ether. It is moisture sensitive and is an **IRRITANT**. [Ashley et al. *J Chem Soc* 103 1942; Fison et al. *J Org Chem* 16 648 1951.]

Cyanoguanidine (dicyanodiamide) [461-58-5] **M 84.1, m 209.5°, pK -0.4**. Crystd from water or EtOH.

5-Cyanoindole [15861-24-2] **M 142.2, m 106-108°, 107-108°, pK <0**. Dissolve in 95% EtOH boil in the presence of charcoal, filter, evaporate to a small volume and add enough H₂O to cause crystallisation and cool. Recrystd directly from aqueous EtOH and dried in a vacuum. UV: λ_{\max} 276 nm (log ϵ 3.6) in MeOH. [Lindwall and Mantell *J Org Chem* 18 345 1953, 20 1458 1955; Thesing et al. *Chem Ber* 95 2205 1962; NMR: Lallemand and Bernath *Bull Soc Chim Fr* 4091 1970.]

p-Cyanophenol [767-00-0] **M 119.1, m 113°, pK²⁵ 7.97**. Crystd from pet ether, *benzene or water and kept under vacuum over P₂O₅. [Bernasconi and Paschelis *J Am Chem Soc* 108 2969 1986.]

3-Cyanopyridine [100-54-9] **M 104.1, m 50°, pK²⁵ 1.38**. Crystd to constant melting point from *o*-xylene/hexane.

4-Cyanopyridine [100-48-1] **M 104.1, m 76-79°, pK²⁵ 1.86**. Crystd from dichloromethane/diethyl ether mixture.

Cyanuric acid (2,4,6-trichloro-1,3,5-triazine) [108-80-5] **M 120.1, m >300°, pK 6.78**. Crystd from water. Dried at room temperature in a desiccator under vacuum.

Cyanuric chloride (TCT, 2,4,6-trichloro-1,3,5-triazine) [108-77-0] **M 184.4, m 146-149°, 154°, b 190°**. Crystd from CCl₄ or pet ether (b 90-100°), and dried under vacuum. Recrystd twice from anhydrous *benzene immediately before use [Abuchowski et al. *J Biol Chem* 252 3582 1977].

Cyclobutane carboxylic acid [3721-95-7] **M 100.1, m 3-4°, -5.4°, b 84-84.5°/10mm, 110°/25mm, 135-138°/110mm, 194°/760mm, d₄²⁰ 1.061, n_D²⁰ 1.453, pK²⁵ 4.79**. Dissolve in aqueous HCO₃⁻ and acidify with HCl and extract into Et₂O, wash with H₂O, dry (Na₂SO₄), concentrate to a small volume, then distil through a glass helices packed column. The *S*-benzylthiouronium salt has *m* 176° (from EtOH), and the *anilide* has *m* 112.5-113°, and the *p*-toluide has *m* 123°. [Payne and Smith *J Org Chem* 22 1680 1957; Kantaro and Gunning *J Am Chem Soc* 73 480 1951.]

trans-Cyclobutane-1,2-dicarboxylic acid [1124-13-6] **M 144.1, m 131°, pK₁²⁵ 4.11, pK₂²⁵ 5.15**. Crystd from *benzene.

Cyclobutanone [1191-95-3] **M 70.1, b 96-97°, d 0.931, n_D²⁰ 1.4189**. Treated with dilute aqueous KMnO₄, dried with molecular sieves and fractionally distd. Purified *via* the semicarbazone, then regenerated, dried with CaSO₄, and distd in a spinning-band column. Alternatively, purified by preparative gas chromatography using a Carbowax 20-M column at 80°. (This treatment removes acetone).

Cyclobutylamine [2516-34-9] **M 71.1, b 82-83°/atm, 83.2-84.2°/760mm, d₄²⁰ 0.839, n_D²⁰ 1.437, pK²⁵ 10.04 (9.34 in 50% aq EtOH)**. It has been purified by steam distn. The aqueous distillate (e.g. 2L) is acidified with 3N HCl (90mL) and evapd to dryness in a vacuum. The *hydrochloride* is treated with a few mL of H₂O, cooled in ice and a slush of KOH pellets ground in a little H₂O is added slowly in portions and keeping the soln very cold. The amine separates as an oil from the strongly alkaline soln. The oil is collected dried over solid KOH and distd using a vac jacketed Vigreux column and protected from CO₂ using a soda lime tube. The fraction boiling at 79-83° is collected, dried over solid KOH for 2 days and redistd over a few pellets of KOH (b 80.5-81.5°). Best distil in a dry N₂ atmosphere. The purity can be checked by GLC using a polyethylene glycol on Teflon column at 72°, 15 psi, flow rate of 102 mL/min of He. The sample can appear homogeneous but because of tailing it is not possible to tell if H₂O is present. The NMR in CCl₄ should show no signals less than 1 ppm from TMS. The *hydrochloride* has a multiplet at *ca* 1.5-2.6ppm (H 2,2,4,3,3,4,4), a quintet at 3.8 ppm (H 1) and a singlet at 4.75 for NH₂ [Roberts and Chambers *J Am Chem*

Soc 73 2509 1951]. The *benzenesulfonamide* has *m* 85-86° (from aq MeOH) and the *benzoyl derivative* has *m* 120.6-121.6° [Roberts and Mazur *J Am Chem Soc* 73 2509 1951; Iffland et al. *J Am Chem Soc* 75 4044 1953; *Org Synth Coll Vol V* 273 1973.]

Cyclodecanone [1502-06-3] *M* 154.2, *m* 21-24°, *b* 100-102°/12mm. Purified by sublimation in a vac.

cis-Cyclodecene [935-31-9] *M* 138.3, *m* -3°, -1°, *b* 73°/15mm, 90.3°/33mm, 194-195°/740mm, 197-199°/atm, d_4^{20} 0.8770, n_D^{20} 1.4854. Purified by fractional distn. It forms an AgNO₃ complex which crystallises from MeOH, *m* 167-187° [Cope et al. *J Am Chem Soc* 77 1628 1955; IR: Blomqvist et al. *J Am Chem Soc* 74 3636 1952; Prelog et al. *Helv Chim Acta* 35 1598 1952].

α-Cyclodextrin (H₂O) [10016-20-3] *M* 972.9, *m* >280°(dec), $[\alpha]_{546}^{20}$ +175° (c 10, H₂O). See entry on p. 524 in Chapter 6.

β-Cyclodextrin (H₂O) [7585-39-9, 68168-23-0] *M* 1135.0, *m* >300°(dec), $[\alpha]_{546}^{20}$ +170° (c 10, H₂O). See entry on p. 524 in Chapter 6.

trans-cis-cis-1,5,9-Cyclododecatriene (cyclododec-1c,5c,9t-triene) [2765-29-9] *M* 162.3, *m* -9°, -8°, *b* 117.5°/2mm, 237-239°/atm, 244°/760mm, d_4^{20} 0.907, n_D^{20} 1.5129. Purified by fractional distn, preferably in a vacuum under N₂, and forms an insoluble AgNO₃ complex. [Breil et al. *Makromol Chemie* 69 28 1963.]

Cyclododecylamine [1502-03-0] *M* 183.3, *m* 27-29°, *b* 140-150°/ca 18mm, 280°/atm, *pK* 9.62 (in 80% methyl cellosolve). It can be purified via the *hydrochloride salt* *m* 274-275° (from EtOH) or the *picrate* *m* 232-234°, and the free base is distd at water-pump vacuum [Prelog et al. *Helv Chim Acta* 33 365 1950].

1,3-Cycloheptadiene [4054-38-0] *M* 94.2, *b* 55°/75mm, 71.5°/150mm, 120-121°/atm, d_4^{20} 0.868, n_D^{20} 1.4972. It was purified by dissolving in Et₂O, wash with 5% HCl, H₂O, dry (MgSO₄), evap and the residue is distd under dry N₂ through a semi-micro column (some foaming occurs) [Cope et al. *J Am Chem Soc* 79 6287 1957; UV: Pesch and Friess *J Am Chem Soc* 72 5756 1950].

Cycloheptane [291-64-5] *M* 98.2, *b* 114.4°, *d* 0.812, *n* 1.4588. Distd from sodium, under nitrogen.

Cycloheptanol [502-41-0] *M* 114.2, *b* 77-81°/11mm, 185°/atm, *d* 0.951, *n* 1.477. Purified as described for cyclohexanol.

Cycloheptanone [502-42-1] *M* 112.2, *b* 105°/80mm, 172.5°/760, *d* 0.952, n_D^{24} 1.4607. Shaken with aq KMnO₄ to remove material absorbing around 230-240nm, then dried with Linde type 13X molecular sieves and fractionally distd.

Cycloheptatriene [544-25-2] *M* 92.1, *b* 114-115°, *d* 0.895, *n* 1.522. Washed with alkali, then fractionally distd.

Cycloheptylamine [5452-35-7] *M* 113.2, *b* 50-52°/11mm, 60°/18mm, d_4^{20} 0.887, n_D^{20} 1.472, *pK*_{Est} ~10.5 (H₂O), *pK*²⁴ 9.99 (in 50% aq methyl cellosolve). It can be purified by conversion to the *hydrochloride* *m* 242-246°, and the free base is distd under dry N₂ in a vacuum [Cope et al. *J Am Chem Soc* 75 3212 1953; Prelog et al. *Helv Chim Acta* 33 365 1950].

1,3-Cyclohexadiene [592-57-4] *M* 80.1, *b* 83-84°/atm, d_4^{20} 0.840, n_D^{20} 1.471. Distd from NaBH₄.

1,4-Cyclohexadiene [628-41-1] M 80.1, b 83-86°/714mm, 88.3°/741mm, 86-88°/atm, 88.7-89°/760mm, d_4^{20} 0.8573, n_4^{20} 1.4725. Dry over CaCl₂ and distil in a vacuum under N₂. [Hückel and Wörrfel *Chem Ber* 88 338 1955; Giovannini and Wegmüller *Helv Chim Acta* 42 1142 1959.]

Cyclohexane [110-82-7] M 84.2, f 6.6°, b 80.7°, d^{24} 0.77410, n 1.42623, n^{25} 1.42354. Commonly, washed with conc H₂SO₄ until the washings are colourless, followed by water, aq Na₂CO₃ or 5% NaOH, and again water until neutral. It is next dried with P₂O₅, Linde type 4A molecular sieves, CaCl₂, or MgSO₄ then Na and distd. Cyclohexane has been refluxed with, and distd from Na, CaH₂, LiAlH₄ (which also removes peroxides), sodium/potassium alloy, or P₂O₅. Traces of *benzene can be removed by passage through a column of silica gel that has been freshly heated: this gives material suitable for ultraviolet and infrared spectroscopy. If there is much *benzene in the cyclohexane, most of it can be removed by a preliminary treatment with nitrating acid (a cold mixture of 30mL conc HNO₃ and 70mL of conc H₂SO₄) which converts *benzene into nitrobenzene. The impure cyclohexane and the nitrating acid are placed in an ice bath and stirred vigorously for 15min, after which the mixture is allowed to warm to 25° during 1h. The cyclohexane is decanted, washed several times with 25% NaOH, then water dried with CaCl₂, and distd from sodium. Carbonyl-containing impurities can be removed as described for chloroform. Other purification procedures include passage through columns of activated alumina and repeated crystn by partial freezing. Small quantities may be purified by chromatography on a Dowex 710-Chromosorb W gas-liquid chromatographic column.

Rapid purification: Distil, discarding the forerun. Stand distillate over Grade I alumina (5% w/v) or 4A molecular sieves.

Cyclohexane butyric acid [4441-63-8] M 170.3, m 31°, 26.5-28.5°, b 136-139°/4mm. 169°/20mm, 188.8°/46mm, pK^{25} 4.95. Distil through a Vigreux column, and the crystalline distillate is recrystd from pet ether. The *S*-benzylthiuronium salt has m 154-155° (from EtOH) [*Acta Chem Scand* 9 1425 1955; English and Dayan *J Am Chem Soc* 72 4187 1950].

Cyclohexane-1,2-diaminetetraacetic acid (H₂O; CDTA) [13291-61-7] M 364.4, pK_1 1.34, pK_2 3.20, pK_3 5.75 (6.12), pK_4 9.26 (12.35). Dissolved in aq NaOH as its disodium salt, then pptd by adding HCl. The free acid was filtered off and boiled with distd water to remove traces of HCl [Bond and Jones *Trans Faraday Soc* 55 1310 1959]. Recrystd from water and dried under vacuum.

trans-Cyclohexane-1,2-dicarboxylic acid [2305-32-0] M 172.2, m 227.5-228°, 228-230.5°, pK_1^{25} 4.30, pK_2^{25} 6.06 [*cis*, pK_1^{25} 4.25, pK_2^{25} 6.74]. It is purified by recrystn from EtOH or H₂O. The dimethyl ester has m 95-96° (from *C₆H₆-pet ether). [Abell *J Org Chem* 22 769 1957; Smith and Byrne *J Am Chem Soc* 72 4406 1950; Linstead et al. *J Am Chem Soc* 64 2093 1942.]

(±)-trans-1,2-Cyclohexanediol [1460-57-7] M 116.2, m 104°, 105°, 120°/14mm. Crystd from Me₂CO and dried at 50° for several days. It can also be recrystd from CCl₄ or EtOAc and can be distilled. The 2,4-dinitrobenzoyl derivative has m 179°. [Winstein and Buckles *J Am Chem Soc* 64 2780 1942.]

trans-1,2-Cyclohexanediol [1R,2R(-)- 1072-86-2; 1S,2S(+)- 57794-08-8] M 116.2, m 107-109°, 109-110.5°, 109-111°, 111-112°, 113-114°, $[\alpha]_D^{22}$ (-) and (+) 46.5° (c 1, H₂O). The enantiomers have been recrystd from *C₆H₆ or EtOAc. The (±) diol has been resolved as the distrychnine salt of the hemisulfate [Hayward, Overton and Whitham *J Chem Soc Perkin Trans 1* 2413 1976]; or the 1-menthoxy acetates. {l-trans- diastereoisomer has m 64°, $[\alpha]_D$ -91.7° (c 1.4 EtOH) from pet ether or aqueous EtOH and yields the (-)-trans-diols } and {d-trans-diastereoisomer has m 126-127°, $[\alpha]_D$ -32.7° (c 0.8 EtOH) from pet ether or aq EtOH and yields the (+)-trans diol}. The bis-4-nitrobenzoate has m 126.5° $[\alpha]_D$ (-) and (+) 25.5° (c 1.1 CHCl₃), and the bis-3,5-dinitrobenzoate has m 160° $[\alpha]_D$ ± -83.0° (c 1.8 CHCl₃) [Wilson and Read *J Chem Soc* 1269 1935].

cis-1,3-Cyclohexanediol [823-18-7] M 116.2, m 86°. Crystd from ethyl acetate and acetone.

trans-1,3-Cyclohexanediol [5515-64-0] M 116.2, m 117°. Crystd from ethyl acetate.

cis-1,4-Cyclohexanediol [556-58-9] M 116.2, m 102.5°. Crystd from acetone (charcoal), then dried and sublimed under vacuum.

Cyclohexane-1,3-dione [504-02-9] M 112.1, m 107-108°. Crystd from *benzene.

Cyclohexane-1,4-dione [637-88-7] M 112.1, m 76-77°, 78°, 79.5°, 79-80°, b 130-133°/20mm, d_4^{25} 1.0861, n_D^{25} 1.4576. Crystd from water, then *benzene. It can also be recrystd from CHCl_3 /pet ether or Et_2O . It has been purified by distn in a vacuum and the pale yellow distillate which solidified is then recrystd from CCl_4 (14.3 g/100 mL) and has m 77-79°. The *di-semicarbazone* has m 231°, the *dioxime HCl* has m 150° (from $\text{MeOH} \cdot \text{C}_6\text{H}_6$) and the *bis-2,4-dinitrophenylhydrazone* m 240° (from PhNO_2). [*Org Synth Coll Vol V* 288 1973; IR: LeFevre and LeFevre *J Chem Soc* 3549 1956.]

Cyclohexane-1,2-dione dioxime (Nioxime) [492-99-9] M 142.2, m 189-190°, pK_1^{25} 10.68, pK_2^{25} 11.92. Crystd from alcohol/water and dried in a vacuum at 40°.

1,4-Cyclohexanedione monoethylene acetal (1,4-dioxa-spiro[4.5]decan-8-one) [4746-97-8] M 156.2, m 70-73°, 73.5-74.5°. Recrystd from pet ether and sublimes slowly on attempted distillation. Also purified by dissolving in Et_2O and adding pet ether (b 60-80°) until turbid and cool. [Gardner et al. *J Am Chem Soc* 22 1206 1957; Britten and Lockwood *J Chem Soc Perkin Trans 1* 1824 1974.]

cis,cis-1,3,5-Cyclohexane tricarboxylic acid [16526-68-4] M 216.2, m 216-218°, $\text{pK}_{\text{Est}(1)} \sim 4.1$, $\text{pK}_{\text{Est}(2)} \sim 5.4$, $\text{pK}_{\text{Est}(3)} \sim 6.8$. Purified by recrystn from toluene + EtOH or H_2O . It forms a 1.5 hydrate with m 216-218°, and a dihydrate at 110°. Purified also by conversion to the triethyl ester b 217-218°/10mm, 151°/1mm and distillate solidifies on cooling, m 36-37° and is hydrolysed by boiling in aq HCl . The trimethyl ester can be distd and recrystd from Et_2O , m 48-49°. [Newman and Lawrie *J Am Chem Soc* 76 4598 1954, Lukes and Galik *Coll Czech Chem Comm* 19 712 1954.]

Cyclohexanol [108-93-0] M 100.2, m 25.2°, b 161.1°, d 0.946, n 1.466, n^{25} 1.437, n^{30} 1.462. Refluxed with freshly ignited CaO , or dried with Na_2CO_3 , then fractionally distd. Redistd from Na. Further purified by fractional crystn from the melt in dry air. Peroxides and aldehydes can be removed by prior washing with ferrous sulfate and water, followed by distillation under nitrogen from 2,4-dinitrophenylhydrazine, using a short fractionating column: water distils as the azeotrope. Dry cyclohexanol is *very hygroscopic*.

Cyclohexanone [108-94-1] M 98.2, f -16.4°, b 155.7°, d 0.947, n^{15} 1.452, n 1.451, pK^{25} -6.8 (aq H_2SO_4), pK^{25} 11.3 (enol), 16.6 (keto). Dried with MgSO_4 , CaSO_4 , Na_2SO_4 or Linde type 13X molecular sieves, then distd. Cyclohexanol and other oxidisable impurities can be removed by treatment with chromic acid or dil KMnO_4 . More thorough purification is possible by conversion to the bisulfite addition compound, or the semicarbazone, followed by decomn with Na_2CO_3 and steam distn. [For example, equal weights of the bisulfite adduct (crystd from water) and Na_2CO_3 are dissolved in hot water and, after steam distn, the distillate is saturated with NaCl and extracted with *benzene which is then dried and the solvent evaporated prior to further distn.]

Cyclohexanone oxime [100-64-1] M 113.2, m 90°. Crystd from water or pet ether (b 60-80°).

Cyclohexanone phenylhydrazone [946-82-7] M 173.3, m 77°. Crystd from EtOH .

Cyclohexene [110-83-8] M 82.2, b 83°, d 0.810, n 1.4464, n^{25} 1.4437. Freed from peroxides by washing with successive portions of dil acidified ferrous sulfate, or with NaHSO_3 soln then with distd water, dried with CaCl_2 or CaSO_4 , and distd under N_2 . Alternative methods of removing peroxides include passage through a column of alumina, refluxing with sodium wire or cupric stearate (then distilling from sodium). Diene is removed by refluxing with maleic anhydride before distg under vac. Treatment with 0.1 moles of MeMgI in 40mL of diethyl ether removes traces of oxygenated impurities. Other purification procedures include washing with aq NaOH , drying and distg under N_2 through a spinning band column; redistg from CaH_2 ; storage with sodium wire; and passage through a column of alumina, under N_2 , immediately before use. Stored in a

refrigerator under argon. [Woon et al. *J Am Chem Soc* **108** 7990 1986; Wong et al. *J Am Chem Soc* **109** 3428 1987.]

(±)-2-Cyclohexen-1-ol (3-hydroxycyclohex-1-ene) [822-67-3] M 242.2, b 63-65°/12mm, 65-66°/13mm, 67°/15mm, 74°/25mm, 85°/35mm, 166°/atm, d_4^{20} 0.9865, n_D^{20} 1.4720. Purified by distillation through a short Vigreux column. The 2,4-dinitrobenzoyl derivative has m 120.5°, and the phenylurethane has m 107°. [*Org Synth* **48** 18 1968, Cook *J Chem Soc* 1774 1938; Deiding and Hartman *J Am Chem Soc* **75** 3725 1953.]

Cyclohexene oxide [286-20-4] M 98.2, b 131-133°/atm, d_4^{20} 0.971, n_D^{20} 1.452. Fractionated through an efficient column. The main impurity is probably H₂O. Dry over MgSO₄, filter and distil several times (b 129-134°/atm). The residue is sometimes hard to remove from the distilling flask. To avoid this difficulty, add a small amount of a mixture of ground NaCl and Celite (1:1) to help break the residue particularly if H₂O is added. [*Org Synth Coll Vol I* 185 1948.]

Cycloheximide [68-81-9] M 281.4, m 119.5-121°, $[\alpha]_{546}^{20}$ +9.5° (c 2, H₂O). Crystd from water/MeOH (4:1), amyl acetate, isopropyl acetate/isopropyl ether or water.

Cyclohexylamine [108-91-8] M 99.2, b 134.5°, d 0.866, d^{25} 0.863, n 1.4593, n^{25} 1.456, pK²⁵ 10.63. Dried with CaCl₂ or LiAlH₄, then distd from BaO, KOH or Na, under N₂. Also purified by conversion to the hydrochloride, several crystns from water, then liberation of the amine with alkali and fractional distn under N₂.

Cyclohexylbenzene [827-52-1] M 160.3, f 6.8°, b 237-239°, d 0.950, n 1.5258. Purified by fractional distn, and fractional freezing.

Cyclohexyl bromide [108-85-0] M 156.3, b 72°/29mm, d 0.902, n^{25} 1.4935. Shaken with 60% aqueous HBr to remove the free alcohol. After separation from excess HBr, the sample was dried and fractionally distd.

Cyclohexyl chloride [542-18-7] M 118.6, b 142-142.5°, d 1.000, n 1.462. See chlorocyclohexane on p. 162.

1-Cyclohexylethylamine [S-(+): 17430-98-7; R-(-): 5913-13-3] M 127.2, b 177-178°/atm, d_4^{20} 0.866, n_D^{20} 1.446, $[\alpha]_D^{15}$ (-) and (+) 3.2° (neat), pK_{Est} ~10.6. Purified by conversion to the bitartrate salt (m 172°), then decomposing with strong alkali and extracting into Et₂O, drying (KOH), filtering, evaporating and distilling. The hydrochloride salt has m 242° (from EtOH-Et₂O), $[\alpha]_D^{15}$ -5.0° (c 10 H₂O; from (+) amine). The oxalate salt has m 132° (from H₂O). The (±)-base has b 176-178°/760mm, and HCl has m 237-238°. [Reihlen, Knöpfle and Sapper *Justus Liebigs Ann Chem* **532** 247 1938; *Chem Ber* **65** 660 1932.]

Cyclohexylidene fulvene [3141-04-6] M 134.2. Purified by column chromatography and eluted with n-hexane [Abboud et al. *J Am Chem Soc* **109** 1334 1987].

Cyclohexyl mercaptan (cyclohexane thiol) [1569-69-3] M 116.2, b 38-39°/12mm, 57°/23mm, 90°/100mm, 157°/763mm, d_4^{20} 0.949, n_D^{20} 1.493, pK_{Est} ~10.8. Possible impurities are the sulfide and the disulfide. Purified by conversion to the Na salt by dissolving in 10% aq NaOH, extract the sulfide and disulfide with Et₂O, and then acidify the aq soln (with cooling and under N₂) with HCl, extract with Et₂O, dry MgSO₄, evaporate and distil in a vacuum (b 41°/12mm). The sulfide has b 74°/0.2mm, $n_D^{18.5}$ 1.5162 and the disulfide has b 110-112°/0.2mm, $n_D^{18.5}$ 1.5557. The Hg-mercaptide has m 77-78° (needles from EtOH). [Naylor *J Chem Soc* 1532 1947.]

Cyclohexyl methacrylate [101-43-9] M 168.2, b 81-86°/0.1mm, d 0.964, n 1.458. Purification as for methyl methacrylate.

1-Cyclohexyl-5-methyltetrazole [7707-57-5] M 166.2, m 124-124.5°. Crystd from absolute EtOH, then sublimed at 115°/3mm.

Cyclononane [3350-30-9] M 140.2, m 142.0-142.8°, b 220-222°. Repeatedly sublimed at 0.05-0.1mm pressure.

cis,cis-1,3-Cyclooctadiene [29965-97-7] M 108.2, m -5°, -49°, b 55°/34mm, 142-144°/760mm, d_4^{20} 0.8690, n_D^{20} 1.48921. Purified by GLC. Fractionally distd through a Widmer column [as a mobile liquid and redistilled with a Claisen flask or through a semi-micro column [Gould, Holzman and Neiman *Anal Chem* 20 361 1948]. NB: It has a strong characteristic disagreeable odour detectable at low concentrations and causes headaches on prolonged exposure. [IR: Cope and Estes *J Am Chem Soc* 72 1128 1950; UV: Cope and Baumgardner *J Am Chem Soc* 78 2812 1956.] [Widmer column. A Dufton column, modified by enclosing within two concentric tubes the portion containing the glass spiral. Vapour passes up the outer tube and down the inner tube before entering the centre portion. In this way flooding of the column, especially at high temperatures, is greatly reduced (Widmer *Helv Chim Acta* 7 59 1924).]

cis-cis-1,5-cyclooctadiene, [1552-12-1] M 108.2, m -69.5°, -70°, b 51-52°/25mm, 97°/144mm, 150.8°/757mm, d_4^{20} 0.880, n_D^{20} 1.4935, . Purified by GLC. It has been purified *via* the AgNO₃ salt. This is prepared by shaking with a soln of 50% aq AgNO₃ w/w several times (e.g. 3 x 50 mL and 4 x 50 mL) at 70° for *ca* 20min to get a good separation of layers. The upper layers are combined and further extracted with AgNO₃ at 40° (2 x 20 mL). The upper layer (19 mL) of original hydrocarbon mixture gives colourless needles AgNO₃ complex on cooling. The adduct is recrystd from MeOH (and cooling to 0°). The hydrocarbon is recovered by steam distilling the salt. The distillate is extracted with Et₂O, dried (MgSO₄), evap and distd. [Jones *J Chem Soc* 312 1954.]

Cyclooctanone [502-49-8] M 126.2, m 42°. Purified by sublimation after drying with Linde type 13X molecular sieves.

1,3,5,7-Cyclooctatetraene [629-20-9] M 104.2, b 141-141.5°, d 1.537, n_D^{25} 1.5350. Purified by shaking 3mL with 20mL of 10% aqueous AgNO₃ for 15min, then filtering off the silver nitrate complex as a ppte. The ppte was dissolved in water and added to cold conc ammonia to regenerate the cyclooctatetraene which was fractionally distd under vacuum onto molecular sieves and stored at 0°. It was passed through a dry alumina column before use [Broadley et al. *J Chem Soc, Dalton Trans* 373 1986].

cis-Cyclooctene [931-88-4] M 110.2, b 32-34°/12mm, 66.5-67°/60mm, 88°/141mm, 140°/170mm, 143°/760mm, d_4^{20} 0.84843, n_D^{20} 1.4702, . The *cis*-isomer was freed from the *trans*-isomer by fractional distn through a spinning-band column, followed by preparative gas chromatography on a Dowex 710-Chromosorb W GLC column. It was passed through a short alumina column immediately before use [Collman et al. *J Am Chem Soc* 108 2588 1986]. It has also been distd in a dry nitrogen glove box from powdered fused NaOH through a Vigreux column and then passed through activated neutral alumina before use [Wong et al. *J Am Chem Soc* 109 4328 1987]. Alternatively it can be purified *via* the AgNO₃ salt. This salt is obtained from crude cyclooctene (40 mL) which is shaken at 70-80° with 50% w/w AgNO₃ (2 x 15 mL) to remove cyclooctadienes (aq layer). Extraction is repeated at 40° (4 x 20 mL, of 50% AgNO₃). Three layers are formed each time. The middle layer contains the AgNO₃ adduct of cyclooctene which crystallises on cooling the layer to room temperature. The adduct (complex 2:1) is highly soluble in MeOH (at least 1g/mL) from which it crystallises in large flat needles when cooled at 0°. It is dried under slight vacuum for 1 week in the presence of CaCl₂ and paraffin wax soaked in the cyclooctene. It has m 51° and loses hydrocarbon on exposure to air. *cis*-Cyclooctene can be recovered by steam distn of the salt, collected, dried (CaCl₂) and distilled in vacuum. [Braude et al. *J Chem Soc* 4711 1957; AgNO₃: Jones *J Chem Soc* 1808 1954; Cope and Estes *J Am Chem Soc* 72 1128 1950.]

cis-Cyclooctene oxide {(1r,8c)-9-oxabicyclo[6.1.0]nonane} [286-62-4] M 126.7, m 56-57°, 57.5-57.8°, 50-60°, b 85-88°/17mm, 82.5°/22mm, 90-93°/37mm, 189-190°/atm. It can be distd in vacuum and the solidified distillate can be sublimed in vacuum below 50°. It has a characteristic odour.

[IR: Cope et al. *J Am Chem Soc* **74** 5884 1952, **79** 3905 1957; Reppe et al. *Justus Liebigs Ann Chem* **560** 1 1948].

Cyclopentadecanone [502-72-7] **M 224.4, m 63°**. Sublimation is better than crystn from aq EtOH.

Cyclopentadiene [542-92-7] **M 66.1, b 41-42°**. Dried with $Mg(ClO_4)_2$ and distd.

Cyclopentane [287-92-3] **M 70.1, b 49.3°, d 0.745, n 1.40645, n²⁵ 1.4340**. Freed from cyclopentene by two passages through a column of carefully dried and degassed activated silica gel.

Cyclopentane carbonitrile [4254-02-8] **M 95.2, m -75.2°, -76°, b 43-44°/7mm, 50-62°/10mm, 67-68°/14mm, 74.5-75°/30mm, d₄²⁰ 0.912, n_D²⁰ 1.441**. Dissolve in Et₂O, wash thoroughly with saturated aqueous K₂CO₃, dry (MgSO₄) and distil through a 10 cm Vigreux column. [McElvain and Stern *J Am Chem Soc* **77** 457 1955, Bailey and Daly *J Am Chem Soc* **81** 5397 1959.]

Cyclopentane-1,1-dicarboxylic acid [5802-65-3] **M 158.1, m 184°, pK₁ 3.23, pK₂ 4.08**. Recrystd from water.

1,3-Cyclopentanedione [3859-41-4] **M 98.1, m 149-150°, 151-152.5°, 151-154°, 151-153°**. Purified by Soxhlet extraction with CHCl₃. The CHCl₃ is evaporated and the residue is recrystd from EtOAc and/or sublimed at 120°/4mm. It has an acidic pK_a of 4.5 in H₂O. [IR: Boothe et al. *J Am Chem Soc* **75** 1732 1953; DePuy and Zaweski *J Am Chem Soc* **81** 4920 1959.]

Cyclopentanone [120-92-3] **M 84.1, b 130-130.5°, d 0.947, n 1.4370, n²⁵ 1.4340**. Shaken with aq KMnO₄ to remove materials absorbing around 230 to 240nm. Dried with Linde type 13X molecular sieves and fractionally distd. Has also been purified by conversion to the NaHSO₃ adduct which, after crystallising four times from EtOH/water (4:1), was decomposed by adding to an equal weight of Na₂CO₃ in hot H₂O. The free cyclopentanone was steam distd from the soln. The distillate was saturated with NaCl and extracted with *benzene which was then dried and evaporated; the residue was distd [Allen, Ellington and Meakins *J Chem Soc* 1909 1960].

Cyclopentene [142-29-0] **M 68.1, b 45-46°, d 0.772, n 1.4228**. Freed from hydroperoxide by refluxing with cupric stearate. Fractionally distd from Na. Chromatographed on a Dowex 710-Chromosorb W GLC column. Methods for **cyclohexene** should be applicable here. Also washed with 1M NaOH soln followed by water. It was dried over anhydrous Na₂SO₄, distd over powdered NaOH under nitrogen, and passed through neutral alumina before use [Woon et al. *J Am Chem Soc* **108** 7990 1986]. It was distd in a dry nitrogen atmosphere from powdered fused NaOH through a Vigreux column, and then passed through activated neutral alumina before use [Wong et al. *J Am Chem Soc* **109** 3428 1987].

1-Cyclopentene-1,2-dicarboxylic anhydride [3205-94-5] **M 138.1, m 42-54°, 46-47°, b 130°/5mm, 133-135°/5mm, n_D²⁰ 1.497**. If IR has OH peaks then some hydrolysis to the diacid (m 178°) must have occurred. In this case reflux with an appropriate volume of Ac₂O for 30min, evaporate the Ac₂O and distil *in vacuo*. The distillate solidifies and can be recrystd from EtOAc-hexane (1:1). The diacid distils without dec due to formation of the anhydride. The *dimethyl ester* has m 120-125°/11mm. [Askain *Chem Ber* **98** 2322 1965.]

Cyclopentylamine [1003-03-8] **M 85.2, m -85.7°, b 106-108°/760mm, 108.5°/760mm, d₄²⁰ 0.869, n_D²⁰ 1.452, pK²⁵ 10.65, (pK²⁵ 4.05 in 50% aq EtOH)**. May contain H₂O or CO₂ in the form of carbamate salt. Dry over KOH pellets and then distil from a few pellets of KOH. Store in a dark, dry CO₂-free atmosphere. It is characterised as the *thiocyanate salt* m 94.5°. The *benzenesulfonyl derivative* has m 68.5-69.5°. [Roberts and Chambers *J Am Chem Soc* **73** 5030 1951; Bollinger et al. *J Am Chem Soc* **75** 1729 1953.]

Cyclopropane [75-19-4] **M 42.1, b -34°**. Washed with a soln of HgSO₄, and dried with CaCl₂, then $Mg(ClO_4)_2$.

Cyclopropanecarbonyl chloride [4023-34-1] M 104.5, b 117.9-118.0°/723mm, 119.5-119.6¹⁰/760mm, d₄²⁰ 1.142, n₄²⁰ 1.453. If the IR shows OH bands then some hydrolysis to the free acid must have occurred. In this case heat with oxalyl chloride at 50° for 2h or SOCl₂ for 30min, then evap and distil three times using a Dufton column. Store in an inert atm, preferably in sealed tubes. Strong IRRITANT. If it is free from OH bands then just distil *in vacuo* and store as before. [Jeffrey and Vogel *J Chem Soc* 1804 1948.]

Cyclopropane-1,1-dicarboxylic acid [598-10-7] M 130.1, m 140°, pK₁²⁵ 1.8, pK₂²⁵ 5.42. Recrystd from CHCl₃.

Cyclopropylamine [765-30-0] M 57.1, b 49-49.5°/760mm, 48-50°/atm, 49-50°/750mm, d₄²⁰ 0.816, n_D²⁰ 1.421, pK²⁵ 9.10 (pK²⁵ 5.33 in 40% aq EtOH). It has been isolated as the *benzamide* m 100.6-101.0° (from aqueous EtOH). It forms a *picrate* m 149° (from EtOH-pet ether) from which the free base can be recovered using a basic ion exchange resin and can then be distd through a Todd column (see p. 174) using an automatic still head which only collects products boiling below 51°/atm. Polymeric materials if present will boil above this temperature. The *hydrochloride* has m 85-86° [Roberts and Chambers *J Am Chem Soc* 73 5030 1951; Jones *J Org Chem* 9 484 1944; Emmons *J Am Chem Soc* 79 6522 1957].

Cyclopropyldiphenylcarbinol [5785-66-0] M 224.3, m 86-87°. Crystd from *n*-heptane.

Cyclopropyl methyl ketone [765-43-5] M 84.1, b 111.6-111.8°/752mm, d 0.850, n 1.4242. Stored with anhydrous CaSO₄, distd under nitrogen. Redistd under vacuum.

Cyclotetradecane [295-17-0] M 192.3, m 56°. Recrystd twice from aq EtOH then sublimed *in vacuo* [Dretloff et al. *J Am Chem Soc* 109 7797 1987].

Cyclotetradecanone [3603-99-4] M 206.3, m 25°, b 145°/10mm, d 0.926, n 1.480. It was converted to the semicarbazone which was recrystd from EtOH and reconverted to the free cyclotetradecanone by hydrolysis [Dretloff et al. *J Am Chem Soc* 109 7797 1987].

Cyclotrimethylenetrinitramine (RDX, 1,3,5-trinitrohexahydro-1,3,5-triazine) [121-82-4] M 222.2, m 203.8°(dec). Crystd from acetone. EXPLOSIVE.

***p*-Cymene** [99-87-6] M 134.2, b 177.1°, d 0.8569, n 1.4909, n²⁵ 1.4885. Washed with cold, conc H₂SO₄ until there is no further colour change, then repeatedly with H₂O, 10% aqueous Na₂CO₃ and H₂O again. Dried with Na₂SO₄, CaCl₂ or MgSO₄, and distd. Further purification steps include steam distn from 3% NaOH, percolation through silica gel or activated alumina, and a preliminary refluxing for several days over powdered sulfur. Stored over CaH₂.

Cystamine dihydrochloride, *S,S*-(*L,L*)-Cystathionine, Cysteamine, Cysteamine hydrochloride, (±)-Cysteic acid, *S*-Cysteic acid (H₂O), *L*-Cysteine hydrochloride (H₂O), (±)-Cysteine hydrochloride and *L*-Cystine, Cytidine, see entries in Chapter 6.

Cytisine (7*R*,9*S*-7,9,10,11,12,13-hexahydro-7,9-methano-12*H*-pyrido[1,2-*a*][1,5]diazocin-8-one, Laburnine, Ulexine) [485-35-8] M 190.3, m 152-153°, 155°, b 218°/2mm, [α]_D¹⁷ -120° (H₂O), [α]_D²³ -115° (c 1, H₂O), pK₁¹⁵ 1.20, pK₂¹⁵ 8.12 [also stated are pK₁ 6.11, pK₂ 13.08]. Crystd from acetone and sublimed in a vacuum. Its solubilities are: 77% (H₂O), 7.7% (Me₂CO), 28.6% (EtOH), 3.3% (*C₆H₆), 50% (CHCl₃) but is insoluble in pet ether. The *tartrate* has m 206-207° [α]_D²⁴ +45.9°, the *N*-*tosylate* has m 206-207°, and the *N*-*acetate* has m 208°. [Bohlmann et al. *Angew Chem* 67 708 1955; van Tamelen and Baran *J Am Chem Soc* 77 4944 1955; Isolation: Ing *J Chem Soc* 2200 1931; Govindachari et al. *J Chem Soc* 3839 1957; Abs config: Okuda et al. *Chem Ind (London)* 1751 1961.] TOXIC.

Cytosine see entry in Chapter 6.

***cis*-Decahydroisoquinoline** [2744-08-3] **M 139.2, b 97-98°/15mm, 208-209°/730mm, pK²⁰ 11.32.** The free base is treated with satd aq picric acid, allowed to stand for 12h, fild, washed with MeOH to remove the more soluble *trans* isomer and recrystd from MeOH to give pure *cis-picrate* **m 149-150°.** The picrate (~5g) is shaken with 5M aq NaOH (50mL) and Et₂O (150mL) while H₂O is added to the aq phase to dissolve insoluble Na picrate. The Et₂O extract is dried over solid NaOH and then shaken with Al₂O₃ (Merck for chromatography) until the yellow color of traces of picric acid disappears (this color cannot be removed by repeated shaking with 5-10 M aq NaOH). The extract is concentrated to 50mL and dry HCl is bubbled through until separatrn of the white crystals of the *cis-HCl* is complete. These are washed with Et₂O, dried at 100° and recrystd from EtOH + EtOAc to yield pure *cis-Hydrochloride* **m 182-183°** (dried in a vac desiccator over KOH) with IR (KBr) ν_{\max} 2920, 2820, 1582, 1470, 1445, 1410, 1395, 1313, 1135, 1080, 990, 870 cm⁻¹. The pure *free base* is prepared by dissolving the *hydrochloride* in 10 M aq NaOH, extracted with Et₂O, dried over solid KOH, fild and distd in vac. It has IR (film) ν_{\max} 2920, 2820, 2720, 2560, 1584, 1470, 1445, 1415, 1395, 1315, 1300, 1135, 1080, 1020, 990, 873 cm⁻¹. The ¹H NMR in CDCl₃ is characteristically different from that of the *trans*-isomer. [Armarego *J Chem Soc (C)* 377 1967; Gray and Heitmeier *J Am Chem Soc* 80 6274 1958; Witkop *J Am Chem Soc* 70 2617 1948; Skita *Chem Ber* 57 1982 1924; Helfer *Helv Chim Acta* 6 7991923.]

***trans*-Decahydroisoquinoline** [2744-09-4] **M 139.2, b 106°/15mm, pK²⁰ 11.32.** This is purified as the *cis*-isomer above. The *trans-picrate* has **m 175-176°**, and the *trans-hydrochloride* has **m 221-222°** and has IR (KBr) ν_{\max} 2930, 3800, 1589, 1450, 1400, 1070, 952, 837 cm⁻¹. The pure *free base* was prepared as above and had IR (film) ν_{\max} 2920, 2820, 2720, 2560, 1584, 1470, 1445, 1415, 1395, 1315, 1300, 1135, 1080, 1020, 990, 873 cm⁻¹. The ¹H NMR in CDCl₃ is characteristically different from that of the *cis*-isomer. (references as above and Helfer *Helv Chim Acta* 9 818 1926).

Decahydronaphthalene (decalin, mixed isomers) [91-17-8] **M 138.2, b 191.7°, d 0.886, n 1.476.** Stirred with conc H₂SO₄ for some hours. Then the organic phase was separated, washed with water, saturated aqueous Na₂CO₃, again with water, dried with CaSO₄ or CaH₂ (and perhaps dried further with Na), filtered and distd under reduced pressure (**b 63-70°/10mm**). Also purified by repeated passage through long columns of silica gel previously activated at 200-250°, followed by distn from LiAlH₄ and storage under N₂. Type 4A molecular sieves can be used as a drying agent. Storage over silica gel removes water and other polar substances.

***cis*-Decahydronaphthalene** [493-01-6] **M 138.2, f -43.2°, b 195.7°, d 0.897, n 1.48113.** Purification methods described for the mixed isomers are applicable. The individual isomers can be separated by very efficient fractional distn, followed by fractional crystn by partial freezing. The *cis*-isomer reacts preferentially with AlCl₃ and can be removed from the *trans*-isomer by stirring the mixture with a limited amount of AlCl₃ for 48h at room temperature, filtering and distilling.

***trans*-Decahydronaphthalene** [493-02-7] **M 138.2, f -30.6°, b 187.3°, d 0.870, n 1.46968.** See purification of *cis*-isomer above.

***cis*-Decahydroquinoline** [10343-99-4] **M 139.2, b 207-208°/708mm, pK²⁰ 11.29.** It is available as a *cis-trans*-mixture (**b 70-73°/10mm**, Aldrich, ~ 18% *cis*-isomer [2051-28-7]), but the isomers can be fractionated in a spinning band column (1~1.5 metre, type E) at atmospheric pressure and collecting 2mL fractions with a distillation rate of 1 drop in 8-10sec. The lower boiling fraction solidifies and contains the *trans*-isomer (see below, **m 48°**). The higher boiling fraction **b 207-208°/708mm**, remains liquid and is mostly the *cis*-isomer. This is reacted with PhCOCl and M aq NaOH to yield the *N-benzoyl derivative* **m 96°** after recrystn from pet ether (**b 80-100°**). It is hydrolysed with 20% aq HCl by refluxing overnight. PhCO₂H is fild off, the filtrate is basified with 5M aq NaOH and extracted with Et₂O. The dried extract (Na₂SO₄) is satd with dry HCl gas and the *cis-decahydroquinoline hydrochloride* which separates has **m 222-224°** after washing with Et₂O and drying at 100°; and has IR (KBr) ν_{\max} 2900, 2780, 2560, 1580, 1445, 1432, 1403, 1165, 1080,

Cytosine see entry in Chapter 6.

***cis*-Decahydroisoquinoline** [2744-08-3] **M 139.2, b 97-98°/15mm, 208-209°/730mm, pK²⁰ 11.32.** The free base is treated with satd aq picric acid, allowed to stand for 12h, fild, washed with MeOH to remove the more soluble *trans* isomer and recrystd from MeOH to give pure *cis-picrate* **m 149-150°.** The picrate (~5g) is shaken with 5M aq NaOH (50mL) and Et₂O (150mL) while H₂O is added to the aq phase to dissolve insoluble Na picrate. The Et₂O extract is dried over solid NaOH and then shaken with Al₂O₃ (Merck for chromatography) until the yellow color of traces of picric acid disappears (this color cannot be removed by repeated shaking with 5-10 M aq NaOH). The extract is concentrated to 50mL and dry HCl is bubbled through until separatrtn of the white crystals of the *cis-HCl* is complete. These are washed with Et₂O, dried at 100° and recrystd from EtOH + EtOAc to yield pure *cis-Hydrochloride* **m 182-183°** (dried in a vac desiccator over KOH) with IR (KBr) ν_{\max} 2920, 2820, 1582, 1470, 1445, 1410, 1395, 1313, 1135, 1080, 990, 870 cm⁻¹. The pure *free base* is prepared by dissolving the *hydrochloride* in 10 M aq NaOH, extracted with Et₂O, dried over solid KOH, fild and distd in vac. It has IR (film) ν_{\max} 2920, 2820, 2720, 2560, 1584, 1470, 1445, 1415, 1395, 1315, 1300, 1135, 1080, 1020, 990, 873 cm⁻¹. The ¹H NMR in CDCl₃ is characteristically different from that of the *trans*-isomer. [Armarego *J Chem Soc (C)* 377 1967; Gray and Heitmeier *J Am Chem Soc* 80 6274 1958; Witkop *J Am Chem Soc* 70 2617 1948; Skita *Chem Ber* 57 1982 1924; Helfer *Helv Chim Acta* 6 799 1923.]

***trans*-Decahydroisoquinoline** [2744-09-4] **M 139.2, b 106°/15mm, pK²⁰ 11.32.** This is purified as the *cis*-isomer above. The *trans-picrate* has **m 175-176°**, and the *trans-hydrochloride* has **m 221-222°** and has IR (KBr) ν_{\max} 2930, 3800, 1589, 1450, 1400, 1070, 952, 837 cm⁻¹. The pure *free base* was prepared as above and had IR (film) ν_{\max} 2920, 2820, 2720, 2560, 1584, 1470, 1445, 1415, 1395, 1315, 1300, 1135, 1080, 1020, 990, 873 cm⁻¹. The ¹H NMR in CDCl₃ is characteristically different from that of the *cis*-isomer. (references as above and Helfer *Helv Chim Acta* 9 818 1926).

Decahydronaphthalene (decalin, mixed isomers) [91-17-8] **M 138.2, b 191.7°, d 0.886, n 1.476.** Stirred with conc H₂SO₄ for some hours. Then the organic phase was separated, washed with water, saturated aqueous Na₂CO₃, again with water, dried with CaSO₄ or CaH₂ (and perhaps dried further with Na), filtered and distd under reduced pressure (**b 63-70°/10mm**). Also purified by repeated passage through long columns of silica gel previously activated at 200-250°, followed by distn from LiAlH₄ and storage under N₂. Type 4A molecular sieves can be used as a drying agent. Storage over silica gel removes water and other polar substances.

***cis*-Decahydronaphthalene** [493-01-6] **M 138.2, f -43.2°, b 195.7°, d 0.897, n 1.48113.** Purification methods described for the mixed isomers are applicable. The individual isomers can be separated by very efficient fractional distn, followed by fractional crystn by partial freezing. The *cis*-isomer reacts preferentially with AlCl₃ and can be removed from the *trans*-isomer by stirring the mixture with a limited amount of AlCl₃ for 48h at room temperature, filtering and distilling.

***trans*-Decahydronaphthalene** [493-02-7] **M 138.2, f -30.6°, b 187.3°, d 0.870, n 1.46968.** See purification of *cis*-isomer above.

***cis*-Decahydroquinoline** [10343-99-4] **M 139.2, b 207-208°/708mm, pK²⁰ 11.29.** It is available as a *cis-trans*-mixture (**b 70-73°/10mm**, Aldrich, ~ 18% *cis*-isomer [2051-28-7]), but the isomers can be fractionated in a spinning band column (1~1.5 metre, type E) at atmospheric pressure and collecting 2mL fractions with a distillation rate of 1 drop in 8-10sec. The lower boiling fraction solidifies and contains the *trans*-isomer (see below, **m 48°**). The higher boiling fraction **b 207-208°/708mm**, remains liquid and is mostly the *cis*-isomer. This is reacted with PhCOCl and M aq NaOH to yield the *N-benzoyl derivative* **m 96°** after recrystn from pet ether (**b 80-100°**). It is hydrolysed with 20% aq HCl by refluxing overnight. PhCO₂H is fild off, the filtrate is basified with 5M aq NaOH and extracted with Et₂O. The dried extract (Na₂SO₄) is satd with dry HCl gas and the *cis-decahydroquinoline hydrochloride* which separates has **m 222-224°** after washing with Et₂O and drying at 100°; and has IR (KBr) ν_{\max} 2900, 2780, 2560, 1580, 1445, 1432, 1403, 1165, 1080,

1036, 990, 867 cm^{-1} . The *free base* is obtained by dissolving the *hydrochloride* salt in 5M aq NaOH, extracting with Et_2O and drying the extract (Na_2SO_4), evaporating and distilling the residue; it has IR (film) ν_{max} 2900, 2840, 2770, 1445, 1357, 1330, 1305, 1140, 1125, 1109, 1068, 844 cm^{-1} . The ^1H NMR in CDCl_3 is characteristically different from that of the *trans*-isomer. [Armarego *J Chem Soc (C)* 377 1967; Hückel and Stepf *Justus Liebigs Ann Chem* 453 163 1927; Bailey and McElvain *J Am Chem Soc* 52 4013 1930.]

***trans*-Decahydroquinoline** [767-92-0] M 139.2, m 48°, b 205-206°/708mm, pK^{20} 11.29. The lower boiling fraction from the preceding spinning band column fractionation of the commercial *cis-trans*-mixture (~ 20:60; see the *cis*-isomer above) solidifies readily (m 48°) and the receiver has to be kept hot with warm water. It is further purified by conversion to the *Hydrochloride* m 285-286° after recrystn from EtOH/AcOEt . This has IR (KBr) ν_{max} 2920, 2760, 2578, 2520, 1580, 1455, 1070, 1050, 975, 950, 833 cm^{-1} . The *free base* is prepared as for the *cis*-isomer above and distd; and has IR (film, at ca 50°) ν_{max} 2905, 2840, 2780, 1447, 1335, 1305, 1240, 1177, 1125, 987, 900, 835 cm^{-1} . The ^1H NMR in CDCl_3 is characteristically different from that of the *cis*-isomer. [Armarego *J Chem Soc (C)* 377 1967; Hückel and Stepf *Justus Liebigs Ann Chem* 453 163 1927; Bailey and McElvain *J Am Chem Soc* 52 4013 1930; Prelog and Szpilfogel *Helv Chim Acta* 28 1684 1945.]

***n*-Decane** [124-18-5] M 142.3, b 174.1°, d 0.770, n 1.41189, n^{25} 1.40967. It can be purified by shaking with conc H_2SO_4 , washing with water, aqueous NaHCO_3 , and more water, then drying with MgSO_4 , refluxing with Na and distilling. Passed through a column of silica gel or alumina. It can also be purified by azeotropic distn with 2-butoxyethanol, the alcohol being washed out of the distillate, using water; the decane is next dried and redistilled. It can be stored with NaH. Further purification can be achieved by preparative gas chromatography on a column packed with 30% SE-30 (General Electric methyl-silicone rubber) on 42/60 Chromosorb P at 150° and 40psig, using helium [Chu *J Chem Phys* 41 226 1964]. Also purified by zone refining.

Decan-1,10-diol [112-47-0] M 174.3, m 72.5-74°. Crystd from dry ethylene dichloride.

***n*-Decanol (*n*-decyl alcohol)** [112-30-1] M 158.3, f 6.0°, b 110-119°/0.1mm, d 0.823, n 1.434. Fractionally distd in an all-glass unit at 10mm pressure (b 110°), then fractionally crystd by partial freezing. Also purified by preparative GLC, and by passage through alumina before use.

***n*-Decyl bromide** [112-29-8] M 221.2, b 117-118°/15.5mm, d 1.066. Shaken with H_2SO_4 , washed with water, dried with K_2CO_3 , and fractionally distd.

Decyltrimethylammonium bromide [2082-84-0] M 280.3, m 239-242°. Crystd from 50% (v/v) $\text{EtOH}/\text{diethyl ether}$, or from acetone and washed with ether. Dried under vacuum at 60°. Also recrystd from EtOH and dried over silica gel. [McDonnell and Kraus *J Am Chem Soc* 73 2170 1952; Dearden and Wooley *J Phys Chem* 91 2404 1987.]

(+)-Dehydroabietylamine (abieta-8,11,13-triene-18-ylamine) [1446-61-3] M 285.5, m 41°, 42.5-45°, b 192-193°/1mm, 250°/12mm, n_D^{40} 1.546, $[\alpha]_{546}^{20} +51^\circ$ (c 1, EtOH), $\text{pK}_{\text{Est}} \sim 10.3$. The crude base is purified by converting 2g of base in toluene (3.3mL) into the acetate salt by heating at 65-70° with 0.46g of AcOH and the crystals are collected and dried (0.96g from two crops; m 141-143°). The acetate salt is dissolved in warm H_2O , basified with aqueous NaOH and extracted with $^*\text{C}_6\text{H}_6$. The dried extract (MgSO_4) is evaporated in vacuum leaving a viscous oil which crystallises and can be distd. [Gottstein and Cheney *J Org Chem* 30 2072 1965.] The *picrate* has m 234-236° (from aq MeOH), and the *formate* has m 147-148° (from heptane).

Dehydro-L(+)-ascorbic acid [490-83-5] M 174.1, m 196°(dec), $[\alpha]_{546}^{20} +42.5^\circ$ (c 1, H_2O), pK 3.90. Crystd from MeOH

7-Dehydrocholesterol [434-16-2] M 384.7, m 142-143°, $[\alpha]_D^{20} -122^\circ$ (c 1, CHCl_3). Crystd from MeOH .

Dehydrocholic acid [81-23-2] M 402.5, m 237°, $[\alpha]_{546}^{20} -159^\circ$ (c 1, in CHCl_3), $\text{pK}_{\text{Est}} \sim 4.9$. Crystd from acetone.

Dehydroepiandrosterone [54-43-0] M 288.4, m 140-141° and 152-153° (dimorphic), $[\alpha]_{\text{D}}^{20} +13^\circ$ (c 3, EtOH). Crystd from MeOH and sublimed in vacuum.

Delphinine [561-07-9] M 559.7, m 197-199°, $[\alpha]_{\text{D}}^{20} +26^\circ$ (EtOH). Crystd from EtOH.

3-Deoxy-D-allose [6605-21-6] M 164.2, $[\alpha]_{\text{D}}^{20} +8^\circ$ (c 0.25 in H_2O). Obtained from diethyl ether as a colourless syrup.

Deoxybenzoin [451-40-1] M 196.3, m 60°, b 177°/12mm, 320°/760mm. Crystd from EtOH.

Deoxycholic acid [83-44-3] M 392.6, m 171-174°, 176°, 176-178°, $[\alpha]_{546}^{20} +64^\circ$ (c 1, EtOH), $[\alpha]_{\text{D}}^{20} +55^\circ$ (c 2.5, EtOH), pK 6.58. Refluxed with CCl_4 (50mL/g), filtered, evaporated under vacuum at 25°, recrystd from acetone and dried under vacuum at 155° [Trenner et al. *J Am Chem Soc* 76 1196 1954]. A soln of (cholic acid-free) material (100mL) in 500mL of hot EtOH was filtered, evaporated to less than 500mL on a hot plate, and poured into 1500mL of cold diethyl ether. The ppte, filtered by suction, was crystd twice from 1-2 parts of absolute EtOH, to give an alcoholate, m 118-120°, which was dissolved in EtOH (100mL for 60g) and poured into boiling water. After boiling for several hours the ppte was filtered off, dried, ground and dried to constant weight [Sobotka and Goldberg *Biochem J* 26 555 1932]. Deoxycholic acid was also freed from fatty acids and cholic acid by silica gel chromatography by elution with 0.5% acetic acid in ethyl acetate [Tang et al. *J Am Chem Soc* 107 4058 1985]. It can also be recrystd from butanone. Its solubility in H_2O at 15° is 0.24g/L but in EtOH it is 22.07g/L.

11-Deoxycorticosterone (21-hydroxyprogesterone) [64-85-7] M 330.5, m 141-142°, $[\alpha]_{546}^{20} +178^\circ$ and $[\alpha]_{546} +223^\circ$ (c 1, EtOH). Crystd from diethyl ether.

2-Deoxy- β -D-galactose [1949-89-9] M 164.2, m 126-128°, $[\alpha]_{\text{D}}^{20} +60^\circ$ (c 4, H_2O). Crystd from diethyl ether.

2-Deoxy- α -D-glucose [154-17-6] M 164.2, m 146°, $[\alpha]_{\text{D}}^{20} +46^\circ$ (c 0.5, H_2O after 45h). Crystd from MeOH/acetone.

6-Deoxy-D-glucose (D-quinovose) [7658-08-4] M 164.2, m 146°, $[\alpha]_{\text{D}}^{20} +73^\circ$ (after 5 min) and $+30^\circ$ (final, after 3h) (c 8.3, H_2O). It is purified by recrystn from EtOAc and is soluble in H_2O , EtOH but almost insoluble in Et_2O and Me_2CO . [Srivastava and Lerner *Carbohydr Res* 64 263 1978; NMR: Angyal and Pickles *Aust J Chem* 25 1711 1972.]

2-Deoxy- β -L-ribose [18546-37-7] M 134.1, m 77°, 80°, $[\alpha]_{\text{D}}^{25} +91.7^\circ$ (c 7, pyridine, 40° final). Crystd from diethyl ether.

2-Deoxy- β -D-ribose [533-67-5] M 134.1, m 86-87°, 87-90°, $[\alpha]_{\text{D}}^{20} -56^\circ$ (c 1, H_2O after 24h). Crystd from diethyl ether.

Desyl bromide (α -bromo-desoxybenzoin, ω -bromo- ω -phenyl acetophenone) [484-50-0] M 275.2, m 57.1-57.5°. Crystd from 95% EtOH.

Desyl chloride (α -chloro-desoxybenzoin, ω -chloro- ω -phenyl acetophenone) [447-31-4] M 230.7, m 62-64°, 66-67°, 67.5°, 68°. For the purification of small quantities recrystallise from pet ether (b 40-60°), but use MeOH or EtOH for larger quantities. For the latter solvent, dissolve 12.5g of chloride in 45mL of boiling EtOH (95%), filter and the filtrate yields colourless crystals (7.5g) on cooling. A further crop (0.9g) can be obtained by cooling in an ice-salt bath. It turns brown on exposure to sunlight but it is

stable in sealed dark containers. [Henley and Turner *J Chem Soc* 1182 1931; *Org Synth Coll Vol II* 159 1943.]

Dexamethasone (9- α -fluoro-16- α -methylprednisolone) [50-02-2] **M 392.5, m 262-264 $^{\circ}$, 268-271 $^{\circ}$, $[\alpha]_D^{25} +77.5^{\circ}$ (c 1, dioxane).** It has been recrystallised from Et₂O or small volumes of EtOAc. Its solubility in H₂O in 10 mg/100mL at 25 $^{\circ}$; and is freely soluble in Me₂CO, EtOH and CHCl₃. [Arth et al. *J Am Chem Soc* 80 3161 1958; for the β -methyl isomer see Taub et al. *J Am Chem Soc* 82 4025 1960.]

Dexamethasone 21-acetate (9- α -fluoro-16- α -methylprednisolone-21-acetate) [1177-87-3] **M 434.5, m 215-225 $^{\circ}$, 229-231 $^{\circ}$, $[\alpha]_D^{25} +77.6^{\circ}$ (c 1, dioxane), +73 $^{\circ}$ (c 1, CHCl₃).** Purified on neutral Al₂O₃ using CHCl₃ as eluent, fraction evaporated, and recrystd from CHCl₃. UV has λ_{\max} at 239nm. [Oliveto et al. *J Am Chem Soc* 80 4431 1958].

Diacetamide [625-77-4] **M 101.1, m 75.5-76.5 $^{\circ}$, b 222-223 $^{\circ}$.** Purified by crystn from MeOH [Arnett and Harrelson *J Am Chem Soc* 109 809 1987].

Diacetoxyiodobenzene (iodobenzenediacetate) [3240-34-4] **M 322.1, m 163-165 $^{\circ}$.** Purity can be checked by treatment with H₂SO₄ then KI and the liberated I₂ estimated with standard thiosulfate. It has been recrystd from 5M acetic acid and dried overnight in a vac desiccator over CaCl₂. The surface of the crystals may become slightly yellow but this does not affect its usefulness. [Sharefkin and Saltzman *Org Synth Coll Vol V* 600 1973.]

1,2-Diacetyl benzene [704-00-7] **M 162.2, m 39-41 $^{\circ}$, 41-42 $^{\circ}$, b 110 $^{\circ}$ /0.1mm, 148 $^{\circ}$ /20mm.** Purified by distn and by recrystn from pet ether. The *bis-2,4-dinitrophenyl hydrazone* has m 221 $^{\circ}$ dec. [Halford and Weissmann *J Org Chem* 17 1646 1952; Riemschneider and Kassahn *Chem Ber* 92 1705 1959.]

1,4-Diacetyl benzene [1009-61-6] **M 162.2, m 113-5-114.2 $^{\circ}$.** Crystd from *benzene and vacuum dried over CaCl₂. Also dissolved in acetone, treated with Norit, evapd and recrystd from MeOH [Wagner et al. *J Am Chem Soc* 108 7727 1986].

(+)-Di-O-acetyl-L-tartaric anhydride [(R,R)-2,3-diacetoxysuccinic anhydride] [6283-74-5] **M 216.2, m 129-132 $^{\circ}$, 133-134 $^{\circ}$, 135 $^{\circ}$, 137.5 $^{\circ}$, $[\alpha]_D^{20} +97.2^{\circ}$ (c 0.5, dry CHCl₃).** If the IR is good, i.e. no OH bands, then keep in a vacuum desiccator overnight (over P₂O₅/paraffin) before use. If OH bands are present then reflux 4g in Ac₂O (12.6mL) containing a few drops of conc H₂SO₄ for 10min (use a relatively large flask), pour onto ice, collect the crystals, wash with dry *C₆H₆ (2 x 2mL), stir with 17mL of cold Et₂O, filter and dry in a vacuum desiccator as above, and store in dark evacuated ampoules under N₂ in small aliquots. It is not very stable in air; the melting point of the crystals drop one degree in the first four days then remains constant (132-134 $^{\circ}$). If placed in a stoppered bottle it becomes gummy and the m falls 100 $^{\circ}$ in three days. Recrystn leads to decomposition. If good quality anhydride is required it should be prepared fresh from tartaric acid. It sublimes in a CO₂ atmosphere. [*Org Synth Coll Vol IV* 242 1963.]

Diallyl amine (N-2-propenyl-2-propen-1-amine) [124-02-7] **M 97.2, b 107-111 $^{\circ}$ /760mm, 112 $^{\circ}$ /760mm, d_4^{20} 0.789, n_D^{20} 1.440, pK^{20} 9.42.** Keep over KOH pellets overnight, decant and distil from a few pellets of KOH at atm pressure (b 108-111 $^{\circ}$), then fractionate through a Vigreux column. [Vliet *J Am Chem Soc* 46 1307 1924; *Org Synth Coll Vol I* 201 1941.] The *hydrochloride* has m 164-165 $^{\circ}$ (from Me₂CO + EtOH). [Butler and Angels *J Am Chem Soc* 79 3128 1957.]

(+)-N,N'-Diallyl tartramide (DATD) [58477-85-3] **M 228.3, m 184 $^{\circ}$, $[\alpha]_{546} +141^{\circ}$ (c 3, MeOH).** Wash with Et₂O containing 10% EtOH until the washings are clear and colourless, and dry *in vacuo*. [*FEBS Lett* 7 293 1970.]

Diamantane [2292-79-7] **M 188.3, m 234-235 $^{\circ}$.** Purified by repeated crystn from MeOH or pentane. Also dissolved in CH₂Cl₂, washed with 5% aq NaOH and water, and dried (MgSO₄). The soln was concentrated to a small volume, an equal weight of alumina was added, and the solvent evaporated. The residue was placed on

an activated alumina column (ca 4 x weight of diamantane) and eluted with pet ether (b 40-60°). Eight sublimations and twenty zone refining experiments gave material **m** 251° of 99.99% purity by differential analysis [*Tetrahedron Lett* 3877 1970; *J Chem Soc (C)* 2691 1972].

3,6-Diaminoacridine hydrochloride [952-23-8] **M** 245.7, **m** 270°(dec), ϵ_{456} 4.3×10^4 , pK_1^{20} 1.5, pK_2^{20} 9.60 (9.65 free base). First purified by pptn of the free base by adding aq NH_3 soln to an aq soln of the hydrochloride or hydrogen sulfate, drying the ppt and subliming at 0.01mm Hg [Müller and Crothers *Eur J Biochem*, 54 267 1975].

3,6-Diaminoacridine sulfate (proflavin sulfate) [1811-28-5] **M** 516.6, **m** >300°(dec), λ_{max} 456nm. An aqueous soln, after treatment with charcoal, was concentrated, chilled overnight, filtered and the ppt was rinsed with a little diethyl ether. The ppt was dried in air, then overnight in a vacuum oven at 70°.

1,3-Diaminoadamantane [10303-95-4] **M** 164.3, **m** 52°, $\text{pK}_{\text{Est}(1)}$ ~8.6, $\text{pK}_{\text{Est}(2)}$ ~10.6. Purified by zone refining.

1,4-Diaminoanthraquinone [128-95-0] **M** 238.3, **m** 268°. Purified by thin-layer chromatography on silica gel using toluene/acetone (9:1) as eluent. The main band was scraped off and extracted with MeOH. The solvent was evaporated and the quinone was dried in a drying pistol [Land, McAlpine, Sinclair and Truscott *J Chem Soc, Faraday Trans 1* 72 2091 1976]. Crystd from EtOH in dark violet crystals.

1,5-Diaminoanthraquinone [129-44-2] **M** 238.3, **m** 319°. Recrystd from EtOH or acetic acid [Flom and Barbara *J Phys Chem* 89 4481 1985].

2,6-Diaminoanthraquinone [131-14-6] **M** 238.3, **m** 310-320°. Crystd from pyridine. Column-chromatographed on Al_2O_3 /toluene to remove a fluorescent impurity, then recrystd from EtOH.

3,3'-Diaminobenzidine tetrahydrochloride (2H₂O) [7411-49-6] **M** 396.1, **m** >300°(dec), $\text{pK}_{\text{Est}(1)}$ ~3.3, $\text{pK}_{\text{Est}(2)}$ ~4.7 (free base). Dissolved in water and pted by adding conc HCl, then dried over solid NaOH.

3,4-Diaminobenzoic acid [619-05-6] **M** 152.2, **m** 213°(dec), 228-229°, pK_1^{25} 2.57 (4-NH₂), pK_2^{25} 3.39 (3-NH₂), $\text{pK}_{\text{Est}(3)}$ ~5.1 (CO₂H). Crystd from H₂O or toluene.

3,5-Diaminobenzoic acid [535-87-5] **M** 152.2, **m** 235-240°(dec), pK^{25} 5.13 (CO₂H). Crystd from water. *Dihydrochloride* has **m** 226-228°(dec).

4,4'-Diaminobenzophenone [611-98-3] **M** 212.3, **m** 242-244°, 243-245°, 246.5-247.5° (after sublimation at 0.0006 mm), pK_1^{25} 1.37, pK_2^{25} 2.92. Purified by recrystn from EtOH and by sublimation in high vacuum. The *dihydrochloride* has **m** 260° dec (from EtOH) and the *thiosemicarbazone* has **m** 207-207.5° dec (from aq EtOH). [Kuhn et al. *Chem Ber* 75 711 1942.]

1,4-Diaminobutane dihydrochloride (putrescine 2HCl) [333-93-7] **M** 161.1, **m** >290°, pK_1^{25} 9.63, pK_2^{25} 10.80. Crystd from EtOH/water.

1,2-Diamino-4,5-dichlorobenzene [5348-42-5] **M** 177.0, **m** 163°, $\text{pK}_{\text{Est}(1)}$ ~0, $\text{pK}_{\text{Est}(2)}$ ~2.9. Refluxed with activated charcoal in CH_2Cl_2 , followed by recrystn from diethyl ether/pet ether or pet ether [Koolar and Kochi *J Org Chem* 52 4545 1987].

2,2'-Diaminodiethylamine (diethylenetriamine) [111-40-0] **M** 103.2, **b** 208°, **d** 0.95, **n** 1.483, pK_1^{25} 4.34, pK_2^{25} 9.13, pK_3^{25} 9.94. Dried with Na and distd, preferably under reduced pressure, or in a stream of N₂. § Polymer-bound diethylenetriamine is commercially available.

4,5-Diamino-2,6-dihydroxypyrimidine (diamino uracil) sulfate [32014-70-3] **M** 382.3, **m** >300°, pK_1^{20} 1.7, pK_2^{20} 3.20, pK_3^{20} 4.56. The salt is quite insoluble in H₂O but can be converted to the

free base which is recrystd from H₂O and converted to the sulfate by addition of the required amount of H₂SO₄. The *hydrochloride* has *m* 300-305° dec and can be used to prepare the sulfate by addition of H₂SO₄; It is more soluble than the sulfate. The *perchlorate* has *m* 252-254°. The free base has λ_{\max} 260nm (log ϵ 4.24) in 0.1M HCl. [Bogert and Davidson *J Am Chem Soc* **86** 1668 1933; Bredereck et al. *Chem Ber* **86** 850 1953; *Org Synth Coll Vol IV* 247 1963; Barlin and Pfeleiderer *J Chem Soc (B)* 1425 1971.]

5,6-Diamino-1,3-dimethyluracil hydrate (5,6-diamino-1,3-dimethyl-2-pyrimidine-2,4-dione hydrate) [5440-00-6] *M* 188.2, *m* 205-208° dec, 209° dec, 210°dec, *pK*₁ 1.7, *pK*₂ 4.6. Recryst from EtOH. The *hydrochloride* has *m* 310° (from MeOH) and the *perchlorate* has *m* 246-248°. [UV: Bredereck et al. *Chem Ber* **92** 583 1959; Taylor et al. *J Am Chem Soc* **77** 2243 1955.]

4,4'-Diamino-3,3'-dinitrobiphenyl [6271-79-0] *M* 274.2, *m* 275°, *pK*_{Est} ~-0.2. Crystd from aqueous EtOH.

4,4'-Diaminodiphenylamine [537-65-5] *M* 199.3, *m* 158°, *pK*_{Est} ~5.0. Crystd from water.

4,4'-Diaminodiphenylmethane [101-77-9] *M* 198.3, *m* 91.6-92°, *pK*_{Est} ~4.9. Crystd from water, 95% EtOH or *benzene.

3,3'-Diaminodipropylamine [56-18-8] *M* 131.2, *b* 152°/50mm, *d* 0.938, *n* 1.481, *pK*₁²⁵ 7.72, *pK*₂²⁵ 9.57, *pK*₃²⁵ 10.65. Dried with Na and distd under vacuum.

6,9-Diamino-2-ethoxyacridine (Ethacridine) [442-16-0] *M* 257.3, *m* 226°, *pK*_{Est} ~11.5. Crystd from 50% EtOH.

2,7-Diaminofluorene [524-64-4] *M* 196.3, *m* 165°, *pK*_{Est} ~4.6. Recrystd from H₂O.

2,4-Diamino-6-hydroxypyrimidine [56-06-4] *M* 126.1, *m* 260-270°(dec), *pK*₁²⁵ 1.34, *pK*₂²⁵ 3.27, *pK*₃²⁵ 10.83. Recrystd from H₂O.

4,5-Diamino-6-hydroxypyrimidine hemisulfate [102783-18-6] *M* 350.3, *m* 268°, 270°, *pK*₁²⁵ 1.34, *pK*₂²⁵ 3.57, *pK*₃²⁵ 9.86. Recrystd from H₂O. The free base also recrystallises from H₂O (*m* 239°). [UV: Mason *J Chem Soc* 2071 1954; Elion et al. *J Am Chem Soc* **74** 411 1952.]

1,5-Diaminonaphthalene [2243-62-1] *M* 158.2, *m* 190°, *pK*²⁵ 4.12. Rerystd from boiling H₂O, but is wasteful due to poor solubility. Boil in chlorobenzene (charcoal), filter hot and cool the filtrate. This gives colourless crystals. Dry in a vac till free from chlorobenzene (odour), and store away from light.

1,8-Diaminonaphthalene [479-27-6] *M* 158.2, *m* 66.5°, *b* 205°/12mm, *pK* 4.44. Crystd from water or aqueous EtOH, and sublimed in a vacuum. *N,N'*-DiMe deriv [20734-56-9] has *m* 103-104° and *pK* 5.61; *N,N,N'*-TriMe deriv [20734-57-0] has *m* 29-30° and *pK* 6.43. [Hodgson et al. *J Chem Soc* 202 1945.]

2,3-Diaminonaphthalene [771-97-1] *M* 158.2, *m* 199°, *pK*²¹ 3.54 (in 50% aq EtOH). Crystd from water, or dissolved in 0.1M HCl, heated to 50°. After cooling, the soln was extracted with decalin to remove fluorescent impurities and centrifuged.

1,8-Diaminooctane [373-44-4] *M* 144.3, *m* 50-52°, 51-52°, 52-53°, *b* 121°/18 mm, 120°/24mm, *pK*₁²⁰ 10.1, *pK*₂²⁰ 11.0. Distil under vacuum in an inert atmosphere(N₂, Ar), cool and store distillate in an inert atmosphere in the dark. The *dihydrochloride* has *m* 273-274°. [Nae and Le *Helv Chim Acta* **15** 55 1955.]

2,4-Diamino-5-phenylthiazole (Amiphenazole) [490-55-1] *M* 191.3, *m* 163-164°(dec). Crystd from aqueous EtOH or water. Stored in the dark under N₂.

1,5-Diaminopentane [462-94-2] M 102.2, m 14-16°, b 78-80°/12mm, 101-103°/35mm, 178-180°/750mm, d_4^{20} 0.869, n_D^{20} 1.458, pK_1^{20} 10.02, pK_2^{20} 10.96. Purified by distn, after standing over KOH pellets (at room temp; i.e. liquid form). It has pK_a^{20} values of 10.02 and 10.96 in H₂O. Its *dihydrochloride* has m 275° (sublimes in vac), and its *tetraphenyl boronate* has m 164°. [Schwarzenbach et al. *Helv Chim Acta* 35 2333 1952.]

***d,l*-2,6-Diaminopimelic acid** [583-93-7] M 190.2, m 313-315°(dec) $pK_{Est(1)}$ ~2.2, $pK_{Est(2)}$ ~9.7. Crystd from water.

1,3-Diaminopropane dihydrochloride [10517-44-9] M 147.1, m 243°, pK_1^{25} 8.29, pK_2^{25} 10.30. Crystd from EtOH/water.

1,3-Diaminopropan-2-ol [616-29-5] M 90.1, m 38-40°, pK_1^{25} 7.94, pK_2^{25} 9.57. Dissolved in an equal amount of water, shaken with charcoal and vacuum distd at 68°/0.1mm. It is too viscous to be distd through a packed column.

***L(S)*-2,3-Diaminopropionic acid monohydrochloride (3-amino-L-alanine hydrochloride)** [1482-97-9] M 140.6, m 132-133°dec, 237°dec, $[\alpha]_D^{25}$ +26.1° (c 5.8, M HCl), pK_1^{25} 1.30, pK_2^{25} 6.79, pK_3^{25} 9.51. Forms needles from H₂O and can be recrystd from aqueous EtOH. [Gmelin et al. *Z Physiol Chem.* 314 28 1959; IR: Koegel et al. *J Am Chem Soc* 77 5708 1977.]

2,3-Diaminopyridine [452-58-4] M 109.1, m 116°, pK_1^{25} -0.50, pK_2^{25} 6.92. Crystd from *benzene and sublimed *in vacuo*.

2,6-Diaminopyridine [141-86-6] M 109.1, m 121.5° $pK_{Est(1)}$ <-6.0, $pK_{Est(2)}$ ~7.3. Crystd from *benzene and sublimed *in vacuo*.

3,4-Diaminopyridine [54-96-6] M 109.1, m 218-219°, pK_1^{20} 0.49, pK_2^{20} 9.14. Crystd from *benzene and stored under H₂ because it is deliquescent and absorbs CO₂.

***meso*-2,3-Diaminosuccinic acid** [23220-52-2] M 148.1, m 305-306°(dec, and sublimes), $pK_{Est(1)}$ ~3.6, $pK_{Est(2)}$ ~9.8. Crystd from water.

Diaminotoluene see **toluenediamine**.

3,5-Diamino-1,2,4-triazole (Guanazole) [1455-77-2] M 99.1, m 206° pK_1^{20} 4.43, pK_2^{20} 12.12. Crystd from water or EtOH.

2,5-Di-*tert*-amylhydroquinone [79-74-3] M 250.4, m 185.8-186.5°. Crystd under N₂ from boiling glacial acetic acid (7mL/g) plus boiling water (2.5mL/g) [Stolow and Bonaventura *J Am Chem Soc* 85 3636 1963].

Di-*n*-amyl phthalate [131-18-0] M 306.4, b 204-206°/11mm, d_4^{25} 1.023, n 1.489. Washed with aqueous Na₂CO₃, then distilled water. Dried with CaCl₂ and distd under reduced pressure. Stored in a vacuum desiccator over P₂O₅.

1,3-Diazaazulene (cycloheptimidazole) [275-94-5] M 130.1, m 120°. Recrystd repeatedly from de-aerated cyclohexane in the dark.

1,5-Diazabicyclo[4.3.0]non-5-ene (DBN, 2,3,4,,6,7,8-hexahydropyrrolo[1,2-*a*]-pyrimidine) [3001-72-7] M 124.2, b 96-98°/11mm, 100-102°/12mm, 118-121°/32mm, d_4^{20} 1.040, n_D^{20} 1.520, pK >13.0. Distd from BaO. It forms a *hydroiodide* by addn of 47% HI, dry and dissolve in MeCN, evaporate and repeat, recrystallise from EtOH then dry at 25°/1mm for 5h, then at 80°/0.03mm for 12h and store and dispense in a dry box, m 154-156° [Jaeger et al. *J Am Chem Soc* 101 717 1979]. The *methiodide* is recrystd from CHCl₃ + Et₂O, m 248-250°, and *hydrogen fumarate* has m 159-160° and is crystd from *iso*-

PrOH [Rokach et al. *J Med Chem* **22** 237 1979; Oediger et al. *Chem Ber* **99** 2012 1966; Reppe et al. *Justus Liebig's Ann Chem* **596** 210 1955].

1,4-Diazabicyclo[2.2.2]octane (DABCO, triethylenediamine TED) [280-57-9] **M 112.2, m 156-157°** (sealed tube), **pK₁²⁵ 2.97, pK₂²⁵ 8.82** Crystd from 95% EtOH, pet ether or MeOH/diethyl ether (1:1). Dried under vacuum over CaCl₂ and BaO. It can be sublimed *in vacuo*, and readily at room temperature. Also purified by removal of water during azeotropic distn of a *benzene soln. It was then recrystd twice from anhydrous diethyl ether under argon, and stored under argon [Blackstock et al. *J Org Chem* **52** 1451 1987].

1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU, 2,3,4,6,7,8,9,10-octahydropyrimidino[1,2-*a*]-azepine) [6674-22-2] **M 152.2, b 115°/11mm, d 1.023, n 1.522, pK_{Est} ~ >13.** Fractionally dist under vac. Also purified by chromatography on Kieselgel and eluting with CHCl₃/EtOH/25% aq NH₃ (15:5:2) and checked by IR and MS. [Oediger et al. *Chem Ber* **99** 2012 1962; *Angew Chem, Int Ed Engl* **6** 76 1967; Guggisberg et al. *Helv Chim Acta* **61** 1057 1978.]

1,8-Diazabiphenylene [259-84-7] **M 154.2, pK_{Est} ~4.4.** Recrystd from cyclohexane, then sublimed in a vacuum.

2,7-Diazabiphenylene [31857-42-8] **M 154.2, pK_{Est} ~4.5.** Recrystd from cyclohexane, then sublimed in a vacuum.

Diazoaminobenzene (1,3-diphenyltriazeno) [136-36-6] **M 197.2, m 99°.** Crystd from pet ether (b 60-80°), 60% MeOH/water or 50% aqueous EtOH (charcoal) containing a small amount of KOH. Also purified by chromatography on alumina/toluene and toluene-pet ether. Stored in the dark.

6-Diazo-5-oxo-L-norleucine [157-03-9] **M 171.2, m 145-155°(dec), [α]_D²⁰ +21° (c 5, EtOH), pK₁ 2.1, pK₂ 8.95.** Crystd from EtOH, aq EtOH or MeOH.

Dibenzalacetone [538-58-9] **M 234.3, m 112°.** Crystd from hot ethyl acetate (2.5mL/g) or EtOH.

Dibenz[*a,h*]anthracene [53-70-3] **M 278.4, m 266-267°.** The yellow-green colour (due to other pentacyclic impurities) has been removed by crystn from *benzene or by selective oxidation with lead tetraacetate in acetic acid [Moriconi et al. *J Am Chem Soc* **82** 3441 1960].

Dibenzo-18-crown-6 [14187-32-7] **M 360.4, m 163-164°.** Crystd from *benzene, *n*-heptane or toluene and dried under vacuum at room temperature for several days. [Szezygiel *J Phys Chem* **91** 1252 1987.]

Dibenzo-18-crown-8 [14174-09-5] **M 448.5, m 103-106°.** Recrystd from EtOH, and vacuum dried at 60° over P₂O₅ for 16hours. [Delville et al. *J Am Chem Soc* **109** 7293 1987.]

Dibenzofuran [132-64-9] **M 168.2, m 82.4°.** Dissolved in diethyl ether, then shaken with two portions of aqueous NaOH (2M), washed with water, separated and dried (MgSO₄). After evaporating the ether, dibenzofuran was crystd from aq 80% EtOH and dried under vacuum. [Cass et al. *J Chem Soc* 1406 1958.] High purity material was obtained by zone refining.

Dibenzopyran (xanthene) [92-83-1] **M 182.2, m 100.5°, b 310-312°.** See entry on p. 386.

Dibenzothiophene [132-65-0] **M 184.3, m 99°.** Purified by chromatography on alumina with pet ether, in a darkened room. Crystd from water or EtOH.

***trans*-1,2-Dibenzoylethylene** [959-28-4] **M 236.3, m 109-112°, 111°.** Recrystd from MeOH or EtOH as yellow needles [Koller et al. *Helv Chim Acta* **29** 512 1946]. The *dioxime* has m 210-211°dec from AcOH. [IR: Kuhn et al. *J Am Chem Soc* **72** 5058 1950; Yates *J Am Chem Soc* **74** 5375 1952; Erickson et al. *J Am Chem Soc* **73** 5301 1951.]

Dibenzoylmethane (1,3-diphenyl-1,3-propanedione) [120-46-7] **M 224.3, m 80°**. Crystd from pet ether or MeOH.

Di-*O*-benzoyl-(*R* and *S*)-tartaric acid (H₂O) [*R*-(+)- 17026-42-5; *S*-(-)- 2743-38-6] **M 376.3, m 88-89° (hydrate), 173° (anhydrous), [α]_D²⁰ (+) and (-) 136° (c 2, EtOH), [α]_D²⁰ (+) and (-) 117° (c 5, EtOH), pK_{Est(1)} ~2.9, pK_{Est(2)} ~4.2**. Crystd from water (18g from 400 mL boiling H₂O) and stir vigorously while cooling in order to obtain crystals; otherwise an oil will separate which solidifies on cooling. Dry in a vacuum desiccator over KOH-H₂SO₄ - yield 16.4g) as monohydrate, **m 88-89°**. It crystallises from xylene as the anhydrous acid, **m 173° (150-153°)**. It does not cryst from *C₆H₆, toluene, *C₆H₆-pet ether (oil), or CHCl₃-pet ether. [Butler and Cretcher *J Am Chem Soc* 55 2605 1933; *Tetrahedron* 41 2465 1085.]

2,3,6,7-Dibenzphenanthrene [222-93-5] **M 276.3, m 257°**. Crystd from xylene.

Dibenzyl amine [103-49-1] **M 197.3, m -26°, b 113-114°/0.1mm, 174-175°/6mm, 270°/250mm, 300° (partial dec), d₄²⁰ 1.027, n_D²⁰ 1.576, pK²⁵ 8.52**. Purified by distn in a vacuum. It causes burns to the skin. The *dihydrochloride* has **m 265-266°** after recrystn from MeOH-HCl, and the *tetraphenyl boronate* has **m 129-133°**. [Bradley and Maisey *J Chem Soc* 247 1954; Hall *J Phys Chem* 60 63 1956; Donetti and Bellora *J Org Chem* 37 3352 1972.]

Dibenzyl disulfide [150-60-7] **M 246.4, m 71-72°**. Crystd from EtOH.

Dibenzylethylenediamine (benzathine, DBED) [140-28-3] **M 240.4, m 26°, b 195°/4mm, d 1.02, n 1.563, pK_{Est(1)} ~ 5.9, pK_{Est(2)} ~ 8.9**. Dissolve in acid, extract with toluene, basify, extract with Et₂O, dry over solid KOH, evap and fractionate *in vacuo*. The *diacetate* cryst from H₂O by addn of EtOH, has **m 111° (sol in H₂O is ~25%)**. [Frost et al., *J Am Chem Soc* 71 3842 1949.]

1,3,4,6-Di-*O*-benzylidene-D-mannitol [28224-73-9] **M 358.4, m 192-195°, 193°, [α]_D²⁰ -11.9° (c 0.7, Me₂CO)**. Recryst from Et₂O in long fine needles. λ_{max} 256nm (ε 435) in 95% EtOH, R_F 0.21 (1:1 CCl₄-EtOAc) on TLC Silica Gel G. [Sinclair *Carbohydr Res* 12 150 1970; ORD, CD, NMR, IR, MS: Brecknell et al. *Aust J Chem* 29 1749 1976.]

Dibenzyl ketone (1,3-diphenyl-2-propanone) [102-04-5] **M 210.3, m 34.0°**. Fractionally crystd from its melt, then crystd from pet ether. Stored in the dark.

Dibenzyl malonate [15014-25-2] **M 284.3, b 188-190°/0.2mm, 193-196°/1mm, d₄²⁰ 1.158, n_D²⁰ 1.5452**. Dissolve in toluene, wash with aqueous NaHCO₃, H₂O, dry over MgSO₄, filter, evaporate and distil. [Ginsburg and Pappo *J Am Chem Soc* 75 1094 1953; Baker et al. *J Org Chem* 17 77 1952.]

Dibenzyl sulfide [538-74-9] **M 214.3, m 48.5°, 50°**. Crystd from EtOH/water (10:1), or repeatedly from Et₂O. Also chromatographed on Al₂O₃ (pentane as eluent), then recrystd from EtOH [Kice and Bowers *J Am Chem Soc* 84 2390 1962]. Vacuum dried at 30° over P₂O₅, fused under nitrogen and re-dried.

2,4-Dibromoaniline [615-57-6] **M 250.9, m 79-80°, pK²⁵ 1.87**. Crystd from aqueous EtOH.

9,10-Dibromoanthracene [523-27-3] **M 336.0, m 226°**. Recrystd from xylene and vacuum sublimed [Johnston et al. *J Am Chem Soc* 109 1291 1987].

***p*-Dibromobenzene** [106-37-6] **M 235.9, m 87.8°**. Steam distd, crystd from EtOH or MeOH and dried in the dark under vacuum. Purified by zone melting.

2,5-Dibromobenzoic acid [610-71-9] **M 279.9, m 157°, pK_{Est} ~1.5**. Crystd from water or EtOH.

4,4'-Dibromobiphenyl [92-86-4] **M 312.0, m 164°, b 355-360°/760mm**. Crystd from MeOH.

***trans*-1,4-Dibromobut-2-ene** [821-06-7] **M 213.9, m 54°, b 85°/10mm**. Crystd from ligroin.

α,α -Dibromodeoxybenzoin [15023-99-1] M 354.0, m 111.8-112.7°. Crystd from acetic acid.

Dibromodichloromethane [594-18-3] M 242.7, m 22°. Crystd repeatedly from its melt, after washing with aqueous $\text{Na}_2\text{S}_2\text{O}_3$ and drying with BaO.

1,3-Dibromo-5,5-dimethylhydantoin [77-48-5] M 285.9, m 190-192°dec, 190-193°dec. Recrystd from H_2O . Solubility in CCl_4 is 0.003 mol/L at 25° and 0.024 mol/L at 76.5°.

1,2-Dibromoethane [106-93-4] M 187.9, f 10.0°, b 29.1°/10mm, 131.7°/760mm, d 2.179, n^{15} 1.54160. Washed with conc HCl or H_2SO_4 , then water, aqueous NaHCO_3 or Na_2CO_3 , more water, and dried with CaCl_2 . Fractionally distd. Alternatively, kept in daylight with excess bromine for 2 hours, then extracted with aqueous Na_2SO_3 , washed with water, dried with CaCl_2 , filtered and distd. It can also be purified by fractional crystn by partial freezing. Stored in the dark.

4',5'-Dibromofluorescein [596-03-2] M 490.1, m 285°. Crystd from aqueous 30% EtOH.

5,7-Dibromo-8-hydroxyquinoline [521-74-4] M 303.0, m 196°, pK_1^{25} 5.84, pK_2^{25} 9.56. Crystd from acetone/EtOH. It can be sublimed.

Dibromomaleic acid [608-37-7] M 273.9, m 123.5°, 125°dec, pK_1^{25} 1.45, pK_2^{25} 4.62. It has been recrystd from Et_2O or $\text{Et}_2\text{O}-\text{CHCl}_3$. It is slightly soluble in H_2O , soluble also in AcOH but insoluble in $^*\text{C}_6\text{H}_6$ and CHCl_3 . [Salmony and Simonis *Chem Ber* 38 2583 1905; Ruggli *Helv Chim Acta* 3 566 1929.]

2,5-Dibromonitrobenzene [3460-18-2] M 280.9, m 84°. Crystd from acetone.

2,6-Dibromo-4-nitrophenol [99-28-5] M 280.9, m 143-144°, pK^{25} 3.39. Crystd from aq EtOH.

2,4-Dibromophenol [615-58-7] M 251.9, m 37°, 41-42°, b 154°/10mm, 239°/atm, pK^{25} 7.79. Crystd from CHCl_3 at -40°.

2,6-Dibromophenol [608-33-3] M 251.9, m 56-57°, b 138°/10mm, 255-256°/740mm, pK^{25} 6.67. Vacuum distd (at 18mm), then crystd from cold CHCl_3 or from EtOH/water.

1,3-Dibromopropane [109-64-8] M 201.9, f -34.4°, b 63-63.5°/26mm, 76-77°/40mm, 90°/80mm, 165°/atm, d 1.977, n 1.522. Washed with dilute aqueous Na_2CO_3 , then water. Dried and fractionally distd under reduced pressure.

2,6-Dibromopyridine [626-05-1] M 236.9, m 117-119°, 118.5-119°, b 249°/757.5mm, $\text{pK}_{\text{Est}} < 0$. Purified by steam distn then twice recrystd from EtOH. Does not form an HgCl_2 salt. [den Hertog and Wibaut *Rec Trav Chim Pays Bas* 51 381 1932.]

meso-2,3-Dibromosuccinic acid [526-78-3] M 275.9, m 288-290° (sealed tube, dec), pK_1^{20} 1.56, pK_2^{20} 2.71. Crystd from distilled water, keeping the temperature below 70°.

1,2-Dibromotetrafluoroethane [124-73-2] M 259.8, b 47.3°/760mm. Washed with water, then with weak alkali. Dried with CaCl_2 or H_2SO_4 and distd. [Locke et al. *J Am Chem Soc* 56 1726 1934.] Also purified by gas chromatography on a silicone DC-200 column.

α,α' -Dibromo-*o*-xylene [91-13-4] M 264.0, m 95°, b 129-130°/4.5mm. Crystd from CHCl_3

α,α' -Dibromo-*m*-xylene [626-15-3] M 264.0, m 77°, b 156-160°/12mm. Crystd from acetone.

α,α' -Dibromo-*p*-xylene [623-24-5] M 264.0, m 145-147°, b 155-158°/12-15mm, 245°/760mm. Crystd from $^*\text{benzene}$ or chloroform.

Di-*n*-butylamine [111-92-2] M 129.3, b 159°, n 1.41766, d 0.761, pK²⁵ 11.25. Dried with LiAlH₄, CaH₂ or KOH pellets, filtered and distd from BaO or CaH₂.

α-Dibutylamino-α-(*p*-methoxyphenyl)acetamide (Ambucetamide) [519-88-0] M 292.4, m 134°. Crystd from EtOH containing 10% diethyl ether.

2,5-Di-*tert*-butyl aniline [21860-03-7] M 205.4, m 103-104°, 103-106°, pK²⁵ 3.34 (50% aq MeOH), 3.58 (90% aq MeOH). Recrystd from EtOH in fine needles after steam distn. It has a pK^{a25} of 3.58 (50% aq EtOH) and 3.34 (90% aq MeOH). The *tosylate* has m 164° (from AcOH). [Bell and Wilson *J Chem Soc* 2340 1956; Carpenter et al. *J Org Chem* 16 586 1951; Bartlett et al. *J Am Chem Soc* 76 2349 1954.]

Di-*tert*-butylazodicarboxylate [870-50-8] M 230.3, m 90-92°. Cryst from ligroin. Best purified by covering the dry solid (22g) with pet ether (b 30-60°, 35-40 mL) and adding ligroin (b 60-90°) to the boiling soln until the solid dissolves. On cooling, large lemon yellow crystals of the ester separate (~20g), m 90.7-92°. Evapn of the filtrate gives a further crop of crystals [Carpino and Crowley *Org Synth* 44 18 1964].

***p*-Di-*tert*-butylbenzene** [1012-72-2] M 190.3, m 80°. Crystd from diethyl ether, EtOH and dried under vacuum over P₂O₅ at 55°. [Tanner et al. *J Org Chem* 52 2142 1987.]

2,6-Di-*tert*-butyl-1,4-benzoquinone [719-22-2] M 220.3, m 66-67°. It can be recrystd from MeOH and sublimed in a vacuum.

3,5-Di-*tert*-butyl-*o*-benzoquinone [3383-21-9] M 220.3, m 112-114°, 113-114°. It can be recrystd from MeOH or pet ether, and forms fine red plates or rhombs. [Flaig et al. *Justus Liebigs Ann Chem* 597 196 1955; IR: Ley and Müller *Chem Ber* 89 1402 1956.]

3,5-Di-*tert*-butyl catechol [1020-31-1] M 222.3, m 99°, 99-100°, pK_{Est(1)} ~11.0, pK_{Est(2)} ~13.1. Recrystd from pet ether. [Ley and Müller *Chem Ber* 89 1402 1956; UV Flaig et al. *Z Naturforschung* 10b 668 1955.] Also crystd three times from pentane [Funabiki et al. *J Am Chem Soc* 108 2921 1986].

Dibutylcarbitol [di(ethyleneglycol)dibutyl ether] [112-73-2] M 218.3, b 125-130°/0.1mm, d 0.883, n 1.424. Freed from peroxides by slow passage through a column of activated alumina. The eluate was shaken with Na₂CO₃ (to remove any remaining acidic impurities), washed with water, and stored with CaCl₂ in a dark bottle [Tuck *J Chem Soc* 3202 1957].

2,6-Di-*tert*-butyl-*p*-cresol (2,6-di-*tert*-butyl-4-methylphenol, butylatedhydroxytoluene, BHT) [128-37-0] M 230.4, m 71.5°, pK²⁵ 12.23. Dissolved in *n*-hexane at room temperature, then cooled with rapid stirring, to -60°. The ppte was separated, redissolved in hexane, and the process was repeated until the mother liquor was no longer coloured. The final product was stored under N₂ at 0° [Blanchard *J Am Chem Soc* 82 2014 1960]. Also crystd from EtOH, MeOH, *benzene, *n*-hexane, methylcyclohexane or pet ether (b 60-80°), and dried under vacuum.

Di-*tert*-butyl dicarbonate (di-*tert*-butyl pyrocarbonate) [24424-99-5] M 218.3, m 23° (21-22°), b 55-56°/0.15mm, 62-65°/0.4mm, d 0.950, n 1.409. Melt by heating at ~35°, and distil in vac. If IR and NMR (ν 1810m 1765 cm⁻¹, δ in CCl₄ 1.50 singlet) suggest very impure then wash with equal vol of H₂O containing citric acid to make the aqueous layer slightly acidic, collect the organic layer and dry over anhyd MgSO₄ and distil in vac. [Pope et al. *Org Synth* 57 45 1977.] **FLAMMABLE.**

2,6-Di-*tert*-butyl-4-dimethylaminomethylphenol [88-27-7] M 263.4, m 93-94°, b 172°/30mm, pK_{Est} ~12.0. Crystd from *n*-hexane.

Di-*tert*-butyldiperphthalate [2155-71-7] M 310.3, m dec explosively, CARE. Crystd from diethyl ether. Dried over H₂SO₄.

2,6-Di-*tert*-butyl-4-ethylphenol [4130-42-1] M 234.4, m 42-44°, pK_{Est} ~12.3. Crystd from aqueous EtOH or *n*-hexane.

***N,N*-Dibutyl formamide** [761-65-9] M 157.3, b 63°/0.1mm, 118-120°/15mm, 244-246°/760mm, d₄²⁰ 0.878, n_D²⁰ 1.445. Purified by fractn distn [Mandel and Hill *J Am Chem Soc* 76 3981 1954].

2,5-Di-*tert*-butylhydroquinone [88-58-4] M 222.3, m 222-223°. Crystd from *C₆H₆ or AcOH.

2,4-Di-*tert*-butyl-4-isopropylphenol [5427-03-2] M 248.4, m 39-41°, pK_{Est} ~12.3. Crystd from *n*-hexane or aq EtOH.

2,6-Di-*tert*-butyl-4-methylpyridine [38222-83-2] M 205.4, m 31-32°, 33-36°, b 148-153°/95mm, 223°/760mm, n₄²⁰ 1.476, pK_{Est} ~5.7. Possible impurity is 2,6-di-*tert*-butyl-4-neopentylpyridine. Attempts to remove coloured impurities directly by distn, acid-base extraction or treatment with activated charcoal were unsuccessful. Pure material can be obtained by dissolving 0.3mole of the alkylpyridine in pentane (150mL) and introducing it at the top of a water jacketed chromatographic column (40 x 4.5cm) the cooling is necessary because the base in pentane reacts exothermically with alumina) containing activated and acidic alumina (300g). The column is eluted with pentane using a 1L constant pressure funnel fitted at the top of the column to provide slight press. All the pyridine is obtained in the first two litres of eluent (the progress of elution is monitored by spotting a fluorescent TLC plate and examining under short wave UV light - a dark blue spot is evidence for the presence of the alkylpyridine). Elution is complete in 1h. Pentane is removed on a rotovap with 90-93% recovery yielding a liquid which solidifies on cooling, m 31-32°, and the base can be distilled. The *HPtCl₆* salt has m 213-314° (dec), and the *CF₃SO₃H* salt has m 202.5-203.5° (from CH₂Cl₂). [*Org Synth* 60 34 1981.]

Di-*tert*-butyl peroxide (*tert*-butyl peroxide) [110-05-4] M 146.2, d 0.794, n 1.389. Washed with aqueous AgNO₃ to remove olefinic impurities, water and dried (MgSO₄). Freed from *tert*-butyl hydroperoxide by passage through an alumina column [Jackson et al. *J Am Chem Soc* 107 208 1985], and if necessary two high vacuum distns from room temp to a liquid-air trap [Offenbach and Tobolsky *J Am Chem Soc* 79 278 1957]. *The necessary protection from EXPLOSION should be used.*

2,6-Di-*tert*-butylphenol [128-39-2] M 206.3, m 37-38°, pK²⁵ 11.70. Crystd from aqueous EtOH or *n*-hexane.

Dibutyl phthalate [84-74-2] M 278.4, b 206°/20mm, 340°/760mm, d 1.4929, d⁵ 1.0426, n²⁵ 1.490. Washed with dilute NaOH (to remove any butyl hydrogen phthalate), aqueous NaHCO₃ (charcoal), then distd water. Dried with CaCl₂, distd under vacuum, and stored in a desiccator over P₂O₅. (See also p. 151.)

2,6-Di-*tert*-butylpyridine [585-48-8] M 191.3, b 100-101°/23mm, d 0.852, n 1.474, pK²⁵ 5.02. Redistd from KOH pellets.

§ Polystyrene supported version is commercially available.

Di-*n*-butyl sulfide [544-40-1] M 146.3, α-form b 182°, β-form 190-230°(dec). Washed with aq 5% NaOH, then water. Dried with CaCl₂ and distd from sodium.

Di-*n*-butyl sulfone [598-04-9] M 162.3, m 43.5°. Purified by zone melting.

***N,N'*-Di-*tert*-butylthiourea** [4041-95-6] M 188.3, m 174-175° (evac capillary). Recrystd from H₂O [Bortnick et al. *J Am Chem Soc* 78 4358 1956].

3,5-Dicarbethoxy-1,4-dihydrocollidine [632-93-9] M 267.3, m 131-132°. Crystd from hot EtOH/water.

Dichloramine-T (*N,N*-dichloro-*p*-toluenesulfonamide) [473-34-7] **M 240.1, m 83°**. Crystd from pet ether (b 60-80°) or CHCl₃/pet ether. Dried in air. (see also chloramine-T in Chapter 5).

Dichloroacetic acid [79-43-6] **M 128.9, m 13.5°, b 95.0-95.5°/17-18mm, d 1.563, n 1.466, pK²⁵ 1.35**. Crystd from *benzene or pet ether. Dried with MgSO₄ and fractionally distd. [Bernasconi et al. *J Am Chem Soc* **107** 3612 1985.]

sym-Dichloroacetone (1,3-dichloropropan-2-one) [534-07-6] **M 127.0, m 41-43°, 45°, b 86-88°/12mm, 75-77°/22mm, 172-172.5°/atm, 170-175° /atm, d 1.383**. Crystd from CCl₄, CHCl₃ and *benzene. Distd under vacuum. [Conant and Quayle *Org Synth Coll Vol* 211 1941; Hall and Sirel *J Am Chem Soc* **74** 836 1952]. It is dimorphic [Daasch and Kagarise *J Am Chem Soc* **77** 6156 1955]. The *oxime* has **m 130-131°, b 106°/25mm** [*Arzneimittel-Forsch* **8** 638 1958].

Dichloroacetonitrile [3018-12-0] **M 110.0, b 110-112°, d 1.369, n 1.440**. Purified by distn and by gas chromatography. **FLAMMABLE**.

2,4-Dichloroaniline [554-00-7] **M 162.0, m 63°, pK²⁵ 2.02**. Crystd from EtOH/water. Also crystd from EtOH and dried *in vacuo* for 6h at 40° [Moore et al. *J Am Chem Soc* **108** 2257 1986; Edidin et al. *J Am Chem Soc* **109** 3945 1987].

3,4-Dichloroaniline [95-76-1] **M 162.0, m 71.5°, pK²⁵ 2.97**. Crystd from MeOH.

9,10-Dichloroanthracene [605-49-1] **M 247.1, m 214-215°**. Purified by crystn from MeOH or EtOH, followed by sublimation under reduced pressure. [Masnori and Kochi *J Am Chem Soc* **107** 7880 1985.]

2,4-Dichlorobenzaldehyde [874-42-0] **M 175.0, m 72°**. Crystd from EtOH or ligroin.

2,6-Dichlorobenzaldehyde [83-38-5] **M 175.0, m 70.5-71.5°**. Crystd from EtOH/water or pet ether (b 30-60°).

***o*-Dichlorobenzene** [95-50-1] **M 147.0, b 81-82°/31-32mm, 180.5°/760mm, d 1.306, n 1.551, n²⁵ 1.549**. Contaminants may include the *p*-isomer and trichlorobenzene [Suslick et al. *J Am Chem Soc* **106** 4522 1984]. It was shaken with conc or fuming H₂SO₄, washed with water, dried with CaCl₂, and distd from CaH₂ or sodium in a glass-packed column. Low conductivity material (*ca* 10⁻¹⁰ mhos) has been obtained by refluxing with P₂O₅, fractionally distilled and passed through a column packed with silica gel or activated alumina: it was stored in a dry-box under N₂ or with activated alumina.

***m*-Dichlorobenzene** [541-73-1] **M 147.0, b 173.0°, d 1.289, n 1.54586, n²⁵ 1.54337**. Washed with aqueous 10% NaOH, then with water until neutral, dried and distd. Conductivity material (*ca* 10⁻¹⁰ mhos) has been prepared by refluxing over P₂O₅ for 8h, then fractionally distilling, and storing with activated alumina. *m*-Dichlorobenzene dissolves rubber stoppers.

***p*-Dichlorobenzene** [106-46-7] **M 147.0, m 53.0°, b 174.1°, d 1.241, n⁶⁰ 1.52849**. *o*-Dichlorobenzene is a common impurity. Has been purified by steam distn, crystn from EtOH or boiling MeOH, air-dried and dried in the dark under vacuum. Also purified by zone refining.

2,2'-Dichlorobenzidine [84-68-4] **M 253.1, m 165°, pK_{Est(1)} ~3.0, pK_{Est(2)} ~4.0**. Crystd from EtOH.

3,3'-Dichlorobenzidine [91-94-1] **M 253.1, m 132-133°, pK_{Est(1)} ~4.8, pK_{Est(2)} ~5.7**. Crystd from EtOH or *benzene. **CARCINOGEN**.

2,4-Dichlorobenzoic acid [50-84-0] **M 191.0, m 163-164°, pK²⁵ 2.68**. Crystd from aqueous EtOH (charcoal), then *benzene (charcoal). It can also be recrystd from water.

2,5-Dichlorobenzoic acid [50-79-3] M 191.0, m 154°, b 301°/760mm, pK²⁵ 2.47. Crystd from water.

2,6-Dichlorobenzoic acid [50-30-6] M 191.0, m 141-142°, pK²⁵ 1.59. Crystd from EtOH and sublimed *in vacuo*.

3,4-Dichlorobenzoic acid [51-44-5] M 191.0, m 206-207°, pK²⁵ 3.64. Crystd from aqueous EtOH (charcoal) or acetic acid.

3,5-Dichlorobenzoic acid [51-36-5] M 191.0, m 188°, pK²⁵ 3.54. Crystd from EtOH and sublimed in a vacuum.

2,6-Dichlorobenzonitrile [1194-65-6] M 172.0, m 145°. Crystd from acetone.

4,4'-Dichlorobenzophenone [90-98-2] M 251.1, m 145-146°. Recrystd from EtOH [Wagner et al. *J Am Chem Soc* 108 7727 1986].

2,5-Dichloro-1,4-benzoquinone [615-93-0] M 177.0, m 161-162°. Recrystd twice from 95% EtOH as yellow needles [Beck et al. *J Am Chem Soc* 108 4018 1986].

2,6-Dichloro-1,4-benzoquinone [697-91-6] M 177.0, m 122-124°. Recrystd from pet ether (b 60-70°) [Carlson and Miller *J Am Chem Soc* 107 479 1985].

2,6-Dichlorobenzoyl chloride [4659-45-4] M 209.5, m 15-17, b 122-124°/15mm, d 1.464. Reflux for 2h with excess of acetyl chloride (3 vols), distil off AcCl followed by the benzoyl chloride. Store away from moisture. It is an **IRRITANT**.

3,4-Dichlorobenzyl alcohol [1805-32-9] M 177.0, m 38-39°. Crystd from water.

2,3-Dichloro-1,3-butadiene [1653-19-6] M 123.0, b 41-43°/85mm, 98°/760mm. Crystd from pentane to constant melting point about -40°. A mixture of *meso* and *d,l* forms was separated by gas chromatography on an 8m stainless steel column (8mm i.d.) with 20% DEGS (diethyleneglycolsilyl chloride) on Chromosorb W (60-80 mesh) at 60° and 80mL He/min. [Su and Ache *J Phys Chem* 80 659 1976.]

(+) and (-) **(8,8-Dichlorocamphorylsulfonyl)oxaziridine** [127184-05-8] M 298.2, m 178-180°, 183-186°, [α]_D²⁰ (+) and (-) 88.3° (c 1.3, CHCl₃), (+) and (-) 91° (c 5, CHCl₃). Recrystd from EtOH [Davis and Weismiller *J Org Chem* 55 3715 1990].

cis-3,4-Dichlorocyclobutene [2957-95-1] M 123.0, b 70-71°/55mm, 74-76°/55mm, d₄²⁰ 1.297, n_D²⁰ 1.499. Distd at 55mm through a 36-in platinum spinning band column, a fore-run b 58-62°/55mm is mainly 1,4-dichlorobutadiene. When the temperature reaches 70° the reflux ratio is reduced to 10:1 and the product is collected quickly. It is usually necessary to apply heat frequently with a sun lamp to prevent any dichlorobutadiene from clogging the exit in the early part of the distn [Pettit and Henery *Org Synth* 50 36 1970].

2,3-Dichloro-5,6-dicyano-p-benzoquinone (DDQ) [84-58-2] M 227.0, m 203° (dec). Crystd from CHCl₃, CHCl₃/*benzene (4:1), or *benzene and stored at 0°. [Pataki and Harvey *J Org Chem* 52 2226 1987.]

β,β' -Dichlorodiethyl ether [111-44-4] M 143.0, b 79-80°/20mm, 176-177.0°/743mm, n 1.457, d 1.219. See bis-(β -dichloroethyl)ether on p. 134.

1,2-Dichloro-1,2-difluoroethane [431-06-1] M 134.9, b 59°, n 1.376. Purified by fract dist [Hazeldine *J Chem Soc* 4258 1952]. For purification of diastereoisomeric mixture, with resolution into *meso* and *rac* forms, see Machulla and Stocklin [*J Phys Chem* 78 658 1974].

Dichlorodifluoromethane (Freon 12) [75-71-8] M 120.9, m -158°, b -29.8°/atm, 42.5°/10atm. Passage through saturated aqueous KOH then conc H₂SO₄, and a tower packed with activated copper on Kiesselguhr at 200° removed CO₂ and O₂. A trap cooled to -29° removed a trace of high boiling material. It is a non-flammable propellant.

1,3-Dichloro-5,5'-dimethylhydantoin [118-52-5] M 197.0, m 132-134°, 136°. Purified by dissolving in conc H₂SO₄ and diluting with ice H₂O, dry and recrystd from CHCl₃. It sublimes at 100° in a vacuum. Exhibits time dependent hydrolysis at pH 9. [Pettersen and Grzeskowiak *J Org Chem* **24** 1414 1959.]

4,5-Dichloro-3H-1,2-dithiol-3-one [1192-52-5] M 187.1, m 52-56°, 61°, b 87°/0.5mm, 125°/11mm. Distd *in vacuo* and then recrystd from pet ether. IR: ν 1650 cm⁻¹ [Boberg *Justus Liebigs Ann Chem* **693** 212 1966].

1,1-Dichloroethane (ethylidene dichloride) [75-34-3] M 99.0, b 57.3°, d¹⁵ 1.18350, d 1.177, n¹⁵ 1.41975. Shaken with conc H₂SO₄ or aqueous KMnO₄, then washed with water, saturated aqueous NaHCO₃, again with water, dried with K₂CO₃ and distd from CaH₂ or CaSO₄. Stored over silica gel.

1,2-Dichloroethane [107-06-2] M 99.0, b 83.4°, d 1.256, n¹⁵ 1.44759. Usually prepared by chlorinating ethylene, so that likely impurities include higher chloro derivatives and other chloro compounds depending on the impurities originally present in the ethylene. It forms azeotropes with water, MeOH, EtOH, trichloroethylene, CCl₄ and isopropanol. Its azeotrope with water (containing 8.9% water, and b 77°) can be used to remove gross amounts of water prior to final drying. As a preliminary purification step, it can be steam distd, and the lower layer was treated as below.

Shaken with conc H₂SO₄ (to remove alcohol added as an oxidation inhibitor), washed with water, then dilute KOH or aqueous Na₂CO₃ and again with water. After an initial drying with CaCl₂, MgSO₄ or by distn, it is refluxed with P₂O₅, CaSO₄ or CaH₂ and fractionally distd. Carbonyl-containing impurities can be removed as described for chloroform.

1,2-Dichloroethylene [cis + trans 540-59-0] M 96.9, b 60° (cis), d 1.284, b 48° (trans), d 1.257. Shaken successively with conc H₂SO₄, water, aqueous NaHCO₃ and water. Dried with MgSO₄ and distn separated the *cis*- and *trans*-isomers.

***cis*-1,2-Dichloroethylene [156-59-2] M 96.9, b 60.4°, d 1.2830, n¹⁵ 1.44903, n 1.4495.** Purified by careful fractional distn, followed by passage through neutral activated alumina. Also by shaking with mercury, drying with K₂CO₃ and distn. from CaSO₄.

***trans*-1,2-Dichloroethylene [156-60-5] M 96.9, b 47.7°, n¹⁵ 1.45189, n 1.4462, d 1.2551.** Dried with MgSO₄, and fractionally distd under CO₂. Fractional crystn at low temperatures has also been used.

5,7-Dichloro-8-hydroxyquinoline [773-76-2] M 214.1, m 180-181°, pK₁ 1.89, pK₂ 7.62. Crystd from acetone/EtOH.

2,3-Dichloromaleic anhydride [1122-17-4] M 167.0, m 105-115°, 120°, 121-121.5°. Purified by sublimation *in vacuo* [Katakis et al. *J Chem Soc, Dalton Trans* 1491 1986]. It has also been purified by Soxhlet extraction with hexane, recrystd from CHCl₃ and sublimed [MS, Relles *J Org Chem* **37** 3630 1972].

Dichloromethane (methylene dichloride) [75-09-2] M 84.9, b 40.0°, d 1.325, n 1.42456, n²⁵ 1.4201. Shaken with portions of conc H₂SO₄ until the acid layer remained colourless, then washed with water, aqueous 5% Na₂CO₃, NaHCO₃ or NaOH, then water again. Pre-dried with CaCl₂, and distd from CaSO₄, CaH₂ or P₂O₅. Stored away from bright light in a brown bottle with Linde type 4A molecular sieves, in an atmosphere of dry N₂. Other purification steps include washing with aq Na₂S₂O₃, passage through a column of silica gel, and removal of carbonyl-containing impurities as described under **Chloroform**. It has also been purified by treatment with basic alumina, distd, and stored over molecular sieves under nitrogen [Puchot et al. *J Am Chem Soc* **108** 2353 1986].

Dichloromethane from Japanese sources contained MeOH as stabiliser which is not removed by distn. It can, however, be removed by standing over activated 3A Molecular Sieves (note that 4A Sieves cause the development of pressure in bottles), passed through activated Al₂O₃ and distd [Gao et al. *J Am Chem Soc* **109** 5771 1987]. It has been fractionated through a platinum spinning band column, degassed, and distd onto degassed molecular sieves, Linde 4A, heated under high vacuum at over 450° until the pressure readings reached the low values of 10⁻⁶ mm — ~1-2h [Mohammad and Kosower *J Am Chem Soc* **93** 2713 1971].

Rapid purification: Reflux over CaH₂ (5% w/v) and distil. Store over 4A molecular sieves.

3,9-Dichloro-7-methoxyacridine [86-38-4] M 278.1, m 160-161°. Crystd from *benzene.

5,7-Dichloro-2-methyl-8-hydroxyquinoline (5,7-dichloro-8-hydroxyquinaldine) [72-80-0] M 228.1, m 114-115°, pK_{Est(1)} ~2.0, pK_{Est(2)} ~8.4. Crystd from EtOH.

2,4-Dichloro-6-methylphenol [1570-65-6] M 177.0, m 55°, b 129-132°/40mm, pK²⁰ 8.14. Crystd from water.

2,4-Dichloro-1-naphthol [2050-76-2] M 213.1, m 106-107°, pK_{Est} ~7.7. Crystd from MeOH.

2,3-Dichloro-1,4-naphthoquinone [117-80-6] M 227.1, m 193°. Crystd from EtOH.

2,5-Dichloro-4-nitroaniline [6627-34-5] M 207.0, m 157-158°, pK²⁵ -1.74 (aq H₂SO₄). Crystd from EtOH, then sublimed.

2,6-Dichloro-4-nitroaniline [99-30-9] M 207.0, m 193°. Crystd from aq EtOH or *benzene/EtOH.

2,5-Dichloro-1-nitrobenzene [89-61-2] M 192.0, m 56°. Crystd from absolute EtOH.

3,4-Dichloro-1-nitrobenzene [99-54-7] M 192.0, m 43°. Crystd from absolute EtOH.

2,4-Dichloro-6-nitrophenol [609-89-2] M 208.0, m 122-123°, pK_{Est} ~5.0. Crystd from AcOH.

2,6-Dichloro-4-nitrophenol [618-00-4] M 208.0, m 125°, pK²⁵ 3.55. Crystd from EtOH and dried *in vacuo* over anhydrous MgSO₄.

4,6-Dichloro-5-nitropyrimidine [4316-93-2] M 194.0, m 100-103°, 101-102°, pK_{Est} <0. If too impure then dissolve in Et₂O, wash with H₂O, dry over MgSO₄, evaporate to dryness and recrystallise from pet ether (b 85-105°) as a light tan solid. It is sol in *ca* 8 parts of MeOH [Boon et al, *J Chem Soc* 96 1951; Montgomery et al. in *Synthetic Procedures in Nucleic Acid Chemistry* (Zorbach and Tipson eds) Wiley & Sons, NY, p76 1968].

Dichlorophen [2,2'-methylenebis(4-chlorophenol)] [97-23-4] M 269.1, b 177-178°, pK_{Est} ~9.7. Crystd from toluene.

2,3-Dichlorophenol [576-24-9] M 163.0, m 57°, pK²⁵ 7.70. Crystd from ether.

2,4-Dichlorophenol [120-83-2] M 163.0, m 42-43°, pK²⁵ 7.89. Crystd from pet ether (b 30-40°). Purified by repeated zone melting, using a P₂O₅ guard tube to exclude moisture. Very *hygroscopic* when dry.

2,5-Dichlorophenol [583-78-8] M 163.0, m 58°, b 211°/744mm, pK²⁵ 7.51. Crystd from ligroin and sublimed.

3,4-Dichlorophenol [95-77-2] M 163.0, m 68°, b 253.5°/767mm, pK²⁵ 8.58. Crystd from pet ether/*benzene mixture.

3,5-Dichlorophenol [591-35-5] M 163.0, m 68°, b 122-124°/8mm, 233-234°/760mm, pK²⁵ 8.81. Crystd from pet ether/*benzene mixture.

2,4-Dichlorophenoxyacetic acid (2,4-D) [94-75-7] M 221.0, m 146°, pK²⁵ 2.90. Crystd from MeOH. TOXIC.

α-(2,4-Dichlorophenoxy)propionic acid (2,4-DP, Dichloroprop) [120-36-5] M 235.1, m 117°, pK²⁰ 2.86, Crystd from MeOH. TOXIC.

2,4-Dichlorophenylacetic acid [19719-28-9] M 205.0, m 131°, 132-133°, pK_{Est} ~4.0. Crystd from aqueous EtOH.

2,6-Dichlorophenylacetic acid [6575-24-2] M 205.0, m 157-158°, pK_{Est} ~3.8. Crystd from aqueous EtOH.

3-(3,4-Dichlorophenyl)-1,1-dimethyl urea (Diuron) [330-54-1] M 233.1. Crystd four times from 95% EtOH [Beck et al. *J Am Chem Soc* 108 4018 1986].

4,5-Dichloro-*o*-phenylenediamine [5348-42-5] M 177.1, m 162°, 162-163°, pK_{Est(1)} ~-1.0, pK_{Est(2)} ~2.9. Recrystd from hexane, *C₆H₆, pet ether or H₂O (Na₂SO₄) and sublimed at 150°/15mm.

4,5-Dichlorophthalic acid [56962-08-4] M 235.0, m 200° (dec to anhydride), pK_{Est(1)} 2.2, pK_{Est(2)} ~4.7. Crystd from water. Can be purified by converting to the anhydride, reacting with boiling EtOH to form the *monoethyl ester* (m 133-134°) and hydrolysing back to the diacid

3,6-Dichlorophthalic anhydride [4466-59-5] M 189-191°, 191-191.5°, b 339°. Boil in xylene (allowing any vapours which would contain H₂O to be removed, e.g. Dean and Stark trap), which causes the acid to dehydrate to the anhydride and cool. Recryst from xylene [Villiger *Chem Ber* 42 3539 1909; Fedorow *Izv Akad Nauk SSSR Otd Khim Nauk* 397 1948, *Chem Abstr* 1585 1948].

1,2-Dichloropropane [78-87-5] M 113°, b 95.9-96.2°, d 1.158, n 1.439. Distd from CaH₂.

2,2-Dichloropropane [594-20-7] M 113.0, b 69.3°, d 1.090, n 1.415. Washed with aqueous Na₂CO₃ soln, then distilled water, dried over CaCl₂ and fractionally distd.

2,6-Dichloropurine [5451-40-1] M 189.0, m 180-181.5°, 181°, 185-195°(dec), 188-189°, pK₁²⁰ 1.16 (aq H₂SO₄), pK₂²⁰ 7.06. It can be recrystd from 150 parts of boiling H₂O and dried at 100° to constant weight. Soluble in EtOAc. The HgCl₂ salt separates from EtOH soln. UV: λ_{max} 275nm (ε 8.9K) at pH 1; and 280nm (ε 8.5K) at pH 11 [Elion and Hitchings *J Am Chem Soc* 78 3508 1956; Schaeffer and Thomas *J Am Chem Soc* 80 3738 1958; Beaman and Robins *J Appl Chem (London)* 12 432 1962; Montgomery *J Am Chem Soc* 78 1928 1956].

2,6-Dichloropyridine [2402-78-0] M 148.0, m 87-88°, pK -2.86 (aq H₂SO₄). Crystd from EtOH.

3,5-Dichloropyridine [2457-47-8] M 148.0, m 64-65°, pK²⁵ 0.67. Crystd from EtOH.

4,7-Dichloroquinoline [86-98-6] M 198.1, m 86.4-87.4°, b 148°/10mm, pK²⁵ 2.80. Crystd from MeOH or 95% EtOH.

2,3-Dichloroquinoxaline [2213-63-0] M 199.0, m 152-153°, 152-154°, pK_{Est} <0. Recrystd from *C₆H₆ and dried in a vacuum [Cheeseman *J Chem Soc* 1804 1955].

2,6-Dichlorostyrene [28469-92-3] M 173.0, b 72-73°/2mm, d 1.4045, n 1.5798. Purified by fractional crystn from the melt and by distn.

2,4-Dichlorotoluene [95-73-8] M 161.1, m -13.5°, b 61-62°/3mm, d 1.250, n 1.5513. Recrystd from EtOH at low temperature or fractionally distd.

2,6-Dichlorotoluene [118-69-4] M 161.1, b 199-200°/760mm, d 1.254, n 1.548. Fractionally distd and collecting the middle fraction.

3,4-Dichlorotoluene [95-75-0] M 161.1, m -16°, b 205°/760mm, d 1.2541, n 1.549. Recrystd from EtOH at very temperature or fractionally distd.

α,α' -Dichloro-*p*-xylene [623-25-6] M 175.1, m 100°. Crystd from *benzene and dried under vacuum.

Dicinnamalacetone (1,9-diphenyl-1,3,6,8-nonatetraen-5-one) [622-21-9] M 314.4, m 146°. Crystd from *benzene/isooctane (1:1).

Dicumyl peroxide [80-43-3] M 270.4, m 39-40°. Crystd from 95% EtOH (charcoal). Stored at 0°. **Potentially EXPLOSIVE.**

9,10-Dicyanoanthracene [1217-45-4] M 228.3, m 340°. Recrystd twice from pyridine [Mattes and Farid *J Am Chem Soc* 108 7356 1986].

1,2-Dicyanobenzene [91-15-6] M 128.1, m 141°. (See phthalonitrile on p. 334.)

1,4-Dicyanobenzene [623-26-7] M 128.1, m 222°. Crystd from EtOH.

1,4-Dicyanonaphthalene [3029-30-9] M 178.2, m 206°. Purified by crystn and sublimed *in vacuo*.

1,3-Dicyclohexylcarbodiimide (DCC) [538-75-0] M 206.3, m 34-35°, b 95-97°/0.2mm, 120-121°/0.6mm, 155°/11mm. It is sampled as a liquid after melting in warm H₂O. It is sensitive to air and *it is a potent skin irritant*. It can be distd in a vacuum and stored in a tightly stoppered flask in a freezer. It is very soluble in CH₂Cl₂ and pyridine where the reaction product with H₂O, after condensation, is dicyclohexyl urea which is insoluble and can be removed by filtration. Alternatively dissolve in CH₂Cl₂ add powdered anhyd MgSO₄ shake 4h, filter, evaporate and distil at 0.6 mm press and oil bath temperature 145°. [*Biochem Prep* 10, 122 1963; *Justus Liebigs Ann Chem* 571 83 1951; *Justus Liebigs Ann Chem* 612 11 1958.]

***cis*-Dicyclohexyl-18-crown-6** [16069-36-6] M 372.5, m 47-50°. Purified by chromatography on neutral alumina and eluting with an ether/hexane mixture [see *Inorg Chem* 14 3132 1975]. Dissolved in ether at *ca* 40°, and spectroscopic grade MeCN was added to the soln which was then chilled. The crown ether pptd and was filtered off. It was dried *in vacuo* at room temperature [Wallace *J Phys Chem* 89 1357 1985]. **SKIN IRRITANT.**

Di-*n*-decylamine [1120-49-6] M 297.6, m 34°. b 153°/1mm, 359°/760mm, pK_{Est} ~11.0. Dissolved in *benzene and pptd as its bisulfate by shaking with 4M H₂SO₄. Filtered, washed with *benzene, separating by centrifugation, then the free base was liberated by treating with aqueous NaOH [McDowell and Allen *J Phys Chem* 65 1358 1961].

Didodecylamine [3007-31-6] M 353.7, m 51.8°, pK²⁵ 11.00. Crystd from EtOH/*C₆H₆ under N₂.

Didodecyltrimethylammonium bromide [3282-73-3] M 463.6, m 157-162°. Recrystd from acetone, acetone/ether mixture, then from ethyl acetate, washed with ether and dried in a vacuum oven at 60° [Chen et al. *J Phys Chem* 88 1631 1984; Rupert et al. *J Am Chem Soc* 107 2628 1985; Halpern et al. *J Am Chem Soc* 108 3920 1986; Allen et al. *J Phys Chem* 91 2320 1987].

Dienestrol [4,4'-(diethylidene-ethylene)diphenol, Dienol] [84-17-3] M 266.3, m 227-228°, 231-233°, $pK_{\text{Est}} \sim 9.8$. Crystd from EtOH or dilute EtOH, sublimes at 130°/1mm. The *diacetate* has m 119-120° (from EtOH) [Hobday and Short *J Chem Soc* 609 1943].

Diethanolamine (2,2'-iminodiethanol) [111-42-2] M 105.1, m 28°, b 154-155°/10mm, 270°/760mm $pK^{25} 8.88$. Fractionally distd twice, then fractionally crystd from its melt.

3,4-Diethoxy-3-cyclobutene-1,2-dione (diethyl squarate) [5321-87-8] M 170.2, b 89-91°/0.4mm, 88-92°/0.4mm, $d_4^{20} 1.162$, $n_D^{25} 1.5000$. Dissolve in Et₂O, wash with Na₂CO₃, H₂O and dry (Na₂SO₄), filter, evaporate and distil using a Kügelrohr or purify by chromatography. Use a Kieselgel column and elute with 20% Et₂O-Pet ether (b 40-60°) then with Et₂O-pet ether (1:1), evaporate and distil *in vacuo*. [Dehmlow and Schell *Chem Ber* 113 1 1980; Perri and Moore *J Am Chem Soc* 112 1897 1990; IR: Cohen and Cohen *J Am Chem Soc* 88 1533 1966.] **It can cause severe dermatitis** [Foland et al. *J Am Chem Soc* 111 975 1989].

N,N-Diethylacetamide [685-91-6] M 157.2, b 86-88°, n 1.474, d 0.994. Dissolved in cyclohexane, shaken with anhydrous BaO and then filtered. The procedure was repeated three times, and the cyclohexane was distd off at 1 atmosphere pressure. The crude amide was also fractionally distd three times from anhydrous BaO.

Diethyl acetamidomalonate [1068-90-2] M 217.2, m 96°. Crystd from *benzene/pet ether.

Diethyl acetylenedicarboxylate [762-21-0] M 170.2, b 60-62°/0.3mm, 107-110°/11mm, 118-120°/20mm, $d_4^{20} 1.0735$, $n_D^{20} 1.4428$. Dissolve in *C₆H₆, wash with NaHCO₃, H₂O, dry over Na₂SO₄, filter, evaporate and distil in a vacuum [IR: Walton and Hughes *J Am Chem Soc* 79 3985 1957; Truce and Kruse *J Am Chem Soc* 81 5372 1959].

Diethylamine [109-89-7] M 73.1, b 55.5°, d 0.707, n 1.38637, $pK^{15} 11.38$. Dried with LiAlH₄ or KOH pellets. Refluxed with, and distd from, BaO or KOH. Converted to the *p*-toluenesulfonamide and crystd to constant melting point from dry pet ether (b 90-120°), then hydrolysed with HCl, excess NaOH was added, and the amine passed through a tower of activated alumina, redistd and dried with activated alumina before use [Swift *J Am Chem Soc* 64 115 1942].

§ A polystyrene diethylaminomethyl supported version is commercially available.

Diethylamine hydrochloride [660-68-4] M 109.6, m 223.5°. Crystd from absolute EtOH. Also crystd from dichloroethane/MeOH. *Hygroscopic*.

trans-4-(Diethylamino)azobenzene [3588-91-8] M 320.5, m 171° $pK_{\text{Est}(1)} \sim 5.4$, $pK_{\text{Est}(2)} \sim 3.0$. Purified by column chromatography [Flamigni and Monti *J Phys Chem* 89 3702 1985].

N,N-Diethylaniline [91-66-7] M 149.2, b 216.5°, d 0.938, n 1.5409 $pK^{25} 6.57$. Refluxed for 4h with half its weight of acetic anhydride, then fractionally distd under reduced pressure (b 92°/10mm).

Diethyl azodicarboxylate (DEAD) [1972-28-7] M 174.2, b 104.5°/12mm, 211-213°/atm, $d_4^{20} 1.110$, $n_D^{20} 1.420$. Dissolve in toluene, wash with 10% NaHCO₃ till neutral (may require several washes if too much hydrolysis had occurred (check IR for OH bands), then wash with H₂O (2 x), dry over Na₂SO₄, filter, evaporate the toluene and distil through a short Vigreux column. Main portion boils at 107-111°/15mm [*Org Synth Coll Vol III* 376 1955].

§ A polystyrene supported DEAD version is commercially available.

5,5-Diethylbarbituric acid (Barbital) [57-44-3] M 184.2, m 188-192°, $pK_1^{25} 8.02$, $pK_2^{25} 12.7$. Crystd from water or EtOH. Dried in a vacuum over P₂O₅.

Diethyl bromomalonate [685-87-0] M 239.1, b 116-118°/10mm, 122-123°/20mm, d_4^{20} 1.420, n_D^{20} 1.4507. Purified by fractional distn in a vacuum. IR: 1800 and 1700 cm^{-1} [Abramovitch *Can J Chem* 37 1146 1959; Bretschneider and Karpitschka *Monatsh Chem* 84 1091 1953].

Diethyl tert-butylmalonate [759-24-0] M 216.3, b 40-42°/0.03, 102-104°/11mm, 109.5-110.5°/17mm, 205-210°/760mm, d_4^{20} 0.980, n_D^{20} 1.425. Dissolve in Et_2O , wash with aqueous NaHCO_3 , H_2O , dry (MgSO_4), filter, evaporate and distil residue. Identified by hydrolysis to the acid and determining the neutralisation equiv (theor: 80.0). The acid has m 155-157° efferv [Hauser, Abramovitch and Adams *J Am Chem Soc* 64 2715 1942; Bush and Beauchamp *J Am Chem Soc* 75 2949 1953].

***N,N'*-Diethylcarbanilide (sym-Diethyldiphenylurea)** [85-98-3] M 268.4, m 79°. Crystd from EtOH.

Diethyl carbonate [105-58-8] M 118.1, b 126.8°, d 0.975, n_D^{25} 1.38287. It was washed (100mL) with an aqueous 10% Na_2CO_3 (20mL) solution, saturated CaCl_2 (20mL), then water (30mL). After drying by standing over solid CaCl_2 for 1h (note that prolonged contact should be avoided because slow combination with CaCl_2 occurs), it should be fractionally distd. Also dried over MgSO_4 and distd.

1,1'-Diethyl-2,2'-cyanine iodide [977-96-8] M 454.4, m 274°(dec). Crystd from EtOH and dried in a vacuum oven at 80° for 4h.

***N,N*-Diethylcyclohexylamine** [91-65-6] M 155.3, b 193°/760mm, d 0.850, n 1.4562, pK^{25} 10.72. Dried with BaO and fractionally distd.

Diethylene glycol [111-46-6] M 106.1, f -10.5°, b 244.3°, d 1.118, n_D^{15} 1.4490, n 1.4475. Fractionally distd under reduced pressure (b 133°/14mm), then fractionally crystd by partial freezing.

Diethylene glycol diethyl ether [112-36-7] M 162.2, b 85-86°/10mm, 188.2-188.3°/751mm, d 0.909, n 1.412. Dried with MgSO_4 , then CaH_2 or LiAlH_4 , under N_2 . If sodium is used the ether should be redistd alone to remove any products which may be formed by the action of sodium on the ether. As a preliminary purification, the crude ether (2L) can be refluxed for 12h with 25mL of conc HCl in 200mL of water, under reduced pressure, with slow passage of N_2 to remove aldehydes and other volatile substances. After cooling, addn of sufficient solid KOH pellets (slowly and with shaking until no more dissolves) gives two liquid phases. The upper of these is decanted, dried with fresh KOH pellets, decanted, then refluxed over, and distd from sodium. Can be passed through (alkaline) alumina prior to purification.

Diethylene glycol ditosylate [7460-82-4] M 414.5, m 86-87°, 87-88°, 88-89°. Purified by recrystn from Me_2CO and dried in a vacuum.

Diethylene glycol mono-*n*-butyl ether (butyl carbitol) [112-34-5] M 162.2, b 69-70°/0.3mm, 230.5°/760mm, d 0.967, n 1.4286. Dried with anhydrous K_2CO_3 or CaSO_4 , filtered and fractionally distd. Peroxides can be removed by refluxing with stannous chloride or a mixture of FeSO_4 and KHSO_4 (or, less completely, by filtration under slight pressure through a column of activated alumina).

Diethylene glycol monoethyl ether [111-90-0] M 134.2, b 201.9°, d 0.999, n 1.4273, n_D^{25} 1.4254. Ethylene glycol can be removed by extracting 250g in 750mL of *benzene with 5mL portions of water, allowing for phase separation, until successive aqueous portions show the same volume increase. Dried, and freed from peroxides, as described for diethylene glycol mono-*n*-butyl ether.

Diethylene glycol monomethyl ether [111-77-3] M 120.2, b 194°, d 1.010, n 1.423. Purified as for diethylene glycol mono-*n*-butyl ether.

Diethylenetriaminepenta-acetic acid (DTPA) [67-43-6] M 393.4, m 219-220°, pK_1^{25} 1.79, pK_2^{25} 2.56, pK_3^{25} 4.42, pK_4^{25} 8.76, pK_5^{25} 10.42. Crystd from water. Dried under vacuum or at 110°. [Bielski and Thomas *J Am Chem Soc* 109 7761 1987].

Diethyl ether (ethyl ether) [60-29-7] **M 74.1, b 34.6°/760mm, d 0.714, n¹⁵ 1.3555, n 1.35272.** Usual impurities are water, EtOH, diethyl peroxide (which is explosive when concentrated), and aldehydes. Peroxides [detected by liberation of iodine from weakly acid (HCl) solutions of KI, or by the blue colour in the ether layer when 1mg of Na₂Cr₂O₇ and 1 drop of dil H₂SO₄ in 1mL of water is shaken with 10mL of ether] can be removed in several different ways. The simplest method is to pass dry ether through a column of activated alumina (80g Al₂O₃/700mL of ether). More commonly, 1L of ether is shaken repeatedly with 5-10mL of a soln comprising 6.0g of ferrous sulfate and 6mL of conc H₂SO₄ in 110mL of water. Aqueous 10% Na₂SO₃ or stannous chloride can also be used. The ether is then washed with water, dried for 24h with CaCl₂, filtered and dried further by adding sodium wire until it remains bright. The ether is stored in a dark cool place, until distd from sodium before use. Peroxides can also be removed by wetting the ether with a little water, then adding excess LiAlH₄ or CaH₂ and leaving to stand for several hours. (This also dried the ether.)

Werner [*Analyst* 58 335 1933] removed peroxides and aldehydes by adding 8g AgNO₃ in 60mL of water to 1L of ether, then 100mL of 4% NaOH and shaking for 6min. Fierz-David [*Chimia* 1 246 1947] shook 1L of ether with 10g of a zinc-copper couple. (This reagent was prepared by suspending zinc dust in 50mL of hot water, adding 5mL of 2M HCl and decanting after 20sec, washing twice with water, covering with 50mL of water and 5mL of 5% cuprous sulfate with swirling. The liquid was decanted and discarded, and the residue was washed three times with 20mL of ethanol and twice with 20mL of diethyl ether).

Aldehydes can be removed from diethyl ether by distn from hydrazine hydrogen sulfate, phenyl hydrazine or thiosemicarbazide. Peroxides and oxidisable impurities have also been removed by shaking with strongly alkaline satd KMnO₄ (with which the ether was left to stand in contact for 24h), followed by washing with water, conc H₂SO₄, water again, then drying (CaCl₂) and distn from sodium, or sodium containing benzophenone to form the ketyl. Other purification procedures include distn from sodium triphenylmethide or butyl magnesium bromide, and drying with solid NaOH or P₂O₅.

Rapid purification: Same as for 1,4-dioxane.

Diethyl ethoxymethylene malonate [87-13-8] **M 216.2, b 014°/0.2mm, 109°/0.5mm, 279-283°/atm, d₄²⁰ 1.079, n_D²⁰ 1.4623.** Likely impurity is diethyl diethoxymethylene malonate which is difficult to separate from diethyl ethoxymethylene malonate by distn and it is necessary to follow the course of the distn by the change in refractive index instead of boiling point. After a low boiling fraction is collected, there is obtained an intermediate fraction (n_D²⁰ 1.414—1.458) the size of which depends on the amount of diethoxymethylene compound. This fraction is fractionated through a 5-inch Vigreux column at low pressure avoiding interruption in heating. Fraction **b** 108-110°/0.25mm was *ca* 10° lower than the submitters' (**b** 97.2°/0.25mm (n_D²⁰ 1.4612—1.4623) [*Org Synth Coll Vol III* 395 1955; Fuson et al. *J Org Chem* 11 197 1946; Duff and Kendal *J Chem Soc* 893 1948].

N,N'-Diethylformamide [617-84-5] **M 101.2, b 29°/0.5mm, 61-63°/10mm, 178.3-178.5°/760mm, d₄²⁰ 0.906, n_D²⁵ 1.4313.** Distd under reduced pressure then at atmospheric pressure [Wintcler et al. *Helv Chim Acta* 37 2370 1954; NMR: Hoffmann *Z Anal Chem* 170 177 1959].

Diethyl fumarate [623-91-6] **M 172.2, b 218°, d 1.052, n 1.441.** Washed with aqueous 5% Na₂CO₃, then with saturated CaCl₂ soln, dried with CaCl₂ and distd.

Di-(2-ethylhexyl)phthalate ('di-iso-octyl' phthalate) [117-81-7] **M 390.6, b 384°, 256-257°/1mm, d 0.9803, n 1.4863.** Washed with Na₂CO₃ soln, then shaken with water. After the resulting emulsion had been broken by adding ether, the ethereal soln was washed twice with water, dried (CaCl₂), and evaporated. The residual liquid was distd several times under reduced pressure, then stored in a vacuum desiccator over P₂O₅ [French and Singer *J Chem Soc* 1424 1956]

Diethyl ketone (3-pentanone) [96-22-0] **M 86.1, b 102.1°, d 0.8099, n 1.392.** Dried with anhydrous CaSO₄ or CuSO₄, and distd from P₂O₅ under N₂ or under reduced pressure. Further purification by conversion to the semicarbazone (recrystd to constant **m** 139°, from EtOH) which, after drying under vacuum over CaCl₂ and paraffin wax, was refluxed for 30min with excess oxalic acid, then steam distd and salted out with K₂CO₃. Dried with Na₂SO₄ and distd [Cowan, Jeffrey and Vogel *J Chem Soc* 171 1940].

Diethyl malonate [105-53-3] M 160.2, b 92°/22mm, 198-199°/760mm, d 1.056, d_4^{25} 1.0507, n 1.413. If too impure (IR, NMR) the ester (250g) has been heated on a steam bath for 36h with absolute EtOH (125mL) and conc H₂SO₄ (75mL), then fractionally distd under reduced pressure. Otherwise fractionally distil under reduced pressure and collect the steady boiling middle fraction.

Diethyl phenyl orthoformate (diethoxy phenoxy ethane) [14444-77-0] M 196.3, b 111°/11mm, 122°/13mm, d_4^{20} 1.0099, n_D^{20} 1.4799. Fractionated through an efficient column under vacuum [Smith *Acta Chem Scand* 10 1006 1956].

Diethyl phthalate [84-66-2] M 222.2, b 172°/12mm, b 295°/760mm, d^{25} 1.1160, n 1.5022. Washed with aqueous Na₂CO₃, then distilled water, dried (CaCl₂), and distd under reduced pressure. Stored in a vacuum desiccator over P₂O₅.

Diethyl phthalimidomalonate [56680-61-5] M 305.3, m 72-74°, 73-74°, pK 9.17. Dissolve in xylene and when the temperature is 30° add pet ether (b 40-60°) and cool to 20° whereby the malonate separates as a pale brown powder [Booth et al. *J Chem Soc* 666 1944]. Alternatively, dissolve in *C₆H₆, dry over CaCl₂, filter, evaporate and the residual oil solidifies. This is ground with Et₂O, filter and wash with Et₂O until white in colour, and dry in a vacuum. The anion has λ_{max} 254nm (ϵ 18.5K) [Clark and Murray *Org Synth Coll Vol I* 271 1941; UV of Na salt: Nnadi and Wang *J Am Chem Soc* 92 4421 1970].

2,2-Diethyl-1,3-propanediol [115-76-4] M 132.2, m 61.4-61.8°. Crystd from pet ether (b 65-70°).

Diethyl pyrocarbonate (DEP) [1609-47-8] M 162.1, b 38-40°/12mm, 160-163°/atm, d_4^{20} 1.119, n_D^{20} 1.398. Dissolve in Et₂O, wash with dilute HCl, H₂O, dry over Na₂SO₄, filter, evaporate and distil the residue first *in vacuo* then at atmospheric pressure. It is soluble in alcohols, esters, ketones and hydrocarbon solvents. A 50% w/w soln is usually prepared for general use. **Treat with great CAUTION as DEP irritates the eyes, mucous membranes and skin.** [Boehm and Mehta *Chem Ber* 71 1797 1938; Thoma and Rinke *Justus Liebigs Ann Chem* 624 30 1959.]

Diethylstilboesterol [56-23-1] M 268.4, m 169-172°. Crystd from *benzene.

Diethyl succinate [123-25-1] M 174.2, b 105°/15mm, d 1.047, n 1.4199. Dried with MgSO₄, and distd at 15mm pressure.

Diethyl sulfate [64-67-5] M 154.2, b 96°/15mm, 118°/40mm, d 1.177, n 1.399. Washed with aqueous 3% Na₂CO₃ (to remove acidic material), then distilled water, dried (CaCl₂), filtered and distd. *Causes blisters to the skin.*

Diethyl disulfide [110-81-6] M 122.3, b 154-155°, d 0.993, n 1.506. Dried with silica gel or MgSO₄ and distd under reduced pressure (optionally from CaCl₂).

Diethyl sulfide [352-93-2] M 90.2, m 0°/15mm, 90.1°/760mm, d 0.837, n 1.443. Washed with aq 5% NaOH, then water, dried with CaCl₂ and distd from sodium. Can also be dried with MgSO₄ or silica gel. Alternative purification is *via* the Hg(II) chloride complex [(Et)₂S.2HgCl₂] (see dimethyl sulfide).

Diethyl (-)-D- (from the non-natural) [13811-71-7] and (+)-L- (from the natural acid) [89-91-2] tartrate M 206.2, m 17°, b 80°/0.5mm, 162°/19mm, 278-282°/atm, d_4^{20} 1.204, n_D^{20} 1.4476, $[\alpha]_D^{20}$ (-) and (+) 26.5° (c 1, H₂O) and (-) and (+) 8.5° (neat), $[\alpha]_{546}^{20}$ (-) and (+) 30° (c 1, H₂O). Distd under high vacuum and stored under vacuum or in an inert atm in a desiccator in round bottomed flasks equipped with a vac stopcock. Have also been dist by Kügelrohr distn and/or by 'wiped-film' molecular distn. Slightly sol in H₂O but miscible in EtOH and Et₂O. [Gao et al. *J Am Chem Soc* 109 5770 (5771) 1987; IR: Pristera *Anal Chem* 25 844 1953.]

Diethyl terephthalate [636-09-0] M 222.2, m 44°, 142°/2mm, 302°/760mm. Crystd from toluene and distd under reduced pressure.

sym-Diethylthiourea [105-55-5] M 132.2, m 76-77°. Crystd from *benzene.

Difluoroacetic acid [381-73-7] M 96.0, m -0.35°, b 67-70°/20mm, 134°/760mm, d_4^{20} 1.530, n_D^{20} 1.3428, pK^{25} 1.28. Purified by distilling over P_2O_5 . The *acid chloride* is a fuming liquid b 25°/atm, and the *amide* has b 108.6°/35mm, m 52° (from * C_6H_6), and the *anilide* has b 90°/1mm, 114°/5mm, m 58° [Henne and Pelley *J Am Chem Soc* 74 1426 1952, Coffman et al. *J Org Chem* 14 749 1949; NMR: Meyer et al. *J Am Chem Soc* 75 4567 1953; pK: Wegscheider *Z Phys Chem* 69 614 1909].

Diglycolic acid (2-oxapentane-1,5-dioic acid) [110-99-6] M 134.1, m 148° (monohydrate), pK_1^{25} 2.97, pK_2^{25} 4.37. Crystd from water.

Diglycyl glycine [556-33-2] M 189.2, m 246°(dec), pK_1^{25} 3.30, pK_2^{25} 7.96. Crystd from H_2O or $H_2O/EtOH$ and dried at 110°.

Diglyme [bis-(2-methoxyethyl) ether, diethylene glycol dimethyl ether] [111-96-6] M 134.2, b 62°/17mm, 75°/35mm, 160°/760mm, d 0.917, n 1.4087. Dried with NaOH pellets or CaH_2 , then refluxed with, and distd (under reduced pressure) from Na, CaH_2 , $LiAlH_4$, $NaBH_4$ or NaH. These operations were carried out under N_2 . The amine-like odour of diglyme has been removed by shaking with a weakly acidic ion-exchange resin (Amberlite IR-120) before drying and distn. Addn of 0.01% $NaBH_4$ to the distillate inhibits peroxidation. Purification as for dioxane. Also passed through a 12-in column of molecular sieves to remove water and peroxides.

Digoxin [20830-75-5] M 781.0, m 265°(dec), $[\alpha]_{546}^{20}$ +14.0° (c 10, pyridine). Crystd from aqueous EtOH or aqueous pyridine.

4,4'-Di-n-heptyloxyazoxybenzene [2635-26-9] M 426.6, m 75°, 95° (smectic → nematic) and 127° (nematic → liquid), pK_{Est} --5. Purified by chromatography on Al_2O_3 (*benzene), recrystd from hexane or 95% EtOH and dried by heating under vacuum. The liquid crystals can be sublimed *in vacuo*. [Mellifiori et al. *Spectrochim Acta Part A* 37(A) 605 1981; Dewar and Schroeder *J Am Chem Soc* 86 5235 1964; Weygand and Glaber *J Prakt Chem* 155 332 1940].

9,10-Dihydroanthracene [613-31-0] M 180.3, m 110-110.5°. Crystd from EtOH [Rabideau et al. *J Am Chem Soc* 108 8130 1986].

2,3-Dihydrobenzofuran (coumaran) [496-16-2] M 120.2, m -21.5°, 72-73°/12mm, 84°/17mm, 188°/atm, d_4^{20} 1.065, n_D^{20} 1.5524. Suspend in aqueous NaOH and steam distil. Saturate the distillate with NaCl and extract with Et_2O , dry extract ($MgSO_4$), filter, evap and distil the residue. It gives a strong violet colour with $FeCl_3 + H_2SO_4$ and forms a yellow *picrate*, m 76°, from EtOH or * C_6H_6 which loses coumaran in a desiccator [Bennett and Hafez *J Chem Soc* 287 1941; Baddeley et al. *J Chem Soc* 2455 1956].

Dihydrochloranil (tetrachloro-1,4-hydroquinone) [87-87-6] M 247.9, m 240.5°. Crystd from EtOH or AcOH+EtOH. Sublimes at 77°/0.6x10⁻³mm. The *dibenzoyl derivative* has m 233°. [Conant and Fieser *J Am Chem Soc* 45 2207 1923 ; Rabideau et al. *J Am Chem Soc* 108 8130 1986.]

Dihydrocodeine [125-28-0] M 301.4, m 112-113°, b 248/14mm°. Crystd from aqueous methanol.

1,4-Dihydro-1,4-epoxynaphthalene [573-57-9] M 144.2, m 53-54.5°, 53-56°, 55-56°. Dissolve in Et_2O , wash with H_2O , dry over K_2CO_3 , filter, evaporate and dry the residue at 15mm, then recrystallise from pet ether (b 40-60°), dry at 25°/0.005mm and sublime (sublimes slowly at room temp)[Wittig and Pohmer *Chem Ber* 89 1334 1956; Gilman and Gorsich *J Am Chem Soc* 79 2625 1957].

Dihdropyran (3,4-dihydro-2H-pyran) [110-87-2] M 84.1, b 84.4°/742mm, 85.4-85.6°/760mm, d_4^{20} 0.9261, n_D^{20} 1.4423, pK_{Est} ~ 4.2. Partially dried with Na_2CO_3 , then fractionally distd. The fraction b 84-85°, was refluxed with Na until hydrogen was no longer evolved when fresh Na was

added. It was then dried, and distd again through a 60 x 1.2cm column packed with glass rings [Brandon et al. *J Am Chem Soc* 72 2120 1950; UV: Elington et al. *J Chem Soc* 2873 1952, NMR: Bushweller and O'Neil *Tetrahedron Lett* 4713 1969]. It has been characterised as the 2,3,5-dinitrobenzoyloxy-tetrahydrofuran derivative, **m** 103° which forms pale yellow crystals from dihydropyran-Et₂O [Woods and Kramer *J Am Chem Soc* 69 2246 1947].

3,4-Dihydro-2H-pyrido[1,2a]-pyrimidin-2-one [5439-14-5] **M** 148.2, **m** 185-187°, 187-188°, 191-191.5°. Dissolve in CHCl₃, filter, evaporate then recrystallise the residue from EtOH-Me₂CO (needles) which can be washed with Et₂O and dried. It can also be recrystd from CHCl₃-pet ether or CHCl₃-hexane. The *hydrochloride* has **m** 295-295° (dec, from EtOH or MeOH-Et₂O), the *hydrobromide* has **m** 299-300°(dec, from MeOH-Et₂O) and the *picrate* has **m** 224-226°(corr), **m** 219-220° from EtOH. [Adams and Pachter *J Am Chem Soc* 74 4906 1952; Lappin *J Org Chem* 23 1358 1958; Hurd and Hayao *J Am Chem Soc* 77 115 1955.]

Dihydrotachysterol [67-96-9] **M** 398.7, **m** 125-127°, $[\alpha]_D^{20} +97^\circ$ (CHCl₃). Crystd from 90% MeOH.

1,8-Dihydroxyanthraquinone [117-10-2] **M** 240.1, **m** 193-197°, **pK₁²⁵** 8.30, **pK₂²⁵** 12.46. Crystd from EtOH and sublimed in a vacuum.

2,4-Dihydroxyazobenzene (Sudan orange G) [2051-85-6] **M** 214.2, **m** 228°, **pK_S(.Est(1)) <0**, **pK_{Est(2)} ~7.3**, **pK_{Est(3)} ~9.3**. Crystd from hot EtOH (charcoal).

2,3-Dihydroxybenzaldehyde [24677-78-9] **M** 138.1, **m** 135-136°, **pK₁²⁰** 7.73, **pK₂²⁰** 10.91. Crystd from water.

2,4-Dihydroxybenzoic acid [89-86-1] **M** 154.1, **m** 226-227°(dec), **pK₁²⁵** 3.30, **pK₂²⁵** 9.12, **pK₃²⁵** 15.6. Crystd from water.

2,5-Dihydroxybenzoic acid [490-79-9] **M** 154.1, **m** 204.5-205°, **pK²⁵** 2.95. Crystd from hot water or *benzene/acetone. Dried in a vacuum desiccator over silica gel.

2,6-Dihydroxybenzoic acid [303-07-1] **M** 154.1, **m** 167°(dec), **pK²⁵** 1.05. Dissolved in aqueous NaHCO₃ and the soln was washed with ether to remove non-acidic material. The acid was pptd by adding H₂SO₄, and recrystd from water. Dried under vacuum and stored in the dark [Lowe and Smith *J Chem Soc, Faraday Trans 1* 69 1934 1973].

2,4-Dihydroxybenzophenone [131-56-6] **M** 214.2, **m** 145.5-147° **pK_{Est(1)} ~7.0**, **pK_{Est(2)} ~12.0**. Recrystd from MeOH.

2,5-Dihydroxybenzyl alcohol (Gentisyl alcohol) [495-08-9] **M** 140.1, **m** 100° **pK_{Est(1)} ~9.3**, **pK_{Est(2)} ~11.3**. Crystd from CHCl₃. Sublimes at ~70° under high vacuum.

2,2'-Dihydroxybiphenyl [1806-29-7] **M** 186.2, **m** 108.5-109.5°, **pK₁²⁵** 7.56, **pK₂²⁵** 11.80. Repeatedly crystd from toluene, then sublimed at 60°/10⁻⁴mm.

3 α ,7 α -Dihydroxycholanolic acid (Chenodeoxycholic acid) [474-25-9] **M** 239.6, **m** 143°, $[\alpha]_{546}^{20} +14^\circ$ (c 2, EtOH), **pK_{Est} ~4.9**. Crystd from ethyl acetate.

7,8-Dihydroxycoumarin (Daphnetin) [486-35-1] **M** 178.2, **m** 256°(dec), **pK_{Est(1)} ~8.5**, **pK_{Est(2)} ~12.3**. Crystd from aqueous EtOH. Sublimed.

2,2'-Dihydroxy-6,6'-dinaphthyl disulfide [6088-51-3] **M** 350.5, **m** 220-223°. See 6-hydroxy-2-naphthyl disulfide on p. 264.

trans-2,3-Dihydroxy-1,4-dioxane [4845-50-5] M 120.1, m 91-95°, 100°. Recryst from Me₂CO. With phenylhydrazine it gives *glyoxal phenylhydrazone* m 175° (from Me₂CO-pet ether). The *diacetyl* derivative has m 105-106° [Head *J Chem Soc* 1036 1955, Raudnitz *Chem Ind (London)* 166 1956].

2,5-Dihydroxy-1,4-dithiane [40018-26-6] M 152.2, m (142-147° ?) 150-152°, 151°. Recrystd from EtOH. The *2,5-diethoxy-dithiane* has m 91° (92-93°) crystallises from pet ether and can be sublimed at 60°/0.001mm [Hormatka and Haber *Monatsh Chem* 85 1088 1954; Thiel et al. *Justus Liebigs Ann Chem* 611 121 1958; Hesse and Jøeder *Chem Ber* 85 924 1952].

(N,N-Dihydroxyethyl)glycine (BICINE) [150-25-4] M 163.2, m 193°(dec), pK₁²⁵ 1.81, pK₂²⁵ 8.27. Dissolved in a small volume of hot water and ppted with EtOH, twice. Repeated once more but with charcoal treatment of the aqueous soln, and filtered before addition of EtOH.

Dihydroxyfumaric (1,2-dihydroxybut-1-ene-1,2-dioic) acid dihydrate [133-38-0] M 184.1, m 155°(dec), pK₁²⁵ 1.57, pK₂²⁵ 3.36. Crystd from water.

3,4'-Dihydroxyisoflavone [578-86-9] M 256.3, m 234-236°. Crystd from aqueous 50% EtOH.

5,7-Dihydroxy-4'-methoxyflavone [480-44-4] M 284.3, m 261°. Crystd from 95% EtOH.

1,8-Dihydroxy-3-methylanthraquinone (chrysophanic acid) [481-74-3] M 245.3, m 196°, pK_{Est(1)} ~8.2, pK_{Est(2)} ~12.4. Crystd from EtOH or *benzene and sublimed in a vacuum.

1,5-Dihydroxynaphthalene [83-56-7] M 160.2, m 260°, 250-261°, pK_{Est} ~9.6. Crystd from nitromethane.

1,6-Dihydroxynaphthalene [575-44-0] M 160.2, m 138-139° (with previous softening), pK_{Est} ~9.4. Crystd from *benzene or *benzene/EtOH after treatment with charcoal.

2,5-Dihydroxyphenylacetic acid (homogentisic acid) [451-13-8] M 168.2, m 152°, 154-152°, pK²⁰ 4.14 (COOH). Crystd from EtOH/CHCl₃ or H₂O (sol 85% at 25°).

3,4-Dihydroxytoluene [452-86-8] M 124.1, m 65-66°, 68°, b 112°/3mm, 241°/760mm, pK₁²⁵ 9.44 (9.7), pK₂²⁵ 10.90 (11.9). Crystd from *C₆H₆. Purity checked by TLC. Crystd from high-boiling pet ether and distd in a vacuum.

1,3-Diiminoisoindoline [3468-11-9] M 145.2, m 193-194° (dec), 196° (dec), pK 8.27. It crystallises from H₂O, MeOH or MeOH-Et₂O (charcoal) in colourless prisms that become green on heating. [Elvidge and Linstead *J Chem Soc* 5000 1952]. IR (nujol): 3150 and 690 cm⁻¹, and UV: λ_{max} 251nm (ε 12.5K), 256nm (ε 12.5K) and 303nm (ε 4.6K) [Elvidge and Golden *J Chem Soc* 700 1957; Clark et al. *J Chem Soc* 3593 1953]. The *thiocyanate* has m 250-255° (dec), the *monohydrochloride* has m 300-301° (turns green) and the *dihydrochloride* has m 326-328° (turns green) and the *picrate* cryst from EtOH has m 299° (dec).

1,4-Diiodobenzene [624-38-4] M 329.9, m 132-133°. Crystd from EtOH or boiling MeOH, then air dried.

1,2-Diiodoethane [624-73-7] M 281.9, m 81-84°, d 2.134. Dissolved in ether, washed with satd aq Na₂S₂O₃, drying it over MgSO₄ and evap the ether *in vacuo* [Molander et al. *J Am Chem Soc* 109 453 1987].

5,7-Diiodo-8-hydroxyquinoline [83-73-8] M 397.0, m 214-215°(dec) pK_{Est(1)} ~3.2, pK_{Est(2)} ~8.2. Crystd from xylene and dried at 70° in a vacuum.

Diiodomethane (methylene diiodide) [75-11-6] M 267.8, m 6.1°, b 66-70°/11-12mm, d 3.325. Fractionally distd under reduced pressure, then fractionally crystd by partial freezing, and stabilized

with silver wool if necessary. It has also been purified by drying over CaCl_2 and fractionally distd from Cu powder.

S-3,5-Diiodotyrosine (iodogorgoic acid) [300-39-0] M 469.0, m 204°(dec), $[\alpha]_{\text{D}} +1.5^\circ$ (in 1M HCl) pK_1 2.12, pK_2 6.48, pK_3 7.82. See 3,5-diiodo-L-tyrosine dihydrate on p. 530 in Chapter 6.

Diisopropanolamine [110-97-4] M 133.2, m 41-44°, d 1.004, $\text{pK}_{\text{Est}} \sim 10.7$. Repeatedly crystd from dry diethyl ether.

Diisopropylamine [108-18-9] M 101.2, b 83.5°/760mm, n 1.39236, d 0.720, pK^{25} 11.20. Distd from NaOH, or refluxed over Na wire or NaH for three minutes and distd into a dry receiver under N_2 .
§ A polystyrene supported version of diisopropylamine is commercially available.

Diisopropylethylamine [7087-68-5] M 129.3, b 127°, $\text{pK}_{\text{Est}} \sim 10.9$. Distd from ninhydrin, then from KOH [Dryland and Sheppard, *J Chem Soc, Faraday Trans 1* 125 1986]. It is a strong base and should be stored in the absence of carbon dioxide.

(-)-2,3:4,6-Di-O-isopropylidene-2-keto-L-gulonic acid monohydrate (- DAG) [18467-77-1] M 292.3, m 100-101°, 103°, $[\alpha]_{\text{D}}^{25} -21.6^\circ$ (c 2.3, MeOH). Dissolve in Et_2O , filter, dry (MgSO_4), filter, evaporate to give a yellow oil. Addition of one drop of H_2O induces crystn to the monohydrate, which also forms rhombic crystals by recrystn from 95% $\text{EtOH-H}_2\text{O}$ at room temperature. [Flatt et al. *Synthesis* 815 1979; Reichstein and Grussner *Helv Chim Acta* 17 311 1934; Takagi and Jeffrey *Acta Crystallogr Sect B* 34 2932 1978; cf *Org Synth* 55 80 1976.]

1,2:5,6-Di-O-isopropylidene-D-mannitol [1707-77-3] M 262.3, m 121-125°, 122°, $[\alpha]_{\text{D}}^{25} +1.2^\circ$ (c 3, H_2O). Although quite soluble in H_2O it gives a purer product from this solvent, forming needles [Baer *J Am Chem Soc* 67 338 1945; NMR: Curtis et al. *J Chem Soc, Perkin Trans 1* 1756 1977].

Diisopropyl ketone (2,4-dimethyl-3-pentanone) [565-80-0] M 114.2, b 124°, d 0.801, n 1.400. Dried with CaSO_4 , shaken with chromatographic alumina and fractionally distd from P_2O_5 under N_2 .

Diketene [674-82-8] M 84.1, m -7°, b 66-68°/90mm, d 1.440, n 1.4376, n^{25} 1.4348. Diketene polymerizes violently in the presence of alkali. Distd at reduced pressure, then fractionally crystd by partial freezing (using as a cooling bath a 1:1 soln of $\text{Na}_2\text{S}_2\text{O}_3$ in water, cooled with Dry-ice until slushy, and stored in a Dewar flask). Freezing proceeds slowly, and takes about a day for half completion. The crystals are separated and stored in a refrigerator under N_2 . See ketene on p. 276.

2,2'-Diketospirilloxanthin [24009-17-4] M 624.9, m 225-227°, $\epsilon_{1\text{cm}}^{1\%}(\lambda_{\text{max}})$ 550(349nm), 820(422nm), 2125(488nm), 2725(516nm), 2130(551nm) in hexane. Purified by chromatography on a column of partially deactivated alumina. Crystd from acetone/pet ether. Stored in the dark, in an inert atmosphere at 0°.

Dilauroyl peroxide [105-74-8] M 398.6, m 53-54°. See lauryl peroxide (dodecyl peroxide) on p. 278.

Dimedone (5,5-dimethylcyclohexane-1,3-dione) [126-81-8] M 140.2, m 148-149°, pK^{25} 5.27. Crystd from acetone (ca 8mL/g), water or aqueous EtOH . Dried in air.

2,3-Dimercapto-1-propanol (BAL, British Anti-Lewisite) [59-52-9] M 124.2, b 82-84°/0.8mm, d 1.239, n 1.5732, pK_1^{25} 8.62, pK_2^{25} 10.75. Ppted as the Hg mercaptide [see Bjöberg *Chem Ber* 75 13 1942], regenerated with H_2S , and distd at 2.7mm [Rosenblatt and Jean *Anal Chem* 951 1955].

1,3-Dimercapto-2-propanol [584-04-3] M 124.2, b 82°/1.5mm. Purified as for 2,3-dimercapto-1-propanol above.

meso-2,3-Dimercaptosuccinic acid [304-55-2] M 182.2, m 191-192° (dec), 210° (dec), 210-211° (dec), pK_1^{25} 2.71, pK_2^{25} 3.48, pK_3^{25} 8.89, pK_4^{25} 10.75. Purified by dissolving in NaOH and precipitating with dilute HCl, dry and recrystallise from MeOH. IR has ν at 2544 (SH) and 1689 (CO₂H) cm^{-1} . The *bis-S-acetyl* deriv has m 183-185° (from EtOAc or Me₂CO) and its *Me ester* has m 119-120° (from pet ether) [Gerecke et al. *Helv Chim Acta* 44 957 1961; Owen and Sultanbawa *J Chem Soc* 3112 1949].

4,4'-Dimethoxyazobenzene [501-58-6] M 242.3, m 162.7-164.7°, pK_{Est} ~0. Chromatographed on basic alumina, eluted with *benzene. Crystd from 2:2:1 (v/v) methanol/ethanol/*benzene.

4,4'-Dimethoxyazoxybenzene [1562-94-3] M 258.3, transition temp 118-121°. See *p,p'*-azoxyanisole on p. 118.

1,2-Dimethoxybenzene (veratrole) [91-16-7] M 137.2, m 23°, b 208.5-208.7, d 1.085, n^{25} 1.53232. Steam distd. Fractionally distd from BaO, CaH₂ or Na. Crystd from *benzene or low-boiling pet ether at 0°. Fractionally crystd from its melt. Stored over anhydrous Na₂SO₄.

1,3-Dimethoxybenzene [151-10-0] M 137.2, b 212-213°, d 1.056, n 1.5215. Extracted with aqueous NaOH, and water, then dried. Fractionally distd from BaO or Na.

1,4-Dimethoxybenzene [150-78-7] M 137.2, m 57.2-57.8°. Steam distd. Crystd from hexane or *benzene, and from MeOH or EtOH but these are wasteful due to high solubilities. Dried under vacuum. Also sublimes under vacuum.

2,4-Dimethoxybenzoic acid [91-52-1] M 182.2, m 109°, pK^{25} 4.36. Crystd from water and dried in a vacuum desiccator over H₂SO₄.

2,6-Dimethoxybenzoic acid [1466-76-8] M 182.2, m 186-187°, pK^{25} 3.44. Crystd from water.

3,4-Dimethoxybenzoic acid (veratric acid) [93-07-2] M 182.2, m 181-182°, pK^{25} 4.43. Crystd from water or aq acetic acid.

3,5-Dimethoxybenzoic acid [1132-21-4] M 182.2, m 185-186°, pK^{25} 3.97. Crystd from water, EtOH or aq acetic acid.

***p,p'*-Dimethoxybenzophenone** [90-96-0] M 242.3, m 144.5°. Crystd from absolute EtOH.

2,6-Dimethoxy-1,4-benzoquinone [530-55-2] M 168.1, m 256°. Crystd from acetic acid. Sublimes in a vacuum.

1,1-Dimethoxyethane (acetaldehyde dimethyl acetal) [534-15-6] M 90.1, b 212°/760mm, d 0.828, n 1.4140. Purification as for acetal on p. 81. Also purified by GLC.

1,2-Dimethoxyethane (glycol dimethyl ether, glyme) [110-71-4] M 90.1, b 84°, d 0.867, n 1.380. Traces of water and acidic materials have been removed by refluxing with Na, K or CaH₂, decanting and distilling from Na, K, CaH₂ or LiAlH₄. Reaction has been speeded up by using vigorous high-speed stirring and molten potassium. For virtually complete elimination of water, 1,2-dimethoxyethane has been dried with Na-K alloy until a characteristic blue colour was formed in the solvent at Dry-ice/cellosolve temperatures: the solvent was kept with the alloy until distd for use [Ward *J Am Chem Soc* 83 1296 1961]. Alternatively, glyme, refluxed with benzophenone and Na-K, was dry enough if, on distn, it gave a blue colour of the ketyl immediately on addition to benzophenone and sodium [Ayscough and Wilson *J Chem Soc* 5412 1963]. Also purified by distn under N₂ from sodium benzophenone ketyl (see above).

5,6-Dimethoxy-1-indanone [2107-69-9] M 192.2, m 118-120°. Crystd from MeOH, then sublimed in a vacuum.

Dimethoxymethane (methylal) [109-87-5] M 76.1, b 42-46°/atm, d_4^{20} 0.8608, n_D^{20} 1.35335. See formaldehyde dimethyl acetal on p. 245.

1,4-Dimethoxynaphthalene [10075-62-4] M 188.2, m 87-88°. Crystd from EtOH.

1,5-Dimethoxynaphthalene [10075-63-5] M 188.2, m 183-184°. Crystd from EtOH.

2,6-Dimethoxyphenol [91-10-1] M 154.2, m 54-56°, pK_{Est} ~9.6. Purified by zone melting or sublimation in a vacuum.

3,5-dimethoxyphenol (phloroglucinol dimethylether) [500-99-2] M 154.2, m 42-43°, b 115°/0.04mm, pK^{25} 9.35. Purified by distn followed by sublimation in a vacuum.

3,4-Dimethoxyphenyl acetic acid (homoveratric acid) [93-40-3] M 196.2, m 97-99°, pK^{25} 4.33. Crystd from water or *benzene/ligroin.

3,5-Dimethoxyphenylacetonitrile [13388-75-5] M 177.1, m 53°. Crystd from MeOH or pet ether (b 90-110°). [Adams et al. *J Am Chem Soc* 70 664 1948; Sankaraman et al. *J Am Chem Soc* 109 5235 1987.]

4,4'-Dimethoxythiobenzophenone [958-80-5] M 258.3, m 120°. Recrystd from a mixture of cyclohexane/dichloromethane (4:1).

2,6-Dimethoxytoluene [5673-07-4] M 152.2, m 39-41°. Sublimed *in vacuo* [Sankaraman et al. *J Am Chem Soc* 109 5235 1987].

4,4'-Dimethoxytrityl chloride (DMT) [40615-36-9] M 338.8, m 114°. Crysts from cyclohexane-acetyl chloride as the hydrochloride and dry over KOH pellets in a desiccator. When dissolved in *C₆H₆ and air is blown through, HCl is removed. It crystallises from Et₂O. [*Justus Liebigs Ann Chem* 370 142 1909; *Chem Ber* 36 2774 1903; Smith et al. *J Am Chem Soc* 84 430 1962; Smith et al. *J Am Chem Soc* 85 3821 1963.] If it had hydrolysed considerably (see OH in IR) then repeat the crystallisation from cyclohexane-acetyl chloride — excess of AcCl is removed in vac over KOH.

N,N-Dimethylacetamide [127-19-5] M 87.1, b 58.0-58.5°/11.4mm, d 0.940, n 1.437. Shaken with BaO for several days, refluxed with BaO for 1h, then fractionally distd under reduced pressure, and stored over molecular sieves.

β,β -Dimethylacrylic acid (senecioic acid) [541-47-9] M 100.1, m 68°, pK^{25} -5.4 (aq H₂SO₄). Crystd from hot water or pet ether (b 60-80°).

Dimethyl adipate [627-93-0] M 174.2, m 9-11°, b 109°/10mm, 121-123°/20mm, 235°/760mm, d_4^{20} 1.0642, n_D^{20} 1.4292. Dissolve in Et₂O, wash with NaHCO₃, H₂O, dry over MgSO₄, filter, evaporate and distil several times until the IR and NMR are consistent with the structure [Lorette and Brown *J Org Chem* 24 261 1959; Hoffmann and Weiss *J Am Chem Soc* 79 4759 1957].

Dimethyl adipimidate dihydrochloride [14620-72-5] M 245.1, m 218-220°, 222-224°. If the salt smells of HCl then wash with MeOH and dry Et₂O (1:3) under N₂ until the HCl is completely removed. Recryst from MeOH-Et₂O (it is very important that the solvents are super dry) [Hartman and Wold *Biochemistry* 6 2439 1967; McElvain and Shroeder *J Am Chem Soc* 71 40 1949].

Dimethylamine [124-40-3] M 45.1, f -92.2°, b 0°/563mm, 6.9°/760mm, pK^{25} 10.73. Dried by passage through a KOH-filled tower, or by standing with sodium pellets at 0° during 18h. § A dimethylaminomethyl polystyrene supported version is commercially available.

Dimethylamine hydrochloride [506-59-2] **M 81.6, m 171°**. Crystd from hot CHCl_3 or abs EtOH. Also recrystd from MeOH/ether soln. Dried in a vacuum desiccator over H_2SO_4 , then P_2O_5 . *Hygroscopic*.

4-*N,N'*-Dimethylaminoazo-benzene-4'-isothiocyanate {DABITC, 4-[(4-isocyanatophenyl)-azo]-*N,N'*-dimethylaniline} [7612-98-8] **M 282.4, m 170-171°, $\text{pK}_{\text{Est}} \sim 2.5$** . Crystd by dissolving 1g in 150mL of boiling Me_2CO , filtering hot and allowing to cool at -20° overnight collecting the solid and drying in vac. Solns in pyridine should be used immediately otherwise it dec. Moisture sensitive. [Chang *Methods Enzymol* 91 79, 455 1983.]

***p*-Dimethylaminoazobenzene (Methyl Yellow)** [60-11-7] **M 225.3, m 118-119°(dec), $\text{pK}_1^{25} -5.34$ (aq H_2SO_4), $\text{pK}_2^{25} 2.96$** . Crystd from acetic acid or isooctane, or from 95% EtOH by adding hot water and cooling. Dried over KOH under vacuum at 50° . **CARCINOGEN**.

***p*-Dimethylaminobenzaldehyde (Ehrlich's Reagent)** [100-10-7] **M 149.2, m 74-75°, $\text{pK}_{\text{Est}} \sim 2.6$** . Crystd from water, hexane, or from EtOH (2mL/g), after charcoal treatment, by adding excess of water. Also dissolved in aqueous acetic acid, filtered, and ppted with ammonia. Finally recrystd from EtOH.

***p*-Dimethylaminobenzoic acid** [619-84-1] **M 165.2, m 242.5-243.5°(dec), $\text{pK}_1 2.51$, $\text{pK}_2 6.03$** . Crystd from EtOH/water.

***p*-Dimethylaminobenzophenone** [530-44-9] **M 225.3, m 92-93°, $\text{pK}_{\text{Est}} \sim 2.7$** . Crystd from EtOH.

***N,N*-Dimethylamino-*p*-chlorobenzene (*p*-chloro-*N,N*-dimethylaniline)** [698-69-1] **M 155.6, m 32-33.5°, 35.5°, b 231°/atm**. Purified by vacuum sublimation [Guarr et al. *J Am Chem Soc* 107 5104 1985]. The *picrate* has **m 126-128°** (from methanol).

2*S*,3*R*-(+)-4-Dimethylamino-1,2-diphenyl-3-methyl-2-butanol [38345-66-3] **M 283.4, m 55-57°, $[\alpha]_{546}^{20} +9.3^\circ$ (c 9.6, EtOH), $[\alpha]_{\text{D}}^{20} +7.7^\circ$ (c 9.6, EtOH), $\text{pK}_{\text{Est}} \sim 10.0$** . Purification of the *hydrochloride* by dissolving 1.5g in 13.5 mL of 5N HCl heating to boiling and evaporate in a vacuum. Recrystn of the *hydrochloride* three times from MeOH-EtOAc gives **m 189-190°**, $[\alpha]_{\text{D}} -33.7^\circ$ (c 1, H_2O) {enantiomer has $+34.2^\circ$ }. The *hydrochloride* in the minimum volume of water is basified with aqueous 5N NaOH and extracted with Et_2O . The extract is dried (K_2CO_3) and evap leaving a residue which is stored in a desiccator over solid KOH as a low melting solid. It can be recovered with these procedures from asymmetric reductions with LAH, and reused. [*J Am Chem Soc* 77 3400 1955; *J Org Chem* 28 2381 2483 1963.]

***dl*-4-Dimethylamino-2,2-diphenylvaleramide** [60-46-8] **M 296.4, m 183-184°, $\text{pK}_{\text{Est}} \sim 9.8$** . Crystd from aqueous EtOH.

(-)-*L*-4-Dimethylamino-2,2-diphenylvaleramide [6078-64-4] **M 296.4, m 136.5-137.5°**. Crystd from pet ether or EtOH.

2-Dimethylaminoethanol [108-01-0] **M 89.1, b 134.5-135.5°, d 1.430, n 1.4362, $\text{pK}^{25} 9.23$** . Dried with anhydrous K_2CO_3 or KOH, and fractionally distd.

1-(3-Dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (EDCI, DEC, 1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide hydrochloride) [25952-53-8] **M 191.7, m 113.5-114.5°, 114-116°, $\text{pK}_{\text{Est}} \sim 10.3$** . An excellent H_2O -soluble peptide coupling reagent. It is purified by dissolving (ca 1g) in CH_2Cl_2 (10mL) at room temperature and then add dry Et_2O (~110mL) dropwise and the crystals that separate are collected, washed with dry Et_2O and recrystd from CH_2Cl_2 - Et_2O and dried in a vacuum over P_2O_5 . It is important to work in a dry atmosphere or work rapidly and then dry the solid as soon as possible. Material is moderately *hygroscopic* but once it becomes wet it reacts slowly with H_2O . Store away from moisture and at -20° to slow down the hydrolysis process. The *free base* has **b 47-48°/0.27mm, 53-54°/0.6mm, $n_{\text{D}}^{25} 1.4582$** . The *methiodide* is recrystallised from CHCl_3 -EtOAc, the crystals are filtered off, washed with dry Et_2O and recrystd from CHCl_3 - Et_2O , and dried *in vacuo* over P_2O_5 , **m 93-95°, 94-95°**. [Sheehan et al. *J Am Chem Soc* 87 2492 1965; Sheehan and Cruickshank *Org Synth Coll Vol V* 555 1973.]

§ A polymer bound version is commercially available.

6-Dimethylaminopurine [938-55-6] M 163.1, m 257.5-258.5°, 259-262°, 263-264°, pK_1^{25} 3.87, pK_2^{25} 10.5. It is purified by recrystn from H₂O, EtOH (0.32g in 10mL) or CHCl₃. [Albert and Brown *J Chem Soc* 2060 1954; UV: Mason *J Chem Soc* 2071 1954.] The *monohydrochloride* crystallises from EtOH-Et₂O, m 253° (dec) [Elion et al. *J Am Chem Soc* 74 411 1952], the *dihydrochloride* has m 225° (dec) and the *picrate* has m 245° (235-236.5°) [Fryth et al. *J Am Chem Soc* 80 2736 1958].

4-Dimethylaminopyridine (DMAP) [1122-58-3] M 122.2, m 108-109°, b 191°, pK^{25} 9.61. Recrystd from toluene [Sadownik et al. *J Am Chem Soc* 108 7789 1986].

§ A polystyrene supported version (PS-DMAP) is commercially available.

***N,N*-Dimethylaniline** [121-69-7] M 121.2, f 2°, b 84°/15mm, 193°/760mm, d 0.956, n^{25} 1.5556, pK^{25} 5.07. Primary and secondary amines (including aniline and monomethylaniline) can be removed by refluxing for some hours with excess acetic anhydride, and then fractionally distilling. Crocker and Jones (*J Chem Soc* 1808 1959) used four volumes of acetic anhydride, then distd off the greater part of it, and took up the residue in ice-cold dil HCl. Non-basic materials were removed by ether extraction, then the dimethylaniline was liberated with ammonia, extracted with ether, dried, and distd under reduced pressure. Metzler and Tobolsky (*J Am Chem Soc* 76 5178 1954) refluxed with only 10% (w/w) of acetic anhydride, then cooled and poured into excess 20% HCl, which, after cooling, was extracted with diethyl ether. (The amine hydrochloride, remains in the aqueous phase.) The HCl soln was cautiously made alkaline to phenolphthalein, and the amine layer was drawn off, dried over KOH and fractionally distd under reduced pressure, under nitrogen. Suitable drying agents for dimethylaniline include NaOH, BaO, CaSO₄, and CaH₂.

Other purification procedures include the formation of the picrate, prepared in *benzene soln and crystd to constant melting point, then decomposed with warm 10% NaOH and extracted into ether: the extract was washed with water, and distd under reduced pressure. The oxalate has also been used. The base has been fractionally crystd by partial freezing and also from aq 80% EtOH then from absolute EtOH. It has been distd from zinc dust, under nitrogen.

2,6-Dimethylaniline [87-62-7] M 121.2, m 11°, b 210-211°/736mm, d 0.974, n 1.5604, pK^{25} 3.95. Converted to its hydrochloride which, after recrystn, was decomposed with alkali to give the free base. Dried over KOH and fractionally distd.

3,4-Dimethylaniline [95-64-7] M 121.2, m 51°, b 116-118°/25mm, b 226°/760mm, pK^{25} 5.17. Crystd from ligroin and distilled under vacuum.

9,10-Dimethylanthracene [781-43-1] M 206.3, m 180-181°. Crystd from EtOH, and by recrystn from the melt.

1,3-Dimethylbarbituric acid [769-42-6] M 156.1, m 123°, pK^{25} 4.56. Crystd from water and sublimed in a vacuum. Also purified by dissolving 10g in 100mL of boiling CCl₄/CHCl₃ (8:2) (1g charcoal), filtered and cooled to 25°. Dried *in vacuo* [Kohn et al. *Anal Chem* 58 3184 1986].

7,12-Dimethylbenz[*a*]anthracene [57-97-6] M 256.4, m 122-123°. Purified by chromatography on alumina/toluene or *benzene. Crystd from acetone/EtOH.

5,6-Dimethylbenzimidazole [582-60-5] M 146.2, m 205-206°, pK_1^{25} 5.96, pK_2^{25} 12.52. Crystd from diethyl ether. Sublimed at 140°/3mm.

2,3-Dimethylbenzoic acid [603-79-2] M 150.2, m 146°, pK^{25} 3.72. Crystd from EtOH and is volatile in steam.

2,4-Dimethylbenzoic acid [611-01-8] M 150.2, m 126-127°, b 267°/727mm, pK^{25} 4.22. Crystd from EtOH, and sublimed in a vacuum.

2,5-Dimethylbenzoic acid [610-72-0] M 150.2, m 134°, b 268°/760mm, pK²⁵ 4.00. Steam distd, and crystd from EtOH.

2,6-Dimethylbenzoic acid [632-46-2] M 150.2, m 117°, pK²⁵ 3.35. Steam distd, and crystd from EtOH.

3,4-Dimethylbenzoic acid [619-04-5] M 150.2, m 166°, pK²⁵ 4.50. Crystd from EtOH and sublimed *in vacuo*.

3,5-Dimethylbenzoic acid [499-06-9] M 150.2, m 170°, pK²⁵ 4.30. Distd in steam, crystd from water or EtOH and sublimed in a vacuum.

4,4'-Dimethylbenzophenone [611-97-2] M 210.3, m 95°, b 333-334°/725mm. Purified by zone refining.

2,5-Dimethyl-1,4-benzoquinone [137-18-8] M 136.1, m 124-125°. Crystd from EtOH.

2,6-Dimethyl-1,4-benzoquinone [527-61-7] M 136.1, m 72° (sealed tube). Crystd from water/EtOH (8:1).

2,3-Dimethylbenzothiophene [31317-17-6] M 212.3, b 123-124°/10mm, n¹⁹ 1.6171. Fractionated through a 90cm Monel spiral column, or other efficient fractionating or spinning band column and collecting the middle fraction.

N,N-Dimethylbenzylamine [103-83-3] M 135.2, b 66-67°/15mm, 181°/760mm, d₄²⁰ 0.898, n_D²⁰ 1.516, pK²⁵ 8.91. See *N*-benzyl dimethylamine on p. 128.

4,4'-Dimethyl-2,2'-bipyridine [1134-35-6] M 184.2, m 175-176°, pK_{Est(1)} ~0.2, pK_{Est(2)} ~4.9. Crystd from ethyl acetate. [Elliott et al. *J Am Chem Soc* 107 4647 1985.]

1,1'-Dimethyl-4,4'-bipyridylium dichloride (3H₂O; Methyl Viologen Dichloride, paraquat dichloride) [1910-42-5] M 311.2, m >300°(dec). Recrystd from MeOH/acetone mixture. Also crystd three times from absolute EtOH [Bancroft et al. *Anal Chem* 53 1390 1981]. Dried at 80° in a vacuum.

N,N-Dimethylbiuret [7710-35-2] M 131.1, m 178°. Purified by repeated crystn from the melt, or H₂O. [Bredereck and Richter *Chem Ber* 99 2461 1968; Dunning and Close *J Am Chem Soc* 75 3615 1953.]

2,3-Dimethyl-1,3-butadiene [513-81-5] M 82.2, m -69-70°, b 68-69°/760mm, d 0.727, n 1.4385. Distd from NaBH₄, and purified by zone melting.

1,3-Dimethylbutadiene sulfone (1,3-dimethylsulfolene) [10033-92-8] M 145.2, m 40.4-41.0°. Crystd from diethyl ether.

2,2-Dimethylbutane [75-83-2] M 86.2, b 49.7°, d 0.649, n²⁵ 1.36595. Distd azeotropically with MeOH, then washed with water, dried and distd.

2,3-Dimethylbutane [79-29-8] M 86.2, b 58.0°, d 1.375, n²⁵ 1.37231. Distd from sodium, passed through a column of silica gel (activated by heating in nitrogen to 350° before use) to remove unsaturated impurities, and again distd from sodium. Also distilled azeotropically with MeOH, then washed with water, dried and redistd.

2,3-Dimethylbut-2-ene [563-79-1] M 84.2, b 72-73°/760mm, d 0.708, n 1.41153. Purified by GLC on a column of 20% squalene on chromosorb P at 50° [Flowers and Rabinovitch *J Phys Chem* 89 563 1985]. Also washed with 1M NaOH soln followed by H₂O. Dried over Na₂SO₄, distd over powdered KOH

under nitrogen and passed through activated alumina before use. [Woon et al. *J Am Chem Soc* **108** 7990 1986; Wong et al. *J Am Chem Soc* **109** 3428 1987.]

Dimethylcarbamoyl chloride [79-44-7] **M 107.5, m -33°, b 34°/0.1mm, d 1.172, n 1.4511.** Must distil under high vacuum to avoid decomposition.

3,3'-Dimethylcarbanilide [620-50-8] **M 240.3, m 225°.** Crystd from ethyl acetate.

Dimethyl carbonate [616-38-5] **M 90.1, m 4.65°, b 90-91°, d 1.070, n 1.369.** Contains small amounts of water and alcohol which form azeotropes. Stood for several days in contact with Linde type 4A molecular sieves, then fractionally distd. The middle fraction was frozen slowly at 2°, several times, retaining 80% of the solvent at each cycle.

cis- and trans-1,4-Dimethylcyclohexane [589-90-2] **M 112.2, b 120°, d 0.788, n 1.427.** Freed from olefins by shaking with conc H₂SO₄, washing with water, drying and fractionally distilling.

1,2-Dimethylcyclohexene [1674-10-8] **M 110.2, b 135-136°/760mm, d 0.826, n 1.4591.** Passed through a column of basic alumina and distd.

1,5-Dimethyl-1,5-diazaundecamethylene polymethobromide (Hexadimethrene, Polybrene) [28728-55-4] **M 5000—10,000 polymer** Purified by chromatography on Dowex 50 and/or by filtration through alumina before use [Frank Hoppe-Seyler's *Z Physiol Chemie* **360** 997 1979]. Hygroscopic, sol in H₂O is 10%.

2,9-Dimethyl-4,7-diphenyl-1,10-phenanthroline [4733-39-5] **M 360.5, m >280°, pK_{Est} ~5.6.** Purified by recrystn from *benzene.

Dimethyl disulfide [624-92-0] **M 94.2, f -98°, b 40°/12mm, 110°/760mm, d 1.0605, n 1.5260.** Passed through neutral alumina before use.

2,2-Dimethylethyleneimine [2658-24-4] **M 71.1, b 70.5-71.0°, pK²⁵ 8.64.** Freshly distd from sodium before use.

N,N-Dimethylformamide (DMF) [68-12-2] **M 73.1, b 40°/10mm, 61°/30mm, 88°/100mm, 153°/760mm, d 0.948, n²⁵ 1.4269, pK -0.3.** Decomposes slightly at its normal boiling point to give small amounts of dimethylamine and carbon monoxide. The decomposition is catalysed by acidic or basic materials, so that even at room temperature DMF is appreciably decomposed if allowed to stand for several hours with solid KOH, NaOH or CaH₂. If these reagents are used as dehydrating agents, therefore, they should not be refluxed with the DMF. Use of CaSO₄, MgSO₄, silica gel or Linde type 4A molecular sieves is preferable, followed by distn under reduced pressure. This procedure is adequate for most laboratory purposes. Larger amounts of water can be removed by azeotropic distn with *benzene (10% v/v, previously dried over CaH₂), at atmospheric pressure: water and *benzene distil below 80°. The liquid remaining in the distn flask is further dried by adding MgSO₄ (previously ignited overnight at 300-400°) to give 25g/L. After shaking for one day, a further quantity of MgSO₄ is added, and the DMF distd at 15-20mm pressure through a 3-ft vacuum-jacketed column packed with steel helices. However, MgSO₄ is an inefficient drying agent, leaving about 0.01M water in the final DMF. More efficient drying (to around 0.001-0.007M water) is achieved by standing with powdered BaO, followed by decanting before distn, with alumina powder (50g/L; previously heated overnight to 500-600°), and distilling from more of the alumina; or by refluxing at 120-140° for 24h with triphenylchlorosilane (5-10g/L), then distilling at ca 5mm pressure [Thomas and Rochow *J Am Chem Soc* **79** 1843 1957]. Free amine in DMF can be detected by colour reaction with 1-fluoro-2,4-dinitrobenzene. It has also been purified by drying overnight over KOH pellets and then distd from BaO through a 10 cm Vigreux column [*Exp Cell Res* **100** 213 1976]. [For efficiency of desiccants in drying dimethylformamide see Burfield and Smithers [*J Org Chem* **43** 3966 1978, and for a review on purification, tests of purity and physical properties, see Juillard *Pure Appl Chem* **49** 885 1977].

It has been purified by distilling from K_2CO_3 under high vac and fractionated in an all-glass apparatus. The middle fraction is collected, degassed (seven or eight freeze-thaw cycles) and redistd under as high a vacuum as possible [Mohammad and Kosower *J Am Chem Soc* **93** 2713 1971].

Rapid purification: Stir over CaH_2 (5% w/v) overnight, filter, then distil at 20mmHg. Store the distd DMF over 3A or 4A molecular sieves. For solid phase synthesis, the DMF used must be of high quality and free from amines.

***d,l*-2,4-Dimethylglutaric acid** [2121-67-7] M 160.2, m 144-145° $pK_{Est(1)} \sim 4.4$, $pK_{Est(2)} \sim 5.4$. Distd in steam and crystd from ether/pet ether.

3,3-Dimethylglutaric acid [4839-46-7] M 160.2, m 103-104°, b 89-90°/2mm, 126-127°/4.5mm, $pK_1^{25} 3.85$, $pK_2^{25} 6.45$. Crystd from water, *benzene or ether/pet ether. Dried in a vacuum.

3,3-Dimethylglutarimide [1123-40-6] M 141.2, m 144-146°. Recrystd from EtOH [Arnett and Harrelson *J Am Chem Soc* **109** 809 1987].

***N,N*-Dimethylglycinehydrazide hydrochloride** [539-64-0] M 153.6, m 181°. Crystd by adding EtOH to a conc aqueous soln.

Dimethylglyoxime [95-45-4] M 116.1, m 240°, $pK_1^{25} 10.60$, $pK_2^{25} 11.85$. Crystd from EtOH (10mL/g) or aqueous EtOH. **TOXIC.**

2,5-Dimethyl-2,4-hexadiene [764-13-6] M 110.2, f 14.5°, b 132-134°, d 0.773, n 1.4796. Distd, then repeatedly fractionally crystd by partial freezing. Immediately before use, the material was passed through a column containing Woelm silica gel (activity I) and Woelm alumina (neutral) in separate layers.

2,2-Dimethylhexane [590-73-8] M 114.2, m -121.2°, b 107°, d 0.695. Dried over type 4A molecular sieves and distd.

2,5-Dimethylhexane [592-13-2] M 114.2, m -91.2°, b 109°, d 0.694. Dried over type 4A molecular sieves and distd.

2,5-Dimethylhexane-2,5-diol [110-03-2] M 146.2, m 88-90°. Purified by fractional crystn. Then the diol was dissolved in hot acetone, treated with activated charcoal, and filtered while hot. The soln was cooled and the diol was filtered off and washed well with cold acetone. The crystn process was repeated several times and the crystals were dried under a vac in a freeze-drying apparatus [Goates et al. *J Chem Soc, Faraday Trans 1* **78** 3045 1982].

5,5-Dimethylhydantoin [77-71-4] M 128.1, m 177-178° $pK^{24} 9.19$. Crystd from EtOH and sublimed *in vacuo*.

1,1-Dimethylhydrazine [57-14-7] M 60.1, b 60.1°/702mm, d 0.790, n 1.408 $pK^{30} 7.21$. Fractionally distd through a 4-ft column packed with glass helices. Pptd as its oxalate from diethyl ether soln. After crystn from 95% EtOH, the salt was decomposed with aqueous saturated NaOH, and the free base was distd, dried over BaO and redistd [McBride and Kruse *J Am Chem Soc* **79** 572 1957]. Distn and storage should be under nitrogen.

4,6-Dimethyl-2-hydroxypyrimidine [108-79-2] M 124.1, m 198-199°, $pK_1^{20} 3.77$, $pK_2^{20} 10.50$. Crystd from absolute EtOH (charcoal).

1,2-Dimethylimidazole [1739-84-0] M 96.1, m 38-40°, b 206°/760mm, d 1.084, $pK_{Est} \sim 8.1$. Crystd from *benzene and stored at 0-4°. [Gorun et al. *J Am Chem Soc* **109** 4244 1987.]

1,1-Dimethyl-1*H*-indene [18636-55-0] M 144.2, b 57°/4.8mm, 115°/20mm. Purified by gas chromatography or by fractional distn.

Dimethyl itaconate [617-52-7] M 158.2, m 38°, b 208°, d 1.124. Crystd from MeOH by cooling to -78°.

Dimethylmaleic anhydride [766-39-2] M 126.1, m 96°, b 225°/760mm. Distd from *benzene/ligroin and sublimed in a vacuum.

Dimethylmalonic acid [595-46-0] M 132.1, m 192-193° pK₁²⁵ 3.03, pK₂²⁵ 5.73. Crystd from *benzene/pet ether and sublimed in a vacuum with slight decomposition.

1,5-Dimethylnaphthalene [571-61-9] M 156.2, m 81-82°, b 265-266°. Crystd from 85% aq EtOH.

2,3-Dimethylnaphthalene [581-40-8] M 156.2, m 104-104.5°. Steam distd and crystd from EtOH.

2,6-Dimethylnaphthalene [581-42-0] M 156.2, m 110-111°, b 122.5-123.5°/10mm, 261-262°/760mm. Distd in steam and crystd from EtOH.

3,3'-Dimethylnaphthidine (4,4'-diamino-3,3'-dimethyl-1,1'-binaphthyl) [13138-48-2] M 312.4, m 213°. Recrystd from EtOH or pet ether (b 60-80°).

***N,N*-Dimethyl-*m*-nitroaniline** [619-31-8] M 166.1, m 60°, pK²⁵ 2.63. Crystd from EtOH.

***N,N*-Dimethyl-*p*-nitroaniline** [100-23-2] M 166.1, m 164.5-165.2°, pK²⁵ 0.61 (0.92). Crystd from EtOH or aqueous EtOH. Dried under vacuum.

Dimethylnitrosamine (*N*-nitrosodimethylamine) [62-75-9] M 74.0, m -28°, b 149-150°/atm, 153°/774mm, d. 1.006, n 1.4370. Dry over anhyd K₂CO₃ or dissolve in Et₂O, dry over solid KOH, filter, evap Et₂O and distil yellow oily residue through a 30cm fractionating column discarding the first fraction which may contain Me₂N. Also dried over CaCl₂ and distd at atm pressure. All should be done in an efficient fume cupboard as the vapors are **TOXIC** and **CARCINOGENIC**. [Fischer *Chem Ber* 8 1588 1875; Romberg *Recl Trav Chim, Pays-Bas* 5 248 1886; Hatt *Org Synth Coll Vol II* 211 1961; Krebs and Mandt *Chem Ber* 108 1130 1975.]

***N,N*-Dimethyl-*p*-nitrosoaniline** [138-89-6] M 150.2, m 86-87°. See 4-nitroso-*N,N*-dimethylaniline on p. 314.

***N,N*-Dimethyl-*p*-nitrosoaniline hydrochloride** [42344-05-8] M 186.7, m 177°. Crystd from hot water in the presence of a little HCl.

2,6-Dimethyl-2,4,6-octatriene [7216-56-0; *cis/trans* mixt 673-84-7; *trans, trans* 3016-19-1] M 136.2, b 80-82°/15mm, ε_{278nm} 42,870. Repeated distn at 15mm through a long column of glass helices, the final distn being from sodium under nitrogen. See *neo*-allocimene on p. 100.

Dimethylolurea [140-95-4] M 120.1, m 137-139°. Crystd from aqueous 75% EtOH.

Dimethyl oxalate [553-90-2] M 118.1, m 54°, b 163-165°, d 1.148. Crystd repeatedly from EtOH. Degassed under nitrogen high vacuum and distd.

3,3-Dimethyloxetane [6921-35-3] M 86.1, b 79.2-80.3°/760mm. Purified by gas chromatography using a 2m silicone oil column.

2,3-Dimethylpentane [565-59-3] M 100.2, b 89.8°, d 0.695, n 1.39197, n²⁵ 1.38946. Purified by azeotropic distn with EtOH, followed by washing out the EtOH with water, drying and distn [Streiff et al. *J Res Nat Bur Stand* 37 331 1946].

2,4-Dimethylpentane [108-08-7] **M 100.2, b 80.5°, d 0.763, n 1.3814, n²⁵ 1.37882.** Extracted repeatedly with conc H₂SO₄, washed with water, dried and distd. Percolated through silica gel (previously heated in nitrogen to 350°). Purified by azeotropic distn with EtOH, followed by washing out the EtOH with water, drying and distn.

4,4-Dimethyl-1-pentene [762-62-9] **M 98.2, b 72.5°/760mm, d 0.6827, n 1.3918.** Purified by passage through alumina before use [Traylor et al. *J Am Chem Soc* **109** 3625 1987].

Dimethyl peroxide [690-02-8] **M 62.1, b 13.5°/760mm, d 0.8677, n 1.3503.** Purified by repeated trap-to-trap fractionation until no impurities could be detected by gas IR spectroscopy [Haas and Oberhammer *J Am Chem Soc* **106** 6146 1984]. *All necessary precautions should be taken in case of EXPLOSION.*

2,9-Dimethyl-1,10-phenanthroline [484-11-7] **M 208.3, m 162-164°, pK²⁵ 5.85.** Purified as hemihydrate from water, and as anhydrous from *benzene.

R-(+)-N,N'-Dimethyl-1-phenethylamine [19342-01-9] and **S-(-)-N,N'-Dimethyl-1-phenethylamine** [17279-31-1] **M 149.2, b 81°/16mm, [α]_D²⁰ (+) and (-) 50.2° (c 1, MeOH), [α]_D²⁶ +61.8° and -64.4° (neat l 1), d 0.908, pK_{Est} ~9.0 (for RS).** The amine is mixed with aqueous 10N NaOH and extracted with toluene. The extract is washed with saturated aqueous NaCl and dried over K₂CO₃, and transferred to fresh K₂CO₃ until the soln is clear, and filtered. The filtrate is distd. If a short column packed with glass helices is used, the yield is reduced but a purer product is obtained. [*Org Synth* **25** 89 1945; *J Am Chem Soc* **71** 291 3929 3931 4165 1949.] The (-)-picrate has m 140-141° (cryst from EtOH). The racemate [1126-71-2] has b 88-89°/16mm, 92-94°/30mm, 194-195°/atm, d₄²⁰ 0.908.

2,3-Dimethylphenol [526-75-0] **M 122.2, m 75°, b 120°/20mm, 218°/760mm, pK²⁵ 10.54.** Crystd from aqueous EtOH.

2,5-Dimethylphenol [95-87-1] **M 122.2, m 73°, b 211.5°/762mm, pK²⁵ 10.41.** Crystd from EtOH/ether.

2,6-Dimethylphenol [576-26-1] **M 122.2, m 49°, b 203°/760mm, pK²⁵ 10.61.** Fractionally distd under nitrogen, crystd from *benzene or hexane, and sublimed at 38°/10mm.

3,4-Dimethylphenol [95-65-8] **M 122.2, m 65°, b 225°/757mm, pK²⁵ 10.36.** Heated with an equal weight of conc H₂SO₄ at 103-105° for 2-3h, then diluted with four volumes of water, refluxed for 1h, and either steam distd or extracted repeatedly with diethyl ether after cooling to room temperature. The steam distillate was also extracted and evaporated to dryness. (The purification process depends on the much slower sulfonation of 3,5-dimethylphenol than most of its likely contaminants.) It can also be crystd from water, hexane or pet ether, and vacuum sublimed. [Kester *Ind Eng Chem (Anal Ed)* **24** 770 1932; Bernasconi and Paschalis *J Am Chem Soc* **108** 2969 1986.]

3,5-Dimethylphenol [108-68-9] **M 122.2, m 68°, b 219°, pK²⁵ 10.19.** Purification as for 3,4-dimethylphenol.

Dimethyl phthalate [131-11-3] **M 194.2, b 282°, n 1.5149, d 1.190, d²⁵ 1.1865.** Washed with aqueous Na₂CO₃, then distilled water, dried (CaCl₂) and distd under reduced pressure (b 151-152°/0.1mm).

2,2-Dimethyl-1,3-propanediol (neopentyl glycol) [126-30-7] **M 104.2, m 128.4-129.4°, b 208°/760mm.** Crystd from *benzene or acetone/water (1:1).

2,2-Dimethyl-1-propanol (neo-pentyl alcohol) [75-84-3] **M 88.2, m 52°, b 113.1°/760mm.** Difficult to distil because it is a solid at ambient temperatures. Purified by fractional crystallisation and sublimation.

***N,N*-Dimethylpropionamide** [758-96-3] M 101.2, b 175-178°, d 0.920, n 1.440. Shaken over BaO for 1-2 days, then distd at reduced pressure.

2,5-Dimethylpyrazine [123-32-0] M 108.1, b 156°, d 0.990, n 1.502, $pK_1^{2.5}$ -4.6 (aq H_2SO_4), $pK_2^{2.5}$ 1.85. Purified *via* its picrate (m 150°) [Wiggins and Wise *J Chem Soc* 4780 1956].

3,5-Dimethylpyrazole [67-51-6] M 96.1, m 107-108°, $pK^{2.0}$ 4.16. Crystd from cyclohexane or water. [Barszez et al. *J Chem Soc, Dalton Trans* 2025 1986.]

2,3-Dimethylquinoxaline [2379-55-7] M 158.2, m 106°, $pK^{2.5}$ -3.84 (aq H_2SO_4). Crystd from distilled water.

2,4-Dimethylresorcinol [634-65-1] M 138.1, m 149-150° $pK_{Est(1)}$ ~9.8, $pK_{Est(2)}$ ~11.7. Crystd from pet ether (b 60-80°).

***meso*- α,β -Dimethylsuccinic acid** [608-40-2] M 146.1, m 211°, $pK_1^{2.5}$ 3.77, $pK^{2.5}$ 5.36. Crystd from EtOH/ether or EtOH/chloroform.

2,2-Dimethylsuccinic acid [597-43-3] M 146.1, m 141°, $pK_1^{2.0}$ 4.15, $pK_2^{2.0}$ 6.40. Crystd from EtOH/ether or EtOH/chloroform.

(\pm)-2,3-Dimethylsuccinic acid [13545-04-5] M 146.1, m 129°, $pK_1^{2.5}$ 3.82, $pK_2^{2.5}$ 5.98. Crystd from water.

Dimethyl sulfide [75-18-3] M 62.1, f -98.27°, b 0°/172mm, 37.5-38°/760mm, d^{21} 0.8458, n^{25} 1.4319. Purified *via* the Hg(II) chloride complex by dissolving 1 mole of Hg(II)Cl₂ in 1250mL of EtOH and slowly adding the boiling alcoholic soln of dimethyl sulfide to give the right ratio for 2(CH₃)₂S.3HgCl₂. After recrystn of the complex to constant melting point, 500g of complex is heated with 250mL conc HCl in 750mL of water. The sulfide is separated, washed with water, and dried with CaCl₂ and CaSO₄. Finally, it is distd under reduced pressure from sodium. Precautions should be taken (*efficient fume hood*) because of its very UNPLEASANT ODOUR.

2,4-Dimethylsulfolane [1003-78-7] M 148.2, b 128°/77mm, d^{25} 1.1314. Vacuum distd.

Dimethyl sulfone [67-71-0] M 94.1, m 109°. Crystd from water. Dried over P₂O₅.

Dimethyl sulfoxide (DMSO) [67-68-5] M 78.1, m 18.0-18.5°, b 75.6-75.8°/12mm, 190°/760mm, d 1.100, n 1.479. Colourless, odourless, very *hygroscopic* liquid, synthesised from dimethyl sulfide. The main impurity is water, with a trace of dimethyl sulfone. The Karl-Fischer test is applicable. It is dried with Linde types 4A or 13X molecular sieves, by prolonged contact and passage through a column of the material, then distd under reduced pressure. Other drying agents include CaH₂, CaO, BaO and CaSO₄. It can also be fractionally crystd by partial freezing. More extensive purification is achieved by standing overnight with freshly heated and cooled chromatographic grade alumina. It is then refluxed for 4h over CaO, dried over CaH₂, and then fractionally distd at low pressure. For efficiency of desiccants in drying dimethyl sulfoxide see Burfield and Smithers [*J Org Chem* 43 3966 1978; Sato et al. *J Chem Soc, Dalton Trans* 1949 1986].

Rapid purification: Stand over freshly activated alumina, BaO or CaSO₄ overnight. Filter and distil over CaH₂ under reduced pressure (~ 12 mm Hg). Store over 4A molecular sieves.

Dimethyl terephthalate [120-61-6] M 194.2, m 150°. Purified by zone melting.

***N,N*-Dimethylthiocarbamoyl chloride** [16420-13-6] M 123.6, m 42-43°, b 64-65°/0.1mm. Crystd twice from pentane.

***N,N*-Dimethyl-*o*-toluidine** [609-72-3] M 135.2, b 68°/10mm, 211-211.5°/760mm, d 0.937, n 1.53664, pK²⁵ 5.85. Isomers and other bases have been removed by heating in a water bath for 100h with two equivalents of 20% HCl and two and a half volumes of 40% aq formaldehyde, then making the soln alkaline and separating the free base. After washing well with water it was distd at 10mm pressure and redistd at ambient pressure [von Braun and Aust *Chem Ber* 47 260 1914]. Other procedures include drying with NaOH, distilling from zinc in an atmosphere of nitrogen under reduced pressure, and refluxing with excess of acetic anhydride in the presence of conc H₂SO₄ as catalyst, followed by fractional distn under vacuum.

***N,N*-Dimethyl-*m*-toluidine** [121-72-2] M 135.2, b 211.5-212.5°, d 0.93, pK²⁵ 5.22. See *m*-methyl-*N,N*-dimethylaniline on p. 291.

***N,N*-Dimethyl-*p*-toluidine** [99-97-8] M 135.2, b 93-94°/11mm, b 211°, d 0.937, n 1.5469, pK²⁵ 4.76. See *p*-methyl-*N,N*-dimethylaniline on p. 291.

1,3-Dimethyluracil [874-14-6] M 140.1, m 121-122°, pK -3.25 (aq H₂SO₄). Crystd from EtOH/ether.

***sym*-Dimethylurea** [96-31-1] M 88.1, m 106°. Crystd from acetone/diethyl ether by cooling in an ice bath. Also crystd from EtOH and dried at 50° and 5mm for 24h [Bloemendahl and Somsen *J Am Chem Soc* 107 3426 1985].

2,2'-Dinaphthylamine [532-18-3] M 269.3, m 170.5°, pK_{Est} <0. Crystd from *benzene.

2,4-Dinitroaniline [97-02-9] M 183.1, m 180°, ε₃₄₈ 12,300 in dil aq HClO₄, pK²⁵ -4.27 (aq H₂SO₄). Crystd from boiling EtOH by adding one-third volume of water and cooling slowly. Dried in a steam oven.

2,6-Dinitroaniline [606-22-4] M 183.1, m 139-140°, pK²⁵ -5.37 (aq H₂SO₄). Purified by chromatography on alumina, then crystd from *benzene or EtOH.

2,4-Dinitroanisole [5327-44-6] M 198.1, m 94-95°. Crystd from aq EtOH.

3,5-Dinitroanisole [119-27-7] M 198.1, m 105-106°. Purified by repeated crystn from water and dried in a vacuum desiccator over P₂O₅.

1,2-Dinitrobenzene [528-29-0] M 168.1, m 116.5°. Crystd from EtOH.

1,3-Dinitrobenzene [99-65-0] M 168.1, m 90.5-91°. Crystd from alkaline EtOH soln (20g in 750mL 95% EtOH at 40°, plus 100mL of 2M NaOH) by cooling and adding 2.5L of water. The ppte, after filtering off, washing with water and sucking dry, was crystd from 120mL, then 80mL of absolute EtOH [Callow, Callow and Emmens *Biochem J* 32 1312 1938]. Has also been crystd from MeOH, CCl₄ and ethyl acetate. Can be sublimed in a vacuum. [Tanner *J Org Chem* 52 2142 1987.]

1,4-Dinitrobenzene [100-25-4] M 168.1, m 173°. Crystd from EtOH or ethyl acetate. Dried under vacuum over P₂O₅. Can be sublimed in a vacuum.

2,4-Dinitrobenzenesulfonyl chloride [528-76-7] M 234.6, m 96°. Crystd from CCl₄.

2,4-Dinitrobenzenesulfonyl chloride [1656-44-6] M 266.6, m 102°. Crystd from *benzene or *benzene/pet ether.

2,4-Dinitrobenzoic acid [610-30-3] M 212.1, m 183°, pK²⁵ 1.42. Crystd from aqueous 20% EtOH (10mL/g), dried at 100°.

- 2,5-Dinitrobenzoic acid** [610-28-6] M 212.1, m 179.5-180°, pK²⁵ 1.62. Crystd from distd water. Dried in a vacuum desiccator.
- 2,6-Dinitrobenzoic acid** [603-12-3] M 212.1, m 202-203°, pK²⁵ 1.14. Crystd from water.
- 3,4-Dinitrobenzoic acid** [528-45-0] M 212.1, m 166°, pK²⁵ 2.81. Crystd from EtOH by addition of water.
- 3,5-Dinitrobenzoic acid** [99-34-3] M 212.1, m 205°, pK²⁵ 2.73 (2.79). Crystd from distilled water or 50% EtOH (4mL/g). Dried in a vacuum desiccator or at 70° over BaO under vacuum for 6h.
- 4,4'-Dinitrobenzoic anhydride** [902-47-6] M 406.2, m 189-190°. Crystd from acetone.
- 3,5-Dinitrobenzoyl chloride** [99-33-2] M 230.6, m 69.5°. Crystd from CCl₄ or pet ether (b 40-60°). It reacts readily with water, and should be kept in sealed tubes or under dry pet ether.
- 2,2'-Dinitrobiphenyl** [2436-96-6] M 244.2, m 123-124°. Crystd from EtOH.
- 2,4'-Dinitrobiphenyl** [606-81-5] M 244.2, m 92.7-93.7°. Crystd from EtOH.
- 4,4'-Dinitrobiphenyl** [1528-74-1] M 244.2, m 240.9-241.8°. Crystd from *benzene, EtOH (charcoal) or acetone. Dried under vacuum over P₂O₅.
- 2,6-Dinitro-*p*-cresol (2,6-dinitro-4-methylphenol)** [609-93-3] M 198.1, m 78-79°, pK_{Est}~3.7. Recrystd from EtOH and is steam volatile. **TOXIC IRRITANT.**
- 4,6-Dinitro-*o*-cresol** [534-52-1] M 198.1, m 85-86°, 87°, pK²⁵4.70. Crystd from aqueous EtOH.
- 2,4-Dinitrodiphenylamine** [961-68-2] M 259.2, m 157°, pK_{Est} <0. Crystd from aqueous EtOH.
- 4,4'-Dinitrodiphenylurea** [587-90-6] M 302.2, m 312°(dec). Crystd from EtOH. Sublimes in vac.
- 2,4-Dinitrofluorobenzene (Sanger's reagent)** [70-34-8] M 186.1, m 25-27°, b 133°/2mm, 140-141°/5mm, d 1.483. Crystd from ether or EtOH. Vacuum distd through a Todd Column (see p. 174). If it is to be purified by distn *in vacuo*, the distn unit must be allowed to cool before air is allowed into the apparatus otherwise the residue carbonises spontaneously and an **EXPLOSION** may occur. The material is a **skin irritant** and may cause serious dermatitis.
- 1,8-Dinitronaphthalene** [602-38-0] M 218.2, m 170-171°. Crystd from *benzene.
- 2,4-Dinitro-1-naphthol (Martius Yellow)** [605-69-6] M 234.2, m 81-82°, pK_{Est} ~3.7. Crystd from *benzene or aqueous EtOH.
- 2,4-Dinitrophenetole** [610-54-8] M 240.2, m 85-86°. Crystd from aqueous EtOH.
- 2,4-Dinitrophenol** [51-28-5] M 184.1, m 114°, pK²⁵ 4.12. Crystd from *benzene, EtOH, EtOH/water or water acidified with dil HCl, then recrystd from CCl₄. Dried in an oven and stored in a vac desiccator over CaSO₄.
- 2,5-Dinitrophenol** [329-71-5] M 184.1, m 108°, pK²⁵ 5.20. Crystd from H₂O with a little EtOH.
- 2,6-Dinitrophenol** [573-56-8] M 184.1, m 63.0-63.7°, pK²⁵ 3.73. Crystd from *benzene/cyclohexane, aqueous EtOH, water or *benzene/pet ether (b 60-80°, 1:1).

3,4-Dinitrophenol [577-71-9] M 184.1, m 138°, pK²⁵ 5.42. Steam distd and crystd from water and air-dried. **CAUTION - EXPLOSIVE** when dry, store with 10% water.

3,5-Dinitrophenol [586-11-8] M 184.1, m 126°, pK²⁵ 6.68. Crystd from *benzene or CHCl₃/pet ether. Should be stored with 10% water because it is **EXPLOSIVE** when dry.

2,4-Dinitrophenylacetic acid [643-43-6] M 226.2, m 179°(dec), pK²⁵ 3.50. Crystd from water.

2,4-Dinitrophenylhydrazine (DNPH) [119-26-6] M 198.1, m 200°(dec), pK_{Est} ~2.0. Crystd from butan-1-ol, dioxane, EtOH, *C₆H₆ or ethyl acetate. HCl has m 186°(dec).

2,2-Dinitropropane [595-49-3] M 162.1, m 53.5°. Crystd from EtOH or MeOH. Dried over CaCl₂ or under vacuum for 1h just above the melting point.

2,4-Dinitroresorcinol [519-44-8] M 200.1, m 149°, pK²⁵ 3.05. Crystd from aq EtOH. **Explosive.**

3,5-Dinitrosalicylic acid [609-99-4] M 228.1, m 173-174°, pK₁²⁵ 0.70, pK₂²⁵ 7.40. Crystd from water.

2,6-Dinitrothymol [303-21-9] M 240.2, m 53-54°. Crystd from aq EtOH.

2,3-Dinitrotoluene [602-01-7] M 182.1, m 63°. Distd in steam and crystd from water or *benzene/pet ether. Stored with 10% water. *Could be EXPLOSIVE when dry.*

2,4-Dinitrotoluene [121-14-2] M 182.1, m 70.5-71.0°. Crystd from acetone, isopropanol or MeOH. Dried under vacuum over H₂SO₄. Purified by zone melting. *Could be EXPLOSIVE when dry.*

2,5-Dinitrotoluene [619-15-8] M 182.1, m 51.2°. Crystd from *benzene.

2,6-Dinitrotoluene [606-20-2] M 182.1, m 64.3°. Crystd from acetone.

3,4-Dinitrotoluene [610-39-9] M 182.1, m 61°. Distil in steam and cryst from *benzene/pet ether. Store with 10% of water to avoid **EXPLOSION**.

3,5-Dinitro-*o*-toluic acid [28169-46-2] M 226.2, m 206°, pK_{Est} ~3.0. Crystd from H₂O or aq EtOH.

2,4-Dinitro-*m*-xylene [603-02-1] M 196.2, m 83-84°. Crystd from EtOH.

Dinonyl phthalate (mainly 3,5,5-trimethylhexyl phthalate isomer) [28553-12-0; 14103-61-8] M 418.6, m 26-29°, b 170°/2mm, d 0.9640, n 1.4825. Washed with aqueous Na₂CO₃, then shaken with water. Ether was added to break the emulsion, and the soln was washed twice with water, and dried (CaCl₂). After evaporating the ether, the residual liquid was distd three times under reduced pressure. It was stored in a vacuum desiccator over P₂O₅.

Diocetyltrimethylammonium bromide [3700-67-2] M 630.9, m 161-163°. Crystd from acetone then MeOH [Lukac *J Am Chem Soc* 106 4387 1984]. Also purified by chromatography on alumina by washing with *C₆H₆ and eluting with Me₂CO, evap and cryst from MeCN [Swain and Kreevoy *J Am Chem Soc* 77 1126 1955].

Diosgenin [512-04-9] M 294.5, m 204-207°, [α]_D²⁵ -129° (in Me₂CO). Crystd from acetone.

1,3-Dioxane [505-22-6] M 88.1, b 104.5°/751mm, d 1.040, n 1.417. Dried with sodium and fractionally distd.

1,4-Dioxane [123-91-1] **M 88.1, f 11.8°, b 101.3°, d²⁵ 1.0292, n¹⁵ 1.4236, n²⁵ 1.42025.**

Prepared commercially either by dehydration of ethylene glycol with H₂SO₄ and heating ethylene oxide or bis(β-chloroethyl)ether with NaOH. Usual impurities are acetaldehyde, ethylene acetal, acetic acid, water and peroxides. Peroxides can be removed (and the aldehyde content decreased) by percolation through a column of activated alumina (80g per 100-200mL solvent), by refluxing with NaBH₄ or anhydrous stannous chloride and distilling, or by acidification with conc HCl, shaking with ferrous sulfate and leaving in contact with it for 24h before filtering and purifying further.

Hess and Frahm [*Chem Ber* 71 2627 1938] refluxed 2L of dioxane with 27mL conc HCl and 200mL water for 12h with slow passage of nitrogen to remove acetaldehyde. After cooling the soln KOH pellets were added slowly and with shaking until no more would dissolve and a second layer had separated. The dioxane was decanted, treated with fresh KOH pellets to remove any aq phase, then transferred to a clean flask where it was refluxed for 6-12h with sodium, then distd from it. Alternatively, Kraus and Vingee [*J Am Chem Soc* 56 511 1934] heated on a steam bath with solid KOH until fresh addition of KOH gave no more resin (due to acetaldehyde). After filtering through paper, the dioxane was refluxed over sodium until the surface of the metal was not further discoloured during several hours. It was then distd from sodium.

The acetal (b 82.5°) is removed during fractional distn. Traces of *benzene, if present, can be removed as the *benzene/MeOH azeotrope by distn in the presence of MeOH. Distn from LiAlH₄ removes aldehydes, peroxides and water. Dioxane can be dried using Linde type 4X molecular sieves. Other purification procedures include distn from excess C₂H₅MgBr, refluxing with PbO₂ to remove peroxides, fractional crystn by partial freezing and the addition of KI to dioxane acidified with aq HCl. Dioxane should be stored out of contact with air, preferably under N₂.

A detailed purification procedure is as follows: Dioxane was stood over ferrous sulfate for at least 2 days, under nitrogen. Then water (100mL) and conc HCl (14mL) / litre of dioxane were added (giving a pale yellow colour). After refluxing for 8-12h with vigorous N₂ bubbling, pellets of KOH were added to the warm soln to form two layers and to discharge the colour. The soln was cooled rapidly with more KOH pellets being added (magnetic stirring) until no more dissolved in the cooled soln. After 4-12h, if the lower phase was not black, the upper phase was decanted rapidly into a clean flask containing sodium, and refluxed over sodium (until freshly added sodium remained bright) for 1h. The middle fraction was collected (and checked for minimum absorbency below 250nm). The distillate was fractionally frozen three times by cooling in a refrigerator, with occasional shaking or stirring. This material was stored in a refrigerator. For use it was thawed, refluxed over sodium for 48h, and distilled into a container for use. All joints were clad with Teflon tape.

Coetzee and Chang [*Pure Appl Chem* 57 633 1985] dried the solvent by passing it slowly through a column (20g/L) of 3A molecular sieves activated by heating at 250° for 24h. Impurities (including peroxides) were removed by passing the effluent slowly through a column packed with type NaX zeolite (pellets ground to 0.1mm size) activated by heating at 400° for 24h or chromatographic grade basic Al₂O₃ activated by heating at 250° for 24h. After removal of peroxides the effluent was refluxed several hours over sodium wire, excluding moisture, distilled under nitrogen or argon and stored in the dark.

One of the best tests of purity of dioxane is the formation of the purple disodium benzophenone complex during reflux and its persistence on cooling. (Benzophenone is better than fluorenone for this purpose, and for the storing of the solvent.) [Carter, McClelland and Warhurst *Trans Faraday Soc* 56 343 1960]. **TOXIC.**

Rapid purification: Check for peroxides (see Chapter 1 and Chapter 2 for test under ethers). Pre-dry with CaCl₂ or better over Na wire. Then reflux the pre-dried solvent over Na (1% w/v) and benzophenone (0.2% w/v) under an inert atmosphere until the blue colour of the benzophenone ketyl radical anion persists. Distil, and store over 4A molecular sieves in the dark.

1,3-Dioxolane [646-06-0] **M 74.1, b 75.0-75.2°, d 1.0600, n²¹ 1.3997.** Dried with solid NaOH, KOH or CaSO₄, and distd from sodium or sodium amalgam. Barker et al. [*J Chem Soc* 802 1959] heated 34mL of dioxolane under reflux with 3g of PbO₂ for 2h, then cooled and filtered. After adding xylene (40mL) and PbO₂ (2g) to the filtrate, the mixture was fractionally distd. Addition of xylene (20mL) and sodium wire to the main fraction (b 70-71°) led to vigorous reaction, following which the mixture was again fractionally distd. Xylene and sodium additions were made to the main fraction (b 73-74°) before it was finally distd.

4,4'-Di-*n*-pentyloxyazoxybenzene [64242-26-8] **M 370.5.** Crystd from Me₂CO, and dried by heating under vacuum.

Diphenic acid [482-05-3] M 242.2, m 228-229°, pK²⁵ 3.46. Crystd from water.

Diphenic anhydride [6050-13-1] M 466.3, m 217°. After removing free acid by extraction with cold aq Na₂CO₃, the residue has been crystd from acetic anhydride and dried at 100°. Acetic anhydride also converts the acid to the anhydride.

***N,N*-Diphenylacetamide** [621-09-0] M 210.3, m 131°. Crystd from EtOH, then sublimed under vacuum at ca 96° onto a "finger" cooled in solid CO₂/MeOH, with continuous pumping to free it from occluded solvent.

Diphenylacetic acid [117-34-0] M 212.3, m 147.4-148.4°, pK²⁵ 3.94. Crystd from *benzene, H₂O or aq 50% EtOH.

Diphenylacetonitrile [86-29-3] M 193.3, m 73-75°. Crystd from EtOH or pet ether (b 90-100°).

Diphenylacetylene (tolan) [501-65-5] M 178.2, m 62.5°, b 90-97°/0.3mm. Crystd from EtOH.

Diphenylamine [122-39-4] M 169.2, m 62.0-62.5°, pK²⁵ 0.77 (aq H₂SO₄). Crystd from pet ether, MeOH, or EtOH/water. Dried under vacuum.

Diphenylamine-2-carboxylic acid [91-40-7] M 213.2, m 184°, pK₁²⁵ -1.28 (aq H₂SO₄), pK₂²⁵ 3.86. See *N*-phenylanthranilic acid on p. 327.

Diphenylamine-2,2'-dicarboxylic acid (2,2'-iminodibenzoic acid) [579-92-0] M 257.2, m 298°(dec), pK_{Est} ~3.7. Crystd from EtOH.

9,10-Diphenylanthracene [1499-10-1] M 330.4, m 248-249°. Crystd from acetic acid or xylene [Baumstark et al. *J Org Chem* 52 3308 1987].

***N,N'*-Diphenylbenzidine** [531-91-9] M 336.4, m 245-247°, 251-252°, pK²⁵ 0.30. Crystd from toluene or ethyl acetate. Stored in the dark.

***trans-trans*-1,4-Diphenylbuta-1,3-diene** [538-81-8] M 206.3, m 153-153.5°. Its soln in pet ether (b 60-70°) was chromatographed on an alumina-Celite column (4:1) and the column was washed with the same solvent. The main zone was cut out, eluted with ethanol and transferred to pet ether, which was then dried and evaporated [Pinckard, Wille and Zechmesiter *J Am Chem Soc* 70 1938 1948]. Recrystd from hexane.

***sym*-Diphenylcarbazine** [140-22-7] M 242.3, m 172°. A common impurity is phenylsemicarbazide which can be removed by chromatography [Willems et al. *Anal Chim Acta* 51 544 1970]. Crystd from EtOH or glacial acetic acid.

1,5-Diphenylcarbazon [538-62-5] M 240.3, m 124-127°. Crystd from EtOH (ca 5mL/g), and dried at 50°. A commercial sample, nominally *sym*-diphenylcarbazon but of m 154-156°, was a mixture of diphenylcarbazine and diphenylcarbazon. The former was removed by dissolving 5g of the crude material in 75mL of warm EtOH, then adding 25g Na₂CO₃ dissolved in 400mL of distd water. The alkaline soln was cooled and extracted six times with 50mL portions of diethyl ether (discarded). Diphenylcarbazon was then pptd by acidifying the alkaline soln with 3M HNO₃ or glacial acetic acid. It was filtered on a Büchner funnel, air dried, and stored in the dark [Gerlach and Frazier *Anal Chem* 30 1142 1958]. Other impurities were phenylsemicarbazide and diphenylcarbodiazone. Impurities can be detected by chromatography [Willems et al. *Anal Chim Acta* 51 544 1970].

Diphenyl carbonate [102-09-0] M 214.2, m 80°. Purified by sublimation, and by preparative gas chromatography with 20% Apiezon on Embacel, and crystn from EtOH.

Diphenylcyclopropenone (Diphencyprone) [886-38-4] M 206.2, m 87-90°(hydrate), 119-121°(anhydr). Crystd from cyclohexane. UV (MeCN): λ_{\max} 226, 282, 297nm.

Diphenyl disulfide (phenyl disulfide) [882-33-7] M 218.3, m 60.5°. Crystd from MeOH. [Alberti et al. *J Am Chem Soc* 108 3024 1986]. Crystd repeatedly from hot diethyl ether, then vac dried at 30° over P₂O₅, fused under nitrogen and re-dried, the whole procedure being repeated, with a final drying under vac for 24h. Also recrystd from hexane/EtOH soln. [Burkey and Griller *J Am Chem Soc* 107 246 1985.]

1,1-Diphenylethanol [599-67-7] M 198.3, m 80-81°. Crystd from *n*-heptane. [Bromberg et al. *J Am Chem Soc* 107 83 1985.]

1,1-Diphenylethylene [530-48-3] M 180.3, b 268-270°, d 1.024, n 1.6088. Distd under reduced pressure from KOH. Dried with CaH₂ and redistd.

***N,N'*-Diphenylethylenediamine (Wanzlick's reagent)** [150-61-8] M 212.3, m 67.5°, b 178-182°/2mm $pK_{\text{Est}(1)} \sim 0.5$, $pK_{\text{Est}(2)} \sim 3.8$. Crystd from aqueous EtOH or MeOH.

***N,N'*-Diphenylformamidine** [622-15-1] M 196.2, m 142°, 137°, 136-139°. Crystd from absolute EtOH, gives the hydrate with aqueous EtOH.

1,3-Diphenylguanidine [102-06-7] M 211.3, m 148°, pK^{25} 10.12. Crystd from toluene, aqueous acetone or EtOH, and vacuum dried.

1,6-Diphenyl-1,3,5-hexatriene [1720-32-7] M 232.3, m 200-203°. Crystd from CHCl₃ or EtOH/CHCl₃ (1:1).

5,5-Diphenylhydantoin [57-41-0] M 252.3, m 293-295°. Crystd from EtOH.

1,1-Diphenylhydrazine (hydrazobenzene) [122-66-7] M 184.2, m 34°, 44°, 175°/10mm, 222°/40mm, $pK_{\text{Est}} \sim 1.7$. Crystd from hot EtOH containing a little ammonium sulfide or H₂SO₃ (to prevent atmospheric oxidation), preferably under nitrogen. Dried in a vacuum desiccator. Also crystd from pet ether (b 60-100°) to constant absorption spectrum. *HCl*, from EtOH has m 163-164°(dec). *Picrate*, from *C₆H₆, has m 123°(dec).

1,3-Diphenylisobenzofuran [5471-63-6] M 270.3, m 129-130°. Recrystd from EtOH or EtOH/CHCl₃ (1:1) under red light (as in photographic dark rooms) or from *benzene in the dark.

Diphenylmethane [101-81-5] M 168.2, m 25.4°. Sublimed under vacuum, or distd at 72-75°/4mm. Crystd from EtOH. Purified by fractional crystn of the melt.

1,1-Diphenylmethanamine [91-00-9] M 183.2, m 34°, $pK_{\text{Est}} \sim 9.1$. Crystd from water.

Diphenylmethyl chloride (benzhydryl chloride) [90-99-3] M 202.7, m 17.0°, b 167°/17mm, n 1.5960. Dried with Na₂SO₄ and fractionally distd under reduced pressure.

***all-trans*-1,8-Diphenyl-1,3,5,8-octatetraene** [3029-40-1] M 258.4, m 235-237°. Crystd from EtOH.

2,5-Diphenyl-1,3,4-oxadiazole (PPD) [725-12-2] M 222.3, m 70° (hydrate), 139-140° (anhydrous), b 231°/13mm, 248°/16mm. Crystd from EtOH and sublimed *in vacuo*.

2,5-Diphenyloxazole (PPO) [92-71-7] M 221.3, m 74°, b 360°/760mm. Distd in steam and crystd from ligroin.

***N,N'*-Diphenyl-*p*-phenylenediamine** [74-31-7] M 260.3, m 148-149°, b 219-224°/0.7mm, pK_{Est} <0. Crystd from EtOH, chlorobenzene/pet ether or *benzene. Has also been crystd from aniline, then extracted three times with absolute EtOH.

1,1-Diphenyl-2-picrylhydrazine [1707-75-1] M 395.3, m 174°(dec), 178-179.5°(dec). Crystd from CHCl₃, or *benzene/pet ether (1:1), then degassed at 100° and <10⁻⁵mm Hg for ca 50h to decompose the 1:1 molar complex formed with *benzene.

2,2-Diphenylpropionic acid [5558-66-7] M 226.3, m 173-174°, pK_{Est} ~3.8. Crystd from EtOH.

3,3-Diphenylpropionic acid [606-83-7] M 226.3, m 155°, pK_{Est} ~4.5. Crystd from EtOH.

Diphenyl sulfide [139-66-2] M 186.3, b 145°/8mm, d 1.114, n 1.633. Washed with aqueous 5% NaOH, then water. Dried with CaCl₂, then with sodium. The sodium was filtered off and the diphenyl sulfide was distd under reduced pressure.

Diphenyl sulfone [127-63-9] M 218.3, m 125°, b 378°(dec). Crystd from diethyl ether. Purified by zone melting.

***sym*-Diphenylthiourea (thiocarbanilide)** [102-08-9] M 228.3, m 154°. Crystd from boiling EtOH by adding hot water and allowing to cool.

1,1-Diphenylurea [603-54-3] M 212.3, m 238-239°. Crystd from MeOH.

Dipicolinic acid (pyridine-2,6-dicarboxylic acid) [499-83-2] M 167.1, m 255°(dec), λ_{max} 270nm, pK₁²⁰ 2.10, pK₂²⁰ 4.68. Crystd from water, and sublimed in a vacuum.

***N,N*-Di-*n*-propylaniline** [2217-07-4] M 177.3, b 127°/10mm, 238-241°/760mm, pK²³ 5.68. Refluxed for 3hr with acetic anhydride, then fractionally distd under reduced pressure.

Dipropylene glycol (octan-4,5-diol) [110-98-5] M 134.2, b 109-110°/8mm, d 1.022, n 1.441. Fractionally distd below 15mm pressure, using packed column and taking precautions to avoid absorption of water.

***Di-n*-propyl ketone** [123-19-3] M 114.2, b 143.5°, d 0.8143, n 1.40732. Dried with CaSO₄, then distd from P₂O₅ under nitrogen.

***Di-n*-propyl sulfide** [111-47-7] M 118.2, b 141-142°, d 0.870, n 1.449. Washed with aqueous 5% NaOH, then water. Dried with CaCl₂ and distd from Na [Dunstan and Griffiths *J Chem Soc* 1344 1962].

Di-(4-pyridoyl)hydrazine [4329-75-3] M 246.2, m 254-255°. Crystd from water.

2,2'-Dipyridylamine [1202-34-2] M 171.2, m 84° and remelts at 95° after solidifying, b 176-178°/13mm, 307-308°/760mm, pK 6.69 (in 20% EtOH). Crystd from *benzene or toluene [Blakley and De Armond *J Am Chem Soc* 109 4895 1987].

2,2'-Dipyridyl disulfide (2,2'dithiopyridine) [2127-03-9] M 220.3, m 53°, 56-58°, 57-58°, pK₁²⁵ 0.35, pK₂²⁵ 2.45. Recrystd from *C₆H₆/pet ether (6:7), ligroin or *C₆H₆. *Picrate* has m 119° (from EtOH). [Walter et al. *Justus Liebigs Ann Chem* 695 7785 1966; Marckwald et al. *Chem Ber* 33 1556 1900; Brocklehurst and Little *Biochem J* 133 67,78 1973.]

1,2-Di-(4-pyridyl)-ethane [4916-57-8] M 184.2, pK_{Est(1)} ~3.8, pK_{Est(2)} ~5.4. Crystd from cyclohexane/*benzene (5:1).

trans-1,2-Di-(4-pyridyl)-ethylene [13362-78-2] M 182.2, m 153-154°, 155.5-156.5°, pK₁²⁵ 3.65, pK₂⁵ 5.6. Crystd from water (1.6g/100mL at 100°). *Di-HCl* has m 347°, from EtOH

1,3-Di-(4-pyridyl)-propane [17252-51-6] M 198.3, m 60.5-61.5°, pK_{Est(1)} ~4.5, pK_{Est(2)} ~5.5. Crystd from *n*-hexane/*benzene (5:1).

S-1,2-Distearin [1188-58-5] M 625.0, m 76-77°, [α]_D²⁰ -2.8° (c 6.3, CHCl₃), [α]₅₄₆²⁰ +1.4° (c 10, CHCl₃/MeOH, 9:1). Crystd from chloroform/pet ether.

2,5-Distyrylpyrazine [14990-02-4] M 284.3, m 219°. Recrystd from xylene; chromatographed on basic silica gel (60-80 mesh) using CH₂Cl₂ as eluent, then vac sublimed on to a cold surface at 10⁻³ torr [Ebied *J Chem Soc, Faraday Trans 1* 78 3213 1982]. Operations should be carried out in the dark.

1,3-Dithiane [505-23-7] M 120.2, m 54°. Crystd from 1.5 times its weight of MeOH at 0°, and sublimed at 40-50°/0.1mm.

2,2'-Dithiobis(benzothiazole) [120-78-8] M 332.2, m 180°. Crystd from *benzene.

4,4'-Dithiodimorpholine [103-34-4] M 236.2, m 124-125°. Crystd from hot aq dimethylformamide.

1,4-Dithioerythritol (DTE, erythro-2,3-dihydroxy-1,4-dithiobutane) [6892-68-8] M 154.3, m 82-84°, pK₁ 9.0, pK₂ 9.9. Crystd from ether/hexane and stored in the dark at 0°.

Dithiooxamide (rubeanic acid) [79-40-3] M 120.2, m >300°. Crystd from EtOH and sublimed in a vacuum.

RS-1,4-Dithiothreitol (DTT, Cleland's reagent) [27565-41-9] M 154.3, m 42-43°, pK₁ 8.3, pK₂ 9.5. Crystd from ether and sublimed at 37°/0.005mm. Should be stored at 0°.

Dithizone (diphenylthiocarbazone) [60-10-6] M 256.3, ratio of ε_{620nm}/ε_{450nm} should be ≥1.65, ε₆₂₀ 3.4 x 10⁴ (CHCl₃), pK₂ 4.6. The crude material is dissolved in CCl₄ to give a concentrated soln. This is filtered through a sintered glass funnel and shaken with 0.8M aq ammonia to extract dithizonate ion. The aqueous layer is washed with several portions of CCl₄ to remove undesirable materials. The aqueous layer is acidified with dil H₂SO₄ to precipitate pure dithizone. It is dried in a vacuum. When only small amounts of dithizone are required, purification by paper chromatography is convenient. [Cooper and Hibbits *J Am Chem Soc* 75 5084 1933.] Instead of CCl₄, CHCl₃ can be used, and the final extract, after washing with water, can be evapd in air at 40-50° and dried in a desiccator. Complexes with Cd, Hg, Ni and Zn.

Di-*p*-tolyl carbonate [621-02-3] M 242.3, m 115°. Purified by GLC with 20% Apiezon on Embacel followed by sublimation *in vacuo*.

***N,N'*-Di-*o*-tolylguanidine** [97-39-2] M 239.3, m 179° (175-176°), pK_{Est} ~10.3. Crystd from aqueous EtOH.

Di-*p*-tolylphenylamine [20440-95-3] M 273.4, m 108.5°, pK_{Est} ~5.0. Crystd from EtOH.

Di-*p*-tolyl sulfone [599-66-6] M 278.3, m 158-159°, b 405°. Crystd repeatedly from diethyl ether. Purified by zone melting.

Djenkolic acid (S,S'-methylene-bis-L-cysteine) [498-59-9] M 254.3, m 300-350°(dec), [α]_D²⁰ -65° (c 2, HCl) [See pK of S-methyl-L-cysteine]. Crystd from a large volume of water (sol 0.5g%).

***cis*-4,7,10,13,16,19-Docosahexaenoic acid** [6217-54-5] M 328.5, m -44/1°, -44.1°, n_D²⁰ 1.5017, pK_{Est} ~4.6. Its solubility in CHCl₃ is 5%. It has been purified from fish oil by GLC using Ar as mobile phase and EGA as stationary phase with an ionisation detector [UV: Stoffel and Ahrens *J Lipid Res* 1

139 1959], and *via* the ester by evaporative "molecular" distillation using a 'continuous molecular still' at 10^{-4} mm with the highest temperature being 110° , and a total contact time with the hot surface being 60sec [Farmer and van den Heuvel *J Chem Soc* 427 1938]. The *methyl ester* has **b** $208-211^{\circ}/2\text{mm}$, d_4^{20} 0.9398, n_D^{20} 1.5035. With Br_2 it forms a *dodecabromide* **m ca** 240° dec. Also the acid was converted to the methyl ester and purified through a three stage molecular still [as described by Sutton *Chem Ind (London)* 11383 1953] at 96° with the rate adjusted so that one third of the material was removed each cycle of three distillations. The distillate (numbered 4) (13g) was dissolved in EtOH (100mL containing 8g of KOH) at -70° and set aside for 4h at 30° with occasional shaking under a vac. Water (100mL) is added and the soln is extracted with pentane, washed with HCl, dried (MgSO_4), filtered and evapd to give a clear oil (11.5g) **m** -44.5° to -44.1° . In the catalytic hydrogenation of the oil six mols of H_2 were absorbed and *docosanoic acid* (*behenic acid*) was produced with **m** $79.0-79.3^{\circ}$ undepressed with an authentic sample (see docosanoic acid below) [Whitcutt *Biochem J* 67 60 1957].

Docosane [629-97-0] **M** 310.6, **m** 47° , **b** $224^{\circ}/15\text{mm}$. Crystd from EtOH or ether.

Docosanoic acid (*behenic acid*) [112-85-6] **M** 340.6, **m** $81-82^{\circ}$, pK_{Est} ~ 4.9 . Crystd from ligroin. [Francis and Piper *J Am Chem Soc* 61 577 1939].

1-Docosanol (*behenyl alcohol*) [661-19-8] **M** 182.3, **m** 70.8° . Crystd from ether or chloroform/ether.

***n*-Dodecane** [112-40-3] **M** 170.3, **b** $97.5-99.5^{\circ}/5\text{mm}$, $216^{\circ}/760\text{mm}$, **d** 0.748, **n** 1.42156. Passed through a column of Linde type 13X molecular sieves. Stored in contact with, and distd from, sodium. Passed through a column of activated silica gel. Has been crystd from diethyl ether at -60° . Unsaturated dry material which remained after passage through silica gel has been removed by catalytic hydrogenation (Pt_2O) at $45\text{lb}/\text{in}^2$ (3.06 atmospheres), followed by fractional distn under reduced pressure [Zook and Goldey *J Am Chem Soc* 75 3975 1953]. Also purified by partial crystn from the melt.

Dodecane-1,10-dioic acid (*decane-1,10-dicarboxylic acid*) [693-23-2] **M** 230.3, **m** 129° , **b** $245^{\circ}/10\text{mm}$, pK_{Est} ~ 4.8 . Crystd from water, 75% or 95% EtOH (sol 10%), or glacial acetic acid.

1-Dodecanol (*dodecyl alcohol*) [112-53-8] **M** 186.3, **m** 24° , **b** $91^{\circ}/1\text{mm}$, $135^{\circ}/10\text{mm}$, $167^{\circ}/40\text{mm}$, $213^{\circ}/200\text{mm}$, $259^{\circ}/\text{atm}$, d_4^{24} 0.8309 (liquid). Crystd from aqueous EtOH, and vacuum distd in a spinning-band column. [Ford and Marvel *Org Synth* 10 62 1930.]

1-Dodecanthiol [112-55-0] **M** 202.4, **b** $111-112^{\circ}/3\text{mm}$, $153-155^{\circ}/24\text{mm}$, **d** 0.844, **n** 1.458, pK_{Est} ~ 10.8 . Dried with CaO for several days, then distd from CaO.

Dodecylammonium butyrate [17615-97-3] **M** 273.4, **m** $39-40^{\circ}$, pK^{25} 10.63 (for free base). Recrystd from *n*-hexane.

Dodecylammonium propionate [17448-65-6] **M** 259.4, **m** $55-56^{\circ}$. Recrystd from hexanol/pet ether (b $60-80^{\circ}$).

Dodecyldimethylamine oxide [1643-20-5] **M** 229.4, **m** 102° . Crystd from acetone or ethyl acetate. [Bunton et al. *J Org Chem* 52 3832 1987].

Dodecyl ether [4542-57-8] **M** 354.6, **m** 33° . Vacuum distd, then crystd from MeOH/*benzene.

1-Dodecylpyridinium chloride [104-74-5] **M** 301.9, **m** $68-70^{\circ}$. Purified by repeated crystn from acetone (charcoal); twice recrystd from EtOH [Chu and Thomas *J Am Chem Soc* 108 6270 1986].

Dodecyltrimethylammonium bromide [1119-94-4] **M** 308.4, **m** 246° (dec). Purified by repeated crystn from acetone. Washed with diethyl ether and dried in a vacuum oven at 60° [Dearden and Wooley *J Phys Chem* 91 2404 1987].

Dodecyltrimethylammonium chloride [112-00-5] M 263.9, m 246°(dec). Dissolved in MeOH, treated with active charcoal, filtered and dried *in vacuo* [Waldenburg *J Phys Chem* 88 1655 1984], or recrystd several times from 10% EtOH in acetone. Also repeatedly crystd from EtOH/ether or MeOH. [Cella et al. *J Am Chem Soc* 74 2062 1952.]

Dulcitol [608-66-2] M 182.2, m 188-189°, b 276-280°/1.1mm. Crystd from water by addition of EtOH.

Duroquinone (tetramethylbenzoquinone) [527-17-3] M 164.2, m 110-111°. Crystd from 95% EtOH. Dried under vacuum.

α -Ecdyson [3604-87-3] M 464.7, m 239-242°, 242°, $[\alpha]_D^{20} +72^\circ$ (c 1, EtOH). Recrystd from tetrahydrofuran-pet ether, and from H₂O as a hydrate. It has been purified by chromatography on Al₂O₃ and elution with EtOAc-MeOH. It has λ_{\max} at 242nm (ϵ 12.400). Its *acetate* has m 214-216° from EtOAc-pet ether, and the *2,4-dinitrophenylhydrazone* has m 170-175° (dec) from EtOAc. [Karlson and Hoffmeister *Justus Liebigs Ann Chem* 662 1 1963; Karlson *Pure Appl Chem* 14 75 1967.]

β -Ecdyson (β -echdysterone) [5289-74-7] M 480.7, m 245-247°, $[\alpha]_D^{20} +66^\circ$ (c 1, MeOH). Crystd from water or tetrahydrofuran/pet ether.

Echinenone [432-68-8] M 550.8, m 178-179°, $A_{1\text{cm}}^{1\%}(\lambda_{\max})$ 2160 (458nm) in pet ether. Purified by chromatography on partially deactivated alumina or magnesia, or by using a thin layer of silica gel G with 4:1 cyclohexane/diethyl ether as the developing solvent. Stored in the dark at -20°.

Eicosane [112-95-8] M 282.6, m 36-37°, b 205°/15mm, $d^{36.7}$ 0.7779, n^{40} 1.43453. Crystd from EtOH.

Elaidic (*trans*-oleic) acid [112-79-8] M 282.5, m 44.5°, pK^{25} 4.9. Crystd from acetic acid, then EtOH.

Ellagic acid (2H₂O) [476-66-4] M 338.2, m >360°, $pK_{\text{Est}(1)} \sim 8$, $pK_{\text{Est}(2)} \sim 11$. Crystd from pyridine.

Elymoclavine (8,9-didehydro-6-methylergoline-8-methanol) [548-43-6] M 254.3, m 249-253°(dec), $[\alpha]_D^{20} -59^\circ$ (c 1, EtOH). Crystd from MeOH.

Embonic acid (Pamoic acid, 4,4'-methylene bis[3-hydroxy-2-naphthalenecarboxylic acid]) [130-85-8] M 388.4, m 295°, >300°, $pK_{\text{Est}(1)} \sim 2.2$, $pK_{\text{Est}(2)} \sim 13.2$. Forms crystals from dilute pyridine which decomposition above 280° without melting. It is almost insoluble in H₂O, EtOH, Et₂O, *C₆H₆, CH₃CO₂H, sparingly soluble in CHCl₃ but soluble in nitrobenzene, pyridine and alkalis [Barber and Gaimster *J Appl Chem (London)* 2 565 1952]. Used for making salts of organic bases.

Emetine hydrochloride hydrate [316-42-7] M 553.6 + aq, m 235-240°, 235-250°, 240-250°, 248-250° (depending on H₂O content), $[\alpha]_D^{20} -49.2^\circ$ (free base, c 4, CHCl₃), +18° (c 6, H₂O, dry salt), pK_1 5.77, pK_2 6.64. It crystallises from MeOH-Et₂O, MeOH or Et₂O-EtOAc. The *free base* has m 104-105°, and the (*-*)-*phenyl thiourea derivative* has m 220-221° [from EtOAc-pet ether, $[\alpha]_D^{25} -29.3^\circ$ (CHCl₃)]. IR: 3413 (OH) and 2611 (NH⁺) cm⁻¹; UV λ_{\max} 230nm (ϵ 16 200) and 282nm (ϵ 6 890) [Brossi et al. *Helv Chim Acta* 42 1515 1959; Barash et al. *J Chem Soc* 3530 1959].

Emodine (1,3,8-trihydroxy-6-methyl-9,10-anthracenedione, archin) [518-82-1] M 270.2, m 253-257°, 255-256°, 256-257°, 262°, 264° (phenolic pKs 7-10). Forms orange needles from EtOH, Et₂O, *C₆H₆, toluene or pyridine. It sublimes above 200° at 12mm. [Tutin and Clewer *J Chem*

Dodecyltrimethylammonium chloride [112-00-5] M 263.9, m 246°(dec). Dissolved in MeOH, treated with active charcoal, filtered and dried *in vacuo* [Waldenburg *J Phys Chem* 88 1655 1984], or recrystd several times from 10% EtOH in acetone. Also repeatedly crystd from EtOH/ether or MeOH. [Cella et al. *J Am Chem Soc* 74 2062 1952.]

Dulcitol [608-66-2] M 182.2, m 188-189°, b 276-280°/1.1mm. Crystd from water by addition of EtOH.

Duroquinone (tetramethylbenzoquinone) [527-17-3] M 164.2, m 110-111°. Crystd from 95% EtOH. Dried under vacuum.

α -Ecdyson [3604-87-3] M 464.7, m 239-242°, 242°, $[\alpha]_D^{20} +72^\circ$ (c 1, EtOH). Recrystd from tetrahydrofuran-pet ether, and from H₂O as a hydrate. It has been purified by chromatography on Al₂O₃ and elution with EtOAc-MeOH. It has λ_{\max} at 242nm (ϵ 12.400). Its *acetate* has m 214-216° from EtOAc-pet ether, and the *2,4-dinitrophenylhydrazone* has m 170-175° (dec) from EtOAc. [Karlson and Hoffmeister *Justus Liebigs Ann Chem* 662 1 1963; Karlson *Pure Appl Chem* 14 75 1967.]

β -Ecdyson (β -echdysterone) [5289-74-7] M 480.7, m 245-247°, $[\alpha]_D^{20} +66^\circ$ (c 1, MeOH). Crystd from water or tetrahydrofuran/pet ether.

Echinenone [432-68-8] M 550.8, m 178-179°, $A_{1\text{cm}}^{1\%}(\lambda_{\max})$ 2160 (458nm) in pet ether. Purified by chromatography on partially deactivated alumina or magnesia, or by using a thin layer of silica gel G with 4:1 cyclohexane/diethyl ether as the developing solvent. Stored in the dark at -20°.

Eicosane [112-95-8] M 282.6, m 36-37°, b 205°/15mm, $d^{36.7}$ 0.7779, n^{40} 1.43453. Crystd from EtOH.

Elaidic (*trans*-oleic) acid [112-79-8] M 282.5, m 44.5°, pK^{25} 4.9. Crystd from acetic acid, then EtOH.

Ellagic acid (2H₂O) [476-66-4] M 338.2, m >360°, $pK_{\text{Est}(1)} \sim 8$, $pK_{\text{Est}(2)} \sim 11$. Crystd from pyridine.

Elymoclavine (8,9-didehydro-6-methylergoline-8-methanol) [548-43-6] M 254.3, m 249-253°(dec), $[\alpha]_D^{20} -59^\circ$ (c 1, EtOH). Crystd from MeOH.

Embonic acid (Pamoic acid, 4,4'-methylene bis[3-hydroxy-2-naphthalenecarboxylic acid]) [130-85-8] M 388.4, m 295°, >300°, $pK_{\text{Est}(1)} \sim 2.2$, $pK_{\text{Est}(2)} \sim 13.2$. Forms crystals from dilute pyridine which decomposition above 280° without melting. It is almost insoluble in H₂O, EtOH, Et₂O, *C₆H₆, CH₃CO₂H, sparingly soluble in CHCl₃ but soluble in nitrobenzene, pyridine and alkalis [Barber and Gaimster *J Appl Chem (London)* 2 565 1952]. Used for making salts of organic bases.

Emetine hydrochloride hydrate [316-42-7] M 553.6 + aq, m 235-240°, 235-250°, 240-250°, 248-250° (depending on H₂O content), $[\alpha]_D^{20} -49.2^\circ$ (free base, c 4, CHCl₃), +18° (c 6, H₂O, dry salt), pK_1 5.77, pK_2 6.64. It crystallises from MeOH-Et₂O, MeOH or Et₂O-EtOAc. The *free base* has m 104-105°, and the (*-*)-*phenyl thiourea derivative* has m 220-221° [from EtOAc-pet ether, $[\alpha]_D^{25} -29.3^\circ$ (CHCl₃)]. IR: 3413 (OH) and 2611 (NH⁺) cm⁻¹; UV λ_{\max} 230nm (ϵ 16 200) and 282nm (ϵ 6 890) [Brossi et al. *Helv Chim Acta* 42 1515 1959; Barash et al. *J Chem Soc* 3530 1959].

Emodine (1,3,8-trihydroxy-6-methyl-9,10-anthracenedione, archin) [518-82-1] M 270.2, m 253-257°, 255-256°, 256-257°, 262°, 264° (phenolic pKs 7-10). Forms orange needles from EtOH, Et₂O, *C₆H₆, toluene or pyridine. It sublimes above 200° at 12mm. [Tutin and Clewer *J Chem*

Soc 99 946 1911; IR: Bloom et al. *J Chem Soc* 178 1959; UV: Birkinshaw *Biochem J* 59 495 1955; Raistrick *Biochem J* 34 159 1940.]

1*R*,2*S*-(-)Ephedrine [299-42-3] *M* 165.2, *m* 40°, *b* 225°, $[\alpha]_{546}^{20}$ -47° and $[\alpha]_{\text{D}}^{20}$ -40° (c 5, 2.2M HCl), pK^{25} 9.57. See (-)-ephedrine (1*R*,2*S*-2-methylamino-1-phenylpropanol) on p. 533 in Chapter 6.

(-)-Ephedrine hydrochloride [50-98-6] *M* 201.7, *m* 218°, $[\alpha]_{546}^{20}$ -48° (c 5, 2M HCl). Crystd from water.

Epichlorohydrin [106-89-8] *M* 92.5, *b* 115.5°, *n* 1.438, *d* 1.180. Distd at atmospheric pressure, heated on a steam bath with one-quarter its weight of CaO, then decanted and fractionally distd.

***R*(-)-Epinephrine (adrenalin)** [51-43-4] *M* 183.2, *m* 215°(dec), $[\alpha]_{546}^{20}$ -61° (c 5, 0.5M HCl), pK_1^{25} 8.75, pK_2^{25} 9.89, pK_3^{25} ~13. Dissolved in dilute aqueous acid, then pptd by addn of dilute aqueous ammonia or alkali carbonates. (Epinephrine readily oxidises in neutral alkaline soln. This can be diminished if a little sulfite is added).

1,2-Epoxybutane [106-88-7] *M* 72.1, *b* 66.4-66.6°, *d* 0.837, *n* 1.3841. Dried with CaSO₄, and fractionally distd through a long (126cm) glass helices-packed column. The first fraction contains a water azeotrope.

(+)-Equilenine [517-09-9] *M* 266.3, *m* 258-259°, $[\alpha]_{\text{D}}^{16}$ +87° (c 7.1, H₂O). Crystd from EtOH and dried in a vacuum.

Ergocornine [564-36-3] *M* 561.7, *m* 182-184°, $[\alpha]_{\text{D}}^{20}$ -176° (c 0.5, CHCl₃). Crystd with solvent of crystn from MeOH.

Ergocristine [511-08-0] *M* 573.7, *m* 165-170°, $[\alpha]_{\text{D}}^{20}$ -183° (c 0.5, CHCl₃). Crystd with 2 moles of solvent of crystn, from *benzene.

Ergocryptine [511-09-1] *M* 575.7, *m* 212-214°, $[\alpha]_{\text{D}}^{20}$ -180° (c 0.5, CHCl₃). Crystd with solvent of crystn, from acetone, *benzene or methanol.

Ergosterol [57-87-4] *M* 396.7, *m* 165-166°, $[\alpha]_{546}^{20}$ -171° (in CHCl₃). Crystd from ethyl acetate, then from ethylene dichloride.

Ergotamine [113-15-5] *M* 581.6, *m* 212-214°(dec), $[\alpha]_{\text{D}}^{20}$ -160° (c 0.5, CHCl₃), pK^{25} 6.40. Crystd from *benzene, then dried by prolonged heating in high vacuum. Very *hygroscopic*.

Ergotamine tartrate [379-79-3] *M* 657.1, *m* 203°(dec). Crystd from MeOH.

Erucic acid (cis-13-docosenoic acid) [112-86-7] *M* 338.6, *m* 33.8°, *b* 358°/400mm, pK_{Est} ~4.9. Crystd from MeOH.

meso-Erythritol [149-32-6] *M* 122.1, *m* 122°. Crystd from distd water and dried at 60° in a vac oven.

Erythrityl tetranitrate [7297-25-8] *M* 302.1, *m* 61°. Crystd from EtOH.

β-Erythroidine [466-81-9] *M* 273.3, $[\alpha]_{\text{D}}^{20}$ +89° (H₂O). Crystd from EtOH.

D-Erythronic acid (3*R*-3,4-dihydroxyfuran-2-one) [15667-21-7] *M* 118.1, *m* 98-100°, 103-104°, 104-105°, 105°, $[\alpha]_{\text{D}}^{20}$ -73.2° (c 0.5, H₂O), $[\alpha]_{546}^{20}$ -87.6° (c 4, H₂O). Recrystd from EtOAc (20 parts) or isoPrOH (3 parts). [Baker and MacDonald *J Am Chem Soc* 82 230 1960; Glattfeld and

Forbrich *J Am Chem Soc* **56** 1209 1934; Weidenhagen and Wegner *Chem Ber* **72** 2010 1939, Musich and Rapoport *J Am Chem Soc* **100** 4865 1978.]

Esculetin (cichorigenin, 6,7-dihydroxycoumarin) [305-01-1] **M 178.2, m 272-275° (dec), 274° (dec), $pK_{Est(1)} \sim 8.7$, $pK_{Est(2)} \sim 12.4$** . Forms prisms from AcOH or aq EtOH and provides leaflets on sublimation in a vacuum. [Sethna and Shah *Chem Rev*; Merz *Arch Pharm (Weinheim Ger)* **270** 486 1932.] *Esculin (the 6-glucoside)* has **m 215° (dec), $[\alpha]_D^{20} -41^\circ$ (c 5, pyridine)**.

Eserine (Physostigmine, Physostol, [(3a*S*-*cis*)-1,2,3,3a,8,8a-hexahydro-1,3a,8-trimethylpyrrolo[2,3-*b*]indol-5-ol methylcarbamate ester] [57-47-6] **M 275.4, m 102-104°, 105-106°, $[\alpha]_D^{17} -67^\circ$ (c 1.3, CHCl₃), $[\alpha]_D^{25} -120^\circ$ (*C₆H₆), $pK_1^{15} 1.96$, $pK_2^{15} 8.08$** . Recrystallises from Et₂O or *C₆H₆ and forms an unstable low melting form **m 86-87°** [Harley-Mason and Jackson *J Chem Soc* 3651 1954; Wijnberg and Speckamp *Tetrahedron* **34** 2399 1978].

1,3,5-Estratrien-3-ol-17-one (Estrone, Folliculin) [53-16-7] **M 270.4, m 260-261°, polymorphic also m 254° and 256°, $[\alpha]_{546}^{20} +198^\circ$ (c 1, dioxane), $pK^{25} 10.91$** . Crystd from EtOH.

1,3,5-Estratrien-3 β ,16a,17 β -triol (Estriol) [50-27-1] **M 288.4, m 283°, $[\alpha]_{546}^{20} +66^\circ$ (c 1, dioxane)**. Crystd from EtOH/ethyl acetate.

Ethane [74-84-0] **M 30.1, f -172°, b -88°, $d_4^0 1.0493$ (air = 1)**. Ethylene can be removed by passing the gas through a sintered-glass disc into fuming H₂SO₄ then slowly through a column of charcoal satd with bromine. Bromine and HBr were removed by passage through firebrick coated with *N,N*-dimethyl-*p*-toluidine. The ethane was also passed over KOH pellets (to remove CO₂) and dried with Mg(ClO₄)₂. Further purification was by several distns of liquified ethane, using a condensing temperature of -195°. Yang and Gant [*J Phys Chem* **65** 1861 1961] treated ethane by standing it for 24h at room temperature in a steel bomb containing activated charcoal treated with bromine. They then immersed the bomb in a Dry-ice/acetone bath and transferred the ethane to an activated charcoal trap cooled in liquid nitrogen. (The charcoal had previously been degassed by pumping for 24h at 450°.) By allowing the trap to warm slowly, the ethane was distd, retaining only the middle third. Removal of methane was achieved using Linde type 13X molecular sieves (previously degassed by pumping for 24h at 450°) in a trap which, after cooling in Dry-ice/acetone, was satd with ethane. After pumping for 10min, the ethane was recovered by warming the trap to room temperature. (The final gas contained less than 10⁻⁴ mole % of either ethylene or methane).

Ethanesulfonyl chloride [594-44-5] **M 128.6, b 55°/9mm, 62°/12mm, 74°/19mm, 76-79°/22mm, 95-98°/50mm, 177°/760mm, $d_4^{20} 1.357$, $n_D^{20} 1.4539$** . Purified by repeated distn to remove HCl formed from hydrolysis. **Fuming, corrosive liquid, handle in a good fumehood**. It is hydrolysed by aq N NaOH at room temperature and is best stored in aliquots in sealed ampules under N₂. [Davies and Dick *J Chem Soc* 484 1932; Klamann and Drahowzal *Monatsh Chem* **83** 463 1952; Saunders et al. *Biochem J* **36** 372 1942.]

Ethanethiol (ethyl mercaptan) [75-08-1] **M 62.1, b 32.9°/704mm, $d^{52} 0.83147$, $pK^{25} 10.61$** . Dissolved in aqueous 20% NaOH, extracted with a small amount of *benzene and then steam distd until clear. After cooling, the alkaline soln was acidified slightly with 15% H₂SO₄ and the thiol was distd off, dried with CaSO₄, CaCl₂ or 4A molecular sieves, and fractionally distd under nitrogen [Ellis and Reid *J Am Chem Soc* **54** 1674 1932].

Ethanol [64-17-5] **M 46.1, b 78.3°, $d^{15} 0.79360$, $d^5 0.78506$, n 1.36139, $pK^{25} 15.93$** . Usual impurities of fermentation alcohol are fusel oils (mainly higher alcohols, especially pentanols), aldehydes, esters, ketones and water. With synthetic alcohol, likely impurities are water, aldehydes, aliphatic esters, acetone and diethyl ether. Traces of *benzene are present in ethanol that has been dehydrated by azeotropic distillation with *benzene. Anhydrous ethanol is very *hygroscopic*. Water (down to 0.05%) can be detected by formation of a voluminous ppt when aluminium ethoxide in *benzene is added to a test portion, Rectified

spirit (95% ethanol) is converted to *absolute* (99.5%) ethanol by refluxing with freshly ignited CaO (250g/L) for 6h, standing overnight and distilling with precautions to exclude moisture.

Numerous methods are available for further drying of *absolute* ethanol for making "Super dry ethanol". Lund and Bjerrum [*Chem Ber* **64** 210 1931] used reaction with magnesium ethoxide, prepared by placing 5g of clean dry magnesium turnings and 0.5g of iodine (or a few drops of CCl₄), to activate the Mg, in a 2L flask, followed by 50-75 mL of *absolute* ethanol, and warming the mixture until a vigorous reaction occurs. When this subsides, heating is continued until all the magnesium is converted to magnesium ethoxide. Up to 1L of ethanol is added and, after an hour's reflux, it is distd off. The water content should be below 0.05%. Walden, Ulich and Laun [*Z Phys Chem* **114** 275 1925] used amalgamated aluminium chips, prepared by degreasing aluminium chips (by washing with Et₂O and drying in a vac to remove grease from machining the Al), treating with alkali until hydrogen was vigorously evolved, washing with H₂O until the washings were weakly alkaline and then stirring with 1% HgCl₂ soln. After 2min, the chips were washed quickly with H₂O, then alcohol, then ether, and dried with filter paper. (The amalgam became warm.) These chips were added to the ethanol, which was then gently warmed for several hours until evolution of hydrogen ceased. The alcohol was distd and aspirated for some time with pure dry air. Smith [*J Chem Soc* 1288 1927] reacted 1L of *absolute* ethanol in a 2L flask with 7g of clean dry sodium, and added 25g of pure ethyl succinate 27g of pure ethyl phthalate was an alternative), and refluxed the mixture for 2h in a system protected from moisture, and then distd the ethanol. A modification used 40g of ethyl formate, instead, so that sodium formate separated out and, during reflux, the excess of ethyl formate decomposed to CO and ethanol.

Drying agents suitable for use with ethanol include Linde type 4A molecular sieves, calcium metal, and CaH₂. The calcium hydride (2g) was crushed to a powder and dissolved in 100mL *absolute* ethanol by gently boiling. About 70mL of the ethanol were distd off to remove any dissolved gases before the remainder was poured into 1L of *ca* 99.9% ethanol in a still, where it was boiled under reflux for 20h, while a slow stream of pure, dry hydrogen (better use nitrogen or Ar) was passed through. It was then distd [Rüber *Z Elektrochem* **29** 334 1923]. If calcium was used for drying, about ten times the theoretical amount should be taken, and traces of ammonia (from some calcium nitride in the Ca metal) would be removed by passing dry air into the vapour during reflux.

Ethanol can be freed from traces of basic materials by distn from a little 2,4,6-trinitrobenzoic acid or sulfanilic acid. *Benzene can be removed by fractional distn after adding a little water (the *benzene/water/ethanol azeotrope distils at 64.9°); the alcohol is then redried using one of the methods described above. Alternatively, careful fractional distn can separate *benzene as the *benzene/ethanol azeotrope (b 68.2°). Aldehydes can be removed from ethanol by digesting with 8-10g of dissolved KOH and 5-10g of aluminium or zinc per L, followed by distn. Another method is to heat under reflux with KOH (20g/L) and AgNO₃ (10g/L) or to add 2.5-3g of lead acetate in 5mL of water to 1L of ethanol, followed (slowly and without stirring) by 5g of KOH in 25mL of ethanol: after 1hr the flask is shaken thoroughly, then set aside overnight before filtering and distilling. The residual water can be removed by standing the distillate over activated aluminium amalgam for 1 week, then filtering and distilling. Distn of ethanol from Raney nickel eliminates catalyst poisons.

Other purification procedures include pre-treatment with conc H₂SO₄ (3mL/L) to eliminate amines, and with KMnO₄ to oxidise aldehydes, followed by refluxing with KOH to resinify aldehydes, and distilling to remove traces of H₃PO₄ and other acidic impurities after passage through silica gel, and drying over CaSO₄. Water can be removed by azeotropic distn with dichloromethane (azeotrope boils at 38.1° and contains 1.8% water) or 2,2,4-trimethylpentane.

Rapid purification: Place degreased Mg turnings (grease from machining the turnings is removed by washing with dry EtOH then Et₂O, and drying in a vac) (5g) in a dry 2L round bottomed flask fitted with a reflux condenser (protect from air with a drying tube filled with CaCl₂ or KOH pellets) and flush with dry N₂. Then add iodine crystals (0.5g) and gently warm the flask until iodine vapour is formed and coats the turnings. Cool, then add EtOH (50mL) and carefully heat to reflux until the iodine disappears. Cool again then add more EtOH (to 1L) and reflux under N₂ for several hours. Distil and store over 3A molecular sieves (pre-heated at 300°-350° for several hours and cooled under dry N₂ or argon).

S-Ethionine [13073-35-3] M 163.2, m 282°(dec), [α]_D²⁵ +23.7° (in 5M HCl), pK²⁵ 9.02 (for RS). Likely impurities are N-acetyl-(R and S)-ethionine, S-methionine, and R-ethionine. Crystd from water by adding 4 volumes of EtOH.

Ethoxycarbonyl isocyanate [19617-43-7] M 115.1, b 51-55°/13mm, 56°/18mm, d_4^{20} 1.15. Fractionally distilled. [*J Heterocycl Chem* 5 837 1968.]

Ethoxycarbonyl isothiocyanate [16182-04-0] M 131.5, b 43°/14mm, 51-55°/13mm, 56°/18mm, d_4^{20} 1.12. Fractionally distd through a short column. It also distils at 83°/30mm with some decomposition liberating CO₂ and sulfurous gases, best distil below 20mm vacuum. [*J Chem Soc* 93 697 1908; 1340, 1948; *J Heterocycl Chem* 5 837 1968.]

3-Ethoxy-N,N-diethylaniline [1846-92-2] M 193.3, b 141-142°/15mm pK_{Est} ~6.1. Refluxed for 3h with acetic anhydride, then fractionally distilled under reduced pressure.

2-Ethoxyethanol [110-80-5] M 90.1, b 134.8°, d 0.931, n 1.40751. Dried with CaSO₄ or K₂CO₃, filtered and fractionally distd. Peroxides can be removed by refluxing with anhydrous SnCl₂ or by filtration under slight pressure through a column of activated alumina.

2-Ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ) [16357-59-8] M 247.3, m 63.5-65°, 66-67°. Dissolve ~180g in CHCl₃, evap to dryness under vac. Add dry Et₂O (20mL) when a white solid separates on standing. Set aside for a few hours, collect solid, wash thoroughly with cold Et₂O and dry in vac (~140g, m 63.5-65°). A further crop of solid (~25g) is obtained from the filtrate on standing overnight. [Fieser and Fieser *Reagents for Organic Synthesis* 2 191 1969; Belleau et al. *J Am Chem Soc* 90 823 1968 and 90 1651 1968.]

2-Ethoxyethyl ether [bis-(2-ethoxyethyl) ether] [112-36-7] M 162.2, b 76°/32mm, d 0.910, n 1.412. See diethyleneglycol diethyl ether on p. 203.

2-Ethoxyethyl methacrylate [2370-63-0] M 158.2, b 91-93°/35mm, d 0.965, n 1.429. Purified as described under methyl methacrylate.

1-Ethoxynaphthalene [5328-01-8] M 172.2, b 136-138°/14mm, 282°/760mm, d 1.061, n 1.604. Fractionally distd (twice) under a vacuum, then dried with, and distd under a vacuum from, sodium.

2-Ethoxynaphthalene [93-18-5] M 172.2, m 35.6-36.0°, b 142-143°/12mm. Crystd from pet ether. Dried under vacuum or distd in a vacuum.

Ethyl acetimidate [1000-84-6] M 87.1, b 92-95°/atm, 89.7-90°/765mm, d 0.8671, n 1.4025, pK_{Est} ~5.5. It is best to prepare it freshly from the *hydrochloride* (see below). Dissolve the hydrochloride (123.5g) by adding it slowly to an ice-cold mixt of H₂O (500mL), K₂CO₃ (276g) and Et₂O (200mL) and stirring rapidly. The Et₂O layer was separated, the aq layer was extd with Et₂O (100mL), the combined Et₂O layers were dried (MgSO₄), evapd and the residual oil distd through a glass helices packed column (70x1.2cm). The yield was 19g (22%). [Glickman and Cope *J Am Chem Soc* 67 1020 1945; Chaplin and Hunter *J Chem Soc* 1118 1937; *Methods Enzymol* 25 585 1972.]

Ethyl acetimidate hydrochloride [2208-07-3] M 123.6, m 98-100°(dec), 110-115° (dec), 112-113°(dec), m 112-114°(dec), pK_{Est} ~5.5. Recrystd by dissolving in the minimum volume of super dry EtOH and addition of dry Et₂O or from dry Et₂O. Dry in vacuum and store in a vacuum desiccator with P₂O₅. Alternatively it could be crystd from EtOH (containing a couple of drops of ethanolic HCl) and adding dry Et₂O. Filter and dry in a vac desiccator over H₂SO₄ and NaOH. [Pinner *Chem Ber* 16 1654 1883.] [Glickman and Cope *J Am Chem Soc* 67 1020 1945; Chaplin and Hunter *J Chem Soc* 1118 1937; McElvain and Schroeder *J Am Chem Soc* 71 40 1949; McElvain and Tate *J Am Chem Soc* 73 2233 1951; *Methods Enzymol* 25 585 1972.]

Ethyl acetate [141-78-6] M 88.1, b 77.1°, d 0.9003, n 1.37239, n_D^{25} 1.36979, pK^{25} -6.93 (aq H₂SO₄). The commonest impurities are water, EtOH and acetic acid. These can be removed by washing with aqueous 5% Na₂CO₃, then with saturated aqueous CaCl₂ or NaCl, and drying with K₂CO₃, CaSO₄ or MgSO₄. More efficient drying is achieved if the solvent is further dried with P₂O₅, CaH₂ or molecular sieves before

distn. CaO has also been used. Alternatively, ethanol can be converted to ethyl acetate by refluxing with acetic anhydride (*ca* 1mL per 10mL of ester); the liquid is then fractionally distd, dried with K_2CO_3 and redistd.

Rapid purification: Distil, dry over K_2CO_3 , distil again and store over 4A molecular sieves.

Ethyl acetoacetate [141-97-9] M 130.1, b 71°/12mm, 100°/80mm, d 1.026, n 1.419, pK²⁵ 10.68. Shaken with small amounts of saturated aqueous $NaHCO_3$ (until no further effervescence), then with water. Dried with $MgSO_4$ or $CaCl_2$. Distd under reduced pressure.

Ethyl acrylate [140-88-5] M 100.1, b 20°/40mm, 99.5°/atm, d 0.922, n 1.406. Washed repeatedly with aqueous NaOH until free from inhibitors such as hydroquinone, then washed with saturated aqueous $CaCl_2$ and distd under reduced pressure. Hydroquinone should be added if the ethyl acrylate is to be stored for extended periods. **LACHRYMATORY.**

Ethylamine [75-04-7] M 45.1, b 16.6°/760mm, d 1.3663, pK²⁰ 10.79. Condensed in an all-glass apparatus cooled by circulating ice-water, and stored with KOH pellets below 0°.

Ethylamine hydrochloride [557-66-4] M 81.5, m 109-110°. Crystd from absolute EtOH or MeOH/ $CHCl_3$.

Ethyl o-aminobenzoate [94-09-7] M 165.2, m 92°, pK²⁵ 2.39. Crystd from EtOH/ H_2O and air dried.

p-Ethylaniline [589-16-2] M 121.2, b 88°/8mm, d 0.975, n 1.554, pK²⁵ 5.00. Dissolved in *benzene, then acetylated. The acetyl derivative was recrystallised from *benzene/pet ether, and hydrolysed by refluxing 50g with 500mL of water and 115mL of conc H_2SO_4 until the soln becomes clear. The amine sulfate was isolated, suspended in water and solid KOH was added to regenerate the free base, which was separated, dried and distd from zinc dust under a vacuum [Berliner and Berliner *J Am Chem Soc* 76 6179 1954].

Ethylbenzene [100-41-6] M 106.2, b 136.2°, d 0.867, n 1.49594, n²⁵ 1.49330. Shaken with cold conc H_2SO_4 until a fresh portion of acid remained colourless, then washed with aqueous 10% NaOH or $NaHCO_3$, followed by distilled water until neutral. Dried with $MgSO_4$ or $CaSO_4$, then dried further with, and distd from, sodium, sodium hydride or CaH_2 . Can also be dried by passing through silica gel. Sulfur-containing impurities have been removed by prolonged shaking with mercury. Also purified by fractional freezing.

Ethyl benzoate [93-89-0] M 150.2, b 98°/19mm, 212.4°/760mm, d 1.046, n¹⁵ 1.5074, n²⁵ 1.5043, pK -7.37 (aq H_2SO_4). Washed with aq 5% Na_2CO_3 , then satd $CaCl_2$, dried with $CaSO_4$ and distd under reduced pressure.

Ethyl bis-(2,4-dinitrophenyl)acetate [5833-18-1] M 358.3, m 150-153°. Crystd from toluene as pale yellow crystals.

Ethyl bixin [6895-43-8] M 436.6, m 138°. Crystd from EtOH.

Ethyl bromide [74-96-4] M 109.0, b 0°/165mm, 38°/745mm, d 1.460, n 1.4241. The main impurities are usually EtOH and water, with both of which it forms azeotropes. Ethanol and unsaturated compounds can be removed by washing with conc H_2SO_4 until no further coloration is produced. The ethyl bromide is then washed with water, aq Na_2CO_3 , and water again, then dried with $CaCl_2$, $MgSO_4$ or CaH_2 , and distd. from P_2O_5 . Olefinic impurities can also be removed by storing the ethyl bromide in daylight with elementary bromine, later removing the free bromine by extraction with dil aq Na_2SO_3 , drying the ethyl bromide with $CaCl_2$ and fractionally distilling. Alternatively, unsaturated compounds can be removed by bubbling oxygen containing *ca* 5% ozone through the liquid for an hour, then washing with aqueous Na_2SO_3 to hydrolyse ozonides and remove hydrolysis products, followed by drying and distn.

Ethyl bromoacetate [105-36-2] M 167.0, b 158-158.5°/758mm, d 1.50, n 1.450. Washed with saturated aqueous Na₂CO₃ (three times), 50% aq CaCl₂ (three times) and saturated aqueous NaCl (twice). Dried with MgSO₄, CaCl₂ or CaCO₃, and distd. **LACHRYMATORY.**

Ethyl 2-(bromomethyl)acrylate [17435-72-2] M 193.1, b 38°/0.8mm, d 1.398, n 1.479. If it contains some free acid, add H₂O, cool, and neutralise with NaHCO₃ until evolution of CO₂ ceases. Extract the mixt with Et₂O (3x) and dry combined extracts (Na₂SO₄, 3h). Evap Et₂O and dist ester collecting fraction b 39-40°/0.9mm and check spectra. [Prep and NMR: Ramarajan et al. *Org Synth Coll Vol VII* 211 1990.]

Ethyl α-bromopropionate [535-11-5] M 181.0, b 69-70°/25mm, d 1.39, n 1.447. Washed with saturated aqueous Na₂CO₃ (three times), 50% aq CaCl₂ (three times) and saturated aqueous NaCl (twice). Dried with MgSO₄, CaCl₂ or CaCO₃, and distd. **LACHRYMATORY.**

Ethyl bromopyruvate [70-23-5] M 195.0, b 47°/0.5mm, 71-73°/5mm, 87°/9mm, 89-104°/14mm, d₄²⁰ 1.561, n_D²⁰ 1.464. Most likely impurity is free carboxylic acid (bromopyruvic or bromoacetic acids). Dissolve in dry Et₂O or dry CHCl₃, stir with CaCO₃ until effecrvescence ceases, filter, (may wash with a little H₂O rapidly), dry (MgSO₄) and distil at least twice. The 2,4-dinitrophenylhydrazone has m 144-145°. [Burros and Holland *J Chem Soc* 672 1947; Letsinger and Laco *J Org Chem* 21 764 1956; Kruse et al. *J Am Chem Soc* 76 5796 1954.] **LACHRYMATORY.**

2-Ethyl-1-butanol [97-95-0] M 102.2, b 146.3°, n¹⁵ 1.4243, n²⁵ 1.4205. Dried with CaSO₄ for several weeks, filtered and fractionally distd.

2-Ethylbut-1-ene [760-21-4] M 84.1, b 66.6°, d 0.833, n 1.423. Washed with saturated aqueous NaOH, then water. Dried with CaCl₂, filtered and fractionally distd.

Ethyl n-butyrate [105-54-4] M 116.2, b 49°/50mm, 119-120°/760mm, d 0.880, n 1.393. Dried with anhydrous CuSO₄ and distd under dry nitrogen.

Ethyl carbamate (urethane) [51-79-6] M 88.1, m 48.0-48.6°. Crystd from *benzene.

Ethyl carbazate (N-ethoxycarbonyl hydrazine) [4114-31-2] M 104.1, m 44-48°, 51-52°, b 95.5°/10m, 92-95°/12mm, 100-102°/11mm. Fractionated using a Vigreux column until the distillate crystallises [Allen and Bell *Org Synth Coll Vol III* 404 1955.]

N-Ethylcarbazole [86-28-2] M 195.3, m 69-70°. Recrystd from EtOH, EtOH/water or isopropanol and dried below 55°.

Ethyl carbonate [105-58-8] M 118.1, b 124-125°, d 0.975, n 1.38287. See diethyl carbonate on p. 203.

Ethyl chloride [75-00-3] M 64.5, b 12.4°, d 0.8978, n 1.3676. Passed through absorption towers containing, successively, conc H₂SO₄, NaOH pellets, P₂O₅ on glass wool, or soda-lime, CaCl₂, P₂O₅. Condensed into a flask containing CaH₂ and fractionally distd. Has also been purified by illumination in the presence of bromine at 0° using a 1000W lamp, followed by washing, drying and distn.

Ethyl chloroacetate [105-39-5] M 122.6, b 143-143.2°, d 1.150, n²⁵ 1.4192. Shaken with satutated aqueous Na₂CO₃ (three times), aqueous 50% CaCl₂ (three times) and saturated aqueous NaCl (twice). Dried with Na₂SO₄ or MgSO₄ and distd. **LACHRYMATORY.**

Ethyl chloroformate [541-41-3] M 108.5, m -81°, b 94-95°, d 1.135, n 1.3974. Washed several times with water, redistd using an efficient fractionating column at atmospheric pressure and a CaCl₂ guard tube to keep free from moisture [Hamilton and Sly *J Am Chem Soc* 47 435 1925; Saunders, Slocombe and Hardy, *J Am Chem Soc* 73 3796 1951]. **LACHRYMATORY AND TOXIC.**

Ethyl chrysanthemate (ethyl \pm 2,2-dimethyl-3{c and t}-[2-methylpropenyl]-cyclopropane carboxylate) [97-41-6] M 196.3, b 98-102°/11mm, 117-121°/20mm. Purified by vacuum distn. The free *trans*-acid has m 54° (from, EtOAc) and the free *cis*-acid has m 113-116° (from EtOAc). The 4-nitrophenyl ester has m 44-45° (from pet ether) [Campbell and Harper *J Chem Soc* 283 1945; IR: Allen et al. *J Org Chem* 22 1291 1957].

Ethyl cinnamate [103-36-6] M 176.2, f 6.7°, b 127°/6mm, 272.7°/768mm, d 1.040, n 1.55983. Washed with aqueous 10% Na₂CO₃, then water, dried (MgSO₄), and distd. The purified ester was saponified with aqueous KOH, and, after acidifying the soln, cinnamic acid was isolated, washed and dried. The ester was reformed by refluxing for 15h the cinnamic acid (25g) with abs EtOH (23g), conc H₂SO₄ (4g) and dry *benzene (100mL), after which it was isolated, washed, dried and distd under reduced pressure [Jeffery and Vogel *J Chem Soc* 658 1958].

Ethyl *trans*-crotonate [623-70-1] M 114.2, b 137°, d 0.917, n 1.425. Washed with aqueous 5% Na₂CO₃, washed with saturated aqueous CaCl₂, dried with CaCl₂ and distd.

Ethyl cyanoacetate [105-56-6] M 113.1, b 206.0°, d 1.061, n 1.41751. Shaken several times with aqueous 10% Na₂CO₃, washed well with water, dried with Na₂SO₄ and fractionally distd.

Ethyl cyanofornate [623-49-4] M 99.1, b 113-114°/740mm, 116.5-116.8°/765.5mm, d₄²⁰ 1.0112, n_D²⁰ 1.3818. Dissolve in Et₂O, dry over Na₂SO₄, filter, evaporate and distil [Malachowsky et al. *Chem Ber* 70 1016 1937; Adickes et al. *J Prakt Chem* [2] 133 313 1932; Grundmann et al. *Justus Liebigs Ann Chem* 577 77 1952].

Ethylcyclohexane [1678-91-7] M 112.2, b 131.8°, d 0.789, n 1.43304, n²⁵ 1.43073. Purified by azeotropic distn with 2-ethoxyethanol, then the alcohol was washed out with water and, after drying, the ethylcyclohexane was redistd.

Ethyl cyclohexanecarboxylate [3289-28-9] M 156.2, b 76-77°/10mm, 92-93°/34mm, d 0.960, n 1.420. Washed with M sodium hydroxide solution, then water, dried with Na₂SO₄ and distd.

Ethyl diazoacetate [623-73-4] M 114.1, m -22°, b 42°/5mm, 45°/12mm, 85-86°/88mm, 140-141°/720mm, 140-143°/atm, d₄^{17.6} 1.0852, n_D^{17.6} 1.4588. A very volatile yellow oil with a strong pungent odour. **EXPLOSIVE [distillation even under reduced pressure is dangerous and may result in an explosion — TAKE ALL THE NECESSARY PRECAUTIONS IF DISTILLATION IS TO BE CARRIED OUT]**. It explodes in contact with conc H₂SO₄ - trace acid causes rapid decomp. It is slightly sol in H₂O, but is miscible with EtOH, *C₆H₆, pet ether and Et₂O. To purify dissolve in Et₂O [using CH₂Cl₂ instead of Et₂O protects the ester from acid], wash with 10% aq Na₂CO₃, dry (MgSO₄), filter and repeat as many times as possible until the Et₂O layer loses its yellow colour, remove the solvent below 20° (vac). Note that prolonged heating may lead to rapid decomp and low yields. It can also be purified by steam distn under reduced pressure but with considerable loss in yield. Place the residual oil in a brown bottle and keep below 10°, and use as soon as possible without distilling. [Womack and Nelson *Org Synth Coll Vol III* 392 1955; UV: Miller and White *J Am Chem Soc* 79 5974 1957; Fieser 1 367 1967.]

Ethyl dibromoacetate [617-33-4] M 245.9, b 81-82°/14.5mm, n²² 1.4973. Washed briefly with conc aqueous NaHCO₃, then with aqueous CaCl₂. Dried with CaSO₄ and distd under reduced pressure.

Ethyl α,β -dibromo- β -phenylpropionate [5464-70-0, *erythro* 30983-70-1] M 336.0, m 75°. Crystd from pet ether (b 60-80°).

Ethyl dichloroacetate [535-15-9] M 157.0, b 131.0-131.5°/401mm, d 1.28, n 1.438. Shaken with aqueous 3% NaHCO₃ to remove free acid, washed with distd water, dried for 3 days with CaSO₄ and distd under reduced pressure.

Ethyl 3,3-diethoxypropionate [10601-80-6] M 190.2, b 58.5°/1.5mm, 65°/2mm, 95-96°/12mm, d_4^{20} 0.78, n_D^{25} 1.4101. Dissolve in dry Et₂O, and dry with solid NaHCO₃, filter and distil and carefully fractionate [Dyer and Johnson *J Am Chem Soc* 56 223 1934].

Ethyl 1,3-dithiane-2-carboxylate [20462-00-4] M 192.3, b 75-77°/0.2mm, 96°/0.4mm, d_4^{20} 1.220, n_D^{25} 1.5379. Dissolve in CHCl₃, wash with aqueous K₂CO₃, 2 x with H₂O, dry over MgSO₄, filter, evaporate and distil. [Eliel and Hartman *J Org Chem* 37 505 1972; Seebach *Synthesis* 1 17 1969.]

Ethyl 1,3-dithiolane-2-carboxylate [20461-99-8] M 178.3, b 85°/0.1mm, d_4^{20} 1.250, n_D^{20} 1.538. Dissolve in CHCl₃, wash with aqueous K₂CO₃, 2 x with H₂O, dry over MgSO₄, filter, evaporate and distil [Hermann et al *Tetrahedron Lett* 2599 1973; Corey and Erickson *J Org Chem* 36 3553 1971].

Ethylene (ethene) [74-85-1] M 28.0, m -169.4°, b -102°/700mm. Purified by passage through a series of towers containing molecular sieves or anhydrous CaSO₄ or a cuprous ammonia soln, then conc H₂SO₄, followed by KOH pellets. Alternatively, ethylene has been condensed in liquid nitrogen, with melting, freezing and pumping to remove air before passage through an activated charcoal trap, followed by a further condensation in liquid air. A sputtered sodium trap has also been used, to remove oxygen.

Ethylene N,N'-bis[(o-hydroxyphenyl)glycine] [1170-02-1] M 360.4, m 249°(dec), $pK_{Est(1)}$ ~1.8, $pK_{Est(2)}$ ~4.8, $pK_{Est(3)}$ ~9.0. Purified by extensive Soxhlet extraction with acetone. [Bonadies and Carrano *J Am Chem Soc* 108 4088 1986].

Ethylene carbonate (1,3-dioxolan-2-one) [96-49-1] M 88.1, m 37°, d 1.32, n^{40} 1.4199. Dried over P₂O₅ then fractionally distd at 10mm pressure. Crystd from dry diethyl ether.

Ethylenediamine (1,2-diaminoethane) [107-15-3] M 60.1, f 11.0°, b 117.0°, d 0.897, n 1.45677, n^{30} 1.4513, pK_1^{25} 6.86, pK_2^{25} 9.92. Forms a constant-boiling (b 118.5°) mixture with water (15%) [hygroscopic and miscible with water]. Recommended purification procedure [Asthana and Mukherjee in J.F.Coetzee (ed), *Purification of Solvents*, Pergamon Press, Oxford, 1982 cf p 53]: to 1L of ethylenediamine was added 70g of type 5A Linde molecular sieves and shaken for 12h. The liquid was decanted and shaken for a further 12h with a mixture of CaO (50g) and KOH (15g). The supernatant was fractionally distd (at 20:1 reflux ratio) in contact with freshly activated molecular sieves. The fraction distilling at 117.2° /760mm was collected. Finally it was fractionally distilled from sodium metal. All distns and storage of ethylenediamine should be carried out under nitrogen to prevent reaction with CO₂ and water. Material containing 30% water was dried with solid NaOH (600g/L), heated on a water bath for 10h. Above 60°, separation into two phases took place. The hot ethylenediamine layer was decanted off, refluxed with 40g of sodium for 2h and distd [Putnam and Kobe *Trans Electrochem Soc* 74 609 1938]. Ethylenediamine is usually distd under nitrogen. Type 5A Linde molecular sieves (70g/L), then a mixture of 50g of CaO and 15g of KOH/L, with further dehydration of the supernatant with molecular sieves has also been used for drying this diamine, followed by distn from molecular sieves and, finally, from sodium metal. A spectroscopically improved material was obtained by shaking with freshly baked alumina (20g/L) before distn.

N,N'-Ethylenediaminediacetic acid (EDDA) [5657-17-0] M 176.2, m 222-224°(dec), pK_1^{25} 6.48, pK_2^{25} 9.57 (for NH groups). Crystd from water.

Ethylenediamine dihydrochloride [333-18-6] M 133.0, pK_1^{25} 6.86, pK_2^{25} 9.92. Crystd from water.

Ethylenediaminetetraacetic acid (EDTA) [60-00-4] M 292.3, m 253°(dec), pK_1^{25} 0.26 pK_2^{25} 0.96, pK_3^{25} 2.60, pK_4^{25} 2.67, pK_5^{25} 6.16, pK_6^{25} 10.26. Dissolved in aqueous KOH or ammonium hydroxide, and pptd with dil HCl or HNO₃, twice. Boiled twice with distd water to remove mineral acid, then recrystd from water or dimethylformamide. Dried at 110°. Also recrystd from boiling 1N HCl, wash crystals with distd H₂O and dried *in vacuo*. [Ma and Ray *Biochemistry* 19 751 1980.]

Ethylene dimethacrylate [97-90-5] M 198.2, b 98-100°/5mm, d 1.053, n 1.456. Distd through a short Vigreux column at about 1mm pressure, in the presence of 3% (w/w) of phenyl-β-naphthylamine.

Ethylene dimyristate [627-84-9] M 482.8, m 61.7°. Crystd from *benzene-MeOH or diethyl ether-MeOH, and dried in a vacuum desiccator.

Ethylene dipalmitate [624-03-3] M 538.9, m 69.1°. Crystd from *benzene-MeOH or diethyl ether-MeOH and dried in a vacuum desiccator.

Ethylene distearate [627-83-8] M 595.0, m 75.3°. Crystd from *benzene-MeOH or diethyl ether-MeOH and dried in a vacuum desiccator.

Ethylene glycol [107-21-1] M 62.1, b 68°/4mm, 197.9°/760mm, d 1.0986, n¹⁵ 1.43312, n²⁵ 1.43056, pK²⁵ 10.6. Very *hygroscopic*, and also likely to contain higher diols. Dried with CaO, CaSO₄, MgSO₄ or NaOH and distd under vacuum. Further dried by reaction with sodium under nitrogen, refluxed for several hours and distd. The distillate was then passed through a column of Linde type 4A molecular sieves and finally distd under nitrogen, from more molecular sieves. Fractionally distd.

Ethylene glycol bis(β-aminoethylether)-N,N'-tetraacetic acid (EGTA) [67-42-5] M 380.4, m >245°(dec), pK₁²⁰ 1.15 (2.40), pK₂²⁰ 2.40 (2.50), pK₃²⁰ 8.40 (8.67), pK₄²⁰ 8.94 (9.22). Dissolved in aq NaOH, pptd by addn of aq HCl, washed with water and dried at 100° *in vacuo*.

Ethylene glycol diacetate [111-55-7] M 146.2, b 190.1°, 79-81°/11mm, d²⁵ 1.4188, n 1.4150. Dried with CaCl₂, filt'd (excluding moisture) fractionally distd under reduced pressure.

Ethylene glycol dibutyl ether [112-48-1] M 174.3, b 78-80°/0.2mm, d 1.105, n 1.42. Shaken with aq 5% Na₂CO₃, dried with MgSO₄ and stored with chromatographic alumina to prevent peroxide formation.

Ethylene glycol diethyl ether (1,2-diethoxyethane) [629-14-1] M 118.2, b 121.5°, d 0.842, n 1.392. After refluxing for 12h, a mixture of the ether (2L), conc HCl (27mL) and water (200mL), with slow passage of nitrogen, the soln was cooled, and KOH pellets were added slowly and with shaking until no more dissolved. The organic layer was decanted, treated with some KOH pellets and again decanted. It was refluxed with, and distd from sodium immediately before use. Alternatively, after removal of peroxides by treatment with activated alumina, the ether has been refluxed in the presence of the blue ketyl formed by sodium-potassium alloy with benzophenone, then distd.

Ethylene glycol dimethyl ether (monoglyme) [110-71-4] M 90.1. See 1,2-dimethoxyethane on p. 210.

Ethylene oxide [75-21-8] M 44.0, b 13.5°/746mm, d¹⁰ 0.882, n⁷ 1.3597. Dried with CaSO₄, then distd from crushed NaOH. Has also been purified by its passage, as a gas, through towers containing solid NaOH.

Ethylene thiourea (2-imidazolidinethione) [96-45-7] M 102.2, m 203-204°. Crystd from EtOH or amyl alcohol.

Ethylene urea (2-imidazolidone) [120-93-4] M 86.1, m 131°. Crystd from MeOH (charcoal).

Ethylenimine (aziridine) [151-56-4] M 43.1, b 55.5°/760mm, d 0.8321, pK²⁵ 8.00. See aziridine on p. 117.

2-Ethylethylenimine [2549-67-9] M 71.1, b 88.5-89°, pK²⁵ 8.31. Freshly distd from sodium before use. **TOXIC**.

Ethyl formate [109-94-4] M 74.1, b 54.2°, d 0.921, d³⁰ 0.909, n 1.35994, n²⁵ 1.3565. Free acid or alcohol is removed by standing with anhydrous K₂CO₃, with occasional shaking, then decanting and

distilling from P_2O_5 . Alternatively, the ester can be stood with CaH_2 for several days, then distd from fresh CaH_2 . Cannot be dried with $CaCl_2$ because it reacts rapidly with the ester to form a crystalline compound.

Ethyl gallate [831-61-8] M 198.2, m 150-151°, 163-165°. Recryst from 1,2-dichloroethane, UV: λ_{max} (neutral species) 275nm (ϵ 10 000); (anion) 235nm (ϵ 10 300), 279nm (ϵ 11 400) and 324nm (ϵ 8 500) [Campbell and Coppinger *J Am Chem Soc* 73 2708 1951].

2-Ethyl-1-hexanol [104-76-7] M 130.2, b 184.3°, d 0.833, n 1.431. Dried with sodium, then fractionally distd.

2-Ethylhexyl vinyl ether [37769-62-3, 103-44-6] M 156.3, b 177-178°/atm. Usually contains amines as polymerization inhibitors. These are removed by fractional distn.

Ethyl hydrocupreine hydrochloride (Optochin) [3413-58-9] M 376.9, m 249-251°, pK_1^{25} 5.50, pK_2^{25} 9.95. Recryst from H_2O [UV: Heidt and Forbes *J Am Chem Soc* 55 2701 1933].

Ethyl iodide (iodoethane) [75-03-6] M 156.0, b 72.4°, d 1.933, n^{15} 1.5682, n^{25} 1.5104. Drying with P_2O_5 is unsatisfactory, and with $CaCl_2$ is incomplete. It is probably best to dry with sodium wire and distil [Hammond et al. *J Am Chem Soc* 82 704 1960]. Exposure of ethyl iodide to light leads to rapid decomposition, with the liberation of iodine. Free iodine can be removed by shaking with several portions of dil aq $Na_2S_2O_3$ (until the colour is discharged), followed by washing with water, drying (with $CaCl_2$, then sodium), and distn. The distd ethyl iodide is stored, over mercury, in a dark bottle away from direct sunlight. Other purification procedures include passage through a 60cm column of silica gel, followed by distn; and treatment with elemental bromine, extraction of free halogen with $Na_2S_2O_3$ soln, followed by washing with water, drying and distn. Free iodine and HI have also been removed by direct distn through a LeBel-Henninger column containing copper turnings. Purification by shaking with alkaline solns, and storage over silver, are reported to be unsatisfactory.

Ethyl isobutyrate [97-62-1] M 116.2, b 110°, d 0.867, n 1.388. Washed with aqueous 5% Na_2CO_3 , then with saturated aqueous $CaCl_2$. Dried with $CaSO_4$ and distd.

Ethyl isocyanate [109-90-0] M 71.1, b 559.8°/759mm, 59-61°/atm, 60-63°/atm, d_4^{20} 0.9031, n_D^{20} 1.3808. Fractionate through an efficient column preferably in an inert atmosphere and store in aliquots in sealed tubes [Bieber *J Am Chem Soc* 74 4700 1952; Slocombe et al. *J Am Chem Soc* 72 1888 1950].

3-Ethylisothionicotinamide [10605-12-6] M 166.2, m 164-166°(dec). Crystd from EtOH.

Ethyl isovalerate [108-64-5] M 130.2, b 134.7°, d 0.8664, n 1.39621, n^{25} 1.3975. Washed with aqueous 5% Na_2CO_3 , then saturated aqueous $CaCl_2$. Dried with $CaSO_4$ and distd.

Ethyl levulinate (4-oxopentanoic acid ethyl ester) [539-88-8] M 144.2, m 37.2°, b 106-108°/2mm, 138.8°/8mm, 203-205°/atm, d_4^{20} 1.012, n_D^{20} 1.423. Stir ester with Na_2CO_3 and charcoal, filter and distil. It is freely soluble in H_2O and EtOH [IR, NMR: Sterk *Monatsh Chem* 99 1770 1968; Thomas and Schuette *J Am Chem Soc* 53 2328 1931; Cox and Dodds *J Am Chem Soc* 55 3392 1933].

Ethyl malonate monoamide [7597-56-0] M 131.1, m 47-50°, 49.5-50°, 50°, b 130-135°/2mm. Crystallise from Et_2O or by slow evaporation of an aqueous soln as colourless crystals [Snyder and Elston *J Am Chem Soc* 76 3039 1954; McAlvain and Schroeder *J Am Chem Soc* 71 45 1949; Rising et al. *J Biol Chem* 89 20 1930].

Ethyl methacrylate [97-63-2] M 114.2, b 59°/100mm, d 0.915, n 1.515. Washed successively with 5% aqueous $NaNO_2$, 5% $NaHSO_3$, 5% $NaOH$, then water. Dried with $MgSO_4$, added 0.2% (w/w) of phenyl- β -naphthylamine, and distd through a short Vigreux column [Schultz *J Am Chem Soc* 80 1854 1958].

Ethyl methyl ether [540-67-0] M 60.1, b 10.8°, d⁰ 0.725. Dried with CaSO₄, passed through an alumina column (to remove peroxides), then fractionally distd.

3-Ethyl-2-methyl-2-pentene [19780-67-7] M 112.2, b 114.5°/760mm. Purified by preparative GLC on a column of 20% squalene on Chromosorb P at 70°.

3-Ethyl-4-methylpyridine [529-21-5] M 121.2, b 76°/12mm, 194.5°/750mm, d 0.947, n 1.510, pK_{Est} ~6.3. Dried with solid NaOH, and fractionally distd.

5-Ethyl-2-methylpyridine [104-90-5] M 121.2, b 178.5°/765mm, d 0.919, n 1.497, pK²⁰ 6.51. Purified by conversion to the picrate, crystn, and regeneration of the free base, then distn.

N-Ethylmorpholine [100-74-3] M 115.2, b 138-139°/763mm, d 0.912, n 1.445, pK²⁵ 7.67. Distd twice, then converted by HCl gas into the hydrochloride (extremely deliquescent) which was crystd from anhydrous EtOH-acetone (1:2) [Herries, Mathias and Rabin *Biochem J* 85 127 1962].

Ethyl nitroacetate [626-35-7] M 133.1, b 42-43°/0.2mm, 71-72°/3mm, 93-96°/9mm, 194-195°/atm, d₄²⁰ 1.1953, n_D²⁰ 1.4260, pK²⁵ 5.82. Purified by repeated distn. IR 1748 (CO₂), 1570 and 1337 (NO₂), and 800cm⁻¹ [Hazeldine *J Chem Soc* 2525 1953]. The *hydrazine salt* crystallises from 95% EtOH or MeOH as yellow crystals m 104-105° [Ungnade and Kissinger *J Org Chem* 22 1661 1957, Emmons and Freeman *J Am Chem Soc* 77 4391 1955].

Ethyl p-nitrobenzoate [99-77-4] M 195.2, m 56°. Dissolved in diethyl ether and washed with aqueous alkali, then the ether was evaporated and the solid recrystd from EtOH.

Ethyl orthoformate [122-51-0] M 148.2, b 144°/760mm, d 0.892, n 1.391. Shaken with aqueous 2% NaOH, dried with solid KOH and distd from sodium through a 20cm Vigreux column.

o-Ethylphenol [90-00-6] M 122.2, f 45.1°, b 210-212°, d 1.020, n 1.537, pK²⁵ 10.20. Purified as for p-ethylphenol below.

p-Ethylphenol [123-07-9] M 122.2, m 47-48°, b 218.0°/762mm, n²⁵ 1.5239, pK²⁵ 10.21. Non-acidic impurities were removed by passing steam through a boiling soln containing 1 mole of the phenol and 1.75 moles of NaOH (as aq 10% soln). The residue was cooled and acidified with 30% (v/v) H₂SO₄, and the free phenol was extracted into diethyl ether. The extract was washed with water, dried with CaSO₄ and the ether was evapd. The phenol was distd at 100mm pressure through a **Stedman** gauze-packed column (see p. 441). It was further purified by fractional crystn by partial freezing, and by zone refining, under nitrogen [Biddiscombe et al. *J Chem Soc* 5764 1963]. Alternative purification is *via* the benzoate, as for phenol.

Ethyl phenylacetate [101-97-3] M 164.2, b 99-99.3°/14mm, d 1.030, n 1.499. Shaken with saturated aqueous Na₂CO₃ (three times), aqueous 50% CaCl₂ (twice) and saturated aqueous NaCl (twice). Dried with CaCl₂ and distd under reduced pressure.

3-Ethyl-5-phenylhydantoin (Ethotoin) [86-35-1] M 204.2, m 94°. Crystd from water.

N-Ethyl-5-phenylisoxazolinium-3'-sulfonate [4156-16-5] M 253.3, m 220°(dec). Crystd from diethyl ether or ethyl acetate/pet ether. [Lamas et al. *J Am Chem Soc* 108 5543 1986.]

3-Ethyl-3-phenyl-2,6-piperidinedione (Glutethimide) [77-21-4] M 217.3, m 84°. Crystd from diethyl ether or ethyl acetate/pet ether.

Ethyl propionate [105-37-3] M 102.1, b 99.1°, d 0.891, n¹⁵ 1.38643, n 1.38394. Treated with anhydrous CuSO₄ and distd under nitrogen.

2-Ethylpyridine [100-71-0] M 107.2, b 148.6°, d 0.942, pK^{25} 5.89. Dried with BaO, and fractionally distd. Purified by conversion to the picrate, recrystn and regeneration of the free base followed by distn.

4-Ethylpyridine [536-75-4] M 107.2, b 168.2-168.3°, d 0.942, pK^{25} 6.02. Dried with BaO, and fractionally distd. Also converted to the picrate, recrystd and the free base regenerated and distd.

4-Ethylpyridine-1-oxide [14906-55-9] M 123.1, m 109-110°, $pK_{Est} \sim 1.1$. Crystd from acetone/ether.

Ethyl pyruvate [617-35-6] M 116.1, m -50°, b 44-45°/10mm, 56°/20mm, 69-71°/42mm, 63°/23mm, 155.5°/760mm, d_4^{20} 1.047, n_D^{20} 1.4052. Shake the ester with 10mL portions of satd aq $CaCl_2$ soln (removes ethyl acetate) and the organic layer is removed by centrifugation, decantation and filtration, and is distilled under reduced pressure. Purification of small quantities is carried out *via* the bisulfite adduct: the ester (2.2mL) is shaken with saturated $NaHSO_3$ (3.6mL), chill in a freezing mixture when crystals separate rapidly (particularly if seeded). After 5min EtOH (10mL) is added and the crystals are filtered off, washed with EtOH and Et_2O and dried. Yield *ca* 3g of *bisulfite adduct*. Then treat the adduct (16g) with saturated aqueous $MgSO_4$ (32mL) and 40% formaldehyde (5mL) and shake, whereby the ester separates as an oil which is extracted with Et_2O , the extract is dried ($MgSO_4$), filtered, evapd and the residue is distd (b 56°/20mm), and then redistd (b 147.5°/750mm) to give 5.5g of pure ester. [Cornforth *Org Synth Coll Vol IV* 467 1963.]

Ethyl Red [2-(4-diethylaminophenylazo)benzoic acid] [76058-33-8] M 197.4, m 150-152°, pK_1 2.5, pK_2 9.5. Crystd from EtOH/diethyl ether or toluene. Indicator: pH 4.4 (red) and 6.2 (yellow)

Ethyl stearate [111-61-5] M 312.5, m 33°, b 213-215°/15mm. The solid portion was separated from the partially solid starting material, then crystd twice from EtOH, dried by azeotropic distn with *benzene, and fractionally distd in a spinning-band column at low pressure [Welsh *Trans Faraday Soc* 55 52 1959].

Ethyl thiocyanate (ethyl rhodanide) [542-90-5] M 87.1, b 144-145°, d 1.011, n 1.462. Fractionally distd at atmospheric pressure. (CARE LACHRYMATOR.)

Ethyl thioglycolate [623-51-8] M 120.2, b 50-51°/10mm, 55°/17mm, 62.5-64°/22mm, 67-68°/24mm, 155-158°/atm, d_4^{20} 1.096, n_D^{20} 1.457. Dissolve in Et_2O , wash with H_2O , dry over Na_2SO_4 , filter, evaporate and distil the residue under reduced pressure [Bredereck et al. *Chem Ber* 90 1837 1957]. The Ni complex $[Ni(SCH_2CO_2Et)_2]$ recrystallised twice from EtOH gives crystals which became black when dried in a vacuum over H_2SO_4 , m 104-105° [Dranet and Cefola *J Am Chem Soc* 76 1975 1954].

N-Ethyl thiourea [625-53-6] M 104.2, m 110°. Crystd from EtOH, MeOH or ether.

Ethyl trichloroacetate [515-84-4] M 191.4, b 100-100.5°/30mm, d 1.383. Shaken with saturated aqueous Na_2CO_3 (three times), aqueous 50% $CaCl_2$ (three times), saturated aqueous NaCl (twice), then distd with $CaCl_2$ and distd under reduced pressure.

Ethyl trifluoroacetate [383-63-1] M 142.1, b 61.3°/750, 60-62°/atm, 62-64°/755mm, d_4^{20} 1.191, n_D^{20} 1.30738. Fractionate through a long Vigreux column. IR has ν at 1800 (CO_2) and 1000 (OCO) cm^{-1} [Fuson et al. *J Chem Phys* 20 1627 1952; Bergman *J Org Chem* 23 476 1958].

Ethyl trifluoromethanesulfonate [425-75-2] M 178.1, b 115°/atm, 118-120°/atm, d_4^{20} 1.378, n_D^{20} 1.336. The ester reacts slowly with H_2O and aqueous alkali. If its IR has no OH bands (~ 3000 cm^{-1}) then purify by redistillation. If OH bands are present then dilute with dry Et_2O and shake (carefully) with aqueous $NaHCO_3$ until effervescence ceases, then wash with H_2O and dry ($MgSO_4$), filter, evaporate and distil the residue under slight vacuum then at atmospheric pressure in a N_2 atmosphere. **IT IS A POWERFUL ALKYLATING AGENT, AND THE FUMES ARE VERY TOXIC - CARRY ALL OPERATIONS IN AN EFFICIENT FUME CUPBOARD.** [Gramstad and Hazeldine *J Chem Soc* 173 1956; Howells and McCown *Chem Rev* 77 69 1977.]

S-Ethyl trifluoroacetate [383-64-2] M 158.1, b 88-90°/atm, 90.5°/760mm, d_4^{20} 1.255, n_D^{20} 1.372. If IR is free of OH bands then fractionate, but if OH bands are present then dilute with dry Et₂O, wash with 5% KOH and H₂O, dry over MgSO₄ and fractionate through an efficient column [Hauptschein et al. *J Am Chem Soc* 74 4005 1952]. *Powerful obnoxious odour.*

Ethyl vinyl ether [109-92-2] M 72.1, b 35.5°, d 0.755. Contains polymerization inhibitors (usually amines, e.g. triethanolamine) which can be removed by fractional distn. Redistd from sodium. **LACHRYMATORY.**

1-Ethynyl-1-cyclohexanol [78-27-3] M 124.2, m 30-33°, 32-33°, b 74°/12mm, 76-78°/17mm, 171-172°/694mm, 180°/atm, d_4^{25} 0.9734, n_D^{25} 1.4801. Dissolve in Et₂O, wash with H₂O, dilute NaHCO₃, H₂O again, dry (Na₂SO₄), filter, evaporate and distil the residue. IR (CCl₄): 3448 (OH), 2941 (CH), 1449-1123 and 956 cm⁻¹; NMR (CCl₄) δ: 3.2 (OH), 2.5 (=CH), 1.70 (m 10H, CH₂) [Hasbrouck and Kiessling *J Org Chem* 38 2103 1972].

Ethynyl *p*-tolylsulfone [13894-21-8] M 180.2, m 73-74°. Recrystd from pet ether and dried in vac.

Etiocholane (5β-androsterone) [438-23-3] M 260.5, m 78-80°. Crystd from acetone.

Etiocolanic acid [438-08-4] M 304.5, m 228-229°, pK_{Est} ~4.7. Crystd from glacial acetic acid and sublimes at 160°/0.002mm. The *methyl ester* has m 99-101°. [Weiland et al. *Z Physiol Chem* 161 80 1926.]

Etioporphyrin I [448-71-5] M 478.7, m 360-363°. Crystd from pyridine or CHCl₃-pet ether.

Eucaliptol (1,8-cineol, 1,8-epoxy-*p*-menthane, 1,3,3-trimethyl-2-oxabicyclo[2.2.2]-octane) [470-82-6] M 154.2, m 1.3°, 1.5°, b 39-39.3°/4mm, 176-176.4°/760mm, d_4^{20} 0.9232, n_D^{20} 1.4575. Purified by dilution with an equal volume of pet ether, then saturated with dry HBr. The ppt was filtered off, washed with small vols of pet ether, then cineole was regenerated by stirring the crystals with H₂O. It can also be purified *via* its *o*-cresol or resorcinol addition compds. Stored over Na until required. Purified by fractional distn. Insoluble in H₂O but soluble in organic solvents. [IR: Kome et al. *Nippon Kagaku Zasshi [J Chem Soc Japan (Pure Chem Sect)]* 80 66 1959; *Chem Abstr* 603 1961.]

Eugenol (4-allyl-2-methoxyphenol) [97-53-0] M 164.2, b 253°/760mm, 255°/760mm, d_4^{20} 1.066, n_D^{20} 1.540, pK²⁵ 10.19. Fractional distn gives a pale yellow liquid which darkens and thickens on in air. Should store under N₂ at -20°. [Waterman and Priedster *Recl Trav Chim Pays-Bas* 48 1272 1929.]

Eugenol methyl ether (4-allyl-1,2-dimethoxybenzene) [93-15-2] M 178.2, m -4°, b 127-129°/11mm, 146°/30mm, 154.7°/760mm, d_4^{20} 1.0354, n_D^{20} 1.53411. Recrystd from hexane at low temp and redistd (preferably *in vacuo*). [Hillmer and Schorning *Z Phys Chem [A]* 167 407 1934; Briner and Fliszár *Helv Chim Acta* 42 2063 1959.]

(+)-α-Fenchol (1R-1,3,3-trimethyl-norbornan-2-ol) [1632-73-1] M 154.3, m 40-43°, 47-47.5°, b 201-202°, [α]_D²⁰ +12.5° (c 10, EtOH). It is prepared by reduction of (-)-fenchone and is purified by recrystallisation from *C₆H₆-pet ether, or distn, or both. The 2-carboxybenzoyl (*monophthalate*) derivative has m 146.5-147.5° [α]_D²⁰ -20.4° (EtOH), and the 2-phenylurethane has m 81°. [Beckmann and Metzger *Chem Ber* 89 2738 1956].

(+)-Fenchone (1S-1,3,3-trimethyl-norbornan-2-one) [4695-62-9] M 152.2, m 5-7°, 6.1°, b 63-65°/13mm, 66°/15mm, 122°/10mm, d_4^{20} 0.9434, n_D^{20} 1.4636, [α]_D²⁰ +66.9° (neat, or in c 1.5, EtOH), [α]₅₄₆²⁰ +60.4° (neat). The oily liquid is purified by distn in a vacuum, and is very soluble in EtOH and Et₂O. [Boyle et al. *J Chem Soc, Chem Commun* 395 1971, Hüchel *Justus Liebigs Ann Chem* 549 186 1941; (±)-isomer: Braun and Jacob *Chem Ber* 66 1461 1933.] It forms two *oximes*; *cis-oxime*: m

167° (cryst from pet ether) $[\alpha]_D^{20} +46.5^\circ$ (c 2, EtOH), *O*-benzoyloxime **m** 81° $[\alpha]_D^{20} +49^\circ$ (EtOH) and *oxime-HCl* **m** 136° (dec). The *trans*-oxime has **m** 123° (cryst from pet ether) $[\alpha]_D^{18} +148^\circ$ (c 2, EtOH) and the *O*-benzoyloxime has **m** 125° $[\alpha]_D^{20} +128.5^\circ$ (c 2, EtOH) [Hückel *Justus Liebigs Ann Chem* **549** 186 1941; Hückel and Sachs *Justus Liebigs Ann Chem* **498** 166 1932].

(-)- **Fenchone** (1*R*-1,3,3-trimethyl-norbornan-2-one) [7787-20-4] **M** 152.2, **m** 5.2°, **b** 67.2°/10mm, 191-195°/atm, d_4^{20} 0.9484, n_D^{20} 1.4630, $[\alpha]_D^{20} -66.8^\circ$ (neat). Purification as for the (+)-enantiomer above and should have the same physical properties except for the optical rotations. UV: λ_{max} 285nm (ϵ 12.29). [Braun and Jacob *Chem Ber* **66** 1461 1933; UV: Ohloff et al. *Chem Ber* **90** 106 1957.]

Flavone (2-phenyl-4*H*-1-benzopyran-4-one) [525-82-6] **M** 222.3, **m** 100°. Crystd from pet ether.

Fluoranthene (benzo[*j,k*]fluorene) [206-44-0] **M** 202.3, **m** 110-111°. Purified by chromatography of CCl₄ solns on alumina, with *benzene as eluent. Crystd from EtOH, MeOH or *benzene. Purified by zone melting. [Gorman et al. *J Am Chem Soc* **107** 4404 1985.]

2-Fluorenamine [153-78-6] **M** 181.2. See 2-aminofluorene on p. 106.

9-Fluorenamine [525-03-1] **M** 181.2, **m** 64-65°, $pK_{Est} \sim 3.5$. Crystd from hexane.

Fluorene [86-73-7] **M** 166.2, **m** 114.7-115.1°, **b** 160°/15mm. Purified by chromatography of CCl₄ or pet ether (b 40-60°) soln on alumina, with *benzene as eluent. Crystd from 95% EtOH, 90% acetic acid and again from EtOH. Crystn using glacial acetic acid retained an impurity which was removed by partial mercuration and pptn with LiBr [Brown, Dubeck and Goldman *J Am Chem Soc* **84** 1229 1962]. Has also been crystd from hexane, or *benzene/EtOH, distd under vacuum and purified by zone refining. [Gorman et al. *J Am Chem Soc* **107** 4404 1985.]

9-Fluorenone [486-25-9] **M** 180.2, **m** 82.5-83.0°, 85-86°, **b** 341°/760mm. Crystd from absolute EtOH, MeOH or *benzene/pentane. [Ikezawa *J Am Chem Soc* **108** 1589 1986.] Also twice recrystd from toluene and sublimed in a vac [Saltiel *J Am Chem Soc* **108** 2674 1986]. Can be distd under high vacuum.

Fluorene-2,7-diamine [525-64-4] **M** 196.3, **m** 165-166°. Crystd from hot H₂O or aq EtOH, dried in a vac and stored in the dark.

9-Fluorenylmethyl chloroformate (Fmoc-Cl) [28920-43-6] **M** 258.7, **m** 61-63°, 61.4-63°. The IR should contain no OH bands (at ~ 3000 cm⁻¹) due to the hydrolysis product 9-fluorenylmethanol. Purify by recrystn from dry Et₂O. IR (CHCl₃) has band at 1770 cm⁻¹ (C=O) and the NMR (CDCl₃) has δ at 4-4.6 (m 2H, CHCH₂) and 7.1-7.8 (m, 8 aromatic H). The *azide* (Fmoc-N₃) has **m** 89-90° (from hexane) and IR (CHCl₃) at 2135 (N₃) and 1730 (C=O) cm⁻¹; and the *carbazate* (Fmoc-NHNH₂) has **m** 171° dec (from nitromethane), IR (KBr) 3310, 3202 (NH) and 1686 (CONH) cm⁻¹. [Caprino and Han *J Org Chem* **37**, 3404 1972 and *J Am Chem Soc* **92** 5748 1970; Koole et al. *J Org Chem* **59** 1657 1989; Fürst et al. *J Chromatogr* **499** 537 1990.]

9-Fluorenylmethyl succinimidyl carbonate [82911-69-1] **M** 337.3, **m** 147-151° (dec), 151° (dec). Recrystd from CHCl₃-Et₂O, or from pet ether (b 40-60°). [Pauet *Can J Chem* **60** 976 1982; Lapatsaris et al. *Synthesis* 671 1983.]

Fluorescein [9-(*o*-carboxyphenyl-6-hydroxy-3*H*-xanthene-3-one)] [2321-07-5] **M** 320.0, ϵ_{495nm} 7.84×10^4 (in 10⁻³M NaOH), pK_1 2.2, pK_2 4.4, pK_3 6.7. Dissolved in dilute aqueous NaOH, filtered and ppted by adding dilute (1:1) HCl. The process was repeated twice more and the fluorescein was dried at 100°. Alternatively, it has been crystd from acetone by allowing the soln to evaporate at 37° in an open beaker. Also recrystd from EtOH and dried in a vacuum oven.

Fluoresceinamine (mixture of 5- and 6-aminofluorescein) [27599-63-9] **M** 347.3, **m** 314-316° (dec, 5-amino) and **m** >200° (dec). Dissolve in EtOH, treat with charcoal, filter, evaporate and dry

residue in vacuum at 100° overnight. Also recrystallise from 6% HCl, then dissolve in 0.5% aqueous NaOH and ppt by acidifying with acetic acid. The separate amines are made from the respective nitro compounds which are best separated *via* their acetate salts. They have similar R_F of 0.26 on Silica Gel Merck F₂₅₄ in 5 mL MeOH + 150 mL Et₂O satd with H₂O. IR (Me₂SO) has a band at 1690 cm⁻¹ (CO₂⁻) and sometimes a weak band at 1750 cm⁻¹ due to lactone. UV (EtOH) of 6-isomer λ_{max} 222 (ϵ 60 000) and 5-isomer λ_{max} 222 (ϵ 60 000) and 285 (ϵ 20.600). [IR: McKinney and Churchill *J Chem Soc (C)* 654 1970; McKinney et al. *J Org Chem* 27 3986 1962; UV: Verbiscar *J Org Chem* 29 490 1964.]

Fluorescein isothiocyanate isomer I (5-isocyanato isomer) [3326-32-7; 27072-45-3 *mixture of 5- and 6-isomers*] **M 389.4, m >160° (slow dec)**. It is made from the pure 5-amino isomer. Purified by dissolving in boiling Me₂CO, filtering and adding pet ether (b 60-70°) until it becomes turbid. If an oil separates then decant and add more pet ether to the supernatant and cool. Orange-yellow crystals separate, collect and dry *in vacuo*. Should give one spot on TLC (silica gel) in EtOAc, pyridine, AcOH (50:1:1) and in Me₂NCHO, CHCl₃, 28% N₄OH (10:5:4). IR (Me₂SO): 2110 (NCS) and 1760 (C=O). The NMR spectra in Me₂CO-d₆ of the 5- and 6-isomers are distinctly different for the protons in the *benzene ring; the UV in phosphate buffer pH 8.0 shows a max at ~490nm. [Sinsheimer et al. *Anal Biochem* 57 227 1974; McKinney et al. *Anal Biochem* 7 74 1964.]

Fluoroacetamide [640-19-7] **M 77.1, m 108°**. Crystd from chloroform.

Fluorobenzene [462-06-6] **M 96.1, b 84.8°, d 1.025, n 1.46573, n³⁰ 1.4610**. Dried for several days with P₂O₅, then fractionally distd.

***o*-Fluorobenzoic acid** [445-29-4] **M 140.1, m 127°, pK²⁵ 3.27**. Crystd from 50% aqueous EtOH, then zone melted or vacuum sublimed at 130-140°.

***m*-Fluorobenzoic acid** [445-38-9] **M 140.1, m 124°, pK²⁵ 3.86**. Crystd from 50% aqueous EtOH, then vacuum sublimed at 130-140°.

***p*-Fluorobenzoic acid** [456-22-4] **M 140.1, m 182°, pK²⁵ 4.15**. Crystd from 50% aqueous EtOH, then zone melted or vacuum sublimed at 130-140°.

3-Fluoro-4-hydroxyphenylacetic acid [458-09-3] **M 170.1, m 33°, pK_{Est(1)}~4.4, pK_{Est(2)}~9.4**. Crystd from water.

1-Fluoro-4-nitrobenzene [350-46-9] **M 141.1, m 27° (stable form), 21.5° (unstable form), b 205.3°/735mm, 95-97.5°/22mm, 86.6°/14mm**. Crystd from EtOH.

1-Fluoro-4-nitronaphthalene [341-92-4] **M 191.2, m 80°**. Recrystd from EtOH as yellow needles [Bunce et al. *J Org Chem* 52 4214 1987].

***o*-Fluorophenol** [367-12-4] **M 112.1, m 16°, b 53°/14mm, d 1.257, n 1.514, pK²⁵ 8.70**. Passed at least twice through a gas chromatographic column for small quantities, or fractionally distd under reduced pressure.

***p*-Fluorophenoxyacetic acid** [405-79-8] **M 170.1, m 106°, pK²⁵ 3.13**. Crystd from EtOH.

4-Fluorophenylacetic acid [405-50-5] **M 154.1, m 86°, pK²⁵ 4.22**. Crystd from heptane.

4-Fluorophenyl isocyanate [1195-45-5] **M 137.1, b 55°/8mm, n_D²⁰ 1.514**. Purify by repeated fractionation through an efficient column. If IR indicated that there is too much urea (in the presence of moisture the symmetrical urea is formed) then dissolve in dry EtOH-free CHCl₃, filter, evaporate and distil. **It is a pungent LACHRYMATORY liquid**. [see Hardy *J Chem Soc* 2011 1934; and Hickinbottom *Reactions of Organic Compounds* Longmans p. 493 1957.]

4-Fluorophenyl isothiocyanate [1544-68-9] M 153.2, m 24-26°, 26-27°, b 66°/2 mm, 215°/atm, 228°/760mm, n_D^{20} 1.6116. Likely impurity is the symmetrical thiourea. Dissolve the isothiocyanate in dry CHCl_3 , filter and distil the residue in a vacuum. It can also be steam distd, the oily layer separated, dried over CaCl_2 and distilled *in vacuo*. *Bis-(4-fluorophenyl)thiourea* has m 145° (from aq EtOH). [Browne and Dyson *J Chem Soc* 3285 1931; Buu Hoi et al. *J Chem Soc* 1573 1955; Olander *Org Synth Coll Vol I* 448 1941].

***p*-Fluorophenyl-*o*-nitrophenyl ether** [448-37-3] M 247.2, m 62°. Crystd from EtOH.

***o*-Fluorotoluene** [95-52-3] M 110.1, b 114.4°, d 1.005, n 1.475. Dried with P_2O_5 or CaSO_4 and fractionally distd through a silvered vacuum-jacketed glass column with 1/8th-in glass helices. A high reflux ratio is necessary because of the closeness of the boiling points of the *o*-, *m*- and *p*- isomers [Potter and Saylor *J Am Chem Soc* 37 90 1915].

***m*-Fluorotoluene** [352-70-5] M 110.1, b 116.5°, d 1.00, n_D^{27} 1.46524. Purification as for *o*-fluorotoluene.

***p*-Fluorotoluene** [352-32-9] M 116.0°, d 1.00, n 1.46884. Purification as for *o*-fluorotoluene.

Formaldehyde [50-00-0] M 30.0, m 92°, b -79.6°/20mm, d_4^{20} 0.815, pK^{25} 13.27 (hydrate). Commonly contains added MeOH. Addition of KOH soln (1 mole KOH: 100 moles HCHO) to 40% formaldehyde soln, or evaporation to dryness, gives paraformaldehyde polymer which, after washing with water, is dried in a vacuum desiccator over P_2O_5 or H_2SO_4 . Formaldehyde is regenerated by heating the paraformaldehyde to 120° under vacuum, or by decomposing it with barium peroxide. The monomer, a gas, is passed through a glass-wool filter cooled to -48° in CaCl_2 /ice mixture to remove particles of polymer, then dried by passage over P_2O_5 and either condensed in a bulb immersed in liquid nitrogen or absorbed in ice-cold conductivity water.

Formaldehyde dimethyl acetal (dimethoxymethane, methylal, formal) [109-87-5] M 76.1, m -108°, b 41-42°/736mm, 41-43°/atm, 42-46°/atm, d_4^{20} 0.8608, n_D^{20} 1.35335. It is a volatile flammable liquid which is soluble in three parts of H_2O , and is readily hydrolysed by acids. Purify by shaking with an equal vol of 20% aq NaOH, stand for 20min, dry over fused CaCl_2 , filter and fractionally distil through an efficient column, store over molecular sieves. [Buchler et al. *Org Synth Coll Vol III* 469 1955; *Ind Eng Chem* 18 1092 1926; Rambaud and Besserre *Bull Soc Chim Fr* 45 1955; IR: *Can J Chem* 36 285 1958.]

Formaldehyde dimethyl mercaptal (bis-[methylthio]methane) [1618-26-4] M 108.2, b 44-47°/13mm, 45.5°/18mm, 148-149°/atm, d_4^{20} 1.0594, n_D^{20} 1.5322. Work in an efficient fume cupboard as the substance may contain traces (or more) of methylmercaptan which has a very bad odour. Dissolve in Et_2O , shake with aqueous alkalis then dry over anhydrous K_2CO_3 , filter and distil over K_2CO_3 under a stream of N_2 . If the odour is very strong then allow all gas effluents to bubble through 5% aqueous NaOH soln which is then treated with dilute KMnO_4 in order to oxidise MeSH to odourless products. UV: λ_{max} 238 nm ($\log \epsilon$ 2.73) [Fehnel and Carmack *J Am Chem Soc* 71 90 1949; Fehér and Vogelbruch *Chem Ber* 91 996 1958; Bøhme and Marz *Chem Ber* 74 1672 1941]. Oxidation with aq KMnO_4 yields *bis-(methylsulfonyl)methane* which has m 142-143° [Fiecchi et al. *Tetrahedron Lett* 1681 1967].

Formamide [75-12-7] M 45.0, f 2.6°, b 103°/9mm, 210.5°/760mm(dec), d 1.13, n 1.44754, n_D^{25} 1.44682. Formamide is easily hydrolysed by acids and bases. It also reacts with peroxides, acid halides, acid anhydrides, esters and (on heating) alcohols; while strong dehydrating agents convert it to a nitrile. It is very *hygroscopic*. Commercial material often contains acids and ammonium formate. Vorhoek [*J Am Chem Soc* 58 2577 1956] added some bromothymol blue to formamide and then neutralised it with NaOH before heating to 80-90° under reduced pressure to distil off ammonia and water. The amide was again neutralised and the process was repeated until the liquid remained neutral on heating. Sodium formate was added, and the formamide was reduced under reduced pressure at 80-90°. The distillate was again neutralised and redistd. It was then fractionally crystd in the absence of CO_2 and water by partial freezing.

Formamide (specific conductance $2 \times 10^{-7} \text{ ohm}^{-1} \text{ cm}^{-1}$) of low water content was dried by passage through a column of 3A molecular sieves, then deionized by treatment with a mixed-bed ion-exchange resin loaded with H^+ and HCONH^- ions (using sodium formamide in formamide) [Notley and Spiro *J Chem Soc (B)* 362 1966].

Formamidine acetate [3473-63-0] **M 104.1, m 159-161°(dec), 164°(dec), $\text{pK}_{\text{Est}} \sim 12$** . Unlike the hydrochloride, the acetate salt is not hygroscopic. It is recrystd from a small volume of acetic acid, by addition of EtOH and the crystals are washed with EtOH then Et₂O and dried in a vac. [Taylor, Ehrhart and Karanisi *Org Synth* 46 39 1966.]

Formamidine sulfinic acid (thiourea-S-dioxide) [1758-73-2] **M 108.1, m 124-126°(dec)**. Dissolved in five parts of aq 1:1% NaHSO₃ at 60-63° (charcoal), then crystd slowly, with agitation, at 10°. Filtered. Dried immediately at 60° [Koniecki and Linch *Anal Chem* 30 1134 1958].

Formanilide [103-70-8] **M 121.1, m 50°, b 166°/14mm, 216°/120mm, d 1.14**. Crystd from ligroin/xylene.

Formic acid [64-18-6] **M 46.0 (anhydr), f 8.3°, b 25°/40mm, 100.7°/760mm, d 1.22, n 1.37140, $n^{25} 1.36938$, $\text{pK}^{25} 3.74$** . Anhydrous formic acid can be obtained by direct fractional distillation under reduced pressure, the receiver being cooled in ice-water. The use of P₂O₅ or CaCl₂ as dehydrating agents is unsatisfactory. Reagent grade 88% formic acid can be satisfactorily dried by refluxing with phthalic anhydride for 6h and then distilling. Alternatively, if it is left in contact with freshly prepared anhydrous CuSO₄ for several days about one half of the water is removed from 88% formic acid: distn removes the remainder. Boric anhydride (prepared by melting boric acid in an oven at a high temperature, cooling in a desiccator, and powdering) is a suitable dehydrating agent for 98% formic acid; after prolonged stirring with the anhydride the formic acid is distd under vacuum. Formic acid can be further purified by fractional crystn using partial freezing.

Forskolin (5-[acetyloxy]-3-ethenyldodecahydro-6,10,10b-trihydroxy-3,4a,7,7,10a-pentamethyl-[3R-{3 α -4 $\alpha\beta$, 5 β , 6 β , 6 $\alpha\alpha$, 10 α , 10 $\alpha\beta$, 10 $\beta\alpha$ }-1H-naphtho[2,1-b]pyran-1-one) [66575-29-9] **M 410.5, m 229-232°, 228-233°**. Recrystd from *C₆H₆-pet ether. It is antihypertensive, positive inotropic, platelet aggregation inhibitory and adenylate cyclase activating properties [*Chem Abstr* 89 1978 244150, de Souza et al. *Med Res Rev* 3 201 1983].

D(-)-Fructose [57-48-7] **M 180.2, m 103-106°, $[\alpha]_{546}^{20} -190^\circ$ (after 1h, c 10, H₂O), $\text{pK}^{25} 12.03$** . Dissolved in an equal weight of water (charcoal, previously washed with water to remove any soluble material), filtered and evaporated under reduced pressure at 45-50° to give a syrup containing 90% of fructose. After cooling to 40°, the syrup was seeded and kept at this temperature for 20-30h with occasional stirring. The crystals were removed by centrifugation, washed with a small quantity of water and dried to constant weight under a vacuum over conc H₂SO₄. For higher purity, this material was recrystd from 50% aqueous ethanol [Tsuzuki, Yamazaki and Kagami *J Am Chem Soc* 72 1071 1950].

D(+)-Fucose [3615-37-0] **M 164.2, m 144°, $[\alpha]_{546}^{20} +89^\circ$ (after 24h, c 10 in H₂O)**. Crystd from EtOH.

Fullerene C₆₀ (Buckminsterfullerene C₆₀, Footballene, Buckyball 60) [99685-96-8] **M 720.66 and Fullerene C₇₀** [115383-22-7] **M 840.77**. Purified from the soluble toluene extract (400mg) of the soot (Fullerite) formed from resistive heating of graphite by adsorption on neutral alumina (100g; Brockmann I; 60 x 8cm). Elution with toluene-hexane (5:95 v/v) gives ca 250mg of quite pure C₆₀. It has characteristic spectral properties (see below). Further elution with toluene-hexane (20:80 v/v; i.e. increased polarity of solvent) provides 50mg of "pure" C₇₀ [*J Am Chem Soc* 113 1050 1991].

Chromatography on alumina can be improved by using conditions which favour adsorption rather than crystn. Thus the residue from toluene extraction (1g) in CS₂ (ca 300mL) is adsorbed on alumina (375g, standard grade, neutral ca 150 mesh, Brockmann I) and loaded as a slurry in toluene-hexanes (5:95 v/v) to a 50 x 8cm column of alumina (1.5Kg) in the same solvent. To avoid crystn of the fullerenes, 10% of toluene in hexanes is added quickly followed by 5% of toluene in hexanes after the fullerenes had left the loading fraction (2-3h). With a flow rate of 15mL/min the purple C₆₀ fraction is eluted during a 3-4h period. Evapn of the eluates gives 550-

630mg of product which, after recrystn from CS₂-cyclohexane yields 520-600mg of C₆₀ which contains adsorbed solvent. On drying at 275°/10⁻³mm for 48h a 2% weight loss is observed although the C₆₀ still contains traces of solvent. Further elution of the column with 20% of toluene in hexanes provides 130mg of C₇₀ containing 10-14% of C₆₀ (by ¹³C NMR). This was rechromatographed as above using a half scale column and adsorbing the 130mg in CS₂ (20mL) on alumina (24g) and gave 105mg of recrystd C₇₀ (containing 2% of C₆₀). The purity of C₆₀ can be improved further by washing the crystalline product with Et₂O and Me₂CO followed by recrystn from *C₆H₆ and vacuum drying at high temperatures. [*J Chem Soc, Chem Commun* 956 1992.]

Carbon soot from resistive heating of a carbon rod in a partial helium atmosphere (0.3bar) under specified conditions is extracted with boiling *C₆H₆ or toluene, filtered and the red-brown soln evapd to give crystalline material in 14% yield which is mainly a mixture of fullerenes C₆₀ and C₇₀. Chromatographic filtration of the 'crude' mixture with *C₆H₆ allows no separation of components, but some separation was observed on silica gel TLC with *n*-hexane or *n*-pentane, but not cyclohexane. Analytical HPLC with hexanes (5µm Econosphere silica) gave satisfactory separation of C₆₀ and C₇₀ (retention times of 6.64 and 6.93min respectively) at a flow rate of 0.5mL/min and using a detector at 256nm. HPLC indicated the presence of minor (<1.5% of total mass) unidentified C_n with species (retention times of 5.86 and 8.31min. Column chromatography on flash silica gel with hexanes gives a few fractions of C₆₀ with ≥95% purity but later fractions contain mixtures of C₆₀ and C₇₀. These can be obtained in 99.85 and >99% purity respectively by column chromatography on neutral alumina. [*J Phys Chem* 94 8630 1990].

Separation of C₆₀ and C₇₀ can be achieved by HPLC on a dinitroanilinopropyl (DNAP) silica (5µm pore size, 300Å pore diameter) column with a gradient from *n*-hexane to 50% CH₂Cl₂ using a diode array detector at wavelengths 330nm (for C₆₀) and 384nm (for C₇₀). [*J Am Chem Soc* 113, 2940, 1991.]

Soxhlet extraction of the "soot" is a good preliminary procedure, or if material of only *ca* 98% purity is required. Soxhlet extraction with toluene is run (20min per cycle) until colourless solvent filled the upper part of the Soxhlet equipment (10h). One third of the toluene remained in the pot. After cooling, the solution was filtered through a glass frit. This solid (purple in toluene) was *ca* 98% C₆₀. This powder was again extracted in a Soxhlet using identical conditions as before and the C₆₀ was recrystd from toluene to give 99.5% pure C₆₀. C₇₀ has greater affinity than C_{h60} for toluene. [*J Chem Soc, Chem Commun* 1402 1992.]

Purification of C₆₀ from a C₆₀/C₇₀ mixture was achieved by dissolving in an aqueous soln of γ (but not β) cyclodextrin (0.02M) upon refluxing. The rate of dissolution (as can be followed by UV spectra) is quite slow and constant up to 10⁻⁵M of C₆₀. The highest concn of C₆₀ in H₂O obtained was 8 x 10⁻⁵M and a 2 γ-cyclodextrin:1 C₆₀ clathrate is obtained. C₆₀ is extracted from this aqueous soln by toluene and C₆₀ of >99 purity is obtained by evaporation. With excess of γ-cyclodextrin more C₆₀ dissolves and the complex precipitates. The ppte is insol in cold H₂O but sol in boiling H₂O to give a yellow soln. [*J Chem Soc, Chem Commun* 604 1922.]

C₆₀ and C₇₀ can also be readily purified by inclusion complexes with *p*-*tert*-butylcalix[6] and [8]arenes. Fresh carbon-arc soot (7.5g) is stirred with toluene (250mL) for 1h and filtered. To the filtrate is added *p*-*tert*-butylcalix[8]arene, refluxed for 10min and filtered. The filtrate is seeded and set aside overnight at 20°. The C₆₀ complex separated as yellow-brown plates and recrystd twice from toluene (1g from 80mL), 90% yield. Addition of CHCl₃ (5mL) to the complex (0.85g) gave C₆₀ (0.28g, 92% from recryst complex).

p-*tert*-Butylcalix[6]arene-(C₆₀)₂ complex is prepared by adding to a refluxing soln of C₆₀ (5mg) in toluene (5mL), *p*-*tert*-butylcalix[6]arene (4.4mg). The hot soln was filtered rapidly and cooled overnight to give prisms (5.5mg, 77% yield). Pure C₆₀ is obtained by decomposing the complex with CHCl₃ as above.

The *p*-*tert*-butylcalix[6]arene-(C₇₀)₂ complex is obtained by adding *p*-*tert*-butylcalix[6]arene (5.8mg) to a refluxing soln of C₇₀ (5mg) in toluene (2mL), filtering hot and slowly cooling. to give red-brown needles (2.5mg, 31% yield) of the complex. Pure C₇₀ is then obtained by decomposing the complex with CHCl₃.

Decomposition of these complexes can also be achieved by boiling a toluene soln over KOH pellets for *ca* 10min. The calixarenes form Na salts which do not complex with the fullerenes. These appear to be the most satisfactory means at present for preparing large quantities of relatively pure fullerene C₆₀ and C₇₀ and is considerably cheaper than previous methods. [*Nature* 368 229 1994.]

Repeated chromatography on neutral alumina yields minor quantities of solid samples of C₇₆, C₈₄, C₉₀ and C₉₄ believed to be higher fullerenes. A stable oxide C₇₀O has been identified. Chromatographic procedures for the separation of these compounds are reported. [*Science* 252 548 1991.]

Physical properties of Fullerene C₆₀: It does not melt below 360°, and starts to sublime at 300° *in vacuo*. It is a mustard coloured solid that appears brown or black with increasing film thickness. It is soluble in common organic solvents, particularly aromatic hydrocarbons which give a beautiful magenta colour. Toluene solutions are purple in colour. Sol in *C₆H₆ (5mg/mL), but dissolves slowly. Crysts of C₆₀ are both needles and plates. UV-Vis in hexanes: λ_{max} nm(log ε) 211(5.17), 227sh(4.91), 256(5.24), 328(4.71), 357sh(4.08), 368sh(3.91), 376sh(3.75), 390(3.52), 395sh(3.30), 403(3.48), 407(3.78), 492sh(2.72) < 540(2.85), 568(2.78), 590(2.86), 598(2.87) and 620(2.60). IR (KBr): ν 1429m, 1182m, 724m, 576m and 527s cm⁻¹. ¹³C NMR: one signal at 142.68ppm.

Physical properties of Fullerene C₇₀: It does not melt below 360°, and starts to sublime at 350° *in vacuo*. A reddish-brown solid, greenish black in thicker films. Solns are port-wine red in colour. Mixtures of C₆₀ and C₇₀ are red due to C₇₀ being more intensely coloured. It is less soluble than C₆₀ in *C₆H₆ but also dissolves slowly. C₇₀ gives orange coloured soln in toluene. Drying at 200-250° is not sufficient to remove All solvent. Samples need to be sublimed to be free from solvent.

UV-Vis in hexanes: λ_{max} nm(log ε) 214(5.05), 235(5.06), 249sh(4.95), 268sh(4.78), 313(4.23), 330(4.38), 359(4.29), 377(4.45), 468(4.16), 542(3.78), 590sh(3.47), 599sh(3.38), 609(3.32), 623sh(3.09), 635sh(3.13) and 646sh(2.80).

IR (KBr): ν 1430m, 1428m, 1420m, 1413m, 1133mw, 1087w, 795s, 674ms, 642ms, 5778s, 566m, 535ms and 458m cm⁻¹.

¹³C NMR [run in the presence of Cr(pentan-2,4-dione)₃ which induces a *ca* 0.12ppm in the spectrum]: Five signals at 150.07, 147.52, 146.82, 144.77 and 130.28ppm, unaffected by proton decoupling.

[Further reading: Kroto, Fischer and Cox *Fullerenes* Pergamon Press, Oxford 1993 ISBN 0080421520; Kadish and Ruoff (Eds) *Fullerenes: Recent Advances in the Chemistry and Physics of Fullerenes and Related Materials* The Electrochemical Soc. Inc, Pennington, NJ, 1994 ISBN 1566770823]

Fumagillin {2,4,6,8-decatetraene-1,10-dioic acid mono[4-(1,2-epoxy-1,5-dimethyl-4-hexenyl)-5-methoxy-1-oxaspiro[2.5]oct-6-yl] ester} [23110-15-8] M 458.5, m 194-195°, [α]_D²⁰ -26.2° (in 95% EtOH), pK_{Est} ~4.5. Forty grams of a commercial sample containing 42% fumagillin, 45% sucrose, 10% antifoam agent and 3% of other impurities were digested with 150mL of CHCl₃. The insoluble sucrose was filtered off and washed with CHCl₃. The combined CHCl₃ extracts were evap almost to dryness at room temperature under reduced pressure. The residue was triturated with 20mL of MeOH and the fumagillin was filtered off by suction. It was crystd twice from 500mL of hot MeOH by standing overnight in a refrigerator (yellow needles). (The long chain fatty ester used as antifoam agent was still present, but was then removed by repeated digestion, on a steam bath, with 100mL of diethyl ether.) For further purification, the fumagillin (10g) was dissolved in 150mL of 0.2M ammonia, and the insoluble residue was filtered off. The ammonia soln (cooled in running cold water) was then brought to pH 4 by careful addn of M HCl with constant shaking in the presence of 150mL of CHCl₃. (Fumagillin is acid-labile and must be removed rapidly from the aq acid soln.) The CHCl₃ extract was washed several times with distd water, dried (Na₂SO₄) and evaporated under reduced pressure. The solid residue was washed with 20mL of MeOH. The fumagillin was filtered by suction, then crystd from 200mL of hot MeOH. [Tarbell et al. *J Am Chem Soc* 77 5610 1955.]

Alternatively, 10g of fumagillin in 100mL CHCl₃ was passed through a silica gel (5g) column to remove tarry material, and the CHCl₃ was evaporated to leave an oil which gave fumagillin on crystn from amyl acetate. It recrystallises from MeOH (charcoal). The fumagillin was stored in dark bottles in the absence of oxygen and at low temperatures. [Schenk, Hargie and Isarasena *J Am Chem Soc* 77 5606 1955.]

Fumaraldehyde bis-(dimethyl acetal) (trans-1,1,4,4-tetramethoxybut-2-ene) [6068-62-8] M 176.2, b 100-103°/15mm, 101-103°/25mm, d₄²⁰ 1.011, n_D²⁰ 1.425. Dry over fused CaCl₂ and dist *in vacuo*. The maleic (cis) isomer has b 112°/11mm, and d₄²³ 0.932 and n_D²⁵ 1.4243. [Zeik and Heusner *Chem Ber* 90 1869 1957; Clauson-Kaas et al. *Acta Chem Scand* 9 111 1955; Clauson-Kaas *Acta Chem Scand* 6 569 1952.]

Fumaric (trans-but-2-ene-1,4-dioic) acid [110-17-8] M 116.1, m 289.5-291.5° (sealed tube), pK₁²⁵ 3.10, pK₂²⁵ 4.60 (4.38). Crystd from hot M HCl or water. Dried at 100°.

Furan [110-00-9] M 68.1, b 31.3°, d 1.42, n 1.4214. Shaken with aqueous 5% KOH, dried with CaSO₄ or Na₂SO₄, then distd under nitrogen, from KOH or sodium, immediately before use. A trace of hydroquinone could be added as an inhibitor of oxidation.

3-(2-Furanyl)acrylic acid [539-47-9] M 138.1, m 141°, pK_{Est} ~3.8. Crystd from H₂O or pet ether (b 80-100°)(charcoal).

Furan-2-carboxylic (2-furoic) acid [88-14-2] M 112.1, m 133-134°, b 141-144°/20mm, 230-232°/760mm, pK₁²⁵ -7.3 (O-protonation), pK₂²⁵ 3.32. Crystd from hot water (charcoal), dried at 120° for 2h, then recrystd from CHCl₃, and again dried at 120° for 2h. For use as a standard in volumetric analysis, good quality commercial acid should be crystd from CHCl₃ and dried as above or sublimed at 130-140° at 50-60mm or less.

Furan-3-carboxylic (3-furoic) acid [488-93-7] M 112.1, m 122-123°, pK²⁵ 4.03 Crystd from water.

Furan-3,4-dicarboxylic acid [3387-26-6] M 156.1, m 217-218° pK₁²⁵ 1.44, pK₂²⁵ 7.84. Crystd from water.

Furfural (2-furfuraldehyde) [98-01-1] M 96.1, b 54-56°/11mm, 59-60°/15mm, 67.8°/20mm, 90°/65mm, 161°/760mm, d₄²⁰ 1.159, n_D²⁰ 1.52608, pK -6.5 (O-protonation). Unstable to air, light and acids. Impurities include formic acid, β-formylacrylic acid and furan-2-carboxylic acid. Distd over an oil bath from 7% (w/w) Na₂CO₃ (added to neutralise acids, especially pyromucic acid). Redistd from 2% (w/w) Na₂CO₃, and then, finally fractionally distd under vacuum. It is stored in the dark. [Evans and Aylesworth *Ind Eng Chem (Anal ed)* 18 24 1926.]

Impurities resulting from storage can be removed by passage through chromatographic grade alumina. Furfural can be separated from impurities other than carbonyl compounds by the bisulfite addition compound. The aldehyde is steam volatile.

It has been purified by distn (using a Claisen head) under reduced pressure. This is essential as is the use of an oil bath with temperatures of no more than 130° are highly recommended. When furfural is distd at atm press (in a stream of N₂), or under reduced pressure with a free flame (caution because the aldehyde is flammable) an almost colourless oil is obtained. After a few days and sometimes a few hours the oil gradually darkens and finally becomes black. This change is accelerated by light but occurs more slowly when kept in a brown bottle. However, when the aldehyde is distd under vacuum and the bath temperature kept below 130° during the distn, the oil develops only a slight colour when exposed to direct sunlight during several days. The distn of very impure material should NOT be attempted at atm pressure otherwise the product darkens rapidly. After one distn under vacuum a distn at atmospheric pressure can be carried out without too much decomposition and darkening. The liquid **irritates mucous membranes**. Store in dark containers under N₂. [Adams and Voorhees *Org Synth Coll Vol I* 280 1941.]

Furfuryl alcohol (2-furylmethanol) [98-00-0] M 98.1, b 68-69°/20mm, 170.0°/750mm, d 1.132, n 1.4873, n³⁰ 1.4801, pK²⁵ 2.61. Distd under reduced pressure to remove tarry material, shaken with aqueous NaHCO₃, dried with Na₂SO₄ and fractionally distd under reduced pressure from Na₂CO₃. Further dried by shaking with Linde 5A molecular sieves.

Furfurylamine (2-aminomethylfuran) [617-89-0] M 97.1, b 142.5-143°/735mm, d 1.059, n 1.489, pK³⁰ 8.89. Distd under nitrogen from KOH through a column packed with glass helices.

Furil [492-94-4] M 190.2, m 165-166°. Crystd from MeOH or *benzene (charcoal).

Furoin [552-86-3] M 192.2, m 135-136°. Crystd from MeOH (charcoal).

[Previous Page](#)

Galactaric Acid (mucic acid) [526-99-6] M 210.1, m 212-213^o(dec) pK₁²⁵ 3.09 (3.29), pK₂²⁵ 3.63 (4.41). Dissolved in the minimum volume of dil aq NaOH, and pptd by adding dil HCl. The temperature should be kept below 25^o.

D-Galactonic acid [576-36-3] M 196.2, m 148^o, pK_{Est} ~3.5. Crystd from EtOH. Cyclises to *D-galactonono-1,4-lactone*, m 134-136^o, [α]₅₄₆³⁰ mutarotates in 1h to -92^o (c 5, H₂O).

D(-)-Galactono-1,4-lactone [2782-07-2] M 178.1, m 134-137^o, [α]_D²⁰ -78^o (in H₂O). Crystd from EtOH.

D(+)-Galactosamine hydrochloride [1772-03-8] M 215.6, m 181-185^o, [α]_D²⁵ +96.4^o (after 24h, c 3.2 in H₂O), pK_{Est} ~7.7 (free base). Dissolved in a small volume of H₂O. Then added three volumes of EtOH, followed by acetone until faintly turbid and stood overnight in a refrigerator. [Roseman and Ludoweig *J Am Chem Soc* 76 301 1954.]

α-D-Galactose [59-23-4] M 180.2, m 167-168^o, [α]_D²⁰ +80.4^o (after 24h, c 4 in H₂O), pK₂²⁵ 12.48. Crystd twice from aqueous 80% EtOH at -10^o, then dried under vacuum over P₂O₅.

Gallic acid (H₂O) (3,4,5-trihydroxybenzoic acid) [5995-86-8 (H₂O), 149-91-7 (anhydr)] M 188.1, m 253^o(dec), pK₁²⁵ 4.27, pK₂²⁵ 8.68. Crystd from water.

Galvinoxyl [2,6-di-*tert*-butyl-α-(3,5-di-*tert*-butyl-4-oxo-2,5-cyclohexadiene-1-ylidene)-*p*-tolylxy] [2370-18-5] M 421.65, m 153.2-153.6^o, 158-159^o. A stable free radical scavenger of short-lived free radicals with odd electrons on C or O. Best prepared freshly by oxidation of 3,3',5,5'-tetra-*tert*-butyl-4,4'-dihydroxydiphenyl-methane [m 154^o, 157.1-157.6^o; obtained by gently heating for 10-15min 2,6-di-*tert*-butylphenol, formaldehyde and NaOH in EtOH and recryst from EtOH (20g/100mL) as colorless plates, Karasch and Joshi *J Org Chem* 22 1435 1957; Bartlett et al. *J Am Chem Soc* 82 1756 1960 and 84 2596 1962.] Oxidation is carried out under N₂ with PbO₂ in Et₂O or isooctane [Galvin A. Coppinger *J Am Chem Soc* 79 501 1957; Bartlett et al. above] or with alkaline potassium ferricyanide [Karasch and Joshi, above], whereby Galvinoxyl separates as deep blue crystals, and is recrystd twice under N₂ from *C₆H₆ soln by suction evaporation at 30^o. The VIS spectrum has λ_{max} 407nm (ε 30,000), 431nm (ε 154,000) and weak absorption at 772.5nm, and IR: ν 1577 and 2967cm⁻¹, and is estimated by iodometric titration. It is sensitive to O₂ in presence of OH⁻ ions and to traces of strong acid in hydroxylic or hydrocarbon solvents. At 62.5^o in a 0.62mM soln in *C₆H₆ the radical decays with a first order *k* = 4 x 10⁻⁸ sec⁻¹ (half life 1.7 x 10¹⁷ sec, ~200 days) as observed by the change in OD at 550nm [see also Green and Adam *J Org Chem* 28 3550 1963].

Genistein (4',5,7-trihydroxyisoflavone) [446-72-0] M 270.2, m 297-298^o, [α]_D²⁰ -28^o (c 0.6, 20mM NaOH), (phenolic pKs 8-10). Crystd from 60% aqueous EtOH or water.

Genistin (genistein-7-D-glucoside) [529-59-9] M 432.4, m 256^o. Crystd from 80% EtOH/water.

α-Gentiobiose (amygdalose, 6-O-α-D-glucopyranosyl-D-glucopyranose) [5995-99-5 (*bi-pyranose*)] M 342.3, m 86^o, [α]₅₄₆²⁰ +11^o (after 24h, c 4, H₂O). Crystd from MeOH (retains solvent of crystn).

β-Gentiobiose (see above) [5996-00-9 (*bi-pyranose*); 554-91-6 (*one open ring*)] M 342.3, m 190-195^o, [α]₅₄₆²⁰ +8^o (after 6h, c 3, H₂O). Crystd from MeOH or EtOH.

Geraniol [106-24-1] M 154.3, b 230^o, d 0.879, n 1.4766. Purified by ascending chromatography or by thin layer chromatography on plates of kieselguhr G with acetone/water/liquid paraffin (130:70:1) as solvent system. Hexane/ethyl acetate (1:4) is also suitable. Also purified by GLC on a silicone-treated column of Carbowax 20M (10%) on Chromosorb W (60-80 mesh). [Porter *Pure Appl Chem* 20 499 1969.] Stored in full, tightly sealed containers in the cool, protected from light.

Gibberillic acid (GA₃) [77-06-5] M 346.4, m 233-235^o(dec), [α]₅₄₆²⁰ +92^o (c 1, MeOH), pK 4.0. Crystd from ethyl acetate.

Girard Reagent T (2-hydrazino-*N,N,N*-trimethyl-2-oxo-ethaniminium chloride) [123-46-6] **M 167.6, m 192°**. Should be crystd from absolute EtOH (slight decomposition) when it has a slight odour, and stored in tightly stoppered containers because it is hygroscopic.

Glucamine [488-43-7] **M 181.2, m 127°, $[\alpha]_D^{20} -8^\circ$ (c 10, H₂O), $pK_{Est} \sim 9.0$** . Crystd from MeOH.

D-Gluconamide [3118-85-2] **M 197.2, m 144°, $[\alpha]_D^{23} +31^\circ$ (c 2, H₂O)**. Crystd from EtOH.

D-Glucono- δ -lactone [90-80-2] **M 178.1, m 152-153°, $[\alpha]_{546}^{20} +76^\circ$ (c 4, H₂O)**. Crystd from ethylene glycol monomethyl ether and dried for 1h at 110°.

D-Glucosamine [3416-24-8] **M 179.2, m 110°(dec), $[\alpha]_D^{20} +28^\circ \rightarrow +48^\circ$ (c 5, H₂O), $pK^{24} 7.71$** . Crystd from MeOH. *N*-Acetyl deriv, **m 205°** from MeOH/Et₂O has $[\alpha]_D^{20} +64^\circ \rightarrow +41^\circ$ (c 5, H₂O).

D-Glucosamine hydrochloride [66-84-2] **M 215.6, m >300°, $[\alpha]_D^{25} +71.8^\circ$ (after 20h, c 4, H₂O)**. Crystd from 3M HCl, water, and finally water/EtOH/acetone as for galactosamine hydrochloride.

α -D-Glucose [492-62-6] **M 180.2, m 146°, $[\alpha]_D^{20} +52.5^\circ$ (after 24h, c 4, H₂O), $pK^{25} 12.46$** . Recrystd slowly from aqueous 80% EtOH, then vacuum dried over P₂O₅. Alternatively, crystd from water at 55°, then dried for 6h in a vacuum oven between 60-70° at 2mm.

β -D-Glucose [50-99-7] **M 180.2, m 148-150°**. Crystd from hot glacial acetic acid.

α -D-Glucose pentaacetate [604-68-2] **M 390.4, m 110-111°, 112°, $[\alpha]_{546}^{20} +119^\circ$ (c 5, CHCl₃)**. Crystd from MeOH or EtOH.

β -D-Glucose pentaacetate [604-69-3] **M 390.4, m 131-132°, $[\alpha]_{546}^{20} +5^\circ$ (c 5, CHCl₃)**. Crystd from MeOH or EtOH.

D-Glucose phenylhydrazone [534-97-4] **M 358.4, m 208°**. Crystd from aqueous EtOH.

D-Glucuronic acid [6556-12-3] **M 194.1, m 165°, $[\alpha]_D^{20} +36^\circ$ (c 3, H₂O), $pK_2^{20} 3.18$** . Crystd from EtOH or ethyl acetate.

D-Glucuronolactone [32449-92-6] **M 176.1, m 175-177°, $[\alpha]_{546}^{20} +22^\circ$ (after 24h, c 10, H₂O)**. Crystd from water.

L-Glutamic acid [56-86-0] **M 147.1, m 224-225°(dec), $[\alpha]_D^{25} +31.4^\circ$ (c 5, 5M HCl), pK_1^{20} , $pK_2^{20} 2.06, 4.35$, $pK_3^{20} 9.85$** . Crystd from H₂O acidified to pH 3.2 by adding 4 volumes of EtOH, and dried at 110°. Likely impurities are aspartic acid and cysteine.

L-Glutamic acid- γ -benzyl ester [1676-73-9] **M 237.3, m 179-181°, $[\alpha]_{589}^{20}$, 19.3° (c 1, HOAc), $pK_1^{25} 2.17$, $pK_2^{25} 9.00$** . Recrystd from H₂O and stored at 0°. [Estrin *Biochem Prep* 13 25 1971.]

L-Glutamine [56-85-9] **M 146.2, m 184-185°, $[\alpha]_D^{25} +31.8^\circ$ (M HCl), $pK_1^{25} 2.17$, $pK_2^{25} 9.13$** . Likely impurities are glutamic acid, ammonium pyroglutamate, tyrosine, asparagine, isoglutamine, arginine. Crystd from water.

Glutaraldehyde [111-30-8] **M 100.1, b 71°/10mm, as 50% aq soln**. Likely impurities are oxidation products - acids, semialdehydes and polymers. It can be purified by repeated washing with activated charcoal (Norit) followed by vacuum filtration, using 15-20g charcoal/100mL of glutaraldehyde soln.

Vacuum distn at 60-65°/15mm, discarding the first 5-10%, was followed by dilution with an equal volume of freshly distilled water at 70-75°, using magnetic stirring under nitrogen. The soln is stored at low temp (3-4°),

in a tightly stoppered container, and protected from light. Standardised by titration with hydroxylamine. [Anderson *J Histochem Cytochem* **15** 652 1967.]

Glutaric acid [110-94-1] **M 132.1, m 97.5-98°, pK₁²⁵ 4.35, pK₂²⁵ 5.40.** Crystd from *benzene, CHCl₃, distilled water or *benzene containing 10% (w/w) of diethyl ether. Dried under vacuum.

dl-Glyceraldehyde [56-82-6] **M 90.1, m 145°.** Crystd from EtOH/diethyl ether.

Glycerol [56-81-5] **M 92.1, m 18.2°, b 182°/20mm, 290°/760mm, d 1.261, n²⁵ 1.47352, pK 14.4.** Glycerol was dissolved in an equal volume of *n*-butanol (or *n*-propanol, amyl alcohol or liquid ammonia) in a water-tight container, cooled and seeded while slowly revolving in an ice-water slurry. The crystals were collected by centrifugation, then washed with cold acetone or isopropyl ether. [Hass and Patterson *Ind Eng Chem (Anal Ed)* **33** 615 1941.] Coloured impurities can be removed from substantially dry glycerol by extraction with 2,2,4-trimethylpentane. Alternatively, glycerol can be decolorised and dried by treatment with activated charcoal and alumina, followed by filtering. Glycerol can be distd at 15mm in a stream of dry nitrogen, and stored in a desiccator over P₂O₅. Crude glycerol can be purified by digestion with conc H₂SO₄ and saponification with a lime paste, then re-acidified with H₂SO₄, filtered, treated with an anion exchange resin and fractionally distd under vacuum.

Glycidol (oxirane-2-methanol) [*RS*-(±)- 556-52-5; *R*-(+)- 57044-25-4; *S*-(-)- 60456-23-7] **M 74.1, b 61-62°/15mm, d 1.117, n 1.433 (±), b 49-50°/7mm, 66-67°/19mm, [α]_D²⁰ -15° (neat) (*S*-isomer, § also available on polymer support), b 56-56.5°/11mm, d 1.117, n 1.429, [α]_D²⁰ +15° (neat).** Purified by fractional distn. The 4-nitrobenzoates have **m 56° (±); m 60-62°, [α]_D²⁰ -37.9° (c 3.38 CHCl₃) for *R*-(-)-isomer [106268-95-5]; m 60-62°, [α]_D²⁰ +38° (c 1 CHCl₃) for *S*-(+)-isomer m 60-62°, [α]_D²⁰ -38° (c 1 CHCl₃) [115459-65-9], and are recrystd from Et₂O or Et₂O/pet ether (b 40-60°) [*S*-isomer: Burgos et al. *J Org Chem* **52** 4973 1987; Sowden and Fischer *J Am Chem Soc* **64** 1291 1942.]**

Glycinamide hydrochloride [1668-10-6] **M 110.5, m 186-189° (207-208°), pK₁²⁵ -6.10, pK₂²⁵ -1.78, pK₃²⁵ 7.95.** Crystd from EtOH.

Glycine see aminoacetic acid.

Glycine ethyl ester hydrochloride [623-33-6] **M 136.9, m 145-146°, pK²⁵ 7.69.** Crystd from absolute EtOH.

Glycine hydrochloride [6000-43-7] **M 111.5, m 176-178°.** Crystd from absolute EtOH.

Glycine methyl ester hydrochloride [5680-79-5] **M 125.6, m 174°(dec), pK²⁵ 7.66.** Crystd from MeOH.

Glycine *p*-nitrophenyl ester hydrobromide. [7413-60-7] **M 277.1, m 214° (dec).** Recryst from MeOH by adding diethyl ether. [Alners et al. *Biochem Preps* **13** 22 1971].

Glycocholic acid (*N*-cholyglycine) [475-31-0] **M 465.6, m 154-155°, 165-168°, [α]₅₄₆²⁰ +37° (c 1, EtOH), pK 4.4.** Crystd from hot water as sesquihydrate. Dried at 100°.

Glycolic (α-hydroxyacetic) acid [79-14-1] **M 76.1, m 81°, pK²⁵ 3.62.** Crystd from diethyl ether.

***N*-Glycylanilide** [555-48-6] **M 150.2, m 62°, pK_{Est}~8.0.** Crystd from water, sol in Et₂O..

Glycylglycine [556-50-3] **M 132.1, m 260-262°(dec), pK²⁰ 8.40, pK³⁰ 8.04.** Crystd from aqueous 50% EtOH or water at 50-60° by addition of EtOH. Dried at 110°.

Glycylglycine hydrochloride [13059-60-4] M 168.6, m 215-220°, 235-236°, 260-262°, pK_1^{25} 3.12, pK_2^{25} 8.17. Crystd from 95% EtOH.

Glycyl-L-proline [704-15-4] M 172.2, m 185°, pK_1^{25} 2.81, pK_2^{25} 8.65. Crystd from water at 50-60° by addition of EtOH.

dl-Glycylserine [687-38-7] M 162.2, m 207°(dec), pK_1^{25} 2.92, pK_2^{25} 8.10. Crystd from H₂O (charcoal) by addition of EtOH.

Glycyrrhizic acid ammonium salt (3H₂O) [53956-04-0] M 823.0, m 210°(dec), $[\alpha]_{546}^{20}$ +60° (c 1, 50% aq EtOH), pK_{Est} ~4.0. Crystd from glacial acetic acid, then dissolved in ethanolic ammonia and evaporated.

Glyoxal bis(2-hydroxyanil) [1149-16-2] M 240.3, m 210-213°, ϵ_{294nm} 9880. Crystd from MeOH or EtOH.

Glyoxylic acid [298-12-4] M 74.0, m 98°(anhydr), 50-52°(monohydrate), pK^{25} 2.98. Crystd from water as the monohydrate.

Gramine (3-dimethylaminoethylindole) [87-52-5] M 174.3, m 134°, pK^{25} 16.00 (NH acidic). Crystd from diethyl ether, ethanol or acetone.

Griseofulvin [126-07-8] M 352.8, m 220°, $[\alpha]_D^{22}$ +365° (c 1, acetone). Crystd from *benzene.

Guaiacic acid [4,4'-(2,3-dimethyl-1-butene-(1,4-diyl)-bis-(2-methoxyphenol)] [500-40-3] M 328.4, m 99-100.5°, pK_{Est} ~10.0. Crystd from EtOH.

Guaiacol (2-methoxyphenol) [90-05-1] M 124.1, m 32°, b 106°/24mm, 205°/746mm, pK^{25} 9.90. Crystd from *benzene/pet ether or distd.

Guaiacol carbonate [553-17-3] M 274.3, m 88.1°. Crystd from EtOH.

Guanidine [113-00-8] M 59.1, m ~50°, pK^{25} 13.6. Crystd from water/EtOH under nitrogen. Very deliquescent and absorbs CO₂ from the air readily.

Guanidine carbonate [593-85-1] M 180.2, m 197°. Crystd from MeOH.

Guanidine hydrochloride [50-01-1] M 95.5, m 181-183°. Crystd from hot methanol by chilling to about -10°, with vigorous stirring. The fine crystals were filtered through fritted glass, washed with cold (-10°) methanol, dried at 50° under vacuum for 5h. (The product is more pure than that obtained by crystn at room temperature from methanol by adding large amounts of diethyl ether.) [Kolthoff et al. *J Am Chem Soc* 79 5102 1957].

Guanosine (H₂O) [118-00-3] M 283.2, m 240-250°(dec), $[\alpha]_{546}^{20}$ -86° (c 1, 0.1M NaOH), pK_1^{25} 1.9, pK_2^{25} 9.24, pK_3^{25} 12.33. Crystd from water. Dried at 110°.

Guanylic acid (guanosine-5'-monophosphoric acid) [85-32-5] M 363.2, m 208°(dec), pK_2^{25} 2.4, pK_3^{25} 6.66 (6.1), pK_4^{25} 9.4. Crystd from water. Dried at 110°.

Harmine [442-51-3] M 212.3, m 261°(dec), pK^{20} 7.61. Crystd from MeOH.

Harmine hydrochloride (hydrate) [343-27-1] M 248.7, m 280°(dec). Crystd from water.

Hecogenine acetate [1915-35-5] M 472.7, m 265-268°, $[\alpha]_D^{23}$ -4.5° (c 1, CHCl₃). Crystd from MeOH.

Heptadecanoic acid (margaric) [506-12-7] M 270.5, m 60-61°, b 227°/100mm, pK_{Est} ~4.9. Crystd from MeOH or pet ether.

1-Heptadecanol [1454-85-9] M 256.5, m 54°. Crystd from acetone.

Heptafluoro-2-iodopropane [677-69-0] M 295.9, b 41°. Purified by gas chromatography on a triacetin (glyceryl triacetate) column, followed by bulb-to-bulb distn at low temperature. Stored over Cu powder to stabilise it.

***n*-Heptaldehyde** [111-71-7] M 114.2, b 40.5°/12mm, 152.8°/760mm, d 0.819, n²⁵ 1.4130. Dried with CaSO₄ or Na₂SO₄ and fractionally distd under reduced pressure. More extensive purification by pptn as the bisulfite compound (formed by adding the aldehyde to saturated aqueous NaHSO₃) which was filtered off and recrystd from hot H₂O. The crystals, after being filtered and washed well with H₂O, were hydrolysed by adding 700mL of aqueous Na₂CO₃ (12.5% w/w of anhydrous Na₂CO₃) per 100g of aldehyde. The aldehyde was then steam distd, separated, dried with CuSO₄ and distd under reduced pressure in a slow stream of nitrogen. [McNesby and Davis *J Am Chem Soc* 76 2148 1954].

***n*-Heptaldoxime** [629-31-2] M 129.2, m 53-55°. Crystd from 60% aqueous EtOH.

***n*-Heptane** [142-18-5] M 100.2, b 98.4°, d 0.684, n 1.38765, n²⁵ 1.38512. Passage through a silica gel column greatly reduces the ultraviolet absorption of *n*-heptane. (The silica gel is previously heated to 350° before use.) For more extensive purification, heptane is shaken with successive small portions of conc H₂SO₄ until the lower (acid) layer remains colourless. The heptane is then washed successively with water, aq 10% Na₂CO₃, water (twice), and dried with CaSO₄, MgSO₄ or CaCl₂. It is distd from sodium. *n*-Heptane can be distd azeotropically with methanol, then the methanol can be washed out with water and, after drying, the heptane is redistd. Other purification procedures include passage through activated basic alumina, drying with CaH₂, storage with sodium, and stirring with 0.5N KMnO₄ in 6N H₂SO₄ for 12h after treatment with conc H₂SO₄. Carbonyl-containing impurities have been removed by percolation through a column of impregnated Celite made by dissolving 0.5g of 2,4-dinitrophenylhydrazine in 6mL of 85% H₃PO₄ by grinding together, then adding 4mL of distilled water and 10g Celite. [Schwartz and Parks *Anal Chem* 33 1396 1961].

Hept-1-ene [592-76-7] M 98.2, b 93°/771mm, d 0.698, n 1.400. Distd from sodium, then carefully fractionally distd using an 18-in gauze-packed column. Can be purified by azeotropic distn with EtOH. Contained the 2- and 3-isomers as impurities. These can be removed by gas chromatography using a Carbowax column at 70°.

***n*-Heptyl alcohol** [111-70-6] M 116.2, b 175.6°, d 0.825, n 1.425. Shaken with successive lots of alkaline KMnO₄ until the colour persisted for 15min, then dried with K₂CO₃ or CaO, and fractionally distd.

***n*-Heptylamine** [111-68-2] M 115.2, b 155°, d 0.775, n 1.434, pK²⁵ 10.66. Dried in contact with KOH pellets for 24h, then decanted and fractionally distd.

***n*-Heptyl bromide** [629-04-9] M 179.1, b 70.6°/19mm, 180°/760mm, d 1.140, n 1.45. Shaken with conc H₂SO₄, washed with water, dried with K₂CO₃, and fractionally distd.

Heptyl-β-D-glucopyranoside [78617-12-6] M 278.4, m 74-77°, 76-77°, $[\alpha]_D^{20}$ -34.2° (c 5, H₂O). Purified by several recrystns from M₂CO which is a better solvent than EtOAc. The *acetate* has m 66-68.5°, $[\alpha]_D^{20}$ -20.5° (c 4, CHCl₃) [Pigman and Richtmyer *J Am Chem Soc* 64 369 1942].

Heptyl-β-D-1-thioglucopyranoside [85618-20-8] M 294.4, m 98-99°. The *tetra-acetyl derivative* is purified by silica gel column chromatography and eluted with a *C₆H₆-Me₂CO (gradient up to 5% of Me₂CO) and recrystd from *n*-hexane as colourless needles m 72-74° (Erbing and Lindberg *Acta Chem Scand*

B30 611 1976 gave **m 69-70°**). Hydrolysis using an equivalent of base in methanol gave the desired glucoside. This is a non-ionic detergent for reconstituting membrane proteins and has a critical micelle concentration of 30 mM. [Shimamoto et al. *J Biochem (Tokyo)* **97** 1807 1985; Saito and Tsuchiya *Chem Pharm Bull Jpn* **33** 503 1985].

Hesperetin (3',5,7-trihydroxy-4'-methoxyflavanone) [520-33-2] **M 302.3, m 227-228°, pK_{Est} ~8.5-10.5 (phenolic)**. Crystd from EtOAc or ethanol. The natural (-) form has $[\alpha]_D^{20} -38^\circ$ (c 2, EtOH). Note that C2 is chiral.

Hesperidin (hesperetin 7-rhamnoside) [520-26-3] **M 610.6, m 258-262°, $[\alpha]_{546}^{20} -82^\circ$ (c 2, pyridine)**. Dissolved in dilute aqueous alkali and ppted by adjusting the pH to 6-7.

Hexachlorobenzene [118-74-1] **M 284.8, m 230.2-231.0°**. Crystd repeatedly from *benzene. Dried under vacuum over P₂O₅.

Hexachloro-1,3-butadiene [87-68-3] **M 260.8**. See perchlorobutadiene on p. 323.

1,2,3,4,5,6-Hexachlorocyclohexane [α -319-84-6; γ -58-89-9] **M 290.8, m 158° (α -), 312° (β -), 112.5° (γ -isomer)**. Crystd from EtOH. Purified by zone melting. **Possible CANCER AGENT, TOXIC**.

Hexachlorocyclopentadiene [77-47-4] **M 272.8, b 80°/1mm, d 1.702, n²⁵ 1.5628**. Dried with MgSO₄. Distd under vacuum in nitrogen.

Hexachloroethane [67-72-1] **M 236.7, m 187°**. Steam distd, then crystd from 95% EtOH. Dried in the dark under vacuum.

Hexacosane [630-01-3] **M 366.7, m 56.4°, b 169°/0.05mm, 205°/1mm, 262°/15mm**. Distd under vacuum and crystd from diethyl ether.

Hexacosanoic acid (cerotinic acid) [506-46-7] **M 396.7, m 86-87°, 88-89°, pK_{Est} ~4.9**. Crystd from EtOH, aq EtOH and pet ether+Me₂CO.

1,14-Hexadecanedioic acid (thaspic acid). [505-54-4] **M 286.4, m 126°, pK_{Est(1)}~4.5, pK_{Est(2)}~5.5**. Crystd from EtOH, ethyl acetate or *C₆H₆.

n-Hexadecane (Cetane) [544-76-3] **M 226.5, m 18.2°, b 105°/0.1mm, d 0.773, n 1.4345, n²⁵ 1.4325**. Passed through a column of silica gel and distd under vacuum in a column packed with Pyrex helices. Stored over silica gel. Crystd from acetone, or fractionally crystd by partial freezing.

Hexadecanoic acid (palmitic acid) [57-10-3] **M 256.4, m 62-63°, b 215°/15mm, pK²⁵ 6.46 (50% aq EtOH), 5.0 (H₂O)**. Purified by slow (overnight) recrystn from hexane. Some samples were also crystd from acetone, EtOH or EtOAc. Crystals were stood in air to lose solvent, or were pumped dry of solvent on a vacuum line. [Iwahashi et al. *J Chem Soc, Faraday Trans 1* **81** 973 1985; pK: White *J Am Chem Soc* **72** 1858 1950].

Hexadecyl 3-hydroxynaphthalene-2-carboxylate [531-84-0] **M 412.6, m 73-74°**. Recrystd from hot EtOH and sublimed in a vacuum. [Oshima and Hayashi *J Soc Chem Ind Jpn* **44** 821 1941.]

1,5-Hexadiene [592-42-7] **M 82.2, b 59.6°, d 0.694, n 1.4039**. Distd from NaBH₄.

Hexaethylbenzene [604-88-6] **M 246.3, m 128.7-129.5°**. Crystd from *benzene or *benzene/EtOH.

Hexafluoroacetone [684-16-2, 34202-69-2 (3H₂O)] **M 166.1, m -129°, (trihydrate m 18-21°), b -28°**. Dehydrated by passage of the vapour over P₂O₅. Ethylene was removed by passing the dried vapour

through a tube containing Pyrex glass wool moistened with conc H_2SO_4 . Further purification was by low temperature distn using Warde-Le Roy stills. Stored in the dark at -78° . [Holmes and Kutschke *Trans Faraday Soc* **58** 333 1962].

Hexafluoroacetylacetone (1,1,1,5,5,5-hexafluoro-2,4-pentanedione) [1522-22-1] M 208.1, b $68^\circ/736\text{mm}$, 70-70.2 $^\circ/760\text{mm}$, 68-71 $^\circ/\text{atm}$, d_4^{20} 1.490, n_D^{20} 1.333. It forms a dihydrate which has no UV spectrum compared with λ_{max} (CHCl_3) 273nm (ϵ 7,800) for the anhydrous ketone. The dihydrate dec at $\sim 90^\circ$. The hydrate (10g) plus anhyd CaSO_4 (Drierite, 30g) are heated and distd; the distillate is treated with more CaSO_4 and redist. When the distillate is treated with aqueous NaOH and heated, the dihydrate crystallises on cooling. The Cu complex has m 135° (after sublimation). [Gilman et al. *J Am Chem Soc* **78** 2790 1956; Belford et al. *J Inorg Nucl Chem* **2** 11 1956].

Hexafluorobenzene [392-56-3] M 186.1, m 5.1° , b $79-80^\circ$, d 1.61, n 1.378. Main impurities are incompletely fluorinated benzenes. Purified by standing in contact with oleum for 4h at room temperature, repeating until the oleum does not become coloured. Washed several times with water, then dried with P_2O_5 . Final purification was by repeated fractional crystn.

Hexafluoroethane [76-16-4] M 138.0, b -79° . Purified for pyrolysis studies by passage through a copper vessel containing CoF_3 at ca 270° , and held for 3h in a bottle with a heated (1300°) platinum wire. It was then fractionally distd. [Steunenberg and Cady *J Am Chem Soc* **74** 4165 1962.]

1,1,1,3,3,3-Hexafluoropropan-2-ol [920-66-1] M 168.1, b $57-58^\circ/760\text{mm}$, d 1.4563, n_D^{22} 1.2750. Distd from 3A molecular sieves, retaining the middle fraction.

Hexahydro-1H-azepine (hexamethyleneimine, Azepane) [111-49-9] M 99.2, b $70-72^\circ/30\text{mm}$, $135-138^\circ/\text{atm}$, d 0.879, n 1.466, pK^{25} 11.10 (pK° 9.71, pK^{75} 9.71). Purified by dissolving in Et_2O and adding ethanolic HCl until all the base separates as the white *hydrochloride*, filter, wash with Et_2O and dry (m 236°). The salt is dissolved in the minimum vol of H_2O and basified to pH ~ 14 with 10N KOH . The soln is extracted with Et_2O , the extract is dried over KOH , evapd and distd. The base is a **FLAMMABLE** and **TOXIC** liquid, and best kept as the salt. The *nitrate* has m $120-123^\circ$, *Picrate* has m $145-147^\circ$, and the *Tosylate* has m 76.5° (ligroin). [Müller and Sauerwald *Monatsh Chem* **48** 727 1027; Hjelt and Agback *Acta Chem Scand* **18** 194 1964].

Hexahydromandelic acid [*R*-(-)-53585-93-6; *S*-(+)-61475-31-8] M 158.2, m $127-129^\circ$, $128-129^\circ$, 129.7° , $[\alpha]_D^{20}$ (-) and (+) 25.5° (c 1, AcOH) and $[\alpha]_D^{20}$ (-) and (+) 13.6° (c 7.6, EtOH). For hexagonal clusters by recrystallisation from CCl_4 or Et_2O . [Wood and ComLey *J Chem Soc* 2638 1924; Lettré et al. *Chem Ber* **69** 1594 1936]. The *racemate* has m $137.2-137.6^\circ$ ($134-135^\circ$) [Smith et al. *J Am Chem Soc* **71** 3772 1949].

Hexamethylbenzene [87-85-4] M 162.3, m $165-165.5^\circ$. Sublimed, then crystd from abs EtOH , *benzene, $\text{EtOH}/\text{*benzene}$ or $\text{EtOH}/\text{cyclohexane}$. Also purified by zone melting. Dried under vac over P_2O_5 .

Hexamethyl(Dewar)benzene [7641-77-2] M 162.3, m 7° , b $60^\circ/20\text{mm}$, d 0.803, n 1.4480. Purified by passage through alumina [Traylor and Miksztal *J Am Chem Soc* **109** 2770 1987].

Hexamethylenediamine [124-09-4] M 116.2, m 42° , b $46-47^\circ/1\text{mm}$, $84.9^\circ/9\text{mm}$, $100^\circ/20\text{mm}$, $204-205^\circ/760\text{mm}$, pK_1^{25} 10.24, pK_2^{25} 11.02. Crystd in a stream of nitrogen. Sublimed in a vacuum.

Hexamethylenediamine dihydrochloride [6055-52-3] M 189.2, m 248° . Crystd from water or EtOH .

Hexamethylene glycol (1,6-hexanediol) [629-11-8] M 118.2, m 41.6° , $43-45^\circ$, b $134^\circ/10\text{mm}$, 250° , n 1.458. Fractionally crystd from its melt or from water. Distils *in vacuo*.

Hexamethylenetetramine (Urotropine, hexamine, HMTA) [100-97-0] M 140.1, m 280° (subln), 290-292° (sealed tube, CARE), d 1.331, pK²⁵ 4.85 (6.30). It is soluble in H₂O (67%), CHCl₃ (10%), EtOH (8%) and Et₂O (0.3%), and a 0.2M soln has a pH of 8.4. Dissolve in hot abs EtOH (reflux, Norite), filter using a heated funnel, cool at room temp first then in ice. Wash crystals with cold Et₂O, dry in air or under a vacuum. A further crop can be obtained by adding Et₂O to the filtrate. It sublimes above 260° without melting. The *picrate* has m 179° (dec). [pK 4.85: Reilley and Schmid *Anal Chem* 30 947 1958; pK 6.30: Pummerer and Hofmann *Chem Ber* 56 1255 1923.]

n-Hexane [110-54-3] M 86.2, b 68.7°, d 0.660, n 1.37486, n²⁵ 1.37226. Purification as for *n*-heptane. Modifications include the use of chlorosulfonic acid or 35% fuming H₂SO₄ instead of conc H₂SO₄ in washing the alkane, and final drying and distn from sodium hydride. Unsatd compounds can be removed by shaking the hexane with nitrating acid (58% H₂SO₄, 25% conc HNO₃, 17% water, or 50% HNO₃, 50% H₂SO₄), then washing the hydrocarbon layer with conc H₂SO₄, followed by H₂O, drying, and distg over sodium or *n*-butyl lithium. Also purified by distn under nitrogen from sodium benzophenone ketyl solubilised with tetraglyme. Also purified by passage through a silica gel column followed by distn [Kajii et al. *J Phys Chem* 91 2791 1987]. FLAMMABLE liquid and possible nerve toxin.

Rapid purification: Distil, discarding the first forerun and stored over 4A molecular sieves.

1,2-Hexanediol [6920-22-5] M 118.2, b 214-215, d 0.951, n 1.442. Fractionally distd.

1-Hexene [592-41-6] M 84.2, b 63°, d 0.674, n 1.388. Purified by stirring over Na/K alloy for at least 6h, then fractionally distd from sodium under nitrogen.

cis-2-Hexene [7688-21-3] M 84.2, b 68-70°, d 0.699, n 1.399. Purification as for 1-hexene above.

trans-2-Hexene [4050-45-7] M 84.2, b 65-67°, n 1.390. Purifn as for 1-hexene above.

trans-3-Hexene [13269-52-8] M 84.2, b 67-69°, d 0.678, n 1.393. Purifn as for 1-hexene above.

meso-Hexoestrol [84-16-2] M 270.4, m 185-188°. Crystd from *benzene or aqueous EtOH.

n-Hexyl alcohol (1-hexanol) [111-27-3] M 102.2, b 157.5°, d 0.818, n¹⁵ 1.4198, n²⁵ 1.4158. Commercial material usually contains other alcohols which are difficult to remove. A suitable method is to esterify with hydroxybenzoic acid, recrystallise the ester and saponify. [Olivier *Recl Trav Chim, Pays-Bas* 55 1027 1936.] Drying agents include K₂CO₃ and CaSO₄, followed by filtration and distn. (Some decomposition to the olefin occurred when Al amalgam was used as drying agent at room temperature, even though the amalgam was removed prior to distn.) If the alcohol is required anhydrous, the redistd material can be refluxed with the appropriate alkyl phthalate or succinate, as described under *Ethanol*.

n-Hexylamine [111-26-2] M 101.2, b 131°, d 0.765, n 1.419, pK²⁵ 10.64. Dried with, and fractionally distd from, KOH or CaH₂.

n-Hexyl bromide [111-25-1] M 165.1, b 87-88°/90mm, 155°/743mm, d 1.176, n 1.448. Shaken with H₂SO₄, washed with water, dried with K₂CO₃ and fractionally distd.

n-Hexyl methacrylate [142-09-6] M 154.2, b 65-66°/4mm, 88-88.5°/14mm, d 0.8849, n 1.4320. Purified as for *methyl methacrylate*. [IR: Hughes and Walton *J Am Chem Soc* 79 3985 1957.]

Hexyltrimethylammonium bromide [2650-53-5] M 224.3, m 186°. Recrystd from acetone. Extremely *hygroscopic* salt. [McDowell and Kraus *J Am Chem Soc* 73 2170 1951.]

1-Hexyne [693-02-7] M 82.2, b 12.5°/75mm, 71°/760mm, d 0.7156, n 1.3989. Distd from NaBH₄ to remove peroxides. Stood with sodium for 24h, then fractionally distd under reduced pressure. Also dried by repeated vac transfer into freshly activated 4A molecular sieves, followed by vacuum transfer into Na/K alloy and stirring for 1h before fractionally distilling.

2-Hexyne [764-35-2] M 82.2, b 83.8°/760mm, d 0.73146, n 1.41382. Purification as for 1-hexyne above.

3-Hexyne [928-49-4] M 82.2, b 81°/760mm, d 0.7231, n 1.4115. Purification as for 1-hexyne above.

Histamine [51-45-6] M 111.2, m 86° (sealed tube), b 167°/0.8mm, 209°/18mm, pK_1^{25} 6.02, pK_2^{25} 9.70. Crystd from *benzene or chloroform.

Histamine dihydrochloride [56-92-8] M 184.1, m 249-252° (244-245°). Crystd from aq EtOH.

S-Histidine [71-00-1] M 155.2, m 287°(dec), $[\alpha]_D^{25}$ -39.7° (c 1, H₂O), +13.0° (6M HCl), pK_1^{25} 1.96, pK_2^{25} 6.12, pK_3^{25} 9.17. Likely impurity is arginine. Adsorbed from aqueous soln on to Dowex 50-H⁺ ion-exchange resin, washed with 1.5M HCl (to remove other amino acids), then eluted with 4M HCl as the dihydrochloride. Histidine is also purified as the dihydrochloride which is finally dissolved in water, the pH adjusted to 7.0, and the free zwitterionic base crystallises out on addition of EtOH. Sol in H₂O is 4.2% at 25°.

S-Histidine dihydrochloride [1007-42-7] M 242.1, m 245°, $[\alpha]_D^{20}$ +47.5° (c 2, H₂O). Crystd from water or aqueous EtOH, and washed with acetone, then diethyl ether. Converted to the histidine di-(3,4-dichlorobenzenesulfonate) salt by dissolving 3,4-dichlorobenzenesulfonic acid (1.5g/10mL) in the aqueous histidine soln with warming, and then the soln is cooled in ice. The resulting crystals (m 280° dec) can be recrystd from 5% aqueous 3,4-dichlorobenzenesulfonic acid, then dried over CaCl₂ under vacuum, and washed with diethyl ether to remove excess reagent. The dihydrochloride can be regenerated by passing the soln through a Dowex-1 (Cl⁻ form) ion-exchange column. The solid is obtained by evapn of the soln on a steam bath or better in a vacuum. [Greenstein and Winitz, *The Amino Acids Vol 3* p. 1976 1961.]

S-Histidine monohydrochloride (H₂O) [5934-29-2 (H₂O); 7048-02-4] M 209.6, m 80° monohydrate, 254°(dec, anhyd), $[\alpha]_D^{25}$ +13.0° (6M HCl). Crystd from aqueous EtOH.

dl-Homocysteine [6027-13-0] M 135.2, pK_2^{25} 8.70, pK_3^{30} 10.46. Crystd from aqueous EtOH.

Homocystine [462-10-2] M 268.4, m 260-265°(dec), pK_1^{25} 1.59, pK_2^{25} 2.54, pK_3^{25} 8.52, pK_4^{25} 9.44. Crystd from water.

Homophthalic acid [89-51-0] M 180.2, m 182-183°, 189-190°, (depends on heating rate) $pK_{Est(1)}$ ~3.5, $pK_{Est(2)}$ ~4.3. Crystd from boiling water (25mL/g). Dried at 100°.

Homopiperazine (1,4-diazepane) [505-66-8] M 100.2, m 38-40°, 43°, b 60°/10mm, 92°/50mm, 169°/atm, pK_1^{20} 6.70, pK_2^{20} 10.41. Purified by fractionation through a column of 10 theoretical plates with a reflux ratio of 3:1. It boiled at 169° and the cool distillate crystallises in plates m 43°. [Poppelsdorf and Myerly *J Org Chem* 26 131 1961.] Its pKa values are 6.89 and 10.65 at 40°, and 6.28 and 9.86 at 40° [Pagano et al. *J Phys Chem* 65 1062 1961]. The 1,4-bis(4-bromobenzoyl) derivative has m 194-198° (from EtOH); the hydrochloride has m 270-290° (from EtOH) and the picrate has m 265° (dec) [Lloyd et al. *J Chem Soc (C)* 780 1966].

L-Homoserine (2-amino-4-hydroxybutyric acid) [672-15-1] M 119.1, m 203°, $[\alpha]_D^{26}$ +18.3° (in 2M HCl), $pK_{Est(1)}$ ~2.1, $pK_{Est(2)}$ ~9.3. Likely impurities are N-chloroacetyl-L-homoserine, N-chloroacetyl-D-homoserine, L-homoserine, homoserine lactone, homoserine anhydride (formed in strong solns of homoserine if slightly acidic). Cyclises to the lactone in strongly acidic soln. Crystd from water by adding 9 volumes of EtOH.

Homoveratronitrile (3,4-dimethoxybenzyl nitrile) [93-17-4] M 177.2, m 62-64°, 68°, b 184°/20mm, 195-196°/2mm, 208°/atm. Its solubility is 10% in MeOH. and has been recrystd from

EtOH or MeOH. Purified by distillation followed by recrystn. [Niederl and Ziering *J Am Chem Soc* **64** 885 1952; Julian and Sturgis *J Am Chem Soc* **57** 1126 1935.]

Homoveratrylamine (2-[3,4-dimethoxyphenyl]ethylamine) [120-20-7] M 181.2, b 99.3-101.3°/0.5mm, 168-170°/15mm, d_4^{20} 1.091, n_D^{20} 1.5460, pK_{Est} ~9.8. Purified by fractionation through an efficient column in an inert atmosphere as it is a relatively strong base. [Horner and Sturm *Justus Liebigs Ann Chem* **608** 12819 1957; Jung et al. *J Am Chem Soc* **75** 4664 1953.] The *hydrochloride* has m 152°, 154°, 156° (from EtOH, Me₂CO or EtOH/Et₂O) and the *picrate* has m 165-167° dec, and the 4-nitrobenzoyl derivative has m 147° [Buck *J Am Chem Soc* **55** 2593 1933].

Hordenine {4-[(2-dimethylamino)ethyl]phenol} [539-15-1] M 165.2, m 117-118°, pK^{25} 9.46 (OH). Crystd from EtOH or water.

Humulon [26472-41-3] M 362.5, m 65-66.5°, $[\alpha]_D^{26}$ -212° (95% EtOH). Crystd from Et₂O.

Hyamine 1622 [(diisobutylphenoxyethoxyethyl)dimethylbenzylammonium chloride, benzethonium chloride] [121-54-0] M 448.1, m 164-166° (sinters at 120°, monohydrate). Crystd from boiling acetone after filtering, or from CHCl₃-pet ether. The ppte was filtered off, washed with diethyl ether and dried for 24h in a vacuum desiccator.

Hydantoin (2,4-dihydroxyimidazole) [461-72-3] M 100.1, m 216°, 220°, pK^{25} 9.15. Crystd from MeOH. The *diacetate* has m 104-105°.

Hydrazine *N,N'*-dicarboxylic acid diamide [110-21-4] M 116.1, m 248°. Crystd from water and dried in vac over P₂O₅.

4-Hydrazinobenzoic acid [619-67-0] M 152.2, m 217° (dec), pK^{25} 4.13. Crystd from water.

1-Hydrazinophthalazine hydrochloride (hydralazine hydrochloride) [304-20-1] M 196.6, m 172-173°, pK^{20} 6.57. Crystd from MeOH.

2-Hydrazinopyridine [4930-98-7] M 109.1, m 41-44°, 46-47°, 49-50°, b 105°/0.5mm, 128-135°/13mm. Purified by distn and by recrystn from Et₂O-hexane. [Kauffmann et al. *Justus Liebigs Ann Chem* **656** 103 1962, Potts and Burton *J Org Chem* **31** 251 1966.] The *mono-hydrochloride* has m 183° (dec) from aq HCl and the *di-hydrochloride* has m 214-215°.

Hydrazobenzene [122-66-7] M 184.2. See 1,1-diphenylhydrazine on p. 225.

Hydrobenzamide [1-phenyl-*N,N'*-bis(phenylmethylene)-methanediamine] [92-29-5] M 298.4, m 101-102°, 107-108°. Crystd from absolute EtOH or cyclohexane/*benzene. Dried under vacuum over P₂O₅. [Pirrone *Gazz Chim Ital* **67** 534 1937.]

***dl*-Hydrobenzoin** [492-70-6] M 214.3, m 120°. Crystd from diethyl ether/pet ether.

***meso*-Hydrobenzoin** [579-43-1] M 214.3, m 139°. Crystd from EtOH or water.

Hydroquinone (1,4-dihydroxybenzene, quinol) [123-31-9] M 110.1, m 175.4, 176.6°, pK_1^{20} 9.91, pK_2^{20} 11.56. Crystd from acetone, *benzene, EtOH, EtOH/*benzene, water or acetonitrile (25g in 30mL), preferably under nitrogen. Dried under vacuum. [Wolfenden et al. *J Am Chem Soc* **109** 463 1987.]

4'-Hydroxyacetanilide [103-90-2] See 4-acetamidophenol on p. 83.

***p*-Hydroxyacetophenone** [99-93-4] M 136.2, m 109°, pK^{25} 8.01. Crystd from diethyl ether, aqueous EtOH or *benzene/pet ether.

- 4-Hydroxyacridine** [18123-20-1] M 195.2, m 116.5° pK₁¹⁵ 5.28, pK₂¹⁵ 9.75. Crystd from EtOH.
- 5-Hydroxyanthranilic acid** [548-93-6] M 153.1, m >240°(dec), λ_{max} 298nm, log ε 3000 (0.1M HCl), pK₁²⁰ 2.7, pK₂²⁰ 5.37, pK₃²⁰ 10.12. Crystd from water. Sublimes below its melting point in a vacuum.
- erythro-3-Hydroxy-RS-aspartic acid** [6532-76-9] M 149.1, pK₁²⁵ 1.91, pK₂²⁵ 3.51, pK₃²⁵ 9.11. Likely impurities are 3-chloromalic acid, ammonium chloride, *threo*-3-hydroxyaspartic acid. Crystd from water.
- m-Hydroxybenzaldehyde** [100-83-4] M 122.1, m 108° pK₁²⁵ 8.98, pK₂²⁵ 15.81. Crystd from water.
- p-Hydroxybenzaldehyde** [123-08-0] M 122.1, m 115-116°, pK₂²⁵ 7.61. Crystd from water (containing some H₂SO₄). Dried over P₂O₅ under vacuum.
- m-Hydroxybenzoic acid** [99-06-9] M 138.1, m 200.8°, pK₁²⁵ 4.08, pK₂²⁵ 9.98. Crystd from absolute EtOH.
- p-Hydroxybenzoic acid** [99-96-7] M 138.1, m 213-214°, pK₁²⁵ 4.50, pK₂²⁵ 9.11. Crystd from water.
- p-Hydroxybenzotrile** [767-00-0] M 119.1, m 113-114°. See *p*-cyanophenol on p. 176.
- 4-Hydroxybenzophenone** [1137-42-4] M 198.2, m 133.4-133.8°, pK₂²⁵ 7.95. See *p*-benzoylphenol on p. 126.
- 2-Hydroxybenzothiazole** [934-34-9] M 183.1, m 117-118° Crystd from aqueous EtOH or water. [Dryland and Sheppard *J Chem Soc Perkin Trans 1* 125 1986.]
- 1-Hydroxybenzotriazole hydrate (HOBt)** [2592-95-2] M 135.1, m 159-160°. Crystd from aqueous EtOH or water. [Dryland and Sheppard *J Chem Soc Perkin Trans 1* 125 1986.]
§ A polystyrene supported version is available.
- 2-Hydroxybenzyl alcohol** [90-01-7] M 124.1, m 87°, pK₂²⁵ 9.84. Crystd from water or *benzene.
- 3-Hydroxybenzyl alcohol** [620-24-6] M 124.1, m 71°, pK_{Est} ~9.8. Crystd from *benzene.
- 4-Hydroxybenzyl alcohol** [623-05-2] M 124.1, m 114-115°, pK₂²⁵ 9.73. Crystd from water.
- 2-Hydroxybiphenyl** [90-43-7] M 170.2, m 56°, b 145°/14mm, 275°/760mm, pK₂²⁰ 10.01. Crystd from pet ether.
- 4-Hydroxybiphenyl (4-phenylphenol)** [92-69-3] M 170.2, m 164-165°, b 305-308°/760mm, pK₂₃ 9.55. Crystd from aqueous EtOH, aq EtOH, *C₆H₆, and vac dried over CaCl₂ [Buchanan et al. *J Am Chem Soc* 108 7703 1986].
- 3-Hydroxy-2-butanone (acetoin)** [513-86-0] M 88.1, b 144-145°, [m 100-105° dimer]. Washed with EtOH until colourless, then with diethyl ether or acetone to remove biacetyl. Air dried by suction and further dried in a vacuum desiccator.
- (±)-α-Hydroxy-γ-butyrolactone** [19444-84-9] M 102.1, b 84°/0.2mm, 133°/10mm, d₄²⁰ 1.310, n_D²⁰ 1.4656. It has been purified by repeated fractionation, forms a colourless liquid. It has to be distd at high vacuum otherwise it will dehydrate. The *acetoxy* derivative has b 94°/0.2mm, [NMR: Daremon and Rambaud *Bull Soc Chim Fr* 294 1971; Schmitz et al. *Chem Ber* 108 1010 1975.]

4-Hydroxycinnamic acid (*p*-coumaric acid) [501-98-4] M 164.2, m 210-213°, 214-215°, 215° pK₁²⁵ 4.64, pK₂²⁵ 9.45. Crystd from H₂O (charcoal). Needles from conc aqueous solutions as the *anhydrous acid*, but from hot dilute solutions the *monohydrate acid* separates on slow cooling. The acid (33g) has been recrystd from 2.5L of H₂O (1.5g charcoal) yielding 28.4g of recrystd acid, m 207°. It is insol in *C₆H₆ or pet ether. The UV in 95% EtOH has λ_{max} 223 and 286nm (ε 14,450 and 19000 M⁻¹cm⁻¹). [UV Wheeler and Covarrubias *J Org Chem* 28 2015 1963; Corti *Helv Chim Acta* 32 681 1949.]

4-Hydroxycoumarin [1076-38-6] M 162.1, m 206°, pK_{Est} ~9.0. Crystd from water and dried in a vacuum desiccator over Sicapent.

3-(4-Hydroxy-3,5-dimethoxyphenyl)acrylic acid [530-59-6] M 234.1, m 204-205°(dec), pK_{Est(1)}~4.6, pK_{Est(2)}~9.3. Crystd from water.

4-Hydroxydiphenylamine [122-37-2] M 185.2, m 72-73°, pK_{Est} ~10.0. Crystd from chlorobenzene/pet ether.

12-Hydroxydodecanoic acid [505-95-3] M 216.3, m 86-88°, pK_{Est} ~4.8. Crystd from toluene [Sadowik et al. *J Am Chem Soc* 108 7789 1986].

2-Hydroxy-4-(*n*-dodecyloxy)benzophenone [2985-59-3] M 382.5, m 50-52°, pK_{Est} ~7.1. Recryst from *n*-hexane and then 10% (v/v) EtOH in acetonitrile [Valenty et al. *J Am Chem Soc* 106 6155 1984].

***N*-[2-Hydroxyethyl]ethylenediamine** [2-(2-aminoethylamino)ethanol] [111-41-1] M 104.1, b 91.2°/5mm, 238-240°/752mm, n 1.485, d 1.030, pK₁²⁰ 3.75, pK₂²⁰ 9.15. Distilled twice through a Vigreux column. Redistilled from solid NaOH, then from CaH₂. Alternatively, it can be converted to the dihydrochloride and recrystallised from water. It is then dried, mixed with excess of solid NaOH and the free base distilled from the mixture. It is finally redistilled from CaH₂. [Drinkard, Bauer and Bailar *J Am Chem Soc* 82 2992 1960.]

***N*-[2-Hydroxyethyl]ethylenediaminetriacetic acid (HEDTA)** [150-39-0] M 278.3, m 212-214°(dec), pK₁²⁰ 2.51, pK₂²⁰ 5.31, pK₃²⁰ 9.86. Crystd from warm H₂O, after filtering, by addition of 95% EtOH and allowing to cool. The crystals, collected on a sintered-glass funnel, were washed three times with cold absolute EtOH, then again crystd from H₂O. After leaching with cold H₂O, the crystals were dried at 100° under vacuum. [Spedding, Powell and Wheelwright *J Am Chem Soc* 78 34 1956.]

***N*-Hydroxyethyliminodiacetic acid (HIMDA)** [93-62-9] M 177.2, m 181°(dec), pK₁²⁵ 2.16, pK₂²⁵ 8.72, pK₃²⁵ 13.7 (OH). Crystd from water.

2-Hydroxyethylimino-tris(hydroxymethyl)methane (MONO-TRIS) [7343-51-3] M 165.2, m 91°, pK_{Est} ~9.8. Crystd twice from EtOH. Dried under vacuum at 25°.

2-Hydroxyethyl methacrylate [868-77-9] M 130.1, b 67°/3.5mm, d 1.071, n 1.452. Dissolved in water and extracted with *n*-heptane to remove ethylene glycol dimethacrylate (checked by gas-liquid chromatography and by NMR) and distilled twice under reduced pressure [Strop, Mikes and Kalal *J Phys Chem* 80 694 1976].

***N*-2-Hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES)** [7365-45-9] M 238.3, pK²⁰ 7.55. Crystd from hot EtOH and water.

3-Hydroxyflavone [577-85-5] M 238.2, m 169-170°, 171-172°. Recrystd from MeOH, EtOH or hexane. Also purified by repeated sublimation under high vacuum, and dried by high vacuum pumping for at least one hour [Bruker and Kelly *J Phys Chem* 91 2856 1987].

β -Hydroxyglutamic acid [533-62-0] M 163.1, m 100°(dec), pK_1^{25} 2.27, pK_2^{25} 4.29, pK_3^{25} 9.66. Crystd from water.

4-Hydroxyindane [1641-41-1] M 134.2, m 49-50°, b 120°/12mm, pK^{25} 10.32. Crystd from pet ether. Acetyl deriv has m 30-32° (from EtOH), b 127°/14mm. [Dallacker et al. *Chem Ber* 105 2568 1972.]

5-Hydroxyindane [1470-94-6] M 134.2, m 55°, b 255°/760mm, pK_{Est} ~10.4. Crystd from pet ether.

5-Hydroxy-L-lysine monohydrochloride [32685-69-1] M 198.7, $[\alpha]_D^{25}$ +17.8° (6M HCl), pK_2^{25} 8.85, pK_3^{25} 9.83. Likely impurities are 5-*allo*-hydroxy-(D and L)-lysine, histidine, lysine, ornithine. Crystd from water by adding 2-9 volumes of EtOH stepwise.

4-Hydroxy-3-methoxyacetophenone [498-02-2] M 166.2, m 115°, pK_{Est} ~7.9. Crystd from water, or EtOH/pet ether.

4-Hydroxy-3-methoxycinnamic acid (ferulic acid) [1135-24-6] M 194.2, m 174°, pK_1^{25} 4.58, pK_2^{25} 9.39. Crystd from H₂O.

1-Hydroxymethyladamantane [770-71-8] M 166.3, m 115°. Dissolve in Et₂O, wash with aqueous 0.1N NaOH and H₂O, dry over CaCl₂, evaporate and recryst residue from aqueous MeOH. [*Chem Ber* 92 1629 1959.]

17 β -Hydroxy-17 α -methyl-3-androsterone (Mestanolone) [521-11-9] M 304.5°, m 192-193°. Crystd from ethyl acetate.

4-Hydroxy-2-methylazobenzene [1435-88-7] M 212.2, m 100-101°, pK_{Est} ~9.5. Crystd from hexane.

4-Hydroxy-3-methylazobenzene [621-66-9] M 212.2, m 125-126°. Crystd from hexane.

3-Hydroxy-4-methylbenzaldehyde [57295-30-4] M 136.1, m 116-117°, b 179°/15mm, pK_{Est} ~10.2. Crystd from water.

***dl*-2-Hydroxy-2-methylbutyric acid** [3739-30-8] M 118.1, m 72-73°, pK^{25} 3.73. Crystd from *benzene, and sublimed at 90°.

***dl*-2-Hydroxy-3-methylbutyric (α -hydroxyisovaleric) acid** [600-37-3] M 118.1, m 86°, pK_{Est} ~3.9. Crystd from ether/pentane.

***R*- γ -Hydroxymethyl- γ -butyrolactone** [52813-63-5] M 116.1, b 101-102°/0.048mm, d_4^{20} 1.2238, n_D^{20} 1.471, $[\alpha]_{546}^{20}$ -38°, $[\alpha]_D^{20}$ -33° (c 3, EtOH), $[\alpha]_D^{30}$ -53.5° (c 3, EtOH). Purified by column chromatography in Silica gel 60 (Merck 70-230 mesh) and eluting with 7% EtOH-73% CHCl₃. IR (film): 3400 (OH), 1765 (C=O) and 1180 (COC) cm⁻¹. [Eguchi and Kakuta *Bull Chem Soc Jpn* 47 1704 1974; IR and NMR: Ravid et al. *Tetrahedron* 34 1449 1978.]

7-Hydroxy-4-methylcoumarin (4-methylumbelliferone) [90-33-5] M 176.2, m 185-186°, pK_{Est} ~10.0. Crystd from absolute EtOH. (See also entry on p. 548 in Chapter 6.)

2-Hydroxymethyl-12-crown-4 [75507-26-5] M 206.2, d_4^{20} 1.186, n_D^{20} 1.480. Purified by chromatography on Al₂O₃ with EtOAc as eluent to give a *hygroscopic* colourless oil with IR 3418 (OH) and 1103 (COC) cm⁻¹, NMR δ 3.70 (s). [Pugia et al. *J Org Chem* 52 2617 1987.]

***S*-(-)-5-Hydroxymethyl-2(5*H*)-furanone** [78508-96-0] M 114.1, 39-42°, 40-44°, b 130°/0.3mm, $[\alpha]_{546}^{20}$ -180°, $[\alpha]_D^{20}$ -148° (c 1.4, H₂O). It has been purified by chromatography on

Silica gel using hexane-EtOAc (1:1) to give a colourless oil which was distd using a Kügelrohr apparatus and the distillate crystallises on cooling. It has R_F 0.51 on Whatman No 1 paper using pentan-1-ol and 85% formic acid (1:1) and developing with ammoniacal $AgNO_3$. [*Boll Acta Chem Scand* **22** 3245 1968; NMR: Oppolzer et al. *Helv Chim Acta* **68** 2100 1985.]

5-(Hydroxymethyl)furfural [67-47-0] M 126.1, m 33.5°, b 114-116°/1mm. Crystd from diethyl ether/pet ether.

3-Hydroxy-3-methylglutaric acid (Meglutol) [503-49-1] M 162.1, m 99-102°, 108-109°, 100°, $pK_{Est(1)} \sim 4.0$, $pK_{Est(2)} \sim 5.0$. Recrystd from diethyl ether/hexane and dried under vac at 60° for 1h.

dl-3-Hydroxy-N-methylmorphinan [297-90-5] M 257.4, m 251-253°. Crystd from anisole + aqueous EtOH.

6-Hydroxy-2-methyl-1,4-naphthaquinone [633-71-6] M 188.2, $pK_{Est} \sim 10.0$. Crystd from aqueous EtOH. Sublimes on heating.

4-Hydroxy-4-methyl-2-pentanone [123-42-2] M 116.2, b 166°, d 0.932, n 1.4235, $n^{25} 1.4213$. Loses water when heated. Can be dried with $CaSO_4$, then fractionally distd under reduced pressure.

17 α -Hydroxy-6 α -methylprogesterone (Medroxyprogesterone) [520-85-4] M 344.5, m 220°, $[\alpha]_D^{25} +75^\circ$. Crystd from chloroform.

2-Hydroxy-2-methylpropionic acid (α -hydroxyisobutyric acid, 2-methylactic acid) [594-61-6] M 104.1, m 79°, b 114°/12mm, 212°/760mm, $pK^{25} 3.78$. Distd in steam, crystd from diethyl ether or *benzene, sublimed at 50° and dried under vacuum.

8-Hydroxy-2-methylquinoline [826-81-3] M 159.2, m 74-75°, b 266-267°, $pK_1^{25} 5.61$, $pK_2^{25} 10.16$. Crystd from EtOH or aqueous EtOH.

2-Hydroxy-1-naphthaldehyde [708-06-5] M 172.2, m 82°, b 192°/27mm, $pK_{Est} \sim 7.8$. Crystd from EtOH (1.5mL/g), ethyl acetate or water.

2-Hydroxy-1-naphthaleneacetic acid [10441-45-9] M 202.2, $pK_{Est(1)} \sim 4.2$, $pK_{Est(2)} \sim 8.3$. Treated with activated charcoal and crystd from EtOH/water (1:9, v/v). Dried under vacuum, over silica gel, in the dark. Stored in the dark at -20° [Gafni, Modlin and Brand *J Phys Chem* **80** 898 1976]. Forms a lactone (m 107°) readily.

6-Hydroxy-2-naphthalenepropionic acid [553-39-9] M 216.2, m 180-181°, $pK_{Est(1)} \sim 4.6$, $pK_{Est(2)} \sim 9.0$. Crystd from aqueous EtOH or aqueous MeOH.

3-Hydroxy-2-naphthalide [92-77-3] M 263.3, m 248.0-248.5°, CI 37505. Crystd from xylene [Schnopper, Broussard and La Forgia *Anal Chem* **31** 1542 1959].

3-Hydroxy-2-naphtho-4'-chloro-o-toluidide [92-76-2] M 311.8, m 243.5-244.5°. Crystd from xylene [Schnopper, Broussard and La Forgia *Anal Chem* **31** 1542 1959].

3-Hydroxy-2-naphthoic-1'-naphthylamide [123-68-3] M 314.3, m 217-.5-218.0°. Crystd from xylene [Schnopper, Broussard and La Forgia *Anal Chem* **31** 1542 1959].

3-Hydroxy-2-naphthoic-2'-naphthylamide [136-64-8] M 305.3, m 243.5-244.5°, and other naphthol AS derivatives. Crystd from xylene [Schnopper, Broussard and La Forgia *Anal Chem* **31** 1542 1959].

2-Hydroxy-1,4-naphthaquinone [83-72-7] M 174.2, m 192°(dec), $pK_1^{2.5}$ -5.6 (C=O protonation), $pK_2^{2.5}$ 4.00 (phenolic OH). Crystd from *benzene.

5-Hydroxy-1,4-naphthaquinone (Juglone) [481-39-0] M 174.2, m 155°, 164-165°, pK 8.7. Crystd from *benzene/pet ether or pet ether.

6-Hydroxy-2-naphthyl disulfide [6088-51-3] M 350.5, m 221-222°, 226-227°, pK_{Est} ~9.0. Crystallises as leaflets from AcOH and is slightly soluble in EtOH, and AcOH, but is soluble in *C₆H₆ and in alkalis to give a yellow soln. [Zincke and Dereser *Chem Ber* 51 352 1918.] The *acetoxy* derivative has m 198-200° (from AcOH or dioxane-MeOH) and the *diacetyl* derivative has m 167-168° (from AcOH). A small amount of impure disulfide can be purified by dissolving in a small volume of Me₂CO and adding a large volume of toluene, filter rapidly and concentrate to one third of its volume. The hot toluene soln is filtered rapidly from any tarry residue, and crystals separate on cooling. After recrystn from hot acetic acid gives crystals m 220-223° [Barrett and Seligman *Science* 116 323 1952].

2-Hydroxy-5-nitrobenzyl bromide [772-33-8] M 232.0, m 147°, pK_{Est} ~8.0. Crystd from *benzene or *benzene/ligroin.

4-Hydroxy-2-n-nonylquinoline N-oxide [316-66-5] M 287.4, m 148-149°, pK_{Est} ~6.0. Crystd from EtOH.

N-Hydroxy-5-norbornene-2,3-dicarboxylic acid imide [21715-90-2] M 179.2, m 165-166°, 166-169°, pK_{Est} ~6. Dissolve in CHCl₃, filter, evaporate and recrystallise from EtOAc. IR (nujol): 1695, 1710 and 1770 (C=O), and 3100 (OH) cm⁻¹. *O*-Acetyl derivative has m 113-114° (from EtOH) with IR bands at 1730, 1770 and 1815 cm⁻¹ only, and the *O*-benzoyl derivative has m 143-144° (from propan-2-ol or *C₆H₆). [Bauer and Miarka *J Org Chem* 24 1293 1959; Fujino et al. *Chem Pharm Bull Jpn* 22 1857 1974].

DL-erythro-3-Hydroxynorvaline (2-amino-3-hydroxypentanoic acid) [34042-00-7] M 133.2, m 257-259° (dec), 263° (dec), $pK_1^{2.0}$ 2.32, $pK_2^{2.0}$ 9.12. Purified by recrystn from aqueous EtOH. The *Cu salt* has m 255-256° (dec), the *benzoyl* derivative has m 181°, and the *N*-phenylcarbamoyl derivative has m 164°. [Buston et al. *J Biol Chem* 204 665 1953].

2-Hydroxyoctanoic acid (2-hydroxycaprylic acid) [617-73-2] M 160.2, m 69.5°, b 160-165°/10mm, pK_{Est} ~3.7. Crystd from EtOH/pet ether or ether/ligroin.

1-Hydroxyphenazine (Hemipyocyanine) [528-71-2] M 196.2, m 157-158°, $pK_1^{2.5}$ 1.61, $pK_2^{1.5}$ 8.33. Chromatographed on acidic alumina with *benzene/ether. Crystd from *benzene/heptane, and sublimed.

2-Hydroxyphenylacetic acid [614-75-5] M 152.2, m 148-149°, b 240-243°/760 mm, $pK_{Est(1)}$ ~4.3, $pK_{Est(2)}$ ~10.1. Crystd from ether or chloroform (m from latter is always lower).

3-Hydroxyphenylacetic acid [621-37-4] M 152.2, m 137°, $pK_{Ext(1)}$ ~4.3, $pK_{Ext(2)}$ ~10. Crystd from *benzene/ligroin.

4-Hydroxyphenylacetic acid [156-38-7] M 152.2, m 150-151°, 152°, pK_1 4.28, pK_2 10.1. Crystd from water or Et₂O/pet ether.

2-(2-Hydroxyphenyl)benzothiazole [3411-95-8] M 227.2, m 132-133°, b 173-179°/3mm. Recrystd several times from aqueous EtOH and by sublimation. [Itoh and Fujiwara *J Am Chem Soc* 107 1561 1985.]

2-(2-Hydroxyphenyl)benzoxazole [835-64-3] M 211.2, m 127°, b 338°/760mm. Recrystd several times from aqueous EtOH and by sublimation. [Itoh and Fujiwara *J Am Chem Soc* 107 1561 1985.]

3-Hydroxy-2-phenylcinchoninic acid [485-89-2] M 265.3, m 206-207°(dec). Crystd from EtOH.

N-(*p*-Hydroxyphenyl)glycine [22818-40-2] M 167.2, m >240°(dec), pK_{Est(1)}~2, pK_{Est(2)}~4.5, pK_{Est(3)}~10.3. Crystd from water.

N-(4-Hydroxyphenyl)-3-phenylsalicylamide [550-57-2] M 305.3, m 183-184°, pK_{Est} ~9.5. Crystd from aqueous MeOH.

L-2-Hydroxy-3-phenylpropionic acid (3-phenyl lactic acid) [20312-36-1] M 166.2, m 125-126°, [α]_D¹⁷ -18.7° (EtOH), pK see below. Crystd from water, MeOH, EtOH or *benzene.

dl-2-Hydroxy-3-phenylpropionic acid [828-01-3] M 166.2, m 97-98°, b 148-150°/15mm, pK_{Est} ~3.7. Crystd from *benzene or chloroform.

3-*p*-Hydroxyphenylpropionic acid (phloretic acid) [501-97-3] M 166.2, m 129-130°, 131-133°, pK_{Est(1)}~4.7, pK_{Est(2)}~10.1. Crystd from ether or H₂O.

p-Hydroxyphenylpyruvic acid [156-39-8] M 180.2, m 220°(dec), pK_{Est} ~2.3. Crystd three times from 0.1M HCl/EtOH (4:1, v/v) immediately before use [Rose and Powell *Biochem J* 87 541 1963], or from Et₂O. The 3,4-Dinitrophenylhydrazone has m 178°.

N-Hydroxyphthalimide [524-38-9] M 163.1, m 230°, ~235° (dec), 237-240°, pK³⁰ 7.0. Dissolve in H₂O by adding Et₃N to form the salt and while hot acidify, cool and pour into a large volume of H₂O. Filter off the solid, wash with H₂O, dry over P₂O₅ in vacuum. [Nefken And Teser *J Am Chem Soc* 83 1263 1961; Fieser 1 485 1976; Nefkens et al. *Recl Trav Chim Pays-Bas* 81 683 1962] The *O*-acetyl derivative has m 178-180° (from EtOH).

3-β-Hydroxy-5-pregnen-20-one (pregnenolone) [145-13-1] M 316.5, m 189-190°, [α]_D²⁰ +30° (EtOH), [α]₅₄₆ +34° (c 1, EtOH). Crystd from MeOH.

17α-Hydroxyprogesterone [604-09-1] M 330.5, m 222-223°, [α]₅₄₆²⁰ +141° (c 2, dioxane), λ_{max} 240nm. Crystd from acetone or EtOH. *Acetate*: m 239-240° and *caproate*: m 119-121° crystallised from CHCl₃/MeOH.

R-(+)-3-Hydroxyprolidine [2799-21-5] M 87.1, b 215-216°, d₄²⁰ 1.078, n_D²⁰ 1.490, [α]_D²⁰ +6.5° (c 1.5, MeOH), pK_{Est} ~10.1. Purify by repeated distn. The *hydrochloride* has -ve rotation and the *dimethiodide* has m 230° and [α]_D²⁴ -8.02°. [Suyama and Kanno *Yakugaku Zasshi (J Pharm Soc Japan)* 85 531 1965; Uno et al. *J Heterocycl Chem* 24 1025 1987; Flanagan and Joullie *Heterocycles* 26 2247 1987.]

trans-L-4-Hydroxyproline [51-35-4] M 131.1, m 274°, [α]_D²⁰ -76.0° (c 5, H₂O), pK₁²⁵ 1.86, pK₂²⁵ 9.79. Crystd from MeOH/EtOH (1:1). Separation from normal *allo*-isomer can be achieved by crystn of the copper salts [see *Biochem Prep* 8 114 1961].

4'-Hydroxypropiophenone [70-70-2] M 150.2, m 149°, pK_{Est} ~10. Crystd from water.

2-(α-Hydroxypropyl)piperidine (2-piperidinepropanol) [24448-89-3] M 143.2, m 121°, b 226°, pK_{Est} ~10.2. Crystd from ether.

7-(2-Hydroxypropyl)theophylline (Proxiphylline) [603-00-9] M 238.2, m 135-136°. Crystd from EtOH.

6-Hydroxypurine (hypoxanthine) [68-94-0] M 136.1, m 150°(dec), pK₁²⁰ 8.96, pK₂²⁰ 12.18. Crystd from hot water. Dried at 105°.

2-Hydroxypyridine (2-pyridone) [142-08-5] M 95.1, m 105-107°, b 181-185°/24mm, ε_{293nm} 5900 (H₂O) pK₁²⁵ 1.25, pK₂²⁵ 11.99. Distd under vacuum to remove coloured impurity, then crystd from

*benzene, CCl₄, EtOH or CHCl₃/diethyl ether. It can be sublimed under high vacuum. [DePue et al. *J Am Chem Soc* **107** 2131 1985.]

3-Hydroxypyridine [109-00-2] M 95.1, m 129° pK₁²⁵ 5.10, pK₂²⁵ 8.6. Crystd from water or EtOH.

4-Hydroxypyridine (4-pyridone) [626-64-2] M 95.1, m 65°(hydrate), 148.5° (anhydr), b >350°/760mm, pK₁²⁰ 3.20, pK₂²⁰ 11.12. Crystd from H₂O. Loses H₂O on drying *in vacuo* over H₂SO₄. Stored over KOH because it is *hygroscopic*.

2(6)-Hydroxypyridine-5(3)-carboxylic acid (6-hydroxynicotinic acid) [5006-66-6] M 139.1, m 304°(dec), pK²⁰ 3.82. Crystd from water.

4-Hydroxypyridine-2,6-dicarboxylic acid (chelidamic acid) [138-60-3] M 183.1, m 254°(dec), pK₁²² 1.9, pK₂²² 3.18, pK₃²² 10.85. Crystd from water.

2-Hydroxypyrimidine [557-01-7] M 96.1, m 179-180°, pK₁²⁰ 2.15, pK₂²⁰ 9.2. Crystd from EtOH or ethyl acetate.

4-Hydroxypyrimidine [4562-27-0] M 96.1, m 164-165°, pK₁²⁰ 1.66, pK₂²⁰ 8.63. Crystd from *benzene or ethyl acetate.

2-Hydroxypyrimidine hydrochloride [38353-09-2] M 132.5, m 205°(dec). Crystd from EtOH.

2-Hydroxyquinoline (carbostyryl) [59-31-4] M 145.2, m 199-200°, pK₁²⁰ -0.31, pK₂²⁰ 11.76. Crystd from MeOH.

8-Hydroxyquinoline (oxine, 8-quinolinol) [148-24-3] M 145.2, m 71-73°, 75-76°, 76°, b ~ 267° pK₁²⁵ 4.91, pK₂²⁵ 9.81. Crystd from hot EtOH, acetone, pet ether (b 60-80°) or water. Crude oxine can be purified by pptn of copper oxinate, followed by liberation of free oxine with H₂S or by steam distn after acidification with H₂SO₄. Stored in the dark. Forms metal complexes. [Manske et al. *Can J Research* **27F** 359 1949; Phillips *Chem Rev* **56** 271 1956.]

8-Hydroxyquinoline-5-sulfonic acid (H₂O) [84-88-8] M 243.3, m >310° pK₁²⁵ 4.09, pK₂²⁵ 8.66. Crystd from water or dil HCl (*ca* 2% by weight).

5-Hydroxysalicylic acid [490-79-9] M 154.1. See 2,5-dihydroxybenzoic acid on p. 207.

trans-4-Hydroxystilbene [6554-98-9] M 196.3, m 189°. Crystd from *benzene or acetic acid.

N-Hydroxysuccinimide [6066-82-6] M 115.1, m 96-98°, pK 6.0. Recrystd from EtOH/ethyl acetate [Manesis and Goodmen *J Org Chem* **52** 5331 1987].

dl-2-Hydroxytetradecanoic acid [2507-55-3] M 244.4, m 81-82°, pK_{Est} ~3.7. Crystd from chloroform.

R-2-Hydroxytetradecanoic acid [26632-17-7] M 244.4, m 88-2-88.5°, [α]_D²⁰ -31° (CHCl₃). Crystd from chloroform.

4-Hydroxy-2,2,6,6-tetramethylpiperidine [2403-88-5] M 157.3, m 130-131°, pK 10.05. Crystd from water as hydrate, and crystd from ether as the anhydrous base.

Hydroxy(tosyloxy)iodobenzene [phenyl(hydroxy)tosyloxyiodine, hydroxy(4-methylbenzenesulfonato-O)phenyliodine, Koser's reagent] [27126-76-7] M 392.2, m 134-136°, 135-138°, 134-136°, 136-138.5°. Possible impurities are tosic acid (removed by washing with Me₂CO) and acetic acid (removed by washing with Et₂O). It is purified by dissolving in the minimum vol of MeOH, adding

Et₂O to cloud point and setting aside for the prisms to separate [Koser and Wettach *J Org Chem* **42** 1476 1977; NMR: Koser et al. *J Org Chem* **41** 3609 1976]. It has also been crystd from CH₂Cl₂ (needles, m 140-142°) [Neiland and Karele *J Org Chem, USSR (Engl Transl)* **6** 889 1970].

4(6)-Hydroxy-2,5,6(2,4,5)-triaminopyrimidine sulfate [35011-47-3] M 257.22, m >340°, pK₁ 2.0, pK₂ 5.1, pK₃ 10.1. This salt has very low solubility in H₂O. It is best purified by conversion into the dihydrochloride salt which is then reconverted to the insoluble sulfate salt. The sulfate salt (2.57g, 10mmoles) is suspended in H₂O (20mL) containing BaCl₂ (10mmoles) and stirred in a boiling water bath for 15min. After cooling the insoluble BaSO₄ is filtered off and washed with boiling H₂O (10mL). The combined filtrate and washings are made acidic with HCl and evaporated to dryness. The residual hydrochloride salt is recrystd from H₂O by adding conc HCl whereby the *dihydrochloride salt* separates as clusters which darken at 260° and dec > 300° [Baugh and Shaw *J Org Chem* **29** 3610 1964; King and Spengley *J Chem Soc* 2144 1952]. The hydrochloride is then dissolved in H₂O and while hot an equivalent of H₂SO₄ is added when the sulfate separates as a white microcrystalline solid which is filtered off washed liberally with H₂O and dried in vacuum over P₂O₅. [Albert and Wood *J Appl Chem London* **3** 521 1953; UV: Cavalieri et al. *J Am Chem Soc* **70** 3875 1948; see also Pfeleiderer *Chem Ber* **90** 2272 1957; Traube *Chem Ber* **33** 1371 1900].

9-Hydroxytryptycene [73597-16-7] M 270.3, m 245-246.5°. Crystd from *benzene/pet ether. Dried at 100° in a vacuum [Imashiro et al. *J Am Chem Soc* **109** 729 1987].

5-Hydroxy-L-tryptophan [4350-09-8] M 220.2, m 273°(dec), [α]_D²² -32.5°, [α]₅₄₆²⁰ -73.5° (c 1, H₂O), pK_{Est(1)}~2.4, pK_{Est(2)}~9.0, pK_{Est(3)}~9.4, pK_{Est(4)} 16 (NH). Likely impurities are 5-hydroxy-D-tryptophan and 5-benzoyloxytryptophan. Crystd under nitrogen from water by adding EtOH. Stored under nitrogen.

Hydroxyurea [127-07-1] M 76.1, m 70-72° (unstable form), m 135-140°, 141° (stable form). See hydroxyurea on p. 431 in Chapter 5.

3-Hydroxyxanthone [3722-51-8] M 212.2, m 246°. Purified by chromatography on SiO₂ gel with pet ether/*benzene). Recrystd from *benzene or EtOH [Itoh et al. *J Am Chem Soc* **107** 4819 1985].

α-Hyodeoxycholic acid [83-49-8] M 392.6, m 196-197°, [α]₅₄₆²⁰ +8° (c 2, EtOH), pK_{Est} ~4.9. Crystd from ethyl acetate.

Hyoscine (scopolamine, atropine) [51-34-3] M 321.4, m 59°, [α]_D²⁰ -18° (c 5, EtOH), -28° (c 2, H₂O), [α]₅₄₆²⁰ -30° (c 5, CHCl₃), pK²⁵ 7.55. Crystd from *benzene/pet ether. Racemate has m 56-57° (H₂O), 37-38° (2H₂O), syrup (anhydr), *l* and *d* isomers can separate as syrups when anhydrous.

Hypericin [548-04-9] M 504.4, m 320°(dec). Crystd from pyridine by addition of methanolic HCl.

Ibogaine [83-74-9] M 300.3, m 152-153°, [α]_D²⁰ -54° (EtOH), pK 8.1 (80% aq MeOCH₂CH₂OH). Crystd from aqueous EtOH and sublimates at 150°/0.01mm.

Imidazole (glyoxaline) [288-32-4] M 68.1, m 89.5-91°, b 256°, pK₁²⁵ 6.99, pK₂²⁵ 14.44. Crystd from *benzene, CCl₄, CH₂Cl₂, EtOH, pet ether, acetone/pet ether and distd deionized water. Dried at 40° under vacuum over P₂O₅. Distd at low pressure. Also purified by sublimation or by zone melting. [Caswell and Spiro *J Am Chem Soc* **108** 6470 1986.] ¹⁵N-imidazole was crystd from *benzene [Scholes et al. *J Am Chem Soc* **108** 1660 1986].

4'-(Imidazol-1-yl)acetophenone [10041-06-2] M 186.2, m 104-107°, pK 4.54. Twice recrystd from CH₂Cl₂/hexane [Collman et al. *J Am Chem Soc* **108** 2588 1986].

Iminodiacetic acid [142-73-4] M 133.1, m 225°(dec), pK₁²⁵ 2.50, pK₂²⁵ 9.40. Crystd from water.

1,3-Indandione [606-23-5] M 146.2, m 129-132°, pK¹⁸ 7.2 (1% aq EtOH). Recrystd from EtOH [Bernasconi and Paschalis *J Am Chem Soc* 108 2969 1986].

Indane [496-11-7] M 118.1, b 177°, d 0.960, n 1.538. Shaken with conc H₂SO₄, then water, dried and fractionally distd.

Indanthrene [81-77-6] M 442.4, m 470-500°. Crystd repeatedly from 1,2,4-trichlorobenzene.

Indazole [271-44-3] M 118.1, m 147°, pK₁²⁰ 1.32, p K₂²⁵ 13.80 (acidic NH). Crystd from water, sublimed under a vacuum, then pet ether (b 60-80°).

Indene [95-13-6] M 116.2, f -1.5°, b 114.5°/100mm, d 0.994, n 1.5763. Shaken with 6M HCl for 24h (to remove basic nitrogenous material), then refluxed with 40% NaOH for 2h (to remove benzonitrile). Fractionally distd, then fractionally crystd by partial freezing. The higher-melting portion was converted to its sodium salt by adding a quarter of its weight of sodamide under nitrogen and stirring for 3h at 120°. Unreacted organic material was distd off at 120°/1mm. The sodium salts were hydrolysed with water, and the organic fraction was separated by steam distn, followed by fractional distn. Before use, the distillate was passed, under nitrogen, through a column of activated silica gel. [Russell *J Am Chem Soc* 78 1041 1956.]

Indigo [482-89-3] M 262.3, sublimes at ~300°, m 390°(dec), and halogen-substituted indigo dyes. Reduced in alkaline soln with sodium hydrosulfite, and filtered. The filtrate was then oxidised by air, and the resulting ppt was filtered off, dried at 65-70°, ground to a fine powder, and extracted with CHCl₃ in a Soxhlet extractor. Evapn of the CHCl₃ gave the purified dye. [Brode, Pearson and Wyman *J Am Chem Soc* 76 1034 1954; spectral characteristics are listed.]

Indole [120-72-9] M 117.2, m 52°, b 124°/5mm, 253-254°/760mm, pK₁²⁵-2.47 (H₀ scale), pK₂²⁵ 16.97 (acidic NH). Crystd from *benzene, hexane, water or EtOH/water (1:10). Further purified by sublimation in a vacuum or zone melting.

Indole-3-acetic acid [87-51-4] M 175.2, m 167-169°, pK₁²⁵-6.13 (aq H₂SO₄), pK₂²⁵ 4.54 (CO₂H). Recrystd from EtOH/water [James and Ware *J Phys Chem* 89 5450 1985].

3-Indoleacetonitrile [771-51-7] M 156.2, m 33-36°, 36-38°, b 157°/0.2mm, 158-160°/0.1mm, viscous oil n_D²⁰ 1.6097. Distil in very high vacuum and the viscous distillate crystallises on standing after a few days; the *picrate* has m 127-128° (from EtOH) [Coker et al. *J Org Chem* 27 850 1962; Thesing and Schülde *Chem Ber* 85 324 1952]. The *N-acetate* has m 118° (from MeOH) and has R_F = 0.8, on Silica Gel F₂₅₄ in CHCl₂-MeOH 19:1 [Buzas et al. *Synthesis* 129 1977].

Indole-3-butanoic acid [133-32-4] M 203.2, m 124-125°. See 3-indolylbutyric acid on p. 543 in Chapter 6.

Indole-3-propionic acid [830-96-6] M 189.2, m 134-135°, pK_{Est} ~4.7. Recrystd from EtOH/water [James and Ware *J Phys Chem* 89 5450 1985].

Indolizine [pyrrocoline, pyrrolo(1,2-*a*)pyridine] [274-40-8] M 117, m 73-74°, 75°, pK²⁰ 3.94 (C-protonation). Purified through an alumina column in *C₆H₆ and eluted with *C₆H₆ (toluene could be used instead). The eluate contained in the fluorescent band (using UV light λ 365nm) was collected, evapd and the cryst residues sublimed twice at 40-50°/0.2-0.5mm. The colourless crystals darkend on standing and should be stored in dark sealed containers. If the original sample is dark in color then it should be covered with water and steam distd. The crystals in the distillate are collected and, dried between filter paper and sublimed. It protonates on C3 in aqueous acid. It should give one fluorescent spot on paper chromatography (Whatman 1) in 3% aq ammonia and in *n*-BuOH, AcOH, H₂O (4:1:1). The *picrate* has m 101° from EtOH. [Armarego *J Chem Soc* 226 1944; Armarego *J Chem Soc (B)* 191 1966; Scholtz *Chem Ber* 45 734 1912.]

(-)-Inosine [58-63-9] M 268.2, m 215°, $[\alpha]_{546}^{20} -76^\circ$ (c 1, 0.1M NaOH), $pK_1^{25} 1.06$, $pK_2^{25} 8.96$, $pK_3^{25} 11.36$. Crystd from aqueous 80% EtOH.

i-Inositol (*myo*) [87-89-8] M 180.2, m 228°. See entry on p. 543 in Chapter 6.

Inositol monophosphate [15421-51-9] M 260.1, m 195-197°(dec). Crystd from water and EtOH.

Iodinine (1,6-phenazine-5,10-dioxide) [68-81-5] M 244.1, m 236°(dec), pK 12.5. Crystd from CHCl₃.

Inulin [9005-80-5] M (162.14)_n. Crystd from water.

Iodoacetamide [144-48-9] M 185.0, m ca 143°(dec). Crystd from water or CCl₄.

Iodoacetic acid [64-69-7] M 160.6, m 78°, $pK^{25} 3.19$. Crystd from pet ether (b 60-80°) or CHCl₃/CCl.

2-Iodoaniline [615-43-0] M 219.0, m 60-61°, $pK^{25} 2.54$. Distd with steam and crystd from *benzene/pet ether.

4-Iodoaniline [540-37-4] M 219.0, m 62-63°, $pK^{25} 3.81$. Crystd from pet ether (b 60-80°) by refluxing, then cooling in an ice-salt bath freezing mixture. Dried in air. Also crystd from EtOH and dried in a vacuum for 6h at 40° [Edidin et al. *J Am Chem Soc* 109 3945 1987].

4-Iodoanisole [696-62-8] M 234.0, m 51-52°, b 139°/35mm, 237°/726mm. Crystd from aqueous EtOH.

Iodobenzene [591-50-4] M 204.0, b 63-65°/10mm, 188°/atm, d 1.829, $n^{25} 1.6169$. Washed with dilute aqueous Na₂S₂O₃, then water. Dried with CaCl₂ or CaSO₄. Decolorised with charcoal. Distd under reduced pressure and stored with mercury or silver powder to stabilise it.

o-Iodobenzoic acid [88-67-5] M 248.4, m 162°, $pK^{20} 2.93$. Crystd repeatedly from water and EtOH. Sublimed under vacuum at 100°.

m-Iodobenzoic acid [618-51-9] M 248.4, m 186.6-186.8°, $pK^{25} 3.85$. Crystd repeatedly from water and EtOH. Sublimed under vacuum at 100°.

p-Iodobenzoic acid [619-58-9] M 248.4, m 271-272°, $pK^{25} 4.00$. Crystd repeatedly from water and EtOH. Sublimed under vacuum at 100°.

4-Iodobiphenyl [1591-31-7] M 280.1, m 113.7-114.3°. Crystd from EtOH/*benzene and dried under vacuum over P₂O₅.

2-Iodobutane [513-48-4] M 184.0, b 120.0, d 1.50, $n^{25} 1.4973$. Purified by shaking with conc H₂SO₄, then washing with water, aq Na₂SO₃ and again with water. Dried with MgSO₄ and distd. Alternatively, passed through a column of activated alumina before distn, or treated with elemental bromine, followed by extraction of the free halogen with aqueous Na₂S₂O₃, thorough washing with water, drying and distilling. It is stored over silver powder and distd before use.

1-Iodo-2,4-dinitrobenzene [709-49-9] M 294.0, m 88°. Crystd from ethyl acetate.

Iodoform [75-47-8] M 393.7, m 119°. Crystd from MeOH, EtOH or EtOH/EtOAc. Steam volatile.

1-Iodo-4-nitrobenzene [636-98-6] M 249.0, m 171-172°. Pptd from acetone by addition of water, then recrystd from EtOH.

o-Iodophenol [533-58-4] M 280.1, m 42°, pK²⁵ 8.51. Crystd from CHCl₃ or diethyl ether.

p-Iodophenol [540-38-5] M 280.1, m 94°, 138-140°/5mm, pK²⁵ 9.30. Crystd from pet ether (b 80-100°) or distd *in vacuo*. If material has a brown or violet color, dissolve in CHCl₃, shake with 5% sodium thiosulfate soln until the CHCl₃ is colorless. Dry (Na₂SO₄), extract, evap and dist residue *in vacuo*. [Dains and Eberly *Org Synth Coll Vol II*, 355 1948.]

5-Iodosalicylic acid (2-hydroxy-5-iodobenzoic acid) [119-30-2] M 264.0, m 197° pK₁²⁵ 2.65, pK₂²⁵ 13.05. Crystd from water.

o-Iodosobenzoic acid [304-91-6] M 264.0, m >200°, pK_{Est} ~2.6. Crystd from EtOH.

N-Iodosuccinimide [516-12-1] M 225.0, m 200-201°. Crystd from dioxane/CCl₄.

p-Iodotoluene [624-31-7] M 218.0, m 35°, b 211-212°. Crystd from EtOH.

3-Iodo-L-tyrosine [70-78-0] M 307.1, m 205-208°(dec), [α]_D²⁵ -4.4° (c 5, 1M HCl), pK_{Est(2)} ~2.1, pK_{Est(3)} ~6.4, pK₄²⁵ 8.7. Likely impurities are tyrosine, diiodotyrosine and iodide. Crystd by soln in dilute ammonia, at room temperature, followed by addition of dilute acetic acid to pH 6. Stored at 0°.

α-Ionone [127-41-3] M 192.3, b 131°/13mm, d 0.931, n 1.520, [α]_D²³ +347° (neat). Purified on a spinning band fractionating column.

β-Ionone [79-77-6] M 192.3, b 150-151°/24mm, d 0.945, n 1.5211, ε_{296nm} 10,700. Converted to the *semicarbazone* (m 149°) by adding 50g of semicarbazide hydrochloride and 44g of potassium acetate in 150mL of water to a soln of 85g of β-ionone in EtOH. (More EtOH was added to redissolve any β-ionone that ppted.) The semicarbazone crystallised on cooling in an ice-bath and was recrystallised from EtOH or 75% MeOH to constant m (148-149°). The semicarbazone (5g) was shaken at room temperature for several days with 20mL of pet ether and 48mL of M H₂SO₄, then the ether layer was washed with water and dilute aqueous NaHCO₃, dried and the solvent was evaporated. The β-ionone was distilled under vacuum. (The customary steam distillation of β-ionone semicarbazone did not increase the purity.) [Young et al. *J Am Chem Soc* 66 855 1944].

Iproniazid (isonicotinic acid 2-isopropylhydrazide) phosphate [305-33-9] M 277.2, m 178-179°, 180-182°, pK_{Est} ~3.5 (free base). Crystd from H₂O and Me₂CO. Free base has m 113-114° from *C₆H₆/pet ether.

(±)-Irone (6-methyl-ionone, ±-*trans*-(α)-4*t*-[2,5,6,6-tetramethyl-cyclohex-2-yl]but-3*t*-en-2-one) [79-69-6] M 206.3, b 85-86°/0.05mm, 109°/0.7mm, d₄²⁰ 0.9340, n_D²⁰ 1.4998. If large amounts are available then fractionate through a Podbielniak column (see p. 141) or an efficient spinning band column, but small amounts are distilled using a Kügelrohr apparatus. The 4-*phenyl-semicarbazone* has m 174-175° (165-165.5°). [IR: Seidel and Ruzocka *Helv Chim Acta* 35 1826 1952; Naves *Helv Chim Acta* 31 1280 1948; Lecomte and Naves *J Chim Phys* 53 462 1956.]

Isatin (indole-2,3-dione) [91-56-5] M 147.1, m 201-203°, 205°, pK >12 (acidic NH). Crystd from amyl alcohol and sublimed at 180°/1mm. In aq NaOH the ring opens to yield sodium *o*-aminobenzoylformate.

Isatoic anhydride (3,1-benzoxazin-2,4[1-*H*]-dione) [118-48-9] M 163.1, m 235-240°, 240-243°, 243°, 243-245°. Recryst from EtOH or 95% EtOH (30mL/g) or dioxane (10mL/g) and dried in a vacuum. [Wagner and Fegley *Org Synth Coll Vol III* 488 1955; Ben-Ishai and Katchalski *J Am Chem Soc* 74 3688 1952; UV: Zentmyer and Wagner *J Org Chem* 14 967 1949.]

Isoamyl acetate (1-butyl-3-methyl acetate) [123-92-2] M 130.2, b 142.0°, d 0.871, n 1.40535. Dried with finely divided K₂CO₃ and fractionally distd.

Isoamyl alcohol (1-butyl-3-methyl alcohol) [123-51-3] M 88.2, b 132°/760mm, d¹⁵ 0.8129, n¹⁵ 1.4085. See 3-methyl-1-butanol on p. 290.

Isoamyl bromide (1-butyl-3-methyl bromide) [107-82-4] M 151.1, f -112°, b 119.2°/737mm, d 1.208, n 1.444. Shaken with conc H₂SO₄, washed with water, dried with K₂CO₃ and fractionally distd.

Isoamyl chloride (1-butyl-3-methyl chloride) [513-36-0] M 106.6, b 99°/734mm, d 0.8704, n 1.4084. Shaken vigorously with 95% H₂SO₄ until the acid layer no longer became coloured during 12h, then washed with water, saturated aq Na₂CO₃, and more water. Dried with MgSO₄, filtered and fractionally distd. Alternatively, a stream of oxygen containing 5% of ozone was passed through the chloride for a time, three times longer than was necessary to cause the first coloration of starch iodide paper by the exit gas. Subsequent washing of the liquid with aqueous NaHCO₃ hydrolysed the ozonides and removed organic acids. After drying and filtering, the isoamyl chloride was distd. [Chien and Willard *J Am Chem Soc* 75 6160 1953.]

Isoamyl ether [diisopentyl ether, di-(1-butyl-3-methyl) ether] [544-01-4] M 158.3, b 173.4°, d 0.778, n 1.40850. This is a mixture of 2- and 3-methylbutyl ether. It is purified by refluxing with sodium for 5h, then distilled under reduced pressure, to remove alcohols. Isoamyl ether can also be dried with CaCl₂ and fractionally distd from P₂O₅.

D(-)-Isoascorbic acid (araboascorbic acid) [89-65-6] M 176.1, m 174°(dec), [α]_D²⁵ -16.8° (c 2, H₂O), pK¹⁸ 3.99. Crystd from H₂O or dioxane.

dl-Isoborneol [124-76-5] M 154.3, m 212° (sealed tube). Crystd from EtOH or pet ether (b 60-80°). Sublimes in a vacuum.

Isobutane [75-28-5] M 58.1, b -10.2°, d 0.557. Olefines and moisture can be removed by passage at 65° through a bed of silica-alumina catalyst which has previously been evacuated at about 400°. Alternatively, water and CO₂ can be taken out by passage through P₂O₅ then asbestos impregnated with NaOH. Treatment with anhydrous AlBr₃ at 0° then removes traces of olefins. Inert gases can be separated by freezing the isobutane at -195° and evacuating out the system.

Isobutene [115-11-7] M 56.1, b -6.6°/760mm. Dried by passage through anhydrous CaSO₄ at 0°. Purified by freeze-pump-thaw cycles and trap-to-trap distn.

Isobutyl alcohol (2-methyl-1-propanol) [78-83-1] M 74.1, b 108°/760mm, d 0.801, n 1.396. Dried with K₂CO₃, CaSO₄ or CaCl₂, filtered and fractionally distd. For further drying, the redistd alcohol can be refluxed with the appropriate alkyl phthalate or succinate as described under *ethanol* (see also p. 271).

Isobutyl bromide (1-bromo-2-methylpropane) [78-77-3] M 137.0, b 91.2°, d 1.260, n 1.437. Partially hydrolysed to remove any tertiary alkyl halide, then fractionally distd, washed with conc H₂SO₄, water and aqueous K₂CO₃, then redistd from dry K₂CO₃. [Dunbar and Hammett *J Am Chem Soc* 72 109 1950.]

Isobutyl chloride (1-chloro-2-methylpropane) [513-36-0] M 92.3, b 68.8°/760mm, d 0.877, n 1.398. Same methods as described under *isoamyl chloride*.

Isobutyl chloroformate [543-27-1] M 136.6, b 123-127°/atm, 128.8°/atm, d 1.053, n 1.4070. It can be dried over CaCl₂ and fractionated at atm press while keeping moisture out. Its purity can be checked by conversion to the *phenyl urethane derivative* with PhNCO [Saunders et al. *J Am Chem Soc* 73 3796 1951.] IR: ν 1780cm⁻¹ [Thompson and Jameson *Spectrochim Acta* 13 236 1959; Röse *Justus Liebigs Ann Chem* 205 227 1880].

Isobutyl formate [542-55-2] M 102.1, b 98.4°, d 0.885, n 1.38546. Washed with saturated aqueous NaHCO₃ in the presence of saturated NaCl, until no further reaction occurred, then with saturated aqueous NaCl, dried (MgSO₄) and fractionally distd.

Isobutyl iodide (1-iodo-2-methylpropane) [513-38-2] M 184.0, b 83°/250mm, 120°/760mm, d 1.60, n 1.495. Shaken with conc H₂SO₄, and washed with water, aqueous Na₂SO₃, and water, dried with MgSO₄ and distd. Alternatively, passed through a column of activated alumina before distn. Stored under nitrogen with mercury in a brown bottle or in the dark.

Isobutyl vinyl ether [109-53-5] M 100.2, b 108-110°, d 0.768, n 1.398. Washed three times with equal volumes of aqueous 1% NaOH, dried with CaH₂, refluxed with sodium for several hours, then fractionally distd from sodium.

Isobutyraldehyde [78-84-2] M 72.1, b 62.0°, d 0.789, n 1.377. Dried with CaSO₄ and used immediately after distn under nitrogen because of the great difficulty in preventing oxidation. Can be purified through its acid bisulfite derivative.

Isobutyramide [563-83-7] M 87.1, m 128-129°, b 217-221°. Crystd from acetone, *benzene, CHCl₃ or water, then dried under vacuum over P₂O₅ or 99% H₂SO₄. Sublimed under vacuum.

Isobutyric acid [79-31-2] M 88.1, b 154-154.5°, d 0.949, n 1.393, pK²⁵ 4.60. Distd from KMnO₄, then redistd from P₂O₅.

Isobutyronitrile (2-methylpropionitrile, isopropyl cyanide) [78-82-0] M 69.1, b 103.6°, d²⁵ 0.7650, n 1.378. Shaken with conc HCl (to remove isonitriles), then with water and aq NaHCO₃. After a preliminary drying with silica gel or Linde type 4A molecular sieves, it is shaken or stirred with CaH₂ until hydrogen evolution ceases, then decanted and distd from P₂O₅ (not more than 5g/L, to minimize gel formation). Finally it is refluxed with, and slowly distd from CaH₂ (5g/L), taking precautions to exclude moisture.

(-)-γ-Isocaryophyllene (8-methylene-4,11,11-trimethylbicyclo[7.2.0]undec-4-ene) [118-65-0] M 204.4, b 122-124°/12mm, 131-133°/16mm, 130-131°/24mm, 271-273°/atm, d₄²⁰ 0.8959, n_D²⁰ 1.496, [α]₅₄₆²⁰ -31°, [α]_D²⁰ -27° (neat). Purified by vac dist or GLC using a nitrile-silicone column [Corey et al. *J Am Chem Soc* 86 485 1964; Ramage and Simonsen *J Chem Soc* 741 1936; Kumar et al. *Synthesis* 461 1976].

L-Isoleucine [73-32-5] M 131.2, m 285-286°(dec), [α]_D²⁰ +40.6° (6M HCl) pK₁²⁵ 2.66, pK₂²⁵ 9.69. Crystd from water by addition of 4 volumes of EtOH.

(-)-β-Isolongifolene (1-R(-)- 2,2,7,7-tetramethyltricyclo[6.2.1.0^{1,6}]undec-5-ene) [1135-66-6] M 204.4, b 82-83°/0.4mm, 144-146°/30mm, 255-256°/atm, d₄²⁰ 0.930, n_D²⁰ 1.4992, [α]₅₄₆²⁰ -166°, [α]_D²⁰ -138° (c 1, EtOH). Refluxed over and distd from Na. [Zeiss and Arakawa *J Am Chem Soc* 76 1653 1954; IR: Reinaecker and Graafe *Angew Chem, Int Ed Engl* 97 348 1985; UV and NMR: Ranganathan et al. *Tetrahedron* 26 621 1970.]

Isolysergic acid [478-95-5] M 268.3, m 218°(dec), [α]_D²⁰ +281° (c 1, pyridine) pK₁²⁴ 3.33, pK₂²⁴ 8.46. Crystd from water.

Isonicotinamide [1453-82-3] M 122.1, m 155.5-156°, pK₁²⁰ -1.0 (protonation of CONH₂), pK₂²⁰ 3.61, pK₃²⁵ 11.47 (acidic CONH₂). Recrystd from hot water.

Isonicotinic acid (pyridine-4-carboxylic acid) [55-22-1] M 123.1, m 320°, pK²⁵ 4.90. Crystd repeatedly from water. Dried under vac at 110°.

Isonicotinic acid hydrazide (isoniazide) [54-85-3] M 137.1, m 172°, pK₁ 1.75 (NHNH₂), pK₂ 3.57 (=N-), pK₃ 10.75 (-NH). Crystd from 95% EtOH.

1-Isonicotinyl-2-isopropylhydrazide [54-92-2] M 179.2, m 112.5-113.5°. Crystd from *benzene/pet ether.

1-Isonicotinyl-2-salicylidenehydrazide [495-84-1] M 241.2, m 232-233°. Crystd from EtOH.

Isonitrosoacetone (anti-pyruvic aldehyde-1-oxime) [31915-82-9] M 87.1, m 69°. Crystd from ether/pet ether or CCl₄.

Isonitrosoacetophenone (phenylglyoxaldoxime) [532-54-7] M 149.2, m 126-128°. Crystd from water.

5-Isonitrosobarbituric acid (violuric acid) [26851-19-9] M 175.1, m 221-223°, 245-250°, pK₁ 4.41, pK₂ 9.66 (10.1). Crystd from water or EtOH. *1,1-Dimethylvioluric acid*, m 144-147° has pK 4.72 [Taylor and Robinson *Talanta* 8 518 1961].

Isononane [34464-40-9] M 128.3, b 142°/760mm. Passed through columns of activated silica gel and basic alumina (activity 1). Distd under high vacuum from Na/K alloy.

Isopentyl formate [110-45-2] M 116.2, b 121-123°/atm, 123-123.6°/atm, 123-124°/atm, d₄²⁰ 0.8713, n_D²⁰ 1.391. Colourless liquid which is soluble in 300 volumes of H₂O and is soluble in common organic solvents. It is purified by repeated distn using an efficient column at atmospheric pressure.

Isophorone [78-59-1] M 138.2, b 94°/16mm, d 0.921, n¹⁸ 1.4778. Washed with aqueous 5% Na₂CO₃ and then distd under reduced pressure, immediately before use. Alternatively, can be purified *via* the semicarbazone. [Erskine and Waight *J Chem Soc* 3425 1960.]

Isophthalic acid (benzene-1,3-dicarboxylic acid) [121-91-5] M 166.1, m 345-348°, pK₁^{2.5} 3.70, pK₂^{2.5} 4.60. Crystd from aqueous EtOH.

Isopinocampheol (pinan-3-ol, 2,6,6-trimethylbicyclo[3.1.1]heptan-3-ol) [1S,2S,3S,5R-(+)-27779-29-9; 1R,2R,3R,5S-(-)-25465-65-0] M 154.25, m 52-55°, 55-56°, 55-57°, b 103°/11mm, n_D²⁰ 1.4832, [α]₅₄₆²⁰ (+) and (-) 43°, [α]_D²⁰ (+) and (-) 36° (c 20, EtOH). Dissolve in Et₂O, dry MgSO₄, filter, evaporate, then recryst from pet ether. Also recryst from aqueous EtOH and has been distd in a vacuum. [Kergomard and Geneix *Bull Soc Chim Fr* 394 1958; Zweifel and Brown *J Am Chem Soc* 86 393 1964.] The 3,4-dinitrobenzoyl deriv has m 100-101°, the phenylcarbamoyl derivative has m 137-138° and the acid -phthalate has m 125-126°.

Isoprene (2-methyl-1,3-butadiene) [78-79-5] M 68.1, b 34.5-35°/762mm, d 0.681, n²⁵ 1.4225. Refluxed with sodium. Distd from sodium or NaBH₄ under nitrogen, then passed through a column containing KOH, CaSO₄ and silica gel. *tert*-Butylcatechol (0.02% w/w) was added, and the isoprene was stored in this way until redistd before use. The inhibitor (*tert*-butylcatechol) in isoprene can be removed by several washings with dil NaOH and water. The isoprene is then dried over CaH₂, distd under nitrogen at atmospheric pressure, and the fraction distilling at 32° is collected. Stored under nitrogen at -15°.

Isopropanol [67-63-0] M 60.1, b 82.5°, d 0.783, n^{25.8} 1.3739, pK^{2.5} 17.1. Isopropyl alcohol is prepared commercially by dissolution of propene in H₂SO₄, followed by hydrolysis of the sulfate ester. Major impurities are water, lower alcohols and oxidation products such as aldehydes and ketones. Purification of isopropanol follows substantially the same procedure as for *n*-propyl alcohol.

Isopropanol forms a constant-boiling mixture, b 80.3°, with water. Most of the water can be removed from this 91% isopropanol by refluxing with CaO (200g/L) for several hours, then distilling. The distillate can be dried further with CaH₂, magnesium ribbon, BaO, CaSO₄, calcium, anhydrous CuSO₄ or Linde type 5A molecular sieves. Distn from sulfanilic acid removes ammonia and other basic impurities. Peroxides [indicated by liberation of iodine from weakly acid (HCl) solns of 2% KI] can be removed by refluxing with solid stannous chloride or with NaBH₄ then fractionally distilling. To obtain isopropanol containing only 0.002M of water, sodium (8g/L) has been dissolved in material dried by distn from CaSO₄, 35mL of isopropyl benzoate has been

added and, after refluxing for 3h, the alcohol has been distd through a 50-cm Vigreux column. [Hine and Tanabe *J Am Chem Soc* **80** 3002 1958.] Other purification steps for isopropanol include refluxing with solid aluminium isopropoxide, refluxing with NaBH₄ for 24h, and the removal of acetone by treatment with, and distn from 2,4-dinitrophenylhydrazine. Peroxides re-form in isopropanol if it is stood for several days.

Isopropenylcyclobutane [3019-22-5] **M 98.1, b 98.7°, d 0.7743, n 1.438.** Purified by preparative chromatography (silicon oil column), or fractionally distd. Dried with molecular sieves.

Isopropyl acetate [108-22-5] **M 102.1, b 88.4°, d 0.873, n 1.3773.** Washed with 50% aq K₂CO₃ (to remove acid), then with saturated aq CaCl₂ (to remove any alcohol). Dried with CaCl₂ and fractionally distd.

Isopropyl bromide (2-bromopropane) [75-26-3] **M 123.0, b 0°/69.2mm, 59.4°/760mm, d 1.31, n¹⁵ 1.42847, n 1.4251.** Washed with 95% H₂SO₄ (conc acid partially oxidised it) until a fresh portion of acid did not become coloured after several hours, then with water, aq NaHSO₃, aq 10% Na₂CO₃ and again with water. (The H₂SO₄ can be replaced by conc HCl.) Prior to this treatment, isopropyl bromide has been purified by bubbling a stream of oxygen containing 5% ozone through it for 1h, followed by shaking with 3% hydrogen peroxide soln, neutralising with aq Na₂CO₃, washing with distilled water and drying. Alternatively, it has been treated with elemental bromine and stored for 4 weeks, then extracted with aq NaHSO₃ and dried with MgSO₄. After the acid treatment, isopropyl bromide can be dried with Na₂SO₄, MgSO₄ or CaH₂, and fractionally distd.

N-Isopropylcarbazole [1484-09-9] **M 209.3, m 120°.** Crystd from isopropanol. Sublimed under vacuum. Zone refined. The *picrate* has **m 143°** after recrystn from EtOH.

Isopropyl chloride (2-chloropropane) [75-29-6] **M 78.5, b 34.8°, d 0.864, n 1.3779, n²⁵ 1.3754.** Purified with 95% H₂SO₄ as described for *isopropyl bromide*, then dried with MgSO₄, P₂O₅ or CaH₂, and fractionally distd from Na₂CO₃ or CaH₂. Alternatively, a stream of oxygen containing *ca* 5% ozone has been passed through the chloride for about three times as long as was necessary to obtain the first coloration of starch iodide paper by the exit gas, and the liquid was then washed with NaHCO₃ soln to hydrolyse ozonides and remove organic acids before drying and distilling.

Isopropyl ether (diisopropyl ether) [108-20-3] **M 102.2, b 68.3°, d 0.719, n 1.3688, n²⁵ 1.36618.** Common impurities are water and peroxides [detected by the liberation of iodine from weakly acid (HCl) solns of 2% KI]. Peroxides can be removed by shaking with aqueous Na₂SO₃ or with acidified ferrous sulfate (0.6g FeSO₄ and 6mL conc H₂SO₄ in 110mL of water, using 5-10g of soln per L of ether), or aqueous NaBH₄ soln. The ether is then washed with water, dried with CaCl₂ and distd. Alternatively, refluxing with LiAlH₄ or CaH₂, or drying with CaSO₄, then passage through an activated alumina column, can be used to remove water and peroxides. Other dehydrating agents used with isopropyl ether include P₂O₅, sodium amalgam and sodium wire. (The ether is often stored in brown bottles, or in the dark, with sodium wire.) Bonner and Goishi (*J Am Chem Soc* **83** 85 1961) treated isopropyl ether with dil sodium dichromate/sulfuric acid soln, followed by repeated shaking with a 1:1 mixture of 6M NaOH and saturated KMnO₄. The ether was washed several times with water, dilute aqueous HCl, and water, with a final washing with, and storage over, ferrous ammonium sulfate acidified with H₂SO₄. Blaustein and Gryder (*J Am Chem Soc* **79** 540 1957), after washing with alkaline KMnO₄, then water, treated the ether with ceric nitrate in nitric acid, and again washed with water. Hydroquinone was added before drying with CaCl₂ and MgSO₄, and refluxing with sodium amalgam (108g Hg/100g Na) for 2h under nitrogen. The distillate (nitrogen atmosphere) was made 2 x 10⁻⁵M in hydroquinone to inhibit peroxide formation (which was negligible if the ether was stored in the dark). Catechol (pyrocatechol) and resorcinol are alternative inhibitors.

4,4'-Isopropylidenediphenol [80-05-7] **M 228.3, m 158°, pK_{Est} ~10.3.** Crystd from acetic acid/water (1:1).

Isopropyl iodide (2-iodopropane) [75-30-9] **M 170.0, b 88.9°, d 1.70, n 1.4987.** Treated with bromine, followed by extraction of free halogen with aqueous Na₂S₂O₃ or NaHSO₃, washing with water, drying (MgSO₄ or CaCl₂) and distn. (The treatment with bromine is optional.) Other purification methods include

passage through activated alumina, or shaking with copper powder or mercury to remove iodine, drying with P_2O_5 and distn. Washing with conc H_2SO_4 or conc HCl (to remove any alcohol), water, aqueous Na_2SO_3 , water and aqueous Na_2CO_3 has also been used. Treatment with silica gel causes some liberation of iodine. Distillations should be carried out at slightly reduced pressure. Purified isopropyl iodide is stored in the dark in the presence of a little mercury.

Isopropyl methyl ether [598-53-8] M 74.1, b 32.5°/777mm, d^{15} 0.724, n 1.3576. Purified by drying with $CaSO_4$, passage through a column of alumina (to remove peroxides) and fractional distn.

Isopropyl *p*-nitrobenzoate [13756-40-6] M 209.2, m 105-106°. Dissolved in diethyl ether, washed with aqueous alkali, then water and dried. Evapn of the ether and recrystn from EtOH gave pure material.

***p*-Isopropyl toluene (*p*-cymene)** [99-87-6] M 134.2, b 176.9°/744mm, d 0.8569, n 1.4902. See entry on p. 183.

Isoquinoline [119-65-3] M 129.2, m 24°, b 120°/18mm, d 1.0986, n 1.6148, pK^{25} 5.40. Dried with Linde type 5A molecular sieves or Na_2SO_4 and fractionally distd at reduced pressure. Alternatively, it was refluxed with, and distd from, BaO. Also purified by fractional crystn from the melt and distd from zinc dust. Converted to its *phosphate* (m 135°) or *picrate* (m 223°), which were purified by crystn and the free base recovered and distd. [Packer, Vaughn and Wong *J Am Chem Soc* 80 905 1958.] The procedure for purifying *via* the picrate comprises the addition of quinoline to picric acid dissolved in the minimum volume of 95% EtOH to yield yellow crystals which are washed with EtOH and air dried before recrystn from acetonitrile. The crystals are dissolved in dimethyl sulfoxide (previously dried over 4A molecular sieves) and passed through a basic alumina column, on which picric acid is adsorbed. The free base in the effluent is extracted with *n*-pentane and distd under vacuum. Traces of solvent are removed by vapour phase chromatography. [Mooman and Anton *J Phys Chem* 80 2243 1976.]

Isovaleric acid [502-74-2] M 102.1, b 176.5°/762mm, d 0.927, n^{15} 1.4064, n 1.40331, pK^{25} 4.77. Dried with Na_2SO_4 , then fractionally distd.

L-Isovaline (2-amino-2-methylbutyric acid) [595-40-4] M 117.2, m *ca* 300° (sublimes in vac), $[\alpha]_D^{25} +10^\circ$ (5M HCl), $pK_{Est(1)} \sim 2.4$, $pK_{Est(2)} \sim 9.7$. Crystd from aqueous acetone.

Isovanillin (3-hydroxy-4-methoxybenzaldehyde) [621-59-0] M 152.2, m 117°, b 175°/14mm, pK^{25} 8.89. Cryst from H_2O or $*C_6H_6$. The *oxime* has m 147°.

Isoviolanthrone [128-64-3] M 456.5, m 510-511°(uncorrected). Dissolved in 98% H_2SO_4 and ppted by adding water to reduce the acid concentration to about 90%. Sublimes *in vacuo*. [Parkyns and Ubbelhode *J Chem Soc* 4188 1960.]

Itaconic acid (2-propen-1,2-dicarboxylic acid) [97-65-4] M 130.1, m 165-166°, pK_1^{25} 3.63, pK_2^{25} 5.00. Crystd from EtOH, EtOH/water or EtOH/*benzene.

Itaconic anhydride (2-propen-1,2-dicarboxylic anhydride) [2170-03-8] M 112.1, m 66-68°, 67-68°, 68°, b 139-140°/30mm. Crystd from $CHCl_3$ /pet ether. Can be distd under reduced press. Distn at atm press, or prolonged distn causes rearrangement to citraconic anhydride (2-methylmaleic anhydride). If the material (as seen in the IR spectrum) contains much free acid then heat with acetyl chloride or $SOCl_2$, evaporate and distil at as high a vacuum as possible. The crude anhydride deposits crystals of itaconic acid on standing probably due to hydrolysis by H_2O — store in sealed ampoules under dry N_2 . [*Org Synth Coll Vol II* 369 1943; IR: Nagai *Bull Chem Soc Jpn* 37 369 1964; Kelly and Segura *J Am Chem Soc* 56 2497 1934.]

Janus Green B (3-dimethylamino-7-[4-dimethylaminoazo]-5-phenylphenazonium chloride) [2869-83-2] M 511.1, m >200°. Dissolves in H_2O to give a bluish violet soln which

[Previous Page](#)

passage through activated alumina, or shaking with copper powder or mercury to remove iodine, drying with P_2O_5 and distn. Washing with conc H_2SO_4 or conc HCl (to remove any alcohol), water, aqueous Na_2SO_3 , water and aqueous Na_2CO_3 has also been used. Treatment with silica gel causes some liberation of iodine. Distillations should be carried out at slightly reduced pressure. Purified isopropyl iodide is stored in the dark in the presence of a little mercury.

Isopropyl methyl ether [598-53-8] M 74.1, b 32.5°/777mm, d^{15} 0.724, n 1.3576. Purified by drying with $CaSO_4$, passage through a column of alumina (to remove peroxides) and fractional distn.

Isopropyl *p*-nitrobenzoate [13756-40-6] M 209.2, m 105-106°. Dissolved in diethyl ether, washed with aqueous alkali, then water and dried. Evapn of the ether and recrystn from EtOH gave pure material.

***p*-Isopropyl toluene (*p*-cymene)** [99-87-6] M 134.2, b 176.9°/744mm, d 0.8569, n 1.4902. See entry on p. 183.

Isoquinoline [119-65-3] M 129.2, m 24°, b 120°/18mm, d 1.0986, n 1.6148, pK^{25} 5.40. Dried with Linde type 5A molecular sieves or Na_2SO_4 and fractionally distd at reduced pressure. Alternatively, it was refluxed with, and distd from, BaO. Also purified by fractional crystn from the melt and distd from zinc dust. Converted to its *phosphate* (m 135°) or *picrate* (m 223°), which were purified by crystn and the free base recovered and distd. [Packer, Vaughn and Wong *J Am Chem Soc* 80 905 1958.] The procedure for purifying *via* the picrate comprises the addition of quinoline to picric acid dissolved in the minimum volume of 95% EtOH to yield yellow crystals which are washed with EtOH and air dried before recrystn from acetonitrile. The crystals are dissolved in dimethyl sulfoxide (previously dried over 4A molecular sieves) and passed through a basic alumina column, on which picric acid is adsorbed. The free base in the effluent is extracted with *n*-pentane and distd under vacuum. Traces of solvent are removed by vapour phase chromatography. [Mooman and Anton *J Phys Chem* 80 2243 1976.]

Isovaleric acid [502-74-2] M 102.1, b 176.5°/762mm, d 0.927, n^{15} 1.4064, n 1.40331, pK^{25} 4.77. Dried with Na_2SO_4 , then fractionally distd.

L-Isovaline (2-amino-2-methylbutyric acid) [595-40-4] M 117.2, m *ca* 300° (sublimes in vac), $[\alpha]_D^{25} +10^\circ$ (5M HCl), $pK_{Est(1)} \sim 2.4$, $pK_{Est(2)} \sim 9.7$. Crystd from aqueous acetone.

Isovanillin (3-hydroxy-4-methoxybenzaldehyde) [621-59-0] M 152.2, m 117°, b 175°/14mm, pK^{25} 8.89. Cryst from H_2O or $*C_6H_6$. The *oxime* has m 147°.

Isoviolanthrone [128-64-3] M 456.5, m 510-511°(uncorrected). Dissolved in 98% H_2SO_4 and ppted by adding water to reduce the acid concentration to about 90%. Sublimes *in vacuo*. [Parkyns and Ubbelhode *J Chem Soc* 4188 1960.]

Itaconic acid (2-propen-1,2-dicarboxylic acid) [97-65-4] M 130.1, m 165-166°, pK_1^{25} 3.63, pK_2^{25} 5.00. Crystd from EtOH, EtOH/water or EtOH/*benzene.

Itaconic anhydride (2-propen-1,2-dicarboxylic anhydride) [2170-03-8] M 112.1, m 66-68°, 67-68°, 68°, b 139-140°/30mm. Crystd from $CHCl_3$ /pet ether. Can be distd under reduced press. Distn at atm press, or prolonged distn causes rearrangement to citraconic anhydride (2-methylmaleic anhydride). If the material (as seen in the IR spectrum) contains much free acid then heat with acetyl chloride or $SOCl_2$, evaporate and distil at as high a vacuum as possible. The crude anhydride deposits crystals of itaconic acid on standing probably due to hydrolysis by H_2O — store in sealed ampoules under dry N_2 . [*Org Synth Coll Vol II* 369 1943; IR: Nagai *Bull Chem Soc Jpn* 37 369 1964; Kelly and Segura *J Am Chem Soc* 56 2497 1934.]

Janus Green B (3-dimethylamino-7-[4-dimethylaminoazo]-5-phenylphenazonium chloride) [2869-83-2] M 511.1, m >200°. Dissolves in H_2O to give a bluish violet soln which

becomes colourless when made 10M in NaOH. Dissolve in EtOH to give a blue-violet colour, filter from insoluble material then add dry Et₂O whereby the dye separates out leaving a small amount of blue colour in soln. Filter off the solid and dry in vacuum. Store in a dark bottle.

Janus Red B {3-[(2-hydroxy-1-naphthol)azo-2-methylphenylazo]N,N,N-trimethylbenzenaminium chloride} [2636-31-9] M 460.0. Crystd from EtOH/H₂O (1:1 v/v) and dry in vacuum. Store in a dark bottle.

Jervine (3β,23β-17,23-epoxy-2-hydroxyvertraman-11-one, a steroidal alkaloid) [469-59-0] M 425.6, m 243-245°, [α]_D²⁰ -150° (in EtOH), pK_{Est}~9.4 Crystd from MeOH/H₂O. The hydrochloride has m 300-302°. [Kutney et al. *Can J Chem* 53 1796 1975.]

Julolidine (2,3,6,7-tetrahydro-1H,5H-benzo[*ij*]quinolizidine) [479-59-4] M 173.3, m 34-36°, 40°, b 105-110°/1mm, 155-156°/17mm, 280° (dec), pK_{Est}~7.0. Purified by dissolving in dilute HCl, steam is bubbled through the soln and the residual acidic soln is basified with 10N NaOH, extracted with Et₂O, washed with H₂O, dried (NaOH pellets), filtered, evaporated and distd *in vacuo*. The distillate crystallises on standing (m 39-40°). On standing in contact with air for several days it develops a red colour. The colour can be removed by distilling or dissolving in 2-3 parts of hexane, adding charcoal, filtering and cooling in Me₂CO-Dry-ice when julolidine crystallises (85-90% yield). The hydrochloride [83646-41-7] has m 218° (239-242°), the picrate has m 165° and the methiodide crystallises from MeOH, m 186° [Glass and Weisberger *Org Synth Coll Vol III* 304 1955.] **Highly TOXIC.**

Kainic acid monohydrate (2S,3S,4S-2-carboxy-4-isoprenyl-3-pyrrolidine-acetic acid) [487-79-6] M 231.4, m 235-245° (dec), 251° (dec), [α]_D²⁰ -14.6° (c 1.46, H₂O), pK₁ 2.09, pK₂ 4.58, pK₃ 10.21. Purified by adsorbing on to a strongly acidic ion exchange resin (Merck), elution of the diacid with aqueous M NaOH, the eluate is evaporated, H₂O is added, and filtered through a weakly acidic ion exchange resin (Merck). The filtrate is then evaporated and recrystd from EtOH. Its solubility is 0.1g in 1mL of 0.5N HCl. (±)-α-Kainic acid recryst from H₂O, m 230-260°. UV (MeOH): λ_{max} 219 (log ε 3.9); ¹H NMR (CCl₄, 100MHz, Me₄Si standard) δ: 1.64 (s 1H), 1.70 (s 3H), 3.24 (d J 7.5, 2H), 3.3-4.2 (1H), 3.70 (s 3H), 3.83 (s 3H), 4.35 (dd J 7.5, J 14.5, 1H), 5.21 (t J 7.5, 1H), 7.26 (t J 7.5, 1H). [Oppolzer and Andres *Helv Chim Acta* 62 2282 1979.]

Kerosene [8008-20-6] (mixture of hydrocarbons) b ~175-325°, d 0.75-0.82, n 1.443. Stirred with conc H₂SO₄ until a fresh portion of acid remains colourless, then washed with water, dried with solid KOH and distd in a Claisen flask. For more complete drying, the kerosene can be refluxed with, and distd from Na.

Ketanserine [3(4-*p*-fluorobenzoylpiperidinyl-*N*-ethyl)quinazolin-2,4-dione] [74050-98-9] M 395.4, m 227-235°, pK 7.5. Solubility is 0.001% in H₂O, 0.038% in EtOH and 2.34 in Me₂NCHO. It has been purified by recrystn from 4-methyl-3-pentanone [Peeters et al. *Cryst Structure Commun* 11 375 1982; Kacprowicz et al. *J Chromatogr* 272 417 1983; Davies et al. *J Chromatogr* 275 232 1983].

Ketene [463-51-4] M 42.0, b 127-130°, d 1.093, n 1.441. Prepared by pyrolysis of acetic anhydride. Purified by passage through a trap at -75° and collected in a liquid-nitrogen-cooled trap. Ethylene was removed by evacuating the ethylene in an isopentane-liquid-nitrogen slush pack at -160°. Stored at room temperature in a suitable container in the dark. See diketene on p. 209.

2-Keto-L-gulonic acid [526-98-7] M 194.1, m 171°. Crystd from water and washed with acetone.

Ketone moschus (4-*tert*-butyl-2,6-dimethyl-3,5-dinitroacetophenone) [81-14-1] M 234.1, m 134-137°, 137-138°. Purified by recryst from MeOH. [Fuson et al. *J Org Chem* 12 587 1947.]

Khellin (4,9-dimethoxy-7-methyl-5-oxofuro[3,2-*g*]-1,2-chromene) [82-02-0] M 260.3, m 154-155°, b 180-200°/0.05mm. Crystd from MeOH or diethyl ether.

Kojic acid [(2-hydroxy-5-hydroxymethyl)-4H-pyran-4-one] [501-30-4] M 142.1, m 154-155°, pK_1^{25} -1.38, pK_2^{25} 7.66. Crystd from MeOH (charcoal) by adding Et₂O. Sublimed at 0.1 torr.

Kynurenic acid (4-hydroxyquinoline-2-carboxylic acid) [492-27-3] M 189.1, m 282-283°, $pK_{Est(1)} \sim 2$, $pK_{Est(2)} \sim 10$. Crystd from absolute EtOH.

L-Kynurenine [343-65-7] M 208.2, m 190°(dec), 210°(dec), $[\alpha]_D^{20}$ -30° (c 0.4, H₂O), $pK_{Est(1)} \sim 2.3$, $pK_{Est(2)} \sim 3.5$, $pK_{Est(3)} \sim 9.2$. Crystd from H₂O or aq AcOH. *Picrate* has m 188.5-189°(dec) after crystn from H₂O.

L-Kynurenine sulfate [16055-80-4] M 306.3, m 194°, monohydrate m 178°, $[\alpha]_D^{25}$ +9.6° (H₂O). Crystd from water by addition of EtOH.

L(+)-Lactic acid [79-33-4] M 90.1, m 52.8°, b 105°/0.1mm, $[\alpha]_D^{20}$ +3.82° (H₂O), pK^{31} 3.83. Purified by fractional distn at 0.1mm pressure, followed by fractional crystn from diethyl ether/isopropyl ether (1:1, dried with sodium). [Borsook, Huffman and Liu *J Biol Chem* 102 449 1933.] The solvent mixture, *benzene/diethyl ether (1:1) containing 5% pet ether (b 60-80°) has also been used.

Lactobionic acid [96-82-2] M 358.3, m 128-130°, $[\alpha]_{546}^{20}$ +28° (c 3, after 24h in H₂O), $pK_{Est} \sim 3.6$. Crystd from water by addition of EtOH.

α -Lactose (H₂O) [63-42-3] M 360.3, m 220°(dec), $[\alpha]_D^{20}$ +52.3° (c 4.2, H₂O), pK 12.2 (OH). Crystd from water below 93.5°.

Lactulose [4618-18-2] M 342.2, m 167-169°(dec), $[\alpha]_{546}^{20}$ -57° (c 1, H₂O). Crystd from MeOH.

Lanatoside A [17575-20-1] M 969.1, m 245-248°, $[\alpha]_D^{20}$ +32° (EtOH). Crystd from MeOH.

Lanatoside B [17575-21-2] M 985.1, m 233°(dec), $[\alpha]_D^{20}$ +35° (MeOH). Crystd from MeOH.

Lanatoside C [17575-22-3] M 297.1, m 246-248°, $[\alpha]_D^{20}$ +34° (EtOH). Crystd from MeOH.

Lanosterol [79-63-0] M 426.7, m 138-140°, $[\alpha]_D^{20}$ +62.0° (c 1, CHCl₃). Recrystd from anhydrous MeOH. Dried *in vacuo* over P₂O₅ for 3h at 90°. Purity checked by proton magnetic resonance.

Lanthanide shift reagents A variety of these reagents are available commercially and they are generally quite stable and should not deteriorate on long storage in a dry state and in the absence of light. [See G.R.Sullivan in *Top Stereochem* (Eliel and Allinger Eds) J Wiley & Sons Vol 10 287 1978; T.C.Morrill Ed. *Lanthanide Shift Reagents* Deerfield Beach Florida 1986, ISBN 0895731193.]

Lapachol [84-79-7] M 226.3, m 140°. Crystd from EtOH or diethyl ether.

dl- and l-Laudanosine [(±) 1699-51-0; (-) 2688-77-9] M 357.4, m 114-115°. Crystd from EtOH. The (-)-isomer has m 83-85° and $[\alpha]_D^{20}$ -85° (c 0.5, EtOH).

Lauraldehyde (1-dodecanal) [112-54-9] M 184.3, b 99.5-100°/3.5mm, $n^{24.7}$ 1.4328. Converted to the addition compound by shaking with saturated aqueous NaHSO₃ for 1h. The ppte was filtered off, washed with ice cold water, EtOH and ether, then decomposed with aqueous Na₂CO₃. The aldehyde was extracted into diethyl ether which, after drying and evap, gave an oil which was fractionally distd under vacuum.

Lauric acid (1-dodecanoic acid) [143-07-7] M 200.3, m 44.1°, b 141-142°/0.6-0.7mm, 225°/100mm, pK^{20} 5.3. Vacuum distd. Crystd from absolute EtOH, or from acetone at -25°.

Alternatively, purified *via* its *methyl ester* (b 140.0°/15mm), as described for *capric acid*. Also purified by zone melting.

Lauryl peroxide (dodecyl peroxide) [105-74-8] M 398.6, m 53-54°. Crystd from *n*-hexane or *benzene and stored below 0°. Potentially **EXPLOSIVE**.

L-Leucine [61-90-5] M 131.2, m 293-295°(dec), $[\alpha]_D^{25} +15.6^\circ$ (5M HCl), $pK_1^{25} 2.33$, $pK_2^{25} 9.74$. Likely impurities are isoleucine, valine, and methionine. Crystd from water by adding 4 volumes of EtOH.

Leucomalachite Green [129-73-7] M 330.5, m 92-93°, $pK^{25} 6.90$ (several pK 's). Crystd from 95% EtOH (10mL/g), then from *benzene/EtOH, and finally from pet ether.

Lithocholic acid [434-13-9] M 376.6, m 184-186°, $[\alpha]_D^{23} +35^\circ$ (c 1, EtOH), $pK_{Est} \sim 4.8$. Crystd from EtOH or acetic acid.

Lumichrome [1086-80-2] M 242.2, m >290°, $pK_{Est(1)} \sim -0.1$ (basic), $pK_{Est(2)} \sim 9.9$ (acidic), Recrystd twice from glacial AcOH and dried at 100° in a vacuum.

Luminol (5-aminophthalazin-1,4-dione) [521-31-3] M 177.2, m 329-332°, $pK_1 3.37$, $pK_2 6.35$. Dissolved in KOH soln, treated with Norit (charcoal), filtered and ppted with conc HCl. [Hardy, Sietz and Hercules *Talanta* 24 297 1977.] Stored in the dark in an inert atmosphere, because its structure changes during its luminescence. It has been recrystd from 0.1M KOH [Merenyi et al. *J Am Chem Soc* 108 77716 1986].

dl-Lupinane [10248-30-3] M 169.3, m 98-99°. Crystd from acetone.

Lupulon [468-28-0] M 414.6, m 92-94°. Crystd from 90% MeOH.

Lutein (α -carotene-3,3'-diol, xanthophyll) [127-40-2] M 568.9, m 196°, $\epsilon_{1cm}^{1\%} 1750$ (423nm), 2560 (446nm), 2340 (477.5nm) in EtOH; λ_{max} in CS₂ 446, 479 and 511nm. Crystd from MeOH (copper-coloured prisms) or from diethyl ether by adding MeOH. Also purified by chromatography on columns of magnesia or calcium hydroxide, and crystd from CS₂/EtOH. May be purified *via* the dipalmitate ester. Stored in the dark, in an inert atmosphere.

Lutidine (mixture). For the preparation of pure 2,3-, 2,4- and 2,5-lutidine from commercial "2,4- and 2,5-lutidine" see Coulson et al. *J Chem Soc* 1934 1959, and Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960.

2,3-Lutidine [583-61-9] M 107.2, f -14.8°, b 160.6°, d 0.9464, n 1.50857, $pK^{25} 6.57$. Steam distd from a soln containing about 1.2 equivalents of 20% H₂SO₄, until *ca* 10% of the base has been carried over with the non-basic impurities. The acid soln was then made alkaline, and the base was separated, dried over NaOH or BaO, and fractionally distd. The distd lutidine was converted to its urea complex by stirring 100g with 40g of urea in 75mL of H₂O, cooling to 5°, filtering at the pump, and washing with 75mL of H₂O. The complex, dissolved in 300mL of H₂O was steam distd until the distillate gave no turbidity with a little solid NaOH. The distillate was then treated with excess solid NaOH, and the upper layer was removed: the aqueous layer was then extracted with diethyl ether. The upper layer and the ether extract were combined, dried (K₂CO₃), and distd through a short column. Final purification was by fractional crystn using partial freezing. [Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960].

2,4-Lutidine [108-47-4] M 107.2, b 157.8°, d 0.9305, n 1.50087, $n^{25} 1.4985$, $pK^{25} 6.77$. Dried with Linde type 5A molecular sieves, BaO or sodium, and fractionally distd. The distillate (200g) was heated with *benzene (500mL) and conc HCl (150mL) in a Dean and Stark apparatus on a water bath until water no longer separated, and the temperature just below the liquid reached 80°. When cold, the supernatant *benzene was decanted and the 2,4-lutidine hydrochloride, after washing with a little *benzene, was dissolved in water (350mL). After removing any *benzene by steam distn, an aqueous soln of NaOH (80g) was added, and the free

lutidine was steam distd. It was isolated by saturating the distillate with solid NaOH, and distd through a short column. The pptn cycle was repeated, then the final distillate was partly frozen in an apparatus at -67.8 - 68.5° (cooled by acetone/ CO_2). The crystals were then melted and distd. [Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960.] Alternative purifications are *via* the picrate [Clarke and Rothwell *J Chem Soc* 1885 1960], or the hydrobromide [Warnhoff *J Org Chem* 27 4587 1962]. The latter is ppted from a soln of lutidine in *benzene by passing dry HBr gas: the salt is recrystd from CHCl_3 /methyl ethyl ketone, then decomposed with NaOH, and the free base is extracted into diethyl ether, dried, evaporated and the residue distd.

2,5-Lutidine [589-93-5] M 107.2, m -15.3° , b $156.7^{\circ}/759\text{mm}$, d 0.927, n^{25} 1.4982, pK^{25} 6.40. Steam distd from a soln containing 1-2 equivalents of 20% H_2SO_4 until about 10% of the base had been carried over with the non-basic impurities, then the acid soln was made alkaline, and the base separated, dried with NaOH and fractionally distd twice. Dried with Na and fractionally distd through a Todd column packed with glass helices (see p. 174).

2,6-Lutidine [108-48-5] M 107.2, m -59° , b 144.0° , d 0.92257, n 1.49779, pK^{25} 6.72. Likely contaminants include 3- and 4-picoline (similar boiling points). However, they are removed by using BF_3 , with which they react preferentially, by adding 4mL of BF_3 to 100mL of dry fractionally distd 2,6-lutidine and redistilling. Distn of commercial material from AlCl_3 (14g per 100mL) can also be used to remove picolines (and water). Alternatively, lutidine (100mL) can be refluxed with ethyl benzenesulfonate (20g) or ethyl *p*-toluenesulfonate (20g) for 1h, then the upper layer is cooled, separated and distd. The distillate is refluxed with BaO or CaH_2 , then fractionally distd, through a glass helices-packed column.

2,6-Lutidine can be dried with KOH or sodium, or by refluxing with (and distilling from) BaO, prior to distn. For purification *via* its picrate, 2,6-lutidine, dissolved in abs EtOH, is treated with an excess of warm ethanolic picric acid. The ppt is filtered off, recrystd from acetone (to give m 163 - 164.5°), and partitioned between ammonia and CHCl_3 /diethyl ether. The organic soln, after washing with dilute aqueous KOH, is dried with Na_2SO_4 and fractionally distd. [Warnhoff *J Org Chem* 27 4587 1962.] Alternatively, 2,6-lutidine can be purified *via* its urea complex, as described under 2,3-lutidine. Other purification procedures include azeotropic distn with phenol [Coulson et al. *J Appl Chem (London)* 2 71 1952], fractional crystn by partial freezing, and vapour-phase chromatography using a 180-cm column of polyethylene glycol-400 (Shell, 5%) on Embacel (May and Baker) at 100° , with argon as carrier gas [Bamford and Block *J Chem Soc* 4989 1961].

3,5-Lutidine [591-22-0] M 107.2, f -6.3° , b $172.0^{\circ}/767\text{mm}$, d 0.9419, n 1.50613, n^{25} 1.5035, pK^{25} 6.15. Dried with sodium and fractionally distd through a Todd column packed with glass helices (see p. 174). Dissolved (100mL) in dil HCl (1:4) and steam distd until 1L of distillate was collected. Excess conc NaOH was added to the residue which was again steam distd. The base was extracted from the distillate, using diethyl ether. The extract was dried with K_2CO_3 , and distd. It was then fractionally crystd by partial freezing.

Lycopene [502-65-8] M 536.9, m 172 - 173° , $\epsilon_{1\text{cm}}^{1\%}$ 2250 (446nm), 3450 (472nm), 3150 (505nm) in pet ether. Crystd from CS_2 /MeOH, diethyl ether/pet ether, or acetone/pet ether, and purified by column chromatography on deactivated alumina, CaCO_3 , calcium hydroxide or magnesia. Stored in the dark, in an inert atmosphere.

Lycorine [476-28-8] M 552.9, m 275 - $280^{\circ}(\text{dec})$ $[\alpha]_{\text{D}}^{20}$ -130° (c 0.16, EtOH). Crystd from EtOH.

Lycoxanthin (Ψ, Ψ -carotene-16-ol) [19891-74-8] M 268.3, m 173 - 174° , $\epsilon_{1\text{cm}}^{1\%}$ 3360 (472.5nm), also λ_{max} 444 and 503nm in pet ether. Crystd from diethyl ether/light petroleum, *benzene/pet ether or CS_2 . Purified by chromatography on columns of CaCO_3 , $\text{Ca}(\text{OH})_2$ or deactivated alumina, washing with *benzene and eluting with 3:1 *benzene/MeOH. Stored in the dark, in an inert atmosphere, at -20° .

Lysergic acid [82-58-6] M 268.3, m $240^{\circ}(\text{dec})$, $[\alpha]_{\text{D}}^{20}$ $+40^{\circ}$ (pyridine), pK_1^{25} 3.32, pK_2^{25} 7.82. Crystd from water.

L-Lysine [56-87-1] M 146.2, m >210°(dec), pK₁ 2.18, pK₂ 8.95, pK₃ 10.53. Crystd from aqueous EtOH.

L-Lysine dihydrochloride [657-26-1] M 219.1, m 193°, [α]_D²⁵ +25.9° (5M HCl). Crystd from MeOH, in the presence of excess HCl, by adding diethyl ether.

L-Lysine monohydrochloride [657-27-2] M 182.7, [α] as above. Likely impurities are arginine, D-lysine, 2,6-diaminoheptanedioic acid and glutamic acid. Crystd from water at pH 4-6 by adding 4 volumes of EtOH. Above 60% relative humidity it forms a dihydrate.

β-D-Lyxose [1114-34-7] M 150.1, m 118-119°, [α]_D²⁰ -14° (c 4, H₂O). Crystd from EtOH or aqueous 80% EtOH. Dried under vacuum at 60°, and stored in a vacuum desiccator over P₂O₅ or CaSO₄.

Malachite Green (carbinol) [510-13-4] M 346.4, m 112-114°, CI 42000, pK²⁴ 6.84. The oxalate was recrystd from hot water and dried in air. The carbinol was pptd from the oxalate (1g) in distd water (100mL) by adding M NaOH (10mL). The ppte was filtered off, recrystd from 95% EtOH containing a little dissolved KOH, then washed with ether, and crystd from pet ether. Dried in a vacuum at 40°. An acid soln (2 x 10⁻⁵M in 6 x 10⁻⁵M H₂SO₄) rapidly reverted to the dye. [Swain and Hedberg *J Am Chem Soc* 72 3373 1950.]

Z-Maleamic acid (cis-maleic acid monoamide) [557-24-4] M 115.1, m 158-161°, 172-173°(dec), pK_{Est} ~2.65. Crystd from EtOH. IRRITANT.

Maleic acid [110-16-7] M 116.1, m 143.5°, pK₁²⁵ 1.91, pK₂²⁵ 6.33. Crystd from acetone/pet ether (b 60-80°) or hot water. Dried at 100°.

Maleic anhydride [108-31-6] M 98.1, m 54°, b 94-96°/20mm, 199°/760mm. Crystd from *benzene, CHCl₃, CH₂Cl₂ or CCl₄. Sublimed under reduced pressure. [Skell et al. *J Am Chem Soc* 108 6300 1986.]

Maleic hydrazide [123-33-1] M 112.1, m 144°(dec), pK₁²⁵ 5.67, pK₂²⁵ 13.3. Crystd from water.

Maleimide (pyrrol-2,5-dione) [541-59-3] M 97.1, m 91-93°, 92.6-93°, d_D^{105.5} 1.2493, n_D^{110.7} 1.49256. Purified by sublimation in a vacuum. The UV has λ_{max} at 216 and 280nm in EtOH. [de Wolf and van de Straete *Bull Soc Chim Belg* 44 288 1935; UV: Rondestvedt et al. *J Am Chem Soc* 78 6115 1956; IR: Chiorboli and Mirone *Ann Chim (Rome)* 42 681 1952.]

Maleuric acid [105-61-3] M 158.1, m 167-168°(dec). Crystd from hot water.

dl-Malic acid [617-48-1 and 6915-15-7] M 134.1, m 128-129°. Crystd from acetone, then from acetone/CCl₄, or from ethyl acetate by adding pet ether (b 60-70°). Dried at 35° under 1mm pressure to avoid formation of the anhydride.

L-Malic acid [97-67-6] M 134.1, m 104.5-106°, [α]_D²⁰ -2.3° (c 8.5, H₂O), pK₁²⁵ 3.46, pK₂²⁵ 5.10. Crystd (charcoal) from ethyl acetate/pet ether (b 55-56°), keeping the temperature below 65°. Or, dissolved by refluxing in fifteen parts of anhydrous diethyl ether, decanted, concentrated to one-third volume and crystd at 0°, repeatedly to constant melting point.

Malonamide [108-13-4] M 102.1, m 170°. Crystd from water.

Malonic acid [141-82-2] M 104.1, m 136°, pK₁²⁵ 2.58, pK₂²⁵ 5.69. Crystd from *benzene/diethyl ether (1:1) containing 5% of pet ether (b 60-80°), washed with diethyl ether, then recrystd from H₂O or acetone. Dried under vac over conc H₂SO₄.

[Previous Page](#)

L-Lysine [56-87-1] M 146.2, m >210°(dec), pK₁ 2.18, pK₂ 8.95, pK₃ 10.53. Crystd from aqueous EtOH.

L-Lysine dihydrochloride [657-26-1] M 219.1, m 193°, [α]_D²⁵ +25.9° (5M HCl). Crystd from MeOH, in the presence of excess HCl, by adding diethyl ether.

L-Lysine monohydrochloride [657-27-2] M 182.7, [α] as above. Likely impurities are arginine, D-lysine, 2,6-diaminoheptanedioic acid and glutamic acid. Crystd from water at pH 4-6 by adding 4 volumes of EtOH. Above 60% relative humidity it forms a dihydrate.

β-D-Lyxose [1114-34-7] M 150.1, m 118-119°, [α]_D²⁰ -14° (c 4, H₂O). Crystd from EtOH or aqueous 80% EtOH. Dried under vacuum at 60°, and stored in a vacuum desiccator over P₂O₅ or CaSO₄.

Malachite Green (carbinol) [510-13-4] M 346.4, m 112-114°, CI 42000, pK²⁴ 6.84. The oxalate was recrystd from hot water and dried in air. The carbinol was pptd from the oxalate (1g) in distd water (100mL) by adding M NaOH (10mL). The ppte was filtered off, recrystd from 95% EtOH containing a little dissolved KOH, then washed with ether, and crystd from pet ether. Dried in a vacuum at 40°. An acid soln (2 x 10⁻⁵M in 6 x 10⁻⁵M H₂SO₄) rapidly reverted to the dye. [Swain and Hedberg *J Am Chem Soc* 72 3373 1950.]

Z-Maleamic acid (cis-maleic acid monoamide) [557-24-4] M 115.1, m 158-161°, 172-173°(dec), pK_{Est} ~2.65. Crystd from EtOH. IRRITANT.

Maleic acid [110-16-7] M 116.1, m 143.5°, pK₁²⁵ 1.91, pK₂²⁵ 6.33. Crystd from acetone/pet ether (b 60-80°) or hot water. Dried at 100°.

Maleic anhydride [108-31-6] M 98.1, m 54°, b 94-96°/20mm, 199°/760mm. Crystd from *benzene, CHCl₃, CH₂Cl₂ or CCl₄. Sublimed under reduced pressure. [Skell et al. *J Am Chem Soc* 108 6300 1986.]

Maleic hydrazide [123-33-1] M 112.1, m 144°(dec), pK₁²⁵ 5.67, pK₂²⁵ 13.3. Crystd from water.

Maleimide (pyrrol-2,5-dione) [541-59-3] M 97.1, m 91-93°, 92.6-93°, d_D^{105.5} 1.2493, n_D^{110.7} 1.49256. Purified by sublimation in a vacuum. The UV has λ_{max} at 216 and 280nm in EtOH. [de Wolf and van de Straete *Bull Soc Chim Belg* 44 288 1935; UV: Rondestvedt et al. *J Am Chem Soc* 78 6115 1956; IR: Chiorboli and Mirone *Ann Chim (Rome)* 42 681 1952.]

Maleuric acid [105-61-3] M 158.1, m 167-168°(dec). Crystd from hot water.

dl-Malic acid [617-48-1 and 6915-15-7] M 134.1, m 128-129°. Crystd from acetone, then from acetone/CCl₄, or from ethyl acetate by adding pet ether (b 60-70°). Dried at 35° under 1mm pressure to avoid formation of the anhydride.

L-Malic acid [97-67-6] M 134.1, m 104.5-106°, [α]_D²⁰ -2.3° (c 8.5, H₂O), pK₁²⁵ 3.46, pK₂²⁵ 5.10. Crystd (charcoal) from ethyl acetate/pet ether (b 55-56°), keeping the temperature below 65°. Or, dissolved by refluxing in fifteen parts of anhydrous diethyl ether, decanted, concentrated to one-third volume and crystd at 0°, repeatedly to constant melting point.

Malonamide [108-13-4] M 102.1, m 170°. Crystd from water.

Malonic acid [141-82-2] M 104.1, m 136°, pK₁²⁵ 2.58, pK₂²⁵ 5.69. Crystd from *benzene/diethyl ether (1:1) containing 5% of pet ether (b 60-80°), washed with diethyl ether, then recrystd from H₂O or acetone. Dried under vac over conc H₂SO₄.

Malononitrile [109-77-3] M 66.1, m 32-34°, b 109°/20mm, 113-118°/25mm, 220°/760mm. Crystd from water, EtOH, *benzene or chloroform. Distd in a vacuum from, and stored over, P₂O₅. [Bernasconi et al. *J Am Chem Soc* 107 7692 1985; Gratenhuis *J Am Chem Soc* 109 8044 1987].

Maltol (3-hydroxy-2-methyl-4-pyrone) [118-71-8] M 126.1, m 161-162°. Crystd from CHCl₃ or aqueous 50% EtOH. Volatile in steam. It can be readily sublimed in a vacuum.

Maltose (H₂O) [6363-53-7] M 360.3, m 118°. Purified by chromatography from aqueous soln on to a charcoal/Celite (1:1) column, washed with water to remove glucose and other monosaccharides, then eluted with aqueous 75% EtOH. Crystd from water, aqueous EtOH or EtOH containing 1% nitric acid. Dried as the monohydrate at room temperature under vacuum over H₂SO₄ or P₂O₅.

Mandelic acid (α-hydroxyphenylacetic acid) [*S*-(+)-17199-29-0; *R*-(-)-611-71-2] M 152.2, m 130-133°, 133°, 133.1° (evacuated capillary), 133-133.5°, [α]₅₄₆²⁰ (+) and (-) 188° (c 5, H₂O), [α]_D²⁰ (+) and (-) 155° (c 5, H₂O) and (+) and (-) 158° (c 5, Me₂CO), pK²⁵ 3.41. Purified by recrystn from H₂O, *C₆H₆ or CHCl₃. [Roger *J Chem Soc* 2168 1932; Jamison and Turner *J Chem Soc* 611 1942.] They have solubilities in H₂O of ca 11% at 25°. [Banks and Davies *J Chem Soc* 73 1938.] The *S*-benzylisothiuronium salt has m 180° (from H₂O) and [α]_D²⁵ (+) and (-) 57° (c 20, EtOH) [El Masri et al. *Biochem J* 68 199 1958].

RS-(±)-Mandelic acid [61-72-3] M 152.2, m 118°, 120-121°. Purified by Soxhlet extraction with *benzene (about 6mL/g), allowing the extract to crystallise. Also crystallises from CHCl₃. The *S*-benzylisothiuronium salt has m 169° (166°) (from H₂O). Dry at room temperature under vacuum.

D-Mannitol [69-65-8] M 182.2, m 166.1°, [α]₅₄₆²⁰ + 29° (c 10, after 1h in 8% borax soln). Crystd from EtOH or distilled water and dried at 100°.

Mannitol hexanitrate [15825-70-4] M 452.2, m 112-113°. Crystd from EtOH. **EXPLOSIVE (on detonation).**

α-D-Mannose [3458-28-4] M 180.2, m 132°, [α]_D²⁰ +14.1° (c 4, H₂O). Crystd repeatedly from EtOH or aq 80% EtOH, then dried under vacuum over P₂O₅ at 60°.

Meconic acid (3-hydroxy-γ-pyrone-2,6-dicarboxylic acid) [497-59-6] M 200.1, m 100° (loses H₂O), pK₁²⁵ 1.83, pK₂²⁵ 2.3, pK₃²⁵ 10.10. Crystd from water and dried at 100° for 20min.

Melamine (2,4,6-triamino-1,3,5-triazine) [108-78-1] M 126.1, m 353°, pK²⁵ 5.00. Crystd from water or dilute aqueous NaOH.

D(+)-Melezitose (H₂O) [597-12-6] M 540.5, m 153-154°(dec), 2H₂O m 160°(dec), [α]_D²⁰ +88° (c 4, H₂O). Crystallises from water as the dihydrate, then dried at 110° (anhydrous).

D(+)-Melibiose (2H₂O) [585-99-9, 66009-10-7] M 360.3, m 84-85°, [α]_D²⁰ +135° (c 5, after 10h H₂O). Crystallises as a hydrate from water or aqueous EtOH.

(±)-Mellein [(±)-3,4-dihydro-8-hydroxy-3-methyl-2-benzopyran-1-one] [1200-93-7] M 178.2, m 37-39°, 39°, pK_{Est} ~9.5. Purified by recrystn from aqueous EtOH. It has UV max at 247 and 314nm. [Arakawa et al. *Justus Liebigs Ann Chem* 728 152 1969; Blair and Newbold *Chem Ind (London)* 93 1955, *J Chem Soc* 2871 1955.] The methyl ether has m 66-67° and UV: λ_{max} 242nm (ε 7,400) and 305nm (ε 4,600).

Melphalan (4-[bis-{2-chloroethyl}amino]-L-phenylalanine) [148-82-3] M 305.2, m 182-183° (dec), 183-185°, [α]_D²⁵ +7.5° (c 1.33, 1.0 N HCl), [α]_D²⁰ -28° (c 0.8, MeOH), pK_{Est} ~6.4. Purified by recrystn from MeOH and its solubility is 5% in 95% EtOH containing one drop of 6N HCl.

It is soluble in EtOH and propylene glycol but is almost insoluble in H₂O. The *RS*-form has *m* 180-181° and the *R*-form crystallises from MeOH with a *m* 181.5-182° and $[\alpha]_D^{21} -7.5^\circ$ (c 1.26, 1.0 N HCl). [Bergel and Stock *J Chem Soc* 2409 1954.]

(-)-Menthol [2216-51-5] *M* 156.3, *m* 44-46.5°, $[\alpha]_D 50^\circ$ (c 10, EtOH). Crystd from CHCl₃, pet ether or EtOH/water.

1*R*-(*-*)-Menthyl chloride (1*S*,2*R*,4*R*-2-chloro-1-isopropyl-1-methylcyclohexane) [16052-42-9] *M* 174.7, *m* -20.1° to -16.5°, *b* 88.5°/12.5mm, 101-105°/21mm, $d_D^{20} 0.936$, $n_D^{20} 1.463$ (neat). Dissolve in pet ether (b 40-60°), wash with H₂O, conc H₂SO₄ until no discoloration of the organic layer occurs (care with the use of conc H₂SO₄ during shaking in a separating funnel), again with H₂O and dry over MgSO₄. Evaporate the pet ether and dist the residual oil through a Claisen head with a Vigreux neck (head) of ca 40 cm length. [Smith and Wright *J Org Chem* 17 1116 1952; Barton et al. *J Chem Soc* 453 1952.]

Meprobamate [2,2-di(carbamoyloxymethyl)pentane] [57-53-4] *M* 246.3, *m* 104-106°. Crystd from hot water. Could be an addictive drug.

2-Mercaptobenzimidazole [583-39-1] *M* 150.2, *m* 302-304°, 312°, *pK*²⁰ 10.24. Crystd from aq EtOH, AcOH or aq ammonia.

2-Mercaptobenzothiazole [149-30-4] *M* 167.2, *m* 182°, *pK*²⁵ 7.5 (50% aq AcOH). Crystd repeatedly from 95% EtOH, or purified by incomplete pptn by dilute H₂SO₄ from a basic soln, followed by several crystns from acetone/H₂O or *benzene. Complexes with Ag, Au, Bi, Cd, Hg, Ir, Pt, and Tl.

2-Mercaptoethanol [60-24-2] *M* 78.1, *b* 44°/4mm, 53.5°/10mm, 58°/12mm, 68°/20mm, 78.5°/40mm, 96-97° (92°)/100mm, 157°/748mm, $d_4^{20} 1.114$, $n_D^{20} 1.500$, *pK*²⁵ 9.72 (9.43). Purified by distn in a vacuum. Distn at atmospheric pressure causes some oxidation and should be done in an inert atmosphere. [Woodward *J Chem Soc* 1892 1948.] It has a foul odour, is irritating to the eyes, nose and skin — should be handled in an efficient fume cupboard. It is miscible with H₂O, EtOH, Et₂O and *C₆H₆ and has a UV max at 235nm. The 2,4-dinitrophenyl thioether has *m* 101-102°(from EtOH or aq MeOH) [Grogen et al. *J Org Chem* 20 50 1955].

2-Mercaptoethylamine [60-23-1] *M* 77.2. See cysteamine p. 525 in Chapter 6.

2-Mercaptoimidazole [872-35-5] *M* 100.1, *m* 221-222°, *pK*₁²⁰ -1.6, *pK*₂²⁰ 11.6. Crystd from H₂O.

2-Mercapto-1-methylimidazole [60-56-0] *M* 114.2, *m* 145-147°, *pK*₁²⁰ -2.0, *pK*₂²⁰ 11.9. Crystd from EtOH.

6-Mercaptopurine (H₂O) [6112-76-1] *M* 170.2, *m* >315°(dec), *pK*₁²⁰ 0.5, *pK*₂²⁰ 7.77, *pK*₃²⁰ 10.8. Crystd from pyridine (30mL/g), washed with pyridine, then triturated with water (25mL/g), adjusting to pH 5 by adding M HCl. Recrystd by heating, then cooling, the soln. Filtered, washed with water and dried at 110°. Has also been crystd from water (charcoal).

8-Mercaptoquinoline (2H₂O, thioxine) [491-33-8] *M* 197.3, *m* 58-59°, *pK*₁²⁵ 2.0, *pK*₂²⁵ 8.40. Easily oxidised in air to give diquinolyl-8,8'-disulfide (which is stable). It is more convenient to make 8-mercaptoquinoline by reduction of the material. [Nakamura and Sekido *Talanta* 17 515 1970.]

Mesaconic acid (methylfumaric acid) [498-24-8] *M* 130.1, *m* 204-205°, *pK*¹⁸ 4.82. Crystd from water or EtOH [Katakis et al. *J Chem Soc, Dalton Trans* 1491 1986].

Mescaline sulfate [2-(3,4,5-trimethoxyphenyl)ethylamine sulfate] [5967-42-0] *M* 309.3, *m* 183-184°, *pK*_{Est} ~9.7. Crystd from water.

Mesitylene (1,3,5-trimethylbenzene) [108-67-8] M 120.2, m -44.7°, b 99.0-99.8°/100mm, 166.5-167°/760mm, n²⁵ 1.4967, d 0.865. Dried with CaCl₂ and distd from Na in a glass helices packed column. Treated with silica gel and redistd. Alternative purifications include vapour-phase chromatography, or fractional distn followed by azeotropic distn with 2-methoxyethanol (which is subsequently washed out with H₂O), drying and fractional distn. More exhaustive purification uses sulfonation by dissolving in two volumes of conc H₂SO₄, precipitating with four volumes of conc HCl at 0°, washing with conc HCl and recrystallising from CHCl₃. The mesitylene sulfonic acid is hydrolysed with boiling 20% HCl and steam distd. The separated mesitylene is dried (MgSO₄ or CaSO₄) and distilled. It can also be fractionally crystd from the melt at low temperatures.

Mesityl oxide [141-79-7] M 98.2, b 112°/760mm, n²⁴ 1.4412, d 0.854, pK²⁰ -5.36 (H₀ scale, aq H₂SO₄). Purified via the semicarbazone (m 165°). [Erskine and Waight *J Chem Soc* 3425 1960.]

Metalphthalein (H₂O) (o-cresolcomplexon) [2411-89-4] M 636.6, m 186°(dec). See o-cresolphthalein complexone on p. 173.

Metanilic acid (3-aminobenzenesulfonic acid) [121-47-1] M 173.2, m <300°(dec), pK₁²⁵ < 1, pK₂²⁵ 3.74. Crystd from water (as the hydrate), under CO₂ in a semi-darkened room. (The soln is photosensitive.) Dried over 90% H₂SO₄ in a vac desiccator.

α-Methacraldehyde [78-85-3] M 68.1, b 68.4°, d 0.849, n 1.416. Fractionally distd under nitrogen through a short Vigreux column. Stored in sealed ampoules. (Slight polymerisation may occur.)

Methacrylamide [79-39-0] M 85.1, m 111-112°. Crystd from *benzene or ethyl acetate and dried under vacuum at room temperature.

Methacrylic acid [79-41-4] M 86.1, b 72°/14mm, 160°/760mm, d 1.015, n 1.431, pK 4.65. Aq methacrylic acid (90%) was satd with NaCl (to remove the bulk of the water), then the organic phase was dried with CaCl₂ and distd under vacuum. Polymerisation inhibitors include 0.25% p-methoxyphenol, 0.1% hydroquinone, or 0.05% N,N'-diphenyl-p-phenylenediamine.

Methacrylic anhydride [760-93-0] M 154.2, b 65°/2mm, d 1.040, n 1.454. Distd at 2mm pressure, immediately before use, in the presence of hydroquinone.

Methacrylonitrile [126-98-7] M 67.1, b 90.3°, d 0.800, n 1.4007, n³⁰ 1.3954. Washed (to remove inhibitors such as p-tert-butylcatechol) with satd aq NaHSO₃, 1% NaOH in saturated NaCl and then with saturated NaCl. Dried with CaCl₂ and fractionally distd under nitrogen to separate from impurities such as methacrolein and acetone.

Methane [74-82-8] M 16.0, m -184°, b -164°/760mm, -130°/6.7atm, d¹⁶⁴ 0.466 (air 1). Dried by passage over CaCl₂ and P₂O₅, then passed through a Dry-ice trap and fractionally distd from a liquid-nitrogen trap. Oxygen can be removed by prior passage in a stream of hydrogen over reduced copper oxide at 500°, and higher hydrocarbons can be removed by chlorinating about 10% of the sample: the hydrocarbons, chlorides and HCl are readily separated from the methane by condensing the sample in the liquid-nitrogen trap and fractionally distilling it. Methane has also been washed with conc H₂SO₄, then solid NaOH and then 30% NaOH soln. It was dried with CaCl₂, then P₂O₅, and condensed in a trap at liquid air temp, then transferred to another trap cooled in liquid nitrogen. CO₂, O₂, N₂ and higher hydrocarbons can be removed from methane by adsorption on charcoal. [Eiseman and Potter *J Res Nat Bur Stand* 58 213 1957.] **HIGHLY FLAMMABLE.**

Methanesulfonic acid [76-75-2] M 96.1, m 20°, b 134.5-135°/3mm, d 1.483, n 1.432, pK²⁵ -1.86 (-1.2). Dried, either by azeotropic removal of water with *benzene or toluene, or by stirring 20g of P₂O₅ with 500mL of the acid at 100° for 0.5h. Then distd under vacuum and fractionally crystd by partial

freezing. Sulfuric acid, if present, can be removed by prior addition of $\text{Ba}(\text{OH})_2$ to a dilute soln, filtering off the BaSO_4 and concentrating under reduced pressure, and is sufficiently pure for most applications.

Methanesulfonyl chloride [124-63-0] M 114.5. b $55^\circ/11\text{mm}$, d 1.474, n 1.452. Distd from P_2O_5 under vacuum.

Methanol [67-56-1] M 32.0, b 64.5° , d^{15} 0.79609, d^{25} 1.32663, n^{15} 1.33057, n^{25} 1.32663, pK^{25} 15.5. Almost all methanol is now obtained synthetically. Likely impurities are water, acetone, formaldehyde, ethanol, methyl formate and traces of dimethyl ether, methylal, methyl acetate, acetaldehyde, carbon dioxide and ammonia. Most of the water (down to about 0.01%) can be removed by fractional distn. Drying with CaO is unnecessary and wasteful. Anhydrous methanol can be obtained from "absolute" material by passage through Linde type 4A molecular sieves, or by drying with CaH_2 , CaSO_4 , or with just a little more sodium than required to react with the water present; in all cases the methanol is then distd. Two treatments with sodium reduces the water content to about $5 \times 10^{-5}\%$. [Friedman, Gill and Doty *J Am Chem Soc* 83 4050 1961.] Lund and Bjerrum [*Chem Ber* 64 210 1931] warmed clean dry magnesium turnings (5g) and iodine (0.5g) with 50-75mL of "absolute" methanol in a flask until the iodine disappeared and all the magnesium was converted to methoxide. Up to 1L of methanol was added and, after refluxing for 2-3h, it was distd off, excluding moisture from the system. Redistn from tribromobenzoic acid removes basic impurities and traces of magnesium oxides, and leaves conductivity-quality material. The method of Hartley and Raikes [*J Chem Soc* 127 524 1925] gives a slightly better product. This consists of an initial fractional distn, followed by distn from aluminium methoxide, and then ammonia and other volatile impurities are removed by refluxing for 6h with freshly dehydrated CuSO_4 (2g/L) while dry air is passed through: the methanol is finally distd. (The aluminium methoxide is prepared by warming with aluminium amalgam (3g/L) until all the aluminium has reacted. The amalgam is obtained by warming pieces of sheet aluminium with a soln of HgCl_2 in dry methanol). This treatment also removes aldehydes.

If acetone is present in the methanol, it is usually removed prior to drying. Bates, Mullaly and Hartley [*J Chem Soc* 401 1923] dissolved 25g of iodine in 1L of methanol and then poured the soln, with constant stirring, into 500mL of M NaOH. Addition of 150mL of water ppted iodoform. The soln was stood overnight, filtered, then boiled under reflux until the odour of iodoform disappeared, and fractionally distd. (This treatment also removes formaldehyde.) Morton and Mark [*Ind Eng Chem (Anal Edn)* 6 151 1934] refluxed methanol (1L) with furfural (50mL) and 10% NaOH soln (120mL) for 6-12h, the refluxing resin carrying down with it the acetone and other carbonyl-containing impurities. The alcohol was then fractionally distd. Evers and Knox [*J Am Chem Soc* 73 1739 1951], after refluxing 4.5L of methanol for 24h with 50g of magnesium, distd off 4L of it, which they then refluxed with AgNO_3 for 24h in the absence of moisture or CO_2 . The methanol was again distd, shaken for 24h with activated alumina before being filtered through a glass sinter and distd under nitrogen in an all-glass still. Material suitable for conductivity work was obtained.

Variations of the above methods have also been used. For example, a sodium hydroxide soln containing iodine has been added to methanol and, after standing for 1day, the soln has been poured slowly into about a quarter of its volume of 10% AgNO_3 , shaken for several hours, then distd. Sulfanilic acid has been used instead of tribromobenzoic acid in Lund and Bjerrum's method. A soln of 15g of magnesium in 500mL of methanol has been heated under reflux, under nitrogen, with hydroquinone (30g), before degassing and distilling the methanol, which was subsequently stored with magnesium (2g) and hydroquinone (4g per 100mL). Refluxing for about 12h removes the bulk of the formaldehyde from methanol: further purification has been obtained by subsequent distn, refluxing for 12h with dinitrophenylhydrazine (5g) and H_2SO_4 (2g/L), and again fractionally distilling.

Rapid purification: Methanol purification is the same as for Ethanol.

Another simple purification procedure consists of adding 2g of NaBH_4 to 1.5L methanol, gently bubbling with argon and refluxing for a day at 30° , then adding 2g of freshly cut sodium (washed with methanol) and refluxing for 1day before distilling. The middle fraction is taken. [Jou and Freeman *J Phys Chem* 81 909 1977.]

dl-Methionine (RS- α -aminohexanoic acid) [59-51-8] M 149.2, m $281^\circ(\text{dec})$. Crystd from hot water.

L-Methionine [63-68-3] M 149.2, m $283^\circ(\text{dec})$, $[\alpha]_D^{25}$ $+21.2^\circ$ (0.2M HCl) pK_1^{25} 2.13, pK_2^{25} 9.73. Crystd from aqueous EtOH.

***dl*-Methionine sulfoxide** [454-41-1, 62697-73-8] M 165.2, m >240°(dec). Likely impurities are *dl*-methionine sulfone and *dl*-methionine. Crystd from water by adding EtOH in excess.

Methoxyacetic acid [625-45-6] M 90.1, b 97°/13-14mm, d 1.175, n 1.417, pK²⁵ 3.57. Fractionally crystd by repeated partial freezing, then fractionally distd under vacuum through a vacuum-jacketed Vigreux column 20cm long.

***p*-Methoxyacetophenone** [100-06-1] M 150.2, m 39°, b 139°/15mm, 264°/736mm. Crystd from diethyl ether/pet ether.

Methoxyamine hydrochloride [593-56-6] M 83.5, m 151-152°, pK²⁵ 4.60. Crystd from absolute EtOH or EtOH by addition of diethyl ether. [Kovach et al. *J Am Chem Soc* 107 7360 1985.]

***p*-Methoxyazobenzene** [2396-60-3] M 212.3, m 54-56°. Crystd from EtOH.

3-Methoxybenzanthrone [3688-79-7] M 274.3, m 173°. Crystd from *benzene, EtOH or Me₂CO as yellow needles.

***m*-Methoxybenzoic acid (*m*-anisic acid)** [586-38-9] M 152.2, m 110°, pK²⁵ 4.09. Crystd from EtOH/water.

***p*-Methoxybenzoic acid (*p*-anisic acid)** [100-09-4] M 152.2, m 184.0-184.5°, pK²⁵ 4.51. Crystd from EtOH, water, EtOH/water or toluene.

4-Methoxybenzyl chloride (anisyl chloride) [824-94-2] M 156.6, m -1°, b 76°/0.1mm, 95°/5mm, 110°/10mm, 117-117.5°/14mm, 117°/18mm, d₄²⁰ 1.15491, n_D²⁰ 1.55478. Purified by fractional distn under vacuum and the middle fraction is redistd at 10⁻⁶ mm at room temperature by intermittent cooling of the receiver in liquid N₂, and the middle fraction is collected. [Mohammed and Kosower *J Am Chem Soc* 93 2709 1971.]

3-Methoxycarbonyl-2,5-dihydrothiophen-1,1-dioxide [67488-50-0] M 176.1, m 57-58°, 60-62°. If IR show CO bands then dissolve in CHCl₂, wash with aqueous Na₂CO₃ and H₂O, dry over MgSO₄, filter, evaporate and wash the residue with cold Et₂O and dry *in vacuo*. NMR (CDCl₃): δ 7.00 (m 1H), 3.98 (bs 4H) and 3.80 (s Me). [Mcintosh and Sieber *J Org Chem* 43 4431 1978.]

"Methoxychlor", 1,1-Bis(*p*-methoxyphenyl)-2,2,2-trichloroethane (dimorphic) [72-43-5] M 345.7, m 78-78.2°, or 86-88°. Freed from 1,1-bis(*p*-chlorophenyl)-2,2,2-trichloroethane by crystn from EtOH.

***trans-p*-Methoxycinnamic acid** [830-09-1, 943-89-5 (*trans*)] M 178.2, m 173.4-174.8°, pK²⁵ 4.54. Crystd from MeOH to constant melting point and UV spectrum.

2-Methoxyethanol (methylcellosolve) [109-86-4] M 76.1, b 124.4°, d 0.964, n 1.4017, pK²⁵ 14.8. Peroxides can be removed by refluxing with stannous chloride or by filtration under slight pressure through a column of activated alumina. 2-Methoxyethanol can be dried with K₂CO₃, CaSO₄, MgSO₄ or silica gel, with a final distn from sodium. Aliphatic ketones (and water) can be removed by making the solvent 0.1% in 2,4-dinitrophenylhydrazine and allowing to stand overnight with silica gel before fractionally distilling.

2-Methoxyethoxymethylchloride (MEMCl) [3970-21-6] M 124.6, b 50-52°/13mm, 140-145°(dec)/atm, d 1.092, n 1.427. Possible impurities are methoxyethanol (b 124°/atm), HCHO and HCl which can be removed below the b of MEMCl. Purify by fractional distn in a vacuum. If too impure, prepare from methoxyethanol (152g) and *s*-trioxane (66g) by bubbling a stream of dry HCl (with stirring) until a clear mixt is obtained. Dilute with pentane (900mL), dry (3h over 100g MgSO₄, at 5°), evaporate and the residue is distd in a vac. It is MOISTURE SENSITIVE and TOXIC. The MEM.NEt₃⁺Cl⁻ salt, prepared by reactn with

1.3 equivs of Et₃N (16h/25°) and dried in vac has m 58-61°, and is moisture sensitive. [Corey et al. *Tetrahedron Lett* 809 1976.]

β-Methoxyethylamine [109-85-3] M 75.1, b 94°, d 0.874, n 1.407, pK²⁵ 9.40. An aqueous 70% soln was dehydrated by azeotropic distn with *benzene or methylene chloride and the amine was distilled twice from zinc dust. Store in a tight container as it absorbs CO₂ from the atmosphere.

6-Methoxy-1-indanone [13623-25-1] M 162.2, m 151-153°. Crystd from MeOH, then sublimed in a high vacuum.

5-Methoxyindole [1006-94-6] M 147.2, m 55°, 57°, b 176-178°/17mm, pK_{Est} ~0. Crystd from cyclohexane pet ether or pet ether/Et₂O.

1-Methoxynaphthalene [2216-69-5] M 158.2, b 268.4-268.5°. See methyl 1-naphthyl ether on p. 295.

2-Methoxynaphthalene [93-04-9] M 158.2, m 73.0-73.6°, b 273°/760mm. Fractionally distd under vacuum. Crystd from absolute EtOH, aqueous EtOH, *benzene or *n*-heptane, and dried under vacuum in an Abderhalden pistol or distd *in vacuo*. [Kikuchi et al. *J Phys Chem* 91 574 1987.]

1-Methoxy-4-nitronaphthalene [4900-63-4] M 203.2, m 85°. Purified by chromatography on silica gel and recrystd from MeOH. [Bunce et al. *J Org Chem* 52 4214 1987.]

***p*-Methoxyphenol** [150-76-5] M 124.1, m 54-55°, b 243°, pK²⁵ 10.21. Crystd from *benzene, pet ether or H₂O, and dried under vacuum over P₂O₅ at room temp. Sublimes *in vacuo*. [Wolfenden et al. *J Am Chem Soc* 109 463 1987.]

α-Methoxyphenylacetic acid (O-methyl mandelic acid), [*R*-(-)- 3966-32-3; *S*-(+)- 26164-26-1] M 166.2, m 62.9°, 62-65°, 65-66°, [α]₅₄₆²⁰ (-) and (+) 179° (169.8°), [α]_D²⁰ (-) and (+) 150.7° (148°) (c 0.5, EtOH), pK_{Est} ~3.1. Purified by recrystn from *C₆H₆-pet ether (b 80-100°). [Neilson and Peters *J Chem Soc* 1519 1962; Weizmann et al. *J Am Chem Soc* 70 1153 1948; Pirie and Smith *J Chem Soc* 338 1932; NMR: Dale and Mosher *J Am Chem Soc* 95 512 1973; for resolution: Roy and Deslongchamps *Can J Chem* 63 651 1985; Trost et al. *J Am Chem Soc* 108 4974 1986.] The *racemic mixture* has m 72°, b 121-122°/0.4mm, 165°/18mm (from pet ether) [Braun et al. *Chem Ber* 63 2847 1930].

***o*-Methoxyphenylacetic acid** [93-25-4] M 166.2, m 71.0-71.2°, pK_{Est} ~4.4. Crystd from H₂O, EtOH or aq EtOH and dried in a vacuum desiccator over Sicapent.

***m*-Methoxyphenylacetic acid** [1798-09-0] M 166.2, m 71.0-71.2°, pK_{Est} ~4.3. Crystd from H₂O, or aq EtOH.

***p*-Methoxyphenylacetic acid** [104-01-8] M 166.2, m 85-87°, pK²⁵ 4.36. Crystd from EtOH/water.

5-(*p*-Methoxyphenyl)-1,2-dithiole-3-thione [42766-10-9] M 240.2, m 111°. Crystd from butyl acetate.

***N*-(*p*-Methoxyphenyl)-*p*-phenylenediamine** [101-64-4] M 214.3, m 102°, b 238°/12mm, pK 6.6 (5.9). Crystd from ligroin.

8-Methoxypsoralen see xanthotoxin p. 577 in Chapter 6.

α-Methoxy-α-trifluoromethylphenylacetic acid (MTPA, Mosher's acid) [*R*-(+)- 20445-31-2; *S*-(-)- 17257-71-5] M 234.2, m 43-45°, 90°/0.1mm, 105-107°/1mm, [α]₅₄₆²⁰ (+) and (-) 87°, [α]_D²⁰ (+) and (-) 73° (c 2, MeOH), pK_{Est} ~2.5. A likely impurity is phenylethylamine from the

resolution. Dissolve acid in ether-benzene (3:1), wash with 0.5N H₂SO₄, then water, dry over magnesium sulfate, filter, evaporate and distil. [Dale et al. *J Org Chem* **34** 2543 1969, *J Am Chem Soc* **75** 512 1973.]

α -Methoxy- α -trifluoromethylphenylacetyl chloride [*R*-(-)- 39637-99-5; *S*-(+)- 20445-33-4] **M 252.6, b 54-56°/1mm, 213-214°/atm, d_4^{20} 1.353, n_D^{20} 1.468, $[\alpha]_{546}^{20}$ (-) and (+) 167°, $[\alpha]_D^{20}$ (-) and (+) 137° (c 4, CCl₄), $[\alpha]_D^{24}$ (-) and (+) 10.0° (neat).** The most likely impurity is the free acid due to hydrolysis and should be checked by IR. If free from acid then distil taking care to keep moisture out of the apparatus. Otherwise add SOCl₂ and reflux for 50h and distil. Note that shorter reflux times resulted in a higher boiling fraction (b 130-155°/1mm) which has been identified as the anhydride. [Dale et al. *J Org Chem* **34** 2543 1969; for enantiomeric purity see *J Am Chem Soc* **97** 512 1973.]

***N*-Methylacetamide** [79-16-3] **M 73.1, m 30°, b 70-71°/2.5-3mm, pK_1^{25} -3.70, pK_2^{25} -0.42.** Fractionally distd under vacuum, then fractionally crystd twice from its melt. Impurities include acetic acid, methyl amine and H₂O. For detailed purification procedure, see Knecht and Kolthoff, *Inorg Chem* **1** 195 1962. Although *N*-methylacetamide is commercially available it is often extensively contaminated with acetic acid, methylamine, water and an unidentified impurity. The recommended procedure is to synthesise it in the laboratory by direct reaction. The gaseous amine is passed into hot glacial acetic acid, to give a partially aq soln of methylammonium acetate which is heated to ca 130° to expel water. Chemical methods of purification such as extractn by pet ether, treatment with H₂SO₄, K₂CO₃ or CaO can be used but are more laborious. Tests for purity include the Karl Fischer titration for water; this can be applied directly. Acetic acid and methylamine can be detected polarographically.

In addition to the above, purification of *N*-methylacetamide can be achieved by fractional freezing, including zone melting, repeated many times, or by chemical treatment with vacuum distn under reduced pressures. For details of zone melting techniques, see Knecht in *Recommended Methods for Purification of Solvents and Tests for Impurities*, Coetzee Ed. Pergamon Press 1982.

***N*-Methylacetanilide** [579-10-2] **M 149.2, m 102-104°.** Crystd from water, ether or pet ether (b 80-100°).

Methyl acetate [79-20-9] **M 74.1, b 56.7-57.2°, d 0.934, n 1.36193, n^{25} 1.3538, pK^{20} -7.28 (H₀ scale, aq H₂SO₄).** Methanol in methyl acetate can be detected by measuring solubility in water. At 20°, the solubility of methyl acetate in water is ca 35g per 100mL, but 1% MeOH confers miscibility. Methanol can be removed by conversion to methyl acetate, using refluxing for 6h with acetic anhydride (85mL/L), followed by fractional distn. Acidic impurities can be removed by shaking with anhydrous K₂CO₃ and distilling. An alternative treatment is with acetyl chloride, followed by washing with conc NaCl and drying with CaO or MgSO₄. (Solid CaCl₂ cannot be used because it forms a crystalline addition compound.) Distn from copper stearate destroys peroxides. Free alcohol or acid can be eliminated from methyl acetate by shaking with strong aq Na₂CO₃ or K₂CO₃ (three times), then with aq 50% CaCl₂ (three times), satd aq NaCl (twice), drying with K₂CO₃ and distn from P₂O₅.

Methyl acetimidate hydrochloride [14777-27-6] **M 109.6, m 93-95°, 105°(dec), pK_{Est} ~ 5.5.** Crystd from methanol by adding dry ether to a ratio of 1:1 and cooled at 0°. Filter off the crystals in a cold room, wash with methanol/ether (1:2), then dry in a vacuum. [Hunter and Ludwig *J Am Chem Soc* **84** 3491 1962.] The free base has b 90-91°/765mm, d 0.867, n 1.403 [Hunter and Ludwig *Methods Enzymol* **25** 585 1973.]

***p*-Methylacetophenone** [122-00-9] **M 134.2, m 22-24°, b 93.5°/7mm, 110°/14mm, d 1.000, n 1.5335.** Impurities, including the *o*- and *m*-isomers, were removed by forming the semicarbazone which, after repeated crystn, was hydrolysed to the ketone. [Brown and Marino *J Am Chem Soc* **84** 1236 1962.] Also purified by distn under reduced pressure, followed by low temperature crystn from isopentane.

Methyl acrylate [96-33-3] **M 86.1, b 80°, d 0.9535, n 1.4040.** Washed repeatedly with aqueous NaOH until free from inhibitors (such as hydroquinone), then washed with distd water, dried (CaCl₂) and fractionally distd under reduced pressure in an all-glass apparatus. Sealed under nitrogen and stored at 0° in the dark. [Bamford and Han *J Chem Soc, Faraday Trans 1* **78** 855 1982.]

1-Methyladamantane [768-91-2] M 150.2, m 103°. Purified by zone melting and sublimes at 90-95°/12mm.

2-Methyladamantane [700-56-1] M 150.2. Purified by zone melting.

Methylamine (gas) [74-89-5] M 31.1, b -7.55°/719mm, pK²⁵ 10.62. Dried with sodium or BaO.

Methylamine hydrochloride [593-51-1] M 67.5, m 231.8-233.4°, b 225-230°/15mm, pK²⁵ 10.62. Crystd from *n*-butanol, absolute EtOH or MeOH/CHCl₃. Washed with CHCl₃ to remove traces of dimethylamine hydrochloride. Dried under vacuum first with H₂SO₄ then P₂O₅. Deliquescent, stored in a desiccator over P₂O₅.

1-Methylaminoanthraquinone [82-38-2] M 237.3, m 166.5°, pK_{Est}~2. Crystd to constant melting point from butan-1-ol, then crystd from EtOH. It can be sublimed under vacuum.

***N*-Methyl-*o*-aminobenzoic acid (*N*-methylantranilic acid)** [119-68-6] M 151.2, m 178.5°, pK₁²⁵ 1.97, pK₂²⁵ 5.34. Crystd from water or EtOH.

***p*-Methylaminophenol sulfate** [55-55-0] M 344.4, m 260°(dec), pK²⁵ 5.9. Crystd from MeOH.

6-Methylaminopurine [443-72-1] M 149.2, m >300°, 312-314° (dec), pK₁²⁰ <1, pK₂²⁰ 4.15, pK₃²⁰ 10.02. Best purified by recrystallising 2g from 50mL of H₂O and 1.2g of charcoal. [UV: Albert and Brown *J Chem Soc* 2060 1954; UV: Mason *J Chem Soc* 2071 1954; see also Elion et al. *J Am Chem Soc* 74 411 1952.] The *picrate* has m 265°(257°) [Bredereck et al. *Chem Ber* 81 307 1948].

Methyl 3-aminopyrazine-2-carboxylate [16298-03-6] M 153.1, m 169-172°, 172°. Forms yellow needles from H₂O (100 parts using charcoal). If it contains the free acid then dissolve in CH₂Cl₂ wash with saturated aqueous Na₂CO₃, brine, dry over MgSO₄ filter, evaporate and recrystallise the residue. The *free acid* has m 203-204° (dec) [UV: Brown and Mason *J Chem Soc* 3443 1956] and pK₁ <1 and pK₂ 3.70. The *ammonium salt* has m 232° (dec) (from aq Me₂CO) and the *amide* has m 239.2° (from H₂O) [Ellingson et al. *J Am Chem Soc* 67 1711 1945].

***N*-Methylaniline** [100-61-8] M 107.2, b 57°/4mm, 81-82°/14mm, d 0.985, n 1.570, pK²⁵ 4.56. Dried with KOH pellets and fractionally distd under vacuum. Acetylated and the acetyl derivative was recrystd to constant melting point (m 101-102°), then hydrolysed with aqueous HCl and distd from zinc dust under reduced pressure. [Hammond and Parks *J Am Chem Soc* 77 340 1955.]

***N*-Methylaniline hydrochloride** [2739-12-0] M 143.7, m 123.0-123.1°. Crystd from dry *benzene/CHCl₃ and dried under vacuum.

Methyl *p*-anisate [121-98-2] M 166.2, m 48°. Crystd from EtOH.

4-Methyl anisole [104-93-8] M 122.2, b 175-176°, d₁₅¹⁵ 0.9757, n 1.512. Dissolved in diethyl ether, washed with M NaOH, water, dried (Na₂CO₃), evaporated and the residue distd under vacuum.

2-Methylantracene [613-12-7] M 192.3, m 204-206° Chromatographed on silica gel with cyclohexane as eluent and recrystd from EtOH [Werst *J Am Chem Soc* 109 32 1987].

4-Methylantracene [779-02-2] M 192.3, m 77-79°, b 196-197°/12mm, d 1.066. Chromatographed on silica gel with cyclohexane as eluent and recrystd from EtOH [Werst *J Am Chem Soc* 109 32 1987].

2-Methylantraquinone [84-54-8] M 222.3, m 176°. Crystd from EtOH, then sublimed.

Methylarenes (see also pentamethyl- and hexamethyl- benzenes). Recrystd from EtOH and sublimed in vacuum [Schlesener et al. *J Am Chem Soc* 106 7472 1984].

Methyl benzoate [93-58-3] M 136.2, b 104-105°/39mm, 199.5°/760mm, d 1.087, n¹⁵ 1.52049, n 1.51701, pK²⁰ -8.11, -6.51 (H₀ scale, aq H₂SO₄). Washed with dilute aqueous NaHCO₃, then water, dried with Na₂SO₄ and fractionally distd under reduced pressure.

p-Methylbenzophenone [134-84-9] M 196.3, m 57°. Crystd from MeOH and pet ether.

Methyl-1,4-benzoquinone [553-97-9] M 122.1, m 68-69°. Crystd from heptane or EtOH, dried rapidly (vacuum over P₂O₅) and stored under vacuum.

Methyl benzoylformate [15206-55-0] M 164.2, m 246-248°. Purified by radial chromatography (diethyl ether/hexane, 1:1), and dried at 110-112° at 6mm pressure. [Meyers and Oppenlaender *J Am Chem Soc* 108 1989 1986.]

2-Methyl-3,4-benzphenanthrene [652-04-0] M 242.3, m 70°. Crystd from EtOH.

dl- α -Methylbenzyl alcohol [13323-81-4] M 122.2, b 60.5-61.0°/3mm. See dl-1-phenylethanol on p. 330.

R-(+)- α -Methylbenzylamine [R(+) 3886-69-9, RS 618-36-0] M 121.2, b 187-188°/atm, [α]_{D²⁰}²⁰ +35° (c 10, EtOH), [α]_{D²⁵}²⁵ +39.7° (neat), pK 9.08 (for RS). Dissolve in toluene, dry over NaOH and distd, fraction boiling at 187-188°/atm is collected. Store under N₂ to avoid forming the carbamate and urea. Similarly for the S(-) enantiomer [2627-86-3]. [*Org Synth Coll Vol II* 503 1943.]

p-Methylbenzyl bromide [104-81-4]. See α -bromo-p-xylene on p. 143.

p-Methylbenzyl chloride [104-82-5] M 140.6, b 80°/2mm, d 1.085, n 1.543. Dried with CaSO₄ and fractionally distd under vacuum.

Methylbixin [26585-94-4] M 408.5, m 163°. Crystd from EtOH/CHCl₃.

Methyl bromide [74-83-9] M 94.9, b 3.6°. Purified by bubbling through conc H₂SO₄, followed by passage through a tube containing glass beads coated with P₂O₅. Also purified by distn from AlBr₃ at -80°, by passage through a tower of KOH pellets and by partial condensation.

Methyl o-bromobenzoate [610-94-6] M 215.1, b 122°/17mm, 234-244°/760mm. Soln in ether is washed with 10% aqueous Na₂CO₃, water, then dried and distd.

Methyl p-bromobenzoate [619-42-1] M 215.1, m 79.5-80.5°. Crystd from MeOH.

2-Methylbutane (isopentane) [78-78-4] M 72.2, b 27.9°, d 0.621, n 1.35373, n²⁵ 1.35088. Stirred for several hours in the cold with conc H₂SO₄ (to remove olefinic impurities), then washed with H₂O, aqueous Na₂CO₃ and H₂O again. Dried with MgSO₄ and fractionally distd using a Todd column packed with glass helices (see p. 174). Material transparent down to 180nm was obtained by distilling from sodium wire, and passing through a column of silica gel which had previously been dried in place at 350° for 12h before use. [Potts *J Phys Chem* 20 809 1952].

2-Methyl-1-butanol [137-32-6; RS 34713-94-5; S(-)- 1565-80-6] M 88.2, b 130°(RS), 128.6°(S), [α]_{D²⁵}²⁵ -5.8° (neat), d 0.809, n⁵² 1.4082. Refluxed with CaO, distd, refluxed with magnesium and again fractionally distd. A small sample of highly purified material was obtained by fractional crystn after conversion into a suitable ester such as the trinitrophthalate or the 3-nitrophthalate. The latter was converted to the cinchonine salt in acetone and recrystd from CHCl₃ by adding pentane. The salt was saponified, extracted with ether, and fractionally distd. [Terry et al. *J Chem Eng Data* 5 403 1960.]

3-Methyl-1-butanol [123-51-3] M 88.2, b 128°/750mm, 132°/760mm, d^{15} 0.8129, n^{15} 1.4085, n 1.4075. Dried by heating with CaO and fractionally distilling, then heating with BaO and redistilling. Alternatively, boiled with conc KOH, washed with dilute H₃PO₄, and dried with K₂CO₃, then anhydrous CuSO₄, before fractionally distilling. If very dry alcohol is required, the distillate can be refluxed with the appropriate alkyl phthalate or succinate as described for *ethanol*. It is separated from 2-methyl-1-butanol by fractional distn, fractional crystn and preparative gas chromatography.

3-Methyl-2-butanol [598-75-4] M 88.2, b 111.5°, d 0.807, n 1.4095, n^{25} 1.4076. Refluxed with magnesium, then fractionally distd.

3-Methyl-2-butanone (methyl isopropyl ketone) [563-80-4] M 86.1, b 93-94°/752mm, d 0.818, n 1.410, pK^{25} -7.1 (aq H₂SO₄). Refluxed with a little KMnO₄. Fractionated on a spinning-band column, dried with CaSO₄ and distd.

2-Methyl-2-butene [513-35-9] M 70.1, f -133.8°, b 38.4°/760mm, d^{15} 0.66708, d 0.6783, d^{25} 0.65694, n^{15} 1.3908. Distd from sodium.

Methyl *n*-butyrate [623-42-7] M 102.1, b 102.3°/760mm, d 0.898, n 1.389. Treated with anhydrous CuSO₄, then distd under dry nitrogen.

***S*-(+)-2-Methylbutyric acid** [1730-91-2] M 102.1, b 64°/2mm, 78°/15mm, 90-94°/23mm, 174-175°/atm, d_4^{20} 0.938, n_D^{20} 1.406, $[\alpha]_{546}^{20} +23^\circ$, $[\alpha]_D^{20} +19.8^\circ$ (neat), $[\alpha]_D^{13} +18.3^\circ$ (c 6, EtOH), pK^{25} 4.76 (for *RS*). Purified by distn *in vacuo* [Sax and Bergmann *J Am Chem Soc* 77 1910 1955; Doering and Aschner *J Am Chem Soc* 75 393 1953]. The *methyl ester* is formed by addition of diazomethane and has b 112-115°/atm, $[\alpha]_D^{27} +21.1^\circ$ (c 1.7, MeOH).

Methyl carbamate [598-55-0] M 75.1, m 54.4-54.8°. Crystd from *benzene.

9-Methylcarbazole [1484-12-4] M 181.2, m 89°. Purified by zone melting.

4-Methylcatechol [452-86-8] M 124.1. See 3,4-dihydroxytoluene on p. 208.

Methyl chloride [74-87-3] M 50.5, b -24.1°. Bubbled through a sintered-glass disc dipping into conc H₂SO₄, then washed with water, condensed at low temperature and fractionally distd. Has been distd from AlCl₃ at -80°. Alternatively, passed through towers containing AlCl₃, soda-lime and P₂O₅, then condensed and fractionally distd. Stored as a gas.

Methyl chloroacetate [96-34-4] M 108.5, b 129-130°, d 1.230, n 1.423. Shaken with satd aq Na₂CO₃ (three times), aq 50% CaCl₂ (three times), satd aq NaCl (twice), dried (Na₂SO₄) and fractionally distd.

***R*-(+) Methyl 2-chloropropionate** [77287-29-7] M 122.6, b 49-50°/35mm, 78-80°/120mm, 132-134°/760mm, d_4^{20} 1.152, n_D^{20} 1.417, $[\alpha]_D^{20} +26^\circ$ (19.0°) (neat). Purified by repeated distillation [Walker *J Chem Soc* 67 916 1895; Walden *Chem Ber* 28 1293 1985; see also *Gless Synth Commun* 16 633 1986].

3-Methylcholanthrene [56-49-5] M 268.4, m 179-180°. Crystd from *benzene and diethyl ether. **CARCINOGEN.**

Methyl cyanoacetate [105-34-0] M 99.1, f -13°, b 205°, d 1.128, n 1.420. Purified by shaking with 10% Na₂CO₃ soln, washing well with water, drying with anhydrous Na₂SO₄, and distilling.

Methyl cyanofornate [17640-15-2] M 85.1, b 81°/47mm, 97°/751mm, 100-101°/760mm, d_4^{20} 1.072, n_D^{20} 1.37378. Purified by fractionation through a 45cm glass helices packed column and with a 30cm spinning band column. [Sheppard *J Org Chem* 27 3756 1962.] It has been distd through a short Vigreux

column, and further purified by recrystn from Et₂O at -40° as white crystals which melt at room temperature. NMR: δ 4.0 (CH₃), and IR: 2250 (CN) and 1750 (CO) cm⁻¹. [Childes and Weber *J Org Chem* **41** 3486 1976.]

Methylcyclohexane [108-87-2] M 98.2, b 100.9°, d²⁵ 0.7650, n 1.4231, n⁵² 1.42058. Passage through a column of activated silica gel gives material transparent down to 220nm. Also purified by passage through a column of activated basic alumina, or by azeotropic distn with MeOH, followed by washing out the MeOH with H₂O, drying and distilling. Methylcyclohexane can be dried with CaSO₄, CaH₂ or sodium. Has also been purified by shaking with a mixture of conc H₂SO₄ and HNO₃ in the cold, washing with H₂O, drying with CaSO₄ and fractionally distilling from potassium. Percolation through a Celite column impregnated with 2,4-dinitrophenylhydrazine (DNPH), phosphoric acid and H₂O (prepared by grinding 0.5g DNPH with 6mL 85% H₃PO₄, then mixing with 4mL of distilled H₂O and 10g of Celite) removes carbonyl-containing impurities.

2-Methylcyclohexanol [583-59-5] M 114.2, b 65°/20mm, 167.6°/760mm, d 0.922, n 1.46085. Dried with Na₂SO₄ and distd under vacuum.

cis- and trans-3-Methylcyclohexanol [591-23-1] M 114.2, b 69°/16mm, 172°/760mm, d 0.930, n 1.45757, n^{25.5} 1.45444. Dried with Na₂SO₄ and distd under vacuum.

4-Methylcyclohexanone [589-92-4] M 112.2, b 165.5°/743mm, d 0.914, n 1.44506. Dried with CaSO₄, then fractionally distd.

1-Methylcyclohexene [591-49-1] M 96.2, b 107.4-108°/atm, 110-111°/760mm, d 0.813, n 1.451. Freed from hydroperoxides by passing through a column containing basic alumina or refluxing with cupric stearate, filtered and fractionally distd from sodium.

Methylcyclopentane [96-37-7] M 84.2, b 71.8°, d 0.749, n 1.40970, n²⁵ 1.40700. Purification procedures include passage through columns of silica gel (prepared by heating in nitrogen to 350° prior to use) and activated basic alumina, distn from sodium-potassium alloy, and azeotropic distn with MeOH, followed by washing out the methanol with water, drying and distilling. It can be stored with CaH₂ or sodium.

3'-Methyl-1,2-cyclopentenophenanthrene [549-38-2] M 232.3, m 126-127°. Crystd from AcOH.

S-Methyl-L-cysteine [1187-84-4] M 135.2, m 207-211°, [α]_D²⁶ -32.0° (c 5, H₂O), pK₁²⁵ 1.94 (COOH), pK₂²⁵ 8.73 (NH₂, 8.97). Likely impurities are cysteine and S-methyl-dl-cysteine. Crystd from water by adding 4 volumes of EtOH.

5-Methylcytosine [4-amino-5-methylpyrimidin-2(1H)-one] [554-01-8] M 125.1, m 270°(dec), pK₁ 4.6, pK₂ 12.4. Crystd from water (sol 3.4%).

Methyl decanoate [110-42-9] M 186.3, b 114°/15mm, 224°/760mm, d 0.874, n 1.426. Passed through alumina before use.

Methyl 2,4-dichlorophenoxyacetate [1928-38-7] M 235.1, m 43°, b 119°/11mm. Crystd from MeOH.

m-Methyl-N,N-dimethylaniline [121-72-2] M 135.2, b 72-74°/5mm, 215°/760mm, pK²⁵ 5.22. Refluxed for 3h with 2 molar equivalents of acetic anhydride, then fractionally distd under reduced pressure. Also dried over BaO, distd and stored over KOH. Methods described for N,N-dimethylaniline are applicable.

p-Methyl-N,N-dimethylaniline [99-97-8] M 135.2, b 76.5-77.5°/4mm, 211°/760mm, pK²⁵ 4.76. Refluxed for 3h with 2 molar equivalents of acetic anhydride, then fractionally distd under reduced pressure. Also dried over BaO, distd and stored over KOH. Methods described for N,N-dimethylaniline are applicable.

2-Methyl-1,3-dithiane [6007-26-7] M 134.3, b 53-54°/1.1mm, 66°/5mm, 79-80°/8-10mm, 85°/12mm, d_4^{20} 1.121, n_D^{20} 1.560. Wash with H₂O, 2.5 M aqueous NaOH, H₂O, brine, dried over K₂CO₃ (use toluene as solvent if volume of reagent is small), filter, evaporate and distil the colourless residue. IR film: 1455, 1371 and 1060 (all medium and CH₃), 1451m, 1422s, 1412m, 1275m, 1236m, 1190m, 1171w, 918m and 866w (all dithiane) cm⁻¹ [Corey and Erickson *J Org Chem* 36 3553 1971; Seebach and Corey *J Org Chem* 40 231 1975].

Methyl dodecanoate [111-82-0] M 214.4, m 5°, b 141°/15mm, d 0.870, n^{50} 1.4199. Passed through alumina before use.

N-Methyleneaminoacetonitrile [109-82-0] M 68.1, m 129°. Crystd from EtOH or acetone.

p,p'-Methylene-bis-(N,N-dimethylaniline) [101-61-1] M 254.4, m 89.5°. See p,p'-tetramethyldiaminodiphenylmethane on p. 364.

Methylene Blue [3,7-bis-(dimethylamino)phenothiazin-5-ium chloride [61-73-4] M 319.9, CI 52015, ϵ_{654} 94,000 (EtOH), ϵ_{664} 81,000 (H₂O), pK²⁵ 3.8. Crystd from 0.1M HCl (16mL/g), the crystals were separated by centrifugation, washed with chilled EtOH and diethyl ether and dried under vacuum. Crystd from 50% aqueous EtOH, washed with absolute EtOH, and dried at 50-55° for 24h. Also crystd from *benzene-MeOH (3:1). Salted out with NaCl from a commercial conc aqueous soln, then crystd from water, dried at 100° in an oven for 8-10h.

3,4-Methylenedioxyaniline [14268-66-7] M 137.1, m 45-46°, b 144°/14mm, pK_{Est} ~3.8. Crystd from pet ether.

3,4-Methylenedioxcinnamic acid [2373-80-0] M 192.2, m 243-244°(dec), pK_{Est} ~4.6. Crystd from glacial acetic acid.

5,5'-Methylenedisalicylic acid [122-25-8] M 372.3, m 238°(dec). Crystd from acetone and *benzene.

Methylene Green [3,7-bis-(dimethylamino)-4-nitrophenothiazin-5-ium chloride] [2679-01-8] M 364.9, m >200°(dec), CI 52020, pK²⁵ 3.2. Crystd three times from water (18mL/g).

N-Methylephedrine (2-dimethylamino-1-phenylpropanol) [1S,2R-(+)- 42151-56-4; 1R,2S-(-)- 552-79-4] M 179.3, m 85-86°, 87-87.5°, 90°, b 115°/2mm, $[\alpha]_{546}^{20}$ (+) and (-) 35°, $[\alpha]_D^{20}$ (+) and (-) 30° (c 4.5, MeOH), pK²⁶ 9.22. It has been recrystd from Et₂O, pet ether, of aq EtOH or aq MeOH and has been distilled under reduced pressure. [Smith *J Chem Soc* 2056 1927; Tanaka and Sugawa *Yakugaku Zasshi (J Pharm Soc Japan)* 72 1548 1952 (*Chem Abstr* 47 8682 1953); Takamatsu *Yakugaku Zasshi (J Pharm Soc Japan)* 76 1227 1956, *Chem Abstr* 51 4304 1957.] The hydrochloride has m 192-193° and $[\alpha]_D^{20}$ +30° (c 5, H₂O) [Prelog and Hüfliger *Helv Chim Acta* 33 2021 1950].

Methyl ether (dimethyl ether) [115-10-6] M 46.1, b -63.5°/96.5mm. Dried by passing over alumina and then BaO, or over CaH₂, followed by fractional distn at low temperatures.

N-Methyl ethylamine hydrochloride [624-60-2] M 95.6, m 126-130°, pK 10.9 (free base). Crystd from absolute EtOH or diethyl ether.

N-Methyl formamide [123-39-7] M 59.1, m -3.5°, b 100.5°/25mm, d 1.005., n^{52} 1.4306. Dried with molecular sieves for 2days, then distd under reduced pressure through a column packed with glass helices. Fractionally crystd by partial freezing and the solid portion was vac distd.

Methyl formate [107-31-3] M 60.1, b 31.5°, d 0.971, n^{15} 1.34648, n 1.34332. Washed with strong aq Na₂CO₃, dried with solid Na₂CO₃ and distd from P₂O₅. (Procedure removes free alcohol or acid.)

2-Methylfuran [534-22-5] M 82.1, b 62.7-62.8°/731mm, d 0.917, n 1.436. Washed with acidified satd ferrous sulfate soln (to remove peroxides), separated, dried with CaSO₄ or CaCl₂, and fractionally distd from KOH immediately before use. To reduce the possibility of spontaneous polymerisation, addition of about one-third of its volume of heavy mineral oil to 2-methylfuran prior to distn has been recommended.

Methyl gallate [99-24-1] M 184.2, m 202°. Crystd from MeOH.

N-Methylglucamine [6284-40-8] M 195.2, m 128-129°, [α]_D²⁰ -19.5° (c 2, H₂O), pK²⁸ 9.62. Crystd from MeOH.

Methyl α -D-glucosamine [97-30-3] M 194.2, m 165°, [α]_D²⁵ +157.8° (c 3.0, H₂O), pK³⁰ 7.1. Crystd from MeOH.

α -Methylglutaric acid [18069-17-5] M 146.1, m 79°, pK₁²⁵ 4.36, pK₂²⁵ 5.37. Crystd from distd water, then dried under vacuum over conc H₂SO₄.

β -Methylglutaric acid [626-51-7] M 146.1, m 87°, pK₁²⁵ 4.35, pK₂²⁵ 5.44. Crystd from distd water, then dried under vacuum over conc H₂SO₄.

Methylglyoxal [78-98-8] M 72.1, b ca 72°/760mm. Commercial 30% (w/v) aqueous soln was diluted to about 10% and distd twice, taking the fraction boiling below 50°/20mm Hg. (This treatment does not remove lactic acid).

Methyl Green [82-94-0, 7114-03-6 (ZnCl₂ salt)] M 458.5, m >200°(dec). Crystd from hot water.

1-Methylguanine [938-85-2] M 165.2, m >300°(dec), pK₁²⁰ 3.13, pK₂²⁰ 10.54. Crystd from 50% aqueous acetic acid.

7-Methylguanine [578-76-7] M 165.2, pK₁²⁰ 3.50, pK₂²⁰ 9.95. Crystd from water.

2-Methylhexane [591-76-4] M 100.2, b 90.1°, d 0.678, n 1.38485, n²⁵ 1.38227. Purified by azeotropic distn with MeOH, then washed with water (to remove the MeOH), dried over type 4A molecular sieves and distd.

3-Methylhexane [589-34-4] M 100.2, b 91.9°, d 0.687, n 1.38864, n²⁵ 1.38609. Purification as for 2-methylhexane.

Methyl hexanoate [106-70-7] M 130.2, b 52°/15mm, 150°/760mm, d 0.885, n 1.410. Passed through alumina before use.

Methylhydrazine [60-34-4] M 46.1, b 87°/745mm, d 0.876, n 1.436, pK³⁰ 7.87. Dried with BaO, then vacuum distd. Stored under nitrogen.

Methyl hydrazinocarboxylate [6294-89-9] M 90.1, m 70-73°. To remove impurities, the material was melted and pumped under vacuum until the vapours were spectroscopically pure [Caminati et al. *J Am Chem Soc* 108 4364 1986].

Methyl 4-hydroxybenzoate [99-76-3] M 152.2, m 127.5°, pK_{Est} ~9.3. Fractionally crystd from its melt, recrystd from *benzene, then from *benzene/MeOH and dried over CaCl₂ in a vacuum desiccator.

Methyl 3-hydroxy-2-naphthoate [883-99-8] M 202.2, m 73-74°, pK_{Est} ~9.0. Crystd from MeOH (charcoal) containing a little water.

N-Methylimidazole [616-47-7] M 82.1, b 81-84°/27mm, 197-198°/760mm, d 1.032, n 1.496, pK²⁵ 7.25. Dried with sodium metal and then distd. Stored at 0° under dry argon.

2-Methylimidazole [693-98-1] M 82.1, m 140-141°, b 267°/760mm, pK²⁵ 7.86. Recrystd from *benzene or pet ether.

4-Methylimidazole [822-36-6] M 82.1, m 47-48°, b 263°/760mm, pK²⁵ 7.61. Recrystd from *benzene or pet ether.

2-Methylindole [95-20-5] M 131.2, m 61°, pK²⁵ -0.28 (C-3 protonation, aq H₂SO₄). Crystd from *benzene. Purified by zone melting.

3-Methylindole (skatole) [83-34-1] M 131.2, m 95°, pK²⁵ -4.55 (C-3-protonation, aq H₂SO₄). Crystd from *benzene. Purified by zone melting.

Methyl iodide [74-88-4] M 141.9, b 42.8°, d 2.281, n 1.5315. Deteriorates rapidly with liberation of iodine if exposed to light. Usually purified by shaking with dilute aqueous Na₂S₂O₃ or NaHSO₃ until colourless, then washed with water, dilute aqueous Na₂CO₃, and more water, dried with CaCl₂ and distd. It is stored in a brown bottle away from sunlight in contact with a small amount of mercury, powdered silver or copper. (Prolonged exposure of mercury to methyl iodide forms methylmercuric iodide.) Methyl iodide can be dried further using CaSO₄ or P₂O₅. An alternative purification is by percolation through a column of silica gel or activated alumina, then distn. The soln can be degassed by using a repeated freeze-pump-thaw cycle.

O-Methylisourea hydrogen sulfate (2-methylpseudourea sulfate) [29427-58-5] M 172.2, m 114-118°, 119°. Recrystd from MeOH-Et₂O (327g of salt dissolved in 1L of MeOH and 2.5L of Et₂O is added) [Fearing and Fox *J Am Chem Soc* 76 4382 1954]. The picrate has m 192° [Odo et al. *J Org Chem* 23 1319 1958].

N-Methyl maleimide [930-88-1] M 111.1, m 94-96°. Crystd three times from diethyl ether.

Methylmalonic acid [516-05-2] M 118.1, m 135°(dec), pK₁²⁵ 3.05, pK₂²⁵ 5.76. Crystallises as the hydrate from water.

3-Methylmercaptoaniline [1783-81-9] M 139.2, b 101.5-102.5°/0.3mm, 163-165°/16mm, d₄²⁰ 1.147, n_D²⁰ 1.641, pK²⁵ 4.05. Purified by fractional distn in an inert atmosphere. It has UV max at 226 and 300. [Bordwell and Cooper *J Am Chem Soc* 74 1058 1952.] The *N*-acetyl derivative has m 78-78.5° (after recrystn from aq EtOH).

4-Methylmercaptoaniline [104-96-1] M 139.2, b 140°/15mm, 151°/25mm, 155°/23mm, d₄²⁰ 1.137, n_D²⁰ 1.639, pK²⁵ 4.40. Purified by fractional distn in an inert atmosphere. [Lumbroso and Passerini *Bull Soc Chim Fr* 311 1957; Mangini and Passerini *J Chem Soc* 4954 1956.]

Methyl methacrylate [80-62-6] M 100.1, f -50°, b 46°/100mm, d 0.937, n 1.4144. Washed twice with aqueous 5% NaOH (to remove inhibitors such as hydroquinone) and twice with water. Dried with CaCl₂, Na₂CO₃, Na₂SO₄ or MgSO₄, then with CaH₂ under nitrogen at reduced pressure. The distillate is stored at low temperatures and redistd before use. Prior to distn, inhibitors such as β-naphthylamine (0.2%) or di-β-naphthol are sometimes added. Also purified by boiling aqueous H₃PO₄ soln and finally with saturated NaCl soln. It was dried for 24h over anhydrous CaSO₄, distd at 0.1mm Hg at room temperature and stored at -30° [Albeck et al. *J Chem Soc, Faraday Trans 1* 1 1488 1978].

Methyl methanesulfonate [66-27-3] M 110.3, b 59°/0.6mm, 96-98°/19mm, d 1.300, n 1.4140. Purified by careful fractionation and collecting the middle fraction. Suspected CARCINOGEN. Note that MeSO₃H has b 134.5-135°, 167-167.5°/10mm and methanesulfonic anhydride has b 138°/10mm)—both are possible impurities.

Methyl methanethiolsulfonate [2949-92-0] M 126.2, b 69-71°/0.4mm, 96-97°/4.5mm, 104-105°/10mm, 119°/16mm, d 1.226, n 1.515. Purified by fractional distn under reduced pressure, IR: ν 1350, 750 cm^{-1} . [Applegate et al. *J Org Chem* 38 943 1973.]

α -Methylmethionine [562-48-1] M 163.0, m 283-284°, pK³⁰ 9.45. Crystd from aqueous EtOH.

S-Methyl-L-methionine chloride (Vitamin U) [1115-84-0] M 199.5, $[\alpha]_D^{23} +33^\circ$ (0.2M HCl), pK₁ 1.9, pK₂ 7.9. Likely impurities are methionine, methionine sulfoxide and methionine sulfone. Crystd from water by adding a large excess of EtOH. Stored in a cool, dry place, protected from light.

N-Methylmorpholine [109-02-4] M 101.2, b 116-117°/764mm, d 0.919, n 1.436, pK²⁵ 7.38. Dried by refluxing with BaO or sodium, then fractionally distd through a helices-packed column.

4-Methylmorpholine-4-oxide monohydrate [7529-22-8] M 135.2, m 71-73°. When dried for 2-3h at high vacuum it dehydrates. Add MeOH to the oxide and distil off the solvent under vacuum until the temp is ca 95°. Then add Me₂CO at reflux then cool to 20°. The crystals are filtered off washed with Me₂CO and dry. The degree of hydration may vary and may be important for the desired reactions. [van Rheen et al. *Tetrahedron Lett* 1973 1076; Schneider and Hanze *US Pat* 2 769 823; see also Sharpless et al. *Tetrahedron Lett* 2503 1976.]

1-Methylnaphthalene [90-12-0] M 142.2, f -30°, b 244.6°, d 1.021, n 1.6108. Dried for several days with CaCl₂ or by prolonged refluxing with BaO. Fractionally distd through a glass helices-packed column from sodium. Purified further by soln in MeOH and pptn of its picrate complex by adding to a saturated soln of picric acid in MeOH. The picrate, after crystn to constant melting point (m 140-141°) from MeOH, was dissolved in *benzene and extracted with aqueous 10% LiOH until the extract was colourless. Evaporation of the *benzene under vacuum gave 1-methylnaphthalene [Kloetzel and Herzog *J Am Chem Soc* 72 1991 1950]. However, neither the picrate nor the styphnate complexes satisfactorily separates 1- and 2-methylnaphthalenes. To achieve this, 2-methylnaphthalene (10.7g) in 95% EtOH (50mL) has been ppted with 1,3,5-trinitrobenzene (7.8g) and the complex has been crystd from MeOH to m 153-153.5° (m of the 2-methyl isomer is 124°). [Alternatively, 2,4,7-trinitrofluorenone in hot glacial acetic acid could be used, and the derivative (m 163-164°) recrystd from glacial acetic acid]. The 1-methylnaphthalene was regenerated by passing a soln of the complex in dry *benzene through a 15-in column of activated alumina and washing with *benzene/pet ether (b 35-60°) until the coloured band of the nitro compound had moved down near the end of the column. The complex can also be decomposed using tin and acetic-hydrochloric acids, followed by extraction with diethyl ether and *benzene; the extracts were washed successively with dilute HCl, strongly alkaline sodium hypophosphite, water, dilute HCl and water. [Soffer and Stewart *J Am Chem Soc* 74 567 1952.] It can be purified from anthracene by zone melting.

2-Methylnaphthalene [91-57-6] M 142.2, m 34.7-34.9°, b 129-130°/25mm. Fractionally crystd repeatedly from its melt, then fractionally distd under reduced pressure. Crystd from *benzene and dried under vacuum in an Abderhalden pistol. Purified *via* its picrate (m 114-115°) as described for 1-methylnaphthalene.

6-Methyl-2-naphthol [17579-79-2] M 158.2, m 128-129°, pK_{Est} ~9.8. Crystd from EtOH or ligroin. Sublimed *in vacuo*.

7-Methyl-2-naphthol [26593-50-0] M 158.2, m 118°, pK_{Est} ~9.7. Crystd from EtOH or ligroin. Sublimed *in vacuo*.

Methyl 1-naphthyl ether [2216-69-5] M 158.2, b 90-91°/2mm, d 1.095, n²⁶ 1.6210. Steam distd from alkali. The distillate was extracted with diethyl ether. After drying (MgSO₄) the extract and evaporating diethyl ether, the methyl naphthyl ether was then fractionated under reduced pressure from CaH₂.

Methyl nitrate [598-58-3] M 77.0, b 65°/760mm, d⁵ 1.2322, d¹⁵ 1.2167, d²⁵ 1.2032. Distd at -80°. The middle fraction was subjected to several freeze-pump-thaw cycles. **VAPOUR EXPLODES ON HEATING.**

Methyl nitrite [624-91-9] M 61.0, b -12° , d^{15} (liq) 0.991. Condensed in a liquid nitrogen trap. Distd under vacuum, first trap containing dry Na_2CO_3 to free it from acid impurities then into further Na_2CO_3 traps before collection.

N-Methyl-4-nitroaniline [100-15-2] M 152.2, m 152.2° , pK^{25} 0.55. Crystd from aqueous EtOH.

2-Methyl-5-nitroaniline [99-55-8] M 152.2, m 109° , pK^{25} 2.35. Acetylated, and the acetyl derivative crystd to constant melting point, then hydrolysed with 70% H_2SO_4 and the free base regenerated by treatment with ammonia [Bevan, Fayiga and Hirst *J Chem Soc* 4284 1956].

4-Methyl-3-nitroaniline [119-32-4] M 152.2, m 81.5° , pK^{25} 3.02. Crystd from hot water (charcoal), then ethanol and dried in a vacuum desiccator.

Methyl 3-nitrobenzoate [618-95-1] M 181.2, m 78° . Crystd from MeOH (1g/mL).

Methyl 4-nitrobenzoate [619-50-1] M 181.2, m $95-95.5^{\circ}$. Dissolved in diethyl ether, then washed with aqueous alkali, the ether was evaporated and the ester was recrystd from EtOH.

2-Methyl-2-nitro-1,3-propanediol [77-49-6] M 135.1, m 145° . Crystd from *n*-butanol.

2-Methyl-2-nitro-1-propanol [76-39-1] M 119.1, m $87-88^{\circ}$. Crystd from pet ether.

N-Methyl-4-nitrosoaniline [10595-51-4] M 136.2, m 118° . Crystd from *benzene.

N-Methyl-N-nitroso-*p*-toluenesulfonamide (diazald) [80-11-5] M 214.2, m 62° . Crystd from *benzene by addition of pet ether, store in a refrigerator.

Methylnorbornene-2,3-dicarboxylic anhydride (5-methylnorborn-5-ene-2-endo-3-endo-dicarboxylic anhydride) [25134-21-8] M 178.2, m $88.5-89^{\circ}$. Purified by thin layer chromatography on Al_2O_3 (previously boiled in EtOAc) and eluted with hexane-* C_6H_6 (1:2) then recrystd from * C_6H_6 -hexane. The free acid has m $118.5-119.5^{\circ}$. [Miranov et al. *Tetrahedron* 19 1939 1963.]

3-Methyloctane [2216-33-3] M 128.3, b $142-144^{\circ}/760\text{mm}$, d 0.719, n 1.407. Passed through a column of silica gel [Klassen and Ross *J Phys Chem* 91 3668 1987].

Methyl octanoate (methyl caprylate) [111-11-5] M 158.2, b $83^{\circ}/15\text{mm}$, $193-194^{\circ}/760\text{mm}$, d 0.877, n 1.419. Passed through alumina before use.

Methyl oleate [112-62-9] M 296.5, f -19.9° , b $217^{\circ}/16\text{mm}$, d 0.874, n 1.4522. Purified by fractional distn under reduced pressure, and by low temperature crystn from acetone.

3-Methyl-2-oxazolidone [19836-78-3] M 101.1, m 15° , b $88-91^{\circ}/1\text{mm}$, d 1.172, n 1.455. Purified by successive fractional freezing, then dried in a dry-box over 4A molecular sieves for 2 days.

3-Methyl-3-oxetanemethanol (3-hydroxymethyl-3-methyloxetane) [3143-02-0] M 102.1, b $80^{\circ}/4\text{mm}$, $92-93^{\circ}/12\text{mm}$, d_4^{20} 1.033, n_D^{25} 1.4449. Purified by fractionation through a glass column [Pattison *J Am Chem Soc* 79 3455 1957].

Methylpentane (mixture of isomers). Passage through a long column of activated silica gel (or alumina) gave material transparent down to 200nm by UV.

2-Methylpentane [107-83-5] M 86.2, b 60.3° , d 0.655, n 1.37145, n_D^{25} 1.36873. Purified by azeotropic distn with MeOH, followed by washing out the MeOH with water, drying (CaCl_2 , then sodium), and distn. [Forziati et al. *J Res Nat Bur Stand* 36 129 1946.]

3-Methylpentane [96-14-0] M 86.2, b 63.3°, d 0.664, n 1.37652, n²⁵ 1.37384. Purified by azeotropic distn with MeOH, as for 2-methylpentane. Purified for ultraviolet spectroscopy by passage through columns of silica gel or alumina activated by heating for 8h at 210° under a stream of nitrogen. Has also been treated with conc (or fuming) H₂SO₄, then washed with water, aqueous 5% NaOH, water again, then dried (CaCl₂, then sodium), and distd through a long, glass helices-packed column.

2-Methyl-2,4-pentanediol [107-41-5] M 118.2, b 107.5-108.5°/25mm, d 0.922, n²⁵ 1.4265. Dried with Na₂SO₄, then CaH₂ and fractionally distd under reduced pressure through a packed column, taking precautions to avoid absorption of water.

2-Methyl-1-pentanol [105-30-6] M 102.2, b 65-66°/60mm, 146-147°/760mm, d 0.827, n 1.420. Dried with Na₂SO₄ and distd.

4-Methyl-2-pentanol [108-11-2] M 102.2, b 131-132°, d 0.810, n 1.413. Washed with aqueous NaHCO₃, dried and distd. Further purified by conversion to the phthalate ester by adding 120mL of dry pyridine and 67g of phthalic anhydride per mole of alcohol, purifying the ester and steam distilling it in the presence of NaOH. The distillate was extracted with ether, and the extract was dried and fractionally distd. [Levine and Walti *J Biol Chem* 94 367 1931].

3-Methyl-3-pentanol carbamate (Emylcamate) [78-28-4] M 145.2, m 56-58.5°. Crystd from 30% EtOH.

4-Methyl-2-pentanone (methyl isobutyl ketone) [108-10-1] M 100.2, b 115.7°, d 0.801, n 1.3958, n²⁵ 1.3938. Refluxed with a little KMnO₄, washed with aqueous NaHCO₃, dried with CaSO₄ and distd. Acidic impurities were removed by passage through a small column of activated alumina.

2-Methyl-1-pentene [763-29-1] M 84.2, b 61.5-62°, d 0.680, n 1.395. Water was removed, and peroxide formation prevented by several vacuum distns from sodium, followed by storage with sodium-potassium alloy.

cis-4-Methyl-2-pentene [691-38-3] M 84.2, m -134.4°, b 57.7-58.5°, d 0.672, n 1.388. Dried with CaH₂, and distd.

trans-4-Methyl-2-pentene [674-76-0] M 84.2, m -140.8°, b 58.5°, d 0.669, n 1.389. Dried with CaH₂, and distd.

5-Methyl-1,10-phenanthroline [3002-78-6] M 194.2, m 113°(anhydr), pK²⁵ 5.28. Crystd from *benzene/pet ether.

N-Methylphenazonium methosulfate see 5-methylphenazinium methyl sulfate on p. 547 in Chapter 6.

N-Methylphenothiazine [1207-72-3] M 213.2, α-form m 99.3° and b 360-365°, β-form m 78-79°. Recrystn (three times) from EtOH gave α-form (prisms). Recrystn from EtOH/*benzene gave the β-form (needles). Also purified by vacuum sublimation and carefully dried in a vacuum line. Also crystd from toluene and stored in the dark [Guarr et al. *J Am Chem Soc* 107 5104 1985; Olmsted et al. *J Am Chem Soc* 109 3297 1987.]

4-Methylphenylacetic acid [622-47-9] M 150.2, m 94°, pK²⁵ 4.37. Crystd from heptane or water.

1-Methyl-1-phenylhydrazine sulfate [33008-18-3] M 218.2, pK²⁵ 4.98 (free base). Crystd from hot H₂O by addition of hot EtOH.

3-Methyl-1-phenyl-5-pyrazolone [89-25-8] M 174.2, m 127°. Crystd from hot H₂O, or EtOH/water (1:1).

N-Methylphthalimide [550-44-7] M 161.1, m 133.8°. Recrystd from absolute EtOH.

2-Methylpiperazine [109-07-9] M 100.2, m 61-62°, 66°, b 147-150°/739mm, pK_1^{25} 5.46, pK_2^{25} 9.90. Purified by zone melting and by distn.

3-Methylpiperidine [626-56-2] M 99.2, b 125°/763mm, d 0.846, n^{25} 1.4448, pK^{25} 11.07. Purified via the *hydrochloride* (m 172°). [Chapman, Isaacs and Parker *J Chem Soc* 1925 1959.]

4-Methylpiperidine [626-58-4] M 99.2, b 124.4°/755mm, d 0.839, n^{25} 1.4430, pK^{25} 10.78. Purified via the *hydrochloride* (m 189°). Freed from 3-methylpyridine by zone melting.

1-Methyl-4-piperidone [1445-73-4] M 113.2, b 53-56°/0.5mm, 54-56°/9mm, 68-71°/17mm, 85-87°/45mm, d_4^{20} 0.972, n_D^{25} 1.4588, pK^{25} 7.9. It is best purified by fractional distn. The *hydrochloride* of the hydrate (4-diol) has m 94.7-95.5°, but the anhydrous *hydrochloride* which crystallises from $CHCl_3$ -Et₂O and has m 165-168° (164-167°) and can also be obtained by sublimation at 120°/2mm. The *oxime* has m 130-132° (from Me₂CO). The *methiodide* crystallises from MeOH and the crystals with 1MeOH has m 189-190°, and the solvent-free *iodide* has m 202-204° dec. [Lyle et al. *J Org Chem* 24 342 1959; Bowden and Green *J Chem Soc* 1164 1952; Tomita *Yakugaku Zasshi (J Pharm Soc Japan)* 71 1053 1951.]

2-Methylpropane-1,2-diamine (1,2-diamino-2-methylpropane) [811-93-8] M 88.2, b 47-48°/17mm, pK_1^{25} 6.25 (6.18), pK_2^{25} 9.82 (9.42). Dried with sodium for 2 days, then distd under reduced pressure from sodium.

2-Methylpropane-1-thiol [513-44-0] M 90.2, b 41.2°/142mm, n^{25} 1.43582, pK_{Est} ~10.8. Dissolved in EtOH, and added to 0.25M Pb(OAc)₂ in 50% aqueous EtOH. The ppted lead mercaptide was filtered off, washed with a little EtOH, and impurities were removed from the molten salt by steam distn. After cooling, dilute HCl was added dropwise to the residue, and the mercaptan was distd directly from the flask. Water was separated from the distillate, and the mercaptan was dried (Na₂CO₃) and distd under nitrogen. [Mathias *J Am Chem Soc* 72 1897 1950.]

2-Methylpropane-2-thiol [75-66-1] M 90.2, b 61.6°/701mm, d^{25} 0.79426, n^{25} 1.41984, pK^{25} 11.22. Dried for several days with CaO, then distd from CaO. Purified as for *2-methylpropane-1-thiol*.

2-Methyl-1-propanol (isobutanol) [78-83-1] M 74.1, b 107.9°, d 0.804, n^{15} 1.39768, n^{25} 1.3939. Dried by refluxing with CaO and BaO for several hours, followed by treatment with calcium or aluminium amalgam, then fractional distn from sulfanilic or tartaric acids. More exhaustive purifications involve formation of phthalate or borate esters. Heating with phthalic anhydride gives the *acid phthalate* which, after crystn to constant melting point (m 65°) from pet ether, is hydrolysed with aqueous 15% KOH. The alcohol is distd as the water azeotrope and dried with K₂CO₃, then anhydrous CuSO₄, and finally magnesium turnings, followed by fractional distn. [Hückel and Ackermann *J Prakt Chem* 136 15 1933.] The borate ester is formed by heating the dried alcohol for 6h in an autoclave at 160-175° with a quarter of its weight of boric acid. After fractional distns under vac the ester is hydrolysed by heating for a short time with aq alkali and the alcohol is dried with CaO and distd. [Michael, Scharf and Voigt *J Am Chem Soc* 38 653 1916.] (see p. 271).

Methyl propiolate [922-67-8] M 84.1, b 100°/atm, 102°/atm, 103-105°/atm, d 0.945, n 1.4080. Purified by fractional distn and collecting the middle fraction; note that propiolic acid has a high b [144°(dec)/atm]. LACHRYMATORY.

N-Methylpropionamide [1187-58-2] M 87.1, f -30.9°, b 103°/12-13mm, d 0.934, n^{25} 1.4356. A colourless, odourless, neutral liquid at room temperature with a high dielectric constant. The amount of water present can be determined directly by Karl Fischer titration; GLC and NMR have been used to detect unreacted propionic acid. Commercial material of high quality is available, probably from the condensation of anhydrous methylamine with 50% excess of propionic acid. Rapid heating to 120-140° with stirring favours the reaction by removing water either directly or as the ternary xylene azeotrope. The quality of the distillate improves during the distn.

The propionamide can be dried over CaO. H₂O and unreacted propionic acid were removed as their xylene azeotropes. It was vacuum dried. Material used as an electrolyte solvent (specific conductance less than 10⁻⁶ ohm⁻¹ cm⁻¹) was obtained by fractional distn under reduced pressure, and stored over BaO or molecular sieves because it readily absorbs moisture from the atmosphere on prolonged storage. [Hoover *Pure Appl Chem* 37 581 1974; *Recommended Methods for Purification of Solvents and Tests for Impurities*, Coetzee Ed., Pergamon Press, 1982.]

Methyl propionate [554-12-1] M 88.1, b 79.7°. Washed with satd aq NaCl, then dried with Na₂CO₃ and distd from P₂O₅. (This removes any free acid and alcohol.) It has also been dried with anhydrous CuSO₄.

Methyl n-propyl ether [557-17-5] M 74.1, b 39°, d 0.736, n¹⁴ 1.3602, pK²⁵ -3.79 (aq H₂SO₄). Dried with CaSO₄, then passed through a column of alumina (to remove peroxides) and fractionally distd.

Methyl n-propyl ketone [107-87-9] M 86.1, b 102.4°, d 0.807, n 1.3903. Refluxed with a little KMnO₄, dried with CaSO₄ and distd. It was converted to its bisulfite addition compound by shaking with excess saturated aqueous NaHSO₃ at room temperature, cooling to 0°, filtering, washing with diethyl ether and drying. Steam distillation gave a distillate from which the ketone was recovered, washed with aq NaHCO₃ and distd water, dried (K₂CO₃) and fractionally distd. [Waring and Garik *J Am Chem Soc* 78 5198 1956.]

3-Methyl-1-propyn-3-ol carbamate [302-66-9] M 141.2, m 55.8-57°. Crystd from ether/pet ether or cyclohexane.

2-Methylpyrazine [109-08-0] M 94.1, b 136-137°, d 1.025, n 1.505, pK₁²⁵ -5.25 (aq H₂SO₄), pK₂²⁵ 1.47. Purified *via* the picrate. [Wiggins and Wise *J Chem Soc* 4780 1956.]

2-Methylpyridine (2-picoline) [109-06-8] M 93.1, b 129.4°, d 0.9444, n 1.50102, pK²⁵ 5.96. Biddiscombe and Handley [*J Chem Soc* 1957 1954] steam distd a boiling soln of the base in 1.2 equivalents of 20% H₂SO₄ until about 10% of the base had been carried over, along with non-basic impurities. Excess aqueous NaOH was then added to the residue, the free base was separated, dried with solid NaOH and fractionally distd.

2-Methylpyridine can also be dried with BaO, CaO, CaH₂, LiAlH₄, sodium or Linde type 5A molecular sieves. An alternative purification is *via* the ZnCl₂ adduct, which is formed by adding 2-methylpyridine (90mL) to a soln of anhydrous ZnCl₂ (168g) and 42mL conc HCl in absolute EtOH (200mL). Crystals of the complex are filtered off, recrystd twice from absolute EtOH (to give m 118.5-119.5°), and the free base is liberated by addition of excess aqueous NaOH. It is steam distd, and solid NaOH added to the distillate to form two layers, the upper one of which is then dried with KOH pellets, stored for several days with BaO and fractionally distd. Instead of ZnCl₂, HgCl₂ (430g in 2.4L of hot water) can be used. The complex, which separates on cooling, can be dried at 110° and recrystd from 1% HCl (to m 156-157°).

3-Methylpyridine (3-picoline) [108-99-6] M 93.1, m -18.5°, b 144°/767mm, d 0.957, n 1.5069, pK²⁵ 5.70. In general, the same methods of purification that are described for 2-methylpyridine can be used. However, 3-methylpyridine often contains 4-methylpyridine and 2,6-lutidine, neither of which can be removed satisfactorily by drying and fractionation, or by using the ZnCl₂ complex. Biddiscombe and Handley [*J Chem Soc* 1957 1954], after steam distn as for 2-methylpyridine, treated the residue with urea to remove 2,6-lutidine, then azeotropically distd with acetic acid (the azeotrope had b 114.5°/712mm), and recovered the base by adding excess of aqueous 30% NaOH, drying with solid NaOH and carefully fractionally distilling. The distillate was then fractionally crystd by slow partial freezing. An alternative treatment [Reithof et al. *Ind Eng Chem (Anal Edn)* 18 458 1946] is to reflux the crude base (500mL) for 20-24h with a mixture of acetic anhydride (125g) and phthalic anhydride (125g) followed by distn until phthalic anhydride begins to pass over. The distillate was treated with NaOH (250g in 1.5L of water) and then steam distd. Addition of solid NaOH (250g) to this distillate (*ca* 2L) led to the separation of 3-methylpyridine which was removed, dried (K₂CO₃, then BaO) and fractionally distd. (Subsequent fractional freezing would probably be advantageous.)

4-Methylpyridine (4-picoline) [108-89-4] M 93.1, m 4.25°, b 145.0°/765mm, d 0.955, n 1.5058, pK²⁵ 4.99. Can be purified as for 2-methylpyridine. Biddescombe and Handley's method for 3-methylpyridine is also applicable. Lidstone [*J Chem Soc* 242 1940] purified via the oxalate (m 137-138°) by heating 100mL of 4-methylpyridine to 80° and adding slowly 110g of anhydrous oxalic acid, followed by 150mL of boiling EtOH. After cooling and filtering, the ppte was washed with a little EtOH, then recrystd from EtOH, dissolved in the minimum quantity of water and distd with excess 50% KOH. The distillate was dried with solid KOH and again distd. Hydrocarbons can be removed from 4-methylpyridine by converting the latter to its hydrochloride, crystallising from EtOH/diethyl ether, regenerating the free base by adding alkali and distilling. As a final purification step, 4-methylpyridine can be fractionally crystd by partial freezing to effect a separation from 3-methylpyridine. Contamination by 2,6-lutidine is detected by its strong absorption at 270nm.

4-Methylpyridine 1-oxide [1003-67-4] M 109.1, m 182-184°. See 4-picoline-*N*-oxide on p. 335.

***N*-Methylpyrrole** [96-54-8] M 81.1, b 115-116°/756mm, d 0.908, n 1.487, pK -3.4 (-2.90). Dried with CaSO₄, then fractionally distd from KOH immediately before use.

1-Methyl-2-pyrrolidinone [872-50-4] M 99.1, f -24.4, b 65-76°/1mm, 78-79°/12mm, 94-96°/20mm, 202°/760mm, d₄²⁰ 1.0328, n_D²⁰ 1.4678, pK -0.17 (also -0.92, and 0.2). Dried by removing water as *benzene azeotrope. Fractionally distd at 10 torr through a 100-cm column packed with glass helices. [Adelman *J Org Chem* 29 1837 1964; McElvain and Voza *J Am Chem Soc* 71 896 1949.] The hydrochloride has m 86-88° (from EtOH or Me₂CO-EtOH) [Reppe et al. *Justus Liebigs Ann Chem* 596 1 1955].

2-Methylquinoline (quinaldine) [91-63-4] M 143.2, b 86-87°/1mm, 155°/14mm, 246-247°/760mm, d 1.058, n 1.6126, pK²⁵ 5.65. Dried with Na₂SO₄ or by refluxing with BaO, then fractionally distd under reduced pressure. Redistd from zinc dust. Purified by conversion to its phosphate (m 220°) or picrate (m 192°) from which after recrystn, the free base was regenerated. [Packer, Vaughan and Wong *J Am Chem Soc* 80 905 1958.] Its ZnCl₂ complex can be used for the same purpose.

4-Methylquinoline (lepidine) [491-35-0] M 143.2, b 265.5°, d 1.084, n 1.61995, pK²⁵ 5.59. Refluxed with BaO, then fractionally distd. Purified via its recrystd dichromate salt (m 138°). [Cumper, Redford and Vogel *J Chem Soc* 1176 1962.]

6-Methylquinoline [91-62-3] M 143.2, b 258.6°, d 1.067, n 1.61606, pK²⁵ 4.92. Refluxed with BaO, then fractionally distd. Purified via its recrystd ZnCl₂ complex (m 190°). [Cumper, Redford and Vogel *J Chem Soc* 1176 1962.]

7-Methylquinoline [612-60-2] M 143.2, m 38°, b 255-260°, d 1.052, n 1.61481, pK²⁵ 5.29. Purified via its dichromate complex (m 149°, after five recrystns from water). [Cumper, Redford and Vogel *J Chem Soc* 1176 1962.]

8-Methylquinoline [611-32-5] M 143.2, b 122.5°/16mm, 247.8°/760mm, d 1.703, n 1.61631, pK²⁵ 4.60. Purified as for 2-methylquinoline. The phosphate and picrate have m 158° and m 201° respectively.

Methyl Red (4-dimethylaminoazobenzene-2'-carboxylic acid) [493-52-7] M 269.3, m 181-182°, CI 13020, pK₁²⁵ 2.30, pK₂²⁵ 4.82. The acid is extracted with boiling toluene using a Soxhlet apparatus. The crystals which separated on slow cooling to room temperature are filtered off, washed with a little toluene and recrystd from glacial acetic acid, *benzene or toluene followed by pyridine/water. Alternatively, dissolved in aq 5% NaHCO₃ soln, and pted from hot soln by dropwise addition of aq HCl. Repeated until the extinction coefficients did not increase.

Methyl salicylate (methyl 2-hydroxybenzoate) [119-36-8] M 152.2, m -8.6°, b 79°/6mm, 104-105°/14mm, 223.3°/atm, d₄²⁰ 1.1149, n_D²⁰ 1.5380, pK²⁵ 10.19. Dilute with Et₂O, wash with satd NaHCO₃ (it may effervesce due to the presence of free acid), brine, dry MgSO₄, filter, evaporate and distil.

Its solubility is 1g/1.5L of H₂O. The *benzoyl* derivative has *m* 92° (b 270-280°/120mm), and the 3,5-*dinitrobenzoate* has *m* 107.5°, and the 3,5-*dinitrocarbamoyl* derivative has *m* 180-181°. [Hallas *J Chem Soc* 5770 1965.]

Methyl stearate [122-61-8] *M* 298.5, *m* 41-43°, *b* 181-182°/4mm. Crystd from pet ether or distd.

α-Methylstyrene (monomer) [98-83-9] *M* 118.2, *b* 57°/15mm, *d* 0.910, *n* 1.5368. Washed three times with aqueous 10% NaOH (to remove inhibitors such as quinol), then six times with distd water, dried with CaCl₂ and distd under vacuum. The distillate is kept under nitrogen, in the cold, and redistd if kept for more than 48h before use. It can also be dried with CaH₂.

trans-β-Methylstyrene [873-66-5] *M* 118.2, *b* 176°/760mm, *d* 0.910, *n* 1.5496. Distd under nitrogen from powdered NaOH through a Vigreux column, and passed through activated neutral alumina before use [Wong et al. *J Am Chem Soc* 109 3428 1987].

4-Methylstyrene [622-97-9] *M* 118.2, *b* 60°/12mm, 106°/10mm, *d*₄²⁰ 0.9173, *n*_D²⁰ 1.542. Purified as the above styrenes and add a small amount of antioxidant if it is to be stored, UV in EtOH λ_{max} 285nm (log ε 3.07), and in EtOH + HCl 295nm (log ε 2.84) and 252nm (log ε 4.23). [Schwartzman and Carson *J Am Chem Soc* 78 322 1956; Joy and Orchin *J Am Chem Soc* 81 305 1959; Buck et al. *J Chem Soc* 23771949.]

Methylsuccinic acid [498-21-5] *M* 132.1, *m* 115.0°, *pK*₁²⁵ 3.88, *pK*₂²⁵ 5.35. Crystd from water.

(±)-3-Methylsulfolane (3-methyl-tetrahydrothiophene-1,1-dioxide) [872-93-5] *M* 134.2, *m* 0.5°, *b* 101°/2mm, 125-130°/12mm, 278-282°/763.5mm, *d*₄²⁰ 1.1885, *n*_D²⁰ 1.4770. Distil under vacuum and recryst from Et₂O at -60° to -70°. IR film has strong bands at 570 and 500 cm⁻¹. [Eigenberger *J Prakt Chem* [2] 131 289 1931; Freaheller and Katon *Spectrochim Acta* 20 10991964.]

17α-Methyltestosterone [58-18-4] *M* 302.5, *m* 164-165°, [α]₅₄₆²⁰ +87° (c 1, dioxane). Crystd from hexane/*benzene.

Methyl tetradecanoate (methyl myristate) [124-10-7] *M* 382.7, *m* 18.5°, *b* 155-157°/7mm. Passed through alumina before use.

2-Methyltetrahydrofuran [96-47-9] *M* 86.1, *b* 80.0°, *d*₄²⁰ 0.856, *n*_D²⁰ 1.4053. Likely impurities are 2-methylfuran, methyl-dihydrofurans and hydroquinone (stabiliser, which is removed by distn under reduced pressures). It was washed with 10% aqueous NaOH, dried, vacuum distd from CaH₂, passed through freshly activated alumina under nitrogen, and refluxed over sodium metal under vacuum. Stored over sodium. [Ling and Kevan *J Phys Chem* 80 592 1976.] Vacuum distd from sodium, and stored with sodium-potassium alloy. (Treatment removes water and prevents the formation of peroxides.) Alternatively, it can be freed from peroxides by treatment with ferrous sulfate and sodium bisulfate, then solid KOH, followed by drying with, and distilling from, sodium, or type 4A molecular sieves under argon. It may be difficult to remove *benzene if it is present as an impurity (can be readily detected by its ultraviolet absorption in the 249-268nm region). [Ichikawa and Yoshida *J Phys Chem* 88 3199 1984.] It has also been purified by percolating through Al₂O₃ and fractionated collecting fraction *b* 79.5-80°. After degassing, the material was distd onto degassed molecular sieves, then distd onto anthracene and a sodium mirror. The solvent was distd from the green soln onto potassium mirror or sodium-potassium alloy, from which it was distilled again. [Mohammad and Kosower *J Am Chem Soc* 93 2713 1971.] It should be stored in the presence of 0.1% of hydroquinone as stabiliser. **HARMFUL VAPOURS.**

N-Methylthioacetamide [5310-10-1] *M* 89.1, *m* 59°. Recrystd from *benzene.

3-Methylthiophene [616-44-4] *M* 98.2, *b* 111-113°, *d* 1.024, *n* 1.531. Dried with Na₂SO₄, then distd from sodium.

6(4)-Methyl-2-thiouracil [56-04-2] M 142.2, m 330°(dec), 299-303° (dec), 323-324° (dec), pK 8.1. Crystd from a large volume of H₂O. Purified by dissolving in base adding charcoal, filtering and acidifying with AcOH. Suspend the wet solid (ca 100g) in boiling H₂O (1L), stir and add AcOH (20mL), stir and refrigerate. Collect the product, wash with cold H₂O (4 x 200mL), drain for several hours then place in an oven at 70° to constant weight. [IR: Short and Thompson *J Chem Soc* 168 1952; Foster and Snyder *Org Synth Coll Vol IV* 638 1063.]

Methyl 4-toluenesulfonate [80-48-8] M 186.2, m 25-28°, 28°, b 144.6-145.2°/5mm, 168-170°/13mm, d₄²⁰ 1.23. It is purified by distn *in vacuo* and could be crystd from pet ether or Et₂O-pet ether at low temperature. It is a powerful methylating agent and is **TOXIC** and a **skin irritant**, so it is better to purify by repeated distn. [IR: Schreiber *Anal Chem* 21 1168 1949; Buehler et al. *J Org Chem* 2 167 1937; Roos et al. *Org Synth Coll Vol I* 145 1948.]

4-Methyl-1,2,4-triazoline-3,5-dione (MTAD) [13274-43-6] M 113.1, m 103-104°, m 107-109°. Obtained as pink needles by sublimation at 40-50°/0.1mm (see 4-phenyl-1,2,4-triazoline-3,5-dione, PTAD below). [Cookson et al. *Org Synth* 51 121 1971; Cheng et al. *J Org Chem* 49 2910 1984.]

2-Methyltricycloquinazoline [2642-52-6] M 334.4, m >300°. Purified by vac sublimation. **CARCINOGEN**.

Methyl trifluoromethanesulfonate (methyl triflate) [333-27-7] M 164.1, b 97-97.5°/736mm, 99°/atm, 100-102°/atm, d₄²⁰ 1.496, n_D²⁵ 1.3238. It is a strong methylating agent but is corrosive and **POISONOUS**. Fractionate carefully and collecting the middle fraction (use efficient fume cupboard) and keep away from moisture. It is **POWERFUL ALKYLATING AGENT** and a strong **IRRITANT**. [IR: Gramstad and Haszeldine *J Chem Soc* 173 1956, 4069 1957.] *Trifluoromethanesulfonic acid* (triflic acid) [1493-13-6] M 151.1, boils higher (b 162°/atm), has a pK_a of 3.10, and is **TOXIC** and hygroscopic. [Hansen *J Org Chem* 30 4322 1965; Kurz and El-Nasr *J Am Chem Soc* 104 5823 1982.]

N-Methyltryptophan (L-abrine) [526-31-8] M 218.3, m 295°(dec), [α]_D²¹ +44.4° (c 2.8, 0.5M HCl), pK_{Est(1)}~2.3, pK_{Est(2)}~9.7. Crystd from water.

dl-5-Methyltryptophan [951-55-3] M 218.3, m 275°(dec) [pK see tryptophan]. Crystd from aqueous EtOH. *Picrate* has m 202° (dec).

6-Methyluracil [626-48-2] M 126.1, m 270-280°(dec), λ_{max} 260_{nm} logε 3.97, pK₁ ~1.1, pK₂ 9.8. Crystd from EtOH or acetic acid.

3-Methyluric acid [39717-48-1] M 182.1, m >350°, pK₁ 5.75 (6.2), pK₂ >12. Crystd from water.

7-Methyluric acid [30409-21-3] M 182.1, m >380°, pK₁ 5.6, pK₂ 10.3. Crystd from water.

9-Methyluric acid [30345-24-5] M 182.1, m >400°. Crystd from water.

Methyl vinyl ketone [78-94-4] M 70.1, b 62-68°/400mm, 79-80°/760mm, d 0.845, n 1.413. Forms an 85% azeotrope with water. After drying with K₂CO₃ and CaCl₂ (with cooling), the ketone is distd at low pressures.

Methyl vinyl sulfone [3680-02-2] M 106.1, b 116-118°/20mm, d 1.215, n 1.461. Passed through a column of alumina, then degassed and distd on a vacuum line and stored at -190° until required.

Methyl Violet 2B [4,4'-bis-(diethylamino)-4''-methyliminotriphenylmethyl hydrochloride] [8004-87-3] M 394.0, m 137°(dec), CI 42535, max ~580nm. Crystd from absolute EtOH by pptn with diethyl ether during cooling in an ice-bath. Filtered off and dried at 105°.

- 1-Methylxanthine** [6136-37-4] M 166.1, m >360° pK₁²⁰ 7.90, pK₂²⁰ 12.23. Crystd from water.
- 3-Methylxanthine** [1076-22-8] M 166.1, m >360° pK₁²⁰ 8.45, pK₂²⁰ 11.92. Crystd from water.
- 7-Methylxanthine** [552-62-5] M 166.1, m >380°(dec) pK₁²⁰ 8.42, pK₂²⁰ >13. Crystd from water.
- 8-Methylxanthine** [17338-96-4] M 166.1, m 292-293°(dec). Crystd from water.
- 9-Methylxanthine** [1198-33-0] M 166.1, m 384°(dec), pK₁²⁰ 2.0, pK₂²⁰ 6.12, pK₃²⁰ 10.5 (>13). Crystd from water.
- Metrazol (Cardiazol, Leptazol, 3a,4,5,6,7,8-hexahydro-1,2,3,3a-tetraaza-azulene, 1,5-pentamethylene-1,2,3,4-tetrazole)** [54-95-5] M 138.2, m 61°, b 194°/12mm, pK_{Est} ~ <0. Crystd from diethyl ether. Dried under vacuum over P₂O₅.
- Michler's ketone [4,4'-bis(dimethylamino)benzophenone]** [90-94-8] M 268.4, m 179°, pK²⁵ 9.84. Dissolved in dilute HCl, filtered and ppted by adding ammonia (to remove water-insoluble impurities such as benzophenone). Then crystd from EtOH or pet ether. [Suppan *J Chem Soc, Faraday Trans I* 71 539 1975.] It was also purified by dissolving in *benzene, then washed with water until the aqueous phase was colourless. The *benzene was evaporated off and the residue recrystd three times from *benzene and EtOH [Hoshino and Kogure *J Phys Chem* 72 417 1988].
- Monensin** [17090-79-8] M 670.9, m 103-105° (1 H₂O), [α]_D +47.7°, pK_{Est} ~ 4.6, pK 6.6 (66% Me₂NCHO). Purified by chromatography, stable in aq alkaline soln. Slightly sol in H₂O but sol in EtOH, EtOAc and Et₂O.
- N-Monobutyl urea** [592-31-4] M 116.2, m 96-98°. Crystd from EtOH/water, then dried under vacuum at room temperature.
- N-Monoethyl urea** [625-52-5] M 88.1, m 92-95°. Crystd from EtOH/water, then dried under vacuum at room temperature.
- N-Monomethyl urea** [598-50-5] M 74.1, m 93-95°. Crystd from EtOH/water, then dried under vacuum at room temperature.
- Monopropyl urea** [627-06-5] M 102.1, m 110°. Crystd from EtOH.
- Morin (hydrate) (2',3,4',5,7-pentahydroxyflavone)** [480-16-0] M 302.2, m 289-292°, pK₁ 5.3, pK₂ 8.74. Stirred at room temperature with ten times its weight of absolute EtOH, then left overnight to settle. Filtered, and evaporated under a heat lamp to one-tenth its volume. An equal volume of water was added, and the ppted morin was filtered off, dissolved in the minimum amount of EtOH and again ppted with an equal volume of water. The ppte was filtered, washed with water and dried at 110° for 1h. (Yield ca 2.5%.) [Perkins and Kalkwarf *Anal Chem* 28 1989 1956.] Complexes with W and Zr.
- Morphine (H₂O)** [57-27-2] M 302.2, m 230°(dec), [α]_D²³ -130.9° (MeOH), pK₁ 8.31, pK₂ 9.51. Crystd from MeOH.
- Morpholine** [110-91-8] M 87.1, f -4.9°, b 128.9°, d 1.0007, n 1.4540, n²⁵ 1.4533, pK²⁵ 8.33. Dried with KOH, fractionally distd, then refluxed with Na, and again fractionally distd. Dermer and Dermer [*J Am Chem Soc* 59 1148 1937] ppted as the oxalate by adding slowly to slightly more than 1 molar equivalent of oxalic acid in EtOH. The ppte was filtered and recrystd twice from 60% EtOH. Addition of the oxalate to conc aq NaOH regenerated the base, which was separated and dried with solid KOH, then sodium, before being fractionally distd.
- § A polystyrene supported morpholine is commercially available.

2-(*N*-Morpholino)ethanesulfonic acid (MES) [4432-31-9] M 213.3, m >300°(dec), pK²⁰ 6.15. Crystd from hot EtOH containing a little water.

Mucochloric acid (2,3-dichloro-4-oxo-2-butenic acid) [87-56-9] M 169.0, m 124-126°, pK²⁵ 4.20. Crystd twice from water (charcoal).

***trans,trans*-Muconic acid (hexa-2,4-dienedioic acid)** [3588-17-8] M 142.1, m 300°, pK²⁵ 4.51, for *cis,cis* pK²⁵ 4.49. Cryst from H₂O.

Muramic acid (H₂O) (3-*O*- α -carboxyethyl-D-glucosamine) [1114-41-6] M 251.2, m 152-154°(dec). See muramic acid on p. 549 in Chapter 6.

Murexide (ammonium purpurate) [3051-09-0] M 284.2, m >300°, λ_{\max} 520nm (ϵ 12,000), pK₂ 9.2, pK₃ 10.9. The sample may be grossly contaminated with uramil, alloxanthine, etc. Difficult to purify. It is better to synthesise it from pure alloxanthine [Davidson *J Am Chem Soc* 58 1821 1936]. Crystd from water.

Myristic acid (tetradecanoic acid) [544-63-8] M 228.4, m 58°, pK²⁰ 6.3 (50% EtOH), pK_{Est} ~4.9 (H₂O). Purified *via* the *methyl ester* (b 153-154°/10mm, n²⁵ 1.4350), as for capric acid. [Trachtman and Miller *J Am Chem Soc* 84 4828 1962.] Also purified by zone melting. Crystd from pet ether and dried in a vacuum desiccator containing shredded wax.

Naphthacene (benz[b]anthracene, 2,3-benzanthracene, rubene) [92-24-0] M 228.3, m >300°, 341° (open capillary), 349°, 357°. Crystd from EtOH or *benzene. Dissolved in sodium-dried *benzene and passed through a column of alumina. The *benzene was evaporated under vacuum, and the chromatography was repeated using fresh *benzene. Finally, the naphthacene was sublimed under vacuum. [Martin and Ubbelohde *J Chem Soc* 4948 1961.] Also recrystd in orange needles from xylene and sublimes *in vacuo* at 186°. [UV: *Chem Ber* 65 517 1932, 69 607 1936; IR: *Spectrochim Acta* 4 373 1951.]

2-Naphthaldehyde [66-99-9] M 156.2, m 59°, b 260°/19mm, pK²⁰ -7.04 (aq H₂SO₄). Distilled with steam and crystd from water or EtOH.

Naphthalene [91-20-3] M 128.2, m 80.3°, b 87.5°/10mm, 218.0°/atm, d 1.0253, d¹⁰⁰ 0.9625, n⁸⁵ 1.5590. Crystd one or more times from the following solvents: EtOH, MeOH, CCl₄, *benzene, glacial acetic acid, acetone or diethyl ether, followed by drying at 60° in an Abderhalden drying apparatus. Also purified by vacuum sublimation and by fractional crystn from its melt. Other purification procedures include refluxing in EtOH over Raney Ni, and chromatography of a CCl₄ soln on alumina with *benzene as eluting solvent. Baly and Tuck [*J Chem Soc* 1902 1908] purified naphthalene for spectroscopy by heating with conc H₂SO₄ and MnO₂, followed by steam distn (repeating the process), and formation of the picrate which, after recrystallisation, was decomposed and the naphthalene was steam distd. It was then crystd from dilute EtOH. It can be dried over P₂O₅ under vacuum. Also purified by sublimation and subsequent crystn from cyclohexane. Alternatively, it has been washed at 85° with 10% NaOH to remove phenols, with 50% NaOH to remove nitriles, with 10% H₂SO₄ to remove organic bases, and with 0.8g AlCl₃ to remove thianaphthalenes and various alkyl derivatives. Then it was treated with 20% H₂SO₄, 15% Na₂CO₃ and finally distd. [Gorman et al. *J Am Chem Soc* 107 4404 1985.]

Zone refining purified naphthalene from anthracene, 2,4-dinitrophenylhydrazine, methyl violet, benzoic acid, methyl red, chrysene, pentacene and indoline.

Naphthalene-2,5-disulfonic acid [92-41-1] M 288.2, pK_{Est} <0. Crystd from conc HCl.

Naphthalene-1-sulfonic acid [85-47-2] M 208.2, m (2H₂O) 90°, (anhydrous) 139-140°, pK²⁰ -0.17. Crystd from conc HCl and twice from water.

2-(*N*-Morpholino)ethanesulfonic acid (MES) [4432-31-9] M 213.3, m >300°(dec), pK²⁵ 6.15. Crystd from hot EtOH containing a little water.

Mucochloric acid (2,3-dichloro-4-oxo-2-butenic acid) [87-56-9] M 169.0, m 124-126°, pK²⁵ 4.20. Crystd twice from water (charcoal).

***trans,trans*-Muconic acid (hexa-2,4-dienedioic acid)** [3588-17-8] M 142.1, m 300°, pK²⁵ 4.51, for *cis,cis* pK²⁵ 4.49. Cryst from H₂O.

Muramic acid (H₂O) (3-*O*- α -carboxyethyl-D-glucosamine) [1114-41-6] M 251.2, m 152-154°(dec). See muramic acid on p. 549 in Chapter 6.

Murexide (ammonium purpurate) [3051-09-0] M 284.2, m >300°, λ_{\max} 520nm (ϵ 12,000), pK₁ 9.2, pK₃ 10.9. The sample may be grossly contaminated with uramil, alloxanthine, etc. Difficult to purify. It is better to synthesise it from pure alloxanthine [Davidson *J Am Chem Soc* 58 1821 1936]. Crystd from water.

Myristic acid (tetradecanoic acid) [544-63-8] M 228.4, m 58°, pK²⁰ 6.3 (50% EtOH), pK_{Est} ~4.9 (H₂O). Purified *via* the *methyl ester* (b 153-154°/10mm, n²⁵ 1.4350), as for capric acid. [Trachtman and Miller *J Am Chem Soc* 84 4828 1962.] Also purified by zone melting. Crystd from pet ether and dried in a vacuum desiccator containing shredded wax.

Naphthacene (benz[b]anthracene, 2,3-benzanthracene, rubene) [92-24-0] M 228.3, m >300°, 341° (open capillary), 349°, 357°. Crystd from EtOH or *benzene. Dissolved in sodium-dried *benzene and passed through a column of alumina. The *benzene was evaporated under vacuum, and the chromatography was repeated using fresh *benzene. Finally, the naphthacene was sublimed under vacuum. [Martin and Ubbelohde *J Chem Soc* 4948 1961.] Also recrystd in orange needles from xylene and sublimes *in vacuo* at 186°. [UV: *Chem Ber* 65 517 1932, 69 607 1936; IR: *Spectrochim Acta* 4 373 1951.]

2-Naphthaldehyde [66-99-9] M 156.2, m 59°, b 260°/19mm, pK²⁰ -7.04 (aq H₂SO₄). Distilled with steam and crystd from water or EtOH.

Naphthalene [91-20-3] M 128.2, m 80.3°, b 87.5°/10mm, 218.0°/atm, d¹⁰⁰ 0.9625, n⁸⁵ 1.5590. Crystd one or more times from the following solvents: EtOH, MeOH, CCl₄, *benzene, glacial acetic acid, acetone or diethyl ether, followed by drying at 60° in an Abderhalden drying apparatus. Also purified by vacuum sublimation and by fractional crystn from its melt. Other purification procedures include refluxing in EtOH over Raney Ni, and chromatography of a CCl₄ soln on alumina with *benzene as eluting solvent. Baly and Tuck [*J Chem Soc* 1902 1908] purified naphthalene for spectroscopy by heating with conc H₂SO₄ and MnO₂, followed by steam distn (repeating the process), and formation of the picrate which, after recrystallisation, was decomposed and the naphthalene was steam distd. It was then crystd from dilute EtOH. It can be dried over P₂O₅ under vacuum. Also purified by sublimation and subsequent crystn from cyclohexane. Alternatively, it has been washed at 85° with 10% NaOH to remove phenols, with 50% NaOH to remove nitriles, with 10% H₂SO₄ to remove organic bases, and with 0.8g AlCl₃ to remove thianaphthalenes and various alkyl derivatives. Then it was treated with 20% H₂SO₄, 15% Na₂CO₃ and finally distd. [Gorman et al. *J Am Chem Soc* 107 4404 1985.]

Zone refining purified naphthalene from anthracene, 2,4-dinitrophenylhydrazine, methyl violet, benzoic acid, methyl red, chrysene, pentacene and indoline.

Naphthalene-2,5-disulfonic acid [92-41-1] M 288.2, pK_{Est} <0. Crystd from conc HCl.

Naphthalene-1-sulfonic acid [85-47-2] M 208.2, m (2H₂O) 90°, (anhydrous) 139-140°, pK²⁰ -0.17. Crystd from conc HCl and twice from water.

Naphthalene-2-sulfonic acid [120-18-3] M 208.2, m 91°, pK_{Est} <1. Crystd from conc HCl.

Naphthalene-1-sulfonyl chloride [85-46-1] M 226.7, m 64-67°, 68°, b 147.5°/0.9 mm, 147.5°/13mm. If the IR indicates the presence of OH then treat with an equal weight of PCl₅ and heat at ca 100° for 3h, cool and pour into ice + H₂O, stir well and filter off the solid. Wash the solid with cold H₂O and dry the solid in a vacuum desiccator over P₂O₅ + solid KOH. Extract the solid with pet ether (b 40-60°) filter off any insoluble solid and cool. Collect the crystalline sulfonyl chloride and recryst from pet ether or *C₆H₆ pet ether. If large quantities are available then it can be distd under high vacuum. [Fierz-Davaid and Weissenbach *Helv Chim Acta* 3 2312 1920.] The *sulfonamide* has m 150° (from EtOH or H₂O).

Naphthalene-2-sulfonyl chloride [93-11-8] M 226.7, m 74-76°, 78°, 79°, b 148°/0.6 mm, 201°/13mm. Crystd (twice) from *benzene/pet ether (1:1 v/v). Purified as the 2-sulfonyl chloride. [Fierz-Davaid and Weissenbach *Helv Chim Acta* 3 2312 1920.] The *sulfonamide* has m 217° (from EtOH).

1,8-Naphthalic acid (naphthalene-1,8-dicarboxylic acid) [518-05-8] M 216.9, m 270°, pK_{Est(1)} ~ 2.1, pK_{Est(2)} ~ 4.5. Crystd from EtOH or aq EtOH.

1,8-Naphthalic anhydride [81-84-5] M 198.2, m 274°. Extracted with cold aqueous Na₂CO₃ to remove free acid, then crystd from acetic anhydride.

Naphthamide [2243-82-5] M 171.2, m 195°, pK²⁰ -2.30 (H₀ scale, aq H₂SO₄). Crystd from EtOH.

Naphthazarin (5,8-dihydroxy-1,4-naphthoquinone) [475-38-7] M 190.2, m ~ 220-230°(dec), m 225-230°, pK_{Est(1)} ~ 9.5, pK_{Est(2)} ~ 11.1. Red-brown needles with a green shine from EtOH. Also recrystd from hexane and purified by vacuum sublimation. [Huppert et al. *J Phys Chem* 89 5811 1985.] It is sparingly soluble in H₂O but soluble in alkalis. It sublimates at 2-10mm. The *diacetate* forms golden yellow prisms from CHCl₃, m 192-193° and the *5,8-dimethoxy* derivative has m 157° (155°) (from pet ether) [Bruce and Thompson *J Chem Soc* 1089 1955; IR: Schmand and Boldt *J Am Chem Soc* 97 447 1975; NMR: Brockmann and Zeeck *Chem Ber* 101 4221 1968]. The *monothiosemicarbazone* has m 168°(dec) from EtOH [Gardner et al. *J Am Chem Soc* 74 2106 1952].

Naphthionic acid (4-aminonaphthalene-1-sulfonic acid) [84-86-6] M 223.3, m > 300°(dec), pK²⁵ 2.68. It crystallises from H₂O as needles of the 0.5 hydrate. Salt solns fluoresce strongly blue.

1-Naphthoic acid [86-55-5] M 172.2, m 162.5-163.0°, pK²⁵ 3.60. Crystd from toluene (3mL/g) (charcoal), pet ether (b 80-100°), or aqueous 50% EtOH.

2-Naphthoic acid [93-09-4] M 172.2, m 184-185°, pK²⁵ 4.14. Crystd from EtOH (4mL/g), or aqueous 50% EtOH. Dried at 100°.

1-Naphthol [90-15-3] M 144.2, m 95.5-96°, pK²⁵ 9.34. Sublimed, then crystd from aqueous MeOH (charcoal), aq 25% or 50% EtOH, *benzene, cyclohexane, heptane, CCl₄ or boiling water. Dried over P₂O₅ under vacuum. [Shizuka et al. *J Am Chem Soc* 107 7816 1985.]

2-Naphthol [135-19-3] M 144.2, m 122.5-123.5°, pK²⁵ 9.57. Crystd from aqueous 25% EtOH (charcoal), water, *benzene, toluene or CCl₄, e.g. by repeated extraction with small amounts of EtOH, followed by dissolution in a minimum amount of EtOH and pptn with distilled water, then drying over P₂O₅ under vacuum. Has also been dissolved in aqueous NaOH, and ppted by adding acid (repeated several times), then ppted from *benzene by addition of heptane. Final purification can be by zone melting or sublimation *in vacuo*. [Bardez et al. *J Phys Chem* 89 5031 1985; Kikuchi et al. *J Phys Chem* 91 574 1987.]

Naphthol AS-D (3-hydroxy-2-naphthoic-o-toluide) [135-61-5] M 277.3, m 1196-198°. Purified by recrystn from xylene. Gives yellow-green fluorescent solutions at pH 8.2-9.5, [IR: Schnopper et al. *Anal Chem* 31 1542 1959.] With AcCl *naphthol AS-D acetate* is obtained m 168-169°, and with

chloroacetyl chloride *naphthol AS-D-chloroacetate* is obtained [Moloney et al. *J Histochem Cytochem* **8** 200 1960; Burstone *Arch Pathology* **63** 164 1957].

α -Naphtholbenzein [bis-(α -{4-hydroxynaphth-1-yl})-benzyl alcohol] [6948-88-5] **M 392.5, m 122-125^o, pK_{Est} ~ 9.3.** Crystd from EtOH, aqueous EtOH or glacial acetic acid.

1-Naphthol-2-carboxylic acid (1-hydroxy-2-naphthoic acid) [86-48-6] **M 188.2, m 203-204^o, pK_{Est(1)}~2.5, pK_{Est(2)}~12.** Successively crystd from EtOH/water, diethyl ether and acetonitrile, with filtration through a column of charcoal and Celite. [Tong and Glesmann *J Am Chem Soc* **79** 583 1957.]

2-Naphthol-3-carboxylic acid (2-hydroxy-3-naphthoic acid) [92-70-6] **M 188.2, m 222-223^o, pK₁²⁵ 2.79, pK₂²⁵ 12.84.** Crystd from water or acetic acid.

1,2-Naphthoquinone [524-42-5] **M 158.2, m 140-142^o(dec).** Crystd from ether (red needles) or *benzene (orange leaflets).

1,4-Naphthoquinone [130-15-4] **M 158.2, m 125-125.5^o.** Crystd from diethyl ether (charcoal). Steam distd. Crystd from *benzene or aqueous EtOH. Sublimed in a vacuum.

β -Naphthoxyacetic acid [120-23-0] **M 202.2, m 156^o, pK_{Est} ~3.0.** Crystd from hot water or *benzene.

β -Naphthoyltrifluoroacetone (4,4,4-trifluoro-2-naphthylbutan-1,3-dione) [893-33-4] **M 266.2, m 70-71^o, 74-76^o, pK²⁰ 6.35.** Crystd from EtOH. The *mono oxime* crystd from H₂O or aq EtOH has m 137-138^o. [Reid and Calvin *J Am Chem Soc* **72** 2948 1950.]

Naphthvalene [34305-47-0] **M 104.1, m dec at 175^o to benzvalene.** Purified by chromatography on alumina and eluting with pentane. It is stable at room temp [Abelt et al. *J Am Chem Soc* **107** 4148 1985]. The ¹H NMR in CCl₄ has τ 3.18 (4H), 6.17 (t J 1.5Hz, 2H), 7.60 (t J 1.5Hz 2H).

1-Naphthyl acetate [830-81-9] **M 186.2, m 45-46^o.** Chromatographed on silica gel and crystd as the 2-isomer below.

2-Naphthyl acetate [1523-11-1] **M 186.2, m 71^o.** Crystd from pet ether (b 60-80^o) or dilute aq EtOH.

1-Naphthylacetic acid [86-87-3] **M 186.2, m 132^o, pK²⁵ 4.23.** Crystd from EtOH or water.

2-Naphthylacetic acid [581-96-4] **M 186.2, m 143.1-143.4^o, pK²⁵ 4.30.** Crystd from water or *benzene.

1-Naphthylamine [134-32-7] **M 143.2, m 50.8-51.2^o, b 160^o, pK²⁵ 3.94.** Sublimed at 120^o in a stream of nitrogen, then crystd from pet ether (b 60-80^o), or abs EtOH then diethyl ether. Dried under vacuum in an Abderhalden pistol. Has also been purified by crystn of its hydrochloride from water, followed by liberation of the free base and distn; finally purified by zone melting. **CARCINOGEN.**

1-Naphthylamine hydrochloride [552-46-5] **M 179.7, m sublimes on heating.** Crystd from water (charcoal).

2-Naphthylamine [91-59-8] **M 143.2, m 113^o, pK²⁵ 4.20.** Sublimed at 180^o in a stream of nitrogen. Crystd from hot water (charcoal) or *benzene. Dried under vacuum in an Abderhalden pistol. **CARCINOGEN.**

1-Naphthylamine-5-sulfonic acid [84-89-9] **M 223.3, m >200^o(dec), pK_{Est(1)}<1, pK₂²⁵ 3.69 (NH₂)** Crystd under nitrogen from boiling water and dried in a steam oven [Bryson *Trans Faraday Soc* **47** 522, 527 1951].

2-Naphthylamine-1-sulfonic acid [81-16-3] M 223.3, m >200°(dec), $pK_1^{25} < 1$, $pK_2^{25} 2.35$ (NH₂). Crystd under nitrogen from boiling water and dried in a steam oven [Bryson *Trans Faraday Soc* 47 522, 527 1951].

2-Naphthylamine-6-sulfonic acid [93-00-5] M 223.3, m >200°(dec). Crystd from a large volume of hot water.

1-(1-naphthyl) ethanol [R-(+)- 42177-25-3; S-(-)- 15914-84-8] M 172.2, m 46°, 45-47.5°, 48°, $[\alpha]_{546}^{20}$ (+) and (-) 94°, $[\alpha]_D^{20}$ (+) and (-) 78° (c 1, MeOH). Purified by recrystn from Et₂O-pet ether, Et₂O, hexane [Balfe et al. *J Chem Soc* 797 1946; IR, NMR: Theisen and Heathcock *J Org Chem* 53 2374 1988; see also Fredga et al. *Acta Chem Scand* 11 1609 1957]. The *RS*-alcohol [57605-95-5] has m 63-65°, 65-66° from hexane.

1-(1-Naphthyl)ethylamine [R-(+)- 3886-70-2; S-(-)- 10420-89-0] M 171.2, b 153°/11mm, 178-181°/20mm, $d_4^{20} 1.067$, $n_D^{20} 1.624$, $[\alpha]_{546}^{20}$ (+) and (-) 65°, $[\alpha]_D^{20}$ (+) and (-) 55° (c 2, MeOH); $[\alpha]_D^{17}$ (+) and (-) 82.8° (neat), $pK_{Est} \sim 9.3$. Purified by distn in a good vacuum. [Mori et al. *Tetrahedron* 37 1343 1981; cf Wilson in *Top Stereochem* (Allinger and Eliel eds) vol 6 135 1971; Fredga et al. *Acta Chem Scand* 11 1609 1957.] The *hydrochlorides* crystallises from H₂O $[\alpha]_D^{18} \pm 3.9^\circ$ (c 3, H₂O) and the *sulfates* recrystallises from H₂O as *tetrahydrates* m 230-232°. The *RS*-amine has b 153°/11mm, 156°/15mm, 183.5°/41mm [Blicke and Maxwell *J Am Chem Soc* 61 1780 1939].

2-Naphthylethylene (2-vinylnaphthalene) [827-54-3] M 154.2, m 66°, b 95-96°/2.1mm, 135-137°/18mm. Crystd from aqueous EtOH.

***N*-(α -Naphthyl)ethylenediamine dihydrochloride** [1465-25-4] M 291.2, m 188-190°, $pK_{Est(1)} \sim 3.8$, $pK_{Est(2)} \sim 9.4$. Crystd from water.

1-Naphthyl isocyanate [86-84-0] M 169.2, m 3-5°, b 269-270°/atm, $d_4^{20} 1.18$. Distd at atmospheric pressure or in a vacuum. Can be crystd from pet ether (b 60-70°) at low temperature. *It has a pungent odour, is TOXIC and is absorbed through the skin.*

1-Naphthyl isothiocyanate [551-06-4] M 185.3, m 58-59°. Crystd from hexane (1g in 9 mL). White needles soluble in most organic solvents but is insoluble in H₂O. *It is absorbed through the skin and may cause dermatitis.* [*Org Synth Coll Vol IV* 700 1963.]

2-Naphthyl lactate [93-43-6] M 216.2. Crystd from EtOH.

2-(2-Naphthyloxy)ethanol [93-20-9] M 188.2, m 76.7°. Crystd from *benzene/pet ether.

***N*-1-Naphthylphthalamic acid** [132-66-1] M 291.3, m 203°. Crystd from EtOH.

2-Naphthyl salicylate [613-78-5] M 264.3, m 95°, $pK_{Est} \sim 10.0$. Crystd from EtOH.

1-Naphthyl thiourea (ANTU) [86-88-4] M 202.2, m 198°. Crystd from EtOH.

1-Naphthyl urea [6950-84-1] M 186.2, m 215-220°. Crystd from EtOH.

2-Naphthyl urea [13114-62-0] M 186.2, m 219-220°. Crystd from EtOH.

1,5-Naphthyridine [254-79-5] M 130.1, m 75°, b 112°/15mm, $pK^{20} 2.84$. Purified by repeated sublimation.

Narcein {6-[6-(2-dimethylaminoethyl)]-2-methoxy-3,4-(methylenedioxy)phenylacetyl]-2,3-dimethoxybenzoic acid} [131-28-2] M 445.4, m 176-177° (145° anhydrous), pK_1^{15} 3.5, pK_2^{15} 9.3. Crystd from water (as trihydrate).

Naringenin (4',5,7-trihydroxyflavanone) [480-41-1] M 272.3, m 251° (phenolic pK_s ~ 8-11). Crystd from aqueous EtOH.

Naringin (naringenin 7-rhamnoglucoside) [10236-47-2] M 580.5, m 171° (2H₂O), $[\alpha]_D^{19}$ -90° (c 1, EtOH), $[\alpha]_{546}^{20}$ -107° (c 1, EtOH). Crystd from water. Dried at 110°(to give the dihydrate).

Neopentane (2,2-dimethylpropane) [463-82-1] M 72.2, b 79.3°, d 0.6737, n 1.38273. Purified from isobutene by passage over conc H₂SO₄ or P₂O₅, and through silica gel.

Neostigmine [(3-dimethylcarbamoylphenyl)trimethylammonium] bromide [114-80-7] M 303.2, m 176°(dec). Crystd from EtOH/diethyl ether. (*Highly TOXIC*).

Neostigmine methyl sulfate [51-60-5] M 334.4, m 142-145°. Crystd from EtOH. (*Highly TOXIC.*)

Nerolidol (3,7,11-trimethyl-1,6,10-dodecatrien-3-ol) M 222.4 [*cis/trans* 7212-44-4] b 122°/3mm, n 1.477, d 0.73, [*cis* 3790-78-1] b 70°/0.1mm, [*trans* 40716-66-3] b 78°/0.2mm, 145-146°/2mm. Purified by TLC on plates of Kieselguhr G [McSweeney *J Chromatogr* 17 183 1965] or silica gel impregnated with AgNO₃, using 1,2-CH₂Cl₂/CHCl₃/EtOAc/PrOH (10:10:1:1) as solvent system. Also by GLC on butanediol succinate (20%) on Chromosorb W. Stored under N₂ at ~5° in the dark.

Neutral Red (2-amino-8-dimethylamino-3-methylphenazine HCl, Basic Red 5, CI 50040) [553-24-2] M 288.8, m 290°(dec), pK^{25} 6.5. Crystd from *benzene/MeOH (1:1). In aq sol it is red at pH 6.8 and yellow at pH 8.0.

Nicotinaldehyde thiosemicarbazone [3608-75-1] M 180.2, m 222-223°. Crystd from water.

Nicotinamide [98-92-0] M 122.1, m 128-131°, pK_1^{20} 0.5, pK_2^{20} 3.33. Crystd from *benzene.

Nicotinic acid (niacin, 2-yridine-3-carboxylic acid) [59-67-6] M 123.1, m 232-234°, pK_1^{25} 2.00, pK_2^{25} 4.82. Crystd from *benzene.

Nicotinic acid hydrazide [553-53-7] M 137.1, m 158-159°, pK_1^{20} 3.3, pK_2^{20} 11.49(NH). Crystd from aqueous EtOH or *benzene.

Nile Blue A (a benzophenoxazinium sulfate dye) [3625-57-8] M 415.5, m >300°(dec), CI 51180, pK^{25} 2.4. Crystd from pet ether.

Ninhydrin (1,2,3-triketohydrindene hydrate) [485-47-2] M 178.1, m 241-243°(dec), pK^{30} 8.82. Crystd from hot water (charcoal). Dried under vacuum and stored in a sealed brown container.

Nitrilotriacetic acid [tris(carboxymethyl)amine, NTA, Complexone 1] [139-13-9] M 191.1, m 247°(dec), pK_1 0.8, pK_2 1.71, pK_3 2.47, pK_4 9.71. Crystd from water. Dried at 110°.

2-Nitroacetanilide [552-32-9] M 180.2, m 93-94°, pK_{Est} <0. Crystd from water.

4-Nitroacetanilide [104-04-1] M 180.2, m 217°, pK_{Est} <0. Ppted from 80% H₂SO₄ by adding ice, then washed with water, and crystd from EtOH. Dried in air.

3-Nitroacetophenone [121-89-1] M 165.2, m 81°, b 167°/18mm, 202°/760mm. Distilled in steam and crystd from EtOH.

4-Nitroacetophenone [100-19-6] M 165.2, m 80-81°, b 145-152°/760mm. Crystd from EtOH or aqueous EtOH.

3-Nitroalizarin (1,2-dihydroxy-3-nitro-9,10-anthraquinone, Alizarin Orange) [568-93-4] M 285.2, m 244° (dec), $pK_{Est(1)} \sim 4.6$, $pK_{Est(2)} \sim 9.6$. Crystd from acetic acid.

***o*-Nitroaniline** [88-74-4] M 138.1, m 72.5-73.0°, $pK^{25} -0.25 (-0.31)$. Crystd from hot water (charcoal), then crystd from water, aqueous 50% EtOH, or EtOH, and dried in a vacuum desiccator. Has also been chromatographed on alumina, then recrystd from *benzene.

***m*-Nitroaniline** [99-09-2] M 138.1, m 114°, $pK^{25} 2.46$. Purified as for *o*-nitroaniline. **Warning: it is absorbed through the skin.**

***p*-Nitroaniline** [100-01-6] M 138.1, m 148-148.5°, $pK^{25} 1.02$. Purified as for *o*-nitroaniline. Also crystd from acetone. Freed from *o*- and *m*-isomers by zone melting and sublimation.

***o*-Nitroanisole (2-methoxynitrobenzene)** [91-23-6] M 153.1, f 9.4°, b 265°/737mm, d 1.251, n 1.563. Purified by repeated vacuum distn in the absence of oxygen.

***p*-Nitroanisole (4-methoxynitrobenzene)** [100-17-4] M 153.1, m 54°. Crystd from pet ether or hexane and dried *in vacuo*.

9-Nitroanthracene [602-60-8] M 223.2, m 142-143°. Purified by recrystn from EtOH or MeOH. Further purified by sublimation or TLC.

5-Nitrobarbituric acid (dilituric acid) [480-68-2] M 173.1, m 176°, 176-183°, $pK^{20} 10.25$. Crystd from water.

***o*-Nitrobenzaldehyde** [552-89-6] M 151.1, m 44-45°, b 120-144°/3-6mm. Crystd from toluene (2-2.5mL/g) by addition of 7mL pet ether (b 40-60°) for 1mL of soln. Can also be distd at reduced pressures.

***m*-Nitrobenzaldehyde** [99-61-6] M 151.1, m 58°. Crystd from water or EtOH/water, then sublimed twice at 2mm pressure at a temperature slightly above its melting point.

***p*-Nitrobenzaldehyde** [555-16-8] M 151.1, m 106°. Purification as for *m*-nitrobenzaldehyde above.

Nitrobenzene [98-95-3] M 123.1, f 5.8°, b 84-86.5°/6.5-8mm, 210.8°/760mm, d 1.206, $n^{15} 1.55457$, $n 1.55257$, $pK^{18} -11.26$ (aq H_2SO_4). Common impurities include nitrotoluene, dinitrothiophene, dinitrobenzene and aniline. Most impurities can be removed by steam distn in the presence of dilute H_2SO_4 , followed by drying with $CaCl_2$, and shaking with, then distilling at low pressure from BaO, P_2O_5 , $AlCl_3$ or activated alumina. It can also be purified by fractional crystn from absolute EtOH (by refrigeration). Another purification process includes extraction with aqueous 2M NaOH, then water, dilute HCl, and water, followed by drying ($CaCl_2$, $MgSO_4$ or $CaSO_4$) and fractional distn under reduced pressure. The pure material is stored in a brown bottle, in contact with silica gel or CaH_2 . It is very *hygroscopic*.

4-Nitrobenzene-azo-resorcinol (magneson II) [74-39-5] M 259.2, m 199-200°. Crystd from EtOH.

2-Nitrobenzenesulfonyl chloride (NPS-Cl) [7669-54-7] M 189.6, m 73-74.5°, 74.5-75°, 74-76°. Recrystd from CCl_4 (2mL/g), filter off the soln at 5° (recovery 75%). Also recrystd from pet ether (b 40-60°), dried rapidly at 50° and stored in a brown glass bottle, sealed well and stored away from moisture. [Hubacher *Org Synth Coll Vol II* 455 1943; Ito et al. *Chem Pharm Bull (Jpn)* 26 296 1978.]

4-Nitrobenzhydrazide [606-26-8] M 181.1, m 213-214°. Crystd from EtOH.

4'-Nitrobenzo-15-crown-5 [60835-69-0] M 313.3, m 84-85°, 93-95°. Recrystd from EtOH, MeOH or *C₆H₆-hexane as for the 18-crown-6 compound below. It complexes with Na⁺, K⁺, NH₄⁺, Ca²⁺, Mg²⁺ and Cd²⁺. NMR (CDCl₃) has δ: 3.6-4.4 (m 16CH₂), 6.8 (d 1H arom), 7.65 (d 1H arom), 7.80 (dd 1H arom *J*_{ab} 9Hz and *J*_{bc} 3Hz) [Shmid et al. *J Am Chem Soc* **98** 5198 1976; Kikukawa et al. *Bull Chem Soc Jpn* **50** 2207 1977; Toke et al. *Justus Liebigs Ann Chem* **349** 349, 761 1988; Lindner et al. *Z Anal Chem* **322** 157 1985].

4'-Nitrobenzo-18-crown-6 [53408-96-1] M 357.4, m 83-84°, 83-84°. If impure and discoloured then chromatograph on Al₂O₃ and eluting with *C₆H₆ (1:1) with 1% MeOH added. The fractions are followed by TLC on Al₂O₃ (using detection with Dragendorff's reagent R_F 0.6 in the above solvent system). Recrystallise the residues from the fractions containing the product from *C₆H₆-hexane to give yellowish leaflets. It complexes with Na or K ions with logK_{Na} 3.95 and logK_K 4.71. [Petranek and Ryba *Collect Chem Czech Chem Commun* **39** 2033 1974.]

2-Nitrobenzoic acid [552-16-9] M 167.1, m 146-148°, pK²⁵ 2.21. Crystd from *benzene (twice), *n*-butyl ether (twice), then water (twice). Dried and stored in a vacuum desiccator. [Le Noble and Wheland *J Am Chem Soc* **80** 5397 1958.] Has also been crystd from EtOH/water.

3-Nitrobenzoic acid [121-92-6] M 167.1, m 143-143.5°, pK²⁵ 3.46. Crystd from *benzene, water, EtOH (charcoal), glacial acetic acid or MeOH/water. Dried and stored in a vacuum desiccator.

4-Nitrobenzoic acid [62-23-7] M 167.1, m 241-242°, pK²⁵ 3.43. Purification as for 3-nitrobenzoic acid above.

4-Nitrobenzoyl chloride [122-04-3] M 185.6, m 75°, b 155°/20mm. Crystd from dry pet ether (b 60-80°) or CCl₄. Distilled under vacuum. **IRRITANT**.

4-Nitrobenzyl alcohol [619-73-8] M 153.1, m 93°. Crystd from EtOH and sublimed in a vacuum. Purity should be at least 99.5%. Sublimed samples should be stored in the dark over anhydrous CaSO₄ (Drierite). If the IR contains OH bands then the sample should be resublimed before use. [Mohammed and Kosower *J Am Chem Soc* **93** 2709 1979.]

4-Nitrobenzyl bromide [100-11-8] M 216.0, m 98.5-99.0°. Recrystd four times from abs EtOH, then twice from cyclohexane/hexane/*benzene (1:1:1), followed by vac sublimation at 0.1mm and a final recrystn from the same solvent mixture. [Lichtin and Rao *J Am Chem Soc* **83** 2417 1961.] Has also been crystd from pet ether (b 80-100°, 10mL/g, charcoal). It slowly decomposes even when stored in a desiccator in the dark. **IRRITANT**.

***m*-Nitrobenzyl chloride** [619-23-8] M 171.6, m 45°. Crystd from pet ether (b 90-120°). **IRRITANT**.

***p*-Nitrobenzyl chloride** [100-14-1] M 171.6, m 72.5-73°. Crystd from CCl₄, dry diethyl ether, 95% EtOH or *n*-heptane, and dried under vacuum. **IRRITANT**.

***p*-Nitrobenzyl cyanide** [555-21-5] M 162.2, m 117°. Crystd from EtOH. **TOXIC**.

4-(4-Nitrobenzyl)pyridine (PNBP) [1083-48-3] M 214.2, m 70-71°, pK_{Est} ~5.0. Crystd from cyclohexane.

2-Nitrobiphenyl [86-00-0] M 199.2, m 36.7°. Crystd from EtOH (seeding required). Sublimed under vacuum.

3-Nitrocinnamic acid [555-68-0] M 193.2, m 200-201°, pK²⁵ 2.58 (*trans*). Crystd from *benzene or EtOH.

4-Nitrocinnamic acid [882-06-4] M 193.2, m 143° (*cis*), 286° (*trans*), $pK_{Est} \sim 2.6$ (*trans*). Crystd from water.

4-Nitrodiphenylamine [836-30-6] M 214.2, m 133-134°, $pK^{25} -2.5$. Crystd from EtOH.

2-Nitrodiphenyl ether [2216-12-8] M 215.2, b 106-108°/0.01mm, 137-138°/0.5mm, 161-162°/4mm, 188-189°/12mm, 195-200°/25mm, $d_4^{20} 1.241$, $n_D^{25} 1.600$. Purified by fractional distn. UV (EtOH): 255, 315nm (ϵ 6200 and 2800); IR (CS₂): 1350 (NO₂) and 1245, 1265 (COC) cm⁻¹ [UV, IR: Dahlgard and Brewster *J Am Chem Soc* **80** 5861 1958; Tomita and Takase *Yakugaku Zasshi (J Pharm Soc Japan)* **75** 1077 1955; Fox and Turner *J Chem Soc* 1115 1930, Henley *J Chem Soc* 1222 1930].

Nitrodurene (1-nitro-2,3,5,6-tetramethylbenzene) [3463-36-3] M 179.2, m 53-55°, b 143-144°/10mm. Crystd from EtOH, MeOH, acetic acid, pet ether or chloroform.

Nitroethane [79-24-3] M 75.1, b 115°, d 1.049, n 1.3920, $n^{25} 1.39015$, $pK^{25} 8.60$ (8.46, pH equilibrium requires *ca* 5 min). Purified as described for *nitromethane*. A spectroscopic impurity has been removed by shaking with activated alumina, decanting and rapidly distilling.

2-Nitrofluorene [607-57-8] M 211.2, m 156°. Crystd from aqueous acetic acid.

Nitroguanidine [556-88-7] M 104.1, m 246-246.5°(dec), 257°, $pK_1^{25} -0.55$, $pK_2^{25} 12.20$. Crystd from water (20mL/g).

5-Nitroindole [6146-52-7] M 162.1, m 141-142°, $pK^{25} -7.4$ (aq H₂SO₄). Decolorised (charcoal) and recrystd twice from aqueous EtOH.

Nitromesitylene (2-nitro-1,3,5-trimethylbenzene) [603-71-4] M 165.2, m 44°, b 255°. Crystd from EtOH.

Nitromethane [75-52-5] M 61.0, f -28.5°, b 101.3°, d 1.13749, $d^{30} 1.12398$, n 1.3819, $n^{30} 1.37730$, $pK^{25} 10.21$. Nitromethane is generally manufactured by gas-phase nitration of methane. The usual impurities include aldehydes, nitroethane, water and small amounts of alcohols. Most of these can be removed by drying with CaCl₂ or by distn to remove the water/nitromethane azeotrope, followed by drying with CaSO₄. Phosphorus pentoxide is not suitable as a drying agent. [Wright et al. *J Chem Soc* 199 1936.] The purified material should be stored by dark bottles, away from strong light, in a cool place. Purifications using extraction are commonly used. For example, Van Looy and Hammett [*J Am Chem Soc* **81** 3872 1959] mixed about 150mL of conc H₂SO₄ with 1L of nitromethane and allowed it to stand for 1 or 2 days. The solvent was washed with water, aqueous Na₂CO₃, and again with water, then dried for several days with MgSO₄, filtered again with CaSO₄. It was fractionally distd before use. Smith, Fainberg and Winstein [*J Am Chem Soc* **83** 618 1961] washed successively with aqueous NaHCO₃, aqueous NaHSO₃, water, 5% H₂SO₄, water and dilute NaHCO₃. The solvent was dried with CaSO₄, then percolated through a column of Linde type 4A molecular sieves, followed by distn from some of this material (in powdered form). Buffagni and Dunn [*J Chem Soc* 5105 1961] refluxed for 24h with activated charcoal while bubbling a stream of nitrogen through the liquid. The suspension was filtered, dried (Na₂SO₄) and distd, then passed through an alumina column and redistd. It has also been refluxed over CaH₂, distd and kept under argon over 4A molecular sieves.

Can be purified by zone melting or by distn under vacuum at 0°, subjecting the middle fraction to several freeze-pump-thaw cycles. An impure sample containing higher nitroalkanes and traces of cyanoalkanes was purified (on the basis of its NMR spectrum) by crystn from diethyl ether at -60° (cooling in Dry-ice)[Parrett and Sun *J Chem Educ* **54** 448 1977].

Fractional crystn was more effective than fractional distn from Drierite in purifying nitromethane for conductivity measurements. [Coetzee and Cunningham *J Am Chem Soc* **87** 2529 1965.] Specific conductivities around 5×10^{-9} ohm⁻¹cm⁻¹ were obtained.

Nitron [1,4-diphenyl-3-phenylamino-(1*H*)-1,2,4-triazolium (hydroxide) inner salt] [2218-94-2] **M 312.4, m 189°(dec)**. Crystd from EtOH, chloroform or EtOH/*C₆H₆.

1-Nitronaphthalene [86-57-7] **M 173.2, m 57.3-58.3°, b 30-40°/0.01mm**. Fractionally distd under reduced pressure, then crystd from EtOH, aqueous EtOH or heptane. Chromatographed on alumina from *benzene/pet ether. Sublimes *in vacuo*.

2-Nitronaphthalene [581-89-5] **M 173.2, m 79°, b 165°/15mm**. Crystd from aqueous EtOH and sublimed in a vacuum.

1-Nitro-2-naphthol [550-60-7] **M 189.2, m 103°, pK²⁵ 5.93**. Crystd (repeatedly) from *benzene-pet ether (b 60-80°)(1:1).

2-Nitro-1-naphthol [607-24-9] **M 189.2, m 127-128°, pK²⁵ 5.89**. Crystd (repeatedly) from EtOH.

5-Nitro-1,10-phenanthroline [4199-88-6] **M 225.2, m 197-198°, pK²⁵ 3.33**. Crystd from *benzene/pet ether, until anhydrous.

2-Nitrophenol [88-75-5] **M 139.1, m 44.5-45.5°, pK²⁵ 7.23**. Crystd from EtOH/water, water, EtOH, *benzene or MeOH/pet ether (b 70-90°). Can be steam distd. Petrucci and Weygandt [*Anal Chem* **33** 275 1961] crystd from hot water (twice), then EtOH (twice), followed by fractional crystn from the melt (twice), drying over CaCl₂ in a vacuum desiccator and then in an Abderhalden drying pistol.

3-Nitrophenol [554-84-7] **M 139.1, m 96°, b 160-165°/12mm, pK²⁵ 8.36**. Crystd from water, CHCl₃, CS₂, EtOH or pet ether (b 80-100°), and dried under vacuum over P₂O₅ at room temperature. Can also be distd at low pressure.

4-Nitrophenol [100-02-7] **M 139.1, m 113-114°, pK²⁵ 7.16**. Crystd from water (which may be acidified, e.g. *N* H₂SO₄ or 0.5*N* HCl), EtOH, aqueous MeOH, CHCl₃, *benzene or pet ether, then dried under vacuum over P₂O₅ at room temperature. Can be sublimed at 60°/10⁻⁴mm.

2-Nitrophenoxyacetic acid [1878-87-1] **M 197.2, m 150-159°, pK²⁵ 2.90**. Crystd from water, and dried over P₂O₅ under vacuum..

***p*-Nitrophenyl acetate** [830-03-5] **M 181.2, m 78-79°**. Recrystd from absolute EtOH [Moss et al. *J Am Chem Soc* **108** 5520 1986].

2-Nitrophenylacetic acid [3740-52-1] **M 181.2, m 120°, pK²⁵ 3.95**. Crystd from EtOH/water and dried over P₂O₅ under vacuum.

4-Nitrophenylacetic acid [104-03-0] **M 181.2, m 80.5°, pK²⁵ 3.92**. Crystd from EtOH/water (1:1), then from sodium-dried diethyl ether and dried over P₂O₅ under vacuum.

4-Nitro-1,2-phenylenediamine [99-56-9] **M 153.1, m 201°, pK₁²⁵ 1.39 (1-NH₂), pK₂²⁵ 2.61 (2-NH₂)**. Crystd from water.

1-(4-Nitrophenyl)ethylamine hydrochloride [*R*-(+)- 57233-86-0; *S*-(-)- 132873-57-5] **M 202.6, m 225°, 240-242° (dec), 243-245° (dec), 248-250°, [α]_D²⁰ (+) and (-) 72° (c 1, 0.05 M NaOH), (+) and (-) 0.3° (H₂O), pK_{Est} ~8.6**. To ensure dryness the hydrochloride (*ca* 175 g) is extracted with EtOH (3x100mL) and evaporated to dryness (any residual H₂O increases the solubility in EtOH and lowers the yield). The hydrochloride residue is triturated with absolute EtOH and dried *in vacuo*. The product is further purified by refluxing with absolute EtOH (200 mL for 83g) for 1h, cool to 10° to give 76.6g of hydrochloride **m 243-245° (dec)**. The *free base* is prepd by dissolving in *N* NaOH, extract with CH₂Cl₂ (3 x 500mL), dry (Na₂CO₃), filter, evaporate and distil, **m 27°, b 119-120°/0.5mm (105-107°/0.5mm, 157-159°/9mm, d₄²⁰**

1.1764, n_D^{20} 1.5688, $[\alpha]_D^{24} \pm 17.7^\circ$ (neat)[Perry et al. *Synthesis* 492 1977; ORD: Nerdel and Liebig *Justus Liebigs Ann Chem* 621 142 1959].

4-Nitrophenylhydrazine [100-16-3] M 153.1, m 158°(dec), pK_1^{25} -9.2 (aq H_2SO_4), pK_2^{25} 3.70. Crystd from EtOH.

3-Nitrophenyl isocyanate [3320-87-4] M 164.1, m 52-54°. Crystd from pet ether (b 28-38°).

4-Nitrophenyl isocyanate [100-28-7] M 164.1, m 53°. Crystd from pet ether (b 28-38°).

2-Nitrophenylpropionic acid [530-85-8] M 191.1, m 157°(dec), pK^{25} 2.83. Crystd from water.

4-Nitrophenyl trifluoroacetate [658-78-6] M 235.1, m 37-39°, b 120°/12mm. Recrystd from $CHCl_3$ /hexane [Margolis et al. *J Biol Chem* 253 7891 1078].

4-Nitrophenyl urea [556-10-5] M 181.2, m 238°. Crystd from EtOH and hot water.

3-Nitrophthalic acid [603-11-2] M 211.1, m 216-218°, pK^{25} 3.93. Crystd from hot water (1.5mL/g). Air dried.

4-Nitrophthalic acid [610-27-5] M 211.1, m 165°, pK^{25} 4.12. Crystd from ether or ethyl acetate.

3-Nitrophthalic anhydride [641-70-3] M 193.1, m 164°. Crystd from *benzene, *benzene/pet ether, acetic acid or acetone. Dried at 100°.

1-Nitropropane [108-03-2] M 89.1, b 131.4°, d 1.004, n 1.40161, n^{25} 1.39936, pK^{25} 8.98. Purified as *nitromethane*.

2-Nitropropane [79-46-9] M 89.1, b 120.3°, d 0.989, n 1.3949, n^{25} 1.39206, pK^{25} 7.68. Purified as *nitromethane*.

5-Nitro-2-*n*-propoxyaniline [553-79-7] M 196.2, m 47.5-48.5°, pK_{Est} ~2.32. Crystd from *n*-propyl alcohol/pet ether.

3-Nitro-2-pyridinesulfonyl chloride [68206-45-1] M 190.2, m 217-222°(dec). Crystallises as yellow needles from CH_2Cl_2 . When pure it is stable for several weeks at room temperature, and no decomposition was observed after 6 months at <0°. UV (MeCN) has λ_{max} at 231nm (ϵ 12,988), 264nm (ϵ 5,784) and 372nm (ϵ 3,117). [NMR and UV: Matsuda and Aiba *Chem Lett* 951 1978; Wagner et al. *Chem Ber* 75 935 1942.]

5-Nitroquinoline [607-34-1] M 174.2, m 70°, pK^{20} 2.69. Crystd from pentane, then from *benzene.

8-Nitroquinoline [706-35-2] M 174.2, m 88-89°, pK^{20} 2.55. Crystd from hot water, MeOH, EtOH or EtOH/diethyl ether (3:1).

4-Nitroquinoline 1-oxide [56-57-5] M 190.2, m 157°. Recrystd from aqueous acetone [Seki et al. *J Phys Chem* 91 126 1987].

2-Nitroresorcinol [601-89-8] M 155.1, m 81-81°, pK_1^{20} 6.37, pK_2^{20} 9.46. Crystd from aq EtOH.

4-Nitrosalicylic acid [619-19-1] M 183.1, m 277-288°, pK^{25} 2.23. Crystd from water.

5-Nitrosalicylic acid [96-97-9] M 183.1, m 233°, pK_1^{25} 2.32, pK_2^{25} 10.34. Crystd from acetone (charcoal), then twice more from acetone alone.

Nitrosobenzene [586-96-9] M 107.1, m 67.5-68°, b 57-59°/18mm. Steam distd, then cryst from a small volume of EtOH with cooling below 0°, dried over CaCl₂ in a dessicator at atm pressure, and stored under N₂ at 0°. Alternatively it can be distd onto a cold finger cooled with brine at ~-10° in a vac at 17mm (water pump), while heating in a water bath at 65-70° [Robertson and Vaughan *J Chem Educ* 27 605 1950].

N-Nitrosodiethanolamine [1116-54-7] M 134.4, b 125°/0.01mm, n 1.485. Purified by dissolving the amine (0.5g) in 1-propanol (10mL) and 5g of anhydrous Na₂SO₄ added with stirring. After standing for 1-2h, it was filtered and passed through a chromatographic column packed with AG 50W x 8 (H⁺form, a strongly acidic cation exchanger). The eluent and washings were combined and evapd to dryness at 35°. [Fukuda et al. *Anal Chem* 53 2000 1981.] Possible **CARCINOGEN**.

4-Nitroso-N,N-dimethylaniline [138-89-6] M 150.2, m 86-87°, 92.5-93.5°, b 191-192°/100mm, pK²⁵ 4.54. Recryst from pet ether or CHCl₃-CCl₄ and dried in air. Alternatively suspend in H₂O, heat to boiling and add HCl until it dissolves. Filter, cool and collect the *hydrochloride* [42344-05-8], m 177° after recrystn from H₂O containing a small amount of HCl. The *hydrochloride* (e.g. 30g) is made into a paste with H₂O (100mL) in a separating funnel. Add cold aq 2.5 NaOH or Na₂CO₃ to a pH of ~ 8.0 (green color due to free base) and extracted with toluene, CHCl₃ or Et₂O. Dry extract (K₂CO₃), filter, distil off the solvent, cool residue and collect the crystalline free base. Recryst as above and dried in air.

N-Nitrosodiphenylamine [156-10-5] M 198.2, m 144-145°(dec). Crystd from *benzene.

1-Nitroso-2-naphthol [131-91-9] M 173.2, m 110.4-110.8°, pK²⁵ 7.63. Crystd from pet ether (b 60-80°, 7.5mL/g).

2-Nitroso-1-naphthol [132-53-6] M 173.2, m 158°(dec), pK²⁵ 7.24. Purified by recrystn from pet ether (b 60-80°) or by dissolving in hot EtOH, followed by successive addition of small volumes of water.

4-Nitroso-1-naphthol [605-60-7] M 173.2, m 198°, pK²⁵ 8.18. Crystd from *benzene.

2-Nitroso-1-naphthol-4-sulfonic acid (3H₂O) [3682-32-4] M 316.3, m 142-146°(dec), pK_{Est} ~6.3 (OH). Crystd from dilute HCl soln. Crystals were dried over CaCl₂ in a vacuum desiccator. Also purified by dissolution in aqueous alkali and pptn by addition of water. Reagent for cobalt.

4-Nitrosophenol [104-91-6] M 123.1, m >124°(dec), pK²⁵ 6.36. Crystd from xylene.

N-Nitroso-N-phenylbenzylamine [612-98-6] M 212.2, m 58°. Crystd from absolute EtOH and dried in air.

trans-β-Nitrostyrene [5153-67-3] M 149.2, m 60°. Crystd from absolute EtOH, or three times from *benzene/pet ether (b 60-80°) (1:1).

4-Nitrostyrene [100-13-0] M 149.2, m 20.5-21°. Crystd from CHCl₃/hexane. Purified by addition of MeOH to ppte the polymer, then crystd at -40° from MeOH. Also crystd from EtOH. [Bernasconi et al. *J Am Chem Soc* 108 4541 1986.]

2-Nitro-4-sulfobenzoic acid [552-23-8] M 247.1, m 111°, pK_{Est} ~1.65. Crystd from dilute HCl. Hygroscopic.

2-Nitrotoluene [88-72-2] M 137.1, m -9.55° (α-form), -3.85° (β-form), b 118°/16mm, d 1.163, 222.3°/760mm, n 1.545. Crystd (repeatedly) from absolute EtOH by cooling in a Dry-ice/alcohol mixture, Further purified by passage of an alcoholic soln through a column of alumina.

3-Nitrotoluene [99-08-1] M 137.1, m 16°, b 113-114°/15mm, 232.6°, d 1.156, n 1.544. Dried with P₂O₅ for 24h, then fractionally distd under reduced pressure. [*Org. Synth Coll Vol I* 416 1948.]

4-Nitrotoluene [99-99-0] M 137.1, m 52°. Crystd from EtOH, MeOH/water, EtOH/water (1:1) or MeOH. Air dried, then dried in a vac desiccator over H₂SO₄. [Wright and Grilliom *J Am Chem Soc* 108 2340 1986.]

5-Nitrouracil (2,4-dihydroxy-5-nitropyrimidine) [611-08-5] M 157.1, m 280-285°, >300°, pK₁²⁰ 0.03, pK₂²⁰ 5.55, pK₃²⁰ 11.3. Recrystallises as prisms from boiling H₂O as the monohydrate and loses H₂O on drying *in vacuo*. [UV: Brown *J Chem Soc* 3647 1959; Brown *J Appl Chem* 2 239 1952; Johnson *J Am Chem Soc* 63 263 1941.]

Nitrourea [556-89-8] M 105.1, m 158.4-158.8°(dec). Crystd from EtOH/pet ether.

5-Nitrovanillin (nitroveratric aldehyde) [6635-20-7] M 197.2, m 172-175°, 176°, 178°. Forms yellow plates from AcOH, and needles from EtOH [Slotta and Szyszke *Chem Ber* 68 184 1935]. With diazomethane, 5-nitro-3,4-dimethoxyacetophenone is formed [Brady and Manjunath *J Chem Soc* 125 1067 1924]. The *methyl ether* crystallises from EtOAc or AcOH, m 88°, 90-91°, and the *phenylhydrazone* has m 108-110° (from aqueous EtOH). [Finger and Schott *J Prakt Chem* [2] 115 288 1927.] For *oxime* m 216° (from EtOH or AcOH) and the *oxime acetate* has m 147° (from aq EtOH) [Vogel *Monatsh Chem* 20 384 1899; Brady and Dunn *J Chem Soc* 107 1861 1915].

n-Nonane [111-84-2] M 126.3, b 150.8°, d 0.719, n 1.40542, n²⁵ 1.40311. Fractionally distd, then stirred with successive volumes of conc H₂SO₄ for 12h each until no further colouration was observed in the acid layer. Then washed with water, dried with MgSO₄ and fractionally distd. Alternatively, it was purified by azeotropic distn with 2-ethoxyethanol, followed by washing out the alcohol with water, drying and distilling. [Forziati et al. *J Res Nat Bur Stand* 36 129 1946].

2,5-Norbornadiene [121-46-0] M 92.1, b 89°, d 0.854, n 1.4707. Purified by distn from activated alumina [Landis and Halpern *J Am Chem Soc* 109 1746 1987].

cis-endo-5-Norbornene-2,3-dicarboxylic anhydride (carbic anhydride, 3 α ,4,7,7, $\alpha\alpha$ -tetrahydro-4 α ,7 α -methanoisobenzofuran-1,3-dione) [129-64-6] M 164.2, m 164.1°, 164-165°, 164-167°, d 1.417. Forms crystals from pet ether, hexane or cyclohexane. It is hydrolysed by H₂O to form the acid [Diels and Alder *Justus Liebigs Ann Chem* 460 98 1928; Maitte *Bull Soc Chim Fr* 499 1959]. The *exo-exo-isomer* has m 142-143° (from *C₆H₆-pet ether) [Alder and Stein *Justus Liebigs Ann Chem* 504 216 1933].

Norbornylene [498-66-8] M 94.2, m 44-46°, b 96°. Refluxed over Na, and distd [Gilliom and Grubbs *J Am Chem Soc* 108 733 1986]. Also purified by sublimation *in vacuo* onto an ice-cold finger [Woon et al. *J Am Chem Soc* 108 7990 1986].

Norcamphor (bicyclo[2.2.1]heptan-2-one) [497-38-1] M 110.2, m 94-95°. Crystd from water.

Norcholanic acid [511-18-2] M 346.5, m 177°, 186°, [α]_D²⁰ +32° (EtOH), pK_{Est} ~4.8. Crystd from acetic acid.

Norcodeine [467-15-2] M 285.3, m 185°, 186°, pK 9.10. Crystd from acetone or ethyl acetate.

Nordihydroguaiaretic [1,4-bis(3,4-dihydroxyphenyl)-2,3-dimethylbutane] acid [500-38-9] M 302.4, m 184-185°, pK_{Est(1)} ~9.7, pK_{Est(2)} ~12. Crystd from dilute acetic acid.

Norleucine (α -amino-n-caproic acid) [R(+)- 327-56-0; S(-)- 327-57-1] M 117.2, m 301° [α]₅₄₆²⁰ (+) and (-) 28° (c 5, 5M HCl); [RS: 616-06-8] m 297-300° (sublimes partially at ~280°), pK₁ 2.39, pK₂ 9.76 (for RS). Crystd from water.

Norvaline (*R*- α -amino-*n*-valeric acid) [*R*(+)- 2031-12-9; *S*(-)- 6600-40-4] M 117.2, m 305°(dec), $[\alpha]_{546}^{20}$ (+) and (-) 25° (c 10, 5M HCl), pK_1^{25} 2.36, pK_2^{25} 9.87 (9.72). Crystd from aqueous EtOH or water.

Nylon powder. Pellets were dissolved in ethylene glycol under reflux. Then ppted as a white powder on addition of EtOH at room temperature. This was washed with EtOH and dried at 100° under vacuum.

***n*-Octacosane** [630-02-4] M 394.8, m 62.5°. Purified by forming its adduct with urea, washing and crystallising from acetone/water. [McCubbin *Trans Faraday Soc* 58 2307 1962.] Crystd from hot, filtered isopropyl ether soln (10mL/g).

***n*-Octacosanol** (octacosyl alcohol) [557-61-9] M 410.8, m 83.4°, 84°. Recryst from large vols of Me₂CO. It sublimes at 200-250°/1mm instead of distilling.

***n*-Octadecane** [593-45-3] M 254.5, m 28.1°, b 173.5°/10mm, 316.1°/760mm, d_4^{20} 0.7768, n 1.4390. Crystd from acetone and distd under reduced pressure from sodium.

Octadecyl acetate [822-23-1] M 312.5, m 32.6°. Distd under vac, then crystd from diethyl ether/MeOH.

***n*-Octadecyl alcohol** (stearyl alcohol) [112-92-5] M 270.5, m 61°, b 153-154°/0.3mm. Crystd from MeOH, or dry diethyl ether and *benzene, then fractionally distd under reduced pressure. Purified by column chromatography. Freed from cetyl alcohol by zone melting.

Octadecyl ether (dioctadecyl ether) [6297-03-6] M 523.0, m 59.4°. Vacuum distd, then crystd from MeOH/*benzene.

Octadecyltrimethylammonium bromide [1120-02-1] M 392.5, m 250°(dec). See entry on p. 446 in Chapter 6.

2,3,7,8,12,13,17,18-Octaethyl-21*H*,23*H*-porphine [2683-82-1] M 534.8, m 322°, 326°. Chromatographed on SiO₂ using CHCl₃ as eluent. It crystallises from CHCl₃ (dark red), MeOH (blue violet), pyridine (m 318°) and *C₆H₆ (deep red). [Fischer and Bämle *Justus Liebigs Ann Chem* 468 58, 85 1929.]

Octafluoropropane (profluorane) [76-19-7] M 188.0, m -183°, b -38°. Purified for pyrolysis studies by passage through a copper vessel containing CoF₃ at about 270°, then fractionally distd. [Steunenberg and Cady *J Am Chem Soc* 74 4165 1952.] Also purified by several trap-to-trap distns at low temperatures [Simons and Block *J Am Chem Soc* 59 1407 1937].

1,2,3,4,6,7,8,9-Octahydroanthracene [1079-71-6] M 186.3, m 78°. Crystd from EtOH, then purified by zone melting.

Octamethylcyclotetrasiloxane [556-67-2] M 296.6, m 17.3°, b 175-176°, d_4^{20} 0.957, n 1.396. Purified by zone melting.

Octan-1,8-diol (octamethylene glycol) [629-41-4] M 146.2, m 59-61°, b 172°/20mm. Recrystd from EtOH and distd in a vac.

***n*-Octane** [111-65-9] M 114.2, b 126.5°, d_4^{20} 0.704, n 1.39743, n^{25} 1.39505. Extracted repeatedly with conc H₂SO₄ or chlorosulfonic acid, then washed with water, dried and distd. Also purified by azeotropic distn with EtOH, followed by washing with water to remove the EtOH, drying and distilling. For further details, see *n*-heptane. Also purified by zone melting.

1-Octanethiol [111-88-6] M 146.3, b 86°/15mm, 197-200°/760mm, d_4^{20} 0.8433, n 1.4540, pK^{25} 10.72 (dil *t*-BuOH). Passed through a column of alumina [Battacharyya et al. *J Chem Soc, Faraday Trans 1* 82 135 1986; Fletcher *J Am Chem Soc* 68 2727 1946].

1-Octene [111-66-0] M 112.2, b 121°/742mm, d_4^{20} 0.716, n 1.4087. Distd under nitrogen from sodium which removes water and peroxides. Peroxides can also be removed by percolation through dried, acid washed alumina. Stored under nitrogen in the dark. [Strukul and Michelin *J Am Chem Soc* 107 7563 1985.]

(trans)-2-Octene [13389-42-9] M 112.2, b 124-124.5°/760mm, d_4^{20} 0.722, n 1.4132. Purification as for 1-octene above.

***n*-Octyl alcohol** [111-87-5] M 130.2, b 98°/19mm, 195.3°/760mm, d 0.828, n 1.43018. Fractionally distd under reduced pressure. Dried with sodium and again fractionally distd or refluxed with boric anhydride and distd (b 195-205°/5mm), the distillate being neutralised with NaOH and again fractionally distd. Also purified by distn from Raney nickel and by preparative GLC.

***n*-Octylammonium 9-anthranilate** [88020-99-9] M 351.5, m 134-135°, pK^{25} 10.65 (for octylamine). Recrystd several times from ethyl acetate.

***n*-Octylammonium hexadecanoate** [88020-97-7] M 385.7, m 52-53°. Purified by several recrystns from *n*-hexane or ethyl acetate. The solid was then washed with cold anhydrous diethyl ether, and dried *in vacuo* over P₂O₅.

***n*-Octylammonium octadecanoate** [32580-92-0] M 413.7, m 56-57°. Purified as for the hexadecanoate above.

***n*-Octylammonium tetradecanoate** [17463-35-3] M 358.6, m 46-48°. Purified as for the hexadecanoate above.

4-Octylbenzoic acid [3575-31-3] M 234.3, m 99-100°, pK^{25} 6.5 (80% aq EtOH), pK_{Est} ~4.5 (H₂O). Crystd from EtOH has m 139°; crystd from aq EtOH has m 99-100°. Forms liquid crystals.

***n*-Octyl bromide** [111-83-1] M 193.1, b 201.5°, d_4^{20} 1.118, n^{25} 1.4503. Shaken with H₂SO₄, washed with water, dried with K₂CO₃ and fractionally distd.

4-(*tert*-Octyl)phenol [140-66-9] M 206.3, m 85-86°, b 166°/20mm, pK_{Est} ~ 10.4. Crystd from *n*-hexane.

1-Octyne [629-05-0] M 110.2, b 126.2°/760mm, d_4^{20} 0.717, n^{25} 1.4159. Distd from NaBH₄ to remove peroxides.

α-Oestradiol [57-91-0] M 272.4, m 220-230°, $[\alpha]_D^{20}$ +55° (c 1, dioxane). Crystd from aq EtOH.

β-Oestradiol-3-benzoate [50-50-0] M 376.5, m 194-195°, $[\alpha]_{546}^{20}$ +70° (c 2, dioxane). Crystd from EtOH

Oleic acid [112-80-1] M 282.5, m 16°, b 360°(dec), d_4^{20} 0.891, n^{30} 1.4571, pK^{25} 6.42 (50% aq EtOH), pK_{Est} ~4.8 (H₂O). Purified by fractional crystn from its melt, followed by molecular distn at 10⁻³mm, or by conversion to its methyl ester, the free acid can be crystd from acetone at -40° to -45° (12mL/g). For purification by the use of lead and lithium salts, see Keffler and McLean [*J Soc Chem Ind (London)* 54 176T 1935]. Purification based on direct crystn from acetone is described by Brown and Shinowara [*J Am Chem Soc* 59 6 1937; *pK White J Am Chem Soc* 72 1857 1950].

Oleyl alcohol [143-28-2] M 268.5, b 182-184°/1.5mm, d_4^{20} 0.847, $n_D^{27.5}$ 1.4582. Purified by fractional crystn at -40° from acetone, then distd under vacuum.

Opianic acid (2-formyl-4,5-dimethylbenzoic acid) [519-05-1] M 210.2, m 150°, pK_1^{25} 3.07. Crystd from water.

Orcinol (5-methylresorcinol) [504-15-4] M 124.2, m 107.5°, m 59-61° (hydrate), pK_1^{20} 9.36 (9.48), pK_2^{20} 11.6 (11.20). Crystd from $CHCl_3$ /*benzene (2:3).

L-Ornithine [70-26-8] M 132.2, m 140°, $[\alpha]_D^{25}$ +16° (c 0.5, H_2O), pK_2^{20} 8.75, pK_3^{25} 10.73. Crystd from water containing 1mM EDTA (to remove metal ions).

L-Ornithine monohydrochloride [3184-13-2] M 168.6, m 233°(dec), $[\alpha]_D^{25}$ +28.3° (5M HCl). Likely impurities are citrulline, arginine and D-ornithine. Crystd from water by adding 4 volumes of EtOH and dried in a vacuum desiccator over fused $CaCl_2$.

Orotic acid (H_2O) [50887-69-9] M 174.1, m 235-346°(dec), pK_1^{25} 1.8, pK_2^{25} 9.55. Crystd from water.

Orthanilic acid (2-aminobenzenesulfonic acid) [88-21-1] M 173.2, m >300°(dec), pK_1^{25} 2.49. Crystd from aqueous soln, containing 20mL of conc HCl per L, then crystd from distilled water, and dried in a vacuum desiccator over Sicapent.

Ouabain {3-[(6-deoxy- α -L-mannopyranosyl)oxy]-1,5,11a,14,19-pentahydroxycard-20(22)-enolide} [630-60-4] M 728.8, m 180°(dec), $[\alpha]_{546}^{20}$ -30° (c 1, H_2O). Crystd from water as the octahydrate. Dried at 130°. Stored in the dark.

Oxalic acid ($2H_2O$) [6153-56-6] M 90.0, m 101.5°; [anhydrous 144-62 -7] m 189.5°, pK_1^{25} 1.08 (1.37), pK_2^{25} 3.55 (3.80). Crystd from distilled water. Dried in vacuum over H_2SO_4 . The anhydrous acid can be obtained by drying at 100° overnight.

Oxaloacetic acid [328-42-7] M 132.1, m 160°(decarboxylates), pK_1^{25} 2.22, pK_2^{25} 3.89, pK_3^{25} 13.0. Crystd from boiling ethyl acetate, or from hot acetone by addition of hot *benzene.

2-Oxoglutaric acid (2-oxopentane-1,5-dioic, α -ketoglutaric acid) [328-50-7] M 146.1, m 114°, 115-117°, (pK_{Est} see oxaloacetic acid above). Crystd repeatedly from Me_2CO /*benzene, EtOAc or ethyl propionate.

Oxalylindigo [2533-00-8] M 316.3. Recrystd twice from nitrobenzene and dried by heating *in vacuo* for several hours. [Sehanze et al. *J Am Chem Soc* 108 2646 1986.]

Oxamide [471-46-5] M 88.1, m >320°(dec). Crystd from water, ground and dried in an oven at 150°.

2-Oxazolidinone [497-25-6] M 87.1, m 89-90°, 91°, b 152°/0.4mm. Crystd from *benzene or dichloroethane.

Oxetane (1.3-trimethylene oxide) [503-30-0] M 58.1, b 45-46°/736mm, 47-49°/atm, 48°/760mm, d_4^{20} 0.892, n_D^{20} 1.395. Distd twice from sodium metal and then fractionated through a small column at atmospheric pressure, b 47.0-47.2°. Also purified by preparative gas chromatography using a 2m silica gel column. Alternatively add KOH pellets (50g for 100g of oxetane) and distil through an efficient column or a column packed with 1/4in Berl Saddles and the main portion boiling at 45-50° is collected and redistilled over fused KOH. [Noller *Org Synth Coll Vol III* 835 1955; Dittmer et al. *J Am Chem Soc* 79 4431 1957.]

Oxine Blue [3-(4-hydroxyphenyl)-3-(8-hydroxy-6-quiniliny)-1(3*H*)-isobenzofuranone] [3733-85-5] **M 369.4, m 134-135°**. Recrystd from EtOH. Dried over H₂SO₄.

Palmitic acid anhydride (hexadecanoic anhydride) [623-65-4] **M 494.9, m 63-64°, 64°, d⁸² 0.838, n⁶⁸ 1.436**. It is moisture sensitive and hydrolyses in water. Purified by refluxing with acetic anhydride for 1hr, evaporating and freeing the residue of acetic acid and anhydride by drying the residue at high vac and crystallising from pet ether at low temperature.

[2.2]-Paracyclophane (tricyclo[8.2.2.2^{4,7}]hexadeca-4,6,10,12,13,15-hexaene) [1633-22-3] **M 208.3, m 284°, 285-287°, 286-288°, 288-290°**. Purified by recrystn from AcOH. ¹H-NMR δ: 1.62 (Ar-H) and -1.71 (CH₂) [Waugh and Fessenden *J Am Chem Soc* **79** 846 1957; IR and UV: Cram et al. *J Am Chem Soc* **76** 6132 1954, Cram and Steinberg *J Am Chem Soc* **73** 5691 1951; complex with unsaturated compounds: Cram and Bauer *J Am Chem Soc* **81** 5971 1959; Syntheses: Brink *Synthesis* 807 1975, Givens et al. *J Org Chem* **44** 16087 1979, Kaplan et al. *Tetrahedron Lett* 3665 1976].

Paraffin (oil) [8012-95-1] **d 0.880, n 1.482**. Treated with fuming H₂SO₄, then washed with water and dilute aqueous NaOH, then percolated through activated silica gel.

Paraffin Wax. Melted in the presence of NaOH, washed with water until all of the base had been removed. The paraffin was allowed to solidify after each wash. Finally, 5g of paraffin was melted by heating on a water-bath, then shaken for 20-30min with 100mL of boiling water and fractionally crystd.

Parafuchsin (4,4',4''-triaminotriptyllium [triphenylmethane] carbonium ion, pararosaniline, paramagenta) [467-62-9] **M 305.4, pK 7.57 and free base has pK >13**. Dissolve in EtOH (1.16g in 30mL), filter and add aqueous NH₃ till neutral and ppte by adding H₂O giving 0.8g **m 247° dec** (sintering at 230°). Dissolve in EtOH neutralise with NH₃ add 0.1g of charcoal filter, and repeat, then add H₂O (100mL) to ppte the colourless *carbinol* dry, **m 257° dec** (sintering at 232°). [Weissberger and Theile *J Chem Soc* 148 1934.] The *carbinol* (pseudo-base) was said to have **m 232°** (186° dec), and is slightly sol in H₂O but sol in acids and EtOH [pK: Goldacre and Phillips *J Chem Soc* 172 1949]. The *perchlorate* (dark red with a green shine) has **m 300°** and explodes at 317° [Dilthey and Diaklage *J Prakt Chem* [2] 129 1931].

Paraldehyde (acetaldehyde trimer, 2,4,6-trimethyl-1,3,5-trioxane) [123-63-7] **M 132.2, m 12.5°, 124°, d 0.995, n 1.407**. Washed with water and fractionally distd.

Patulin {4-hydroxy-4*H*-furo[3.2-*c*]pyran-2(6*H*)-one} [149-29-1] **M 154.1, m 110°**. Crystd from diethyl ether or chloroform. (*Highly TOXIC*).

Pavatrine hydrochloride [548-65-2] **M 333.7, m 143-144°**. Recrystd from isopropanol, and dried over P₂O₅ under vacuum.

Pelargonic acid (nonanoic acid) [112-05-0] **M 158, m 15°, b 98.9°/1mm, 225°/760mm, pK²⁵ 4.96**. Esterified with ethylene glycol and distd. (This removes dibasic acids as undistillable residues.) The acid was regenerated by hydrolysing the ester.

Pelargonitrile (octyl cyanide) [2243-27-8] **M 139.2, m -34°, b 92°/10mm, 224°, d 0.818, n 1.4255**. Stirred with P₂O₅ (~5%), distd from it and redistd under vac. IR should have CN but no OH bands.

Pelargonyl chloride (nonanoyl chloride) [764-85-2] **M 176.7, b 88°/12mm, d 0.941, n 1.436**. Refluxed with acetyl chloride (~ 3 vols) for 1h, then distil off the AcCl followed by the nanoyl chloride under reduced pressure. It is moisture sensitive and should be stored in sealed ampules.

Penicillic acid [90-65-3] **M 158.2, m 58-64° (monohydrate), 83-84° (anhydrous, lactone)**. Crystd from water as the monohydrate, or from pet ether. Free acid is in equilibrium with the lactone.

[Previous Page](#)

Oxine Blue [3-(4-hydroxyphenyl)-3-(8-hydroxy-6-quiniliny)-1(3*H*)-isobenzofuranone] [3733-85-5] **M 369.4, m 134-135°**. Recrystd from EtOH. Dried over H₂SO₄.

Palmitic acid anhydride (hexadecanoic anhydride) [623-65-4] **M 494.9, m 63-64°, 64°, d⁸² 0.838, n⁶⁸ 1.436**. It is moisture sensitive and hydrolyses in water. Purified by refluxing with acetic anhydride for 1hr, evaporating and freeing the residue of acetic acid and anhydride by drying the residue at high vac and crystallising from pet ether at low temperature.

[2.2]-Paracyclophane (tricyclo[8.2.2.2^{4,7}]hexadeca-4,6,10,12,13,15-hexaene) [1633-22-3] **M 208.3, m 284°, 285-287°, 286-288°, 288-290°**. Purified by recrystn from AcOH. ¹H-NMR δ: 1.62 (Ar-H) and -1.71 (CH₂) [Waugh and Fessenden *J Am Chem Soc* **79** 846 1957; IR and UV: Cram et al. *J Am Chem Soc* **76** 6132 1954, Cram and Steinberg *J Am Chem Soc* **73** 5691 1951; complex with unsaturated compounds: Cram and Bauer *J Am Chem Soc* **81** 5971 1959; Syntheses: Brink *Synthesis* 807 1975, Givens et al. *J Org Chem* **44** 16087 1979, Kaplan et al. *Tetrahedron Lett* 3665 1976].

Paraffin (oil) [8012-95-1] **d 0.880, n 1.482**. Treated with fuming H₂SO₄, then washed with water and dilute aqueous NaOH, then percolated through activated silica gel.

Paraffin Wax. Melted in the presence of NaOH, washed with water until all of the base had been removed. The paraffin was allowed to solidify after each wash. Finally, 5g of paraffin was melted by heating on a water-bath, then shaken for 20-30min with 100mL of boiling water and fractionally crystd.

Parafuchsin (4,4',4''-triaminotriptyllium [triphenylmethane] carbonium ion, pararosaniline, paramagenta) [467-62-9] **M 305.4, pK 7.57 and free base has pK >13**. Dissolve in EtOH (1.16g in 30mL), filter and add aqueous NH₃ till neutral and ppte by adding H₂O giving 0.8g **m 247° dec** (sintering at 230°). Dissolve in EtOH neutralise with NH₃ add 0.1g of charcoal filter, and repeat, then add H₂O (100mL) to ppte the colourless *carbinol* dry, **m 257° dec** (sintering at 232°). [Weissberger and Theile *J Chem Soc* 148 1934.] The *carbinol* (pseudo-base) was said to have **m 232° (186° dec)**, and is slightly sol in H₂O but sol in acids and EtOH [pK: Goldacre and Phillips *J Chem Soc* 172 1949]. The *perchlorate* (dark red with a green shine) has **m 300°** and explodes at 317° [Dilthey and Diaklage *J Prakt Chem* [2] 129 1931].

Paraldehyde (acetaldehyde trimer, 2,4,6-trimethyl-1,3,5-trioxane) [123-63-7] **M 132.2, m 12.5°, 124°, d 0.995, n 1.407**. Washed with water and fractionally distd.

Patulin {4-hydroxy-4*H*-furo[3.2-*c*]pyran-2(6*H*)-one} [149-29-1] **M 154.1, m 110°**. Crystd from diethyl ether or chloroform. (*Highly TOXIC*).

Pavatrine hydrochloride [548-65-2] **M 333.7, m 143-144°**. Recrystd from isopropanol, and dried over P₂O₅ under vacuum.

Pelargonic acid (nonanoic acid) [112-05-0] **M 158, m 15°, b 98.9°/1mm, 225°/760mm, pK²⁵ 4.96**. Esterified with ethylene glycol and distd. (This removes dibasic acids as undistillable residues.) The acid was regenerated by hydrolysing the ester.

Pelargonitrile (octyl cyanide) [2243-27-8] **M 139.2, m -34°, b 92°/10mm, 224°, d 0.818, n 1.4255**. Stirred with P₂O₅ (~5%), distd from it and redistd under vac. IR should have CN but no OH bands.

Pelargonyl chloride (nonanoyl chloride) [764-85-2] **M 176.7, b 88°/12mm, d 0.941, n 1.436**. Refluxed with acetyl chloride (~ 3 vols) for 1h, then distil off the AcCl followed by the nanoyl chloride under reduced pressure. It is moisture sensitive and should be stored in sealed ampules.

Penicillic acid [90-65-3] **M 158.2, m 58-64° (monohydrate), 83-84° (anhydrous, lactone)**. Crystd from water as the monohydrate, or from pet ether. Free acid is in equilibrium with the lactone.

Pentabromoacetone [79-49-2] M 452.6, m 76°, pK 8.0 (MeOH), pK_{Est} ~4.6 (H₂O). Crystd from diethyl ether or EtOH and sublimes.

Pentabromophenol [608-71-9] M 488.7, m 229°, pK_{Est} ~ 4.5. Purified by crystn (charcoal) from toluene then from CCl₄. Dried for 2 weeks at ca 75°.

1-Pentacene [135-48-8] M 278.4, m 300°. Crystd from *benzene.

Pentachloroethane (pentalin) [76-01-7] M 202.3, b 69°/37mm, 152.2°/64mm, 162.0°, d 1.678, n¹⁵ 1.50542. Usual impurities include trichloroethylene. Partially decomposes if distd at atmospheric pressure. Drying with CaO, KOH or sodium is unsatisfactory because of the elimination of the elements of HCl. It can be purified by steam distn, or by washing with conc H₂SO₄, water, and then aqueous K₂CO₃, drying with solid K₂CO₃ or CaSO₄, and fractionally distd under reduced pressure.

Pentachloronitrobenzene [82-68-8] M 295.3, m 146°. Crystd from EtOH.

Pentachlorophenol [87-86-5] M 266.3, m 190-191°, pK²⁵ 4.8. Twice crystd from toluene/EtOH. Sublimed *in vacuo*.

Pentachloropyridine [2176-62-7] M 251.3, m 122-124°, 123°, 124°, 124-125°, 125-126°, b 279-280°/atm, pK -6.02 (aq H₂SO₄). Purified by recryst from EtOH or aqueous EtOH. It sublimes at 150°/3mm. [den Hertog et al. *Recl Trav Chim Pays-Bas* 69 673 1950; Schikh et al. *Chem Ber* 69 2604 1936.]

Pentachlorothiophenol [133-49-3] M 282.4, m between 228° and 235°, pK_{Est} ~1.1. Crystd from *benzene.

Pentadecafluoro octanoic acid (perfluorocaprylic acid) [335-67-1] M 414.1, m 54.9-55.6°, b 189°/736mm, pK_{Est} <0. Recrystd from CCl₄ and toluene, and can be distd. It forms micelles in H₂O and the solubility is 1% in H₂O. [Bernett and Zisman *J Phys Chem* 63 1911 1959; IR: Bro and Sperati *J Polym Sci* 38 289 1959.]

Pentadecanoic acid [1002-84-2] M 242.4, m 51-53°, 80°, b 158°/1mm, 257°/760mm, d⁸⁰ 0.8424, pK_{Est} ~5.0. Cryst from Et₂O and distd. Very hygroscopic. See purification of palmitic acid.

Pentadecanolide (1-oxacyclohexadecan-2-one, pentadecanoic- ω -lactone, 15-hydroxypentadecanoic lactone, exaltolide, Tibetolide) [106-02-5] M 240.4, m 34-36°, 37-37.5°, 37-38°, b 102-103°/0.03mm, 112-114°/0.2mm, 137°/2mm, 169°/10-11mm, d₄⁴⁰ 0.9401. It has been recrystd from MeOH (4 parts) at -15°. [Hundiecker and Erlbach *Chem Ber* 80 135 1947; Galli and Mandolini *Org Synth* 58 100 1978; Demole and Enggist *Helv Chim Acta* 11 2318 1978.]

Penta-1,3-diene [*cis*: 1574-41-0; *trans*: 2004-70-8] M 68.1, b 42°, d 0.680, n 1.4316. Distd from NaBH₄. Purified by preparative gas chromatography. [Reimann et al. *J Am Chem Soc* 108 5527 1986.]

Penta-1,4-diene [591-93-5] M 68.1, b 25.8-26.2°/756mm, d 0.645, n 1.3890. Distd from NaBH₄. Purified by preparative gas chromatography. [Reimann et al. *J Am Chem Soc* 108 5527 1986.]

Pentaerythritol [115-77-5] M 136.2, m 260.5°. Refluxed with an equal volume of MeOH, then cooled and the ppte dried at 90°. Crystd from dil aq HCl. Sublimed under vacuum at 200°.

Pentaerythritol tetraacetate [597-71-7] M 304.3, m 78-79°. Crystd from hot water, then leached with cold water until the odour of acetic acid was no longer detectable.

Pentaerythrityl laurate [13057-50-6] M 864.6, m 50°. Crystd from pet ether.

Pentaerythritol tetranitrate. [78-11-5] M 316.2, m 140.1°. Crystd from acetone or acetone/EtOH. **EXPLOSIVE.**

Pentaethylenehexamine [4067-16-7] M 232.4, d 0.950, n_D^{20} 1.510, pK_1 1.2, pK_2 2.7, pK_3 4.3, pK_4 7.8, pK_5 9.1, pK_6 9.9 (all estimated). Fractionally distd twice at 10-20mm, the fraction boiling at 220-250° being collected. Its soln in MeOH (40mL in 250mL) was cooled in an ice-bath and conc HCl was added dropwise with stirring. About 50mL was added, and the ppted hydrochloride was filtered off, washed with acetone and diethyl ether, then dried in a vacuum desiccator. [Jonassen et al. *J Am Chem Soc* 79 4279 1957.]

Pentafluorobenzene [363-72-4] M 168.1, b 85°/atm, 85-86°/atm, 88-89°/atm, d_4^{20} 1.524, n_D^{20} 1.3931. Purified by distn and by gas chromatography. IR film: 1535 and 1512 cm^{-1} (*benzene ring). [UV: Stephen and Tatlow *Chem Ind (London)* 821 1957 ; Nield et al. *J Chem Soc* 166 1959.] See triethylenepentamine

2,3,4,5,6-Pentafluorobenzoic acid [602-94-8] M 212.1, m 101-103°, 103-104°, 104-105°, 106-107°, pK^{25} 1.75. Dissolve in Et₂O, treat with charcoal, filter, dry (CaSO₄), filter, evaporate and recrystallise residue from pet ether (b 90-100°) after adding a little toluene to give large colourless plates. UV (H₂O): λ_{max} 265nm (ϵ 761). The *S*-benzylisothiuronium salt has m 187° after recrystn from H₂O. [McBee and Rapkin *J Am Chem Soc* 73 1366 1951; Nield et al. *J Chem Soc* 166, 170 1959.]

O-(2,3,4,5,6-Pentafluorobenzyl)hydroxylamine hydrochloride (PFBHA) [57981-02-9] M 249.6, m 215°, 215-216°, pK_{Est} ~1.1. Recrystd from EtOH to form colourless leaflets. Drying the compound at high vacuum and elevated temperature will result in losses by sublimation. [Youngdale *J Pharm Sci* 65 625 1976; Wehner and Handke *J Chromatog* 177 237 1979; Nambara et al. give incorrect m as 115-116° *J Chromatogr* 114 81 1975.]

2,3,4,5,6-Pentafluorophenol [771-61-9] M 184.1, m 33-35°, 38.5-39.5°, b 72-74°/48mm, 142-144°/atm, 143°/atm, n_D^{20} 1.4270 (liquid prep), pK^{25} 5.53. A *hygroscopic* low melting solid not freely soluble in H₂O. Purified by distn, preferably in a vacuum [Forbes et al. *J Chem Soc* 2019 1959; IR and pK_a : Birchall and Haszeldine *J Chem Soc* 13 1959]. IR film: 3600 (OH) and 1575 (fluoroaromatic breathing) cm^{-1} . The *benzoyl* derivative has m 74-75°, *3,4-dinitrobenzoyl* derivative has m 107°, the *tosylate* has m 64-65° (from EtOH) and the *K salt* crystallises from Me₂CO, m 242° dec, with 1H₂O salt the m is 248° dec and the 2H₂O salt has m 245° dec.

1-(Pentafluorophenyl)ethanol [*R*-(+)- 104371-21-3; *S*-(-)- 104371-20-2] M 212.1, m 41-42°, 42°, 42.5-43°, $[\alpha]_{546}^{20}$ (+) and (-) 9°, $[\alpha]_D^{20}$ (+) and (-) 7.5° (c 1, *n*-pentane). Recrystd from *n*-pentane at -40° and vacuum sublimed at room temp at 0.3mm (use ice cooled cold finger). It has also been purified by column chromatography through Kieselgel 60 (0.063-0.2mm mesh, Merck), eluted with EtOAc-*n*-hexane (1:5), then recrystd from *n*-pentane and vacuum sublimed. It has R_F on Kieselgel 60 F₂₅₄ TLC foil and eluting with EtOAc-*n*-hexane (1:5). [Meese *Justus Liebigs Ann Chem* 2004 1986.] The *racemate* [75853-08-6] has m 32-34°, b 77-79°/8mm, 80-82°/37mm, n_D^{20} 1.4426 and the *3,4-dinitrobenzoate* has m 83° [Nield et al. *J Chem Soc* 166 1959].

2,2,3,3,3-Pentafluoropropan-1-ol [422-05-9] M 150.1, b 80°, d 1.507, n 1.288, pK^{25} 12.74. Shaken with alumina for 24h, dried with anhydrous K₂CO₃, and distd, collecting the middle fraction (b 80-81°) and redistilling.

Pentafluoropyridine [700-16-3] M 169.1, m -41.5°, b 83.5°, 84°, 83-85°, d_4^{20} 1.609, n_D^{20} 1.3818, pK_{Est} ~<0. Distd through a concentric tube column; has λ max in cyclohexane at 256.8nm. [Chambers et al. *J Chem Soc* 3573 1964]; ¹⁹F NMR: Bell et al. *J Fluorine Chem* 1 51 1971.] The *hexafluoroantimonate* has m 98-102° dec.

Pentamethylbenzene [700-12-9] M 148.3, m 53.5-55.1°. Successively crystd from absolute EtOH, toluene and MeOH, and dried under vacuum. [Rader and Smith *J Am Chem Soc* **84** 1443 1962.] It has also been crystd from *benzene or aqueous EtOH, and sublimed.

n-Pentane [109-66-0] M 72.2, b 36.1°, d 0.626, n²⁵ 1.35472. Stirred with successive portions of conc H₂SO₄ until there was no further coloration during 12h, then with 0.5N KMnO₄ in 3M H₂SO₄ for 12h, washed with water and aqueous NaHCO₃. Dried with MgSO₄ or Na₂SO₄, then P₂O₅ and fractionally distd through a column packed with glass helices. It was also purified by passage through a column of silica gel, followed by distn and storage with sodium hydride. An alternative purification is by azeotropic distn with MeOH, which is subsequently washed out from the distillate (using water), followed by drying and distn. For removal of carbonyl-containing impurities, see *n-heptane*. Also purified by fractional freezing (ca 40%) on a copper coil through which cold air was passed, then washed with conc H₂SO₄ and fractionally distd.

Pentane-1-thiol [110-66-7] M 104.2, m -76°, b 122.9°/697.5mm, d²⁵ 0.8375, pK_{Est} ~10.1. Dissolved in aqueous 20% NaOH, then extracted with a small amount of diethyl ether. The soln was acidified slightly with 15% H₂SO₄, and the thiol was distd out, dried with CaSO₄ or CaCl₂, and fractionally distd under nitrogen. [Ellis and Reid *J Am Chem Soc* **54** 1674 1932.]

Pentan-2-ol [6032-29-7] M 88.2, b 119.9°, d 0.810, n 1.41787, n²⁵ 1.4052. Refluxed with CaO, distd, refluxed with magnesium and again fractionally distd.

Pentan-3-ol [584-02-1] M 88.2, b 116.2°, d 0.819, n²⁵ 1.4072. Refluxed with CaO, distd, refluxed with magnesium and again fractionally distd.

Pentan-3-one (diethyl ketone) [96-22-0] M 86.1, b 102.1°, d 0.8099, n 1.392. See diethyl ketone on p. 204.

Pentaquine monophosphate [5428-64-8] M 395.6, m 189-190°, pK⁷⁰ 8.22. Crystd from 95% EtOH.

Pent-2-ene (mixed isomers) [109-68-2] M 70.1, b 36.4°, d 0.650, n 1.38003, n²⁵ 1.3839. Refluxed with sodium wire, then fractionally distd twice through a Fenske (glass helices packing) column.

cis-Pent-2-ene [627-20-3] M 70.1, b 37.1°, d 0.657, n²⁵ 1.3798. Dried with sodium wire and fractionally distd, or purified by azeotropic distn with MeOH, followed by washing out the MeOH with water, drying and distilling. Also purified by chromatography through silica gel and alumina [Klassen and Ross *J Phys Chem* **91** 3668 1987].

trans-Pent-2-ene [646-04-8] M 70.1, b 36.5°, d 0.6482, n 1.3793. It was treated as above and washed with water, dried over anhydrous Na₂CO₃, and fractionally distd. The middle cut was purified by two passes of fractional melting.

Pentobarbital (5-ethyl-5-1'-methylbutyl barbituric acid, Nembutal) [76-74-4] M 226.4, m ~127°(dec), pK_{Est(1)} ~ 8.0, pK_{Est(2)} ~12.7. Soln of the sodium salt in 10% HCl was prepared and the acid was extracted by addition of ether. Then purified by repeated crystn from CHCl₃. [Bucket and Sandorfy *J Phys Chem* **88** 3274 1984.]

Pentyl acetate (n-amyl acetate) [628-63-7] M 130.2, b 147-149°/atm, 149.55°, 149.2°/atm, d₄²⁰ 0.8753, n_D²⁰ 1.4028. Purified by repeated fractional distn through an efficient column or spinning band column. [Timmermann and Hennant-Roland *J Chim Phys* **52** 223 1955; Mumford and Phillips *J Chem Soc* **75** 1950; ¹H NMR: Crawford and Foster *Can J Phys* **34** 653 1956.]

Pent-2-yne [627-21-4] M 68.1, b 26°/2.4mm, d 0.710, n²⁵ 1.4005. Stood with, then distd at low pressure from, sodium or NaBH₄.

Perbenzoic acid [93-59-4] M 138.1, m 41-43°, b 97-110°/13-15mm, $pK_{\text{Est}} \sim 7.7$. Crystd from *benzene or pet ether. Readily sublimed and is steam volatile. Sol in CHCl_3 , CCl_4 and Et_2O . [*Org Synth Coll Vol I* (2nd Edn) 431 1948.] **EXPLOSIVE**.

Perchlorobutadiene [87-68-3] M 260.8, b 144.1°/100mm, 210-212°/760mm, d 1.683, n 1.5556. Washed with four or five 1/10th volumes of MeOH (or until the yellow colour has been extracted), then stirred for 2h with H_2SO_4 , washed with distilled water until neutral and filtered through a column of P_2O_5 . Distd under reduced pressure through a packed column. [Rytner and Bauer *J Am Chem Soc* 82 298 1960.]

Perfluorobutyric acid [375-22-4] M 214.0, m -17.5°, b 120°/735mm, d 1.651, n^{16} 1.295, pK^{25} -0.17. Fractionally distd twice in an Oldershaw column with an automatic vapour-dividing head, the first distn in the presence of conc H_2SO_4 as a drying agent.

Perfluorocyclobutane [115-25-3] M 200.0, m -40°, b -5°, d^{-20} 1.654, d^0 1.72. Purified by trap-to-trap distn, retaining the middle portion.

Perfluorocyclohexane [355-68-0] M 300.1, m 51° (sublimes), b 52°. Extracted repeatedly with MeOH, then passed through a column of silica gel (previously activated by heating at 250°).

Perfluoro-1,3-dimethylcyclohexane [335-27-3] M 400.1, b 101°, d 1.829, n 1.300. Fractionally distd, then 35mL was sealed with about 7g KOH pellets in a borosilicate glass ampoule and heated at 135° for 48h. The ampoule was cooled and opened, and the liquid was resealed with fresh KOH in another ampoule and heated as before. This process was continued until no further decomposition was observed. The substance was then washed with distilled water, dried (CaSO_4) and distd. [Grafstein *Anal Chem* 26 523 1954.]

Perfluoroheptane [335-57-9] M 388.1, b 99-101°, d^{25} 1.7200. Purified as for *perfluorodimethylhexane*. Other procedures include shaking with H_2SO_4 , washing with water, drying with P_2O_5 for 48h and fractionally distilling. Alternatively, it has been refluxed for 24h with saturated acid KMnO_4 (to oxidise and remove hydrocarbons), then neutralised, steam distd, dried with P_2O_5 , and passed slowly through a column of dry silica gel. It has been purified by fractional crystn, using partial freezing.

Perfluoro-*n*-hexane [355-42-0] M 338.1, m -4°, b 58-60°, d 1.684. Purified by fractional freezing. The methods described for *perfluoroheptane* should be applicable here.

Perfluoro(methylcyclohexane) [355-02-2] M 350.1, b 76.3°, d^{25} 1.7878. Refluxed for 24h with saturated acid KMnO_4 (to oxidise and remove hydrocarbons), then neutralised, steam distd, dried with P_2O_5 and passed slowly through a column of dry silica gel. [Glew and Reeves *J Phys Chem* 60 615 1956.] Also purified by percolation through a 1m neutral activated alumina column, and ^1H -impurities checked by NMR.

Perfluorononane [375-96-2] M 488.1, b 126-127°, d 1.80, n 1.275. Purified as for *perfluorodimethylcyclohexane*.

Perfluoropropyl iodide [754-34-7] M 295.9, b 41°, d 2.13, n 1.339. Purified by fractional distn.

Perfluorotributylamine (heptacosafuorotributylamine) [311-89-7] M 671.1, b 177.6°/760mm, d 1.881, n 1.291, $pK_{\text{Est}} \sim 5.0$. Purified as for *perfluorodimethylcyclopropane*, see also *perfluorotripropylamine* [Hazardine *J Chem Soc* 102 1951].

Perfluorotripropylamine (heneicosafuorotripropylamine) [338-83-0] M 521.1, b 130°/atm, 129.5-130.5°/atm, d 1.822, n 1.279, $pK_{\text{Est}} \sim 5.6$. Purified as for *perfluorodimethylcyclopropane*. [Hazardine *J Chem Soc* 102 1951, for azeotropes see Simons and Linevsky *J Am Chem Soc* 74 4750 1972.] **IRRITANT**.

Pericyazine [10-{3-(4-hydroxy-1-piperidinyl)-propyl}-10*H*-phenothiazine-2-carbonitrile] [2622-26-6] M 365.4, m 116-117°. Recrystd from a saturated soln in cyclohexane. Antipsychotic and is a reagent for Pd and Rh.

Perylene [198-55-0] M 252.3, m 273-274°. Purified by silica-gel chromatography of its recrystd picrate. [Ware *J Am Chem Soc* 83 4374 1961.] Crystd from *benzene, toluene or EtOH and sublimed in a flow of oxygen-free nitrogen. [Gorman et al. *J Am Chem Soc* 107 4404 1985; Johansson et al. *J Am Chem Soc* 109 7374 1987.]

Petroleum ether [8032-32-4] b 35-60°, d 0.640, n 1.363. Shaken several times with conc H₂SO₄, then 10% H₂SO₄ and conc KMnO₄ (to remove unsatd, including aromatic, hydrocarbons) until the permanganate colour persists. Washed with water, aqueous Na₂CO₃ and again with water. Dried with CaCl₂ or Na₂SO₄, and distd. It can be dried further using CaH₂ or sodium wire. Passage through a column of activated alumina, or treatment with CaH₂ or sodium, removes peroxides. For the elimination of carbonyl-containing impurities without using permanganate, see *n-heptane*. These procedures could be used for all fractions of pet ethers.

Rapid purification: Pass through an alumina column and fractionally distilling, collecting the desired boiling fraction.

R(-)- α -Phellandrene (*p-menta-1,5-diene*) [4221-98-1] M 136.2, b 61°/11mm, 175-176°/760mm, d 0.838, n 1.471. Purified by gas chromatography on an Apiezon column.

Phenacylamine hydrochloride [5468-37-1] M 171.6, m 194°(dec). See 2-aminoacetophenone hydrochloride on p. 103.

Phenanthrene [85-01-8] M 178.2, m 98°. Likely contaminants include, anthracene, carbazole, fluorene and other polycyclic hydrocarbons. Purified by distn from sodium, boiling with maleic anhydride in xylene, crystn from acetic acid, sublimation and zone melting. Has also been recrystd repeatedly from EtOH, *benzene or pet ether (b 60-70°), with subsequent drying under vacuum over P₂O₅ in an Abderhalden pistol. Feldman, Pantages and Orchin [*J Am Chem Soc* 73 4341 1951] separated from most of the anthracene impurity by refluxing phenanthrene (671g) with maleic anhydride (194g) in xylene (1.25L) under nitrogen for 22h, then filtered. The filtrate was extracted with aqueous 10% NaOH, the organic phase was separated, and the solvent was evaporated. The residue, after stirring for 2h with 7g of sodium, was vacuum distd, then recrystd twice from 30% *benzene in EtOH, then dissolved in hot glacial acetic acid (2.2mL/g), slowly adding an aqueous soln of CrO₃ (60g in 72mL H₂O added to 2.2L of acetic acid), followed by slow addition of conc H₂SO₄ (30mL). The mixture was refluxed for 15min, diluted with an equal volume of water and cooled. The ppt was filtered off, washed with water, dried and distd, then recrystd twice from EtOH. Further purification is possible by chromatography from CHCl₃ soln on activated alumina, with *benzene as eluent, and by zone refining.

Phenanthrene-9-aldehyde [4707-71-5] M 206.3, m 102-103°, pK -6.39 (aq H₂SO₄). Crystd from EtOH and sublimed at 95-98°/0.07mm.

9,10-Phenanthrenequinone [84-11-7] M 208.2, m 208°, pK -7.1 (aq H₂SO₄). Crystd from dioxane or 95% EtOH and dried under vacuum.

Phenanthridine [229-87-8] M 179.2, m 106.5°, 108-109°, b 350°, pK²⁰ 4.61 (4.48). Purified via the HgCl₂ addition compound formed when phenanthridine (20g) in 1:1 HCl (100mL) was added to aq HgCl₂ (60g in 3L), and the mixture was heated to boiling. Conc HCl was then added until all of the solid had dissolved. The compound separated on cooling, and was dec with aq NaOH (ca 5M). Phenanthridine was extracted with Et₂O and crystd from pet ether (b 80-100°) or EtOAc. [Cumper et al. *J Chem Soc* 45218 1962.] Also purified by zone melting; sublimes in vac. [Slough and Ubbelohde *J Chem Soc* 911 1957.] See p. 124.

1,10-Phenanthroline (*o-phenanthroline*) [66-71-7 (*anhydr*); 5144-89-8 (*H₂O*)] M 198.2, m 98-101°, 108-110° (*hydr*), 118° (*anhydr*), b >300°, pK₁²⁵ -0.7 (aq HClO₄), pK₂²⁵ 4.86 (4.96). Crystd as its *picrate* (m 191°) from EtOH, then the free base was liberated, dried at 78°/8mm over P₂O₅ and crystd from pet ether (b 80-100°). [Cumper, Ginman and Vogel *J Chem Soc* 1188 1962.] It can be purified by zone melting. Also crystd from hexane, *benzene/pet ether (b 40-60°) or sodium-dried *benzene, dried and stored over H₂SO₄. The monohydrate is obtained by crystn from aqueous EtOH or ethyl acetate. It has been crystd from H₂O (300 parts) to give the *monohydrate* m 102-103° and sublimes at 10⁻³mm [Fielding and LeFevre *J*

Chem Soc 1811 1951.] The *anhydrous* compound has *m* 118° (after drying at high vacuum at 80°), also after recrystn from pet ether or *C₆H₆ (70 parts) and drying at 78°/8mm. [UV: Badger et al. *J Chem Soc* 3199 1951.] It has a pKa in H₂O of 4.857 (25°) or 5.02 (20°) and 4.27 in 50% aq EtOH (20°) [Albert et al. *J Chem Soc* 2240 1948].

1,10-Phenanthroline hydrochloride (o-phenanthroline hydrochloride) [3829-86-5] *M* 243.7, *m* 212-219°. It crystallises from 95% EtOH, *m* 212-219° as the *monohydrate*, the *half hydrate* has *m* 217°. The *3HCl* has *m* 143-145° (sinters at 128°) [Thevenet et al. *Acta Cryst Sect B* 33 2526 1977].

4,7-Phenanthroline-5,6-dione [84-12-8] *M* 210.2, *m* 295°(dec). Crystd from MeOH.

Phenazine [92-82-0] *M* 180.2, *m* 171°, *pK*₁²⁰ -4.9 (aq H₂SO₄), *pK*₂²⁰ 1.21. Crystd from EtOH, CHCl₃ or ethyl acetate, after pre-treatment with activated charcoal. It can be sublimed *in vacuo*, and zone refined.

Phenazine methosulfate [299-11-6] *M* 306.3. See 5-methylphenazinium methyl sulfate on p. 547 in Chapter 6.

Phenethylamine [64-04-0] *M* 121.2, *b* 87°/13mm, *d* 0.962, *n* 1.535, *pK*²⁵ 9.88. Distd from CaH₂, under reduced pressure, just before use.

Phenethyl bromide [103-63-9] *M* 185.1, *b* 92°/11mm, *d* 1.368, *n* 1.557. Washed with conc H₂SO₄, water, aq 10% Na₂CO₃ and water again, then dried with CaCl₂ and fractionally distd just before use.

N-2-Phenethyl urea [2158-04-5] *M* 164.2, *m* 173-174°. Crystd from water.

Phenetole [103-73-1] *M* 122.2, *b* 60°/9mm, 77.5°/31mm, 170.0°/760mm, *d* 0.967, *n* 1.50735, *n*⁴⁶ 1.50485. Small quantities of phenol can be removed by shaking with NaOH, but this is not a very likely contaminant of commercial material. Fractional distn from sodium, at low pressures, probably gives adequate purification. It can be dissolved in diethyl ether and washed with 10% NaOH (to remove phenols), then water. The ethereal soln was evaporated and the phenetole fractionally distd under vacuum.

Phenocoll hydrochloride (4-ethoxyaniline, p-phenetidine HCl) [536-10-6] *M* 230.7, *m* 234°, *pK*²⁸ 5.20. Crystd from water. Sublimes *in vacuo*.

Phenol [108-95-2] *M* 94.1, *m* 40.9°, *b* 85.5-86.0°/20mm, 180.8°/760mm, *d* 1.06, *n*⁴¹ 1.54178, *n*⁴⁶ 1.53957, *pK*²⁵ 9.86 (10.02). Steam was passed through a boiling soln containing 1mole of phenol and 1.5-2.0moles of NaOH in 5L of H₂O until all non-acidic material had distd. The residue was cooled, acidified with 20% (v/v) H₂SO₄, and the phenol was separated, dried with CaSO₄ and fractionally distd under reduced pressure. It was then fractionally crystd several times from its melt [Andon et al. *J Chem Soc* 5246 1960]. Purification *via* the benzoate has been used by Berliner, Berliner and Nelidow [*J Am Chem Soc* 76 507 1954]. The benzoate was crystd from 95% EtOH, then hydrolysed to the free phenol by refluxing with two equivalents of KOH in aq EtOH until the soln became homogeneous. It was acidified with HCl and extracted with diethyl ether. The ether layer was freed from benzoic acid by thorough extraction with aqueous NaHCO₃, and, after drying and removing the ether, the phenol was distd.

Phenol has also been crystd from a 75% w/w soln in water by cooling to 11° and seeding with a crystal of the hydrate. The crystals were centrifuged off, rinsed with cold water (0-2°) satd with phenol, and dried. It can be crystd from pet ether [Berasconi and Paschalis *J Am Chem Soc* 108 2969 1986].

Draper and Pollard [*Science* 109 448 1949] added 12% water, 0.1% aluminium (can also use zinc), and 0.05% NaHCO₃ to phenol, and distd at atmospheric pressure until the azeotrope was removed. The phenol was then distd at 25mm. Phenol has also been dried by distn from the *benzene soln to remove the water-*benzene azeotrope and the excess *benzene, followed by distn of the phenol at reduced pressure under nitrogen. Processes such as this are probably adequate for analytical grade phenol which has as its main impurity water. Phenol has also been crystd from pet ether/*benzene or pet ether (*b* 40-60°). Purified material is stored in a vacuum desiccator over P₂O₅ or CaSO₄.

Phenol-2,4-disulfonic acid [96-77-5] M 254.2, $pK_1 < 1$, $pK_2 < 1$, $pK_3 \sim 8.3$. Crystd from EtOH/diethyl ether.

Phenolphthalein [77-09-8] M 319.2, m 263°, $pK_{Est(1)} \sim 4.2$, $pK_{Est(2)} \sim 9.8$. Dissolved in EtOH (7mL/g), then diluted with eight volumes of cold water. Filtered. Heated on a water-bath to remove most of the alcohol and the pptd phenolphthalein was filtered off and dried under vacuum.

Phenolphthalol [81-92-5] M 306.3, m 201-202°, $pK_{Est} \sim 9.8$. Crystd from aqueous EtOH.

Phenosafranin (3,7-diamino-5-phenylphenazinium chloride) [81-93-6] M 322.8, m $> 300^\circ$, λ_{max} 530nm (H₂O). Crystd from dilute HCl.

Phenothiazine [92-84-2] M 199.3, m 184-185°. Crystd from *benzene or toluene (charcoal) after boiling for 10min under reflux. Filtered on a suction filter. Dried in an oven at 100°, then in a vacuum desiccator over paraffin chips. Also twice recrystd from water and dried in an oven at 100° for 8-10h.

Phenoxazine [135-67-1] M 199.2, m 156°, 156-158°, 158-159°, b 215°/4mm. Crystd from EtOH and sublimed *in vacuo*. If too impure then extract in a Soxhlet using toluene. Evaporate the solvent and dissolve residue (*ca* 100g) in *C₆H₆ (1L) **CARCINOGEN**, use an efficient fume cupboard) and chromatograph through an Al₂O₃ column (50 x 450 mm). The eluent (*ca* 3L) is evaporated to *ca* 150mL and cooled when *ca* 103g of phenoxazine m 149-153° is obtained. Sublimation yields platelets m 158-159°. It forms a green *picrate* m 141.5-142°. [Gilman and Moore *J Am Chem Soc* 79 3485 1957; Müller et al. *J Org Chem* 24 37 1959.]

Phenoxyacetic acid [122-59-8] M 152.2, m 98-99°, $pK^{25} 3.18$. Crystd from water or aq EtOH.

Phenoxyacetyl chloride [701-99-5] M 170.6, b 112°/10mm, 102°/16mm, 225-226°/atm, $d_4^{20} 1.235$, $n_D^{20} 1.534$. If it has no OH band in the IR then distil in a vacuum, taking precautions for the moisture-sensitive compound. If it contains free acid (due to hydrolysis, OH bands in the IR) then add an equal volume of redistilled SOCl₂, reflux for 2-3h, evaporate and distil the residue in a vacuum as before. The *amide* has m 101°. [McElvain and Carney *J Am Chem Soc* 68 2592 1946.]

4-Phenoxyaniline [139-59-3] M 185.2, m 95°, $pK^{20} 4.44$ (50% aq EtOH). Crystd from water.

Phenoxybenzamine [*N*-(2-chloroethyl)-*N*-(1-methyl-2-phenoxyethyl)benzylamine] [59-96-1] M 303.5, m 38-40°, **hydrochloride** [63-92-3] M 340.0, m 137.5-140°, $pK_{Est} \sim 4.2$. The free base is crystd from pet ether and the *HCl* is crystd from EtOH/diethyl ether.

2-Phenoxybenzoic acid [2243-42-7] M 214.2, m 113°, b 355°/760mm, $pK^{15} 3.53$. Crystd from aqueous EtOH.

3-Phenoxybenzoic acid [3739-38-6] M 214.2, m 145°, $pK 3.59$. Crystd from aqueous EtOH.

Phenoxybutyric acid [6303-58-8] M 180.2, m 64°, 65-66°, 82-83°, 99°, b 180-185°/12mm, $pK 3.17$. It has been purified by recrystn from pet ether, *C₆H₆, Et₂O-pet ether, EtOH and from H₂O. It can be steam distd or distd in a good vac. [UV: Ramart-Lucas and Hoch *Bull Soc Chim Fr* [4] 51 824 1932; Dann and Arndt *Justus Liebigs Ann Chem* 587 38 1954.] The *acid chloride* has b 154-156°/20mm [Hamford and Adams *J Am Chem Soc* 57 921 1935]; and the *amide* crystallises from *C₆H₆ as needles m 113°.

2-Phenoxypropionic acid (lactic acid *O*-phenylether) [940-31-8] M 166.2, m 115-116°, b 105-106°/5mm, 265-266°/758mm, $pK 3.11$. Crystd from water.

Phensuximide (*N*-methyl-2-phenylsuccinimide) [86-34-0] M 189.2, m 71-73°. Crystd from hot 95% EtOH.

Phenylacetamide [103-81-1] M 135.2, m 158.5°. Crystd repeatedly from absolute EtOH. Dried under vacuum over P₂O₅.

Phenyl acetate [122-79-2] M 136.2, b 78°/10mm, d 1.079, n²² 1.5039. Freed from phenol and acetic acid by washing (either directly or as a soln in pentane) with aqueous 5% Na₂CO₃, then with saturated aqueous CaCl₂, drying with CaSO₄ or Na₂SO₄, and fractional distn at reduced pressure.

Phenylacetic acid [103-82-2] M 136.2, m 76-77°, b 140-150°/20mm, pK₁ -7.59 (aq H₂SO₄), pK₂ 4.31. Crystd from pet ether (b 40-60°), isopropyl alcohol, aq 50% EtOH or hot water. Dried under vac. It can be distd under reduced pressure.

Phenylacetone (1-phenylpropan-2-one) [103-79-9] M 134.2, b 69-71°/3mm, d 1.00, n 1.516. Converted to the semicarbazone and crystd three times from EtOH (m 186-187°). The semicarbazone was hydrolysed with 10% phosphoric acid and the ketone was distd. [Kumler, Strait and Alpen *J Am Chem Soc* 72 1463 1950.]

4'-Phenylacetophenone [92-91-1] M 196.3, m 120.3-121.2°, b 196-210°/18mm, 325-327°/760mm. Crystd from EtOH or acetone. Can also be distd under reduced or atmospheric pressure.

Phenylacetylene [536-74-3] M 102.1, b 75°/80mm, d 0.930, n²⁵ 1.5463, pK -19. Distd through a spinning band column. Should be filtered through a short column of alumina before use [Collman et al. *J Am Chem Soc* 108 2988 1986; for pK see Brandsma *Preparative Acetylenic Chemistry*, 1st Edn Elsevier 1971, p. 15, ISBN 0444409475].

dl-Phenylalanine [150-30-1] M 165.2, m 162°, pK₁²⁵ 2.58, pK₂²⁵ 9.24. Crystd from water and dried under vacuum over P₂O₅.

L-Phenylalanine [63-91-2] M 165.2, m 280°(dec), [α]_D²⁵ -34.0° (c 2, H₂O). Likely impurities are leucine, valine, methionine and tyrosine. Crystd from water by adding 4 volumes of EtOH. Dried under vac over P₂O₅. Also crystd from satd refluxing aq solns at neutral pH, or 1:1 (v/v) EtOH/water soln, or conc HCl.

Phenylalaninol (2-amino-3-phenylpropan-1-ol) [*R*-(+)- 5267-64-1; *S*-(-)- 3182-95-4] M 151.2, m 91-92°, 91.5°, 92-94°, b 80°/11mm (Kügelrohr), [α]_D²⁰ (+) and (-) 28°, [α]_D²⁰⁻²⁵ (+) and (-) 23-28.7° (c 1-5, EtOH), pK_{Est} ~9.3. It can be recrystd from Et₂O, *C₆H₆-pet ether (b 40-60°) or toluene and distd in a vacuum. Has been purified by dissolving in Et₂O, drying over K₂CO₃, filtering, evaporating to a small volume, cooling in ice and collecting the plates. Store in the presence of KOH (i.e. CO₂-free atm). [Karrer and Ehrhardt *Helv Chim Acta* 34 3203 1951; Oeda *Bull Chem Soc Jpn* 13 465 1938.] The *picrate* has m 141-141.5° (from EtOH-pet ether). The *hydrogen oxalate* has m 177°, 161-162° [Hunt and McHale *J Chem Soc* 2073 1957]. The *racemate* has m 87-88° from *C₆H₆-pet ether (75-77° from Et₂O), and the *hydrochloride* has m 139-141° [Fodor et al. *J Chem Soc* 1858 1951].

3-Phenylallyl chloride (cinnamyl chloride) [*E*: 18685-01-3; *Z*: 39199-93-4] M 152.6, b 92-93°/3mm. Distd under vacuum three times from K₂CO₃.

Phenyl 4-aminosalicylate [133-11-9] M 229.2, m 153°, pK_{Est(1)} ~2.0 (NH₂), pK_{Est(2)} ~9.7 (OH). Crystd from isopropanol.

4-Phenylanisole (4-methoxybiphenyl) [361-37-6] M 184.2, m 89.9-90.1°. Crystd from *benzene/pet ether. Dried under vacuum in an Abderhalden pistol.

9-Phenylanthracene [602-55-1] M 254.3, m 153-154°, b 417°. Chromatographed on alumina in *benzene and crystd from acetic acid.

***N*-Phenylanthranilic acid** [91-40-7] M 213.2, m 182-183°, pK₁²⁵ -1.28 (aq H₂SO₄), pK₂²⁵ 3.86 (CO₂H). Crystd from EtOH (5mL/g) or acetic acid (2mL/g) by adding hot water (1mL/g).

2-Phenyl-1-azaindolizine [56983-95-0] M 194.2, m 140°, pK_{Est} ~1.9. Crystd from EtOH or *benzene/pet ether.

p-Phenylazobenzoyl chloride [104-24-5] M 244.7, m 93°. Crystd from pet ether (b 60-80°).

4-Phenylazo-1-naphthylamine [131-22-6] M 247.3, m 125-125.5°. Crystd from cyclohexane or aq EtOH. [Brode et al. *J Am Chem Soc* 74 4641 1952.]

1-Phenylazo-2-naphthylamine [85-84-7] M 247.3, m 102-104°. See 1-benzeneazo-2-naphthylamine on p. 120.

4-Phenylazophenacyl bromide [62625-24-5] M 317.3, m 103-104°. Purified on a column of silica gel, using pet ether/diethyl ether (9:1 v/v) as solvent.

4-Phenylazophenol (4-hydroxyazobenzene) [1689-82-3] M 198.2, m 155°, pK₁²⁵ -0.93, pK₂²⁵ 8.2. Crystd from *benzene or 95% EtOH.

Phenyl benzenethiosulfonate (diphenyldisulfoxide) [1212-08-4] M 250.3, m 36-37°, 45-46°, 45-47°. Recrystd from EtOH or MeOH. Also purified from phenylsulfide impurities by dissolving in CHCl₃, washing with aq satd NaHCO₃, drying (Na₂SO₄) evaporating and the residual oil was passed through a silica gel column (600g) and eluted with hexane-*C₆H₆ (1L, 4:1, eluting PhSSPh) then *C₆H₆ (1L) which elutes PhSSO₂Ph. [Trost and Massiot *J Am Chem Soc* 99 4405 1977; Knoevenagel and Römer *Chem Ber* 56 215 1923.]

Phenyl benzoate [93-99-2] M 198.2, m 69.5°, b 198-199°. Crystd from EtOH using *ca* twice the volume needed for complete soln at 69°.

Phenyl-1,4-benzoquinone [363-03-1] M 184.2, m 114-115°. Crystd from heptane or pet ether (b 60-70°) and sublimed *in vacuo*. [Carlson and Miller *J Am Chem Soc* 107 479 1985.]

1-Phenylbiguanide [102-02-3] M 177.2, m 144-146°, pK₁³² 2.16, pK₂³² 10.74. Crystd from water or toluene.

S-(-)-1-Phenylbutanol [22135-49-5] M 150.2, m 46-47°, 46-48°, 49°, b 90-92°/2mm. [α]_D¹⁸ -51.4° (c 5, CHCl₃), -44.7° (c 5.13, *C₆H₆). Purified by distn and crystallises on cooling. The *hydrochloride* has [α]_D²⁰ +45.1° (c 4.8, *C₆H₆). The (-)-*hydroperoxide* has b 58°/0.005mm, n_D²⁰ 1.5123, α _D¹⁸ -2.14°, (l = 0.5dcm, neat). [Holding and Ross *J Chem Soc* 145 1954; Davies and Feld *J Chem Soc* 4637 1958.] The (\pm)-*racemate* has b 73°/0.05mm, and its 4-nitrophenylhydrazone has m 58°.

Phenylbutazone (4-butyl-1,2-diphenylpyrazolidin-3,5-dione) [50-33-9] M 308.4, m 105°, 106-108°. Crystd from EtOH.

2-Phenylbutyramide [90-26-6] M 163.2, m 86°. Crystd from water.

2-Phenylbutyric acid [R-(-)- 938-79-4; S-(+)- 4286-15-1] M 164.2, b 102-104°/atm, d₄²⁰ 1.056, n_D²⁰ 1.521, [α]_D²⁰ (-) and (+) 96° (c 2.5, *C₆H₆), [α]_D²³ (-) and (+) 5.8° (neat), pK_{Est} ~4.3. Purified by distn at atmospheric pressure using an efficient column. The *acid chlorides* have b 106-107°/20mm, [α]_D¹⁸ (-) and (+) 108° (c 2, *C₆H₆). [Levene et al. *J Biol Chem* 100 589 1933, Gold and Aubert *Helv Chim Acta* 41 1512 1958; ORD in heptane: Rothen and Levene *J Chem Phys* 7 975 1939.]

3-Phenylbutyric acid [R-(-)- 772-14-5; S-(+)- 772-15-6] M 164.2, b 94-95°/3mm, 134°/4mm, d₄²⁶ 1.066, n_D²⁵ 1.5167, [α]_D²⁰ (-) and (+) 57° (c 1, *C₆H₆), pK₂₅ 4.40. Purified as the 2-isomer above, i.e. by distn, but under a good vacuum. [Prelog and Scherrer *Helv Chim Acta* 42 2227 1959; Levene and Marker *J Biol Chem* 93 761 1932, 100 685 1933; Cram *J Am Chem Soc* 74 2137 1952.] The *R-amide*

crystallises from H₂O, **m** 101.5-102°, [α]_D²⁰ -16.5° (c 1.2, EtOH). The *racemic acid* has **m** 39-40°, **b** 134-136°/6mm, 158°/12mm [Marvel et al. *J Am Chem Soc* **62** 3499 1940].

4-Phenylbutyric acid [1821-12-1] **M** 164.2, **m** 50°, **pK**²⁵ 4.76. Crystd from pet ether (b 40-60°).

***o*-(Phenylcarbamoyl)-1-scopolamine methobromide** [138-10-3] **M** 518.4, **m** 200.5-201.5° (**dec**). Crystd from 95% EtOH.

9-Phenylcarbazole [1150-62-5] **M** 243.3, **m** 94-95°. Crystd from EtOH or isopropanol and sublimed *in vacuo*.

***O*-Phenyl chlorothionoformate** [1005-56-7] **M** 172.6, **b** 81-83°/6mm, 91°/10mm, **d**₄²⁰ 1.276, **n**_D²⁰ 1.585. Purified by dissolving in CHCl₃, washing with H₂O, drying (CaCl₂), filtering, evaporating and distilling twice under vacuum to give a clear yellow liquid. **It is reactive and POISONOUS - work in a fume cupboard.** Store in sealed ampoules under N₂. Possible impurity is *O,O'*-diphenyl thiocarbonate which has **m** 106° which remains behind in the distilling flask. [Bögemann et al. in *Methoden Der Organischen Chemie (Houben-Weyl)* 4th edn (E. Müller Ed.) **Vol 9 Schwefel-Selen-Tellur Verbindungen** pp. 807-808 1955; Rivier and Schalch *Helv Chim Acta* **6** 612 1932; Kalson *Chem Ber* **20**, 2384 1987; Rivier and Richard *Helv Chim Acta* **8** 490 1925; Schönberg and Varga *Justus Liebigs Ann Chem* **483** 176 1930; *Chem Ber* **64** 1390 1931.]

Phenyl cinnamate [2757-04-2] **M** 224.3, **m** 75-76°, **b** 205-207°/15mm. Crystd from EtOH (2mL/g). It can also be distd under reduced pressure.

α -Phenylcinnamic acid [91-48-5] **M** 224.3, **m** 174°(*cis*), **m** 138-139°(*trans*), **pK** 4.8 (60% aq EtOH). Crystd from ether/pet ether.

***o*-Phenylenediamine** [95-54-5] **M** 108.1, **m** 100-101°, **pK**₁²⁵ 0.67 (aq H₂SO₄), **pK**₂²⁵ 4.47 (4.85). Crystd from aqueous 1% sodium hydrosulfite (charcoal), washed with ice-water and dried in a vacuum desiccator, or sublimed *in vacuo*. It has been purified by recrystn from toluene and zone refined [Anson et al. *J Am Chem Soc* **108** 6593 1986]. Purification by refluxing a CH₂Cl₂ solution containing charcoal was also carried out followed by evaporation and recrystn [Koola and Kochi *J Org Chem* **52** 4545 1987], protect from light.

***m*-Phenylenediamine** [108-45-2] **M** 108.1, **m** 61-63°, 62-63°, 62.5°, 63-64°, **b** 146°/22mm, 282-284°/760mm, 284-287°/atm, **d**₁₀¹⁰ 1.1422, **n**_D^{57.7} 1.6340, **pK**₁²⁵ 2.41, **pK**₂²⁵ 4.98. Purified by distn under vac followed by recryst from EtOH (rhombs) and if necessary redistn. It should be protected from light otherwise it darkens rapidly. [Neilson et al. *J Chem Soc* 371 1962; IR: Katritzky and Jones *J Chem Soc* 3674, 2058 1959; UV: Forbes and Leckie *Can J Chem* **36** 1371 1958.] The *hydrochloride* has **m** 277-278°, and the *bis-4-chlorobenzenesulfonyl* derivative has **m** 220-221° from H₂O (214-215°, from MeOH-H₂O) [Runge and Pfeiffer *Chem Ber* **90** 1737 1957].

***p*-Phenylenediamine** [106-50-3] **M** 108.1, **m** 140°, **pK**₁²⁵ 2.89, **pK**₂²⁵ 6.16. Crystd from EtOH or *benzene, and sublimed *in vacuo*, protect from light.

***o*-Phenylenediamine dihydrochloride** [615-28-1] **M** 181.1, **m** 180°. Crystd from dilute HCl (60mL conc HCl, 40mL water, with 2g stannous chloride), after treatment of the hot soln with charcoal by adding an equal volume of conc HCl and cooling in an ice-salt mixture. The crystals were washed with a small amount of conc HCl and dried in a vacuum desiccator over NaOH.

2-Phenyl-1,3-diazahexahydroazulene [2161-31-1] **M** 212.3. Recrystd three times from de-aerated cyclohexane in the dark.

1,4-Phenylene diisothiocyanate (bitoscanate) [4044-65-9] M 192.3, m 129-131°, 130-131°, 132°. Purified by recrystn from AcOH, pet ether (b 40-60°), Me₂CO or aq Me₂CO. [van der Kerk et al. *Recl Trav Chim Pays-Bas* 74 1262 1955; Leiber and Slutkin *J Org Chem* 27 2214 1962.]

1-Phenyl-1,2-ethanediol [*R*-(-)- 16355-00-3; *S*-(+)- 25779-13-9] M 138.2, m 64-67°, 65-66°, $[\alpha]_D^{24}$ (-) and (+) 40.5° (c 2.8, H₂O), $[\alpha]_D^{20}$ (-) and (+) 39° (c 3, EtOH). Purified by recryst from *C₆H₆-ligroin and sublimed at 1-2mm. [Arpesella et al. *Gazetta* 85 1354 1955; Prelog et al. *Helv Chim Acta* 37 221 1954.]

***dl*-1-Phenylethanol** [13323-81-4] M 122.2, b 60.5-61.0°/3mm, 106-107°/22-23mm, d 1.01, n²⁵ 1.5254. Purified *via* its hydrogen phthalate. [See Houssa and Kenyon *J Chem Soc* 2260 1930.] Shaken with a soln of ferrous sulfate, and the alcohol layer was washed with distilled H₂O, dried (MgSO₄) and fractionally distd.

2-Phenylethanol [60-12-8] M 122.2, b 215-217°, d 1.020. Purified by shaking with a soln of ferrous sulfate, and the alcohol layer was washed with distd water and fractionally distd.

Phenyl ether (diphenyl ether) [101-84-8] M 170.2, m 27.0°, d 1.074, n^{30.7} 1.57596. Crystd from 90% EtOH. Melted, washed with 3M NaOH and water, dried with CaCl₂ and fractionally distd under reduced pressure. Fractionally crystd from its melt and stored over P₂O₅.

1-Phenylethyl isocyanate (α-methylphenyl isocyanate) [*R*-(+)- 33375-06-3; *S*(-)- 14649-03-7] M 147.2, b 82-83°/12-14mm, d₄²⁰ 1.045, n_D²⁰ 1.513, $[\alpha]_D^{24}$ (+) and (-) 2° (c 3.5, *C₆H₆), (+) and (-) 10.5° (neat). Purified by fractional distn under vacuum. With ammonia it gives the *ureido* derivative which crystallises from H₂O, m 121-122°, $[\alpha]_D^{25}$ (+) and (-) 48.8°. [Cairns *J Am Chem Soc* 63 870 1941.] The *racemate* has b 90-94°/3mm, 96°/18mm [Seifan *Justus Liebigs Ann Chem* 562 75 1949].

***p*-α-Phenylethylphenol** [1988-89-2] M 198.3, m 56.0-56.3°, pK_{Est} ~10.3. Crystd from pet ether.

5-(α-Phenylethyl)semioxamazine [93-95-8] M 207.1, m 167-168° (*l*-), 157° (*dl*-). Crystd from EtOH.

9-Phenyl-3-fluorone [975-17-7] M 320.3, m >300°(dec), λ_{max} 462nm (ε 4.06 x 10⁴, in 1M HCl aq EtOH). Recrystd from warm, acidified EtOH by addition of ammonia. The crude material (1g) can be extracted with EtOH (50mL) in a Soxhlet apparatus for 10h to remove impurities. Impurities can be detected by paper electrophoresis. [Petrova et al. *Anal Lett* 5 695 1972.]

***L*-α-Phenylglycine** [2935-35-5] M 151.2, m 305-310°, $[\alpha]_{546}$ +185° (c 1, M HCl), pK₁^{2.5} 1.83, pK₂^{2.5} 4.39 (for *dl*). Crystd from EtOH.

Phenylglycine-*o*-carboxylic acid [612-42-0] M 195.2, m 208°. Crystd from hot water (charcoal).

Phenylhydrazine [100-63-0] M 108.1, m 23°, b 137-138°/18mm, 241-242°/760mm, d 1.10, n 1.607, pK₁^{2.0} -5.2 (aq H₂SO₄), pK₂^{2.5} 5.27. Purified by chromatography, then crystd from pet ether (b 60-80°)/*benzene. [Shaw and Stratton *J Chem Soc* 5004 1962.]

Phenylhydrazine hydrochloride [59-88-1] M 144.5, m 244°. One litre of boiling EtOH was added to 100g of phenylhydrazine hydrochloride dissolved during 1-3h (without heating) in 200mL of warm water (60-70°). The soln was filtered off, while still hot, through Whatman No 2 filter paper and cooled in a refrigerator. The ppt was collected on a medium sintered-glass filter and recrystd twice this way, then washed with cold EtOH, dried thoroughly and stored in a stoppered brown bottle. [Peterson, Karrer and Guerra *Anal Chem* 29 144 1957.] Hough, Powell and Woods [*J Chem Soc* 4799 1956] boiled the hydrochloride with three times its weight of water, filtered hot (charcoal), added one-third volume of conc HCl and cooled to 0°. The crystals were washed with acetone, and dried over P₂O₅ under vacuum. The salt has also been crystd from 95% EtOH.

Phenylhydroxylamine (N-hydroxyaniline) [100-65-2] M 109.1, m 82°, pK 3.2. Impure base deteriorates rapidly. Crystd from H₂O, *C₆H₆ or *C₆H₆/pet ether (40-60°). Picrate has m 186°.

2-Phenyl-1,3-indandione [83-12-5] M 222.2, m 149-151°, pK²⁰ 4.12 (1% aq MeOH). Crystd from EtOH.

2-Phenylindolizine [25379-20-8] M 193.2, m 214°(dec), pK_{Est} ~4.4. Crystd from EtOH.

Phenylisocyanate [103-71-9] M 119.1, b 45-47°/10mm, d 1.093, n 1.536. Distd under reduced pressure from P₂O₅.

Phenylisothiocyanate (phenyl mustard oil) [103-72-0] M 135.2, m -21°, b 95°/12mm, 117.1°/33mm, 221°/760mm, d₄²⁵ 1.1288, n_D^{23.4} 1.64918. It is insol in H₂O, but sol in Et₂O and EtOH. If impure (due to formation of thiourea) then steam dist into a receiver containing 5-10mL of N H₂SO₄. Separate the oil, dry over CaCl₂ and distil under vacuum. [Dains et al. *Org Synth Coll Vol I* 447 1941.]

8-Phenylmenthol [1R,2S,5R(-)- 65253-04-5; 1S,2R,5S(+)- 57707-91-2] M 232.4, [α]_D²⁰ (-) and (+) 26° (c 2, EtOH). Dissolve in toluene, dry (Na₂SO₄), evap and chromatograph on a silica gel column and eluting with 5% Et₂O in pet ether to give an oil with the desired rotation. IR has ν 3420cm⁻¹ (OH) with consistent ¹H NMR [Corey and Ensley *J Org Chem* 43 1610 1978; Whitesell et al. *Tetrahedron* 42 2993 1986; Bednarski and Danishefsky *J Am Chem Soc* 108 7060 1986.]

1-Phenyl-5-mercaptotetrazole [86-93-1] M 178.2, m 150° (dec), 155° (dec), 157-158°, pK²⁵ 3.65 (5% aq EtOH). Purified by recryst from EtOH or CHCl₃ (m 152°) [Tautomerism: Kauer and Sheppard *J Org Chem* 32 3580 1967; UV: Leiber et al. *Can J Chem* 37 563 1959]. The ammonium salt crystallises from EtOH and dec at 176°, and the sodium salt crystallises from EtOH-*C₆H₆, melts at 96° and dec at 145° [Stollé *J Prakt Chem* [2] 133 60 1932].

Phenyl methanesulfonate [16156-59-5] M 172.1, m 61-62°. Crystd from MeOH.

2-Phenyl-naphthalene [612-94-2] M 204.3, m 103-104°. Chromatographed on alumina in *benzene and crystd from aqueous EtOH.

N-Phenyl-1-naphthylamine [90-30-2] M 219.3, m 63.7-64.0°, pK_{Est} ~0.1. Crystd from EtOH, pet ether or *benzene/EtOH. Dried under vacuum in an Abderhalden pistol.

N-Phenyl-2-naphthylamine [135-88-6] M 219.3, m 107.5-108.5°, 110°, pK_{Est} ~0.5. Crystd from EtOH, MeOH, glacial acetic acid or *benzene/hexane.

4-Phenylphenacyl bromide [135-73-9] M 275.2, m 126°. Crystd (charcoal) from EtOH (15mL/g), or ethyl acetate/pet ether (b 90-100°).

2-Phenylpropanal [93-53-8] M 134.2, b 206°/760mm, d 1.001, n 1.5183. May contain up to 15% of acetophenone. Purified via the bisulfite addition compound [Lodge and Heathcock *J Am Chem Soc* 109 3353 1987] and see Chapter 2 for prepn, and decompn, of bisulfite adduct.

Phenylpropionic acid [637-44-5] M 146.2, m 137.8-138.4°, pK²⁵ 2.23. Crystd from *benzene, CCl₄ or aqueous EtOH.

RS-2-Phenylpropionic acid [492-37-5] M 150.2, m 16-16.5°, b 153-155°/20mm, 189°/48mm, 260-262°/atm, d 1.10, n 1.522, pK²⁵ 4.3. Fractionally distd, or recrystd from pet ether (b 40-60°) strong cooling (see references below).

2-Phenylpropionic acid [R(-)- 7782-26-5; S(+)- 7782-24-3] M 150.2, m 30.3-31°, 30-32°, b 115°/1-2mm, 142°/12mm, [α]_D²⁰ (-) and (+) 99.7° (l = 1 dcm, neat), (-) and (+) 89.1° (c

1.7, EtOH), (-) and (+) 75° (c 1.6, CHCl₃). Purified by vacuum distn and by recrystn from pet ether. The *S*-anilide has **m 103-104°** (from H₂O or CHCl₃/*C₆H₆), $[\alpha]_D^{25} +47^\circ$ (c 9, Me₂CO) [Argus and Kenyon *J Chem Soc* 916 1939; Campbell and Kenyon *J Chem Soc* 25 1946; Levene et al. *J Biol Chem* 88 27, 34 1930; Beilstein 9, 3rd Suppl p 2417].

3-Phenylpropionic acid (hydrocinnamic acid) [501-52-0] M 150.2, m 48-48.5°, pK²⁵ 4.56. Crystd from *benzene, CHCl₃ or pet ether (b 40-60°). Dried in a vacuum.

3-Phenylpropyl bromide [637-59-2] M 199.1, b 110°/12mm, 128-129°/29mm, d 1.31. Washed successively with conc H₂SO₄, water, 10% aqueous Na₂CO₃ and again with water, then dried with CaCl₂ and fractionally distd just before use.

Phenyl 2-pyridyl ketoxime [1826-28-4] M 198.2, m 151-152°. Crystd from EtOH (charcoal).

Phenylpyruvic acid [156-06-9] M 164.2, m 150-154°, 158-159°, pK_{Est} ~2.1. Recrystd from *C₆H₆. The *phenylhydrazone* has **m 173°** [Zeller *Helv Chim Acta* 26 1614 1943; Hopkins and Chisholm *Can J Research [B]* 24 89 1946]. The *2,4-dinitrophenylhydrazone* has **m 162-164°** (189°, 192-194°) [Fones *J Org Chem* 17 1952].

6-Phenylquinoline [612-95-3] M 205.3, m 110.5-111.5°, pK_{Est} ~5.2. Crystd from EtOH (charcoal).

2-Phenylquinoline-4-carboxylic acid (cinchophen) [132-60-5] M 249.3, m 215°, pK_{Est(1)}~0.5 (CO₂H), pK_{Est(2)}~5.1 (N). Crystd from EtOH (*ca* 20mL/g).

Phenyl salicylate (Salol) [118-55-8] M 214.2, m 41.8-42.6°, pK_{Est} ~9.9. Fractionally crystd from its melt, then crystd from *benzene.

3-Phenylsalicylic acid [304-06-3] M 214.3, m 186-187.5°, pK_{Est(1)}~2.8 (CO₂H), pK_{Est(2)}~11.0 (OH). Dissolved in *ca* 1 equivalent of saturated aqueous Na₂CO₃, filtered and ppted by adding 0.8 equivalents of M HCl. Crystd from ethylene dichloride (charcoal), and sublimed at 0.1mm. [Brooks, Eglington and Norman *J Chem Soc* 661 1961.]

1-Phenylsemicarbazide [103-03-7] M 151.2, m 172°. Crystd from water and dried in vac over KOH.

4-Phenylsemicarbazide [537-47-3] M 151.2, m 122°. Crystd from water and dried in vac over KOH.

Phenylsuccinic acid [R(-)- 46292-93-7; S-(+)- 4036-30-1] M 194.2, m 173-176°, 178.5-179°, 179-180°, $[\alpha]_D^{25}$ (-) and (+) 171° (c 2, Me₂CO), $[\alpha]_D^{26-30}$ (-) and (+) 148° (c 0.27-5, EtOH), pK₁²⁵ 3.78, pK₂²⁵ 5.55. Purified by repptn from alkali and recrystn from H₂O. [Naps and Johns *J Am Chem Soc* 62 2450 1940; Fredga and Matell *Bull Soc Chim Belg* 62 47 1953; Wren and Williams *J Chem Soc* 109 572 1916.] The *racemate* [635-51-8] has **m 166-168°**, 168° after recrystn from H₂O or MeCN; its *S*-benzylthiouronium salt has **m 164-165°** (from EtOH) [Griediger and Pedersen *Acta Chem Scand* 9 1425 1955].

1-Phenyl-5-sulfanilamidopyrazole [526-08-9] M 314.3, m 179-183°. Crystd from EtOH.

1-Phenylthiosemicarbazide [645-48-7] M 167.2, m 200-201°(dec). Crystd from EtOH.

4-Phenylthiosemicarbazide [5351-69-9] M 167.2, m 140°. Crystd from EtOH.

1-Phenyl-2-thiourea [103-85-5] M 152.1, m 154°. Crystd from water and dried at 100° in air.

Phenyltoloxamine [2-(2-dimethylaminoethoxy)-diphenylmethane] hydrochloride [6152-43-8] M 291.8, m 119-120°, pK²⁵ 9.3 (free base). Crystd from isobutyl methyl ketone.

[Previous Page](#)

Phenyl 4-toluenesulfonate [640-60-8] M 248.2, m 94.5-95.5°. Crystd from MeOH or glacial acetic acid.

Phenyl 4-tolylcarbonate [13183-20-5] M 228.2, m 67°. Purified by preparative GLC with 20% Apiezon on Embacel, and sublimed *in vacuo*.

4-Phenyl-1,2,4-triazole-3,5-diol (4-phenylurazole) [15988-11-1] M 175.2, m 207-209°. Crystd from water.

4-Phenyl-1,2,4-triazoline-3,5-dione (PTAD) [4233-33-4] M 175.2, m 165-170°(dec), 170-177°(dec). Carmine red needles obtained by sublimation (ice cold finger) at 100°/0.1mm, and/or by recrystn from EtOH. IR: ν 1760, 1780 cm^{-1} . [Cookson et al. *Org Synth* 51 121 1971; Moore et al. *J Org Chem* 39 3700 1974.]

1-Phenyl-2,2,2-trifluoroethanol [*R*-(-)- 10531-50-7; *S*-(+)- 340-06-7] M 176.1, b 74-76°/10mm, 125-127°/760mm, d_4^{20} 1.301, n_D^{20} 1.4632, $[\alpha]_D^{20}$ (-) and (+) 31° (neat). Purified by fractional distn preferably in a vacuum. [Morrison and Ridgeway *Tetrahedron Lett* 573 1969; NMR: Pirkle and Beare *J Am Chem Soc* 90 6250 1968.] The racemate [340-05-6] has b 52-54°/2mm, 57-59°/2mm, 64-65°/5mm, d_4^{20} 1.293, n_D^{20} 1.457, and the 2-carbobenzoyl derivative has m 137-138° [Mosher et al. *J Am Chem Soc* 78 4374 1956].

Phenylurea [64-10-8] M 136.2, m 148°, pK^{25} -1.45 (aq H_2SO_4). Crystd from boiling water (10mL/g). Dried in a steam oven at 100°.

9-Phenyl-9-xanthenol (hydroxypixyl) [596-38-3] M 274.3, m 158-161°, 158.5-159°, 159°. Dissolve in AcOH and add H_2O whereby it separates as colourless prisms. It is slightly soluble in CHCl_3 , soluble in C_6H_6 but insoluble in pet ether. It sublimes on heating. UV in H_2SO_4 : λ_{max} 450nm (ϵ 5620) and 370nm (ϵ 24,900) and the HClO_4 salt in CHCl_3 has λ_{max} 450 (ϵ 404) and 375nm (ϵ 2420). [Sharp *J Chem Soc* 2558 1958; Bünzly and Decker *Chem Ber* 37 2983 1904; Chattopadhyaya and Reece *J Chem Soc, Chem Commun* 639 1978; Gomberg and Cone *Justus Liebigs Ann Chem* 370 142 1909.]

Phloretin [2',4',6'-trihydroxy-3-(*p*-hydroxyphenyl)propiophenone] [60-82-2] M 274.3, m 264-271°(dec), $\text{pK}_{\text{Est}(1)}$ ~7.5, $\text{pK}_{\text{Est}(2)}$ ~8.0, $\text{pK}_{\text{Est}(3)}$ ~10, $\text{pK}_{\text{Est}(4)}$ ~12 (phenolic OH's). Crystd from aqueous EtOH.

Phloridzin (2H₂O) [phloretin 2'-*O*- β -D-glucoside] [60-81-1] M 472.5, m 110°, $[\alpha]_{546}^{20}$ -62° (c 3.2, EtOH). Crystd as dihydrate from water.

Phloroacetophenone (2H₂O) (2',4',6'-trihydroxyacetophenone) [480-66-0] M 186.2, m 218-219°, $\text{pK}_{\text{Est}(1)}$ ~7.9, $\text{pK}_{\text{Est}(2)}$ ~12.0. Crystd from hot water (35mL/g).

Phloroglucinol (2H₂O) (benzene-1,3,5-triol) [6099-90-7 (2H₂O); 108-73-6 (anhydr)] M 126.1, m 217-219°, 117° (anhydrous), pK_1^{25} -7.74 (HClO_4), pK_2^{20} 7.97, pK_3^{20} 9.23. Crystd from water, and stored in the dark under nitrogen.

Phorone (2,6-dimethylhepta-2,5-dien-4-one) [504-20-1] M 138.2, m 28°, b 197°/743mm. Crystd repeatedly from EtOH.

"Phosphine" [dye CI 793, Chrysaniline mononitrate, 3-amino-9-(4-aminophenyl)-acridinium mononitrate] [10181-37-0] M 348.4, m >250°(dec), pK_{Est} ~8.0. Crystd from *benzene/EtOH.

Phthalaldehyde [643-79-8] M 134.1, m 54-56°, 55.5-56°, 58°, b 83-84°/0.8mm. Purified by steam distillation better by using super heated steam (at 175-180°) and efficient cooling. The distillate is saturated with Na_2SO_4 extracted exhaustively with EtOAc, dried (Na_2SO_4), filtered and evaporated. The residue

is recrystd from pet ether (b 90-100°) [Beill and Tarbell *Org Synth Coll Vol IV* 808 1963]. It can be distd under vacuum. The bis-2,4-dinitrophenylhydrazone has m 278-280° [Hatt and Stephenson *J Chem Soc* 199 1952].

Phthalazine [253-52-1] M 130.2, m 90-91°, pK²⁰ 3.47. Crystd from diethyl ether or *benzene, and sublimed under vacuum.

Phthalazine-1,4-dione (phthalhydrazide) [1445-69-8] M 162.2, m 330-333°, 336°, 346°, pK₁²⁰ -3.29 pK₂²⁰ -0.99, pK₃²⁰ 5.67, pK₄²⁰ 13.0. Twice recrystd from 0.1M KOH [Merenyi et al. *J Am Chem Soc* 108 7716 1986], EtOH or dimethylformamide and sublimes >300°.

Phthalazone (1-hydroxyphthalazine) [119-39-1] M 146.2, m 183-184°, 186-188°, b 337°/760mm, pK₁²⁰ -2.2, pK₂²⁰ -1.4, pK₃²⁰ 11.99. Crystd from H₂O or EtOH and sublimed *in vacuo*.

o-Phthalic acid [88-99-3] M 166.1, m 211-211.5°, pK₁²⁵ 2.76 (3.05), pK₂²⁵ 4.92 (4.73). Crystd from water.

Phthalic anhydride [85-44-9] M 148.1, m 132°, b 295°. Distd under reduced pressure. Purified from the acid by extracting with hot CHCl₃, filtering and evaporating. The residue was crystd from CHCl₃, CCl₄ or *benzene, or sublimed. Fractionally crystd from its melt. Dried under vacuum at 100°. [Saltiel *J Am Chem Soc* 108 2674 1986.]

Phthalide [87-41-2] M 134.1, m 72-73°, pK -7.98 (aq H₂SO₄). Crystd from water (75mL/g) and dried in air on filter paper.

Phthalimide [85-41-6] M 147.1, m 235°, 238°, pK 8.30. Crystd from EtOH (20mL/g) (charcoal), or by sublimation. For potassium phthalimide see entry in Chapter 5.

Phthalimidoglycine [4702-13-0] M 205.2, m 192-193°, pK_{Est} ~3. Crystd from water or EtOH.

Phthalonitrile [91-15-6] M 128.1, m 141°. Crystd from EtOH, toluene or *benzene. Can also be distd under high vacuum.

Phthalylsulfacetamide [131-69-1] M 362.3, m 196°. Crystd from water.

Phthiocol (2-hydroxy-3-methylnaphthaquinone) [483-55-6] M 188.1, m 173-174°, pK_{Est} ~4.2. Crystd from diethyl ether/pet ether.

Physalien (all trans β-carotene-3,3'-diol dipalmitate) [144-67-2] M 1044, m 98.5-99.5°, A_{1m}^{1%} (λ_{max}) 1410 (449nm), 1255 (478nm) in hexane. Purified by chromatography on water-deactivated alumina, using hexane/diethyl ether (19:1) to develop the column. Crystd from *benzene/EtOH. Stored in the dark, in inert atmosphere, at 0°.

Physodic acid [4,4',6'-trihydroxy-6-(2-oxoheptyl)-2'-pentyl-2,3'-oxydibenzoic acid 1,5-lactone] [84-24-2] M 470.5, m 205°, pK_{Est(1)} ~3.0, pK_{Est(2)} ~10, pK_{Est(3)} ~13. Crystd from MeOH. The diacetate has m 155-156° (from Me₂CO/CS₂).

Phytoene (7,7',8,8',11,11',12,12'-octahydro-ψ,ψ-carotene) [540-04-5] M 544.9, A_{1m}^{1%} (λ_{max}) 850 (287nm) in hexane, λ_{max} 275, 287 and 297nm nm. Purified by chromatography on columns of magnesium oxide-Supercel (a diatomaceous filter aid) or alumina [Rabourn et al. *Arch Biochem Biophys* 48 267 1954]. Stored as a solution in pet ether under nitrogen at -20°.

Phytofluene [540-05-6] M 549.0, b 140-185°(bath temp)/0.0001m A_{1m}^{1%} (λ_{max}) 1350 (348nm) in pet ether, λ_{max} 331, 348, 267. Purified by chromatography on partially deactivated alumina [Kushwaha et al. *J Biol Chem* 245 4708 1970]. Stored as a soln in pet ether under nitrogen at -20°.

Picein (*p*-acetylphenyl- β -D-glucopyranoside) [530-14-3] M 298.3, m 195-196°, $[\alpha]_D^{20}$ -88° (c 1, H₂O). Crystd from MeOH or (as monohydrate) from water.

Picene [213-14-3] M 278.3, m 364°. Crystd from isopropylbenzene/xylene. Can also be sublimed.

2-Picoline-N-oxide (2-methylpyridine-1-oxide) [931-19-1] M 109.1, m 41-45°, b 89-90°/0.8-0.9mm, 90-100°/1mm, 110°/4mm, 135°/5mm, 123°/9mm, 123-124°/15mm, 259-261°/atm, n_D^{25} 1.5854 (supercooled), pK^{25} 1.10. Purified by fractional distillation and could be recrystd from *C₆H₆-hexane but is *hygroscopic*. [Bullitt and Maynard *J Am Chem Soc* 76 1370 1954; Ross et al. *J Am Chem Soc* 78 3625 1956; IR: Wiley and Slaym Aker *J Am Chem Soc* 79 2233 1957.] The *picrate* has m 125-126.5° (from EtOH) [Boekelheide and Linn *J Am Chem Soc* 76 1286 1954]. The *phthalate* has m 115-116° (from EtOH) [den Hertog et al. *Recl Trav Chim Pays-Bas* 70 591 1951.]

3-Picoline-N-oxide (3-methylpyridine-1-oxide) [1003-73-2] M 109.1, m 37-39°, 37-38° (evac capillary), 84-85°/0.3mm, 101-103°/0.7-0.8mm, 114-115°/1.5mm, 118°/2mm, pK^{25} 1.08. Purified by careful fractionation *in vacuo*. The distillate remains supercooled for several days before solidifying. It is a slightly *hygroscopic* solid which could melt in the hand. The *picrate* has m 149-151° (from EtOH). [Taylor and Corvetti *Org Synth Coll Vol IV* 654 1963; IR: Katritzky et al. *J Chem Soc* 3680 1959; Jaffé and Doak *J Am Chem Soc* 77 4441, 4481 1955; Boekelheide and Linn *J Am Chem Soc* 76 1286 1954].

4-Picoline-N-oxide (4-methylpyridine-1-oxide) [1003-67-4] M 109.1, m 182-184°, 185-186°, 186-188°, pK^{25} 1.29. Recryst from EtOH-EtOAc, Me₂CO-Et₂O or *C₆H₆. [Bullitt and Maynard *J Am Chem Soc* 76 1370 1954; Boekelheide and Linn *J Am Chem Soc* 76 1286 1954].

Picolinic acid (pyridine-2-carboxylic acid) [98-98-6] M 123.1, m 138°, pK_1^{25} 1.03 (1.36), pK_2^{25} 5.30 (5.80). Crystd from water or *benzene.

α -Picolinium chloride [14401-91-3] M 129.6, m 200°. 1:1 Mixture of α -picoline and HCl, distd at 275°. Then vacuum sublimed at 91-91.5°.

N-Picolinoylbenzimidazole [100312-29-6] M 173.3, m 105-107°. Recrystd three times from hexane [Fife and Przystas *J Am Chem Soc* 108 4631 1986].

Picric acid [88-89-1] M 229.1, m 122-123°, pK^{25} 0.33 (0.37). Crystd first from acetic acid then acetone, toluene, CHCl₃, aqueous 30% EtOH, 95% EtOH, MeOH or H₂O. Dried in a vacuum oven at 80° for 2h. Alternatively, dried over Mg(ClO₄)₂ or fused and allowed to freeze under vacuum three times. Because it is **EXPLOSIVE**, picric acid should be stored moistened with H₂O, and only small portions should be dried at any one time. The dried acid should **NOT** be heated.

Picrolonic acid [3-methyl-4-nitro-1-(4-nitrophenyl)-2-pyrazolin-5-one, picrolonic acid] [550-74-3] M 264.2, m 120°(dec), 116.5°(dec at 125°) 125°. Crystd from water or EtOH (Solubility is 0.123% at 15° and 1.203% at 100° in H₂O; and 1.107% at 0° and 11.68% at 81° in EtOH). It forms Ca, Cu Hg, Mg, Na, Sr, and Pb complexes [Maquestian et al *Bull Soc Chim Belg* 82 233 1973; Isaki et al. *Chem Ber* 74 1420 1941].

Picrotoxin [124-87-8] M 602.6, m 203°, $[\alpha]_{546}^{20}$ -40° (c 1, EtOH). Crystd from water.

Picryl chloride [88-88-0] M 226.3, m 83°. Crystd from CHCl₃ or EtOH.

Picryl iodide [4436-27-5] M 340.0, m 164-165°. Crystd from *benzene.

Pimelic acid (heptane-1,7-dioic acid) [111-16-0] M 160.2, m 105-106°, pK_1^{25} 4.46, pK_2^{25} 5.58. Crystd from water or from *benzene containing 5% diethyl ether.

Pinacol (hexahydrate) [6091-58-3 (6H₂O); 76-09-5 (anhydr)] M 194.3, m 46.5°, b 59°/4mm. Distd then crystd repeatedly from water.

Pinacol (anhydrous) [76-09-5] M 118.1, m 41.1°, b 172°. The hydrate is rendered anhydrous by azeotropic distn of water with *benzene. Recrystd from *benzene or toluene/pet ether, absolute EtOH or dry diethyl ether. Recrystn from water gives the hexahydrate.

Pinacolone oxime [2475-93-6] M 115.2, m 78°. Crystd from aqueous EtOH.

Pinacyanol chloride [2768-90-3] M 388.9, m 270°(dec). Crystd from EtOH/diethyl ether.

R- α -Pinene [7785-70-8] M 136.2, b 61°/30mm, 156.2°/760mm, d 0.858, n¹⁵ 1.4634, n 1.4658, $[\alpha]_D^{25}$ +47.3°. Isomerised by heat, acids and certain solvents. Should be distd under reduced pressure under nitrogen and stored in the dark. Purified *via* the nitrosochloride [Waterman et al. *Recl Trav Chim Pays-Bas* 48 1191 1929]. For purification of optically active forms see Lynn [*J Am Chem Soc* 91 361 1919]. Small quantities (0.5mL) have been purified by GLC using helium as carrier gas and a column at 90° packed with 20 wt% of polypropylene sebacate on a Chromosorb support. Larger quantities were fractionally distd under reduced pressure in a column packed with stainless steel gauze spirals. Material could be dried with CaH₂ or sodium, and stored in a refrigerator: CaSO₄ and silica gel were not satisfactory because they induced spontaneous isomerisation. [Bates, Best and Williams *J Chem Soc* 1521 1962.]

S- α -Pinene [7785-26-4] M 136.2, b 155-156°/760mm, d 0.858, n 1.4634, $[\alpha]_D^{20}$ -47.2°. Purification as for R- α -Pinene above.

dl-Pipecolic acid (piperidine-2-carboxylic acid) [4043-87-2] M 129.1, m 264°, pK₁²⁵ 2.29, pK₂²⁵ 10.77. Crystd from water.

Piperazine [110-85-0] M 86.1, m 110-112°, 44° (hexahydrate 142-63-2) b 125-130°/760mm, pK₁²⁵ 5.33, pK₂²⁵ 9.73. Crystd from EtOH or anhydrous *benzene, and dried at 0.01mm. It can be sublimed under vacuum and purified by zone melting.

§ Piperazine on polystyrene support is commercially available.

Piperazine-N,N'-bis(2-ethanesulfonic acid) (PIPES) [5625-37-6] M 302.4, pK₁²⁵ <3, pK₂²⁵ 6.82 (7.82). Crystd from boiling water (maximum solubility is about 1g/L) or as described for ADA [*N*-(2-acetamido)iminodiacetic acid, see above].

Piperazine dihydrochloride (H₂O) [142-64-3 (2HCl); 6094-40-2 (xHCl)] M 177.1, m 82.5-83.5°. Crystd from aqueous EtOH. Dried at 110°.

Piperazine phosphate (H₂O) [18534-18-4] M 197.6. Crystd twice from water, air-dried and stored for several days over Drierite. The salt dehydrates slowly if heated at 70°.

Piperic acid [*trans,trans*-5-(3,4-methylenedioxyphenyl)-2,4-pentadieneoic acid] [136-72-1] M 218.2, m 217°, pK_{Est} ~4.7. Crystd from EtOH. Turns yellow in light. Sublimes with partial dec.

Piperidine [110-89-4] M 85.2, f -9°, b 35.4°/40mm, 106°/760mm, d 0.862, n 1.4535, n²⁵ 1.4500, pK₂²⁵ 11.20. Dried with BaO, KOH, CaH₂, or sodium, and fractionally distd (optionally from sodium, CaH₂, or P₂O₅). Purified from pyridine by zone melting.

§ Piperidine on polystyrene support is commercially available.

Piperidinium chloride [6091-44-7] M 121.6, m 244-245°. Crystd from EtOH/diethyl ether in the presence of a small amount of HCl.

Piperidinium nitrate [6091-45-8] M 145.2, m 110°. Crystd from acetone/ethyl acetate.

Piperine (1-piperoylpiperidine) [94-62-2] M 285.4, m 129-129.5°, pK¹⁵ 1.98. Crystd from EtOH or *benzene/ligroin.

Piperonal [120-57-0] M 150.1, m 37°, b 140°/15mm, 263°/760mm. Crystd from aqueous 70% EtOH or EtOH/water.

Piperonylic acid [94-53-1] M 166.1, m 229°, pK²⁵ 4.50. Crystd from EtOH or water.

Pivalic acid (trimethylacetic acid) [75-98-9] M 102.1, m 35.4°, b 71-73°/0.1mm, pK²⁵ 5.03. Fractionally distd under reduced pressure, then fractionally crystd from its melt. Recrystd from *benzene.

Pivaloyl chloride (trimethylacetyl chloride) [3282-30-2] M 120.6, b 57.6°/150mm, 70.5-71/250mm, 104°/754mm, 104-105°/atm, 105-108°/atm, d₄²⁰ 1.003, n_D²⁰ 1.4142. First check the IR to see if OH bands are present. If absent, or present in small amounts, then redistil under moderate vac. If present in large amounts then treat with oxalyl chloride or thionyl chloride and reflux for 2-3h, evap and distil residue. **Strongly LACHRYMATORY - work in a fumecupboard.** Store in sealed ampoules under N₂. [Traynham and Battiste *J Org Chem* 22 1551 1957; Grignard reactns: Whitmore et al. *J Am Chem Soc* 63 647 1941.]

Plumbagin (5-hydroxy-2-methyl-1,4-naphthaquinone) [481-42-5] M 188.1, m 78-79°, pK_{Est(1)}~9.5, pK_{Est(2)}~11.0. Crystd from aqueous EtOH and sublimed in a vac. Steam distils.

Polyacrylonitrile [25014-41-9]. Ppted from dimethylformamide by addition of MeOH.

Poly(diallyldimethylammonium) chloride [26062-79-3]. Ppted from water in acetone, and dried in vacuum for 24h. [Hardy and Shriner *J Am Chem Soc* 107 3822 1985.]

Polyethylene [9002-88-4]. Crystd from thiophen-free *benzene and dried over P₂O₅ under vacuum.

Polymethyl acrylate [9003-21-8]. Ppted from a 2% soln in acetone by addition of water.

Polystyrene [9003-53-6]. Ppted repeatedly from CHCl₃ or toluene soln by addition of MeOH. Dried *in vacuo* [Miyasaka et al. *J Phys Chem* 92 249 1988].

Polyvinyl acetate [9003-20-7]. Ppted from acetone by addition of *n*-hexane.

Poly(*N*-vinylcarbazole) [25067-59-8]. Ppted seven times from tetrahydrofuran with MeOH, with a final freeze-drying from *benzene. Dried under vacuum.

Polyvinyl chloride [9002-81-2]. Ppted from cyclohexanone by addition of MeOH.

Poly(4-vinylpyridine) [25232-41-1] M (105.1)_n. Purified by repeated pptn from solns in EtOH and dioxane, and then EtOH and ethyl acetate. Finally, freeze-dried from *tert*-butanol.

Poly(*N*-vinylpyrrolidone) [9003-39-8] M (111.1)_n, crosslinked [25249-54-1] m >300°. Purified by dialysis, and freeze-dried. Also by pptn from CHCl₃ soln by pouring into ether. Dried in a vacuum over P₂O₅. For the crosslinked polymer purification is by boiling for 10min in 10% HCl and then washing with glass-distilled water until free from Cl ions. Final Cl ions were removed more readily by neutralising with KOH and continued washing.

Prednisone [53-03-2] M 358.5, m 238°(dec), [α]_D²⁰ +168° (c 1, dioxane), λ_{max} 238nm (log ε 4.18) in MeOH. Crystd from acetone/hexane.

Pregnane [481-26-5] M 300.5, m 83.5°, [α]_D²⁰ +21° (CHCl₃). Crystd from MeOH.

5 β -Pregnane-3 α ,20 α -diol [80-92-2] M 320.5, m 243-244 $^{\circ}$, $[\alpha]_{546}^{20}$ +31 $^{\circ}$ (c 1, EtOH). Crystd from acetone.

5 β -Pregnane-3 α ,20 β -diol [80-91-1] M 320.5, m 244-246 $^{\circ}$, $[\alpha]_{546}^{20}$ +22 $^{\circ}$ (c 1, EtOH). Crystd from EtOH.

Procaine [4-(2-diethylaminomethoxycarbonyl)aniline] [59-46-1] M 236.3, m 51 $^{\circ}$ (dihydrate), 61 $^{\circ}$ (anhydrous), pK_1^{15} 2.45, pK_2^{15} 8.91. Crystd as the dihydrate from aqueous EtOH and as anhydrous material from pet ether or diethyl ether. The latter is *hygroscopic*.

Proclavine (3,6-diaminoacridine) [92-62-6] M 209.2, m 284-286 $^{\circ}$, pK^{25} 9.60. Crystd from aqueous MeOH. For proflavin see 3,6-diaminoacridine hydrochloride

Progesterone [57-83-0] M 314.5, m 128.5 $^{\circ}$, $[\alpha]_{546}^{20}$ +220 $^{\circ}$ (c 2, dioxane). Crystd from EtOH. When crystd from pet ether m is 121 $^{\circ}$, λ_{\max} 240nm, log ϵ 4.25 (EtOH).

L-Proline [147-85-3] M 115.1, m 215-220 $^{\circ}$ (dec)(D-isomer), 220-222 $^{\circ}$ (dec) (L-form), 205 $^{\circ}$ (dec)(DL-isomer), $[\alpha]_D^{25}$ (H₂O, L-isomer), pK_1^{25} 1.95, pK_2^{25} 10.64. Likely impurity are hydroxyproline. Purified *via* its picrate which was crystd twice from water, then decomposed with 40% H₂SO₄. The picric acid was extracted with diethyl ether, the H₂SO₄ was pptd with Ba(OH)₂, and the filtrate evapd. The residue was crystd from hot absolute EtOH [Mellan and Hoover *J Am Chem Soc* 73 3879 1951] or EtOH/ether. *Hygroscopic*. Stored in a desiccator.

Polycopene (all Z- ψ , ψ -carotene) [2361-24-2] M 536.5, m 111 $^{\circ}$, λ_{\max} 443.5, 470nm in pet ether. Purified by chromatography on deactivated alumina [Kushwaha et al. *J Biol Chem* 245 4708 1970]. Crystd from pet ether. Stored in the dark, in an inert atmosphere at -20 $^{\circ}$.

L-Prolylglycine [2578-57-6] M 172.2, m 236 $^{\circ}$, $[\alpha]_D^{20}$ +21.1 $^{\circ}$ (c 4, H₂O), pK_1^{25} 3.19, pK_2^{25} 8.97. Crystd from water at 50-60 $^{\circ}$ by addition of EtOH.

Proneurosporene (3,4,7',8'-tetrahydrolycopene) [10467-46-6] M 538.9, λ_{\max} 408, 432, 461 nm, $\epsilon_{1\text{cm}}^{1\%}$ 2040 (432nm) in hexane. Purified by chromatography on deactivated alumina [Kushwaha et al. *J Biol Chem* 245 4708 1970]. Stored in the dark, in an inert atmosphere at 0 $^{\circ}$.

Propane [74-98-6] M 44.1, m -189.7, b -42.1 $^{\circ}$ /760mm, d 0.5005, n 1.2898. Purified by bromination of the olefinic contaminants. Propane was treated with bromine for 30min at 0 $^{\circ}$. Unreacted bromine was quenched, and the propane was distd through two -78 $^{\circ}$ traps and collected at -196 $^{\circ}$ [Skell et al. *J Am Chem Soc* 108 6300 1986].

Propane-1,2-diamine (propylenediamine) [78-90-0] M 74.1, b 120.5 $^{\circ}$, d 0.868, n 1.446, pK_1^{25} 6.61, pK_2^{25} 9.82. Purified by azeotropic distn with toluene. [Horton, Thomason and Kelly *Anal Chem* 27 269 1955.]

Propane-1,2-diol (propyleneglycol) [57-55-6] M 76.1, b 104 $^{\circ}$ /32mm, d 1.040, n 1.433. Dried with Na₂SO₄, decanted and distd under reduced pressure.

Propane-1,3-diol [504-63-2] M 76.1, b 110-122 $^{\circ}$ /12mm, d 1.053, $n^{18.5}$ 1.4398. Dried with K₂CO₃ and distd under reduced pressure. More extensive purification involved conversion with benzaldehyde to 2-phenyl-1,3-dioxane (m 47-48 $^{\circ}$) which was subsequently decomposed by shaking with 0.5M HCl (3mL/g) for 15min and standing overnight at room temperature. After neutralisation with K₂CO₃, the benzaldehyde was removed by distn and the diol was recovered from the remaining aqueous soln by continuous extraction with CHCl₃ for 1day. The extract was dried with K₂CO₃, the CHCl₃ was evaporated and the diol was distd. [Foster, Haines and Stacey *Tetrahedron* 6 177 1961.]

Propane-1-thiol [107-03-9] M 76.1, b 65.3°/702mm, d²⁵ 0.83598, n²⁵ 1.43511, pK²⁰ 10.82. Purified by soln in aqueous 20% NaOH, extraction with a small amount of *benzene and steam distn until clear. After cooling, the soln was acidified slightly with 15% H₂SO₄, and the thiol was distd out, dried with anhydrous CaSO₄ or CaCl₂, and fractionally distd under nitrogen. [Mathias and Filho *J Phys Chem* 62 1427 1958.] Also purified by liberation of the mercaptan by adding dilute HCl to the residue remaining after steam distn. After direct distn from the flask, and separation of the water, the mercaptan was dried (Na₂SO₄) and distd under nitrogen.

Propane-2-thiol (Isopropyl mercaptan) [75-33-2] M 76.1, b 49.8°/696mm, d²⁵ 0.80895, n²⁵ 1.42154, pK²⁵ 10.86. Purification as for propane-1-thiol above.

Propargyl alcohol (2-propyn-1-ol) [107-19-7] M 56.1, b 54°/57mm, 113.6°/760mm, d 0.947, n 1.432. Commercial material contains a stabiliser. An aqueous soln of propargyl alcohol can be concentrated by azeotropic distn with butanol or butyl acetate. Dried with K₂CO₃ and distd under reduced pressure, in the presence of about 1% succinic acid, through a glass helices-packed column.

Propargyl chloride (3-chloropropyne) [624-65-7] M 74.5, b 58°/760mm, 65°/760mm, d 1.03, n 1.435. Purified by fractional distn at atm press. Note that a possible impurity propargyl alcohol has b 114-115°/atm. [Henry *Chem Ber* 8 398 1875.] **HIGHLY TOXIC and FLAMMABLE.**

Propene [115-07-1] M 42.1, m -185.2°, b -47.8°/750mm, d 0.519, n⁷¹ 1.357. Purified by freeze-pump-thaw cycles and trap-to-trap distn.

p-(1-Propenyl)phenol [cis/trans 6380-21-8; 539-12-8] M 134.2, m 93-94°, pK_{Est} ~10.2. Crystd from water.

β-Propiolactone [57-57-8] M 72.1, b 83°/45mm, d 1.150, n²⁵ 1.4117. Fractionally distd under reduced pressure, from sodium. **CARCINOGEN.**

Propionaldehyde [123-38-6] M 58.1, b 48.5-48.7°, d 0.804, n 1.3733, n²⁵ 1.37115. Dried with CaSO₄ or CaCl₂, and fractionally distd under nitrogen or in the presence of a trace of hydroquinone (to retard oxidation). Blacet and Pitts [*J Am Chem Soc* 74 3382 1952] repeatedly vacuum distd the middle fraction until no longer gave a solid polymer when cooled to -80°. It was stored with CaSO₄.

Propionamide [79-05-0] M 73.1, m 79.8-80.8°, pK²⁴ -0.9 (H₀ scale, aq H₂SO₄). Crystd from acetone, *benzene, CHCl₃, water or acetone/water, then dried in a vacuum desiccator over P₂O₅ or conc H₂SO₄.

Propionic acid [79-09-4] M 74.1, b 141°, d 0.992, n 1.3865, n²⁵ 1.3843, pK₁²⁵ -6.8 (H₀ scale, aq H₂SO₄), pK₂²⁵ 4.88. Dried with Na₂SO₄ or by fractional distn, then redistd after refluxing with a few crystals of KMnO₄. An alternative purification uses the conversion to the ethyl ester, fractional distn and hydrolysis. [Bradbury *J Am Chem Soc* 74 2709 1952.] Propionic acid can also be heated for 0.5h with an amount of benzoic anhydride equivalent to the amount of water present (in the presence of CrO₃ as catalyst), followed by fractional distn. [Cham and Israel *J Chem Soc* 196 1960.]

Propionic anhydride [123-62-6] M 130.2, b 67°/18mm, 168°/780mm, d 1.407, n 1.012. Shaken with P₂O₅ for several minutes, then distd.

Propionitrile [107-12-0] M 55.1, b 97.2°, d 1.407, n¹⁵ 1.36812, n³⁰ 1.36132. Shaken with dil HCl (20%), or with conc HCl until the odour of isonitrile has gone, then washed with water, and aqueous K₂CO₃. After a preliminary drying with silica gel or Linde type 4A molecular sieves, it is stirred with CaH₂ until hydrogen evolution ceases, then decanted and distd from P₂O₅ (not more than 5g/L, to minimise gel formation). Finally, it is refluxed with, and slowly distd from CaH₂ (5g/L), taking precautions to exclude moisture.

***n*-Propyl acetate** [109-60-4] M 102.1, b 101.5°, d 0.887, n 1.38442, pK²⁵ -7.18 (H₀ scale, aq H₂SO₄). Washed with satd aqueous NaHCO₃ until neutral, then with satd aqueous NaCl. Dried with MgSO₄ and fractionally distd.

***n*-Propyl alcohol (1-propanol)** [71-23-8] M 60.1, b 97.2°, d²⁵ 0.79995, n 1.385, pK²⁵ 16.1. The main impurities in *n*-propyl alcohol are usually water and 2-propen-1-ol, reflecting the commercial production by hydration of propene. Water can be removed by azeotropic distn either directly (azeotrope contains 28% water) or by using a ternary system, e.g. by adding *benzene. Alternatively, for gross amounts of water, refluxing over CaO for several hours is suitable, followed by distn and a further drying. To obtain more nearly anhydrous alcohol, suitable drying agents are firstly NaOH, CaSO₄ or K₂CO₃, then CaH₂, aluminium amalgam, magnesium activated with iodine, or a small amount of sodium. Alternatively, the alcohol can be refluxed with *n*-propylsuccinate or phthalate in a method similar to the one described under EtOH. Allyl alcohol is removed by adding bromine (15mL/L) and then fractionally distilling from a small amount of K₂CO₃. Propionaldehyde, also formed in the bromination, is removed as the 2,4-dinitrophenylhydrazone. *n*-Propyl alcohol can be dried down to 20ppm of water by passage through a column of pre-dried molecular sieves (type 3 or 4A, heated for 3h at 300°) in a current of nitrogen. Distn from sulfanilic or tartaric acids removes impurities. Albrecht [*J Am Chem Soc* 82 3813 1960] obtained spectroscopically pure material by heating with charcoal to 50-60°, filtering and adding 2,4-dinitrophenylhydrazine and a few drops of conc H₂SO₄. After standing for several hours, the mixture was cooled to 0°, filtered and vac distd. Gold and Satchell [*J Chem Soc* 1938 1963] heated *n*-propyl alcohol with 3-nitrophthalic anhydride at 76-110° for 15h, then recrystd the resulting ester from H₂O, *benzene/pet ether (b 100-120°)(3:1), and *benzene. The ester was hydrolysed under reflux with aq 7.5M NaOH for 45min under nitrogen, followed by distn (also under nitrogen). The fraction (b 87-92°) was dried with K₂CO₃ and stirred under reduced pressure in the dark over 2,4-dinitrophenylhydrazine, then freshly distilled. Also purified by adding 2g NaBH₄ to 1.5L alcohol, gently bubbling with argon and refluxing for 1day at 50°. Then added 2g of freshly cut sodium (washed with propanol) and refluxed for one day. Distd, taking the middle fraction [Jou and Freeman *J Phys Chem* 81 909 1977].

***n*-Propylamine** [107-10-8] M 59.1, b 48.5°, d 0.716, n 1.38815, pK²⁵ 10.69. Distd from zinc dust, at reduced pressure, in an atmosphere of nitrogen.

***n*-Propyl bromide.** [106-94-5] M 123.0, b 71.0°, d 1.354., n¹⁵ 1.43695, n²⁵ 1.43123. Likely contaminants include *n*-propyl alcohol and isopropyl bromide. The simplest purification procedure uses drying with MgSO₄ or CaCl₂ (with or without a preliminary wash of the bromide with aq NaHCO₃, then water), followed by fractional distn away from bright light. Chien and Willard [*J Am Chem Soc* 79 4872 1957] bubbled a stream of oxygen containing 5% ozone through *n*-propyl bromide for 1h, then shook with 3% hydrogen peroxide soln, neutralised with aq Na₂CO₃, washed with distilled water and dried. Then followed vigorous stirring with 95% H₂SO₄ until fresh acid did not discolour within 12h. The propyl bromide was separated, neutralised, washed dried with MgSO₄ and fractionally distd. The centre cut was stored in the dark. Instead of ozone, Schuler and McCauley [*J Am Chem Soc* 79 821 1957] added bromine and stored for 4 weeks, the bromine then being extracted with aq NaHSO₃ before the sulfuric acid treatment was applied. Distd. Further purified by preparative gas chromatography on a column packed with 30% SE-30 (General Electric ethylsilicone rubber) on 42/60 Chromosorb P at 150° and 40psi, using helium. [Chu *J Phys Chem* 41 226 1964.]

***n*-Propyl chloride** [540-54-5] M 78.5, b 46.6°, d 0.890, n 1.3880. Dried with MgSO₄ and fractionally distd. More extensively purified using extraction with H₂SO₄ as for *n*-propyl bromide. Alternatively, Chien and Willard [*J Am Chem Soc* 75 6160 1953] passed a stream of oxygen containing about 5% ozone through the *n*-propyl chloride for three times as long as was needed to cause the first coloration of starch iodide paper by the exit gas. After washing with aqueous NaHCO₃ to hydrolyse ozonides and remove organic acids, the chloride was dried with MgSO₄ and fractionally distd.

1-Propyl-3-(*p*-chlorobenzenesulfonyl) urea [94-20-2] M 260.7, m 127-129°. Crystd from aqueous EtOH.

Propylene carbonate [108-32-7] M 102.1, b 110°/0.5-1mm, 238-239°/760mm, d 1.204, n 1.423. Manufactured by reaction of 1,2-propylene oxide with CO₂ in the presence of a catalyst (quaternary

ammonium halide). Contaminants include propylene oxide, carbon dioxide, 1,2- and 1,3-propanediols, allyl alcohol and ethylene carbonate. It can be purified by percolation through molecular sieves (Linde 5A, dried at 350° for 14h under a stream of argon), followed by distn under vac. [Jasinski and Kirkland *Anal Chem* **39** 163 1967.] It can be stored over molecular sieves under an inert gas atmosphere. When purified in this way it contains less than 2ppm water. Activated alumina and dried CaO have been also used as drying agents prior to fractional distn under reduced pressure. It has been dried with 3A molecular sieves and distd under nitrogen in the presence of *p*-toluenesulfonic acid. Then redistilled and the middle fraction collected.

***dl*-Propylene oxide** [75-56-9] M 58.1, b 34.5°, d 0.829, n 1.3664. Dried with Na₂SO₄ or CaH₂, and fractionally distilled through a packed column (glass helices), after refluxing with Na, CaH₂, or KOH pellets.

***n*-Propyl ether (dipropyl ether)** [111-43-3] M 102.2, b 90.1°, d 0.740, n¹⁵ 1.38296, n 1.3803, pK -4.40 (aq H₂SO₄). Purified by drying with CaSO₄, by passage through an alumina column (to remove peroxides), and by fractional distn.

Propyl formate [110-74-7] M 88.1, b 81.3°, d 0.9058, n 1.3779. Distd, then washed with satd aq NaCl, and with satd aq NaHCO₃ in the presence of solid NaCl, dried with MgSO₄ and fractionally distd.

***n*-Propyl gallate** [121-79-9] M 212.2, m 150°. Crystd from aqueous EtOH.

***n*-Propyl iodide (1-iodopropane)** [107-08-4] M 170.0, b 102.5°, d 1.745, n 1.5041. Should be distd at reduced pressure to avoid decomposition. Dried with MgSO₄ or silica gel and fractionally distd. Stored under nitrogen with mercury in a brown bottle. Prior to distn, free iodine can be removed by shaking with copper powder or by washing with aq Na₂S₂O₃ and drying. Alternatively, the *n*-propyl iodide can be treated with bromine, then washed with aq Na₂S₂O₃ and dried. See also *n*-butyl iodide.

***n*-Propyl propionate** [106-36-5] M 120.2, b 122°, d 0.881, n 1.393. Treated with anhydrous CuSO₄, then distd under nitrogen.

6-Propyl-2-thiouracil (propacil, propylail) [51-52-5] M 170.2, m 218-220°, 218-220°, pK₁²¹ -6.54 (aq H₂SO₄), pK₂²¹ -4.22 (aq H₂SO₄), pK₃²¹ 8.25 (4% aq EtOH). Purified by recrystn from H₂O (sol in 900 parts at 20°, and 100 parts at 100°). UV, MeOH: λ_{max} 277nm. [Anderson et al. *J Am Chem Soc* **67** 2197 1945; Vanderhaegue *Bull Soc Chim Belg* **59** 689 1950.]

Propyne [74-99-7] M 40.1, m -101.5°, b -23.2°/760mm, d⁵⁰ 0.7062, n⁴⁰ 1.3863. Purified by preparative gas chromatography.

Protocatechualdehyde [139-85-5] M 138.1, m 153°. Crystd from water or toluene and dried in a vacuum desiccator over KOH pellets or shredded wax respectively.

Protopine [fumarine, macleine, 4,6,7,14-tetrahydro-5-methyl-bis[1,3]-benzodioxolo[4,5-*c*:5',6'-*g*]azecine-13(5*H*)-one] [130-86-9] M 353.4, m 208°, pK 5.99. Crystd from EtOH/CHCl₃.

1*S*,2*S*-Pseudoephedrine (1-hydroxy-1-phenyl-2-methylaminopropane) [90-82-4] M 165.2, m 118-119°, [α]_D²⁰ +53.0° (EtOH), +40.0° (H₂O), pK²⁵ 9.71. Crystd from dry diethyl ether, or from water and dried in a vacuum desiccator.

1*S*,2*S*-Pseudoephedrine hydrochloride [345-78-8] M 210.7, m 181-182°, 185-188°, [α]_D²⁰ +61° (c 1 H₂O). Crystd from EtOH.

Pteridine [91-18-9] M 132.2, m 139.5-140°, pK₁²⁰ 4.05 (equilibrium, hydrate), pK₂²⁰ 11.90 (OH of hydrate). Crystd from EtOH, *benzene, *n*-hexane, *n*-heptane or pet ether. It sublimes at 120-130°/20mm. Stored at 0°, in the dark; turns green in the presence of light and on long standing in the dark.

2,4-(1H,3H)-Pteridinedione H₂O (lumazine) [487-21-8] M 182.1, m >350°, pK₁²⁰ <1.0, pK₂²⁰ 7.94. Crystd from water.

Pterin (2-aminopteridin-4(3H)-one) [2236-60-4] M 163.1, m >300°, pK₁²⁰ 2.27 (basic), pK₂²⁰ 7.96 (acidic). It was dissolved in hot 1% aqueous ammonia, filtered, and an equal volume of hot 1M aqueous formic acid was added. The soln was allowed to cool at 0-2° overnight. The solid was collected and washed with distilled water several times by centrifugation and dried *in vacuo* over P₂O₅ overnight, and then at 100° overnight.

Pterocarpin {(6a*R*-*cis*)-6a,12a-dihydro-3-methoxy-6H-[1,3]dioxolo[5,6]benzofuro[3,2c][1]-benzopyran} [524-97-0] M 298.3, m 165°, [α]₅₄₆²⁰ -215° (c 0.5, CHCl₃). Crystd from EtOH, or pet ether.

Pteronic acid (2-amino-6-*p*-carboxyanilinomethylpteridin-4(3H)-one) [119-24-4] M 312.3, m >300°(dec), pK_{Est(1)}~ 2.3 (basic, N1), pK_{Est(2)}~ 2.6 (basic, CH₂NH), pK_{Est(3)}~ 4.5 (COOH), pK_{Est(4)}~ 7.9 (acidic 4-OH). Crystd from dilute HCl. Hygroscopic IRRITANT

R(+)-Pulegone [89-82-7] M 152.2, b 69.5°/5mm, n 1.4849, d 0.935, [α]₅₄₆²⁰ +23.5°(neat). Purified *via* the semicarbazone. [Erskine and Waight *J Chem Soc* 3425 1960.]

Purine [120-73-0] M 120.1, m 216-217°, pK₁²⁰ 2.30, pK₂²⁰ 9.86. Crystd from toluene or EtOH.

Purpurin (1,2,4-trihydroxy-5,10-anthraquinone) [81-54-9] M 256.2, m 253-256°, pK_{Est(1)}~7.0 (2-OH), pK_{Est(2)}~9.0 (4-OH), pK_{Est(3)}~11.1 (1-OH). Cryst from aq EtOH, dry at 100°.

Purpurogallin (2,3,4,6-tetrahydroxy-5H-benzocyclohepten-5-one) [569-77-7] M 220.2, m 274° (rapid heating) (pK 7-10, phenolic OH). Crystd from acetic acid.

Pyocyanine (1-hydroxy-5-methylphenazinium zwitterion) [85-66-5] M 210.2, m 133° (sublimes and dec on further heating). Crystd from H₂O as dark blue needles. *Picrate* has m 190° dec.

Pyrazine [290-37-9] M 80.1, m 47°, b 115.5-115.8°, pK₁²⁰ -6.25 (aq H₂SO₄), pK₂²⁵ 1.1 (0.51 at 20°). Distd in steam and crystd from water. Purified by zone melting.

Pyrazinecarboxamide [98-96-4] M 123.1, m 189-191° (sublimes slowly at 159°), pK -0.5. Crystd from water or EtOH.

Pyrazinecarboxylic acid [98-97-5] M 124.1, m 225-229°(dec), pK²⁵ 2.92. Crystd from water.

Pyrazine-2,3-dicarboxylic acid [89-01-0] M 168.1, m 183-185°(dec), pK₁ <-2.0, pK₂ 0.9, pK₃ 2.77 (2.20). Crystd from water. Dried at 100°.

Pyrazole [288-13-1] M 68.1, m 70°, pK²⁵ 2.48. Crystd from pet ether, cyclohexane, or water. [Barszcz et al. *J Chem Soc, Dalton Trans* 2025 1986.]

Pyrazole-3,5-dicarboxylic acid [3112-31-0] M 174.1, m 287-289°(dec), pK_{Est(1)}~1.2 (CO₂H), pK_{Est(2)}~3.7 (CO₂H), pK_{Est(3)}~12 (NH). Crystd from water or EtOH.

Pyrene [129-00-0] M 202.3, m 149-150°. Crystd from EtOH, glacial acetic acid, *benzene or toluene. Purified by chromatography of CCl₄ solns on alumina, with *benzene or *n*-hexane as eluent. [Backer and Whitten *J Phys Chem* 91 865 1987.] Also zone refined, and purified by sublimation. Marvel and Anderson [*J Am Chem Soc* 76 5434 1954] refluxed pyrene (35g) in toluene (400mL) with maleic anhydride (5g) for 4days, then added 150mL of aqueous 5% KOH and refluxed for 5h with occasional shaking. The toluene layer was separated, washed thoroughly with H₂O, concentrated to about 100mL and allowed to cool. Crystalline pyrene was filtered off and recrystd three times from EtOH or acetonitrile. [Chu and Thomas *J Am Chem Soc* 108

6270 1986; Russell et al. *Anal Chem* **50** 2961 1986.] The material was free from anthracene derivatives. Another purification step involved passage of pyrene in cyclohexane through a column of silica gel. It can be sublimed in a vacuum and zone refined. [Kano et al. *J Phys Chem* **89** 3748 1985.]

Pyrene-1-aldehyde [3029-19-4] **M 230.3, m 125-126°**. Recrystd three times from aqueous EtOH.

1-Pyrenebutyric acid [3443-45-6] **M 288.4, m 184-186°, pK_{Est} ~4.1**. Crystd from *benzene, EtOH, EtOH/water (7:3 v/v) or *C₆H₆/AcOH. Dried over P₂O₅. [Chu and Thomas *J Am Chem Soc* **108** 6270 1986.]

1-Pyrenecarboxylic acid [19694-02-1] **M 230.3, m 126-127°, pK_{Est} ~3.2**. Crystd from *C₆H₆ or 95% EtOH.

1-Pyrenesulfonic acid [26651-23-0] **M 202.2, m >350°, pK_{Est} <0**. Crystd from EtOH/water. The tetra-Na salt cryst from H₂O and the sulfonyl chloride has **m 120°(dec)**. [Vollmann et al. *Justus Liebigs Ann Chem* **531** 32 1937 and *Justus Liebigs Ann Chem* **540** 189 1939.]

1,3,6,8-Pyrenetetrasulfonic acid [6528-53-6] **M 522.2, m >400°, pK_{Est} <0** Crystd from water [Tietz and Bayer *Justus Liebigs Ann Chem* **540** 189 1939.]

Pyridine [110-86-1] **M 79.1, f -41.8°, b 115.6°, d 0.9831, n 1.51021, pK²⁵ 5.23**. Likely impurities are H₂O and amines such as the picolines and lutidines. Pyridine is *hygroscopic* and is miscible with H₂O and organic solvents. It can be dried with solid KOH, NaOH, CaO, BaO or sodium, followed by fractional distn. Other methods of drying include standing with Linde type 4A molecular sieves, CaH₂ or LiAlH₄, azeotropic distn of the H₂O with toluene or *benzene, or treated with phenylmagnesium bromide in ether, followed by evaporation of the ether and distn of the pyridine. A recommended [Lindauer and Mukherjee *Pure Appl Chem* **27** 267 1971] method dries pyridine over solid KOH (20g/Kg) for 2weeks, and fractionally distils the supernatant over Linde type 5A molecular sieves and solid KOH. The product is stored under CO₂-free nitrogen. Pyridine can be stored in contact with BaO, CaH₂ or molecular sieves. Non-basic materials can be removed by steam distilling a soln containing 1.2 equivalents of 20% H₂SO₄ or 17% HCl until about 10% of the base has been carried over along with the non-basic impurities. The residue is then made alkaline, and the base is separated, dried with NaOH and fractionally distd.

Alternatively, pyridine can be treated with oxidising agents. Thus pyridine (800mL) has been stirred for 24h with a mixture of ceric sulfate (20g) and anhydrous K₂CO₃ (15g), then filtered and fractionally distd. Hurd and Simon [*J Am Chem Soc* **84** 4519 1962] stirred pyridine (135mL), water (2.5L) and KMnO₄ (90g) for 2h at 100°, then stood for 15h before filtering off the ppted manganese oxides. Addition of solid KOH (*ca* 500g) caused pyridine to separate. It was decanted, refluxed with CaO for 3h and distd.

Separation of pyridine from some of its homologues can be achieved by crystn of the oxalates. Pyridine is ppted as its oxalate by adding it to the stirred soln of oxalic acid in acetone. The ppte is filtered, washed with cold acetone, and pyridine is regenerated and isolated. Other methods are based on complex formation with ZnCl₂ or HgCl₂. Heap, Jones and Speakman [*J Am Chem Soc* **43** 1936 1921] added crude pyridine (1L) to a soln of ZnCl₂ (848g) in 730mL of water, 346mL of conc HCl and 690mL of 95% EtOH. The crystalline ppte of ZnCl₂.(pyridine)₂ was filtered off, recrystd twice from absolute EtOH, then treated with a conc NaOH soln, using 26.7g of solid NaOH to 100g of the complex. The ppte was filtered off, and the pyridine was dried with NaOH pellets and distd. Similarly, Kyte, Jeffery and Vogel [*J Chem Soc* 4454 1960] added pyridine (60mL) in 300mL of 10% (v/v) HCl to a soln of HgCl₂ (405g) in hot water (2.3L). On cooling, crystals of pyridine-HgCl₂ (1:1) complex separated and were filtered off, crystd from 1% HCl (to **m 178.5-179°**), washed with a little EtOH and dried at 110°. The free base was liberated by addition of excess aq NaOH and separated by steam distn. The distillate was saturated with solid KOH, and the upper layer was removed, dried further with KOH, then BaO and distd. Another possible purification step is fractional crystn by partial freezing.

Small amounts of pyridine have been purified by vapour-phase chromatography, using a 180-cm column of polyethyleneglycol-400 (Shell 5%) on Embacel (May and Baker) at 100°, with argon as carrier gas. The Karl Fischer titration can be used for determining water content. A colour test for pyrrole as a contaminant is described by Biddiscombe et al. [*J Chem Soc* 1957 1954].

§ Polystyrene supported pyridine is commercially available.

Pyridine-2-aldehyde [1121-60-4] M 107.1, b 81.5°/25mm, d 1.121, n 1.535, pK_1^{25} 3.84, pK_2^{25} 12.68. Sulfur dioxide was bubbled into a soln of 50g in 250mL of boiled water, under nitrogen, at 0°, until pptn was complete. The addition compound was filtered off rapidly and, after washing with a little water, it was refluxed in 17% HCl (200mL) under nitrogen until a clear soln was obtained. Neutralisation with $NaHCO_3$ and extraction with ether separated the aldehyde which was recovered by drying the extract, then distilling twice, under nitrogen. [Kyte, Jeffery and Vogel *J Chem Soc* 4454 1960.]

Pyridine-3-aldehyde [500-22-1] M 107.1, b 89.5°/14mm, d 1.141, n 1.549, pK_1^{20} 3.80, pK_2^{20} 13.10. Purification as for pyridine-2-aldehyde.

Pyridine-4-aldehyde [872-85-5] M 107.1, b 79.5°/12mm, d 1.137, n 1.544, pK_1^{20} 4.77, pK_2^{20} 12.20. . Purification as for pyridine-2-aldehyde.

Pyridine-2-aldoxime (pyridine-2-carboxaldoxime) [873-69-8] M 122.1, m 111-113°, 114°, pK_1^{25} 3.56, pK_2^{25} 10.17. Recrystd from Et_2O -pet ether or H_2O . The *picrate* has m 169-171° (from aqueous EtOH). It is used in peptide synthesis. [UV: Grammaticakis *Bull Chem Soc Fr* 109, 116 1956; Ginsberg and Wilson *J Am Chem Soc* 79 481 1957; Hanania and Irvine *Nature* 183 40 1959; Green and Saville *J Chem Soc* 3887 1956.]

Pyridine-3-aldoxime [1193-92-6] M 122.1, m 150°, pK_1^{20} 4.07, pK_2^{20} 10.39. Crystd from water.

Pyridine-4-aldoxime [696-54-8] M 122.1, m 129°, pK_1^{20} 4.73, pK_2^{20} 10.03. Crystd from water.

2,6-Pyridinedialdoxime [2851-68-5] M 165.1, m 212°, $pK_{Est(1)}$ ~3.0, $pK_{Est(2)}$ ~10. Crystd from water.

Pyridine-2,5-dicarboxylic acid [100-26-5] M 167.1, m 254°, pK_2^{25} 2.49, pK_3^{25} 5.12. Crystd from dil HCl.

Pyridine-3,4-dicarboxylic acid [490-11-9] M 167.1, m 256°, pK_1^{25} 2.43, pK_2^{25} 4.78. Crystd from dilute aqueous HCl.

Pyridine hydrobromide perbromide (pyridinium bromide perbromide) [39416-48-3] M 319.9, m 130° (dec), 132-134° (dec). It is a very good brominating agent - liberating one mol. of Br_2 . Purified by recrystn from glacial acetic acid (33g from 100mL of AcOH). [Fieser and Fieser *Reagents for Organic Chemistry* Vol 1 967 1967.]

Pyridine hydrochloride [628-13-7] M 115.6, m 144°, b 218°. Crystd from $CHCl_3$ /ethyl acetate and washed with diethyl ether.

Pyridine N-oxide [694-59-7] M 95.1, m 67°, pK^{24} 0.79. Purified by vacuum sublimation.

Pyridine 3-sulfonic acid [636-73-7] M 159.2, m 365-366° (dec), 357°, pK^{25} 2.89 (12% aq EtOH), 3.22 (H_2O)(protonation on N). Purified by recrystn from H_2O or aqueous EtOH as needles or plates. [pKa: Evans and Brown *J Org Chem* 27 3127 1962; IR: Arnett and Chawla *J Am Chem Soc* 100 214 1978.] UV in 50% aqueous EtOH: λ_{max} at 208 and 262nm. The *ammonium salt* has m 243° (from H_2O), the *sulfonyl chloride* has m 133-134° (from pet ether), the *amide* has m 110-111° (from H_2O), the *hydrochloride* has m > 300° (dec), and the *N-methyl betaine* has m 130° (from H_2O). [Gastel and Wibaut *Recl Trav Chim Pays Bas* 53 1031 1934; McIlvain and Goese *J Am Chem Soc* 65 2233 1943; Machek *Monatsh Chem* 72 77 1938.]

2-Pyridinethiol (2-mercaptopyridine) [2637-34-5] M 111.2, m 127.4°, 127-130°, 130-132°, pK_1^{20} -1.07, pK_2^{20} 9.97. If impure, dissolve in $CHCl_3$, wash with dil AcOH, H_2O , dry ($MgSO_4$), evaporate under reduced press and recryst residue from C_6H_6 or H_2O . *2-Methylmercaptopyridine* (b 100-104°/33mm) was

formed by treatment with MeI/NaOH. [Albert and Barlin *J Chem Soc* 2394 1959; Phillips and Shapiro *J Chem Soc* 584 1942.]

Pyridoxal hydrochloride, pyridoxamine hydrochloride and pyridoxine hydrochloride (vitamin B₆) see entries in Chapter 6.

1-(2-Pyridylazo)-2-naphthol (PAN) [85-85-8] M 249.3, m 140-142°, pK₁³⁰⁻³⁶ 2.9, pK₂³⁰⁻³⁶ 11.2. Purified by repeated crystn from MeOH. It can also be purified by sublimation under vacuum. Purity can be checked by TLC using a mixed solvent (pet ether, diethyl ether, EtOH; 10:10:1) on a silica gel plate.

4-(2-Pyridylazo)resorcinol (PAR) [1141-59-9] M 215.2, m >195°(dec), λ_{max} 415nm, ε 2.59 x 10⁴ (pH 6-12), pK₁²⁵ 2.69, pK₂²⁵ 5.50. Purified as the sodium salt by recrystn from 1:1 EtOH/water. Purity can be checked by TLC using a silica gel plate and a mixed solvent (*n*-BuOH:EtOH:2M NH₃; 6:2:2).

Pyridyldiphenyltriazine [1046-56-6] M 310.4, m 191-192°. Purified by repeated recrystn from EtOH/dimethylformamide.

1-(4-pyridyl)ethanol [*R*-(+)- 27854-88-2; *S*-(-)- 54656-96-1] M 123.2, m 63-65°, 67-69°, [α]_D²⁰ (+) and (-) 49.8° (c 0.5, EtOH), pK_{Est} ~5.4. Purified by recrystn from pet ether. The (-)-*di-O-benzoyl tartrate salt* has m 146-148° (from EtOH). [UV, ORD: Harelli and Samori *J Chem Soc Perkin Trans 2* 1462 1974.] The *racemate* recrystallises from Et₂O m 74-76°, b 90-94°/1mm [Ferles and Attia *Collect Czech Chem Commun* 38 611 1973; UV, NMR: Nielson et al. *J Org Chem* 29 2898 1964.]

Pyrogallol [87-66-1] M 126.1, m 136°, pK₁²⁰ 9.05, pK₂²⁰ 11.19, pK₃²⁰ 14. Crystd from EtOH*/benzene.

***R*-pyroglutamic acid (5-oxo-D-proline, *R*-2-Pyrrolidone-5-carboxylic acid)** [4042-36-8] M 129.1, m 156-158°, [α]₅₄₆²⁰ +11° (c 5, H₂O). Crystd from EtOH/pet ether.

***S*-Pyroglutamic acid (5-oxo-L-proline)** [98-79-3] M 129.1, m 156-158°, 162-164°, [α]₅₄₆²⁰ -11° (c 5, H₂O), pK 12.7 (by electron spin resonance). Crystd from EtOH by addition of pet ether. *NH₄* salt has m 184-186° (from EtOH).

Pyromellitic acid (benzene-1,2,4,5-tetracarboxylic acid) [89-05-4] M 254.2, m 276°, 281-284°, pK₁²⁵ 1.87, pK₂²⁵ 2.72, pK₃²⁵ 4.30, pK₄²⁵ 5.52. Dissolved in 5.7 parts of hot dimethylformamide, decolorised and filtered. The ppt obtained on cooling was separated and air dried, the solvent being removed by heating in an oven at 150-170° for several hours. Crystd from water.

Pyromellitic dianhydride [89-32-7] M 218.1, m 286°. Crystd from ethyl methyl ketone or dioxane. Dried, and sublimed *in vacuo*.

α-Pyrone (2*H*-pyran-2-one) [504-31-4] M 96.1, m 5°, 8-9°, b 103-111°/19-22mm, 110°/26mm, 104°/30mm, 115-118°/37mm, 206-207°/atm, d₄²⁰ 1.1972, n_D²⁰ 1.5298, pK -1.14 (aq H₂SO₄). Dissolve in Et₂O, wash with brine, dry (Na₂SO₄), filter, evaporate, distil residue under vacuum and redistil. It is a colourless liquid. [Zimmermann et al. *Org Synth Coll Vol V* 982 1973; Nakagawa and Saegusa *Org Synth* 56 49 1977; Elderfield *J Org Chem* 6 566 1941.] The *picrate* has m 106-107° (from EtOH).

γ-Pyrone (4*H*-pyran-4-one) [108-97-4] M 96.1, m 32.5-32.6°, 33°, 32-34°, b 88.5°/7mm, 91-91.5°/9mm, 95-97°/13mm, 105°/23mm, 215°/atm, pK₂₅ 0.10. Purified by vacuum distn, the distillate crystallises and is *hygroscopic*. It is non-steam volatile. The *hydrochloride* has m 139° (from EtOH), and the *picrate* has m 130.2-130.3° (from EtOH or H₂O). [Mayer *Chem Ber* 90 2362 1957; IR: Jones et al. *Can J Chem* 37 2007 1959; Neelakatan *J Org Chem* 22 1584 1957.]

Pyronin Y [3,6-bis(dimethylamino)xanthylium chloride] [92-32-0] M 302.8, m 250-260°, CI 45005, λ_{\max} 522nm, $pK_{\text{Est}} \sim 7.6$. Commercial material contained a large quantity of zinc. Purified by dissolving 1g in 50mL of hot water containing 5g NaEDTA. Cooled to 0°, filtered, evapd to dryness and the residue extracted with EtOH. The soln was evaporated to 5-10mL, filtered, and the dye pptd by addition of excess dry diethyl ether. It was centrifuged and the crystals were washed with dry ether. The procedure was repeated, then the product was dissolved in CHCl_3 , filtered and evapd. The dye was stored in a vacuum.

Pyrrole [109-97-7] M 67.1, m 23.4°, b 66°/80mm, 129-130°/atm, d 0.966, n 1.5097, $pK_1^{25} -4.4$ (Protonation on carbon), $pK_2^{25} 17.51$ (aq KOH, H. scale). Dried with NaOH, CaH_2 or CaSO_4 . Fractionally distd under reduced pressure from CaH_2 . Stored under nitrogen, turns brown in air. Redistd immediately before use.

Pyrrolidine [123-75-1] M 71.1, b 87.5-88.5°, d 0.860, n 1.443, $pK^{25} 11.31$. Dried with BaO or sodium, then fractionally distd, under N_2 , through a Todd column packed with glass helices (see p. 174).

Pyrrolidine-1-carbodithioic acid ammonium salt [5108-96-3] M 164.3, m 128-130°, $pK^{25} 3.25$ (free acid). Purified by recryst twice by dissolving in MeOH and adding Et_2O . Also by recrystn from EtOH. [Synth and Polarography: Kitagawa and Taku *Bull Chem Soc Jpn* 64 2151 1973; Malissa and Schöffmann *Mikrochim Acta* 187 1955.]

Pyruvic acid [127-17-3] M 88.1, m 13°, b 65°/10mm, $pK^{25} 2.39$ (2.60). Distd twice, then fractionally crystd by partial freezing.

***p*-Quaterphenyl** [135-70-6] M 306.4, m 312-314°. Recrystd from dimethyl sulfoxide at *ca* 50°.

Quercetin (2H₂O) (3,3',4',5,6-pentahydroxyflavone) [6151-25-3 (2H₂O); 117-39-3 (anhydr)] M 338.3, m *ca* 315°(dec), (phenolic pK_s 7-10). Crystd from aq EtOH and dried at 100°.

Quercitrin (quercetin glycoside) [522-12-3] M 302.2, m 168°, 176-178°. Crystd from aq EtOH and dried at 135° to give the higher melting form.

Quinaldic (quinoline-2-carboxylic) acid [93-10-7] M 173.2, m 156-157°, $pK_1^{25} 1.45$, $pK_2^{25} 2.49$ (2.97). Crystd from *benzene.

Quinalizarin (1,2,5,8-tetrahydroxy-9,10-anthraquinone) [81-61-8] M 272.2, m 275°, $pK_{\text{Est}(1)} \sim 7.1$ (1-OH), $pK_{\text{Est}(2)} \sim 9.9$ (8-OH), $pK_{\text{Est}(3)} \sim 11.1$ (5-OH), $pK_{\text{Est}(4)} \sim 11.8$ (2-OH). Crystd from acetic acid or nitrobenzene. It can be sublimed *in vacuo*.

Quinazoline [253-82-7] M 130.2, m 48.0-48.5°, b 120-121°/17-18mm, $pK_1^{20} -4.51$ (aq H_2SO_4 , anhydrous dication), $pK_2^{20} 2.01$ (anhydrous monocation), $pK_3^{20} 4.3$ (equilibrium with 3,4-hydrated species), $pK_4^{20} 12.1$ (hydrated anion). Purified by passage through an activated alumina column in *benzene or pet ether (b 40-60°). Distd under reduced pressure, sublimed under vacuum and crystd from pet ether. [Armarego *J Appl Chem* 11 70 1961.]

Quinhydrone [106-34-3] M 218.2, m 168°. Crystd from H_2O at 65°, then dried in a vac desiccator.

1R,3R,4R,5R-Quinic acid (1,3,4,5-tetrahydroxy-cyclohexane-carboxylic acid) [77-95-2] M 192.3, m 172°(dec), $[\alpha]_{546}^{20} -51^\circ$ (c 20, H_2O), $pK^{25} 3.58$. Crystd from water.

Quinidine [56-54-2] M 324.4, m 171°, $[\alpha]_{546}^{20} +301.1^\circ$ (CHCl_3 contg 2.5% (v/v) EtOH), $pK_1^{15} 4.13$, $pK_2^{15} 8.77$. Crystd from *benzene or dry CHCl_3 /pet ether (b 40-60°), discarding the initial, oily crop of crystals. Dried under vacuum at 100° over P_2O_5 .

Pyronin Y [3,6-bis(dimethylamino)xanthylium chloride] [92-32-0] M 302.8, m 250-260°, CI 45005, λ_{\max} 522nm, $pK_{\text{Est}} \sim 7.6$. Commercial material contained a large quantity of zinc. Purified by dissolving 1g in 50mL of hot water containing 5g NaEDTA. Cooled to 0°, filtered, evapd to dryness and the residue extracted with EtOH. The soln was evaporated to 5-10mL, filtered, and the dye pptd by addition of excess dry diethyl ether. It was centrifuged and the crystals were washed with dry ether. The procedure was repeated, then the product was dissolved in CHCl_3 , filtered and evapd. The dye was stored in a vacuum.

Pyrrole [109-97-7] M 67.1, m 23.4°, b 66°/80mm, 129-130°/atm, d 0.966, n 1.5097, $pK_1^{25} -4.4$ (Protonation on carbon), $pK_2^{25} 17.51$ (aq KOH, H. scale). Dried with NaOH, CaH_2 or CaSO_4 . Fractionally distd under reduced pressure from CaH_2 . Stored under nitrogen, turns brown in air. Redistd immediately before use.

Pyrrolidine [123-75-1] M 71.1, b 87.5-88.5°, d 0.860, n 1.443, $pK^{25} 11.31$. Dried with BaO or sodium, then fractionally distd, under N_2 , through a Todd column packed with glass helices (see p. 174).

Pyrrolidine-1-carbodithioic acid ammonium salt [5108-96-3] M 164.3, m 128-130°, $pK^{25} 3.25$ (free acid). Purified by recryst twice by dissolving in MeOH and adding Et_2O . Also by recrystn from EtOH. [Synth and Polarography: Kitagawa and Taku *Bull Chem Soc Jpn* 64 2151 1973; Malissa and Schöffmann *Mikrochim Acta* 187 1955.]

Pyruvic acid [127-17-3] M 88.1, m 13°, b 65°/10mm, $pK^{25} 2.39$ (2.60). Distd twice, then fractionally crystd by partial freezing.

***p*-Quaterphenyl** [135-70-6] M 306.4, m 312-314°. Recrystd from dimethyl sulfoxide at *ca* 50°.

Quercetin (2H₂O) (3,3',4',5,6-pentahydroxyflavone) [6151-25-3 (2H₂O); 117-39-3 (anhydr)] M 338.3, m *ca* 315°(dec), (phenolic pK_s 7-10). Crystd from aq EtOH and dried at 100°.

Quercitrin (quercetin glycoside) [522-12-3] M 302.2, m 168°, 176-178°. Crystd from aq EtOH and dried at 135° to give the higher melting form.

Quinaldic (quinoline-2-carboxylic) acid [93-10-7] M 173.2, m 156-157°, $pK_1^{25} 1.45$, $pK_2^{25} 2.49$ (2.97). Crystd from *benzene.

Quinalizarin (1,2,5,8-tetrahydroxy-9,10-anthraquinone) [81-61-8] M 272.2, m 275°, $pK_{\text{Est}(1)} \sim 7.1$ (1-OH), $pK_{\text{Est}(2)} \sim 9.9$ (8-OH), $pK_{\text{Est}(3)} \sim 11.1$ (5-OH), $pK_{\text{Est}(4)} \sim 11.8$ (2-OH). Crystd from acetic acid or nitrobenzene. It can be sublimed *in vacuo*.

Quinazoline [253-82-7] M 130.2, m 48.0-48.5°, b 120-121°/17-18mm, $pK_1^{20} -4.51$ (aq H_2SO_4 , anhydrous dication), $pK_2^{20} 2.01$ (anhydrous monocation), $pK_3^{20} 4.3$ (equilibrium with 3,4-hydrated species), $pK_4^{20} 12.1$ (hydrated anion). Purified by passage through an activated alumina column in *benzene or pet ether (b 40-60°). Distd under reduced pressure, sublimed under vacuum and crystd from pet ether. [Armarego *J Appl Chem* 11 70 1961.]

Quinhydrone [106-34-3] M 218.2, m 168°. Crystd from H_2O at 65°, then dried in a vac desiccator.

1R,3R,4R,5R-Quinic acid (1,3,4,5-tetrahydroxy-cyclohexane-carboxylic acid) [77-95-2] M 192.3, m 172°(dec), $[\alpha]_{546}^{20} -51^\circ$ (c 20, H_2O), $pK^{25} 3.58$. Crystd from water.

Quinidine [56-54-2] M 324.4, m 171°, $[\alpha]_{546}^{20} +301.1^\circ$ (CHCl_3 contg 2.5% (v/v) EtOH), $pK_1^{15} 4.13$, $pK_2^{15} 8.77$. Crystd from *benzene or dry CHCl_3 /pet ether (b 40-60°), discarding the initial, oily crop of crystals. Dried under vacuum at 100° over P_2O_5 .

Quinine [130-95-0] M 324.4, m 177°(dec), $[\alpha]_{546}^{20} -160^\circ$ (c 1, CHCl₃), pK₁²⁰ 4.13 (quinoline N), pK₂²⁰ 8.52 (piperidine N). Crystd from abs EtOH.

Quinine bisulfate [6183-68-2 (7H₂O); 549-56-4 (anhydr)] M 422.4, m 160° (anhydrous). Crystd from 0.1M H₂SO₄, forms heptahydrate when crystd from water

Quinine sulfate (2H₂O) [6119-70-6 (H₂O); 804-63-7 (anhydr)] M 783.0, m 205°. Crystd from water, dried at 110°.

Quinizarin (1,4-dihydroxy-9,10-anthraquinone) [81-64-1] M 240.2, m 200-202°, pK₁²⁵ 9.90 (9.5), pK₂²⁵ 11.18. Crystd from glacial acetic acid.

Quinoline [91-22-5] M 129.2, m -16°, b 111.5°, 236°/758mm, d 1.0937, n 1.625, pK²⁵ 4.80 (4.93). Dried with Na₂SO₄ and vac distd from zinc dust. Also dried by boiling with acetic anhydride, then fractionally distilling. Calvin and Wilmarth [*J Am Chem Soc* 78 1301 1956] cooled redistd quinoline in ice and added enough HCl to form its hydrochloride. Diazotization removed aniline, the diazo compound being broken down by warming the soln to 60°. Non-basic impurities were removed by ether extraction. Quinoline was liberated by neutralising the hydrochloride with NaOH, then dried with KOH and fractionally distd at low pressure. Addition of cuprous acetate (7g/L of quinoline) and shaking under hydrogen for 12h at 100° removed impurities due to the nitrous acid treatment. Finally the hydrogen was pumped off and the quinoline was distd. Other purification procedures depend on conversion to the phosphate (m 159°, pptd from MeOH soln, filtered, washed with MeOH, then dried at 55°) or the picrate (m 201°) which, after crystn were reconverted to the amine. The method using the picrate [Packer, Vaughan and Wong *J Am Chem Soc* 80 905 1958] is as follows: quinoline is added to picric acid dissolved in the minimum volume of 95% EtOH, giving yellow crystals which were washed with EtOH, air-dried and crystd from acetonitrile. These were dissolved in dimethyl sulfoxide (previously dried over 4A molecular sieves) and passed through basic alumina, on which the picric acid is adsorbed. The free base in the effluent is extracted with *n*-pentane and distd under vacuum. Traces of solvent can be removed by vapour-phase chromatography. [Moonaw and Anton *J Phys Chem* 80 2243 1976.] The ZnCl₂ and dichromate complexes have also been used. [Cumper, Redford and Vogel *J Chem Soc* 1176 1962.]

2-Quinolinealdehyde [5470-96-2] M 157.2, m 71°, pK_{Est} ~3.3. Steam distd. Crystd from H₂O. Protected from light.

8-Quinolinecarboxylic acid [86-59-9] M 173.2, m 186-187.5°, pK₁²⁵ 1.82, pK₂²⁵ 6.87. Crystd from water.

Quinoline ethiodide (1-ethylquinolinium iodide) [634-35-5] M 285.1, m 158-159°. Crystd from aqueous EtOH.

Quinoxaline [91-19-0] M 130.2, m 28° (anhydr), 37°(H₂O), b 108-110°/0.1mm, 140°/40mm, pK₁²⁰-5.52 (-5.8, dication), pK₂²⁵ 2.08 (monocation). Crystd from pet ether. Crystallises as the monohydrate on addition of water to a pet ether soln.

Quinoxaline-2,3-dithiol [1199-03-7] M 194.1, m 345°(dec), pK₁ 6.9, pK₂ 9.9. Purified by repeated dissolution in alkali and re-pptn by acetic acid.

***p*-Quinquephenyl (*p*-pentaphenyl)** [61537-20-0] M 382.5, m 388.5°. Recrystd from boiling dimethyl sulfoxide (b 189°, lowered to 110°). The solid obtained on cooling was filtered off and washed repeatedly with toluene, then with conc HCl. The final material was washed repeatedly with hot EtOH. It was also recrystd from pyridine, then sublimed *in vacuo*.

Quinuclidine (1-azabicyclo[2.2.2]octane) [100-76-5] M 111.2, m 158°(sublimes), pK²⁵ 10.95. Crystd from diethyl ether.

D-Raffinose ($5H_2O$) [17629-30-0 ($5H_2O$); 512-69-6 (*anhydr*)] M 594.5, m 80°, $[\alpha]_{546}^{20} +124^\circ$ (c 10, H_2O). Crystd from aqueous EtOH.

Rauwolfscine hydrochloride [6211-32-1] M 390.0, m 278-280°. Crystd from water.

Reductic acid (1,2-dihydroxycyclopent-1,2-en-3-one) [80-72-8] M 114.1, m 213°, $pK^{20} 4.72$. Crystd from ethyl acetate.

Rescinnamine [24815-24-5] M 634.7, m 238-239°(vac), $[\alpha]_D^{20} -97^\circ$ (c 1, $CHCl_3$), $pK_{Est(1)} < 0$ (carbazole N), $pK_{Est(2)} \sim 7.0$ (quinolizidine N). Crystd from *benzene or MeOH.

Reserpic acid [83-60-3] M 400.5, m 241-243°, $pK_{Est(1)} < 0$ (carbazole N), $pK_{Est(2)} \sim 4.0$ (CO_2H), $pK_{Est(3)} \sim 7.4$ (quinolizidine N). Crystd from MeOH. The *hydrochloride* $0.5H_2O$ has m 257-259°, $[\alpha]_D^{23} -81^\circ$ (H_2O).

Reserpine [50-55-5] M 608.7, m 262-263°, $[\alpha]_{546}^{20} -148^\circ$ (c 1, $CHCl_3$), $pK_{Est(1)} < 0$ (carbazole N), $pK_2 6.6$ (7.4)(quinolizidine N). Crystd from aq acetone.

Resorcinol [108-46-3] M 110.1, m 111.2-111.6°, $pK_1^{25} 9.23$, $pK_2^{25} 13.05$. Crystd from *benzene, toluene or *benzene/diethyl ether.

Retene [483-65-8] M 234.3, m 99°. Crystd from EtOH.

Retinal (vitamin A aldehyde), **Retinoic acid (vitamin A acid)**, **Retinol (vitamin A alcohol)** see entries in Chapter 6.

Retinyl acetate [127-47-9] M 328.5, m 57°. Separated from retinol by column chromatography, then crystd from MeOH. See Kofler and Rubin [*Vitamins and Hormones (NY)* 18 315 1960] for review of purification methods. Stored in the dark, under N_2 or Ar, at 0°. See Vitamin A acetate p. 574 in Chapter 6.

Retinyl palmitate [79-81-2] M 524.9, m 28-29°, $\epsilon_{1cm}^{1\%}$ (*all-trans*) 1000 (325 nm) in EtOH. Separated from retinol by column chromatography on water-deactivated alumina with hexane containing a very small percentage of acetone. Also chromatographed on TLC silica gel G, using pet ether/isopropyl ether/acetic acid/water (180:20:2:5) or pet ether/acetonitrile/acetic acid/water (190:10:1:15) to develop the chromatogram. Then recrystd from propylene at low temperature.

Rhamnetin (3,3'-4',5-tetrahydroxy-7-methoxy flavone, 7-methyl quercitin) [90-19-7] M 316.3, m $>300^\circ$ (dec), several phenolic $pKs \sim 7-10.5$. Crystd from EtOH.

L- α -Rhamnose (H_2O) [10030-85-0 (H_2O); 3615-41-6 (*anhydr*)] M 182.2, m 105°, $[\alpha]_D^{15} +9.1^\circ$ (c 5, H_2O). Crystd from water or EtOH.

Rhodamine B chloride [3,5-bis-(diethylamino)-9-(2-carboxyphenyl)xanthylium chloride] [81-88-9] M 479.0, m 210-211°(dec), CI 45170, λ_{max} 543nm, {Free base [509-34-2] CI 749}, $pK 5.53$. Major impurities are partially dealkylated compounds not removed by crystn. Purified by chromatography, using ethyl acetate/isopropanol/ammonia (conc)(9:7:4, R_F 0.75 on Kieselgel G). Also crystd from conc soln in MeOH by slow addition of dry diethyl ether; or from EtOH containing a drop of conc HCl by slow addition of ten volumes of dry diethyl ether. The solid was washed with ether and air dried. The dried material has also been extracted with *benzene to remove oil-soluble material prior to recrystn. Store in the dark.

Rhodamine 6G [Basic Red 1, 3,5-bis-(ethylamino)-9-(2-ethoxycarbonylphenyl)-2,7-dimethylxanthylium chloride] [989-38-8] M 479.3, CI 45160, λ_{max} 524nm, $pK 5.58$. Crystd from MeOH or EtOH, and dried in a vac oven.

Rhodanine (2-mercaptothiazolidin-4-one) [141-84-4] M 133.2, m 168.5° (capillary), pK²⁰ 5.18. Crystd from glacial acetic acid or water.

Riboflavin, riboflavin-5'-phosphate (Na salt, 2H₂O) and **ribonucleic acid** see entries in Chapter 6.

α-D-Ribose [50-69-1] M 150.1, m 90°, [α]₅₄₆²⁰ -24° (after 24h, c 10, H₂O), pK²⁵ 12.22. Crystd from aqueous 80% EtOH, dried under vacuum at 60° over P₂O₅ and stored in a vacuum desiccator.

Ricinoleic acid (dl 12-hydroxyoleic acid) [141-22-0] M 298.5, m 7-8° (α-form), 5.0° (γ-form), n 1.4717, pK_{Est} ~4.5. Purified as methyl acetylricinoleate [Rider *J Am Chem Soc* 53 4130 1931], fractionally distilling at 180-185°/0.3mm, then 87g of this ester was refluxed with KOH (56g), water (25mL), and MeOH (250mL) for 10min. The free acid was separated, crystd from acetone at -50°, and distd in small batches, b 180°/0.005mm. [Bailey et al. *J Chem Soc* 3027 1957.]

Rosaniline HCl (Magenta I, Fuschin) [632-99-5] M 337.9, m >200°(dec). Purified by dissolving in EtOH, filtering and adding H₂O. Filter or centrifuge and wash the ppte with Et₂O and dry in air. Could be crystd from H₂O. Also recrystd from water and dried *in vacuo* at 40°. Crystals have a metallic green lustre. UV max in EtOH is at 543nm (ε 93,000). Solubility in H₂O is 0.26%. A carmine red colour is produced in EtOH. [Scalan *J Am Chem Soc* 57 887 1937.]

p-Rosolic acid (4-[bis-(4-hydroxyphenyl)methylene]-2,5-cyclohexadien-one, 4',4''-di-hydroxy-fuschson, aurin, corallin) [603-45-2] M 290.3, m 292°, 295-300° (dec with liberation of phenol), 308-310°(dec), pK₁ 3.11, pK₂ 8.62. It forms green crystals with a metallic lustre but the colour depends on the solvent used. When recrystd from brine (satd aqueous NaCl) acidified with HCl it forms red needles, but when recrystd from EtOH-AcOH the crystals have a beetle iridescent green colour. It has been recrystd from Me₂CO (although it dissolves slowly), methyl ethyl ketone, 80-95% AcOH and from AcOH·C₆H₆. An aq KOH soln is golden yellow and a 70% H₂SO₄ soln is deep red in colour. An alternative purification is to dissolve this triphenylmethane dye in 1.5% of aq NH₃, filter, and heat to 70-80°, then acidify with dilute AcOH by adding it slowly with vigorous stirring, whereby the aurin separates as a brick-red powder or as purplish crystals depending on the temperature and period of heating. Filter off the solid, wash with H₂O and a little dilute AcOH then H₂O again. Stir this solid with Et₂O to remove any ketones and allow to stand overnight in the Et₂O, then filter and dry in air then in a vacuum. [Gomberg and Snow *J Am Chem Soc* 47 202 1925; Baines and Driver *J Chem Soc* 123 1216 1923; UV: Burawoy *Chem Ber* 64 462 1941; Neuk and Schmid *J Prakt Chem* [2] 23 549 1881.]

Rubijervine (slanid-5-ene-3β-12α-diol) [79-58-3] M 413.6, m 240-246°, [α]_D²⁰ +19° (EtOH), pK_{Est} ~7.0. Crystd from EtOH. It has solvent of crystn.

Rubrene [517-51-1] M 532.7, m >315°. See 5,6,11,12-tetraphenylnaphthacene on p. 366.

(+)-Rutin (quercetin-3-rubinoside) [153-18-4] M 610.5, m 188-189, [α]₅₄₆²⁰ +13° (c 5, EtOH) (polyphenolic flavone pKs 7-10). Crystd from MeOH or water/EtOH, air dried, then dried for several hours at 110°.

Saccharic acid (D-glucaric acid) [87-73-0] M 210.1, m 125-126°, [α]_D²⁰ +6.9° → +20.6° (H₂O), pK₁ 3.01, pK₂ 3.94 (D-isomer). Crystd from 95% EtOH.

Safranine O [477-73-6] M 350.9, λ_{max} 530nm, pK²⁵ 6.4. Crystd from *benzene/MeOH (1:1) or water. Dried under vacuum over H₂SO₄.

Safrole (5-allyl-1,3-benzodioxole, 4-allyl-1,2-methylenedioxybenzene) [94-59-7] M 162.1, m~ 11°, b 69-70°/1.5mm, 104-105°/6mm, 231.5-232°/atm, 235-237°/atm, d_4^{20} 1.0993, n_D^{20} 1.53738. It has been purified by fractional distn, although it has also been recrystd from low boiling pet ether at low temperatures. [IR: Briggs et al. *Anal Chem* 29 904 1957; UV: Patterson and Hibbert *J Am Chem Soc* 65 1962 1943.] The *maleic anhydride adduct* forms yellow crystals from toluene m 257° [Hickey *J Org Chem* 13 443 1948], and the *picrate* forms orange-red crystals from CHCl₃ [Baril and Magrdichian *J Am Chem Soc* 58 1415 1936].

D(-)-Salicin [138-52-3] M 286.3, m 204-208°, $[\alpha]_D^{25}$ -63.5° (c ca 3, H₂O). Crystd from EtOH.

Salicylaldehyde (*o*-hydroxybenzaldehyde) [90-02-8] M 122.1, b 93°/25mm, 195-197°/760mm, d 1.167, n 1.574, pK^{25} 8.37. Pptd as the bisulfite addition compound by pouring the aldehyde slowly and with stirring into a 25% soln of NaHSO₃ in 30% EtOH, then standing for 30min. The ppte, after filtering at the pump, and washing with EtOH, was decomposed with aq 10% NaHCO₃, and the aldehyde was extracted into diethyl ether, dried with Na₂SO₄ or MgSO₄, and distd, under reduced pressure. Alternatively, salicylaldehyde can be pptd as its copper complex by adding it to warm, satd soln of copper acetate, shaking and then standing in ice. The ppte was filtered off, washed thoroughly with EtOH, then with diethyl ether, and decomposed with 10% H₂SO₄, the aldehyde was extracted into diethyl ether, dried and distd. It has also been purified by repeated vacuum distn, and by dry column chromatography on Kieselgel G [Nishiya et al. *J Am Chem Soc* 108 3880 1986]. The *acetyl* derivative has m 38-39° (from pet ether or EtOH) and b 142°/18mm, 253°/atm.

Salicylaldoxime [94-67-7] M 137.1, m 57°, pK_{Est} ~ 8.3. Crystd from CHCl₃/pet ether (b 40-60°).

Salicylamide [65-45-2] M 137.1, m 142-144°, pK^{20} 8.37. Crystd from water or repeatedly from chloroform [Nishiya et al. *J Am Chem Soc* 108 3880 1986].

Salicylanilide [87-17-2] M 213.2, m 135°, pK_{Est} ~8.3. Crystd from water.

Salicylhydroxamic acid [89-73-6] M 153.1, m 179-180°(dec), pK_1^{30} 2.15, pK_2^{30} 7.46, pK_3^{30} 9.72. Crystd from acetic acid.

Salicyclic acid (2-hydroxybenzoic acid) [69-72-7] M 138.1, m 157-159°, 158-160°, 159.5°, 159-160°, 162°, b 211°/20mm, pK_1^{25} 3.01, pK_2^{25} 13.43 (13.01). It has been purified by steam distn, by recrystn from H₂O (solubility is 0.22% at room temp and 6.7% at 100°), absolute MeOH, or cyclohexane and by sublimation in a vacuum at 76°. The *acid chloride* (needles) has m 19-19.5°, b 92°/15mm, *amide* m 133° (yellow needles from H₂O), and *anilide* (prisms from H₂O) m 135°. The *O-acetyl* derivative has m 135° (rapid heating and the liquid resolidifies at 118°) and the *o-benzoyl* derivative has m 132° (aq EtOH). [IR: Hales et al. *J Chem Soc* 3145 1954; UV: Bergmann et al. *J Chem Soc* 2351 1950].

Sarcosine [107-97-1] M 89.1, m 212-213°(dec), pK_1^{20} 2.12, pK_2^{20} 10.19. Crystd from abs EtOH.

Scopoletin (7-hydroxy-6-methoxycoumarin) [92-61-5] M 192.2, m 206°, 208-209°, pK 8.96 (70%aq EtOH). Crystd from water, acetic acid or *C₆H₆/MeOH.

Sebacic acid [111-20-6] M 202.3, m 134.5°, pK_1^{25} 4.58, pK_2^{25} 5.54. Purified *via* the disodium salt which, after crystn from boiling water (charcoal), was again converted to the free acid. The free acid was crystd repeatedly from hot distd water or from Me₂CO+pet ether and dried under vacuum.

Sebacic acid monomethyl ester [818-88-2] M 216.3, m 42-43°, b 169-171°/4mm. Recrystd from Me₃CO+pet ether or pet ether at low temperature and distd in a vacuum.

Sebaconitrile (decanedinitrile) [1871-96-1] M 164.3, m 8°, b 199-200°. Mix with P₂O₅ (10% by wt) and distilled from it, then redistilled.

Secobarbital (5-allyl-5-1'-methylbutylbarbituric acid) [76-73-3] M 260.3, m 100°, $pK_{Est(1)} \sim 3.5$, $pK_{Est(2)} \sim 12.0$. A soln of the salt in 10% HCl was ppted and the acid form was extracted by the addition of ether. Then purified by repeated crystn from $CHCl_3$. [Buchet and Sandorfy *J Phys Chem* 88 3274 1984.]

Semicarbazide hydrochloride (hydrazine carboxamide hydrochloride) [563-41-7] M 111.5, m 173°(dec), 175°(dec), $pK^{24} 3.66$. Crystd from aqueous 75% EtOH and dried under vacuum over $CaSO_4$. Also crystd from a mixture of 3.6 mole % MeOH and 6.4 mole % of water. [Kovach et al. *J Am Chem Soc* 107 7360 1985.] IR: ν 700, 3500 cm^{-1} [*Org Synth Coll Vol I* 485 1941; Davison and Christie *J Chem Soc* 3389 1955; Thiele and Stange *Chem Ber* 27 33 1894; pK: Bartlett *J Am Chem Soc* 54 2853 1923]. The free base crystd as prisms from abs EtOH, m 96° [Curtius and Heidenreich *Chem Ber* 27 55 1894]. **TOXIC ORALLY**, possible **CARCINOGEN** and **TERATOGEN**.

Sennoside A [81-27-6] M 862.7, m 220-240°(dec), $[\alpha]_D^{20} -147^\circ$ (c 5, Me_2CO/H_2O 7:1). Crystd from aq acetone or large vols of H_2O .

Sennoside B [128-57-4] M 962.7, m 182-190°(dec), $[\alpha]_D^{20} -100^\circ$ (c 2, Me_2CO/H_2O 7:3). Crystd from aq acetone or large vols of H_2O .

L-Serine [56-45-1] M 105.1, m 228°(dec), $[\alpha]_D^{25} +14.5^\circ$ (1M HCl), $[\alpha]_{546}^{20} +16^\circ$ (c 5, 5M HCl), $pK_1^{25} 2.15$, $pK_2^{25} 9.21$. Likely impurity is glycine. Crystd from water by adding 4 volumes of EtOH. Dried. Stored in a desiccator.

Serotonin creatinine sulfate (H_2O) [971-74-4] M 405.4, m 220°(dec), $pK_1 10.1$, $pK_2 11.1$, $pK_3 18.25$ (NH) for serotonin, $pK 4.9$ for creatinine. Crystd (as monohydrate) from water.

Shikimic acid [138-59-0] M 174.2, m 183-184.5°, 190°, $[\alpha]_{546}^{20} -210^\circ$ (c 2, H_2O), $pK^{14} 4.15$. Crystd from water or MeOH/AcOEt and sublimes in a vac.

Sinomenine hydrochloride (7,8-didehydro-4-hydroxy-3,7-dimethoxy-17-methyl-9 α ,13 α ,14 α -morphan-6-one HCl) [6080-33-7] M 365.9, m 231°, $[\alpha]_D^{17} -83^\circ$ (c 4, H_2O), $pK_{Est(1)} \sim 10.0$ (N), $pK_{Est(2)} \sim 10.4$ (OH). Crystd from water.

β -Sitosterol [83-46-5] M 414.7, m 136-137°, $[\alpha]_{546}^{20} -42^\circ$ (c 2, $CHCl_3$). Crystd from MeOH. Also purified by zone melting.

Skellysolve A is essentially *n*-pentane, b 28-30°,

Skellysolve A is essentially *n*-hexane, b 60-68°,

Skellysolve C is essentially *n*-heptane, b 90-100°,

Skellysolve D is mixed heptanes, b 75-115°,

Skellysolve E is mixed octanes, b 100-140°,

Skellysolve F is pet ether, b 30-60°,

Skellysolve G is pet ether, b 40-75°,

Skellysolve H is hexanes and heptanes, b 69-96°,

Skellysolve L is essentially octanes, b 95-127°. For methods of purification, see **petroleum ether**.

Smilagenin [126-18-1] M 416.6, m 185°, $[\alpha]_D^{25} -69^\circ$ (c 0.5, $CHCl_3$). Crystd from acetone.

Solanidine [80-78-4] M 397.6, m 218-219°, $[\alpha]_D^{20} -29^\circ$ (c 0.5, $CHCl_3$), $pK^{15} 6.66$. Crystd from $CHCl_3/MeOH$.

α -Solanine [20562-02-1] M 868.1, m 286°(dec), $[\alpha]_D^{20} -58^\circ$ (c 0.8, pyridine), $pK^{15} 6.66$.

See α -solanine on p. 566 in Chapter 6.

Solanone [*S(+)-trans-2-methyl-5-isopropyl-1,3-nonan-8-one*] [1937-54-8] M 194.3, b 60°/1mm, $[\alpha]_{\text{D}}^{20} +14^{\circ}$ (neat). Purified by high vacuum distillation and stored in sealed ampules [Kohda and Sato *J Chem Soc, Chem Commun* 951 1981]. It has UV (hexane) at λ_{max} 230nm (ϵ 11,800).

Solasodine [126-17-0] M 413.6, m 202°, $[\alpha]_{\text{D}}^{25} -100^{\circ}$ (c 2, MeOH), pK 7.7. Crystd (as monohydrate) from MeOH or aq 80% EtOH, and sublimes in a vac.

Solasonine (solasodine-3-O-mannoglucoside) [19121-58-5] M 884.0, m 279°, $[\alpha]_{\text{D}}^{20} -75^{\circ}$ (c 0.5, MeOH), pK_{Est} ~ 7.7. Crystd from aq 80% dioxane or MeOH.

Solochrome Violet R [4-hydroxy-3-(2-hydroxynaphthyl-1-ylazo)benzenesulfonic acid] [2092-55-9] M 367.3, CI 15670, λ_{max} 501nm, pK₂²⁵ 7.22 (OH), pK₃²⁵ 13.39 (OH). Converted to the monosodium salt by pptn with NaOAc/AcOH buffer of pH 4, then purified by pptn of the free acid from aq soln with conc HCl, washing and extracting with EtOH in a Soxhlet extractor. The acid ppted on evaporating the EtOH and was reconverted to the sodium salt as described for *Chlorazole Sky Blue FF*. Dried at 110°. It is *hygroscopic*. [Coates and Rigg *Trans Faraday Soc* 57 1088 1961.]

Sorbic acid (2,4-hexadienoic acid) [110-44-1] M 112.1, m 134°, pK²⁵ 4.76. Crystd from water.

Sorbitol [50-70-4] M 182.2, m 89-93° (hemihydrate), 110-111° (anhydrous), $[\alpha]_{546}^{20} -1.8^{\circ}$ (c 10, H₂O), pK⁶⁰ 13.00. Crystd (as hemihydrate) several times from EtOH/water (1:1), then dried by fusing and storing over MgSO₄.

(-)-Sparteine sulfate pentahydrate [6160-12-9] M 422.5, m loses H₂O at 100° and turns brown at 136° (dec), $[\alpha]_{\text{D}}^{20} -22^{\circ}$ (c 5, H₂O), $[\alpha]_{\text{D}}^{21} -16^{\circ}$ (c 10, EtOH for free base), pK₁²⁰ 2.24, pK₂²⁰ 9.46. Recrystd from aq EtOH or H₂O although the solubility in the latter is high. The *free (-)-base* has b 173°/8mm and is steam volatile but resinifies in air. The *dipicrate* forms yellow needles from EtOH-Me₂CO, m 205-206° [Clemo et al. *J Chem Soc* 429 1931; see also Bolnmann and Schuman *The Alkaloids* (Ed Manske) Vol 9 175 1967]. The *free (±)-base* has m 71-72.5° [van Tamelen and Foltz *J Am Chem Soc* 82 2400 1960].

Spermidine [*N*-(3-aminopropyl)-1,4-diaminobutane] [124-20-9] M 145.3, m 23-25°, b 128-131°/15mm, d 0.918, n 1.482, pK₁²⁵ 8.25, pK₂²⁵ 9.64, pK₃²⁵ 10.43. It is a strong base with an alkylamine odour and absorbs CO₂ from the atmosphere. It is purified by shaking with solid K₂CO₃ or NaOH, decanting and distilling from K₂CO₃ in a vacuum. Store in the dark under N₂.

Spermidine trihydrochloride [334-50-9] M 245.3, m ~250°(dec), 256-258°, for pKa see *free base* above. Recrystd from dry 3% HCl in ethanol adding dry Et₂O if necessary. Filter rapidly and dry in a vac desiccator. Alternatively centrifuge the crystals off wash them with dry Et₂O and dry in a vacuum.

Spermine 4HCl (*N,N*-bis(3-aminopropyl)-1,4-butanediamine 4HCl) [306-67-2] M 348.2, m 313-315°. The pKs are similar to spermidine above. Purification as for spermidine trihydrochloride above.

Squalene (*all-trans-2,6,10,15,19,23-hexamethyltetracosahexa-2,6,10,14,18,22-ene, spinacen*) [111-02-4] M 410.7, m ~75°, b 203°/0.15mm. See squalene on p. 567 in Chapter 6.

Squaric acid (3,4-dihydroxy-3-cyclobutene-1,2-dione) [2892-51-5] M 114.1, m 293°(dec), 294°(dec), >300°, pK₁²⁰ 1.50, pK₂²⁰ 2.93. Purified by recryst from H₂O — this is quite simple because the acid is ~ 7% soluble in boiling H₂O and only 2% at room temperature. It is not soluble in Me₂CO or Et₂O hence it can be rinsed with these solvents and dried in air or a vacuum. It is not hygroscopic and gives an intense purple colour with FeCl₃. It has IR ν at 1820 (C=O) and 1640 (C=C) cm⁻¹; and UV λ_{max} at 269.5nm (ϵ 37K M⁻¹cm⁻¹.) [Cohn et al. *J Am Chem Soc* 81 3480 1959; Park et al. *J Am Chem Soc* 84 2919 1962] See also pKa values of 0.59 ± 0.09 and 3.48 ± 0.023 [Scwartz and Howard *J Phys Chem* 74 4374 1970].

Starch [9005-84-9] M (162.1)n. See entry on p. 567 in Chapter 6.

Stearic acid (octadecanoic acid) [57-11-4] M 284.5, m 71.4°, 72°, b 144-145°/27mm, 383°/760mm, d 0.911, n 1.428, $pK_{\text{Est}} \sim 5.0$. Crystd from acetone, acetonitrile, EtOH (5 times), aq MeOH, ethyl methyl ketone or pet ether (b 60-90°), or by fractional pptn by dissolving in hot 95% EtOH and pouring into distd water, with stirring. The ppte, after washing with distd water, was dried under vacuum over P₂O₅. It has also been purified by zone melting and partial freezing. [Tamai et al. *J Phys Chem* 91 541 1987.]

Stigmasterol [83-48-7] M 412.7, m 170°, $[\alpha]_{\text{D}}^{22} -51^\circ$ (CHCl₃), $[\alpha]_{546}^{20} -59^\circ$ (c 2, CHCl₃). Crystd from hot EtOH. Dried in vacuum over P₂O₅ for 3h at 90°. Purity was checked by NMR.

cis-Stilbene [645-49-8] M 180.3, b 145°/12mm. Purified by chromatography on alumina using hexane and distd under vacuum. (The final product contains ca 0.1% of the *trans*-isomer). [Lewis et al. *J Am Chem Soc* 107 203 1985; Saltiel *J Phys Chem* 91 2755 1987.]

trans-Stilbene [103-30-0] M 180.3, m 125.9°, b 305-307°/744mm, d 0.970. Purified by vac distn. (The final product contains about 1% of the *cis* isomer). Crystd from EtOH. Purified by zone melting. [Lewis et al. *J Am Chem Soc* 107 203 1985; Bollucci et al. *J Am Chem Soc* 109 515 1987; Saltiel *J Phys Chem* 91 2755 1987.]

(-)-Strychnine [57-24-9] M 334.4, m 268°, $[\alpha]_{546}^{20} -139^\circ$ (c 1, CHCl₃), $pK_1^{20} 2.50$, $pK_2^{20} 8.2$. Crystd as the hydrochloride from water, then neutralised with ammonia.

Styphnic acid (2,4,6-trinitroresorcinol) [82-71-3] M 245.1, m 177-178°, 179-180°, $pK_1^{25} 0.06$ (1.74), $pK_2^{25} 4.23$ (4.86). Crystd from ethyl acetate or water containing HCl [EXPLODES violently on rapid heating.]

Styrene [100-42-5] M 104.2, b 41-42°/18mm, 145.2°/760mm, d 0.907, n 1.5469, $n^{25} 1.5441$. Styrene is difficult to purify and keep pure. Usually contains added inhibitors (such as a trace of hydroquinone). Washed with aqueous NaOH to remove inhibitors (e.g. *tert*-butanol), then with water, dried for several hours with MgSO₄ and distd at 25° under reduced pressure in the presence of an inhibitor (such as 0.005% *p-tert*-butylcatechol). It can be stored at -78°. It can also be stored and kept anhydrous with Linde type 5A molecular sieves, CaH₂, CaSO₄, BaO or sodium, being fractionally distd, and distd in a vacuum line just before use. Alternatively styrene (and its deuterated derivative) were passed through a neutral alumina column before use [Woon et al. *J Am Chem Soc* 108 7990 1986; Collman *J Am Chem Soc* 108 2588 1986].

(±)-Styrene glycol (±-1-phenyl-1,2-ethanediol) [93-56-1] M 138.2, m 67-68°. Crystd from pet ether.

Styrene oxide [96-09-3] M 120.2, b 84-86°/16.5mm, d 1.053, n 1.535. Fractional distn at reduced pressure does not remove phenylacetaldehyde. If this material is present, the styrene oxide is treated with hydrogen under 3 atmospheres pressure in the presence of platinum oxide. The aldehyde, but not the oxide, is reduced to β-phenylethanol) and separation is now readily achieved by fractional distn. [Schenck and Kaizermen *J Am Chem Soc* 75 1636 1953.]

Suberic acid (hexane-1,6-dicarboxylic acid) [505-48-6] M 174.2, m 141-142°, $pK_1^{25} 4.12$, $pK_2^{25} 5.40$. Crystd from acetone and sublimes at 300° without dec.

Succinamic acid (succinic acid amide) [638-32-4] M 117.1, m 155°, 156-157°, $pK_1^{25} 4.54$. Crystd from Me₂CO or H₂O and dried in vac. Not v sol in MeOH. Converted to succinimide above 200°.

Succinamide [110-14-5] M 116.1, m 262-265°(dec). Crystd from hot water.

Succinic acid [110-15-6] M 118.1, m 185-185.5°, $pK_1^{25} 4.21$, $pK_2^{25} 5.72$. Washed with diethyl ether. Crystd from acetone, distd water, or *tert*-butanol. Dried under vacuum over P₂O₅ or conc H₂SO₄. Also

purified by conversion to the disodium salt which, after crystn from boiling water (charcoal), is treated with mineral acid to regenerate the succinic acid. The acid is then recrystd and vacuum dried.

Succinic anhydride [108-30-5] M 100.1, m 119-120°. Crystd from redistd acetic anhydride or CHCl_3 , then filtered, washed with diethyl ether and dried under vacuum.

Succinimide [123-56-8] M 99.1, m 124-125°, pK^{25} 9.62. Crystd from EtOH (1mL/g) or water.

Succinonitrile [110-61-2] M 80.1, m 57.9°, b 108°/1mm, 267°/760mm. Purified by vacuum sublimation, also crystd from acetone.

D(+)-Sucrose (β -D-fructofuranosyl- α -D-glucopyranoside) [57-50-1] M 342.3, m 160-186°, 186-188°, $[\alpha]_{546}^{20} +78^\circ$ (c 10, H_2O), $[\alpha]_{\text{D}}^{20} +66^\circ$ (c 26, H_2O), pK 12.62. Crystd from water (solubility: 1g in 0.5mL H_2O at 20°, 1g in 0.2mL in boiling H_2O). Sol in EtOH (0.6%) and MeOH (1%).
Sucrose diacetate hexaisobutyrate is purified by melting and, while molten, treated with NaHCO_3 and charcoal, then filtered.

D-Sucrose octaacetate [126-14-7] M 678.6, m 83-85°, $[\alpha]_{546}^{20} +70^\circ$ (c 1, CHCl_3). Crystd from EtOH.

Sudan I (Solvent Yellow 14, 1-phenylazo-2-naphthol) [824-07-9] M 248.3, m 135°, CI 12055, λ_{max} 476nm, $\text{pK}_{\text{Est}} \sim 9.0$. Crystd from EtOH.

Sudan III [Solvent Red 23, 1-(*p*-phenylazo-phenylazo)-2-naphthol] [85-86-9] M 352.4, m 199°(dec), CI 26100, λ_{max} 354, 508 nm, $\text{pK}_{\text{Est}} \sim 9.0$. Crystd from EtOH, EtOH/water or *benzene/abs EtOH (1:1).

Sudan IV [Solvent Red 24, 1-(4-*o*-tolylazo-*o*-tolylazo)-2-naphthol] [85-83-6] M 380.5, m ~184°(dec), CI 26105, λ_{max} 520nm, $\text{pK}_{\text{Est}} \sim 9.0$. Crystd from EtOH/water or acetone/water.

Sulfaguanidine [57-67-0] M 214.2, m 189-190°, pK_1 0.48, pK_2 2.75. Crystd from hot water (7mL/g).

Sulfamethazine [57-68-1] M 278.3, m 198-200°, pK_1 2.65, pK_2 7.4. Crystd from dioxane.

Sulfanilamide (*p*-aminobenzenesulfonamide) [63-74-1] M 172.2, m 166°, pK_1^{20} 2.30, pK_2^{20} 10.26. Crystd from water or EtOH.

Sulfanilic acid (4-aminobenzenesulfonic acid) [121-57-3] M 173.2, $\text{pK}_1^{25} <1$, pK_2^{25} 3.23. Crystd (as dihydrate) from boiling water. Dried at 105° for 2-3h, then over 90% H_2SO_4 in a vacuum desiccator.

Sulfapyridine [144-83-2] M 349.2, m 193°, pK^{20} 8.64. Crystd from 90% acetone and dried at 90°.

***o*-Sulfobenzoic acid (H_2O)** [123333-68-6 (H_2O); 632-25-7] M 202.2, m 68-69°, $\text{pK}_{\text{Est}(1)} <1$, $\text{pK}_{\text{Est}(2)} \sim 3.1$ (CO_2H). Crystd from water.

***o*-Sulfobenzoic acid (monoammonium salt)** [6939-89-5] M 219.5. Crystd from water.

***o*-Sulfobenzoic anhydride** [81-08-3] M 184.2, m 128°, b 184-186°/18mm. See also 2,1-benzoxathiol-3-one-1,1-dioxide on p. 126.

Sulfolane (tetramethylenesulfone) [126-33-0] M 120.2, m 28.5°, b 153-154°/18mm, 285°/760mm, d 1.263, n^{30} 1.4820. Prepared commercially by Diels-Alder reaction of 1,3-butadiene and sulfur dioxide, followed by Raney nickel hydrogenation. The principle impurities are water, 3-sulfolene, 2-sulfolene and 2-isopropyl sulfolanyl ether. It is dried by passage through a column of molecular sieves. Distd

under reduced pressure through a column packed with stainless steel helices. Again dried with molecular sieves and distd. [Cram et al. *J Am Chem Soc* **83** 3678 1961; Coetzee *Pure Appl Chem* **49** 211 1977.]

Also, it was stirred at 50° and small portions of solid KMnO_4 were added until the colour persisted during 1h. Dropwise addition of MeOH then destroyed the excess KMnO_4 , the soln was filtered, freed from potassium ions by passage through an ion-exchange column and dried under vacuum. It has also been vacuum distd from KOH pellets. It is *hygroscopic*. [See Sacco et al. *J Phys Chem* **80** 749 1976; *J Chem Soc, Faraday Trans 1* **73** 1936 1977; **74** 2070 1978; *Trans Faraday Soc* **62** 2738 1966.] Coetzee has reviewed the methods of purification of sulfolane, and also the removal of impurities. [Coetzee in *Recommended Methods of Purification of Solvents and Tests for Impurities*, Coetzee Ed. Pergamon Press, 1982.]

5-Sulfosalicylic acid [5965-83-3] M 254.2, m 108-110°, $\text{pK}_1^{25} < 0$, $\text{pK}_2^{25} 2.67$, $\text{pK}_3^{25} 11.67$. Crystd from water. Alternatively, it was converted to the monosodium salt which was crystd from water and washed with a little water, EtOH and then diethyl ether. The free acid was recovered by acidification.

Syringaldehyde (3,5-dimethoxy-4-hydroxybenzaldehyde) [134-96-3] M 182.2, m 113°, $\text{pK}_{\text{Est}} \sim 8$. Crystd from pet ether.

Syringic acid (3,5-dimethoxy-4-hydroxybenzoic acid) [530-57-4] M 198.2, m 204-205°, 206.5°, 206-209°, 209-210°, $\text{pK}_1^{25} 4.34$, $\text{pK}_2^{25} 9.49$. Recrystd from H_2O using charcoal [Bogert and Coyne *J Am Chem Soc* **51** 571 1929; Anderson and Nabenhauer *J Am Chem Soc* **48** 3001 1926.] The *methyl ester* has m 107° (from MeOH), the *4-acetyl* derivative has m 190° and the *4-benzoyl* derivative has m 229-232°. [Hahn and Wassmuth *Chem Ber* **67** 2050 1934; UV: Lemon *J Am Chem Soc* **69** 2998 1947 and Pearl and Beyer *J Am Chem Soc* **72** 1743 1950.]

D(-)-Tagatose [87-81-0] M 180.2, m 134-135°, $[\alpha]_{546} -6.5^\circ$ (c 1, H_2O). Crystd from aqueous EtOH.

d-Tartaric acid [147-71-7] M 150.1, m 169.5-170° (2*S*,3*S*-form, natural) $[\alpha]_{546}^{20} -15^\circ$ (c 10, H_2O); m 208° (2*RS*,3*RS*-form), $\text{pK}_1^{25} 3.03$, $\text{pK}_2^{25} 4.46$, $\text{pK}_3^{25} 14.4$. Crystd from distilled H_2O or *benzene/diethyl ether containing 5% of pet ether (b 60-80°) (1:1). Soxhlet extraction with diethyl ether has been used to remove an impurity absorbing at 265nm. It has also been crystd from absolute EtOH/hexane, and dried in a vacuum for 18h [Kornblum and Wade *J Org Chem* **52** 5301 1987].

meso-Tartaric acid [147-73-9] M 150.1, m 139-141°, $\text{pK}_1^{25} 3.17$, $\text{pK}_2^{25} 4.91$. Crystd from water, washed with cold MeOH and dried at 60° under vacuum.

Taurocholic acid [81-24-3] M 515.6, m 125°(dec), $[\alpha]_{\text{D}} +38.8$ (c 2, EtOH), $\text{pK} 1.4$. Crystd from EtOH/diethyl ether.

Terephthalaldehyde [623-27-8] M 134.1, m 116°, b 245-248°/771mm. Crystd from water.

Terephthalic acid (benzene-1,4-dicarboxylic acid) [100-21-0] M 166.1, m sublimes >300° without melting, $\text{pK}_1^{20} 3.4$, $\text{pK}_2^{20} 4.34$. Purified *via* the sodium salt which, after crystn from water, was reconverted to the acid by acidification with mineral acid.

Terephthaloyl chloride [100-20-9] M 203.0, m 80-82°. Crystd from dry hexane.

o-Terphenyl [84-15-1] M 230.3, m 57-58°. Crystd from EtOH. Purified by chromatography of CCl_4 solns on alumina, with pet ether as eluent, followed by crystn from pet ether (b 40-60°) or pet ether/*benzene. They can also be distd under vacuum.

m-Terphenyl [92-06-8] M 230.3, m 88-89°. Purification as for *o*-terphenyl above.

[Previous Page](#)

under reduced pressure through a column packed with stainless steel helices. Again dried with molecular sieves and distd. [Cram et al. *J Am Chem Soc* **83** 3678 1961; Coetzee *Pure Appl Chem* **49** 211 1977.]

Also, it was stirred at 50° and small portions of solid KMnO_4 were added until the colour persisted during 1h. Dropwise addition of MeOH then destroyed the excess KMnO_4 , the soln was filtered, freed from potassium ions by passage through an ion-exchange column and dried under vacuum. It has also been vacuum distd from KOH pellets. It is *hygroscopic*. [See Sacco et al. *J Phys Chem* **80** 749 1976; *J Chem Soc, Faraday Trans 1* **73** 1936 1977; **74** 2070 1978; *Trans Faraday Soc* **62** 2738 1966.] Coetzee has reviewed the methods of purification of sulfolane, and also the removal of impurities. [Coetzee in *Recommended Methods of Purification of Solvents and Tests for Impurities*, Coetzee Ed. Pergamon Press, 1982.]

5-Sulfosalicylic acid [5965-83-3] M 254.2, m 108-110°, $\text{pK}_1^{25} < 0$, $\text{pK}_2^{25} 2.67$, $\text{pK}_3^{25} 11.67$. Crystd from water. Alternatively, it was converted to the monosodium salt which was crystd from water and washed with a little water, EtOH and then diethyl ether. The free acid was recovered by acidification.

Syringaldehyde (3,5-dimethoxy-4-hydroxybenzaldehyde) [134-96-3] M 182.2, m 113°, $\text{pK}_{\text{Est}} \sim 8$. Crystd from pet ether.

Syringic acid (3,5-dimethoxy-4-hydroxybenzoic acid) [530-57-4] M 198.2, m 204-205°, 206.5°, 206-209°, 209-210°, $\text{pK}_1^{25} 4.34$, $\text{pK}_2^{25} 9.49$. Recrystd from H_2O using charcoal [Bogert and Coyne *J Am Chem Soc* **51** 571 1929; Anderson and Nabenhauer *J Am Chem Soc* **48** 3001 1926.] The *methyl ester* has m 107° (from MeOH), the *4-acetyl* derivative has m 190° and the *4-benzoyl* derivative has m 229-232°. [Hahn and Wassmuth *Chem Ber* **67** 2050 1934; UV: Lemon *J Am Chem Soc* **69** 2998 1947 and Pearl and Beyer *J Am Chem Soc* **72** 1743 1950.]

D(-)-Tagatose [87-81-0] M 180.2, m 134-135°, $[\alpha]_{546} -6.5^\circ$ (c 1, H_2O). Crystd from aqueous EtOH.

d- Tartaric acid [147-71-7] M 150.1, m 169.5-170° (2*S*,3*S*-form, natural) $[\alpha]_{546}^{20} -15^\circ$ (c 10, H_2O); m 208° (2*R**S*,3*R**S*-form), $\text{pK}_1^{25} 3.03$, $\text{pK}_2^{25} 4.46$, $\text{pK}_3^{25} 14.4$. Crystd from distilled H_2O or *benzene/diethyl ether containing 5% of pet ether (b 60-80°) (1:1). Soxhlet extraction with diethyl ether has been used to remove an impurity absorbing at 265nm. It has also been crystd from absolute EtOH/hexane, and dried in a vacuum for 18h [Kornblum and Wade *J Org Chem* **52** 5301 1987].

meso-Tartaric acid [147-73-9] M 150.1, m 139-141°, $\text{pK}_1^{25} 3.17$, $\text{pK}_2^{25} 4.91$. Crystd from water, washed with cold MeOH and dried at 60° under vacuum.

Taurocholic acid [81-24-3] M 515.6, m 125°(dec), $[\alpha]_{\text{D}} +38.8$ (c 2, EtOH), $\text{pK} 1.4$. Crystd from EtOH/diethyl ether.

Terephthalaldehyde [623-27-8] M 134.1, m 116°, b 245-248°/771mm. Crystd from water.

Terephthalic acid (benzene-1,4-dicarboxylic acid) [100-21-0] M 166.1, m sublimes >300° without melting, $\text{pK}_1^{20} 3.4$, $\text{pK}_2^{20} 4.34$. Purified *via* the sodium salt which, after crystn from water, was reconverted to the acid by acidification with mineral acid.

Terephthaloyl chloride [100-20-9] M 203.0, m 80-82°. Crystd from dry hexane.

o-Terphenyl [84-15-1] M 230.3, m 57-58°. Crystd from EtOH. Purified by chromatography of CCl_4 solns on alumina, with pet ether as eluent, followed by crystn from pet ether (b 40-60°) or pet ether/*benzene. They can also be distd under vacuum.

m-Terphenyl [92-06-8] M 230.3, m 88-89°. Purification as for *o*-terphenyl above.

***p*-Terphenyl** [92-94-4] M 230.3, m 212.7°. Crystd from nitrobenzene or trichlorobenzene. It was purified by chromatography on alumina in a darkened room, using pet ether, and then crystallizing from pet ether (b 40-60°) or pet ether/*benzene.

Terpin hydrate [2451-01-6] M 190.3, m 105.5° (*cis*), 156-158° (*trans*). Crystd from H₂O or EtOH.

2,2':6',2''-Terpyridyl (2,2':6',2''-terpyridyl) [1148-79-4] M 233.3, m 91-92°, pK₁²³ 2.64, pK₂²³ 4.33. Crystd from diethyl ether, toluene or from pet ether, then aqueous MeOH, followed by vacuum sublimation at 90°.

Terreic acid (2-hydroxy-3-methyl-1,4-benzoquinone-5,6-epoxide) [121-40-4] M 154.1, m 127-127.5°, [α]_D²² +74° (pH 4, phosphate buffer), -17° (CHCl₃), pK 4.5. Crystd from *benzene or hexane. Sublimed *in vacuo*.

Terthiophene (2,5-di[thienyl]thiophene; α-terthienyl) [1081-34-1] M 248.4, m 94-95.5°, 94-96°. Possible impurities are bithienyl and polythienyls. Suspend in H₂O and steam distil to remove bithienyl. The residue is cooled and extracted with CHCl₃, dried (MgSO₄), filtered, evaporated and the residue chromatographed on Al₂O₃ using pet ether-3% Me₂CO as eluant. The terphenyl zone is then eluted from the Al₂O₃ with Et₂O, the extract is evaporated and the residue is recrystd from MeOH (40mL per g). The platelets are washed with cold MeOH and dried in air. [UV: Sease and Zechmeister *J Am Chem Soc* 69 270 1947; Uhlenbroek and Bijloo *Recl Trav Chim Pays-Bas* 79 1181 1960.] See also entry on p. 568 in Chapter 6.

Testosterone [58-22-0] M 288.4, m 155°, [α]₅₄₆²⁰ +130° (c 1, dioxane). Crystd from aq acetone.

Testosterone propionate [57-85-2] M 344.5, m 118-122°, [α]₅₄₆²⁰ +100° (c 1, dioxane). Crystd from aqueous EtOH.

2,4,5,6-Tetraaminopyrimidine sulfate [5392-28-9] M 238.2, m 255° (dec), >300°, >350° (dec), pK²⁰ 6.82. Purified by recrystn from H₂O, 2N H₂SO₄ (20 parts, 67% recovery) or 0.1N H₂SO₄ (40 parts, 62% recovery), and dried in air. [UV: Konrad and Pfeleiderer *Chem Ber* 103 722 1970; Malletta et al. *J Am Chem Soc* 69 1814 1947; Cavalieri et al. *J Am Chem Soc* 70 3875 1948.]

Tetra-*n*-amylammonium bromide [866-97-7] M 378.5, m 100-101°. Crystd from pet ether, *benzene or acetone/ether mixtures and dried in vacuum at 40-50° for 2 days.

Tetra-*n*-amylammonium iodide [2498-20-6] M 425.5, m 135-137°. Crystd from EtOH and dried at 35° under vac. Also purified by dissolving in acetone and pptd by adding diethyl ether; and dried at 50° for 2 days.

1,4,8,11-Tetraazacyclotetradecane [295-37-4] M 200.33, m 173° (closed capillary and sublimes at 125°), 183-185°, 185°, pK_{Est(1)}~3.8, pK_{Est(2)}~6.0, pK_{Est(3)}~9.0, pK_{Est(4)}~9.6. Purified by recrystn from dioxane (white needles) and sublimes above 120°. It has been distilled, b 132-140°/4-8mm. It forms complexes with metals and gives a sparingly soluble nitrate salt, m 205° (dec), which crystallises from H₂O and is dried at 150°. [UV: Bosnich et al. *Inorg Chem* 4 1102 1963, van Alphen *Recl Trav Chim Pays-Bas* 56 343 1937.]

Tetrabenazine (2-oxo-3-isobutyl-9,10-dimethoxy-1,2,3,4,6,7,-hexahydro-11bH-benzo[a]-quinolizine) [58-46-8] M 317.4, m 127-128°, pK_{Est} ~ 8. Crystd from MeOH. The *hydrochloride* has m 208-210° and the *oxime* has m 158° (from EtOH).

1,1,2,2,-Tetrabromoethane [79-27-6] M 345.7, f 0.0°, b 243.5°, d 2.965, n 1.63533. Washed successively with conc H₂SO₄ (three times) and H₂O (three times), dried with K₂CO₃ and CaSO₄ and distd.

3',3'',5',5''-Tetrabromophenolphthalein ethyl ester [1176-74-5] M 662.0, m 212-214°. Crystd from *benzene, dried at 120° and kept under vacuum.

Tetra-*n*-butylammonium bromide [1643-19-2] M 322.4, m 119.6°. Crystd from *benzene (5mL/g) at 80° by adding hot *n*-hexane (three volumes) and allowing to cool. Dried over P₂O₅ or Mg(ClO₄)₂, under vacuum. The salt is *very hygroscopic*. It can also be crystd from ethyl acetate or dry acetone by adding diethyl ether and dried *in vacuo* at 60° for 2 days. It has been crystd from acetone by addition of diethyl ether. So *hygroscopic* that all manipulations should be carried out in a dry-box. Purified by precipitation of a saturated solution in dry CCl₄ by addition of cyclohexane or by recrystallisation from ethyl acetate, then heating in vacuum to 75° in the presence of P₂O₅. [Symons et al. *J Chem Soc, Faraday Trans 1* 76 2251 1908.] Also recrystallised from CH₂Cl₂/diethyl ether and dried in a vacuum desiccator over P₂O₅. [Blau and Espenson *J Am Chem Soc* 108 1962 1986.]

Tetra-*n*-butylammonium chloride [1112-67-0] M 277.9, m 15.7°. Crystd from acetone by addition of diethyl ether. Very *hygroscopic* and forms crystals with 34H₂O.

Tetra-*n*-butylammonium fluoroborate [429-42-5] M 329.3, m 161-163°. See tetrabutylammonium fluoroborate on p. 480 in Chapter 5.

Tetra-*n*-butylammonium hexafluorophosphate [3109-63-5] M 387.5, m 239-241°. Recrystd from satd EtOH/water and dried for 10h in vac at 70°. It was also recrystd three times from abs EtOH and dried for 2 days in a drying pistol under vac at boiling toluene temperature [Bedard and Dahl *J Am Chem Soc* 108 5933 1986].

Tetra-*n*-butylammonium hydrogen sulfate [32503-27-8] M 339.5, m 171-172°. Crystd from acetone.

Tetra-*n*-butylammonium iodide [311-28-4] M 369.4, m 146°. Crystd from toluene/pet ether (see entry for the corresponding bromide), acetone, ethyl acetate, EtOH/diethyl ether, nitromethane, aq EtOH or water. Dried at room temperature under vac. It has also been dissolved in MeOH/acetone (1:3, 10mL/g), filtered and allowed to stand at room temperature to evaporate to *ca* half its original volume. Distilled water (1mL/g) was then added, and the ppte was filtered off and dried. It was also dissolved in acetone, ppted by adding ether and dried in vac at 90° for 2 days. It has also been recrystallised from CH₂Cl₂/pet ether or hexane, or anhydrous methanol and stored in a vacuum desiccator over H₂SO₄. [Chau and Espenson *J Am Chem Soc* 108 1962 1986.]

Tetra-*n*-butylammonium nitrate [1941-27-1] M 304.5, m 119°. Crystd from *benzene (7mL/g) or EtOH, dried in a vacuum over P₂O₅ at 60° for 2 days.

Tetra-*n*-butylammonium perchlorate [1923-70-2] M 341.9°, m 210°(dec). Crystd from EtOH, ethyl acetate, from *n*-hexane or diethyl ether/acetone mixture, ethyl acetate or hot CH₂Cl₂. Dried in vacuum at room temperature over P₂O₅ for 24h. [Anson et al. *J Am Chem Soc* 106 4460 1984; Ohst and Kochi *J Am Chem Soc* 108 2877 1986; Collman et al. *J Am Chem Soc* 108 2916 1986; Blau and Espenson *J Am Chem Soc* 108 1962 1986; Gustowski et al. *J Am Chem Soc* 108 1986; Ikezawa and Kutal *J Org Chem* 52 3299 1987.]

Tetra-*n*-butylammonium picrate [914-45-4] M 490.6, m 89°. Crystd from EtOH. Dried in a vacuum desiccator over P₂O₅.

Tetra-*n*-butylammonium tetrabutylborate (Bu₄N⁺ Bu₄B⁻) [23231-91-6] M 481.7, m 109.5°. Dissolved in MeOH or acetone, and crystd by adding distd water. Dried in vacuum at 70°. It has also been successively recrystd from isopropyl ether, isopropyl ether/acetone (50:1) and isopropyl ether/EtOH (50:1) for 10h, then isopropyl ether/acetone for 1h, and dried at 65° under reduced pressure for 1 week. [Kondo et al. *J Chem Soc, Faraday Trans 1* 76 812 1980.]

- 2,3,4,5-Tetrachloroaniline** [634-83-3] M 230.9, m 119-120°, pK_{Est} ~-0.26. Crystd from EtOH.
- 2,3,5,6-Tetrachloroaniline** [3481-20-7] M 230.9, m 107-108°, pK_{Est} ~-1.8. Crystd from EtOH.
- 1,2,3,4-Tetrachlorobenzene** [634-66-2] M 215.9, m 45-46°, b 254°/760mm. Crystd from EtOH.
- 1,2,3,5-Tetrachlorobenzene** [634-90-2] M 215.9, m 51°, b 246°/760mm. Crystd from EtOH.
- 1,2,4,5-Tetrachlorobenzene** [95-94-3] M 215.9, m 139.5-140.5°, b 240°/760mm. Crystd from EtOH, ether, *benzene, *benzene/EtOH or carbon disulfide.
- 3,4,5,6-Tetrachloro-1,2-benzoquinone** [2435-53-2] M 245.9, m 130°. Crystd from AcOH. Dry in vacuum desiccator over KOH.
- 1,1,2,2-Tetrachloro-1,2-difluoroethane** [72-12-0] M 203.8, f 26.0°, b 92.8°/760 mm. Purified as for trichlorotrifluoroethane.
- sym-Tetrachloroethane** [79-34-5] M 167.9, b 62°/100mm, 146.2°/atm, d 1.588, n¹⁵ 1.49678. Stirred, on a steam-bath, with conc H₂SO₄ until a fresh portion of acid remained colourless. The organic phase was then separated, distd in steam, dried (CaCl₂ or K₂CO₃), and fractionally distd in a vac.
- Tetrachloroethylene** [127-18-4] M 165.8, b 62°/80mm, 121.2°, d¹⁵ 1.63109, d 1.623, n¹⁵ 1.50759, n 1.50566 It decomposes under similar conditions to CHCl₃, to give phosgene and trichloroacetic acid. Inhibitors of this reaction include EtOH, diethyl ether and thymol (effective at 2-5ppm). Tetrachloroethylene should be distd under a vac (to avoid phosgene formation), and stored in the dark out of contact with air. It can be purified by washing with 2M HCl until the aq phase no longer becomes coloured, then with water, drying with Na₂CO₃, Na₂SO₄, CaCl₂ or P₂O₅, and fractionally distilling just before use. 1,1,2-Trichloroethane and 1,1,1,2-tetrachloroethane can be removed by counter-current extraction with EtOH/water.
- Tetrachloro-N-methylphthalimide** [14737-80-5] M 298.9, m 209.7°. Crystd from absolute EtOH.
- 2,3,4,6-Tetrachloronitrobenzene** [879-39-0] M 260.9, m 42°. Crystd from aqueous EtOH.
- 2,3,5,6-Tetrachloronitrobenzene** [117-18-0] M 260.9, m 99-100°. Crystd from aqueous EtOH.
- 2,3,4,5-Tetrachlorophenol** [4901-51-3] M 231.9, m 116-117°, pK_{Est} ~6.2. Crystd from pet ether.
- 2,3,4,6-Tetrachlorophenol** [58-90-2] M 231.9, m 70°, b 150°/15mm, pK_{Est} ~5.4. Crystd from pet ether.
- 2,3,5,6-Tetrachlorophenol** [935-95-5] M 231.9, m 115°, pK_{Est} ~5.0. Crystd from pet ethers.
- Tetrachlorophthalic anhydride** [117-08-8] M 285.9, m 255-257°. Crystd from chloroform or *benzene, then sublimed.
- 2,3,4,6-Tetrachloropyridine** [14121-36-9] M 216.9, m 74-75°, b 130-135°/16-20mm, pK_{Est} ~-5.7. Crystd from 50% EtOH.
- Tetracosane** [646-31-1] M 338.7, m 54°, b 243-244°/15mm. Crystd from diethyl ether.
- Tetracosanoic (lignoceric) acid** [557-59-5] M 368.7, m 84°, 87.5-88°, pK_{Est} ~5.0. Crystd from acetic acid, Me₂CO, toluene, pet ether/Me₂CO, *C₆H₆/Me₂CO.

1,2,4,5-Tetracyanobenzene [712-74-3] M 178.1, m 270-272° (280°). Crystd from EtOH and sublimed *in vacuo*. [Lawton and McRitchie *J Org Chem* 24 26 1959; Bailey et al. *Tetrahedron* 19 161 1963.]

Tetracyanoethylene [670-54-2] M 128.1, m 199-200° (sealed tube). Crystd from chlorobenzene, dichloroethane, or dichloromethane [Hall et al. *J Org Chem* 52 5528 1987]. Stored at 0° in a desiccator over NaOH pellets. (It slowly evolves HCN on exposure to moist air.) It can also be sublimed at 120° under vacuum. Also purified by repeated sublimation at 120-130°/0.5mm. [Frey et al. *J Am Chem Soc* 107 748 1985; Traylor and Mikszal *J Am Chem Soc* 109 2778 1987.]

7,7,8,8-Tetracyanoquinodimethane [1518-16-7] M 204.2, m 287-290°(dec). Recrystd from distd, dried, acetonitrile.

Tetradecane [629-59-4] M 198.4, m 6°, b 122°/10mm, 252-254°, d 0.763, n 1.429. Washed successively with 4M H₂SO₄ and water. Dried over MgSO₄ and distd several times under reduced pressure [Poë et al. *J Am Chem Soc* 108 5459 1986].

1-Tetradecanol [112-72-1] M 214.4, m 39-39.5°, b 160°/10mm, 170-173°/20mm. Crystd from aq EtOH. Purified by zone melting.

Tetradecyl ether (di-tetradecyl ether) [5412-98-6] M 410.7. Distd under vac and then crystd repeatedly from MeOH/*benzene.

Tetradecyltrimethylammonium bromide [1119-97-7] M 336.4, m 244-249°. Crystd from acetone or a mixture of acetone and >5% MeOH. Washed with diethyl ether and dried in a vacuum oven at 60°. [Dearden and Wooley *J Phys Chem* 91 2404 1987.]

Tetraethoxymethane [78-09-1] M 192.3, b 159°. See tetraethyl orthocarbonate on p. 360.

Tetraethylammonium bromide [71-91-0] M 210.2, m 269°(dec), 284°(dec). Recrystd from EtOH, CHCl₃ or diethyl ether, or, recrystd from acetonitrile, and dried over P₂O₅ under reduced pressure for several days. Also recrystd from EtOH/diethyl ether (1:2), EtOAc, water or boiling MeOH/acetone (1:3) or by adding equal volume of acetone and allowing to cool. Dried at 100° *in vacuo* for 12 days, and stored over P₂O₅.

Tetraethylammonium chloride hydrate [56-34-8] M 165.7, m dec>200°. Crystd from EtOH by adding diethyl ether, from warm water by adding EtOH and diethyl ether, from dimethylacetamide or from CH₂Cl₂ by addition of diethyl ether. Dried over P₂O₅ in vacuum for several days. Also crystd from acetone/CH₂Cl₂/hexane (2:2:1) [Blau and Espenson *J Am Chem Soc* 108 1962 1986; White and Murray *J Am Chem Soc* 109 2576 1987].

Tetraethylammonium iodide [68-05-3] M 257.2, m 302°, >300°(dec). Crystd from acetone/MeOH, EtOH/water, dimethylacetamide or ethyl acetate/EtOH (19:1). Dried under vacuum at 50° and stored over P₂O₅.

Tetraethylammonium perchlorate [2567-83-1] M 229.7, m 345°(dec). Crystd repeatedly from water, aqueous MeOH, acetonitrile or acetone, and dried at 70° under vacuum for 24h. [Cox et al. *J Am Chem Soc* 106 5965 1984; Liu et al. *J Am Chem Soc* 108 1740 1986; White and Murray *J Am Chem Soc* 109 2576 1987.] Also twice crystd from ethyl acetate/95% EtOH (2:1) [Lexa et al. *J Am Chem Soc* 109 6464 1987].

Tetraethylammonium picrate [741-03-7] M 342.1, m >300°(dec). Purified by successive crystns from water or 95% EtOH followed by drying in vacuum at 70°.

Tetraethylammonium tetrafluoroborate [429-06-1] M 217.1, m 235°, 275-277°, 289-291°. Recrystd three times from a mixture of ethyl acetate/hexane (5:1) or MeOH/pet ether, then stored at 95° for 48h

under vacuum [Henry and Faulkner *J Am Chem Soc* **107** 3436 1985; Huang et al. *Anal Chem* **58** 2889 1986]. See entry on p. 481 in Chapter 5.

Tetraethylammonium tetraphenylborate [12099-10-4] M 449.4. Recrystd from aqueous acetone. Dried in a vacuum oven at 60° for several days. *Similarly for the propyl and butyl homologues.*

Tetraethyl 1,1,2,2-ethanetetracarboxylate [632-56-4] M 318.3, m 73-74°. Twice recrystd from EtOH by cooling to 0°.

Tetraethylene glycol dimethyl ether [143-24-8] M 222.3, b 105°/1mm, d 1.010, n 1.435. Stood with CaH₂, LiAlH₄ or sodium, and distd when required.

Tetraethylenepentamine [112-57-2] M 189.3, b 169-171°/0.05mm, d 0.999, n 1.506, pK₁²⁵ 2.98, pK₂²⁵ 4.72, pK₃²⁵ 8.08, pK₄²⁵ 9.10, pK₅²⁵ 9.68. Distd under vacuum. Purified *via* its pentachloride, nitrate or sulfate. Jonassen, Frey and Schaafsma [*J Phys Chem* **61** 504 1957] cooled a soln of 150g of the base in 300mL of 95% EtOH, and added dropwise 180mL of conc HCl, keeping the temperature below 20°. The white ppt was filtered, crystd three times from EtOH/water, then washed with diethyl ether and dried by suction. Reilley and Holloway [*J Am Chem Soc* **80** 2917 1958], starting with a similar soln cooled to 0°, added slowly (keeping the temperature below 10°) a soln of 4.5g-moles of HNO₃ in 600mL of aqueous 50% EtOH (also cooled to 0°). The ppt was filtered by suction, recrystd five times from aqueous 5% HNO₃, then washed with acetone and absolute EtOH and dried at 50°. [For purification *via* the sulfate see Reilley and Vavoulis (*Anal Chem* **31** 243 1959), and for an additional purification step using the Schiff base with benzaldehyde see Jonassen et al. *J Am Chem Soc* **79** 4279 1957].

Tetraethyl orthocarbonate (ethyl orthocarbonate, tetraethoxy ethane) [78-09-1] M 192.3, b 59.6-60°/14mm, 158°/atm, 159°/atm, 160-161°/atm, d₄²⁰ 0.9186, n_D²⁰ 1.3932. Likely impurities are hydrolysis products. Shake with brine (satd NaCl; dilute with a little Et₂O if amount of material is small) and dry (MgSO₄). The organic layer is filtered off and evaporated, and the residue is distd through a helices packed fractionating column with a total reflux partial take-off head. All distns can be done at atmospheric pressure in an inert atmosphere (e.g. N₂). [Roberts and McMahon *Org Synth Coll Vol IV* 457 1963; Connolly and Dyson *J Chem Soc* 828 1937; Tieckelmann and Post *J Org Chem* **13** 266 1948.]

1,1,2,2-Tetrafluorocyclobutane [374-12-9] M 128.1, b 50-50.7°, d 1.275, n 1.3046. Purified distn or by preparative gas chromatography using a 2m x 6mm(i.d.) column packed with β,β'-oxydipropionitrile on Chromosorb P at 33°. [Conlin and Fey *J Chem Soc, Faraday Trans 1* **76** 322 1980; Coffmann et al. *J Am Chem Soc* **71** 490 1949.]

Tetrafluoro-1,3-dithietane [1717-50-6] M 164.1, m -6°, b 47-48°/760mm, d²⁵ 1.6036, n²⁵ 1.3908. Purified by preparative gas chromatography or by distn through an 18in spinning band column. Also purified by shaking vigorously *ca* 40mL with 25mL of 10% NaOH, 5mL of 30% H₂O₂ until the yellow colour disappeared. The larger layer was separated, dried over silica gel to give a colourless liquid boiling at 48°. It had a singlet at -1.77ppm in the NMR spectrum. [Middleton, Howard and Sharkey, *J Org Chem* **30** 1375 1965.]

2,2,3,3-Tetrafluoropropanol [76-37-9] M 132.1, b 106-106.5°, pK²⁵ 12.74. Tetrafluoropropanol (450mL) was added to a soln of 2.25g of NaHSO₃ in 90mL of water, shaken vigorously and stood for 24h. The fraction distilling at or above 99° was refluxed for 4h with 5-6g of KOH and rapidly distd, followed by a final fractional distn. [Kosower and Wu *J Am Chem Soc* **83** 3142 1961.] Alternatively, shaken with alumina for 24h, dried overnight with anhydrous K₂CO₃ and distd, taking the middle fraction (b 107-108°).

Tetra-*n*-heptylammonium bromide [4368-51-8] M 490.7, m 89-91°. Crystd from *n*-hexane, then dried in a vacuum oven at 70°.

Tetra-*n*-heptylammonium iodide [3535-83-9] M 537.7. Crystd from EtOH.

Tetra-*n*-hexylammonium bromide [4328-13-6] M 434.6, m 99-100°. Washed with ether, and dried in a vacuum at room temperature for 3 days.

Tetra-*n*-hexylammonium chloride [5922-92-9] M 390.1. Crystd from EtOH.

Tetra-*n*-hexylammonium iodide [2138-24-1] M 481.6, m 99-101°. Washed with diethyl ether and dried at room temperature *in vacuo* for 3 days.

Tetrahexylammonium perchlorate [4656-81-9] M 454.1, m 104-106°. Crystd from acetone and dried *in vacuo* at 80° for 24h.

Tetrahydrofuran [109-99-9] M 72.1, b 25°/176mm, 66°/760mm, d_4^{20} 0.889, n_D^{20} 1.4070, pK -2.48 (aq H₂SO₄). It is obtained commercially by catalytic hydrogenation of furan from pentosan-containing agricultural residues. It was purified by refluxing with, and distilling from LiAlH₄ which removes water, peroxides, inhibitors and other impurities [Jaeger et al. *J Am Chem Soc* 101 717 1979]. Peroxides can also be removed by passage through a column of activated alumina, or by treatment with aq ferrous sulfate and sodium bisulfate, followed by solid KOH. In both cases, the solvent is then dried and fractionally distd from sodium. Lithium wire or vigorously stirred molten potassium have also been used for this purpose. CaH₂ has also been used as a drying agent.

Several methods are available for obtaining the solvent almost anhydrous. Ware [*J Am Chem Soc* 83 1296 1961] dried vigorously with sodium-potassium alloy until a characteristic blue colour was evident in the solvent at Dry-ice/cellosolve temperatures. The solvent was kept in contact with the alloy until distd for use. Worsfold and Bywater [*J Chem Soc* 5234 1960], after refluxing and distilling from P₂O₅ and KOH, in turn, refluxed the solvent with sodium-potassium alloy and fluorenone until the green colour of the disodium salt of fluorenone was well established. [Alternatively, instead of fluorenone, benzophenone, which forms a blue ketyl, can be used.] The tetrahydrofuran was then fractionally distd, degassed and stored above CaH₂. *p*-Cresol or hydroquinone inhibit peroxide formation. The method described by Coetzee and Chang [*Pure Appl Chem* 57 633 1985] for 1,4-dioxane also applies here. Distns should always be done in the presence of a reducing agent, e.g. FeSO₄. **It irritates the skin, eyes and mucous membranes and the vapour should never be inhaled. It is HIGHLY FLAMMABLE and the necessary precautions should be taken.**

Rapid purification: Purification as for diethyl ether.

***l*-Tetrahydropalmatine (2,3,9,10-tetramethoxy-6*H*-dibenzo[*a,g*]quinolizidine)** [10097-84-4] M 355.4, m 148-149°, $[\alpha]_D^{20}$ -291° (EtOH). Crystd from MeOH by addition of water [see *J Chem Soc (C)* 530 1967].

Tetrahydropyran [142-68-7] M 86.1, b 88.0°, n 1.4202, d 0.885, pK -2.79 (aq H₂SO₄). Dried with CaH₂, then passed through a column of silica gel to remove olefinic impurities and fractionally distd. Freed from peroxides and moisture by refluxing with sodium, then distilling from LiAlH₄. Alternatively, peroxides can be removed by treatment with aqueous ferrous sulfate and sodium bisulfate, followed by solid KOH, and fractional distn from sodium.

Tetrahydro-4*H*-pyran-4-one [29943-42-8] M 100.1, b 57-59°/11mm, 65-66°/15mm, 67-68°/18mm, 73°/20mm, 164.7°/atm, 166-166.5°/atm, d_4^{20} 1.0844, n_D^{20} 1.4551. Purified by repeated distn preferably in a vacuum. [Baker *J Chem Soc* 296 1944; IR: Olsen and Bredoch *Chem Ber* 91 1589 1958.] The *oxime* has m 87-88° and b 110-111°/13mm [Cornubert et al. *Bull Soc Chim Fr* 36 1950]. The 4-nitrophenylhydrazone forms orange-brown needles from EtOH, m 186° [Cawley and Plant *J Chem Soc* 1214 1938].

Tetrahydrothiophene [110-01-0] M 88.2, m -96°, b 14.5°/10mm, 120.9°/760mm, d 0.997, n 1.5289. Crude material was purified by crystn of the mercuric chloride complex to a constant melting point. It was then regenerated, washed, dried, and fractionally distd. [Whitehead et al. *J Am Chem Soc* 73 3632 1951.] It has been dried over Na₂SO₄ and distd in a vacuum [Roberts and Friend *J Am Chem Soc* 108 7204 1986].

Tetrahydro-4H-thiopyran-4-one [1072-72-6] *M* 116.2, *m* 60-62°, 61-62°, 64-65°, 65-67°. Purified by recrystn from diisopropyl ether or pet ether and dried in air. If too impure then dissolve in Et₂O, wash with aq NaHCO₃, then H₂O, dried (MgSO₄), filt'd, evap'd and the residue recryst'd as before. [Cardwell *J Chem Soc* 715 1949.] The *oxime* can be recryst'd from CHCl₃-pet ether (at -20°) and has *m* 84-85° [Barkenbus et al. *J Org Chem* 20 871 1955]. The *2,4-dinitrophenylhydrazone* has *m* 186° (from EtOAc) [Barkenbus et al. *J Org Chem* 16 232 1951]. The *S-dioxide* is recryst'd from AcOH, *m* 173-174° [Fehnel and Carmack *J Am Chem Soc* 70 1813 1948].

Tetrahydroxy-*p*-benzoquinone (2H₂O) [5676-48-2] *M* 208.1, *pK*₁³⁰ 4.80, *pK*₂³⁰ 6.8. Crystd from water.

Tetrakis(dimethylamino)ethylene [996-70-3] *M* 300.2, *b* 60°/1mm, *d* 0.861, *n* 1.4817, *pK*_{Est(1)}<0, *pK*_{Est(2)}<0, *pK*_{Est(3)}~1.5, *pK*_{Est(4)} 5.1. Impurities include tetramethylurea, dimethylamine, tetramethylethanediamine and tetramethyloxamide. It was washed with water while being flushed with nitrogen to remove dimethylamine, dried over molecular sieves, then passed through a silica gel column (previously activated at 400°) under nitrogen. Degassed on a vacuum line by distn from a trap at 50° to one at -70°. Finally, it was stirred over sodium-potassium alloy for several days. [Holroyd et al. *J Phys Chem* 89 4244 1985; Wiberg *Angew Chem Int Ed Engl* 7 766 1968.]

Tetralin (1,2,3,4-tetrahydronaphthalene) [119-64-2] *M* 132.2, *b* 65-66°/5mm, 207.6°/760 mm, *d* 0.968, *n* 1.5413. It was washed with successive portions of conc H₂SO₄ until the acid layer no longer became coloured, then washed with aq 10% Na₂CO₃, and then dist'd water. Dried (CaSO₄ or Na₂SO₄), filtered, refluxed and fractionally dist'd at under reduced pressure from sodium or BaO. It can also be purified by repeated fractional freezing.

Bass [*J Chem Soc* 3498 1964] freed tetralin, purified as above, from naphthalene and other impurities by conversion to ammonium tetralin-6-sulfonate. Conc H₂SO₄ (150mL) was added slowly to stirred tetralin (272mL) which was then heated on a water bath for about 2h to give complete soln. The warm mixture, when poured into aq NH₄Cl soln (120g in 400mL water), gave a white ppt which, after filtering off, was crystd from boiling water, washed with 50% aq EtOH and dried at 100°. Evap'n of its boiling aq soln on a steam bath removed traces of naphthalene. The pure salt (229g) was mixed with conc H₂SO₄ (266mL) and steam dist'd from an oil bath at 165-170°. An ether extract of the distillate was washed with aq Na₂SO₄, and the ether was evap'd, prior to distilling the tetralin from sodium. Tetralin has also been purified *via* barium tetralin-6-sulfonate, conversion to the sodium salt and decomposition in 60% H₂SO₄ using superheated steam.

Tetralin hydroperoxide [771-29-9] *M* 164.2, *m* 56°. Crystd from hexane.

α-Tetralone (1,2,3,4-tetrahydro-1-oxonaphthalene) [529-34-0] *M* 146.2, *m* 2-7°, 7.8-8.0°, *b* 75-85°/0.3mm, 89°/0.5mm, 94-95°/2mm, 132-134°/15mm, 143-145°/20mm, *d*₄²⁰ 1.0695, *n*_D²⁰ 1.5665. Check the IR first. Purify by dissolving 20mL in Et₂O (200mL), washing with H₂O (100mL), 5% aq NaOH (100mL), H₂O (100mL), 3% aq AcOH (100mL), 5% NaHCO₃ (100mL) then H₂O (100mL) and dry the ethereal layer over MgSO₄. Filter, evap and fractionate the residue through a 6in Vigreux column under reduced pres to give a colourless oil (~17g) with *b* 90-91°/0.5-0.7mm. [Snyder and Werber *Org Synth Coll Vol III* 798 1955.] It has also been fractionated through a 0.5metre packed column with a heated jacket under reflux using a partial take-off head. [Olson and Bader *Org Synth Coll Vol IV* 898 1963.]

β-Tetralone (1,2,3,4-tetrahydro-2-oxonaphthalene) [530-93-8] *M* 146.2, *m* 17-18°, ~18°, *b* 93-95°/2mm, 104-105°/4mm, 114-115°/4-5mm, 140°/18mm, *d*₄²⁰ 1.1000, *n*_D²⁰ 1.5598. If reasonably pure then fractionate through an efficient column. Otherwise purify *via* the *bisulfite adduct*. To a soln of NaHSO₃ (32.5g, 0.31mol) in H₂O (57mL) is added 95% EtOH (18mL) and set aside overnight. Any bisulfite-sulfate that separated is removed by filtration and the filtrate is added to the tetralone (14.6g, 0.1mol) and shaken vigorously. The adduct separates in a few minutes as a white ppt and kept on ice for ~3.5h with occasional shaking. The ppt is collected, washed with 95% EtOH (13mL), then with Et₂O (4 x 15mL, by stirring the suspension in the solvent, filtering and repeating the process). The colourless product is dried in air and stored in air tight containers in which it is stable for extended periods (yield is ~17g). This bisulfite (5g) is suspended in H₂O (25mL) and Na₂CO₃.H₂O (7.5g) is added (pH of soln is ~10). The mixture is then extracted

with Et₂O (5 x 10mL, i.e. until the aqueous phase does not test for tetralone — see below). Wash the combined extracts with 10% aqueous HCl (10mL), H₂O (10mL, i.e. until the washings are neutral), dry (MgSO₄), filter, evaporate and distil the residual oil using Claisen flask under reduced pressure and in a N₂ atm. The pure tetralone is a colourless liquid **b** 70-71°/0.25mm (see also above). The yield is ~2g. **Tetralone test:** Dissolve a few drops of the tetralone soln (ethereal or aqueous) in 95% EtOH in a test tube and add 10 drops of 25% NaOH down the side of the tube. A deep blue colour develops at the interface with air. [Soffer *Org Synth Coll Vol IV* 903 1963; Cornforth et al. *J Chem Soc* 689 1942; UV: Soffer et al. *J Am Chem Soc* 1556 1952.] The *phenylhydrazone* has **m** 108° [Crawley and Robinson *J Chem Soc* 2001 1938].

Tetramethylammonium bromide [64-20-0] **M 154.1, sublimes with dec >230°**. Crystd from EtOH, EtOH/diethyl ether, MeOH/acetone, water or from acetone/MeOH (4:1) by adding an equal volume of acetone. It was dried at 110° under reduced pressure or at 140° for 24h.

Tetramethylammonium chloride [75-57-0] **M 109.6, m >230°(dec)**. Crystd from EtOH, EtOH/CHCl₃, EtOH/diethyl ether, acetone/EtOH (1:1), isopropanol or water. Traces of the free amine can be removed by washing with CHCl₃.

Tetramethylammonium hydroxide (5H₂O) [10424-65-4 (5H₂O); 75-59-2 (aq soln)] **M 181.2, m 63°, 65-68°**. Freed from chloride ions by passage through an ion-exchange column (Amberlite IRA-400, prepared in its OH⁻ form by passing 2M NaOH until the effluent was free from chloride ions, then washed with distilled H₂O until neutral). A modification, to obtain carbonate-free hydroxide, uses the method of Davies and Nancollas [*Nature* 165 237 1950].

Tetramethylammonium iodide [75-58-1] **M 201.1, m >230°(dec)**. Crystd from water or 50% EtOH, EtOH/diethyl ether, ethyl acetate, or from acetone/MeOH (4:1) by adding an equal volume of acetone. Dried in a vacuum desiccator.

Tetramethylammonium perchlorate [2537-36-2] **M 173.6, m >300 °(dec)**. Crystd from acetone and dried *in vacuo* at 60° for several days.

Tetramethylammonium tetraphenylborate [15525-13-0] **M 393.3**. Recrystd from acetone, acetone/CCl₄ and from acetone/1,2-dichloroethane. Dried over P₂O₅ in vacuum, or in a vacuum oven at 60° for several days.

1,2,3,4-Tetramethylbenzene (prehnitine) [488-23-3] **M 134.2, m -6.3°, b 79.4°/10mm, 204-205°/760mm, d 0.905, n 1.5203**. Dried over sodium and distd under reduced pressure.

1,2,3,5-Tetramethylbenzene (isodurene) [527-53-7] **M 134.2, m -23.7°, b 74.4°/10mm, 198°/760mm, d 0.890, n 1.5130**. Refluxed over sodium and distd under reduced pressure.

1,2,4,5-tetramethylbenzene (durene) [95-93-2] **M 134.2, m 79.5-80.5°**. Chromatographed on alumina, and recrystd from aqueous EtOH or *benzene. Zone-refining removes duroaldehydes. Dried under vacuum. [Yamauchi et al. *J Phys Chem* 89 4804 1985.] It has also been sublimed *in vacuo* [Johnston et al. *J Am Chem Soc* 109 1291 1987].

N,N,N',N'-Tetramethylbenzidine [366-29-0] **M 240.4, m 195.4-195.6°, pK_{Est(1)}~3.4, pK_{Est(2)}~4.5**. Crystd from EtOH or pet ether, then from pet ether/*benzene, and sublimed in a vacuum. [Guarr et al. *J Am Chem Soc* 107 5104 1985.] Dried under vac in an Abderhalden pistol, or carefully on a vacuum line.

2,2,4,4-Tetramethylcyclobutan-1,3-dione [933-52-8] **M 140.2, m 114.5-114.9°**. Crystd from *benzene and dried under vacuum over P₂O₅ in an Abderhalden pistol.

3,3,5,5-Tetramethylcyclohexanone [14376-79-5] **M 154.3, m 11-12°, 13.2°, b 59-61°, 80-82°/13mm, 196°/760mm, 203.8-204.8°/760mm, d_D²⁰ 0.8954, n_D²⁰ 1.4515**. Purified first through a

24in column packed with Raschig rings then a 40cm Vigreux colum under reduced pressure (b 69-69.3°/7mm, see above). The *oxime* has m 144-145° (from 60% EtOH) and the *semicarbazone* has m 196-197°, 197-198° (214.5°, 217-218°) [Karasch and Tawney *J Am Chem Soc* 63 2308 1941; UV: Sandris and Ourisson *Bull Soc Chim Fr* 958 1956].

***p,p'*-Tetramethyldiaminodiphenylmethane [bis(*p*-dimethylaminophenyl)methane, Michler's base]** [101-61-1] M 254.4, m 89-90°, b 155-157°/0.1mm, pK_{Est(1)}~5.8, pK_{Est(2)}~5.1. Crystd from EtOH (2mL/g) or 95% EtOH (ca 12mL/g). It sublimes on heating.

Tetramethylene sulfoxide (tetrahydrothiophen 1-oxide) [1600-44-8] M 104.2, b 235-237°, d 1.175, n 1.525. Shaken with BaO for 4 days, then distd from CaH₂ under reduced pressure.

***N,N,N',N'*-Tetramethylethylenediamine (TMEDA, TEMED)** [110-18-9] M 116.2, b 122°, d 1.175, n²⁵ 1.4153, pK₁²⁵ 5.90, pK₂²⁵ 9.14. Partially dried with molecular sieves (Linde type 4A), and distd in vacuum from butyl lithium. This treatment removes all traces of primary and secondary amines and water. [Hay, McCabe and Robb *J Chem Soc, Faraday Trans 1* 68 1 1972.] Or, dried with KOH pellets. Refluxed for 2h with one-sixth its weight of *n*-butyric anhydride (to remove primary and secondary amines) and fractionally distd. Refluxed with fresh KOH, and distd under nitrogen. [Cram and Wilson *J Am Chem Soc* 85 1245 1963.] Also distd from sodium.

Tetramethylethylenediamine dihydrochloride [7677-21-8] M 198.2, m ~300°. Crystd from 98% EtOH/conc HCl. Hygroscopic. [Knorr *Chem Ber* 37 3510 1904.]

1,1,3,3-Tetramethylguanidine [80-70-6] M 115.2, b 159-160°, d 0.917 n 1.470, pK²⁵ 13.6. Refluxed over granulated BaO, then fractionally distd.

***N,N,N',N'*-Tetramethyl-1,8-naphthalenediamine** [20734-58-1] M 214.3, m 45-48°, 47-48°, b 144-145°/4mm, pK₁ -10.5 (aq H₂SO₄, diprotonation), pK₂ 12.34 (monoprotonation). It is prepared by methylating 1,8-diaminonaphthalene and likely impurities are methylated products. The tetramethyl compound is a stronger base than the unmethylated, di and trimethylated derivatives. The pK_a values are: 1,8-(NH₂)₂ = 4.61, 1,8-(NHMe)₂ = 5.61, 1-NHMe-8-NHMe₂ = 6.43 and 1,8-(NMe₂)₂ = 12.34. The mixture is then treated H₂O at pH 8 (where all but the required base are protonated) and extracted with Et₂O or CHCl₃. The dried extract (K₂CO₃) yields the tetramethyldiamine on evapn which can be distd. It is a strong base with weak nucleophilic properties, e.g. it could not be alkylated by refluxing with EtI in MeCN for 4 days and on treatment with methyl fluorosulfonate only the fluorosulfonate salt of the base is obtained. [NMR: Adler et al. *J Chem Soc, Chem Commun* 723 1968; *J Am Chem Soc* 63 358 1941.] See Proton sponge p. 134.

Tetramethyl orthocarbonate (methyl orthocarbonate, tetramethoxy methane) [1850-14-2] M 136.2, m -5.6°, -5°, -2°, b 113.5°/760mm, 113.5-114°/755mm, 112-114°/atm, d₄²⁰ 1.0202, n_D²⁰ 1.3860. Purified in the same way as for tetraethyl orthocarbonate. [Smith *Acta Chem Scand* 10 1006 1956; Tiekemann and Post *J Org Chem* 13 266 1948.]

2,6,10,14-Tetramethylpentadecane (pristane, norphytane) [1921-70-6] M 268.5, b 68° (bath temp)/0.004mm, 158°/10mm, 296°/atm, d₄²⁰ 0.7827, n_D²⁰ 1.4385. Purified by shaking with conc H₂SO₄ (care with this acid, if amount of pristane is too small then it should be diluted with pet ether *not* Et₂O which is quite sol in the acid), the H₂O (care as it may heat up if in contact with conc H₂SO₄), dried (MgSO₄) evaporated and distd over Na metal. [Sörensen and Sörensen *Acta Chem Scand* 3 939 1949.]

***N,N,N',N'*-Tetramethyl-1,4-phenylenediamine** [100-22-1] M 164.3, m 51°, b 260°/760mm, pK₁²⁰ 2.29, pK₂²⁰ 6.35. Crystd from pet ether or water. It can be sublimed or dried carefully in a vacuum line, and stored in the dark under nitrogen. Also recrystd from its melt.

***N,N,N',N'*-Tetramethyl-1,4-phenylenediamine dihydrochloride** [637-01-4] M 237.2, m 222-224°. Crystd from isopropyl or *n*-butyl alcohols, satd with HCl. Treated with aq NaOH to give the free base which was filtered, dried and sublimed in a vacuum. [Guarr et al. *J Am Chem Soc* 107 5104 1985.]

2,2,6,6-Tetramethylpiperidinyl-1-oxy (TEMPO) [2564-83-2] M 156.3, m 36-38°. Purified by sublimation (33°, water aspirator) [Hay and Fincke *J Am Chem Soc* 109 8012 1987].

2,2,6,6-Tetramethyl-4-piperidone hydrochloride (triacetoneamine) [33973-59-0] M 191.7, m 190° (dec), 198-199° (dec), pK^{25} 7.90. Purified by recrystn from EtOH/Et₂O, MeCN or Me₂CO/MeOH. The *free base* has m 37-39° (after sublimation), b 102-105°/18mm, and *hydrate* m 56-58° (wet Et₂O); the *hydrobromide* has m 203° (from EtOH-Et₂O) and the *picrate* has m 196° (from aq EtOH). [Sandris and Ourisson *Bull Soc Chim Fr* 345 1958.]

Tetramethylthiuram disulfide [bis-(dimethylthiocarbamyl)-disulfide] [137-26-8] M 240.4, m 146-148°, 155-156°. Crystd (three times) from boiling CHCl₃, then recrystd from boiling CHCl₃ by adding EtOH dropwise to initiate pptn, and allowed to cool. Finally it was pptd from cold CHCl₃ by adding EtOH (which retained the monosulfide in soln). [Ferington and Tobolsky *J Am Chem Soc* 77 4510 1955.]

1,1,3,3-Tetramethyl urea [632-22-4] M 116.2, f -1.2°, b 175.2°/760mm, d 0.969, n 1.453. Dried over BaO and distd under nitrogen.

Tetramethyl uric acid [2309-49-1] M 224.2, m 225°, 228°, $pK_{Est} < 0$. Crystd from H₂O or MeOH.

1,3,5,5-Tetranitrohexahydropyrimidine [81360-42-1] M 270.1, m 153-154°. Crystd from EtOH (5x), and sublimed (~65°/0.05mm) [*J Org Chem* 47 2474 1982; *J Labelled Comp Radiopharm* 29 1197 1991].

Tetranitromethane [509-14-8] M 196.0, m 14.2°, b 21-23°/23mm, 126°/760mm, d 1.640, n 1.438. Shaken with dilute NaOH, washed, steam distd, dried with Na₂SO₄ and fractionally crystd by partial freezing. The melted crystals were dried with MgSO₄ and fractionally distd under reduced pressure. Shaken with a large volume of dilute NaOH until no absorption attributable to the *aci*-nitro anion (from mono- di- and trinitromethanes) is observable in the water. Then washed with distilled water, and distilled at room temperature by passing a stream of air or nitrogen through the liquid and condensing in a trap at -80°. It can be dried with MgSO₄ or Na₂SO₄, fractionally crystd from the melt, and fractionally distd under reduced pressure.

Tetra(*p*-nitrophenyl)ethylene [47797-98-8] M 512.4, m 306-307°. Crystd from dioxane or AcOH and dried at 150°/0.1mm. [Gorvin *J Chem Soc* 678 1959.]

4,7,13,18-Tetraoxa-1,10-diazabicyclo[8.5.5]eicosane (Cryptand 211) [31250-06-3] M 288.1, $pK_{Est} \sim 7.9$. Redistd, dried under high vacuum over 24h, and stored under nitrogen.

1,7,10,16-Tetraoxa-4,13-diazacyclooctadecane (4,13-diaza-18-crown-6) [23978-55-4] M 262.3, m 118-116°, $pK_{Est} \sim 8.8$. Twice recrystd from *benzene/*n*-heptane, and dried for 24h under high vacuum [E.Weber and F.Vögtle *Top Curr Chem* (Springer Verlag, Berlin) 98 1 1981; D'Aprano and Sesta *J Phys Chem* 91 2415 1987].

Tetrapentylammonium bromide [866-97-7] M 378.5, m 100-101°. See tetra-*n*-amylammonium bromide on p. 356.

Tetraphenylethylene [632-51-9] M 332.4, m 223-224°, b 415-425°/760mm. Crystd from dioxane or from EtOH/*benzene. Sublimed under high vacuum.

Tetraphenylhydrazine [632-52-0] M 336.4, m 147°, $pK_{Est} \sim 0$. Crystd from 1:1 CHCl₃/toluene or CHCl₃/EtOH. Stored in a refrigerator, in the dark.

***trans*-1,1,4,4-Tetraphenyl-2-methylbutadiene** [20411-57-8] M 372.5. Crystd from EtOH.

1,2,3,4-Tetraphenylnaphthalene [751-38-2] M 432.6, m 199-201°, 204-204.5°. Crystd from MeOH or as EtOH. [Fieser and Haddadin *Org Synth* 46 107 1966.]

5,6,11,12-Tetraphenylnaphthacene (Rubrene) [517-51-1] M 532.7, $m > 315^\circ$, 322° , d 1.255. Orange crystals by sublimation at $250-260^\circ/3-4\text{mm}$ [UV Badger and Pearce *Spectrochim Acta* 4 280 1950]. Also recrystd from *benzene under red light because it is chemiluminescent and light sensitive.

5,10,15,20-Tetraphenylporphyrin (TPP) [917-23-7] M 614.7, λ_{max} 482nm. Purified by chromatography on neutral (Grade I) alumina, and recrystd from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ [Yamashita et al. *J Phys Chem* 91 3055 1987].

Tetra-*n*-propylammonium bromide [1941-30-6] M 266.3, $m > 280^\circ(\text{dec})$. Crystd from ethyl acetate/EtOH (9:1), acetone or MeOH. Dried at 110° under reduced pressure.

Tetra-*n*-propylammonium iodide [631-40-3] M 313.3, $m > 280^\circ(\text{dec})$. Purified by crystallization from EtOH, EtOH/diethyl ether (1:1), EtOH/water or aqueous acetone. Dried at 50° under vacuum. Stored over P_2O_5 in a vacuum desiccator.

Tetra-*n*-propylammonium perchlorate [15780-02-6] M 285.8, m 239-241°. See tetrapropylammonium perchlorate on p. 483 in Chapter 5.

5,10,15,20-Tetra-4'-pyridinylporphyrin [16834-13-2] M 618.7, $m > 300^\circ(\text{dec})$. Purified by chromatography on alumina (neutral, Grade I), followed by crystallization from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ [Yamashita et al. *J Phys Chem* 91 3055 1987].

Tetrathiafulvalene [31366-25-3] M 204.4, m 122-124°. Recrystd from cyclohexane/hexane under an argon atmosphere [Kauzlarich et al. *J Am Chem Soc* 109 4561 1987].

1,2,3,4-Tetrazole [288-94-8] M 70.1, m 156°, pK 4.89 (acidic). Crystd from EtOH, sublimed under high vacuum at ca 120° (care should be taken due to possible EXPLOSION).

Thebaine [115-37-7] M 311.4, m 193°, $[\alpha]_{\text{D}}^{25}$ -219° (EtOH), pK¹⁵ 8.15. Sublimed at $170-180^\circ$.

2-Thenoyltrifluoroacetone [326-91-0] M 222.2, m 42-44°, b 96-98°/9mm. Crystd from hexane or *benzene. (Aqueous solutions slowly decompose).

2-Thenylamine [27757-85-3] M 113.1, b 78.5°/15mm, pK³⁰ 8.92. Distd under reduced pressure (nitrogen), from BaO, through a column packed with glass helices.

Theobromine [83-67-0] M 180.2, m 337°, pK₁⁴⁰ -0.16, pK₂²⁵ 9.96. Crystd from water.

Theophylline [58-55-9] M 180.2, m 270-274°, pK₁⁴⁰ -0.24, pK₂ 2.5, pK₃⁴⁰ 8.79. Crystd from H₂O.

Thevetin [11018-93-2] M 858.9, m softens at 194° , m 210°. Crystd (as trihydrate) from isopropanol. Dried at $100^\circ/10\text{mm}$ to give the hemihydrate (very hygroscopic).

Thianthrene [92-85-3] M 216.3, m 158°. Crystd from Me₂CO (charcoal), AcOH or EtOH. Sublimes in a vacuum.

ϵ -[2-(4-Thiazolidone)]hexanoic acid [539-35-5] M 215.3, m 140°, pK_{Est} ~4.7. Crystd from H₂O, Me₂CO or MeOH.

Thiazoline-2-thiol [96-53-7] M 119.2, m 106-107°, 106-108°, pK_{Est} ~13.0. Purified by dissolution in alkali, pptn by addition of HCl and then recrystd from H₂O as needles. [IR: Flett *J Chem Soc* 347 1953 and Mecke et al. *Chem Ber* 90 975; Gabriel and Stelzner *Chem Ber* 28 2931 1895.]

4-(2-Thiazolylazo)resorcinol [2246-46-0] M 221.2, m 200-202°(dec), λ_{\max} 500 nm, $\text{pK}_1^{2.5}$ 1.25, $\text{pK}_2^{2.5}$ 6.53, $\text{pK}_3^{2.5}$ 10.76. Dissolved in alkali, extracted with diethyl ether, and re-ppted with dil HCl. The purity was checked by TLC on silica gel using pet ether/diethyl ether/EtOH (10:10:1) as the mobile phase.

Thietane (trimethylene sulfide) [287-27-4] M 74.1, m -64°, -73.2°, b 93.8-94.2°/752mm, 95°/atm, d_4^{20} 1.0200, n_D^{20} 1.5020. Purified by preparative gas chromatography on a dinonyl phthalate column. It has also been purified by drying over anhydrous K_2CO_3 , and distd through a 25 cm glass helices packed column (for 14g of thietane), then dried over CaSO_4 before sealing in a vac. [Haines et al. *J Phys Chem* 58 270 1954.] It is characterised as the *dimethylsulfonium iodide* m 97-98° [Bennett and Hock *J Chem Soc* 2496 1927]. The *S-oxide* has b 102°/25mm, n_D^{21} 1.5075 [Tamres and Searles *J Am Chem Soc* 81 2100 1959].

Thioacetamide [62-55-5] M 75.1, m 112-113°, $\text{pK}^{2.5}$ 13.4. Crystd from absolute diethyl ether or *benzene. Dried at 70° in vacuum and stored over P_2O_5 at 0° under nitrogen. (*Develops an obnoxious odour on storage*, and absorption at 269nm decreases, hence it should be freshly crystd before use).

Thioacetanilide [677-53-6] M 151.2, m 75-76°, pK_{Est} ~13.1. Crystd from H_2O and dried *in vacuo*.

Thiobarbituric acid [504-17-6] M 144.2, m 235°(dec), $\text{pK}_1^{2.5}$ 2.25, $\text{pK}_2^{2.5}$ 10.72 (2% aq ETOH). Crystd from water.

Thiobenzanilide [636-04-4] M 213.2, m 101.5-102°, pK_{Est} ~12.6. Crystd from MeOH at Dry-ice temperature.

(1R)-(-)-Thiocamphor (1R-bornane-2-thione, 1R-(-)-1,7,7-trimethylbicyclo[2.2.1]heptane-2-thione) [53402-10-1] M 168.3, m 136-138°, 146°, $[\alpha]_D^{22}$ -22° (c 3, EtOAc). Forms red prisms from EtOH and sublimes under vacuum. It possesses a sulfurous odour and is volatile as camphor. [Sen *J Indian Chem Soc* 12 647 1935; 18 76 1941.] The *racemate* crystallises from * C_6H_6 and has m 145° [138.6-139°, White and Bishop *J Am Chem Soc* 62 10 1940].

1,1'-Thiocarbonyldiimidazole [6160-65-2] M 178.1, m 100-102°, 105-106°. It forms yellow crystals by recrystn from tetrahydrofuran or by sublimation at 10^{-3} Torr (bath temp 70-80°). Hydrolysed by H_2O , store dry. [Staab and Walther *Justus Liebigs Ann Chem* 657 98 1962; Pullukat et al. *Tetrahedron Lett* 1953 1967.]

Thiochrome {2,7-dimethyl-5H-thiachromine-8-ethanol; 3,8-dimethyl-2-hydroxyethyl-5H-thiazolo[2,3:1',2']pyrimido[4',5'-d]pyrimidine} [92-35-3] M 262.3, m 227-228°, pK_{Est} ~5.8 (thiazol-N protonation). Crystd from chloroform.

Thiodiglycollic acid [123-93-3] M 150.2, m 129°, $\text{pK}_1^{2.5}$ 3.15 (3.24), $\text{pK}_2^{2.5}$ 4.13 (4.56). Crystd from water.

3,3'-Thiodipropionic acid [111-17-1] M 178.2, $\text{pK}_1^{2.5}$ 3.84, $\text{pK}_2^{2.5}$ 4.66. Crystd from water.

Thioflavine T [2390-54-7] M 318.9, $\text{pK}^{2.5}$ 2.7. Crystd from *benzene/EtOH (1:1).

Thioformamide [115-08-2] M 61.0, m 29°, pK_{Est} ~12.4. Crystd from ethyl acetate or ether/pet ether.

Thioglycollic acid [68-11-1] M 92.1, b 95-96°/8mm, d 1.326, n 1.505, $\text{pK}_1^{2.5}$ 3.42, $\text{pK}_2^{2.5}$ 10.20. Mixed with an equal volume of *benzene, the *benzene is then distd to dehydrate the acid. After heating to 100° to remove most of the *benzene, the residue was distd under vacuum and stored in sealed ampoules at 3°. [Eshelman et al. *Anal Chem* 22 844 1960.]

Thioguanosine (2-amino-6-mercapto-9- β -D-ribofuranosylpurine) [85-31-4] M 299.3, m 230-231°(dec), $[\alpha]_D^{20}$ -64° (c 1.3, 0.1N NaOH), pK 8.33. Crystd (as hemihydrate) from water.

[Previous Page](#)

Thioindigo [522-75-8] M 296.2, m >280°. Adsorbed on silica gel from CCl₄/*benzene (3:1), eluted with *benzene, crystd from CHCl₃ and dried at 60-65°. [Wyman and Brode *J Am Chem Soc* 73 1487 1951.] This paper also gives details of purification of other thioindigo dyes.

Thiomalic (mercaptosuccinic) acid [70-49-5] M 150.2, m 153-154°, pK₁²⁵ 3.64 (3.17), pK₂²⁵ 4.64 (4.67), pK₃²⁵ 10.37 (10.52). Extracted from aqueous soln several times with diethyl ether, and the aqueous soln freeze-dried.

Thio-Michler's Ketone [1226-46-6] M 284.6, λ_{max} 457 nm (ε 2.92 x 10⁴ in 30% aq *n*-propanol). Purified by recrystn from hot EtOH or by triturating with a small volume of CHCl₃, followed by filtration and washing with hot EtOH [Terbell and Wystrade *J Phys Chem* 68 2110 1964].

Thionanthone (thioxanthone) [492-22-8] M 212.3, m 212-213°, b 371-373°/712mm. See 9*H*-thioxanthene-9-one on p. 369.

2-Thionaphthol [91-60-1] M 160.2, m 81°, 82°, b 153.5°/15mm, 286°/760mm, pK_{Est} ~6.1. Crystd from EtOH.

Thionine (3,7-diaminophenothiazine) [135-59-1; 581-64-6 (HCl)] M 263.7, ε₅₉₀ 6.2 x 10⁴ M⁻¹ cm⁻¹, pK¹⁵ 6.9. The standard biological stain is highly pure. It can be crystd from water or 50% EtOH, then chromatographed on alumina using CHCl₃ as eluent [Shepp, Chaberek and McNeil *J Phys Chem* 66 2563 1962]. Dried overnight at 100° and stored in a vacuum. The *hydrochloride* can be crystd from 50% EtOH or dilute HCl and aqueous *n*-butanol. Purified also by column chromatography and washed with CHCl₃ and acetone. Dried *in vacuo* at room temperature.

Thioxine hydrochloride (8-mercaptoquinoline hydrochloride) [34006-16-1] M 197.7, m 170-175° (dec), pK₁²⁵ 2.16, pK₂²⁵ 8.38. Crystallises from EtOH and the crystals are yellow in colour. It has pK_a²⁰ values of 2.05 and 8.29 in H₂O. [UV: Albert and Barlin *J Chem Soc* 2384 1959.]

Thiophane (tetrahydrothiophene) [110-01-0] M 88.2, b 40.3°/39.7mm. See tetrahydrothiophene on p. 361.

Thiophene [110-02-1] M 84.1, f -38.5°, b 84.2°, d 1.525, n 1.52890, n³⁰ 1.5223. The simplest purification procedure is to dry with solid KOH, or reflux with sodium, and fractionally distd through a glass-helices packed column. More extensive treatments include an initial wash with aq HCl, then water, drying with CaSO₄ or KOH, and passage through columns of activated silica gel or alumina. Fawcett and Rasmussen [*J Am Chem Soc* 67 1705 1945] washed thiophene successively with 7M HCl, 4M NaOH, and distd water, dried with CaCl₂ and fractionally distd. *Benzene was removed by fractional crystn by partial freezing, and the thiophene was degassed and sealed in Pyrex flasks. [Also a method is described for recovering the thiophene from the *benzene-enriched portion.]

Thiophene-2-acetic acid [1918-77-0] M 142.2, m 76°, pK²⁵ 3.89. Crystd from ligroin.

Thiophene-3-acetic acid [6964-21-2] M 142.2, m 79-80°, pK_{Est} ~3.1. Crystd from ligroin.

2-Thiophenecarboxaldehyde [98-03-3] M 112.2, b 106°/30mm, d 1.593, n 1.222. Washed with 50% HCl and distd under reduced pressure just before use.

Thiophene-2-carboxylic acid [527-72-0] M 128.2, m 129-130°, pK²⁵ 3.89. Crystd from water.

Thiophene-3-carboxylic acid [88-31-1] M 128.1, m 137-138°, pK²⁵ 6.23. Crystd from water.

Thiophenol (benzenethiol) [108-98-5] M 110.2, f -14.9°, b 46.4°/10mm, 168.0°/760mm, d 1.073, n 1.5897, pK²⁵ 6.62. Dried with CaCl₂ or CaSO₄, and distd at 10mm pressure or at 100mm (b 103.5°) in a stream of nitrogen.

Thiopyronine (2,7-dimethylaminothioxanthane) [2412-14-8] M 318.9, λ_{\max} 564nm (ϵ 78,500) H₂O, pK_{Est} ~ 7. Purified as the hydrochloride by recrystn from hydrochloric acid. [Fanghanel et al. *J Phys Chem* 91 3700 1987.]

Thiosalicylic (2-mercaptobenzoic) acid [147-93-3] M 154.2, m 164-165°, pK₁²⁵ 3.54, pK₂²⁵ 8.80. Crystd from hot EtOH (4mL/g), after adding hot distd water (8mL/g) and boiling with charcoal. The hot soln was filtered, cooled, the solid collected and dried *in vacuo* (P₂O₅). Cryst from AcOH and sublimes *in vacuo*.

Thiosemicarbazide [79-19-6] M 91.1, m 181-183°, pK₁²⁵ 1.88, pK₂²⁵ 12.81. Crystd from water.

Thiothienoyltrifluoroacetone [4552-64-1] M 228.2, m 61-62°. Easily oxidised and has to be purified before use. This may be by recrystd from *benzene or by dissolution in pet ether, extraction into 1M NaOH soln, acidification of the aqueous phase with 1-6M HCl soln, back extraction into pet ether and final evapn of the solvent. The purity can be checked by TLC. It was stored in ampoules under nitrogen at 0° in the dark. [Muller and Rother *Anal Chim Acta* 66 49 1973.]

Thiouracil [141-90-2] M 128.2, m 240°(dec), pK²⁵ 7.52. Crystd from water or EtOH.

Thiourea [62-56-6] M 76.1, m 179°, pK²⁰ -1.19 (aq H₂SO₄). Crystd from absolute EtOH, MeOH, acetonitrile or water. Dried under vacuum over H₂SO₄ at room temperature.

9H-Thioxanthene-9-one (thioxanthone) [492-22-8] M 212.3, m 200-202°, 209°, 212-214°, b 371-373°/712mm. Yellow needles from CHCl₃ or EtOH and sublimes *in vacuo*. Sol in CS₂, hot AcOH and soln in concn H₂SO₄ to give a yellow color with green fluorescence in VIS light. The *sulfone* has m 187° (from EtOH), and the *hydrazone* has m 115° (yellow leaflets from EtOH/*C₆H₆). [Szmant et al. *J Org Chem* 18 745 1953; Ullmann et al. *Chem Ber* 49 2509 1916; NMR: Sharpless et al. *Org Magn Res* 6 115 1974.]

L-Threonine [72-19-5] M 119.1, m 251-253°, [α]_D²⁶ -28.4° (H₂O), pK₁²⁵ 2.17, pK₂²⁵ 9.00. Likely impurities are *allo*-threonine and glycine. Crystd from water by adding 4 volumes of EtOH. Dried and stored in a desiccator.

Thymidine [50-89-5] M 242.2, m 185°, pK₂²⁵ 9.65. Crystd from ethyl acetate.

Thymine [65-71-4] M 126.1, m 326°, pK²⁵ 9.82. Crystd from ethyl acetate or water. Purified by preparative (2mm thick) TLC plates of silica gel, eluting with ethyl acetate/isopropanol/water (75:16:9, v/v; R_F 0.75). Spot localised by uv lamp, cut from plate, placed in MeOH, shaken and filtered through a millipore filter, then rotary evapd. [Infante et al. *J Chem Soc, Faraday Trans 1* 68 1586 1973.]

Thymolphthalein complexone [1913-93-5] M 720.8, m 190°(dec), pK₁^{18.2} 7.35, pK₂^{18.2} 12.25. Purification as for phthalein complexone except that it was synthesised from thymolphthalein instead of cresolphthalein.

Tiglic acid [80-59-1] M 100.1, m 63.5-64°, b 198.5°, pK¹⁸ 4.96. Crystd from water.

Tinuvin P (2-[2H-benzotriazol-2-yl]-p-cresol) [50936-05-5] M 225.3, m 131-133°, pK_{Est(1)} ~ 1.6 (N protonation), pK_{Est(2)} ~ 8 (phenolic OH). Recrystd from *n*-heptane or Me₂CO/pentane. [Woessner et al. *J Phys Chem* 81 3629 1985.]

***o*-Tolidine (3,3'-dimethylbenzidine)** [119-93-7] M 212.3, m 131-132°, pK²⁵ 4.45. Dissolved in *benzene, percolated through a column of activated alumina and crystd from *benzene/pet ether.

***p*-Tolualdehyde** [104-87-0] M 120.2, b 83-85°/0.1mm, 199-200°/760mm, d 1.018, n 1.548. Steam distd, dried with CaSO₄ and fractionally distd.

***o*-Toluamide** [527-85-5] M 135.2, m 141°. Crystd from hot water (10mL/g) and dried in air.

Toluene [108-88-3] M 92.1, b 110.6°, d¹⁰ 0.87615, d²⁵ 0.86231, n 1.49693, n²⁵ 1.49413. Dried with CaCl₂, CaH₂ or CaSO₄, and dried further by standing with sodium, P₂O₅ or CaH₂. It can be fractionally distd from sodium or P₂O₅. Unless specially purified, toluene is likely to be contaminated with methylthiophenes and other sulfur containing impurities. These can be removed by shaking with conc H₂SO₄, but the temperature must be kept below 30° if sulfonation of toluene is to be avoided. A typical procedure consists of shaking toluene twice with cold conc H₂SO₄ (100mL of acid per L), once with water, once with aqueous 5% NaHCO₃ or NaOH, again with H₂O, then drying successively with CaSO₄ and P₂O₅, with final distn from P₂O₅ or over LiAlH₄ after refluxing for 30min. Alternatively, the treatment with NaHCO₃ can be replaced by boiling under reflux with 1% sodium amalgam. Sulfur compounds can also be removed by prolonged shaking of the toluene with mercury, or by two distns from AlCl₃, the distillate then being washed with water, dried with K₂CO₃ and stored with sodium wire. Other purification procedures include refluxing and distn of sodium dried toluene from diphenylpicrylhydrazyl, and from SnCl₂ (to ensure freedom from peroxides). It has also been co-distd with 10% by volume of ethyl methyl ketone, and again fractionally distd. [Brown and Pearsall *J Am Chem Soc* 74 191 1952.] For removal of carbonyl impurities see **benzene*. Toluene has been purified by distn under nitrogen in the presence of sodium benzophenone ketyl. Toluene has also been dried with MgSO₄, after the sulfur impurities have been removed, and then fractionally distd from P₂O₅ and stored in the dark [Tabushi et al. *J Am Chem Soc* 107 4465 1985]. Toluene can be purified by passage through a tightly packed column of Fuller's earth.

Rapid purification: Alumina, CaH₂ and 4A molecular sieves (3% w/v) may be used to dry toluene (6h stirring and standing). Then the toluene is distd, discarding the first 5% of distillate, and is stored over molecular sieves (3A, 4A) or Na wire.

Toluene-2,4-diamine [95-80-7] M 122.2, m 99°, b 148-150°/8mm, 292°/760mm, pK_{Est(1)}~2.5, pK_{Est(2)}~4.4. Recrystd from water containing a very small amount of sodium dithionite (to prevent air oxidation), and dried under vacuum. Also cryst from **benzene*.

***o*-Toluenesulfonamide** [88-19-7] M 171.2, m 155.5°. Crystd from hot water, then from EtOH or Et₂O-pet ether.

***p*-Toluenesulfonamide** [70-55-3] M 171.2, m 137-137.5°, 138°. Crystd from hot water, then from EtOH or Et₂O-pet ether.

***p*-Toluenesulfonic acid** [6192-52-5] M 190.2, m 38° (anhydrous), m 105-107° (monohydrate), pK 1.55. Purified by pptn from a satd soln at 0° by introducing HCl gas. Also crystd from conc HCl, then crystd from dilute HCl (charcoal) to remove benzenesulfonic acid. It has been crystd from EtOH/water. Dried in a vacuum desiccator over solid KOH and CaCl₂. *p*-Toluenesulfonic acid can be dehydrated by azeotropic distn with **benzene* or by heating at 100° for 4h under water-pump vacuum. The anhydrous acid can be crystd from **benzene*, CHCl₃, ethyl acetate, anhydrous MeOH, or from acetone by adding a large excess of **benzene*. It can be dried under vacuum at 50°.

Toluenesulfonic acid hydrazide (tosylhydrazide) [1576-35-8] M 186.2, m 108-110°, 109-110°. Dissolve in hot MeOH (~1g/4mL), filter through Celite and ppte material by adding 2-2.5 vols of distd H₂O. Air or vac dry. [Fiedman et al. *Org Synth Coll Vol V* 1055 1973.]

***p*-Toluenesulfonyl chloride (tosyl chloride)** [98-59-9] M 190.7, m 66-69°, 67.5-68.5°, 69°, b 138-139°/9mm, 146°/15mm, 167°/36mm. Material that has been standing for a long time contains tosic acid and HCl and has m ca 65-68°. It is purified by dissolving (10g) in the minimum volume of CHCl₃ (ca 25mL) filtered, and diluted with five volumes (i.e. 125mL) of pet ether (b 30-60°) to precipitate impurities. The soln is filtered, clarified with charcoal and concentrated to 40mL by evaporation. Further evaporation to a very small volume gave 7g of white crystals which were analytically pure, m 67.5-68.5°. (The insoluble material was largely tosic acid and had m 101-104°). [Pelletier *Chem Ind (London)* 1034 1953.] Also crystd from toluene/pet ether in the cold, from pet ether (b 40-60°) or **benzene*. Its soln in diethyl ether has been washed with aqueous 10% NaOH until colourless, then dried (Na₂SO₄) and crystd by cooling in

powdered Dry-ice. It has also been purified by dissolving in *benzene, washing with aqueous 5% NaOH, then dried with K_2CO_3 or $MgSO_4$, and distd under reduced pressure and can be sublimed at high vacuum [Ebel *Chem Ber* **60** 2086/1927].

p-Toluenethiol [106-45-6] M 124.2, m 43.5-44°, pK^{25} 6.82. Crystd from pet ether (b 40-70°).

Tolhydroquinone (2-methylbenzene-1,4-diol) [95-71-6] M 124.1, m 128-129°, pK_1^{20} 10.15, pK_2^{20} 11.75. Crystd from EtOH.

o-Toluic acid [118-90-1] M 136.2, m 102-103°, pK^{25} 3.91. Crystd from *benzene (2.5mL/g) and dried in air.

m-Toluic acid [99-04-7] M 136.2, m 111-113°, pK^{25} 4.27. Crystd from water.

p-Toluic acid [99-94-5] M 136.2, m 178.5-179.5°, pK^{25} 4.37. Crystd from water, water/EtOH (1:1), MeOH/water or *benzene.

o-Toluidine (2-methylaniline) [95-53-4] M 107.2, f -16.3°, b 80.1°/10mm, 200.3°/760mm, d 0.999, n 1.57246, n^{25} 1.56987, pK^{25} 4.45. In general, methods similar to those for purifying aniline can be used, e.g. distn from zinc dust, at reduced pressure, under nitrogen. Berliner and May [*J Am Chem Soc* **49** 1007 1927] purified *via* the oxalate. Twice-distd *o*-toluidine was dissolved in four times its volume of diethyl ether and the equivalent amount of oxalic acid needed to form the dioxalate was added as its soln in diethyl ether. (If *p*-toluidine is present, its oxalate pptes and can be removed by filtration.) Evapn of the ether soln gave crystals of *o*-toluidine dioxalate. They were filtered off, recrystd five times from water containing a small amount of oxalic acid (to prevent hydrolysis), then treated with dilute aqueous Na_2CO_3 to liberate the amine which was separated, dried ($CaCl_2$) and distd under reduced pressure.

m-Toluidine (3-methylaniline) [108-44-1] M 107.2, f -30.4°, b 82.3°/10mm, 203.4°/760mm, d 0.989, n 1.56811, n^{25} 1.56570, pK^{25} 4.71. It can be purified as for aniline. Twice-distd, *m*-toluidine was converted to the hydrochloride using a slight excess of HCl, and the salt was fractionally crystd from 25% EtOH (five times), and from distd water (twice), rejecting, in each case, the first material that crystd. The amine was regenerated and distd as for *o*-toluidine. [Berliner and May *J Am Chem Soc* **49** 1007 1927.]

p-Toluidine (4-methylaniline) [106-49-0] M 107.2, m 44.8°, b 79.6°/10mm, 200.5°/760mm, d 0.962, n 1.5636, $n^{59.1}$ 1.5534, pK^{25} 5.08. In general, methods similar to those for purifying aniline can be used. It can be separated from the *o*- and *m*-isomers by fractional crystn from its melt. *p*-Toluidine has been crystd from hot water (charcoal), EtOH, *benzene, pet ether or EtOH/water (1:4), and dried in a vacuum desiccator. It can also be sublimed at 30° under vacuum. For further purification, use has been made of the oxalate, the sulfate and acetylation. The oxalate, formed as described for *o*-toluidine, was filtered, washed and recrystd three times from hot distd water. The base was regenerated with aq Na_2CO_3 and recrystd three times from distd water. [Berliner and May *J Am Chem Soc* **49** 1007 1927.] Alternatively, *p*-toluidine was converted to its acetyl derivative which, after repeated crystn from EtOH, was hydrolysed by refluxing (50g) in a mixture of 500mL of water and 115mL of conc H_2SO_4 until a clear soln was obtained. The amine sulfate was isolated, suspended in water, and NaOH was added. The free base was distd twice from zinc dust under vacuum. The *p*-toluidine was then recrystd from pet ether and dried in a vacuum desiccator or in a vacuum for 6h at 40°. [Berliner and Berliner *J Am Chem Soc* **76** 6179 1954; Moore et al. *J Am Chem Soc* **108** 2257 1986.]

Toluidine Blue O [93-31-9] M 305.8, CI 52040, λ_{max} 626nm, pK^{25} 7.5. Crystd from hot water (18mL/g) by adding one and a half volumes of alcohol and chilling on ice. Dried at 100° in an oven for 8-10h.

p-Toluidine hydrochloride [540-23-8] M 143.6, m 245.9-246.1°. Crystd from MeOH containing a few drops of conc HCl. Dried under vacuum over paraffin chips.

2-*p*-Toluidinylnaphthalene-6-sulfonic acid [7724-15-4] M 313.9, $pK_{Est} \sim 0$. Crystd twice from 2% aqueous KOH and dried under high vacuum for 4h at room temperature. Crystd from water. Tested for purity by TLC on silica gel with isopropanol as solvent. The free acid was obtained by acidifying a satd aqueous soln.

***o*-Tolunitrile** [529-19-1] M 117.2, b 205.2°, d 0.992, n 1.5279. Fractionally distd, washed with conc HCl or 50% H₂SO₄ at 60° until the smell of isonitrile had gone (this also removed any amines), then washed with saturated NaHCO₃ and dilute NaCl solns, then dried with K₂CO₃ and redistd.

***m*-Tolunitrile** [620-22-4] M 117.2, b 209.5-210°/773mm, d 0.986, n 1.5250. Dried with MgSO₄, fractionally distd, then washed with aqueous acid to remove possible traces of amines, dried and redistd.

***p*-Tolunitrile** [104-85-8] M 117.2, m 29.5°, b 104-106°/20mm. Melted, dried with MgSO₄, fractionally crystd from its melt, then fractionally distd under reduced pressure in a 6-in spinning band column. [Brown *J Am Chem Soc* 81 3232 1959.] It can also be crystd from *benzene/pet ether (b 40-60°).

4-Tolyl-2-benzoic acid (4'-methylbiphenyl-2-carboxylic acid) [7148-03-0] M 196.2, m 138-139°, $pK^{25} 3.64$. Crystd from toluene.

***p*-Tolylacetic acid** [622-47-9] M 150.2, m 94°. See 4-methylphenylacetic acid on p. 297.

***p*-Tolyl carbinol (4-methylbenzyl alcohol)** [589-18-4] M 122.2, m 61°, b 116-118°/20mm, 217°/760mm. Crystd from pet ether (b 80-100°, 1g/mL). It can also be distd under reduced pressure.

***p*-Tolyl disulfide** [103-19-5] M 246.4, m 45-46°. Purified by chromatography on alumina using hexane as eluent, then crystd from MeOH. [Kice and Bowers *J Am Chem Soc* 84 2384 1962.]

***p*-Tolylsulfonylmethyl isocyanide (tosylmethyl isocyanide, TOSMIC)** [36635-61-7] M 195.2, m 114-115°(dec), 116-117°(dec). Use an efficient fume cupboard. Purify by dissolving TOSMIC (50g) in CH₂Cl₂ (150mL) and pass through a column (40x3cm) containing neutral alumina (100g) in CH₂Cl₂ and eluting with CH₂Cl₂. A nearly colorless soln (700mL) is collected, evaporated *in vacuo* and the residue (42-47g) of TOSMIC (m 113-114° dec) is recrystd once from MeOH (m 116-117° dec). [Hoogenboom et al. *Org Synth* 57 102 1977; Lensen *Tetrahedron Lett* 2367 1972..] Also recrystd from EtOH (charcoal) [Saito and Itano, *J Chem Soc, Perkin Trans 1* 1 1986].

***p*-Tolyl urea** [622-51-5] M 150.2, m 181°. Crystd from EtOH/water (1:1).

***trans*-Traumatic acid (2-dodecene-1,12-dioic acid)** [6402-36-4] M 228.3, m 165-166°, 150-160°/0.001mm, $pK_{Est(1)} \sim 4.2$, $pK_{Est(2)} \sim 4.6$. Crystd from EtOH, acetone or glyme.

α, α' -Trehalose (2H₂O) [6138-23-4] M 378.3, m 96.5-97.5°, 203° (anhydrous). Crystd (as the dihydrate) from aqueous EtOH. Dried at 13°.

1,2,3-Triaminopropane trihydrochloride [free base 21291-99-6] M 198.7, m 250°, $pK_1^{20} 3.72$, $pK_2^{20} 7.95$, $pK_3^{20} 9.59$. Crystd from EtOH.

1,5,7-Triazabicyclo[4.4.0]dec-5-ene (TBD, 1,3,4,6,7,8-hexahydro-2*h*-pyrimido[1,2-*a*]-pyrimidine) [5807-14-7] M 139.2, m 125-130°, $pK \sim 16$ Cryst from Et₂O but readily forms white crystals of the carbonate. It is a strong base (see pK , i.e. about 100 times more basic than tetramethylguanidine. The *picrate* has m 220.5-222° (from EtOH). Forms the *5-nitro derivative* m 14.5-160° that gives a *5-nitro nitrate* salt m 100-101° (from EtOH-Et₂O) and a *5-nitro picrate* m 144-145° (from H₂O). [McKay and Kreling *Can J Chem* 35 1438 1957; Schwesinger *Chimia* 39 369 1985; Hilpert et al. *J Chem Soc, Chem Commun* 1401 1983; Kamfen and Eschenmoser *Helv Chim Acta* 72 185 1989].

1,2,4-Triazole [288-88-0] M 69.1, m 121°, 260°, $pK_1^{25} 2.27$ (basic), $pK_2^{25} 10.26$ (acidic). Crystd from EtOH or water [Barszcz et al. *J Chem Soc, Dalton Trans* 2025 1986].

Tribenzylamine [620-40-6] M 287.4, m 93-94°, 230°/13mm, $pK_{\text{Est}} < 0$. Crystd from abs EtOH or pet ether. Dried in a vacuum over P_2O_5 at room temperature. *HCl* has m 226-228° (from EtOH) and *picrate* has m 191° (from H_2O or aq EtOH).

2,4,6-Tribromoacetanilide [607-93-2] M 451.8, m 232°. Crystd from EtOH.

2,4,6-Tribromoaniline [147-82-0] M 329.8, m 120°, $pK_{\text{Est}} \sim -0.5$ (aq H_2SO_4). Crystd from MeOH.

sym-Tribromobenzene [626-39-1] M 314.8, m 122°. Crystd from glacial acetic acid/water (4:1), then washed with chilled EtOH and dried in air.

Tribromochloromethane [594-15-0] M 287.2, m 55°. Melted, washed with aqueous $Na_2S_2O_3$, dried with BaO and fractionally crystd from its melt.

2,4,6-Tribromophenol [118-79-6] M 330.8, m 94°, pK^{25} 6.00. Crystd from EtOH or pet ether. Dried under vacuum over P_2O_5 at room temperature.

Tri-*n*-butylamine [102-82-9] M 185.4, b 68°/3mm, 120°/44mm, d 0.7788, n 1.4294, pK^{25} 9.93. Purified by fractional distn from sodium under reduced pressure. Pegolotti and Young [*J Am Chem Soc* 83 3251 1961] heated the amine overnight with an equal volume of acetic anhydride, in a steam bath. The amine layer was separated and heated with water for 2h on the steam bath (to hydrolyse any remaining acetic anhydride). The soln was cooled, solid K_2CO_3 was added to neutralize any acetic acid that had been formed, and the amine was separated, dried (K_2CO_3) and distd at 44mm pressure. Davis and Nakshbendi [*J Am Chem Soc* 84 2085 1926] treated the amine with one-eighth of its weight of benzenesulfonyl chloride in aqueous 15% NaOH at 0-5°. The mixture was shaken intermittently and allowed to warm to room temperature. After a day, the amine layer was washed with aq NaOH, then water and dried with KOH. (This treatment removes primary and secondary amines.) It was further dried with CaH_2 and distd under vacuum.

Tri-*n*-butylammonium hydrobromide [37026-85-0] M 308.3, m 75.2-75.9°. Crystd from ethyl acetate.

Tri-*n*-butylammonium nitrate [33850-87-2] M 304.5. Crystd from mixtures of *n*-hexane and acetone (95:5). Dried over P_2O_5 .

Tri-*n*-butylammonium perchlorate [14999-66-7] M 285.5. Recrystd from *n*-hexane.

sym-Tri-*tert*-butylbenzene [1460-02-2] M 246.4, m 73.4-73.9°. Crystd from EtOH.

2,4,6-Tri-*tert*-butylphenol [732-26-3] M 262.4, m 129-132°, 131°/1mm, 147°/10mm, 278°/760mm, pK^{25} 12.19. Crystd from *n*-hexane or several times from 95% EtOH until the EtOH soln was colourless [Balasubramanian and Bruice *J Am Chem Soc* 108 5495 1986]. It has also been purified by sublimation [Yuan and Bruice *J Am Chem Soc* 108 1643 1986; Wong et al. *J Am Chem Soc* 109 3428 1987]. Purification has been achieved by passage through a silica gel column followed by recrystn from *n*-hexane [Kajii et al. *J Phys Chem* 91 2791 1987].

Tricarballic acid (propane-1,2,3-tricarboxylic acid) [99-14-9] M 176.1, m 166°, pK_1^{25} 3.47, pK_2^{25} 4.54, pK_3^{25} 5.89. Crystd from diethyl ether.

Trichloroacetamide [594-65-0] M 162.4, m 139-141°, b 238-240°. Its xylene soln was dried with P_2O_5 , then fractionally distd.

Trichloroacetanilide [2563-97-5] M 238.5, m 95°. Crystd from *benzene.

Trichloroacetic acid [76-03-9] M 163.4, m 59.4-59.8°, pK²⁵ 0.51. Purified by fractional crystn from its melt, then crystd repeatedly from dry *benzene and stored over conc H₂SO₄ in a vac desiccator. It can also be crystd from CHCl₃ or cyclohexane, and dried over P₂O₅ or Mg(ClO₄)₂ in a vac desiccator. Trichloroacetic acid can be fractionally distd under reduced pressure from MgSO₄. Layne, Jaffé and Zimmer [*J Am Chem Soc* 85 435 1963] dried trichloroacetic acid in *benzene by distilling off the *benzene-water azeotrope, then crystd the acid from the remaining *benzene soln. Manipulations were carried out under nitrogen. [Use a well ventilated fume cupboard].

2,3,4-Trichloroaniline [634-67-3] M 196.5, m 67.5°, b 292°/774mm, pK_{Est} ~1.3. Crystd from ligroin.

2,4,5-Trichloroaniline [636-30-6] M 196.5, m 96.5°, b 270°/760mm, pK 1.09. Crystd from ligroin.

2,4,6-Trichloroaniline [634-93-5] M 196.5, m 78.5°, b 127°/14mm, 262°/746mm, pK 0.03. Crystd from ligroin.

1,2,3-Trichlorobenzene [87-61-6] M 181.5, m 52.6°. Crystd from EtOH.

1,2,4-Trichlorobenzene [120-82-1] M 181.5, m 17°, b 210°. Separated from a mixture of isomers by washing with fuming H₂SO₄, then water, drying with CaSO₄ and slowly fractionally distilling. [Jensen, Marino and Brown *J Am Chem Soc* 81 3303 1959.]

1,3,5-Trichlorobenzene [108-70-3] M 181.5, m 64-65°. Recrystd from dry *benzene or toluene.

3,4,5-Trichloro-*o*-cresol (3,4,5-trichloro-2-methylphenol) [608-92-4] M 211.5, m 77°, pK_{Est} ~7.6. Crystd from pet ether.

2,3,5-Trichloro-*p*-cresol [608-91-3] M 211.5, m 66-67°, pK_{Est} ~6.9. Crystd from pet ether.

1,1,1-Trichloroethane [71-55-6] M 133.4, f -32.7°, b 74.0°, d 1.337, n 1.4385. Washed successively with conc HCl (or conc H₂SO₄), aq 10% K₂CO₃ (Na₂CO₃), aq 10% NaCl, dried with CaCl₂ or Na₂SO₄, and fractionally distd. It can contain up to 3% dioxane as preservative. This is removed by washing successively with 10% aq HCl, 10% aq NaHCO₃ and 10% aq NaCl; and distd over CaCl₂ before use.

1,1,2-Trichloroethane [79-00-5] M 133.4, f -36.3°, b 113.6°, d 1.435, n 1.472. Purification as for 1,1,1-trichloroethane above.

Trichloroethylene [79-01-6] M 131.4, f -88°, b 87.2°, d 1.463, n²¹ 1.4767. Undergoes decomposition in a similar way to CHCl₃, giving HCl, CO, COCl₂ and organic products. It reacts with KOH, NaOH and 90% H₂SO₄, and forms azeotropes with water, MeOH, EtOH, and acetic acid. It is purified by washing successively with 2M HCl, water and 2M K₂CO₃, then dried with K₂CO₃ and CaCl₂, and fractionally distd immediately before use. It has also been steam distd from 10% Ca(OH)₂ slurry, most of the water being removed from the distillate by cooling to -30° to -50° and filtering off the ice through chamois skin: the trichloroethylene was then fractionally distd at 250mm pressure and collected in a blackened container. [Carlisle and Levine *Ind Eng Chem (Anal Ed)* 24 1164 1932.]

2,4,5-Trichloro-1-nitrobenzene [89-69-0] M 226.5, m 57°. Crystd from EtOH.

3,4,6-Trichloro-2-nitrophenol [82-62-2] M 242.4, m 92-93°, pK_{Est} ~4.1. Crystd from pet ether or EtOH.

2,4,5-Trichlorophenol [95-95-4] M 197.5, m 67°, b 72°/1mm, pK²⁵ 7.0. Crystd from EtOH or pet ether.

2,4,6-Trichlorophenol [88-06-2] M 197.5, m 67-68°, pK²⁵ 6.23. Crystd from *benzene, EtOH or EtOH/water.

3,4,5-Trichlorophenol [609-19-8] M 197.5, m 100°, pK²⁵ 7.84. Crystd from pet ether/*benzene mixture.

2,4,5-Trichlorophenoxyacetic acid (2,4,5-T) [93-76-5] M 255.5, m 153°, 155-158°, pK²⁵ 2.83. Crystd from *benzene. (CANCER SUSPECT)

1,1,2-Trichlorotrifluoroethane [76-13-1] M 187.4, b 47.6°/760mm, d 1.576, n 1.360. Washed with water, then with weak alkali. Dried with CaCl₂ or H₂SO₄ and distd. [Locke et al. *J Am Chem Soc* 56 1726 1934.]

Tricycloquinazoline [195-84-6] M 230.3, m 322-323°. Crystd repeatedly from toluene, followed by vac sublimation at 210° at a pressure of 0.15-0.3 Torr in subdued light.

Tridecanoic acid [638-53-9] M 214.4, m 41.8°, 44.5-45.5° (several forms), b 199-200°/24mm, pK_{Est} ~5.0. Crystd from acetone.

7-Tridecanone [462-18-0] M 198.4, m 33°, b 255°/766mm. Crystd from EtOH.

Tri-*n*-dodecylammonium nitrate [2305-34-2] M 585.0. Crystd from *n*-hexane/acetone (95:5) and kept in a desiccator over P₂O₅.

Tri-*n*-dodecylammonium perchlorate [5838-82-4] M 622.4. Recrystd from *n*-hexane or acetone and kept in a desiccator over P₂O₅.

Triethanolamine hydrochloride [637-39-8] M 185.7, m 177°, pK²⁵ 7.92 (free base). Crystd from EtOH. Dried at 80°.

1,1,2-Triethoxyethane [4819-77-6] M 162.2, b 164°, d 0.897, n 1.401. Dried with Na₂SO₄, and distd.

Triethylamine [121-44-8] M 101.2, b 89.4°, d 0.7280, n 1.4005, pK²⁵ 10.82. Dried with CaSO₄, LiAlH₄, Linde type 4A molecular sieves, CaH₂, KOH, or K₂CO₃, then distd, either alone or from BaO, sodium, P₂O₅ or CaH₂. It has also been distd from zinc dust, under nitrogen. To remove traces of primary and secondary amines, triethylamine has been refluxed with acetic anhydride, benzoic anhydride, phthalic anhydride, then distd, refluxed with CaH₂ (ammonia-free) or KOH (or dried with activated alumina), and again distd. Another purification involved refluxing for 2h with *p*-toluenesulfonyl chloride, then distd. Grovenstein and Williams [*J Am Chem Soc* 83 412 1961] treated triethylamine (500mL) with benzoyl chloride (30mL), filtered off the ppte, and refluxed the liquid for 1h with a further 30mL of benzoyl chloride. After cooling, the liquid was filtered, distd, and allowed to stand for several hours with KOH pellets. It was then refluxed with, and distd from, stirred molten potassium. Triethylamine has been converted to its hydrochloride, crystd from EtOH (to m 254°), then liberated with aq NaOH, dried with solid KOH and distd from sodium under nitrogen.

Triethylammonium hydrobromide [636-70-4] M 229.1, m 248°. Equimolar portions of triethylamine and aqueous solutions of HBr in acetone were mixed. The ppted salt was washed with anhydrous acetone and dried in vacuum for 1-2h. [Odiakov et al. *J Chem Soc, Faraday Trans 2* 80 899 1984.] Recrystd from CHCl₃ or EtOH.

Triethylammonium hydrochloride [554-68-7] M 137.7, m 257-260°(dec). Purified like the bromide above.

Triethylammonium hydroiodide [4636-73-1] M 229.1, m 181°. Purified as for triethylammonium bromide, except the soln for pptn was precooled acetone at -10° and the ppte was twice recrystd from a cooled acetone/hexane mixture at -10°.

Triethylammonium trichloroacetate [4113-06-8] M 263.6. Equimolar solns of triethylamine and trichloroacetic acid in *n*-hexane were mixed at 10°. The solid so obtained was recrystd from CHCl₃/*benzene.

Triethylammonium trifluoroacetate [454-49-9] M 196.2. Purified as for the corresponding trichloroacetate. The salt was a colourless liquid at ambient temperature.

1,2,4-Triethylbenzene [877-44-1] M 162.3, b 96.8-97.1°/12.8mm, d 0.8738, n 1.5015. For separation from a commercial mixture see Dillingham and Reid [*J Am Chem Soc* 60 2606 1938].

1,3,5-Triethylbenzene [102-25-0] M 162.3, b 102-102.5°, d 0.8631, n 1.4951. For separation from a commercial mixture see Dillingham and Reid [*J Am Chem Soc* 60 2606 1938].

Triethylene glycol [112-27-6] M 150.2, b 115-117°/0.1mm, 278°/760mm, n¹⁵ 1.4578, d¹⁵ 1.1274. Dried with CaSO₄ for 1 week, then repeatedly and very slowly fractionally distd under vacuum. Stored in a vacuum desiccator over P₂O₅. It is very *hygroscopic*.

Triethylene glycol dimethyl ether (triglyme) [112-49-2] M 178.2, b 225°, d 0.987, n 1.425. Refluxed with, and distd from sodium hydride or LiAlH₄.

Triethylenetetramine (TRIEN, TETA, trientine) [112-24-3] M 146.2, m 12°, b 157°/20mm, d 0.971, n 1.497, pK₁^{2.5} 3.32, pK₂^{2.5} 6.67, pK₃^{2.5} 9.20, pK₄^{2.5} 9.92. Dried with sodium, then distd under vac. Further purification has been *via* the nitrate or the chloride. For example, Jonassen and Strickland [*J Am Chem Soc* 80 312 1958] separated TRIEN from admixture with TREN (38%) by soln in EtOH, cooling to approximately 5° in an ice-bath and adding conc HCl dropwise from a burette, keeping the temperature below 10°, until all of the white crystalline ppte of TREN.HCl had formed and been removed. Further addition of HCl then pptd thick creamy white TRIEN.HCl which was crystd several times from hot water by adding an excess of cold EtOH. The crystals were finally washed with Me₂CO, then Et₂O and dried in a vacuum desiccator.

Triethylenetetramine tetrahydrochloride (TRIEN.HCl) [4961-10-4] M 292.1, m 266-270°. Crystd repeatedly from hot water by pptn with cold EtOH or EtOH/HCl. Washed with acetone and abs EtOH and dried in a vacuum oven at 80° (see TRIEN above).

Triethyl orthoformate (1,1,1-triethoxymethane) [122-51-0] M 148.2, m 30°, b 60°/30mm, 144-146°, d 0.891, n 1.392. Fractionate first at atm press, then in a vac. If impure, then wash with H₂O, dry over anhyd K₂CO₃, filter and fractionate through a Widmer column. [Sah and Ma *J Am Chem Soc* 54 2964 1932; Ohme and Schmitz *Justus Liebigs Ann Chem* 716 207 1968.] **IRRITANT** and **FLAMMABLE**.

Triethyloxonium fluoroborate [368-39-8] M 190.0, m 92-93°(dec). Crystd from diethyl ether. *Very hygroscopic*, and should be handled in a dry box and stored at 0°. [*Org Synth* 46 113 1966.] Pure material should give a clear and colourless soln in dichloromethane (1 in 50, w/v).

Trifluoroacetic acid [76-05-1] M 114.0, f -15.5°, b 72.4°, d 1.494, n 1.2850, pK^{2.5} 0.52. The purification of trifluoroacetic acid, reported in earlier editions of this work, by refluxing over KMnO₄ for 24h and slowly distilling has resulted in very **SERIOUS EXPLOSIONS** on various occasions, but not always. This apparently depends on the source and/or age of the acid. The method is **NOT RECOMMENDED**. Water can be removed by adding trifluoroacetic anhydride (0.05%, to diminish water content) and distd. [Conway and Novak *J Phys Chem* 81 1459 1977]. It can be refluxed and distd from P₂O₅. It is further purified by fractional crystn by partial freezing and again distd. **Highly TOXIC vapour**.

Trifluoroacetic anhydride [407-25-0] M 210.0, b 38-40°/760mm, d 1.508. Purification by distilling over KMnO_4 , as for the acid above is **EXTREMELY DANGEROUS** due to the possibility of **EXPLOSION**. It is best purified by distilling from P_2O_5 slowly, and collecting the fraction boiling at 39.5°. Store in a dry atmosphere. **Highly TOXIC vapour**.

1,1,1-Trifluoro-2-bromoethane [421-06-7] M 163.0. See 1-bromo-3,3,3-trifluoroethane on p. 142.

2,2,2-Trifluoroethanol [75-89-8] M 100.0, b 72.4°/738mm, d 1.400, pK^{25} 12.8. Dried with CaSO_4 and a little NaHCO_3 (to remove traces of acid). **Highly TOXIC vapour**.

Trifluoromethanesulfonic anhydride (triflic anhydride) [358-23-6] M 282.1, b 82-85°, 84°, d 1.71, n 1.322. Distil through a short Vigreux column. Could be freshly prepd from the anhydrous acid (11.5g) and P_2O_5 (11.5g, or half this weight) by setting aside at room temp for 1h, distilling off volatile products then through a short Vigreux column. Readily hydrolysed by H_2O and decomposes appreciably after a few days to liberate SO_2 and produce a viscous liquid. Store dry at low low temp. [Burdon et al. *J Chem Soc* 2574 1957; Beard et al. *J Org Chem* 38 373 1973.] **Highly TOXIC vapour**.

4-(Trifluoromethyl)acetophenone [709-63-7] M 188.2, m 31-33°, b 79-81°/9mm. Purified by distillation or sublimation *in vacuo*.

3-Trifluoromethyl-4-nitrophenol [88-30-2] M 162.1, m 81°, pK_{Est} ~6.1. Crystd from *benzene or from pet ether/*benzene mixture.

α,α,α -Trifluorotoluene (benzotrifluoride) [98-08-8] M 144.1, b 102.5°, d 1.190, n^{30} 1.4100. Purified by repeated treatment with boiling aqueous Na_2CO_3 (until no test for chloride ion was obtained), dried with K_2CO_3 , then with P_2O_5 , and fractionally distd.

Triglycyl glycine (tetraglycine) [637-84-3] M 246.2, m 270-275°(dec). Crystd from distilled water (optionally, by the addition of EtOH).

Trigonelline (1-methylnicotinic acid zwitterion) [535-83-1] M 137.1, m 218°(dec). Crystd (as monohydrate) from aqueous EtOH, then dried at 100°.

2,3,4-Trihydroxybenzoic acid [610-02-6] M 170.1, m 207-208°, $\text{pK}_{\text{Est}(1)}$ ~3.4, $\text{pK}_{\text{Est}(2)}$ ~7.8, $\text{pK}_{\text{Est}(3)}$ >12. Crystd from water.

2,4,6-Trihydroxybenzoic acid [83-30-7] M 170.1, m 205-212°(dec), $\text{pK}_{\text{Est}(1)}$ ~1.5, $\text{pK}_{\text{Est}(2)}$ ~8.0, $\text{pK}_{\text{Est}(3)}$ >12. Crystd from water.

4',5,7-Trihydroxyflavone (apigenin) [520-36-5] M 270.2, m 296-298°, 300-305°, 345-350° (pK 's 7—10, for phenolic OH). Crystd from aq pyridine or aq EtOH. Dyes wool yellow when with added Cr.

3,4,5-Triiodobenzoic acid [2338-20-7] M 499.8, m 289-290°, 293°, pK^{25} 0.65. Crystd from aqueous EtOH or water.

3,4,5-Triiodobenzyl chloride [52273-54-8] M 504.3, m 138°. Crystd from CCl_4 /pet ether (charcoal).

3,3',5-Triiodo-S-thyronine [6893-02-3] M 651.0, m 236-237°(dec), $[\alpha]_{\text{D}}^{29.5} + 21.5^\circ$ (EtOH/1M aq HCl, 2:1), pK_1^{25} 6.48, pK_2^{25} 7.62, pK_3^{25} 7.82. Likely impurities are as in *thyroxine*. Purified by dissolving in dilute NH_3 at room temperature, then crystd by addition of dilute acetic acid to pH 6.

Trimellitic (benzene-1,2,4-tricarboxylic) acid [528-44-9] M 210.1, m 218-220°, pK_1^{25} 2.42, pK_2^{25} 3.71, pK_3^{25} 5.01. Crystd from acetic acid or aqueous EtOH.

1,2,3-Trimethoxybenzene [634-36-6] M 168.2, m 45-46°. Sublimed under vacuum.

1,3,5-Trimethoxybenzene [621-23-8] M 168.2, m 53°. Sublimed under vacuum.

Trimethylamine [75-50-3] M 59.1, b 3.5°, pK²⁵ 9.80. Dried by passage of the gas through a tower filled with solid KOH. Water and impurities containing labile hydrogen were removed by treatment with freshly sublimed, ground, P₂O₅. Has been refluxed with acetic anhydride, and then distd through a tube packed with HgO and BaO. [Comyns *J Chem Soc* 1557 1955.] For more extensive purification, trimethylamine has been converted to the hydrochloride, crystd (see below), and regenerated by treating the hydrochloride with excess aq 50% KOH, the gas passing through a CaSO₄ column into a steel cylinder containing sodium ribbon. After 1-2 days, the cylinder was cooled at -78° and hydrogen and air were removed by pumping. [Day and Felsing *J Am Chem Soc* 72 1698 1950.] Trimethylamine has also been trap-to-trap distd and then freeze-pump-thaw degassed [Halpern et al. *J Am Chem Soc* 108 3907 1986].

Trimethylamine hydrochloride [593-81-7] M 95.7, m >280°(dec). Crystd from CHCl₃, EtOH or *n*-propanol, and dried under vacuum. It has also been crystd from *benzene/MeOH, MeOH/diethyl ether and dried under vacuum over paraffin wax and H₂SO₄. Stood over P₂O₅. It is *hygroscopic*.

Trimethylamine hydroiodide [20230-89-1] M 186.0, m 263°. Crystd from MeOH.

1,2,4-Trimethylbenzene (pseudocumene) [95-63-6] M 120.2, m -43.8°, b 51.6°/10mm, 167-168°/760mm, d 0.889, n 1.5048. Refluxed over sodium and distd under reduced pressure.

2,4,6-Trimethylbenzoic acid (mesitoic acid) [480-63-7] M 164.2, m 155°, pK²⁵ 3.45. Crystd from water, ligroin or carbon tetrachloride [Ohwada et al. *J Am Chem Soc* 108 3029 1986].

Trimethyl-1,4-benzoquinone [935-92-2] M 150.1, m 29-30°, 36°, b 98°/10mm, 108°/18mm. Distd in a vac or sublimed *in vacuo* before use. [Smith et al. *J Am Chem Soc* 60 318 1939.]

R-(-)-2,2,6-Trimethyl-1,4-cyclohexanedione [60046-49-3] M 154.2, m 88-90°, 91-92°, [α]_D²⁰ -270° (c 0.4%, MeOH), [α]_D²⁰ -275° (c 1, CHCl₃). Obtained from fermentation and purified by recrystn from diisopropyl ether. [ORD: Leuenberger et al. *Helv Chim Acta* 59 1832 1976.] The *racemate* has m 65-67° and the 4-(4-phenyl)semicarbazone has m 218-220° (from CH₂Cl₂-MeOH) [Isler et al. *Helv Chim Acta* 39 2041 1956.]

2,2,5-Trimethylhexane [3522-94-9] M 128.3, m -105.8°, b 124.1°, d 0.716, n 1.39971, n²⁵ 1.39727. Extracted with conc H₂SO₄, washed with H₂O, dried (type 4A molecular sieves), and fractionally distd.

Trimethyl-1,4-hydroquinone (2,3,5-trimethylbenzene-1,4-diol) [700-13-0] M 152.2, m 173-174°, pK_{Est(1)}~ 11.1, pK_{Est(2)}~ 12.7. Recrystd from water, under anaerobic conditions.

1',3',3'-Trimethyl-6-nitrospiro[2H-benzopyran-2,2'-indoline] [1498-88-0] M 322.4, m 180°. Recrystd from absolute EtOH [Hinnen et al. *Bull Soc Chim Fr* 2066 1968; Ramesh and Labes *J Am Chem Soc* 109 3228 1987].

Trimethylolpropane [77-99-6] M 134.2, m 57-59°. Crystd from acetone and ether.

2,2,3-Trimethylpentane [564-02-3] M 114.2, b 109.8°, d 0.7161, n 1.40295, n²⁵ 1.40064. Purified by azeotropic distn with 2-methoxyethanol, which was subsequently washed out with water. The trimethylpentane was then dried and fractionally distd. [Forziati et al. *J Res Nat Bur Stand* 36 129 1946.]

2,2,4-Trimethylpentane (isooctane) [540-84-1] M 114.2, b 99.2°, d 0.693, n 1.39145, n²⁵ 1.38898. Distd from sodium, passed through a column of silica gel or activated alumina (to remove traces of

olefins), and again distd from sodium. Extracted repeatedly with conc H_2SO_4 , then agitated with aqueous KMnO_4 , washed with water, dried (CaSO_4) and distd. Purified by azeotropic distn with EtOH, which was subsequently washed out with water, and the trimethylpentane was dried and fractionally distd. [Forziati et al. *J Res Nat Bur Stand* **36** 126 1946.] Also purified by fractional crystn.

2,3,5-Trimethylphenol [697-82-5] M 136.2, m 95-96°, b 233°/760mm, pK^{25} 10.67. Crystd from water or pet ether.

2,4,5-Trimethylphenol [496-78-6] M 136.2, m 70.5-71.5°, pK^{25} 10.57. Crystd from water.

2,4,6-Trimethylphenol [527-60-6] M 136.2, m 69°, b 220°/760mm, pK^{25} 10.86. Crystd from water and sublimed *in vacuo*.

3,4,5-Trimethylphenol [527-54-8] M 136.2, m 107°, b 248-249°/760mm, pK^{25} 10.25. Crystd from pet ether.

Trimethylphenylammonium benzenesulfonate [16093-66-6] M 293.3. Crystd repeatedly from MeOH (charcoal).

2,2,4-Trimethyl-6-phenyl-1,2-dihydroquinoline [3562-69-4] M 249.3, m 102°. Vacuum distd, then crystd from absolute EtOH.

2,4,6-Trimethylpyridine (sym-collidine) [108-75-8] M 121.2, m -46°, b 10°/2.7mm, 36-37°/2mm, 60.7°/13mm, 65°/31mm, 170.4°/760mm, 175-178°/atm, d^{25} 0.9100, n_D^{20} 1.4939, 1.4981, n^{25} 1.4959, pK^{25} 6.69. Commercial samples may be grossly impure. Likely contaminants include 3,5-dimethylpyridine, 2,3,6-trimethylpyridine and water. Brown, Johnson and Podall [*J Am Chem Soc* **76** 5556 1954] fractionally distd 2,4,6-trimethylpyridine under reduced pressure through a 40-cm Vigreux column and added to 430mL of the distillate slowly, with cooling to 0°, 45g of BF_3 -diethyl etherate. The mixture was again distd, and an equal volume of dry *benzene was added to the distillate. Dry HCl was passed into the soln, which was kept cold in an ice-bath, and the hydrochloride was filtered off. It was recrystd from abs EtOH (1.5mL/g) to m 286-287°(sealed tube). The free base was regenerated by treatment with aq NaOH, then extracted with *benzene, dried (MgSO_4) and distd under reduced pressure. Sisler et al. [*J Am Chem Soc* **75** 446 1953] pted trimethylpyridine as its phosphate from a soln of the base in MeOH by adding 85% H_3PO_4 , shaking and cooling. The free base was regenerated as above. Garrett and Smythe [*J Chem Soc* 763 1903] purified the trimethylpyridine via the HgCl_2 complex. It is more soluble in cold than hot H_2O [sol 20.8% at 6°, 3.5% at 20°, 1.8% at 100°].

Also purified by dissolving in CHCl_3 , adding solid K_2CO_3 and Drierite, filtering and fractionally distilling through an 8in helix packed column. The *sulfate* has m 205°, and the *picrate* (from hot H_2O) has m 155-156°. [Frank and Meikle *J Am Chem Soc* **72** 4184 1950.]

Trimethylsulfonium iodide [2181-42-2] M 204.1, m 215-220°(dec). Crystd from EtOH.

1,3,7-Trimethyluric acid [5415-44-1] M 210.2, m 345°(dec), pK 6.0. Crystd from water.

1,3,9-Trimethyluric acid [7464-93-9] M 210.2, m 340°, 347°, pK^{20} 9.39. Crystd from water.

1,7,9-Trimethyluric acid [55441-82-2] M 210.2, m 316-318 (dec)°, 345°, pK_{Est} ~9.0. Crystd from water or EtOH, and sublimed *in vacuo*.

Trimyristin [555-45-3] M 723.2, m 56.5°. Crystd from diethyl ether.

2,4,6-Trinitroanisole [606-35-9] M 243.1, m 68°. Crystd from EtOH or MeOH. Dried under vac.

1,3,6-Trinitrobenzene [99-35-4] M 213.1, m 122-123°. Crystd from glacial acetic acid, CHCl₃, CCl₄, EtOH aq EtOH or EtOH/*benzene, after (optionally) heating with dil HNO₃. Air dried. Fused, and crystd under vacuum.

2,4,6-Trinitrobenzenesulfonic acid hydrate (TNBS, picrylsulfonic acid) [2508-19-2] M 293.2, m 180°, λ_{max} 240nm (ε 650 M⁻¹cm⁻¹), pK_{Est} ~ <0. It is also available as 0.1M and 5%w/v solns in H₂O. Recrystd from 1M HCl and dried at 100° or a mixt of EtOH (50mL), H₂O (30mL) and conc HCl (70mL) for 65g of acid. The *diethanolamine salt* had m 182-183° [Golumbic *J Org Chem* 11 518 1946].

2,4,6-Trinitrobenzoic acid [129-66-8] M 225.1, m 227-228°, pK²⁵ 0.65. Crystd from distilled water. Dried in a vacuum desiccator.

2,4,6-Trinitro-*m*-cresol [602-99-3] M 243.1, m 107.0-107.5°, pK 2.8. Crystd successively from H₂O, aq EtOH and *benzene/cyclohexane, then dried at 80° for 2h. [Davis and Paabo *J Res Nat Bur Stand* 64A 533 1960.]

2,4,7-Trinitro-9-fluorenone [129-79-3] M 315.2, m 176°. Crystd from nitric acid/water (3:1), washed with water and dried under vacuum over P₂O₅, or recrystd from dry *benzene.

2,4,6-Trinitrotoluene (TNT) [118-96-7] M 227.1, m 81.0-81.5°. Crystd from *benzene and EtOH. Then fused and allowed to cryst under vacuum. Gey, Dalbey and Van Dolah [*J Am Chem Soc* 78 1803 1956] dissolved TNT in acetone and added cold water (1:2:15), the ppte was filtered, washed free from solvent and stirred with five parts of aq 8% Na₂SO₃ at 50-60° for 10min. It was filtered, washed with cold water until the effluent was colourless, and air dried. The product was dissolved in five parts of hot CCl₄, washed with warm water until the washings were colourless and TNT was recovered by cooling and filtering. It was recrystd from 95% EtOH and carefully dried over H₂SO₄. The dry solid should not be heated without taking precautions for a possible **EXPLOSION**.

2,4,6-Trinitro-*m*-xylene [632-92-8] M 241.2, m 182.2°. Crystd from ethyl methyl ketone.

Tri-*n*-octylamine [1116-76-3] M 353.7, b 164-168°/0.7mm, 365-367°/760mm, d 0.813, n 1.450, pK²⁵ 10.65. It was converted to the amine hydrochloride etherate which was recrystd four times from diethyl ether at -30° (see below). Neutralisation of this salt regenerated the free amine. [Wilson and Wogman *J Phys Chem* 66 1552 1962.] Distd at 1-2mm pressure.

Tri-*n*-octylammonium chloride [1188-95-0] M 384.2, m 78-79°, pK 8.35 (in 70% aq EtOH). Crystd from Et₂O, then *n*-hexane (see above). [Burrows et al. *J Chem Soc* 200 1947.]

Tri-*n*-octylammonium perchlorate [2861-99-6] M 454.2, m >300°(dec). Crystd from *n*-hexane.

1,3,5-Trioxane [110-88-3] M 90.1, m 64°, b 114.5°/759mm. Crystd from sodium-dried diethyl ether or water, and dried over CaCl₂. Purified by zone refining.

Trioxsalen (2,5,9-trimethyl-7H-furo[3,2-*g*]benzopyran-7-one) [3902-71-4] M 228.3, m 233-235°, 234.5-235°. Purified by recrystn from CHCl₃. If too impure it is fractionally crystd from CHCl₃-pet ether (b 30-60°) using Norit and finally crystd from CHCl₃ alone to give colourless prisms, m 234.5-235°. It is a photosensitiser so it should be stored in the dark. [UV: Kaufmann *J Org Chem* 26 117 1961; Baeme et al. *J Chem Soc* 2976 1949.]

Tripalmitin [555-44-2] M 807.4, m 66.4°. Crystd from acetone, diethyl ether or EtOH.

Triphenylamine [603-34-9] M 245.3, m 127.3-127.9°, pK -5.0 (in fluorosulfuric acid). Crystd from EtOH or from *benzene/abs EtOH, diethyl ether and pet ether. It was sublimed under vacuum and carefully dried in a vacuum line. Stored in the dark under nitrogen.

1,3,5-Triphenylbenzene [612-71-5] M 306.4, m 173-175°. Purified by chromatography on alumina using *benzene or pet ether as eluents.

Triphenylene [217-59-4] M 228.3, m 198°, b 425°. Purified by zone refining or crystn from EtOH or CHCl₃, and sublimed.

1,2,3-Triphenylguanidine [101-01-9] M 287.3, m 144°, pK 9.10. Crystd from EtOH or EtOH/water, and dried under vacuum.

Triphenylmethane [519-73-3] M 244.3, m 92-93°. Crystd from EtOH or *benzene (with one molecule of *benzene of crystallisation which is lost on exposure to air or by heating on a water bath). It can also be sublimed under vacuum. It can also be given a preliminary purification by refluxing with tin and glacial acetic acid, then filtered hot through a glass sinter disc, and pptd by addition of cold water.

Triphenylmethanol (triphenylcarbinol) [76-84-6] M 260.3, m 164°, b 360-380° (without dec), pK²⁵ -6.63 (aq H₂SO₄). Crystd from EtOH, MeOH, CCl₄ (4mL/g), *benzene, hexane or pet ether (b 60-70°). Dried at 90°. [Ohwada et al. *J Am Chem Soc* 108 3029 1986.]

Triphenylmethyl chloride (trityl chloride) [76-83-5] M 278.9, m 111-112°. Crystd from iso-octane. Also crystd from 5 parts of pet ether (b 90-100°) and 1 part of acetyl chloride using 1.8g of solvent per g of chloride. Dried in a desiccator over soda lime and paraffin wax. [*Org Synth Coll Vol III* 841 1955; Thomas and Rochow *J Am Chem Soc* 79 1843 1957; Moisel et al. *J Am Chem Soc* 108 4706 1986.]

2,3,5-Triphenyltetrazolium chloride (TTC) [298-96-4] M 334.8, m 243°(dec). Crystd from EtOH or CHCl₃, and dried at 105°.

Tri-*n*-propylamine [102-69-2] M 143.3, b 156.5°, d 0.757, n 1.419, pK²⁵ 10.66. Dried with KOH and fractionally distd. Also refluxed with toluene-*p*-sulfonyl chloride and with KOH, then fractionally distd. The distillate, after addn of 2% phenyl isocyanate, was redistd and the residue fractionally distd from sodium. [Takahashi et al. *J Org Chem* 52 2666 1987.]

Tripyridyl triazine [3682-35-7] M 312.3, m 245-248°. Purified by repeated crystn from aq EtOH.

Tris-(2-aminoethyl)amine (TREN) [4097-89-6] M 146.2, b 114°/15mm, 263°/744mm, d 0.977, n 1.498, pK₁²⁵ 8.42, pK₂²⁵ 9.44, pK₃²⁵ 10.13. For a separation from a mixture containing 62% TRIEN, see entry under triethylenetetramine. Also purified by conversion to the hydrochloride (see below), recrystn and regeneration of the free base [Xie and Hendrickson *J Am Chem Soc* 109 6981 1987].

Tris-(2-aminoethyl)amine trihydrochloride [14350-52-8] M 255.7, m 300°(dec). Crystd several times by dissolving in a minimum of hot water and precipitating with excess cold EtOH. The ppte was washed with acetone, then diethyl ether and dried in a vacuum desiccator.

Tris(d,d-dicamopholylmethanato)europium (III) [52351-64-1] M 108.5, m 220-227.5°, 229-232°, [α]_D²⁵ +28.6° (c 5.4, CCl₄; and varies markedly with concentration). Dissolve in pentane, filter from any insol material, evaporate to dryness and dry the residue (white powder) at 100°/0.1mm for 36h. The IR has ν 1540cm⁻¹. [McCreary et al. *J Am Chem Soc* 96 1038 1974.]

Tris-(dimethylamino)methane (N,N,N',N',N'',N''-hexamethylmethanetriamine) [5762-56-1] M 145.3, b 42-43°/12mm, n 1.4349, pK_{Est} ~ 10. Dry over KOH and dist through a Vigreux column at waterpump vacuum. Store in absence of CO₂. [Bredereck et al. *Chem Ber* 101 1885 1968 and *Angew Chem, Int Ed Engl* 5 132 1966.]

Tris-(hydroxymethyl)methylamine (TRIS) [77-86-1] M 121.1, m 172°, pK²⁵ 8.07. Tris can ordinarily be obtained in highly pure form suitable for use as an acidimetric standard. If only impure material is available, it should be crystd from 20% EtOH. Dry in a vacuum desiccator over P₂O₅ or CaCl₂.

Alternatively, it is dissolved in twice its weight of water at 55-60°, filtered, concd to half its volume and poured slowly, with stirring, into about twice the volume of EtOH. The crystals which separate on cooling to 3-4° are filtered off, washed with a little MeOH, air dried by suction, then finally ground and dried in a vacuum desiccator over P₂O₅. It has also been crystd from water, MeOH or aq MeOH, and vacuum dried at 80° for 2 days.

Tris-(hydroxymethyl)methylamonium hydrochloride (TRIS-HCl) [1185-53-1] M 157.6, m 149-150°(dec). Crystd from 50% EtOH, then from 70% EtOH. Tris-hydrochloride is also available commercially in a highly pure state. Otherwise, crystd from 50% EtOH, then 70% EtOH, and dried below 40° to avoid risk of decomposition.

1,1,1-Tris-(hydroxymethyl)ethane (2-hydroxymethyl-2-methyl-1,3-propanediol) [77-85-0] M 120.2, m 200°. Dissolved in hot tetrahydrofuran, filtered and ppted with hexane. It has also been crystd from acetone/water (1:1). Dried in vacuum.

N-Tris-(hydroxymethyl)methyl-2-aminomethanesulfonic acid (TES) [7365-44-8] M 229.3, m 224-226°(dec), pK²⁰ 7.50. Crystd from hot EtOH containing a little water.

N-Tris-(hydroxymethyl)methylglycine (TRICINE) [5704-04-1] M 179.2, m 186-188°(dec), pK₁²⁰ ~2.3, pK₂²⁰ 8.15. Crystd from EtOH and water.

Tris-(hydroxymethyl)nitromethane [2-(hydroxymethyl)-2-nitro-1,3-propanediol] [126-11-4] M 151.1, m 174-175°(dec, tech grade), 214°(pure). Crystd from CHCl₃/ethyl acetate or ethyl acetate/*benzene. It is an acid and a 0.1M sol in H₂O has pH 4.5. **IRRITANT.**

Tris-[(3-trifluoromethylhydroxymethylene)-d-camphorato] europium (III) [Eu(tfc)₃] [34830-11-0] M 893.6, m 195-299° (dec), ~220°, [α]_D²⁴ +152° (c 2, CCl₄; and varies markedly with concentration). Purified by extraction with pentane, filtered and filtrate evapd and the residual bright yellow amorphous powder is dried at 100°/0.1mm for 36h. A sample purified by fractional molecular distn at 180-200°/0.004mm gave a liquid which solidified and softened at ~130° and melted at ~180° and was analytically pure. IR (CCl₄) v: 1630-1680cm⁻¹ and NMR (CCl₄) δ broad: -1.3 to 0.5, -0.08 (s), 0.41 (s), 1.6-2.3 and 3.39 (s). [McCreary et al. *J Am Chem Soc* 96 1038 1974; ; Goering et al. *J Am Chem Soc* 93 5913 1971.]

1,3,5-Trithiane [291-21-4] M 138.3, m 220°(dec). Crystd from acetic acid.

Triuret (1,3-dicarbamoylurea) [556-99-0] M 146.1, m 233°(dec). Crystd from aq ammonia. Gives mono and dipotassium salts.

Tropaeolin 00, [554-73-4] M 316.3, pK_{Est(2)} ~ 5.8, pK_{Est(3)} ~ 10.3. Recrystd twice from water [Kolthoff and Gus *J Am Chem Soc* 60 2516 1938].

Tropaeolin 000 (see Orange II p. 477 in Chapter 5). Purified by salting out from hot distilled water using sodium acetate, then three times from distilled water and twice from EtOH.

3-Tropanol (Tropine) [120-29-6] M 141.2, m 63°, b 229°/760mm, pK¹⁵ 3.80. Distd in steam and crystd from diethyl ether. *Hygroscopic.*

dl-Tropic (3-hydroxy-2-phenylpropionic) acid [529-64-6] M 166.2, m 118°, pK²⁵ 4.12. Crystd from water or *benzene.

Tropolone [533-75-5] M 122.1, m 49-50°, b 81-84°/0.1mm, pK₁ -0.53 (protonation of CO, aq H₂SO₄), pK₂ 6.67 (acidic OH). Crystd from hexane or pet ether and sublimed at 40°/4mm.

Tryptamine [(3-2-aminoethyl)indole] [61-54-1] M 160.1, m 116°, pK₁²⁵ -6.31 (aq H₂SO₄, diprotonation), pK_{Est(2)} ~4.9, pK₃²⁵ 16.60 (acidic indole NH). Crystd from *benzene.

Tryptamine hydrochloride [343-94-2] M 196.7, m 252-253°. Crystd from EtOH/water.

L-Tryptophan [73-22-3] M 204.3, m 278°, $[\alpha]_D^{20}$ -33.4° (EtOH), $[\alpha]_{546}^{20}$ -36° (c 1, H₂O), pK_1^{25} -6.23 (aq H₂SO₄), pK_2^{25} 2.46, pK_3^{25} 9.41, pK_4^{25} 14.82 (acidic NH, in aq NaOH). Crystd from water/EtOH, washed with anhydrous diethyl ether and dried at room temperature under vac over P₂O₅.

Tryptophol [3-(2-hydroxyethyl)indole] [526-55-6] M 161.2, m 59°, b 174°/2mm. Crystd from diethyl ether/pet ether, *C₆H₆, *C₆H₆/pet ether. The *picrate* has m 98-100° (from *C₆H₆).

(+)-Tubocurarine chloride (5H₂O) [57-94-3] M 771.7, m 274-275°(dec) (anhydrous), $[\alpha]_{546}^{20}$ +235° (c 0.5, H₂O), $pK_{Est(1)}$ ~8.5, $pK_{Est(2)}$ ~8.8. Crystd from water and forms various hydrates.

D(+)-Turannose [547-25-1] M 342.3, m 168-170°, $[\alpha]_D^{20}$ +88° (c 4, H₂O). Crystd from water by addition of EtOH.

Tyramine (4-hydroxybenzylamine) [51-67-2] M 137.2, m 164-165°, pK_1^{25} 9.74 (OH), pK_2^{25} 10.52 (NH₂). Crystd from *benzene or EtOH.

Tyramine hydrochloride [60-19-5] M 173.6, m 274-276°. Crystd from EtOH by addition of diethyl ether, or from conc HCl.

Tyrocidine A (cyclic decapeptide antibiotic with two D-Phe amino acids) [1481-70-5] M 1268.8, m 240°(dec), $[\alpha]_D^{25}$ -115° (c 0.91, MeOH). Crystd as hydrochloride from MeOH or EtOH and HCl. [Paladin and Craig *J Am Chem Soc* 76 688 1954; King and Craig *J Am Chem Soc* 77 6624 1955; Okamoto et al. *Bull Chem Soc Jpn* 50 231 1977.]

L-Tyrosine [60-18-4] M 181.2, m 290-295°(dec), $[\alpha]_D^{25}$ -10.0° (5M HCl), pK_1^{25} 2.18 (CO₂H), pK_2^{25} 9.21 (OH), pK_3^{20} 10.47 (NH₂). Likely impurities are L-cysteine and the ammonium salt. Dissolved in dilute ammonia, then crystd by adding dilute acetic acid to pH 5. Also crystd from water or EtOH/water, and dried at room temperature under vacuum over P₂O₅.

Umbelliferone (7-hydroxycoumarin) [93-35-6] M 162.2, m 225-228°, pK_{Est} ~8.0. Crystd from water.

Undecan-1-ol [112-42-5] M 172.3, m 16.5°. Purified by repeated fractional crystn from its melt or by distn in a vacuum.

Undec-10-enoic acid [112-38-9] M 184.3, m 25-25.5°, b 131°/1mm, 168°/15mm, pK_{Est} ~5.0. Purified by repeated fractional crystn from its melt or by distn in a vacuum.

Uracil [66-22-8] M 122.1, m 335°(dec), pK_1^{25} 9.43, pK_2^{25} 13.3-14.2. Crystd from water.

Uramil (5-aminobarbituric acid) [118-78-5] M 143.1, m 310-312°, 320°, >400°(dec), $pK_{Est(1)}$ ~3.9, $pK_{Est(2)}$ ~8.0, $pK_{Est(3)}$ ~12.5. Crystd from water.

Urea [57-13-6] M 60.1, m 132.7-132.9°, pK^{25} 0.12. Crystd twice from conductivity water using centrifugal drainage and keeping the temperature below 60°. The crystals were dried under vacuum at 55° for 6h. Levy and Margouls [*J Am Chem Soc* 84 1345 1962] prepared a 9M soln in conductivity water (keeping the temperature below 25°) and, after filtering through a medium-porosity glass sinter, added an equal volume of absolute EtOH. The mixture was set aside at -27° for 2-3 days and filtered cold. The ppt was washed with a small amount of EtOH and dried in air. Crystn from 70% EtOH between 40° and -9° has also been used. Ionic impurities such as ammonium isocyanate have been removed by treating the conc aqueous soln at 50° with

Tryptamine hydrochloride [343-94-2] M 196.7, m 252-253°. Crystd from EtOH/water.

L-Tryptophan [73-22-3] M 204.3, m 278°, $[\alpha]_D^{20}$ -33.4° (EtOH), $[\alpha]_{546}^{20}$ -36° (c 1, H₂O), pK_1^{25} -6.23 (aq H₂SO₄), pK_2^{25} 2.46, pK_3^{25} 9.41, pK_4^{25} 14.82 (acidic NH, in aq NaOH). Crystd from water/EtOH, washed with anhydrous diethyl ether and dried at room temperature under vac over P₂O₅.

Tryptophol [3-(2-hydroxyethyl)indole] [526-55-6] M 161.2, m 59°, b 174°/2mm. Crystd from diethyl ether/pet ether, *C₆H₆, *C₆H₆/pet ether. The *picrate* has m 98-100° (from *C₆H₆).

(+)-Tubocurarine chloride (5H₂O) [57-94-3] M 771.7, m 274-275°(dec) (anhydrous), $[\alpha]_{546}^{20}$ +235° (c 0.5, H₂O), $pK_{Est(1)}$ ~8.5, $pK_{Est(2)}$ ~8.8. Crystd from water and forms various hydrates.

D(+)-Turannose [547-25-1] M 342.3, m 168-170°, $[\alpha]_D^{20}$ +88° (c 4, H₂O). Crystd from water by addition of EtOH.

Tyramine (4-hydroxybenzylamine) [51-67-2] M 137.2, m 164-165°, pK_1^{25} 9.74 (OH), pK_2^{25} 10.52 (NH₂). Crystd from *benzene or EtOH.

Tyramine hydrochloride [60-19-5] M 173.6, m 274-276°. Crystd from EtOH by addition of diethyl ether, or from conc HCl.

Tyrocidine A (cyclic decapeptide antibiotic with two D-Phe amino acids) [1481-70-5] M 1268.8, m 240°(dec), $[\alpha]_D^{25}$ -115° (c 0.91, MeOH). Crystd as hydrochloride from MeOH or EtOH and HCl. [Paladin and Craig *J Am Chem Soc* 76 688 1954; King and Craig *J Am Chem Soc* 77 6624 1955; Okamoto et al. *Bull Chem Soc Jpn* 50 231 1977.]

L-Tyrosine [60-18-4] M 181.2, m 290-295°(dec), $[\alpha]_D^{25}$ -10.0° (5M HCl), pK_1^{25} 2.18 (CO₂H), pK_2^{25} 9.21 (OH), pK_3^{20} 10.47 (NH₂). Likely impurities are L-cysteine and the ammonium salt. Dissolved in dilute ammonia, then crystd by adding dilute acetic acid to pH 5. Also crystd from water or EtOH/water, and dried at room temperature under vacuum over P₂O₅.

Umbelliferone (7-hydroxycoumarin) [93-35-6] M 162.2, m 225-228°, pK_{Est} ~8.0. Crystd from water.

Undecan-1-ol [112-42-5] M 172.3, m 16.5°. Purified by repeated fractional crystn from its melt or by distn in a vacuum.

Undec-10-enoic acid [112-38-9] M 184.3, m 25-25.5°, b 131°/1mm, 168°/15mm, pK_{Est} ~5.0. Purified by repeated fractional crystn from its melt or by distn in a vacuum.

Uracil [66-22-8] M 122.1, m 335°(dec), pK_1^{25} 9.43, pK_2^{25} 13.3-14.2. Crystd from water.

Uramil (5-aminobarbituric acid) [118-78-5] M 143.1, m 310-312°, 320°, >400°(dec), $pK_{Est(1)}$ ~3.9, $pK_{Est(2)}$ ~8.0, $pK_{Est(3)}$ ~12.5. Crystd from water.

Urea [57-13-6] M 60.1, m 132.7-132.9°, pK^{25} 0.12. Crystd twice from conductivity water using centrifugal drainage and keeping the temperature below 60°. The crystals were dried under vacuum at 55° for 6h. Levy and Margouls [*J Am Chem Soc* 84 1345 1962] prepared a 9M soln in conductivity water (keeping the temperature below 25°) and, after filtering through a medium-porosity glass sinter, added an equal volume of absolute EtOH. The mixture was set aside at -27° for 2-3 days and filtered cold. The ppt was washed with a small amount of EtOH and dried in air. Crystn from 70% EtOH between 40° and -9° has also been used. Ionic impurities such as ammonium isocyanate have been removed by treating the conc aqueous soln at 50° with

Amberlite MB-1 cation- and anion-exchange resin, and allowing to crystallise. [Benesch, Lardy and Benesch *J Biol Chem* **216** 663 1955.] Also crystd from MeOH or EtOH, and dried under vacuum at room temperature.

Urea nitrate [124-47-0] M 123.1, m 152°(dec). Crystd from dilute HNO₃.

Uric acid [69-93-2] M 168.1, m >300°, pK₁ 5.75, pK₂ 10.3. Crystd from hot distilled water.

Uridine [58-96-8] M 244.2, m 165°, [α]_D²⁰ +4.0° (H₂O), pK²⁵ 9.51 (9.25). Crystd from aqueous 75% MeOH.

Urocanic acid (4-imidazolylacrylic acid) [104-98-3] M 138.1, m 225°, 226-228°, pK_{Est(1)}~2.5, pK_{Est(2)}~6, pK_{Est(3)}~11. Crystd from water and dried at 100°.

Ursodeoxycholic acid [128-13-2] M 392.5, m 203°, [α]_D²⁰ +60° (c 0.2, EtOH), pK_{Est} ~4.8. Crystd from EtOH.

(+)-Usnic acid [2,6-diacetyl-3,7,9-trihydroxy-8,9b-dimethyldibenzofuran-1(2H)-one] [7562-61-0] M 344.3, m 204°. See (+)-usnic acid on p. 573 in Chapter 6.

***trans*-Vaccenic acid** (octadec-11-enoic acid) [693-72-1] M 282.5, m 43-44°, pK_{Est} ~4.9. Crystd from acetone. The *methyl ester* has b 174-175°/5mm.

***n*-Valeraldehyde** [110-62-3] M 86.1, b 103°, d 0.811, n²⁵ 1.40233. Purified *via* the bisulfite derivative. [Birrell and Trotman-Dickinson *J Chem Soc* 2059 1960.]

***n*-Valeramide (pentanamide)** [626-97-1] M 101.1, m 115-116°. Crystd from EtOH.

Valeric acid (*n*-pentanoic acid) [109-52-4] M 102.1, b 95°/22mm, 186.4°/atm, d 0.938, n 1.4080, pK²⁵ 4.81. Water was removed from the acid by distn using a Vigreux column, until the boiling point reached 183°. A few crystals of KMnO₄ were added, and after refluxing, the distn was continued, [Andrews and Keefer *J Am Chem Soc* **83** 3708 1961.]

δ-Valerolactam (2-piperidone) [675-20-7] M 99.1, m 38.5-39.5°, 39-40°, 40°, b 81-82°/0.1mm, 136-137°/15mm, pK 0.75 (in AcOH). Purified by repeated fractional distn. [Cowley *J Org Chem* **23** 1330 1958; Reppe et al. *Justus Liebigs Ann Chem* **596** 198 1955; IR: Huisgen et al. *Chem Ber* **90** 1437 1957.] The *hydrochloride* has m 183-184° (from isoPrOH or EtOH-Et₂O) [Hurd et al. *J Org Chem* **17** 865 1952], and the *oxime* has m 122.5° (from pet ether) [Behringer and Meier *Justus Liebigs Ann Chem* **607** 67 1957]. *Picrate* has m 92-93°.

γ-Valerolactone (± 4,5-dihydro-5-methyl-2(3H)-furanone) [108-29-2] M 100.1, m -37°, 36°, b 82-85°/10mm, 102-103°/28mm, 125.3°/68mm, 136°/100mm, 205.75-206.25°/754mm, d₄²⁰ 1.072, n_D²⁰ 1.4322. Purified by repeated fractional distillation [Boorman and Linstead *J Chem Soc* 577, 580 1933]. IR v: 1790 (CS₂), 1775 (CHCl₃) cm⁻¹ [Jones et al. *Can J Chem* **37** 2007 1959]. The BF₃-complex distils at 110-111°/20mm [Reppe et al. *Justus Liebigs Ann Chem* **596** 179 1955]. It is characterised by conversion to γ-hydroxy-*n*-valeramide by treatment with NH₃, m 51.5-52° (by slow evapn of a CHCl₃ soln).

δ-Valerolactone (tetrahydro-2H-pyran-2-one) [542-28-9] M 100.1, m -13°, -12°, b 88°/4mm, 97°/10mm, 124°/24mm, 145-146°/40mm, 229-229.5°/atm, d₄²⁰ 1.1081, n_D²⁰ 1.4568. Purified by repeated fractional distn. IR v: 1750 (in CS₂), 1732 (in CHCl₃), 1748 (in CCl₄) and 1733 (in MeOH) cm⁻¹. [Huisgen and Ott *Tetrahedron* **6** 253 1959; Linstead and Rydon *J Chem Soc* 580 1933; Jones et al. *Can J Chem* **37** 2007 1959.]

Valeronitrile [110-59-8] M 83.1, b 142.3°, d 0.799, n¹⁵ 1.39913, n³⁰ 1.39037. Washed with half its volume of conc HCl (twice), then with saturated aqueous NaHCO₃, dried with MgSO₄ and fractionally distd from P₂O₅.

L-Valine [72-18-4] M 117.2, m 315°, [α]_D²⁰ +266.7° (6M HCl), pK₁ 2.26, pK₂ 9.68. Crystd from water by addition of EtOH.

Vanillin (4-hydroxy-3-methoxybenzaldehyde) [121-33-5] M 152.2, m 83°, b 170°/15mm, pK²⁵ 7.40. Crystd from water or aqueous EtOH, or by distn *in vacuo*.

Veratraldehyde [120-14-9] M 166.2, m 42-43°. Crystd from diethyl ether, pet ether, CCl₄ or toluene.

Variamine Blue RT [4-(phenylamino)benzenediazonium sulfate (1:1)] [4477-28-5] M 293.3, CI 37240, λ_{max} 377 nm. Dissolved 10g in 100mL of hot water. Sodium dithionite (0.4g) was added, followed by active carbon (1.5g) and filtered hot. To the colourless or slightly yellow filtrate a soln of saturated NaCl was added and the mixture cooled. The needles were filtered off, washed with cold water, dried at room temperature, and stored in a dark bottle (light sensitive). [Erdey *Chem Analyst* 48 106 1959.]

Vicine (2,4-diamino-5-β-D-glucopyranosidoxy-6-hydroxypyrimidine) [152-93-2] M 304.3, m 243-244°, [α]_D²⁰ -12° (c 4, 0.2N NaOH). Crystd from water or aqueous 85% EtOH, and dried at 135°.

Vinyl acetate [108-05-4] M 86.1, b 72.3°, d 0.938, n 1.396. Inhibitors such as hydroquinone, and other impurities are removed by drying with CaCl₂ and fractionally distilling under nitrogen, then refluxing briefly with a small amount of benzoyl peroxide and redistilling under nitrogen. Stored in the dark at 0°.

9-Vinylanthracene [2444-68-0] M 204.3, m 65-67°, b 61-66°/10mm. Purified by vacuum sublimation. Also by chromatography on silica gel with cyclohexane as eluent, and recrystd from EtOH [Werst et al. *J Am Chem Soc* 109 32 1987].

Vinyl butoxyethyl ether [4223-11-4] M 144.2. Washed with aqueous 1% NaOH, dried with CaH₂, then refluxed with and distd from, sodium.

N-Vinylcarbazole [1484-13-5] M 193.3, m 66°. Crystd repeatedly from MeOH in amber glassware. Vacuum sublimed.

Vinylene carbonate [872-36-6] M 86.1, m 22°. Purified by zone melting.

Vinyl chloroformate [5130-24-5] M 106.5, b 46.5°/80mm, 67-69°/atm, 109-110°/760mm, d₄²⁰ 1.136, n_D²³ 1.420. It has been fractionated through a Todd column (Model A with ~60 plates, see p. 174) under atmospheric pressure and purity can be checked by gas chromatography. It has IR with ν at 3100 + 2870 (CH₂), 1780 (C=O), 1640 (C=C) and 940 (CH₂ out-of-plane) and 910 (CH₂ wagging) cm⁻¹. [IR: Lee *J Org Chem* 30 3943 1965; Levaillant *Ann Chim (Paris)* 6 504 1936.] Used for protecting NH₂ groups in peptide synthesis [Olofson et al. *Tetrahedron Lett* 1563 1977].

1-Vinylnaphthalene [826-74-4] M 154.2, b 124-125°/15mm. Fractionally distd under reduced pressure on a spinning-band column, dried with CaH₂ and again distd under vacuum. Stored in sealed ampoules in a freezer.

2-Vinylpyridine monomer [100-69-6] M 105.1, b 79-82°/29mm, d 0.974, n 1.550, pK²⁵ 4.92. Steam distd, then dried with MgSO₄ and distd under vacuum.

4-Vinylpyridine monomer [100-43-6] M 105.1, b 40-41°/1.4mm, 54°/5mm, 58-61°/12mm, 68°/18mm, 79°/33mm, d₄²⁰ 0.9836, n_D²⁰ 1.5486, pK²⁵ 5.62. Purified by fractional distillation under a good vacuum and in a N₂ atmosphere and stored in sealed ampoules under N₂, and kept in the dark at -20°.

The *picrate* has **m** 175-176°. [UV: Coleman and Fuoss *J Am Chem Soc* **77** 5472 1955; Overberger et al. *J Polymer Sci* **27** 381 1958; Petro and Smyth *J Am Chem Soc* **79** 6142 1957.] Used for alkylating SH groups in peptides [Anderson and Friedman *Can J Biochem* **49** 1042 1971; Cawins and Friedman *Anal Biochem* **35** 489 1970].

Vinyl stearate [111-63-7] **M 310.5**, **m** 35°, **b** 166°/1.5mm. Vacuum distd under nitrogen, then crystd from acetone (3mL/g) or ethyl acetate at 0°.

Violanthrene (dibenzanthrene, 5,10-dihydroviolanthrene A) [81-31-2] **M 428.5**. Purified by vacuum sublimation over Cu in a muffle furnace at 450°/25mm in a CO₂ atmosphere [Scholl and Meyer *Chem Ber* **67** 1229 1934]. *Violanthrene A* (anthro[9,1,2-cde]benzo[rst]pentaphene [188-87-4]) **M 426.5** has **m** 506°. [Clar *Chem Ber* **76** 458 1943.]

Viologen (4,4'-dipyridyl dihydrochloride) [27926-72-3] **M 229.1**, **m** >300°. Purified by pptn on adding excess of acetone to a concentrated solution in aqueous MeOH. It has also been recrystd several times from MeOH and dried at 70° under vacuum for 24h [Prasad et al. *J Am Chem Soc* **108** 5135 1986], and recrystd three times from MeOH/isopropanol [Stramel and Thomas *J Chem Soc, Faraday Trans* **82** 799 1986].

Visnagin (4-methoxy-7-methyl-5H-furo[3,2-g][1]benzopyran-5-one) [82-57-5] **M 230.2**, **m** 142-145°. Crystd from water.

dl-Warfarin (4-hydroxy-3-(3-oxo-1-phenylbutyl)-2H-1-benzopyran-2-one) [81-81-2] **M 308.3**, **m** 161°. Crystd from MeOH. The *acetate* has **m** 182-183° and *2,4-dinitrophenylhydrazone* has **m** 215-216°. Effective anticoagulant and rodenticide.

Xanthatin (3-methylene-7-methyl-6-[3-oxo-1-buten-1-yl]cyclohept-5-ene-[10,11-b]furan-2-one) [26791-73-1] **M 246.3**, **m** 114.5-115°, [α]_D -20° (EtOH). Crystd from MeOH or EtOH. UV: λ_{\max} 213 and 275nm (ϵ 22800 and 7300).

Xanthene [92-83-1] **M 182.2**, **m** 100.5°, **b** 310-312°/760mm. Crystd from *benzene or EtOH.

9-Xanthenone (xanthone) [90-47-1] **M 196.2**, **m** 175.6-175.4°. Crystd from EtOH (25mL/g) and dried at 100°. It has also been recrystd from *n*-hexane three times and sublimed *in vacuo*. [Saltiel *J Am Chem Soc* **108** 2674 1986].

Xanthopterin (H₂O) see entry on p. 576 in Chapter 6.

Xanthorhamnin [1324-63-6] **M 770.7**, **m** 195°, [α]_D²⁰ +3.75° (EtOH). Crystd from a mixture of ethyl and isopropyl alcohols, air dried, then dried for several hours at 110°.

Xanthosine (2H₂O) [9-(β -D-ribose)purin-2,6(1H,3H)-dione] [5968-90-1] **M 320.3**, [α]_D²⁰ -53° (c 8, 0.3M NaOH), pK₁^{2.5} <2.5, pK₂^{2.5} 5.67, pK₃^{2.5} 12.85. Crystd from EtOH or water (as dihydrate).

Xanthurenic acid (5,8-dihydroxyquiniline-2-carboxylic acid) [59-00-7] **M 205.2**, **m** 286°, 290-295°(dec), pK_{Est(1)}~ 1.5, pK_{Est(2)}~ 4.9, pK_{Est(3)}~ 9.8. Ppted by the addition of 2N formic acid to its soln in hot 2M ammonia (charcoal). Filter solid off, dry in a vac at ~80° in the dark. UV (H₂O) has λ_{\max} nm (ϵ M⁻¹cm⁻¹): 243 (30,000) and 342 (6,500). The *methyl ester* has **m** 262° (from MeOH).

Xanthydrol [90-46-0] **M 198.2**, **m** 123-124°. Crystd from EtOH and dried at 40-50°.

Xylene [1330-20-7] M 106.1 (mixed isomers). Usual impurities are ethylbenzene, paraffins, traces of sulfur compounds and water. It is not practicable to separate the *m*-, and *p*-isomers of xylene by fractional distn, although, with a sufficiently efficient still, *o*-xylene can be fractionally distd from a mixture of isomers. Purified (and dried) by fractional distn from LiAlH₄, P₂O₅, CaH₂ or sodium. This treatment can be preceded by shaking successively with conc H₂SO₄, water, aqueous 10% NaOH, water and mercury, and drying with CaCl₂ for several days. Xylene can be purified by azeotropic distn with 2-ethoxyethanol or 2-methoxyethanol, the distillate being washed with water to remove the alcohol, then dried and fractionally distilled.

***o*-Xylene** [95-47-6] M 106.2, f -25.2°, b 84°/14mm, 144.4°/760mm, d²⁵ 0.88020, d²⁵ 0.87596, n 1.50543, n²⁵ 1.50292. The general purification methods listed under xylene are applicable [Clarke and Taylor *J Am Chem Soc* 45 831 1923]. *o*-Xylene (4.4Kg) is sulfonated by stirring for 4h with 2.5L of conc H₂SO₄ at 95°. After cooling, and separating the unsulfonated material, the product was diluted with 3L of water and neutralised with 40% NaOH. On cooling, sodium *o*-xylene sulfonate separated and was recrystd from half its weight of water. [A further crop of crystals was obtained by concentrating the mother liquor to one-third of its volume]. The salt was dissolved in the minimum amount of cold water, then mixed with the same amount of cold water, and with the same volume of conc H₂SO₄ and heated to 110°. *o*-Xylene was regenerated and steam distd. It was then dried and redistd.

***m*-Xylene** [108-38-3] M 106, f -47.9°, b 139.1°, d 0.86417, d²⁵ 0.85990, n 1.49721, n²⁵ 1.49464. The general purification methods listed under xylene are applicable. The *o*- and *p*-isomers can be removed by their selective oxidation when a *m*-xylene sample containing them is boiled with dilute HNO₃ (one part conc acid to three parts water). After washing with water and alkali, the product can be steam distd, then distd and purified by sulfonation. [Clarke and Taylor *J Am Chem Soc* 45 831 1923.] *m*-Xylene is selectively sulfonated when a mixture of xylenes is refluxed with the theoretical amount of 50-70% H₂SO₄ at 85-95° under reduced pressure. By using a still resembling a Dean and Stark apparatus, water in the condensate can be progressively withdrawn while the xylene is returned to the reaction vessel. Subsequently, after cooling, then adding water, unreacted xylenes are distd off under reduced pressure. The *m*-xylene sulfonic acid is subsequently hydrolysed by steam distn up to 140°, the free *m*-xylene being washed, dried with silica gel and again distd. Stored over molecular sieves Linde type 4A.

***p*-Xylene** [106-42-3] M 106.2, f 13.3, b 138.3°, d 0.86105, d²⁵ 0.85669, n 1.49581, n²⁵ 1.49325. The general purification methods listed for xylene are applicable. *p*-Xylene can readily be separated from its isomers by crystn from such solvents as MeOH, EtOH, isopropanol, acetone, butanone, toluene, pentane or pentene. It can be further purified by fractional crystn by partial freezing, and stored over sodium wire or molecular sieves Linde type 4A. [Stokes and French *J Chem Soc, Faraday Trans 1* 76 537 1980.]

Xylenol Orange {3*H*-2,1-benzoxathiol-3-ylidene-bis-[(6-hydroxy-5-methyl-*m*-phenylene)-methylnitro]tetraacetic acid, S,S-dioxide} [1611-35-4] M 672.6, m 210°(dec), ε₅₇₈ 6.09 x 10⁴ (pH 14), ε₄₃₅ 2.62 x 10⁴ (pH 3.1), pK₁ -1.74, pK₂ -1.09 (aq H₂SO₄-HNO₃), pK₃ 2.58, pK₄ 3.23, pK₅ 6.46, pK₆ 10.46, pK₇ 12.28. Generally contaminated with starting material (cresol red) and semixylenol orange. Purified by ion-exchange chromatography using DEAE-cellulose, eluting with 0.1M NaCl soln which will give the sodium salt. Cresol Red, semixylenol orange and iminodiacetic acid bands elute first. This procedure will give the sodium salt of the dye. To obtain the free acid dissolve the salt in H₂O and acidify with AcOH. Filter off, wash with H₂O and dry first in air and then in a vac desiccator over P₂O₅ in the dark [Sato, Yokoyama and Momoki *Anal Chim Acta* 94 317 1977].

α-D-Xylose [58-86-6] M 150.1, m 146-147°, [α]_D²⁰ -18.8° (c 4, H₂O). Purified by slow crystn from aq 80% EtOH or EtOH, then dried at 60° under vac over P₂O₅. Stored in a vacuum desiccator over CaSO₄.

***m*-Xylylene diisocyanate** [3634-83-1] M 188.2, b 88-89°/0.02mm, 130°/2mm, d₄²⁰ 1.204, n_D²⁰ 1.4531. Purified by repeated distn through a 2 plate column. [Ferstundig and Scherrer *J Am Chem Soc* 81 4838 1959.]

α -Yohimbine [146-48-5] M 354.5, m 278°(dec), $[\alpha]_D^{20} +55.6^\circ$ (c 2, EtOH), $pK_1^{22} 3.0$, $pK_2^{22} 7.45$. Crystd from EtOH, and dried to remove EtOH of crystn. For γ -Yohimbine see ajmalicine on p. 98.

Zeaxanthin [all *trans*- β -carotene-3,3'(*R,R'*)-diol] [144-68-3] M 568.9, m 207°, 215.5°, λ_{max} 275 (log ϵ 4.34), 453 (log ϵ 5.12), 480 (log ϵ 5.07) in EtOH. Yellow plates (with a blue lustre) from MeOH or EtOH.

CHAPTER 5

PURIFICATION OF INORGANIC AND METAL ORGANIC CHEMICALS

(Including Organic compounds of B, Bi, P, Se, Si, and ammonium and metal salts of organic acids)

The most common method of purification of inorganic species is by recrystallisation, usually from water. However, especially with salts of weak acids or of cations other than the alkaline and alkaline earth metals, care must be taken to minimise the effect of hydrolysis. This can be achieved, for example, by recrystallising acetates in the presence of dilute acetic acid. Nevertheless, there are many inorganic chemicals that are too insoluble or are hydrolysed by water so that no general purification method can be given. It is convenient that many inorganic substances have large temperature coefficients for their solubility in water, but in other cases recrystallisation is still possible by partial solvent evaporation.

Organo-metallic compounds, on the other hand, behave very much like organic compounds, e.g. they can be redistilled and may be soluble in organic solvents. A note of **caution** should be made about handling organo-metallic compounds, e.g. arsines, because of their **potential toxicities**, particularly when they are volatile. Generally the suppliers of such compounds provide details about their safe manipulation. These should be read carefully and adhered to closely. If in any doubt always assume that the materials are lethal and treat them with utmost care. The same **safety precautions** about the handling of substances as stated in Chapter 4 should be followed here (see Chapter 1).

For information on **ionization (pK)** see Chapter 1, p. 7, and Chapter 4, p. 80. In order to avoid repetition, the literature (or predicted) pK values of anionic and/or cationic species are usually reported at least once, and in several cases is entered for the free acid or free base, e.g. Na_2SO_4 will have a pK value for Na^+ at the entry for NaOH and the pK values for SO_4^{2-} at the entries for H_2SO_4 . When the pK values of the organic counter-ions are not given in this Chapter, as in case of sodium benzoate, the reader is referred to the value(s) in Chapter 4, e.g. of benzoic acid.

Abbreviations of titles of periodical are defined as in the Chemical Abstracts Service Source Index (CASSI). A note on other abbreviations is in Chapter 1, p. 30.

Benzene, which has been used as a solvent successfully and extensively in the past for reactions and purification by chromatography and crystallisation is now considered a **very dangerous substance** so it has to be used with extreme care. We emphasised that an alternative solvent to benzene (e.g. toluene, toluene-petroleum ether, or a petroleum ether to name a few) should be used first. However, if benzene has to be used then all operations have to be performed in a well ventilated fumehood and precautions taken to avoid inhalation and contact with skin and eyes. Whenever benzene is mentioned in the text and asterisk e.g. $^*\text{C}_6\text{H}_6$ or $^*\text{benzene}$, is inserted to remind the user that special precaution should be adopted.

Organic dyes which are *not* complexed or are salts of metals are included in Chapter 4 (use the CAS Registry Numbers to find them). Commercially available polymer supported reagents are indicated with § under the appropriate reagent.

Acetarsol see *N*-Acetyl-4-hydroxy-*m*-arsanilic acid.

Acetyltriphenylphosphonium chloride [1235-21-8] **M 354.8, m 237-238°, 244-246° (dec)**. Recrystd from $\text{CHCl}_3 + *C_6H_6 + \text{pet ether}$ (b 60-80°) and by dissolving in CHCl_3 and pouring it into dry Et_2O . $\lambda_{\text{max}}^{\text{EtOH}}$ nm(ϵ) 255(3,600), 262(3,700), 268(4,000) and 275(3,100). The *iodide salt* crystallises from H_2O and has **m 207-209°**. [*J Org Chem* 22 41 1957.] **IRRITANT and hygroscopic**. When shaken with a 10% aqueous soln of Na_2CO_3 (8h) it gives **acetylmethylene triphenyl phosphorane** which is recrystd from $\text{MeOH-H}_2\text{O}$ and after drying at 70°/0.1mm has **m 205-206°**. UV: λ_{max} nm(ϵ) 268 (6600), 275 (6500) and 288 (5700); IR: ν (cm^{-1}) 1529 (s), 1470 (m), 1425 (s), 1374 (m), 1105 (s) and 978 (s). [*J Org Chem* 22 41, 44 1957.]

3R,4R,1'R-4-Acetoxy-3-[1-(tert-butylmethylsilyloxy)ethyl]-2-azetinone [76855-69-1] **M 287.4, m 107-108°, $[\alpha]_{\text{D}}^{20} +55^\circ$ (c 0.5, toluene), $[\alpha]_{\text{D}}^{20} +53.7^\circ$ (c 1.04, CHCl_3)**. Purified by chromatography on silica gel (3 x 14cm) for 50g of ester using 20% EtOAc in *n*-hexane. The eluate is evaporated and the residue recrystd from hexane as white fluffy crystals. [*Tetrahedron* 39 2505 1983.]

Acetylenedicarboxylic acid monopotassium salt [928-04-1] **M 152.2**. Very soluble in H_2O , but can be crystd from small volume of H_2O in small crystals. These are washed with EtOH and dried over H_2SO_4 at 125°. [*Chem Ber* 10 841 1877; *Justus Liebigs Ann Chem* 272 133 1893.]

Acetylferrocene (ferrocenyl methylketone) [1271-55-2] **M 228.1, m 86°, 86-87°**. Orange-red crystals, recrystd from isooctane and sublimed at 100°/1mm. The *oxime* has **m 167-170°** (from Et_2O or aq EtOH). The *semicarbazone* has **m 198-201°** (from EtOH). [*J Am Chem Soc* 77 2022 3009 1955; *J Chem Soc* 650 1958.]

N-Acetyl-4-hydroxy-m-arsanilic acid [97-44-9] **M 275.1, pK₁ 3.73, pK₂ 7.9, pK₃ 9.3**. Crystd from water.

Allyl trimethylsilane (2-propenyltrimethylsilane) [762-72-1] **M 114.3, b 83.0-84.5°, 84-88°, 85.5-86.0°, d 0.713, n 1.405**. Fractionate through an efficient column at atm pressure. If impure dissolve in THF, shake with H_2O (2x), dry (Na_2SO_4) and fractionate. [Cudlin and Chvalovský *Collect Czech Chem Commun* 27 1658 1962.]

Allyl tri-n-butylstannane (allyl tributyl tin) [24850-33-7] **M 331.1, b 88-92°/0.2mm, 115°/17mm, d 1.068, n 1.487**. A possible impurity is tributylchlorostannane — test for Cl as Cl ion after hydrolysing. Dissolve in $*C_6H_6$ (or toluene), shake with dil aq NaOH , dry (CaCl_2), and dist in a vac [Jones et al. *J Chem Soc* 1446 1947; *Aldrichimica Acta* 17 75 1984 and 20 45 1987].

Alizarin Red S (3,4-dihydroxy-9,10-dioxo-2-anthracene sulfonic acid, Na salt. H_2O) [130-22-3] **M 360.4, pK₁^{2.5} <1, pK₂^{2.5} 5.49, pK₃^{2.5} 10.85 (11.01)**. Commercial samples contain large amounts of sodium and potassium chlorides and sulfates. It is purified by passing through a Sephadex G-10 column, followed by elution with water, then 50% aq EtOH [King and Pruden *Analyst (London)* 93 601 1968]. Finally dissolve in EtOH and ppt with Et_2O several times [*J Phys Chem* 54 829 1950; polarography *J Am Chem Soc* 70 3055 1948].

Alumina (neutral) [1344-28-1] **M 102.0 (anhyd.)**. Stirred with hot 2M HNO_3 , either on a steam bath for 12h (changing the acid every hour) or three times for 30min, then washed with hot distilled water until the washings had pH 4, followed by three washings with hot MeOH . The product was dried at 270° [Angyal and Young *J Am Chem Soc* 81 5251 1959]. For the preparation of alumina for chromatography see Chapter 1.

Aluminum acetylacetonate [13963-57-0] **M 324.3, m 192-194°, 195°**. Crystd several times from $*\text{benzene}$ or aqueous MeOH , λ_{max} 216 and 286nm. [*J Phys Chem* 62 440 1958.] It can be purified by sublimation and has the following solubilities in g per cent: $*C_6H_6$ 35.9 (20°), 47.6 (40°), toluene 15.9 (20°), 22.0 (40°) and acetylacetone 6.6 (20°), 10.4 (40°). [*Inorg Synth* 5 105 1957.]

Aluminium ammonium sulfate (10H₂O) [7784-26-1] M 453.3, m 93°, pK₁²⁵ 4.89, pK₂²⁵ 5.43, pK₃²⁵ 5.86 (fAl³⁺ aquo), pK₄²⁵ 11.22 [aluminate Al(OH)₄⁻]. Crystd from hot H₂O and cool in ice.

Aluminium bromide [7727-15-3] M 266.7, m 97°, b 114°/10mm. Refluxed and then distilled from pure aluminium chips in a stream of nitrogen into a flask containing more of the chips. It was then distd under vacuum into ampoules [Tipper and Walker *J Chem Soc* 1352 1959]. Anhydrous conditions are essential, and the white to very light brown solid distillate can be broken into lumps in a dry-box (under nitrogen). Fumes in moist air.

Aluminium caesium sulfate (12H₂O) [7784-17-0 (12H₂O); 14284-36-7] M 568.2. Crystd from hot water (3mL/g).

Aluminium chloride (anhydrous) [7446-70-0] M 133.3, m 192.6°. Sublimed several times in an all glass system under nitrogen at 30-50mm pressure. Has also been sublimed in a stream of dry HCl and has been subjected to a preliminary sublimation through a section of granular aluminium metal [for manipulative details see Jensen *J Am Chem Soc* 79 1226 1957]. Fumes in moist air.

Aluminum ethoxide [555-75-9] M 162.2, m 154-159°, 146-151°, b 187-190°/7mm, 210-214°/13mm. Crystd from CS₂ [m 139°, CS₂ complex] and distd in a vacuum. Molecular weight corresponds to [Al(OEt)₃]₄ [*J Phys Chem* 39 1127 1935; *J Am Chem Soc* 69 2605 1947].

Aluminium fluoride (anhydrous) [7784-18-4] M 84.0, m 250°. Technical material may contain up to 15% alumina, with minor impurities such as aluminium sulfate, cryolite, silica and iron oxide. Reagent grade AlF₃ (hydrated) contains only traces of impurities but its water content is very variable (may be up to 40%). It can be dried by calcining at 600-800° in a stream of dry air (some hydrolysis occurs), followed by vacuum distn at low pressure in a graphite system, heated to approximately 925° (condenser at 900°) [Henry and Dreisbach *J Am Chem Soc* 81 5274 1959].

Aluminium isopropoxide [555-31-7] M 204.3, m 119°, b 94°/0.5mm, 135°/10mm. Distd under vacuum. *Hygroscopic*.

Aluminium nitrate (9H₂O) [7784-27-2 (9H₂O); 13473-90-0] M 375.1. Crystd from dilute HNO₃, and dried by passing dry nitrogen through the crystals for several hours at 40°. After 2 recrystns of ACS grade it had S, Na and Fe at 2.2, 0.01 and 0.02 ppm resp.

Aluminium potassium sulfate (12H₂O, alum) [7784-24-9] M 474.4, m 92°. Crystd from weak aqueous H₂SO₄ (ca 0.5mL/g).

Aluminium rubidium sulfate (12H₂O) [7784-29-4] M 496.2. Crystd from aq H₂SO₄ (ca 2.5mL/g).

Aluminium sulfate (anhydrous) [10043-01-3] M 342.2, m 765°(dec); Al₂O₃ 14-18 H₂O [17927-65-0]; Al₂O₃ 18 H₂O [7784-31-8]. Crystd from hot dilute H₂SO₄ (1 mL/g) by cooling in ice. When a soln of alumina (Al₂O₃) in conc H₂SO₄ is slowly cooled, Al₂SO₄ 17 or 18H₂O deposits as a crystalline mass. Al₂SO₄ 17H₂O is the stable form in equilibrium with its saturated aqueous soln at 25° [Smith *J Am Chem Soc* 64 41 1942]. This is purified by dissolving in a small vol of H₂O and adding EtOH until the sulfate readily crystallises from the oily supersaturated soln. It forms Al₂O₃ 16H₂O between 0-112°. On gradual heating the hydrate melts giving the anhydrous salt at ca 250°. Several hydrates up to 27H₂O have been described. Further heating to red heat (~ 600-800) causes decomposition to Al₂O₃ + SO₃ + SO₂ and O₂ [Cobb *J Soc Chem Ind* 29 250 1910]. ACS reagent is Al₂O₃ 18H₂O (98+%).

Aluminum triethyl (triethyl aluminum) [97-93-8] M 114.2, b 69°/1.5mm, 76°/2.5mm, 129-131°/55mm, d₄²⁰ 0.695, n_D²⁰ 1.394. Purified by fractionation in an inert atmosphere under vacuum in a 50cm column containing a heated nichrome spiral, taking the fraction 112-114°/27mm. It is very sensitive to H₂O and should be stored under N₂. It should not contain chloride which can be shown by hydrolysis and testing with AgNO₃. [*J Am Chem Soc* 75 4828 5193 1953; NMR: *J Am Chem Soc* 81 3826 1959.]

Aluminium tri-*tert*-butoxide [556-91-2] M 246.3, m 208-210°(dec). Crystd from *benzene and sublimed at 180°.

Aluminium trimethanide (trimethyl aluminium) [75-24-1] M 72.1, m 15.2°, b 111.5°/488.2mm, 124.5°/atm, d_4^{20} 0.725. Distd through a 10-20 theoretical plates column under 1 atm of N₂ (better with very slow take-off). Attacks grease (use glass joints). Also vac distd over Al in absence of grease, into small glass vials and sealed under N₂. Purity is measured by freezing point. Reacts with H₂O, is non-conducting in *C₆H₆ and is **HIGHLY FLAMMABLE**. [*J Chem Soc* 4681946; *J Am Chem Soc* 68 2204 1946.]

4-Aminophenylmercuric acetate [6283-24-5] M 371.8, m 168°, 175°(dec), 180°(dec). Recrystd from hot dilute AcOH and dried in air. [*J Indian Chem Soc* 32 613 1955; *Justus Liebigs Ann Chem* 465 269 1928.]

Ammonia (gas) [7664-41-7] M 17.0, pK²⁵ 9.25. Major contaminants are water, oil and non-condensable gases. Most of these impurities are removed by passing the ammonia through a trap at -22° and condensing it at -176° under vacuum. Water is removed by distilling the ammonia into a tube containing a small lump of sodium. Also dried by passage through porous BaO, or over alumina followed by glass wool impregnated with sodium (prepared by soaking the glass wool in a solution of sodium in liquid ammonia, and evaporating off the ammonia). It can be rendered oxygen-free by passage through a soln of potassium in liquid ammonia.

Ammonia (liquid) [7664-41-7] M 17.0, m -77.7°, b -33.4°, n_D 1.325, d 0.597, d⁻⁷⁹ 0.817g/mL. Dried, and stored, with sodium in a steel cylinder, then distd and condensed by means of liquid air, the non-condensable gases being pumped off. In order to obtain liquid NH₃ from a cylinder turn the cylinder up-side-down (i.e. with the valve at the bottom, use a metal stand to secure it in this position) and lead a plastic tube from the tap to a measuring cylinder placed in an efficient fume cupboard which is kept running. Turn the tap on and allow the ammonia to be released. At first, gas and liquid will splatter out (make sure that the plastic tube is secure) but soon the liquid will drip into the measuring cylinder. The high latent heat of evaporation will cool the ammonia so that the liquid will remain cool and not boil vigorously. If the ammonia is required dry the necessary precautions should be taken, i.e. the gas is allowed to flow through tubes packed with coarse CaO pellets.

Ammonia (aqueous) [7664-41-7] M 17.0 + H₂O, d 0.90 (satd, 27% w/v, 14.3 N), pK²⁵ 9.25. Obtained metal-free by saturating distilled water, in a cooling bath, with ammonia (from tank) gas. Alternatively, can use isothermal distn by placing a dish of conc aq ammonia and a dish of pure water in an empty desiccator and leaving for several days. **AMMONIA (gas, liquid or aq soln) is very irritating and should not be inhaled in large volumes as it can lead to olfactory paralysis (temporary and partially permanent).**

Ammonium acetate [631-61-8] M 77.1, m 112-114°, d 1.04. Crystd twice from anhydrous acetic acid, dried under vacuum for 24h at 100° [Proll and Sutcliff *Trans Faraday Soc* 57 1078 1961].

Ammonium benzoate [1863-63-4] M 139.2, m 198°, 200°(dec), d 1.26. Crystd from EtOH.

Ammonium bisulfate (ammonium hydrogen sulfate) [7803-63-6] M 115.1°, m ~147°, d 1.79, pK²⁵ 1.96 (HSO₄⁻). Crystd from water at room temperature (1mL/g) by adding EtOH and cooling.

Ammonium bromide [12124-97-9] M 98.0, m 450°(sublimes), d 2.43. Crystd from 95% EtOH.

Ammonium chloride [12125-02-9] M 53.5, m 338°(sublime point, without melting), d 1.53. Crystd several times from conductivity water (1.5mL/g) between 90° and 0°. Sublimes. After one crystn, ACS grade had: metal(ppm) As (1.2), K (1), Sb (7.2), V (10.2).

Ammonium chromate [7788-98-9] M 152.1, m 185°(dec), d 1.81, pK₁²⁵ 0.74, pK₂²⁵ 6.49 (for H₂CrO₄). Crystd from weak aqueous ammonia (ca 2.5mL/g) by cooling from room temperature.

Ammonium dichromate [7789-09-5] M 252.1, m 170°(dec), d 1.26. Crystd from weak aq HCl (ca 1mL/g). (Possible carcinogen)

Ammonium dihydrogen arsenate [13462-93-6] M 159.0, m 300°(dec). Crystd from water (1mL/g).

Ammonium dihydrogen orthophosphate [7722-76-1] M 115.0, m 190°, d 1.80. Crystd from water (0.7mL/g) between 100° and 0°.

Ammonium dodecylsulfate [2235-54-3] M 283.4. Recrystd first from 90% EtOH and then twice from abs EtOH, finally dried in a vacuum.

Ammonium ferric oxalate (3H₂O) [13268-42-3] M 428.1, m ~160°(dec), d 1.77. Crystd from hot water (0.5mL/g).

Ammonium ferric sulfate (12H₂O) [7783-83-7 (12H₂O); 10138-04-2 (anhydr)] M 482.2, m ~37°, d 1.71. Crystd from aqueous ethanol.

Ammonium ferrous sulfate (6H₂O) [7783-85-9 (6H₂O); 10045-89-3 (anhydr)] M 392.1, m 100°(dec), d 1.86. A soln in warm water (1.5mL/g) was cooled rapidly to 0°, and the resulting fine crystals were filtered at the pump, washed with cold distilled water and pressed between sheets of filter paper to dry.

Ammonium formate [540-69-2] M 63.1, m 116°, 117.3°, d₄⁴⁵ 1.280. Heat solid in NH₃ vapour and dry in vacuum till NH₃ odour is faint. Recryst from abs EtOH and then keep in a desiccator over 99% H₂SO₄ *in vacuo*. It is very *hygroscopic*. Exists in two forms, stable needles and less stable plates. Also forms acid salts, i.e. HCO₂NH₄.3HCO₂H and HCO₂NH₄.HCO₂H. [*J Am Chem Soc* 43 1473 1921; 63 3124 1941 .]

Ammonium hexachloroiridate (IV) [16940-92-4] M 441.0. Ppted several times from aqueous soln by saturation with ammonium chloride. This removes any palladium and rhodium. Then washed with ice-cold water and dried over conc H₂SO₄ in a vacuum desiccator. If osmium or ruthenium is present, it can be removed as the tetroxide by heating with conc HNO₃, followed by conc HClO₄, until most of the acid has been driven off. (This treatment is repeated). The near-dry residue is dissolved in a small amount of water and added to excess NaHCO₃ soln and bromine water. On boiling, iridic (but not platonic) hydroxide is ppted. It is dissolved in HCl and ppted several times, then dissolved in HBr and treated with HNO₃ and HCl to convert the bromides to chlorides. Saturation with ammonium chloride and cooling precipitates ammonium hexachloroiridate which is filtered off and purified as above [Woo and Yost *J Am Chem Soc* 53 884 1931].

Ammonium hexacyanoferrate II hydrate [14481-29-9] M 284.1, m dec on heating. The pale yellow trihydrate powder can be washed with 10% aq NH₃, fildt, then washed several times with EtOH and Et₂O, and dried at room temp. Decomposes in vacuum above 100° and should be stored away from light and under N₂. In light and air it decomposes by losing NH₃. [*Handbook of Preparative Inorganic Chem (Ed. Brauer)* Vol II 1509 1965.]

Ammonium hexafluorophosphate [16941-11-0] M 163.0, d₄¹⁸ 2.181, pK₁²⁵~ 0.5, pK₂²⁵ 5.12 (for fluorophosphoric acid H₂PO₃F). Crystallises from H₂O in square plates. Decomposes on heating before melting. Soluble in H₂O at 20° (74.8% w/v), also very soluble in Me₂CO, MeOH, EtOH and MeOAc and is decomposed by boiling acids. [*Chem Ber* 63 1063 1930.]

Ammonium hexafluorosilicate [16919-19-0] M 178.1, pK₂ 1.92 (for H₂SiF₆). Crystd from water (2mL/g). After 3 recrystns of Tech grade it had Li, Na, K and Fe at 0.3, 0.2, 0.1 and 1.0 ppm resp.

Ammonium hypophosphite [7803-65-8] M 83.0. Crystd from hot EtOH.

Ammonium iodate [13446-09-8] **M 192.9, pK²⁵ 0.79 (IO³⁺)**. Crystd from water (8mL/g) between 100° and 0°.

Ammonium iodide [12027-06-4] **M 144.9, sublimes with dec ~405°, d 2.51**. Crystd from EtOH by addition of ethyl iodide. *Very hygroscopic*. Stored in the dark.

Ammonium ionophore I (Nonactin) [6833-86-7] **M 736.9, m 147-148°, [α]_D²⁰ 0° (c 1.2, CHCl₃)**. Crystd from MeOH in colourless needles and is dried at 20° in high vac. A selectophore with high sensitivity for NH₄⁺ ions. [*Helv Chim Acta* **38** 1445 1955, **45** 129 1962, **55** 1371 1972; *Acta Cryst* **27B** 1680 1971.]

Ammonium magnesium chloride (6H₂O) [60314-43-4] **M 256.8**. Crystd from water (6mL/g) by partial evapn in a desiccator over KOH (deliquescent).

Ammonium magnesium sulfate (6H₂O) [20861-69-2] **M 360.6**. Crystd from water (1mL/g) between 100° and 0°.

Ammonium manganous sulfate (6H₂O) [13566-22-8] **M 391.3**. Crystd from water (2mL/g) by partial evapn in a desiccator.

Ammonium metavanadate [7803-55-6] **M 117.0, m 200°(dec)**. See ammonium (meta) vanadate on p. 395.

Ammonium molybdate [13106-76-8] **M 196.0, pK₁²⁵ 0.9 (proton addition), pK₂²⁵ 3.57, pK₃²⁵ 4.08 (for H₂MoO₄)**. Crystd from water (2.5mL/g) by partial evapn in a desiccator.

Ammonium nickel sulfate (6H₂O) [7785-20-8 (6H₂O); 15699-18-0 (anhydr)] **M 395.0, d 1.923**. Crystd from water (3mL/g) between 90° and 0°.

Ammonium nitrate [6484-52-2] **M 80.0, m 210°(dec explosively), d 1.72**. Crystd twice from distilled water (1mL/g) by adding EtOH, or from warm water (0.5mL/g) by cooling in an ice-salt bath. Dried in air, then under vacuum. After 3 recrystns of ACS grade it contained Li and B at 0.03 and 0.74 ppm resp.

Ammonium oxalate (H₂O) [6009-70-7] **M 142.1, d 1.50**. Crystd from water (10mL/g) between 50° and 0°.

Ammonium perchlorate [7790-98-9] **M 117.5, d 1.95, pK²⁵ -2.4 to -3.1 (for HClO₄)**. Crystd twice from distilled water (2.5mL/g) between 80° and 0°, and dried in a vacuum desiccator over P₂O₅. Drying at 110° might lead to slow decomposition to chloride. **POTENTIALLY EXPLOSIVE**.

Ammonium peroxydisulfate [7727-54-0] **M 228.2, m dec when heated wet liberating oxygen, d 1.98**. Recrystd at room temperature from EtOH/water.

Ammonium picrate [131-74-8] **M 246.1, m EXPLODES above 200°**. Crystd from EtOH and acetone.

Ammonium reineckate (Reineckate salt) [13573-16-5] **M 345.5, m 270-273°(dec)**. Crystd from water, between 30° and 0°, working by artificial light. Solns of reineckate decompose slowly at room temperature in the dark and more rapidly at higher temperatures or in diffuse sunlight.

Ammonium selenate [7783-21-3] **M 179.0, d 2.19, m dec on heating**. Crystd from water at room temperature by adding EtOH and cooling.

Ammonium sulfamate [7773-06-0] **M 114.1, m 132-135°, dec at 160°**. Crystd from water at room temperature (1mL/g) by adding EtOH and cooling.

Ammonium sulfate [7783-20-2] **M 132.1, m 230°(dec), 280°(dec), d 1.77.** Crystd twice from hot water containing 0.2% EDTA to remove metal ions, then finally from distilled water. Dried in a desiccator for 2 weeks over $\text{Mg}(\text{ClO}_4)_2$. After 3 recrystns ACS grade had Ti, K, Fe, Na at 11, 4.4, 4.4, 3.2 ppm resp.

Ammonium tetrafluoroborate [13826-83-0] **M 104.8, pK^{25} 2.77 (for HBF_4).** Crystd from conductivity water (1mL/g) between 100° and 0°.

Ammonium tetraphenylborate [14637-34-4] **M 337.3, m ca 220°(dec).** Dissolve in aqueous Me_2CO and allow crystn to proceed slowly otherwise very small crystals are formed. No trace of Me_2CO was left after drying at 120° [*Trans Faraday Soc* 53 19 1957]. The salt was pptd from dilute AcOH soln of sodium tetraphenylborane in the presence of NH_4^+ ions. After standing for 5min, the ppte was filtered off onto a sintered porcelain crucible, washed with very dilute AcOH and dried at room temp for at least 24h [*Anal Chem* 28 1001 1956]. Alternatively a soln of sodium tetraphenylborane (5% excess) in H_2O is added to NH_4Cl soln. After 5min the ppte is collected, washed several times with H_2O and recryst from aqueous Me_2CO . [*Analyt Chim Acta* 19 342 1958.]

Ammonium thiocyanate [1762-95-4] **M 76.1, m 138°(dec), 149°(dec), pK^{25} -1.85 (for HSCN), 149.** Crystd three times from dilute HClO_4 , to give material optically transparent at wavelengths longer than 270nm. Has also been crystd from absolute MeOH and from acetonitrile.

Ammonium tungstate (VI) [11120-25-5] **M 283.9, pK_1^{25} 2.20, pK_2^{25} 3.70 (for tungstic acid, H_2WO_4).** Crystd from warm water by adding EtOH and cooling.

Ammonium (meta) vanadate [7803-55-6] **M 117.0, d_{10}^{20} 2.326.** Wash with H_2O until free from Cl⁻ and dry in air. It is soluble in H_2O (5.18g/100mL at 15°, 10.4g/100mL at 32°) but is more soluble in dilute NH_3 . Crystd from conductivity water (20mL/g). When heated at relatively low temperatures it loses H_2O and NH_3 to give vanadium oxide (V_2O_5) and at 210° it forms lower oxides. [*Inorg Synth* 3 117 1950.] After washing Tech grade with H_2O it had Na, Mn and U at 0.06, 0.2 and 0.1 ppm resp.

n-Amylmercuric chloride [544-15-0] **M 307.2, m 110°.** Crystd from EtOH.

Anthraquinone Blue B (Acid Blue 45, 1,5-diamino-4,8-dihydroxy-9,10-anthraquinone-3,7-disulfonic acid di-Na salt) [2861-02-1] **M 474.3, m >300°, CI 63010, λ_{max} 595nm, $\text{pK}_{\text{Est}(1)} \sim <0$, $\text{pK}_{\text{Est}(2)} \sim 2$, $\text{pK}_{\text{Est}(3)} \sim 9$.** Purified by salting out three times with sodium acetate, followed by repeated extraction with EtOH [McGrew and Schneider *J Am Chem Soc* 72 2547 1950].

Anthraquinone Blue RXO [4403-89-8] **M 445.5.** Purified by salting out three times with sodium acetate, followed by repeated extraction with EtOH [McGrew and Schneider *J Am Chem Soc* 72 2547 1950].

Anthraquinone Green G [Acid Green 25, Alizarin Cyanine Green F, 1,4-bis-(4-methyl-2-sulfophenyl-1-amino)-9,10-anthraquinone di-Na salt] [4403-90-1] **M 624.6, m 235-238°, CI 61570, λ_{max} 642nm, $\text{pK} >0$.** Purified by salting out three times with sodium acetate, followed by repeated extraction with EtOH [McGrew and Schneider *J Am Chem Soc* 72 2547 1950]. It is a green powder that slightly sol in Me_2CO , EtOH and pyridine. Sol in conc H_2SO_4 to give a blue soln which becomes turquoise on dilution. [Allen et al. *J Org Chem* 7 63 1942.]

9,10-Anthraquinone-2,6-disulfonic acid (disodium salt) [853-68-9] **M 412.3, m >325°, $\text{pK}_{\text{Est}} \sim <0$ (for SO_3H).** Crystd three times from water, in the dark [Moore et al. *J Chem Soc. Faraday Transl* 82 745 1986].

9,10-Anthraquinone-2-sulfonic acid (Na salt, H_2O) [131-08-8] **M 328.3, $\text{pK}_{\text{Est}} \sim <0$ (SO_3H).** Crystd from H_2O or MeOH (charcoal). [Costa and Bookfield *J Chem Soc, Faraday Trans 1* 82 991 1986].

Antimony (V) pentafluoride [7783-70-2] **M 216.7, m 7.0°, 8.3°, b 141°, 150°, 148-150°, d 2.99, pK^{25} 2.55 [for $\text{HSb}(\text{OH})_6 = \text{Sb}(\text{OH})_6^- + \text{H}^+$].** Purified by vacuum distillation preferably in a

quartz apparatus, and stored in quartz or aluminum bottles. It is a *hygroscopic* viscous liquid which reacts *violently* with H₂O and is hydrolysed by alkalis. *It is POISONOUS and attacks the skin.* [*J Chem Soc* 2200 1950; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 200 1965.]

Antimony trichloride [10025-91-9] M 228.1, m 73°, b 283°, pK₁²⁵ 1.4, pK₂²⁵ 11.0 (11.8), pK₃²⁵ 12.95 (for Sb³⁺ aquo). Dried over P₂O₅ or by mixing with toluene or xylene and distilling (water is carried off with the organic solvent), then distd twice under dry nitrogen at 50mm, degassed and sublimed twice in a vacuum into ampoules. Can be crystd from CS₂. Deliquescent. Fumes in moist air.

Antimony trifluoride [7783-56-4] M 178.8, m 292°. Crystd from MeOH to remove oxide and oxyfluoride, then sublimed under vacuum in an aluminium cup on to a water-cooled copper condenser [*Woolf J Chem Soc* 279 1955].

Antimony triiodide [7790-44-5] M 502.5, m 167°. Sublimed under vacuum.

Antimony trioxide [1309-64-4] M 291.5, m 656°. Dissolved in minimum volume of dilute HCl, filtered, and six volumes of water were added to ppte a basic antimonous chloride (free from Fe and Sb₂O₅). The ppte was redissolved in dilute HCl, and added slowly, with stirring, to a boiling soln (containing a slight excess) of Na₂CO₃. The oxide was filtered off, washed with hot water, then boiled and filtered, the process being repeated until the filtrate gave no test for chloride ions. The product was dried in a vacuum desiccator [*Schuhmann J Am Chem Soc* 46 52 1924]. After on cryst(pptn?), the oxide from a Chinese source had: metal (ppm) Al (8), Ag (0.2), As (56), Cr (6), Ge (0.4), Mn (0.2), Na (16), Ni (2.2) Pb (2.4), Sn (0.4) and V (32).

Aqua regia. This is prepared by adding slowly concentrated HNO₃ (1 vol) to concentrated hydrochloric acid (3 vols) in a glass container. This mixture is used to dissolve metals, including noble metals and alloys, as well as minerals and refractory substances. It is done by suspending the material and boiling (**EFFICIENT FUME CUPBOARD — EYE PROTECTION**) to dryness and repeating the process until the residue dissolves in H₂O. If the aqua regia is to be stored for long periods it is advisable to dilute it with one volume of H₂O which will prevent it from releasing chlorine and other chloro and nitrous compounds which are objectionable. Store cool in a fume cupboard. However, it is good laboratory practice to prepare it freshly and dispose of it down the fume cupboard sink with copious amounts of water.

Argon [7440-37-1] M 39.95, b -185.6°. Rendered oxygen-free by passage over reduced copper at 450°, or by bubbling through alkaline pyrogallol and H₂SO₄, then dried with CaSO₄, Mg(ClO₄)₂, or Linde 5A molecular sieves. Other purification steps include passage through Ascarite (asbestos impregnated with sodium hydroxide), through finely divided uranium at about 800° and through a -78° cold trap. Alternatively the gas is passed over CuO pellets at 300° to remove hydrogen and hydrocarbons, over Ca chips at 600° to remove oxygen and, finally, over titanium chips at 700° to remove nitrogen. Also purified by freeze-pump-thaw cycles and by passage over sputtered sodium [*Arnold and Smith J Chem Soc, Faraday Trans 2* 77 861 1981].

***o*-Arsanilic acid** [2045-00-3] M 216.1, m 153°, pK₁²² 3.77 (AsO₃H₂), pK₂²² 8.66 (AsO₃H⁻). Crystd from water or ethanol/ether. **POISONOUS.**

***p*-Arsanilic acid** [98-50-0] M 216.1, m 232°, pK₁²² 4.05 (AsO₃H₂), pK₂²² 8.66 (AsO₃H⁻). Crystd from water or ethanol/ether. **POISONOUS.**

Arsenazo I [3(2-*arsonophenylazo*)-4,5-dihydroxy-2,7-naphthalenedisulfonic acid di Na salt] [66019-20-3] M 614.3, ε 2.6 x 10⁴ at 500nm, pH 8.0; pK₁ 0.6(0.8), pK₂ 3.52, pK₃ 2.97(AsO₃H₂), pK₄ 8.20(AsO₃H⁻), pK₅ 9.98(OH), pK₆ 15.0. A saturated aqueous soln of the free acid was slowly added to an equal volume of conc HCl. The orange ppte was filtered, washed with acetonitrile and dried for 1-2h at 110° [*Fritz and Bradford Anal Chem* 30 1021 1958].

Arsenazo III [3,6-bis(2-*arsonophenylazo*)-4,5-dihydroxy-2,7-naphthalenedisulfonic acid di Na salt] [62337-00-2] M 776.4, pK₁ -2.7, pK₂ -2.7, pK₃ 0.6, pK₄ 0.8, pK₅ 1.6, pK₆ 3.4,

pK₇ 6.27, pK₈ 9.05, pK₉ 11.98, pK₁₀ 15.1. Contaminants include monoazo derivatives, starting materials for synthesis and by-products. Partially purified by pptn of the dye from aqueous alkali by addition of HCl. More thorough purification by taking a 2g sample in 15-25mL of 5% aq NH₃ and filter. Add 10mL HCl (1:1) to the filtrate to ppt the dye. Repeat procedure and dissolve solid dye (0.5g) in 7mL of a 1:1:1 mixture of *n*-propanol:conc NH₃:water at 50°. After cooling, filter soln and treat the filtrate on a cellulose column using 3:1:1 mixture of *n*-propanol:conc NH₃:water as eluent. Collect the blue band and evaporate to 10-15mL below 80°, then add 10mL conc HCl to ppt pure Arsenazo III. Wash with EtOH and air-dry [Borak et al. *Talanta* 17 215 1970]. The purity of the dye can be checked by paper chromatography using M HCl as eluent.

Arsenic [7440-38-2] **M 74.9, m 816°.** Heated under vacuum at 350° to sublime oxides, then sealed in a Pyrex tube under vacuum and sublimed at 600°, the arsenic condensing in the cooler parts of the tube. Stored under vacuum [Shih and Peretti *J Am Chem Soc* 75 608 1953]. **POISONOUS.**

Arsenic acid (arsenic pentoxide hydrate, arsenic V oxide hydrate, orthoarsenic acid) [12044-50-7] **M 229.8 + xH₂O, pK₁²⁵ 2.26, pK₂²⁵ 6.76, pK₃²⁵ 11.29 (H₃AsO₄).** Cryst from conc solns of boiling conc HNO₃ as rhombic crystals. Dried in vac to give hemihydrate (hygroscopic). Heating above 300° yields As₂O₅. [Thaler *Z Anorg Allg Chem* 246 19 1941.] **POISONOUS.**

Arsenic tribromide [7784-33-0] **M 314.6, m 31.1°, b 89°/11mm, 221°/760mm.** Distd under vacuum. **POISONOUS.**

Arsenic trichloride (butter of arsenic) [7784-34-1] **M 181.3, b 25°/11mm, 130.0°.** Refluxed with arsenic for 4h, then fractionally distd. The middle fraction was stored with sodium wire for two days, then again distd [Lewis and Sowerby *J Chem Soc* 336 1957]. Fumes in moist air and readily hydrolysed by H₂O. **POISONOUS.**

Arsenic triiodide [7784-45-4] **M 455.6, m 146°, b 400°/atm.** Crystd from acetone, sublimes below 100°. **POISONOUS**

Arsenic III oxide (arsenic trioxide, arsenious oxide) [1327-53-3] **M 197.8, three forms: m ~200°(amorphous glass), m 275°(sealed tube, octahedral, common form, sublimes > 125° without fusion but melts under pressure), m ~312°, pK₁²⁰ 9.27, pK₂²⁰ 13.54, pK₃²⁰ 13.99 (for H₃AsO₃).** Crystd in octahedral form from H₂O or from dil HCl (1:2), washed, dried and sublimed (193°/760mm). Analytical reagent grade material is suitable for use as an analytical standard after it has been dried by heating at 105° for 1-2h or has been left in a desiccator for several hours over conc H₂SO₄. **POISONOUS (particulary the vapour, handle in a ventilated fume cupboard).**

Aurothioglucose (gold thioglucose) [12192-57-3] **M 392.2.** Purified by dissolving in H₂O (0.05g in 1mL) and ppting by adding EtOH. Yellow cryst with slight mercaptan odour. Decomposes slowly in H₂O, sol in propylene glycol but insol in EtOH and other common organic solvents. [*FEBS Lett* 98 351 1970.]

Barium (metal) [7440-39-3] **M 137.3, m 727°.** Cleaned by washing with diethyl ether to remove adhering paraffin, then filed in an argon-filled glove box, washed first with ethanol containing 2% conc HCl, then with dry ethanol. Dried under vacuum and stored under argon [Addison, Coldrey and Halstead *J Chem Soc* 3868 1962]. Has also been purified by double distn under 10mm argon pressure.

Barium acetate [543-80-6] **M 255.4.** Crystd twice from anhydrous acetic acid and dried under vacuum for 24h at 100°.

Barium bromate [13967-90-3] **M 265.3.** Crystd from hot water (20mL/g). The monohydrate melts at 260°(dec).

Barium bromide (2H₂O) [7791-28-8] M 333.2, m at 75° loses first H₂O and at 120° loses the second H₂O. Crystd from water (1mL/g) by partial evaporation in a desiccator.

Barium chlorate (H₂O) [10294-38-9 (hydrate); 13477-00-4 (anhydr)] M 322.3, m 414°. Crystd from water (1mL/g) between 100° and 0°.

Barium chloride (2H₂O) [10326-27-9] M 244.3, m ~120°(dec, hydrate), 963° (anhyd). Twice crystd from water (2mL/g) and oven dried to constant weight.

Barium dithionate (2H₂O) [13845-17-5] M 333.5, m >150° loses SO₂, pK²⁵ 0.49 (for H₂S₂O₆, theory pK₁ -3.4, pK₂ -0.2). Crystd from water.

Barium ferrocyanide (6H₂O) [13821-06-2] M 594.8, m 80°(dec), pK₃²⁵ 2.57, pK₄²⁵ 4.35 (for ferrocyanide). Crystd from hot water (100mL/g).

Barium fluoride [7787-32-8] M 175.3, m 1353°, 1368°, b 2260°, d 4.83. Washed well with distd H₂O and dried in vacuum Sol in H₂O [1.6g (10°), 1.6g (20°) and 1.62g (30°) per L], mineral acids and aq NH₄Cl. May be stored in glass bottles. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer) Vol I 234 1963.*]

Barium formate [541-43-5] M 277.4, pK²⁵ 3.74 (for HCO₂H). Crystd from warm water (4mL/g) by adding EtOH and cooling.

Barium hydroxide (8H₂O) [12230-71-6] M 315.5, m 78°, pK₁²⁵ 13.13, pK₂²⁵ 13.36. Crystd from water (1mL/g).

Barium hypophosphite (H₂O) [14871-79-5] M 285.4. Pptd from aq soln (3mL/g) by adding EtOH.

Barium iodate (H₂O) [7787-34-0] M 487.1, m 130°(loses H₂O), 476°(dec). Crystd from a large volume of hot water by cooling.

Barium iodide (2H₂O) [7787-33-9 (2H₂O); 13718-50-8 (anhydr)] M 427.2, m 740°(dec). Crystd from water (0.5mL/g) by partial evapn in a desiccator. **POISONOUS.**

Barium ionophore I [*N,N,N',N'*-tetracyclohexyloxy-bis-(*o*-phenyleneoxy)diacetamide] [96476-01-6] M 644.9, m 156-158°. Purified by chromatography on a Kieselgel column and eluted with CH₂Cl₂-EtOAc (5:1), and recryst from EtOH-Me₂CO as colourless crystals. It is an electrically neutral ionophore with high selectivity for Ba²⁺ ions and with high lipophilicity. [*Chem Ber 118 1071 1985.*]

Barium manganate (barium permanganate) [7787-35-1] M 256.3, d 3.77. Wash with conductivity H₂O by decantation until the supernatant gives a faint test for Ba²⁺. Remove excess H₂O in vac (IMPORTANT), then heat at 100° and the last traces of H₂O are removed in a vac desiccator over P₂O₅. Store over KOH. It disproportionates in hot H₂O or dil acid to Ba(MnO₂)₂ and MnO₂, and is a mild oxidant. [*J Am Chem Soc 44 1965 1924; Inorg Synth 11 56 1960.*]

Barium nitrate [10022-31-8] M 261.4, m 593°(dec). Crystd twice from water (4mL/g) and dried overnight at 110°. **POISONOUS.**

Barium nitrite (H₂O) [7787-38-4] M 247.4, m 217°(dec). Crystd from water (1mL/g) by cooling in an ice-salt bath. **POISONOUS.**

Barium perchlorate [13465-95-7] M 336.2, m 505°, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd twice from water.

Barium propionate (H₂O) [5908-77-0] M 301.5, pK²⁴ 4.88 (for propionic acid). Crystd from warm water (50mL/g) by adding EtOH and cooling.

Barium sulfate [7722-43-7] M 233.4, m >1580°. Washed five times by decantation with hot distilled water, dialysed against distd water for one week, then freeze-dried and oven dried at 105° for 12h.

Barium tetrathionate [82203-66-5] M 361.6. Purified by dissolution in a small volume of water and ppted with EtOH below 5°. After drying the salt was stored in the dark at 0°.

Barium thiocyanate (2 H₂O) [2092-17-3] M 289.6, pK²⁵ -1.85 (for HSCN). Crystd from water (2.5mL/g) by partial evaporation in a desiccator.

Barium thiosulfate [35112-53-9] M 249.5, m 220°(dec), pK₁²⁵ 0.6, pK₂²⁵ 1.74 (for H₂S₂O₃). Very slightly soluble in water. Washed repeatedly with chilled water and dried in air at 40°.

Benzaldehyde-2-sulfonic acid sodium salt [1008-72-6] M 208.2, m dec on heating. Forms prisms or plates by extracting with boiling EtOH, filtering, evaporate to dryness and recrystallise the Na salt from a small volume of H₂O. The *N*-phenylhydrazone sodium salt recrystd from H₂O, m 174.5°. [Gnehm and Schüle *Justus Liebigs Ann Chem* 299 363 1898.]

Benzenechromium tricarbonyl [12082-08-5] M 214.1, m 163-166°. Purified by sublimation *in vacuo*.

Benzeneselenenyl bromide (phenylselenenyl bromide) [34837-55-3] M 236.0, m 58-62°, 60°, 62°, b 107-108°/15mm, 134°/35mm. Dist in a vac, recryst from pet ether, CHCl₃ (EtOH free), or Et₂O (cooling mixture) as dark red or orange crystals, sublimes at 25°/0.001mm [Behaghel et al. *Chem Ber* 65 815 1932; Pitteloud and Petrzilka *Helv Chim Acta* 62 1319 1979]. **HIGHLY TOXIC**

Benzeneselenenyl chloride (benzeneselenenyl chloride, phenylselenenyl chloride) [5707-04-0] M 191.5, m 59-60°, 64-65°, b 92°/5mm, 120°/20mm. Purified by distn in vac and recrystn (orange needles) from hexane [Foster *J Am Chem Soc* 55 822 1933, Foster et al. *Recl Trav Chim, Pays-Bas* 53 405, 408 1934; Behaghel and Seibert *Chem Ber* 66 714 1933.] **HIGHLY TOXIC.**

Benzeneseleninic acid [6996-92-5] M 189.1, m 122-124°, pK²⁵ 4.70. Add 10% excess of 15M NH₃ to the solid acid and stir until the solid dissolves, filter, decolorise with charcoal (2x, Norite) and acidify by slow addn of 6M HCl, filter solid and wash with H₂O. Dissolve the acid in the minimum vol of MeOH and this soln is added dropwise to boiling H₂O until cloudiness appears. At this point add 25% more boiling H₂O, filter hot (decolorise if necessary) and cool rapidly with scratching to 0°. After 30min the solid is filt'd off and recryst as before but with very slow cooling. The colorless needles are filtered off and dried in a vac desiccator (CaCl₂) before the melting point is measured [McCullough and Gould *J Am Chem Soc* 71 674 1949].

Benzeneseleninic anhydride [17697-12-0] M 360.1, m 124-126°, 164-165°, 170-173°. When the anhydride is recrystd from *C₆H₆ it has m 124-126° but when this is heated at 140°/1h in a vac or at 90°/2h it had m 164-165° and gives a solid m 124-126° when recrystd from *C₆H₆. Both depress the melting point of the acid PhSeO₂H. If the high melting anhydride is dissolved in *C₆H₆ and seeded with the high melting anhydride, the high melting anhydride crystallises out. It readily absorbs H₂O to form the acid (PhSeO₂H, m 122-124°). Because of this the commercial anhydride could contain up to 30% of the acid. Best purified by converting to the HNO₃ complex (m 112°) and heating this *in vacuo* at 120°/72h to give the anhydride as a white powder m 164-165°. Alternatively heat the anhydride *in vacuo* at 120°/72 h until IR shows no OH band. [Ayvrey et al. *J Chem Soc* 2089 1962; Barton et al. *J Chem Soc, Perkin Trans 1* 567 1977.] **TOXIC solid.**

Benzeneselenol (phenylselenol, selenophenol) [645-96-5] M 157.1, b 57-59°/8mm, 71-72°/18mm, 84-86°/25mm, d 1.480, n 1.616. Dissolve in aq N NaOH, acidify with conc HCl and extract with Et₂O, dry over CaCl₂, filter, evap on a steam bath and distil from a Claisen flask or through a short

column collecting the middle fraction and seal immediately in a glass vial, otherwise the colourless liquid becomes yellow. The alkali insol materials consist of diphenylselenide (**b** 167°/16mm) and diphenyldiselenide, **m** 63° (from EtOH). **TOXIC**, use rubber gloves. It has a foul odour. [Foster *Org Synth Coll Vol III* 771 1955.]

Benzeneselenonic acid (benzeneselenoic acid) [39254-48-3] **M 205.1**, **m 64°**, **pK²⁵ 4.79**. Purified by dissolving in H₂O and passing through a strong cation exchange resin (H⁺ form). Evap under reduced pressure and dry in a high vac to give colourless hygroscopic crystals [Dostal et al. *Z Chem* 6 153 1966 and IR: *Chem Ber* 104 2044 1971].

Benzenestibonic acid [535-46-6] **M 248.9**, **m >250°(dec)**. Crystd from acetic acid, or from EtOH-CHCl₃ mixture by addition of water.

Benzenesulfinic acid Na salt [873-55-2] **M 164.2**, **m >300°**, **pK²⁵ 2.16 (2.74; for PhSO₂H)**. Dissolve in the minimum vol of O₂ free H₂O (prepared by bubbling N₂ through for 2 h) and adding O₂ free EtOH (prepared as for H₂O), set aside at 4° overnight under N₂, filt'd, washed with abs EtOH, then Et₂O and dried in a vac. The Na salt is relatively stable to air oxidation, but is best kept under N₂ in the dark. Also recrystd from EtOH and dried at 120° for 4h in a vacuum [Kornblum and Wade *J Org Chem* 52 5301 1987].

Benzopurpurin 4B {3,3'-[(3,3'-dimethyl[1,1'-biphenyl]-4,4'-diyl)bis(azo)]bis[4-amino-1-naphthalenesulfonic acid di-Na salt, Direct red 2} [992-59-6] **M 724.7**, **λ_{max} 500nm**, **CI 23500**, **pK<0**. Crystd from H₂O. It is a biological stain that is violet at pH 1.2 and red at pH 4.0 and is used for detecting Al, Mg, Hg, Au and U.

Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent) [56602-33-6] **M 442.29**, **m > 130° (dec)**, **147-149° (dec)**. Dissolve in CH₂Cl₂, dry (MgSO₄), filter, conc under vac then add dry Et₂O and filter off first crop. Add CH₂Cl₂ to the filtrate and concentrate again to give a second crop. Solid is washed with dry Et₂O and dried in vac. Also recryst from dry Me₂CO/Et₂O and check purity by NMR. Store in the dark. [Castro et al. *Synthesis* 751 1976.]

Benzylidene-bis-(tricyclohexylphosphine) dichlororuthenium (Grubbs catalyst) [172222-30-9] **M 823.0**. Repeatedly wash with Me₂CO and MeOH and dry in vac. Alternatively dissolve in CH₂Cl₂ concentrate to half vol, filter, add MeOH to ppt it as purple microcrystals. Filter off, wash several times with Me₂CO and MeOH and dry in a vac for several hours. [Scwab, Grubbs and Ziller *J Am Chem Soc* 118 100 1996; Miller, Blackwell and Grubbs *J Am Chem Soc* 118 9606 1966.]

§ A polymer supported version is available.

Benzyl Orange [4-(4-benzylaminophenylazo)benzenesulfonic acid potassium salt] [589-02-6] **M 405.5**, **pK_{Est(1)}~<0**, **pK_{Est(2)}~3.8**. Crystd from H₂O.

Benzyltriphenylphosphonium chloride [1100-88-5] **M 388.9**, **m 280° (sintering)**, **287-288°**. Wash with Et₂O and crystallise from EtOH (six sided plates). *Hygroscopic* and forms crystals with one mol H₂O. [Justus Liebigs *Ann Chem* 229 320 1885; *Chem Ber* 83 291 1950.]

Beryllium acetate (basic) [Be₄O(OAc)₆] [1332-52-1] **M 406.3**, **m 285-286°**. Crystd from chloroform.

Beryllium potassium fluoride [7787-50-0] **M 105.1**. Crystd from hot water (25mL/g).

Beryllium sulfate (4H₂O) [7787-56-6] **M 177.1**, **m ~100°(dec)**, **pK₁²⁵ 3.2**, **pK₂²⁵ ~6.5 (Be²⁺)**. Crystd from weak aqueous H₂SO₄.

Bicyclo[2.2.1]hepta-2,5-diene rhodium (I) chloride dimer (norbornadiene rhodium chloride complex dimer) [12257-42-0] **M 462**, **m 240°(dec)**. Recrystd from hot CHCl₃-pet ether as fine crystals soluble in CHCl₃ and *C₆H₆ but almost insoluble in Et₂O or pet ether. [*J Chem Soc* 3178 1959.]

R-(-)-1,1'-Binaphthyl-2,2'-diylhydrogen phosphate [39648-67-4] M 348.3, m 217°, $[\alpha]_D^{20}$ -608° (c 1, MeOH), pK²⁰ 0.74. Recrystallise from EtOH. Reflux for 3h in N NaOH is required to hydrolyse the cyclic phosphate. [*Tetrahedron Lett* 4617 1971 ; *Tetrahedron Lett* 24, 343 1983.]

S-(+)-1,1'-Binaphthyl-2,2'-diylhydrogen phosphate [35193-64-7] M 348.3, $[\alpha]_D^{20}$ +608° (c 1, MeOH), pK²⁰ 0.74. Recrystallise from EtOH. Reflux for 3h in N NaOH is required to hydrolyse the cyclic phosphate. [*Tetrahedron Lett* 4617 1971 ; *Tetrahedron Lett* 24, 343 1983.]

2-Biphenyl diphenyl phosphate [132-29-6] M 302.4, n²⁵ 1.5925. Vacuum distd, then percolated through an alumina column. Passed through a packed column maintained at 150° to remove residual traces of volatile materials by a counter-current stream of nitrogen at reduced pressure. [Dobry and Keller *J Phys Chem* 61 1448 1957.]

2,2'-Bipyridinium chlorochromate [76899-34-8] M 292.6. Washed with cold conc HCl then H₂O (sintered glass funnel) and dried in vacuum (CaCl₂) to a free flowing yellow-brown powder. Stored in the dark. [*Synthesis* 691 1980; *Synth Commun* 10 951 1980.] **SUSPECTED CARCINOGEN.**

2,2'-Biquinolin-4,4'-dicarboxylic acid dipotassium salt [63451-34-3] M 420.51. Recryst from H₂O. The Cu salt has λ_{\max} at 562nm. [*Anal Biochem* 56 4409 1973.]

Bis-(p-tert-butylphenyl)phenyl phosphate [115-87-7] M 438.5, b 281°/5mm, n²⁵ 1.5412. Same as for 2-biphenyl diphenyl phosphate (above).

Bis-(2-chlorophenyl) phenyl phosphate [597-80-8] M 395, b 254°/4mm, n²⁵ 1.5767. Same as for 2-biphenyl diphenyl phosphate above.

Bis-(1,5-cyclooctadiene)nickel (0) [1295-35-8] M 275.0, m 142° (dec). Available in sealed ampoules under N₂. All procedures should be carried out in a dry box and in an atmosphere of N₂ or Argon in subdued light because the complex is light and oxygen sensitive and flammable. The solid is washed with dry Et₂O (under Ar) and separates from toluene as yellow crystals. Filter under Ar gas pressure, place the crystals in a container and dry under a vac of 0.01 mm to remove adhered toluene, flush with Ar and seal under Ar or N₂ in glass ampoules. [Semmelhack *Org Reactions* 19 115 and 178 1972; Wilke et al. *Justus Liebigs Ann Chem* 699 1 1966.] **SUSPECTED CARCINOGEN.**

Bis(2,9-dimethyl-1,10-phenanthroline) copper(I) perchlorate [54816-44-5] M 579.6, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from acetone.

2,2'-Bis-(diphenylphosphino)-1,1'-binaphthyl (BINAP) [RS 98327-87-8] M 622.7, m 283-286°, [R-(+)-76189-55-4] m 241-242°, [S-(-)-76189-56-5] m 241-242°, $[\alpha]_D^{20}$ (+) and (-) 233° (c 0.3 toluene). Dissolve the individual enantiomers in toluene, wash with 30% aq NaOH, three times with H₂O, dry (Na₂SO₄), evap to ~15% of its vol and add an equal vol of degassed MeOH. Collect the solid, wash with MeOH and dry at 80°/0.005mm for 6h. Recryst from 1:1 mixt of toluene-EtOH to optical purity (m 241-242°)[Takaya et al. *Org Synth* 67 20 1989].

§ A polymer supported version is available.

1,4-Bis-(diphenylphosphino)butane [7688-25-7] M 426.5, m 135-136°. Recrystd from EtOH [Trippett *J Chem Soc* 4263 1961].

2R,3R-(+)-2,3-Bis(diphenylphosphino)butane (R,R-CHIRAPHOS) [74839-84-2], **2S,3S-(-)-2,3-bis(di-phenylphosphino)butane (S,S-CHIRAPHOS)** [64896-28-2] M 426.5, m 108-109°, $[\alpha]_D^{20}$ (+) and (-) 200° (c 1.5 CHCl₃). Recrystd from abs EtOH (~6g in 60mL) as colorless plates [Fryzuk and Bosnich *J Am Chem Soc* 99 6262 1977 and 101 3043 1979].

1,2-Bis-(diphenylphosphino)ethane (DIPHOS) [1663-45-2] **M 398.4, m 139-140°, 140-142°, 143-144°, $pK_{Est} \sim 4.5$** . Recrystd from aq EtOH or *C_6H_6 . The *dimethiodide* recrystd from MeOH has **m 305-307°** and the *dioxide* recrystd from toluene or DMF (needles), or *C_6H_6 (plates) has **m 252-254° (276-278°)** [Isslieb et al. *Chem Ber* **92** 3175 1959; NMR: Aquiar et al. *J Org Chem* **29** 1660 1964; Bäckvall et al. *J Org Chem* **52** 5430 1987].

1,1'-Bis-(diphenylphosphino)ferrocene [12150-46-8] **M 554.4, m 181-183°, 184-194°**. Wash with distilled H_2O and dry in a vacuum. Dissolve in *ca* 5 parts of hot dioxane and cool to give orange crystals **m 181-183°**. Recrystn from *C_6H_6 -heptane (1:2) gives product with **m 183-184°**. [*J Organomet Chem* **27** 241 1971.]

Bis-(2-ethylhexyl) 2-ethylhexyl phosphonate [25103-23-5] **M 434.6, $n^{25} 1.4473$** . Purified by stirring an 0.4M soln in * benzene with an equal volume of 6M HCl at *ca* 60° for 8h. The * benzene layer was then shaken successively with equal volumes of water (twice), aqueous 5% Na_2CO_3 (three times), and water (eight times), followed by evaporation of the * benzene and distilled under reduced pressure at room temperature (using a rotating evacuated flask). Stored in dry, dark conditions [Peppard et al. *J Inorg Nucl Chem* **24** 1387 1962]. Vacuum distilled, then percolated through an alumina column before finally passed through a packed column maintained at 150° where residual traces of volatile materials were removed by a counter-current stream of N_2 at reduced pressure [Dobry and Keller *J Phys Chem* **61** 1448 1957].

Bis-(2-ethylhexyl) phosphoric acid [298-07-7] **M 322.4**. See di-(2-ethylhexyl) phosphoric acid on p. 418.

Bis(ethyl)titanium(IV) chloride [2247-00-9] **M 177.0**. Crystd from boiling toluene.

Bis(ethyl)zirconium(IV) chloride [92212-70-9] **M 220.3**. Crystd from boiling toluene.

2,4-Bis-(methylthio)-1,3,2 λ^5 ,4 λ^5 -dithiadiphosphetane-2,4-dithione (Davy's reagent) [82737-61-9] **M 284.4, m 160°**. Recrystd from *C_6H_6 in yellow plates or from hot trichlorobenzene. The low **m** observed in the literature (112° with gradual softening at 68-102°) has been attributed to the presence of elemental sulfur in the crystals. [*Tetrahedron* **40** 2663 1984; *J Org Chem* **22** 789 1957.]

Bismuth [7440-69-9] **M 209.0, m 271-273°**. Melted in an atmosphere of dry helium and filtered through dry Pyrex wool to remove any bismuth oxide present [Mayer, Yosim and Topol *J Phys Chem* **64** 238 1960].

Bismuthiol I (2,5-dimercapto-1,3,4-thiadiazole) potassium salt [4628-94-8] **M 226.4, m 275-276°(dec), $pK_{Est(1)} \sim 4.1$** . Usually contaminated with disulfide. Purified by crystn from EtOH. Reagent for detection of Bi, Cu, Pb and Sb.

Bismuth trichloride [7787-60-2] **M 315.3, m 233.6°, $pK^{25} 1.58 (Bi_3^+ = BiOH_2^+ + H^+)$** . Sublimed under high vacuum, or dried under a current of HCl gas, followed by fractional distn, once under HCl and once under argon.

***N,N'*-Bis-(salicylidene)ethylenediamine cobalt (II) [Co(SALEN)₂, salcomine]** [14167-18-1] **M 325.2**. The powder should have an oxygen capacity of 4.7-4.8% as measured by the increase in wt under O_2 at 100 pounds pressure at *ca* 20°. The O_2 is expelled on heating the material to 65°. Recryst from pyridine, $CHCl_3$ or *C_6H_6 , and the solvent may be removed by heating at 120° in a vac. However this heating may mean reduced O_2 capacity. In the dry state it absorbs O_2 turning from maroon colour to black. [Diehl and Hack *Inorg Synth* **3** 196 1950.]

Bis-(tetrabutylammonium) dichromate [56660-19-6] **M 700.9, m 139-142°**. Wash with water and dry in a vacuum. Crystallises from hexane (**m 79-80°**). [*Synth Commun* **10** 75 1980.] (Possible CARCINOGEN).

Bis-[4-(1,1,3,3-tetramethylbutyl)phenyl]phosphate calcium salt (Selectophore) [40835-97-0] **M 987.3**. The Ca diester salt is washed with H₂O (x3) and MeOH (x3) alternately and dried in a vacuum oven at 50°. If the Ca salt is contaminated with much Ca salt of the monoester then it (10g) is converted to the free acid by adding 6N HCl (*ca* 10vols) and Et₂O (> 50vols) to it and stirred vigorously to form the free acids. When no white ppte remained (*ca* 5min), the Et₂O is separated, washed with H₂O (2 x > 50 mL) and dried by filtering through a bed of anhydrous Na₂SO₄ (11 x 5 cm) which is then washed with Et₂O (2 x > 50 mL). Evapn gives an oil (TLC R_F 0.81 for diester and 0.50 for monoester). The oil is dissolved in *benzene (*ca* 25mL) and extracted with ethane-1,2-diol (25mL, 10x). After ten washings, a small sample of the *benzene layer is washed twice with H₂O to remove the diol and showed that it is pure bis-[4-(1,1,3,3-tetramethylbutyl)phenyl]phosphoric acid by TLC, i.e. no monophosphate. To form the Ca salt the oil is dissolved in MeOH and to it is added the equivalent amount of CaCl₂ together with aq NaOH to keep the pH >10. The resulting white ppte is collected washed alternately with 3 batches of H₂O and MeOH and dried in a vacuum oven at 50°. [*J Inorg Nucl Chem* **40** 1483 1978.]

2,4-Bis-(*p*-tolylthio)-1,3,2λ⁵,4λ⁵-dithiadiphosphetane-2,4-dithione (Heimgartner's reagent) [114234-09-2] **M 436.6, m 175-176°**. Recrystallise from toluene (light yellow solid), wash with Et₂O and dry in a vacuum. [*Helv Chim Acta* **70** 1001 1987.]

***N,O*-Bis-(trimethylsilyl)acetamide (BSA)** [10416-59-8] **M 203.4, b 71-73°/35mm, d 0.836, 1.4150**. Fractionate through a spinning band column and collect liquid **b 71-73°/35mm**, and not higher because the main impurity MeCONHSiMe₃ distills at **b 105-107°/35mm**. Used for derivatising alcohols and sugars [Klebe et al. *J Am Chem Soc* **88** 3390 1966, see *Carbohydr Res* **241** 209 1993 and **237** 313 1992]. It is **FLAMMABLE** and **TOXIC**.

Bis-(trimethylsilyl)acetylene [14630-40-1] **M 170.4, m 26°, b 134-136°/atm**. Dissolve in pet ether, wash with ice-cold dilute HCl. The pet ether extract is dried (MgSO₄), evaporated and fractionated at atmospheric pressure. [*J Organomet Chem* **37** 45 1972.]

Bis-(trimethylsilyl) sulfide (hexamethyldisilathiane) [3385-94-2] **M 178.5, b 65-67°/16mm, 162.5-163.5°/750mm corr, 164°/760mm, d 0.85, n 1.4598**. Dissolve in pet ether (*b ca* 40°), remove solvent and distilled. Redistilled under atmospheric pressure of dry N₂. It is collected as a colourless liquid which solidifies to a white solid in Dry-ice. On standing for several days it turns yellow possibly due to liberation of sulfur. Store below 4° under dry N₂. [*J Chem Soc* 3077 1950.]

Bis-(triphenylphosphine)nickel(II) chloride [14264-16-5] **M 654.2, m 225°(dec)**. Wash with glacial AcOH and dry in vac over H₂SO₄ and KOH until AcOH is removed. [*J Chem Soc* 719 1958.]

Boric acid (boracic acid) [10043-35-3] **M 61.8, m 171°, pK²⁵ 9.23**. Crystd three times from H₂O (3mL/g) between 100° and 0°, after filtering through sintered glass. Dried to constant weight over metaboric acid in a desiccator. It is steam volatile. After 2 recrystns of ACS grade it had Ag at 0.2 ppm.

9-Borabicyclo[3.3.1]nonane (9BBN) [monomer 280-64-8] [dimer 21205-91-4 or 70658-61-6] [1:1 coordination compound with tetrahydrofuran 76422-63-4] **M 122.0 (monomer), 244.0 (dimer), m 141-143° (monomer), 150-152°, 154-155° (dimer), b 195°/12mm**. Available as the solid dimer or in tetrahydrofuran soln. The solid is relatively stable and can be purified by distn in a vacuum (as dimer) and by recrystn from tetrahydrofuran (solubility at room temp is 9.5%, 0.78M), filter solid under N₂ wash with dry pentane and dry *in vacuo* at *ca* 100°. The solid is a dimer (IR 1567cm⁻¹), stable in air (for *ca* 2 months), and can be heated for 24h at 200° in an inert atmosphere without loss of hydride activity. It is a dimer in tetrahydrofuran soln also (IR 1567cm⁻¹). It is sensitive to H₂O and air (O₂) in soln. Concentration in soln can be determined by reaction with MeOH and measuring the vol of H₂ liberated, or it can be oxidised to *cis*-cyclooctane-1,5-diol (**m 73.5-74.5°**). [IR: *J Am Chem Soc* **90** 5280 1968, **96** 7765 1974; *J Org Chem* **41** 1778 1976, **46** 3978 1981.]

Borane pyridine complex [110-51-0] **M 92.9, m 8-10°, 10-11°, b 86°/7mm, 100-101°/12mm, d₄²⁰ 0.785**. Dissolve in Et₂O and wash with H₂O in which it is insol. Evap Et₂O and distil

(gives better than 99.8% purity). Its vap pressure is less than 0.1mm at room temp. [*J Am Chem Soc* **77** 1506 1955.]

Borane triethylamine complex [1722-26-5] **M 115.0, b 76°/4mm, 97.0°/12mm, d_4^{20} 0.78.** Distil in a vacuum using a 60cm glass helices packed column. [*J Am Chem Soc* **64** 325 1942, **84** 3407 1962; *Tetrahedron Lett* 4703 1968.]

Borane trimethylamine complex [75-22-9] **M 73.0, m 94-94.5°, b 171°/atm.** Sublimed using equipment described in *J Am Chem Soc* **59** 780 1937. Its vapour pressure is 86mm at 100°. Colourless hexagonal crystals varying from needles to short lumps, slightly soluble in H₂O (1.48% at 30°), EtOH (1%), hexane (0.74%) but very soluble in Et₂O, *C₆H₆ and AcOH. Stable at 125°. [*J Am Chem Soc* **59** 780 1939, **104** 325 1942.]

Boron trichloride (trichloroborane) [10294-34-5] **M 117.2, b 0°/476mm.** Purified (from chlorine) by passage through two mercury-filled bubblers, then fractionally distd under vacuum. In a more extensive purification the nitrobenzene addition compound is formed by passage of the gas over nitrobenzene in a vacuum system at 10°. Volatile impurities are removed from the crystalline yellow solid by pumping at -20°, and the BCl₃ is recovered by warming the addition compound at 50°. Passage through a trap at -78° removes entrained nitrobenzene; the BCl₃ finally condensing in a trap at -112° [Brown and Holmes *J Am Chem Soc* **78** 2173 1956]. Also purified by condensing into a trap cooled in acetone/Dry-ice, where it was pumped for 15min to remove volatile impurities. It was then warmed, recondensed and again pumped.

Boron trifluoride [7637-07-2] **M 67.8, b -101°/760mm.** The usual impurities - bromine, BF₅, HF and non-volatile fluorides - are readily separated by distn. Brown and Johannesen [*J Am Chem Soc* **72** 2934 1950] passed BF₃ into benzonitrile at 0° until the latter was satd. Evacuation to 10⁻⁵mm then removed all traces of SiF₄ and other gaseous impurities. [A small amount of the BF₃-benzonitrile addition compound sublimed and was collected in a U-tube cooled to -80°]. Pressure was raised to 20mm by admitting dry air, and the flask containing the BF₃ addition compound was warmed with hot water. The BF₃ evolved was passed through a -80° trap (to condense any benzonitrile) into a tube cooled in liquid air. The addition compound with anisole can also be used. For drying, BF₃ can be passed through H₂SO₄ saturated with boric oxide. Fumes in moist air. [Commercially available as a 1.3M soln in MeOH or PrOH.]

Boron trifluoride diethyl etherate [109-63-7] **M 141.9, b 67°/43mm, b 126°/760mm, d 1.154, n 1.340.** Treated with a small quantity of diethyl ether (to remove an excess of this component), and then distd under reduced pressure, from CaH₂. Fumes in moist air. **TOXIC.**

Bromine [7726-95-6] **M 159.8, b 59°, d 3.102, n 1.661.** Refluxed with solid KBr and distd, dried by shaking with an equal volume of conc H₂SO₄, then distd. The H₂SO₄ treatment can be replaced by direct distn from BaO or P₂O₅. A more extensive purification [Hildenbrand et al. *J Am Chem Soc* **80** 4129 1958] is to reflux about 1L of bromine for 1h with a mixture of 16g of CrO₃ in 200mL of conc H₂SO₄ (to remove organic material). The bromine is distd into a clean, dry, glass-stoppered bottle, and chlorine is removed by dissolving ca 25g of freshly fused CsBr in 500mL of the bromine and standing overnight. To remove HBr and water, the bromine was then distd back and forth through a train containing alternate tubes of MgO and P₂O₅. **HIGHLY TOXIC.**

Bromine pentafluoride [7789-30-2] **M 174.9, m -60.5°, b 41.3°, d^{25} 2.466.** Purified via its KF complex, as described for chlorine trifluoride. **HIGHLY TOXIC.**

2-Bromoallyltrimethylsilane [81790-10-5] **M 193.2, b 64-66°/10mm, 82-85°/58-60mm, d_4^{20} 1.13.** Fractionally distd through an efficient column. It is flammable. [*J Am Chem Soc* **104** 3733 6879 1982.]

2-Bromo-1,3,2-benzodioxaborole [51901-85-0] **M 198.8, m 47°, 51-53°, b 76°/9mm.** Keep at 20°/15mm for some time and then fractionally distil. [*J Chem Soc* 1529 1959.]

IR(endo,anti)-3-Bromocamphor-8-sulfonic acid ammonium salt [55870-50-3] M 328.2, m 284-285°(dec), $[\alpha]_D^{25} +84.8^\circ$ (c 4, H₂O). Passage of a hot aqueous soln through an alumina column removed water-soluble coloured impurities which remained on the column when the ammonium salt was eluted with hot water. The salt was crystd from water and dried over CaCl₂ [Craddock and Jones *J Am Chem Soc* 84 1098 1962; Kauffmann *J Prakt Chem* 33 295 1966].

Bromopyrogallol Red [16574-43-9] M 576.2, ϵ 5.45 x 10⁴ at 538nm (water pH 5.6-7.5). See Bromopyrogallol Red (5,5'-dibromopyrogallolsulfonephthalein) on p. 141 in Chapter 4.

Bromosulfalein (phenoltetrabromophthalein 3',3'-disulfonic acid disodium salt) [71-67-0] M 838.0. Purified by TLC on silica Gel G (Merck 250 μ particle size) in two solvent systems (BuOH-AcOH-H₂O 30:7.5:12.5 v/v; and BuOH-propionic acid-H₂O 30:20:7.5 v/v). When the solvent reached a height of 10cm the plate was removed, dried in air and developed with NH₃ vapour giving blue coloured spots. Also the dye was chromatographed on MN Silica Gel with *t*-BuOH-H₂O-*n*-BuOH (32:10:5 v/v and visualised with a dilute KOH (or NaOH if the Na salt is required) spray. The product corresponding to bromosulfalein was scraped off and eluted with H₂O, filtered and evap to dryness in a vacuum. It was dissolved in H₂O and filtered through Sephadex G-25 and evaporated to dryness. [UV and IR identification: *J Pharm Sci* 57 819 1968; NMR: *Chem Pharm Bull Jpn* 20 581 1972; *Anal Biochem* 83 75 1977.]

Bromtrimethylsilane (trimethylbromosilane, trimethylsilyl bromide) [2857-97-8] M 153.1, m -43.5° to -43.2°; b 40.5°/200mm, 77.3°/735mm, 79°/744mm, 79.8-79.9°/754mm, d_4^{20} 1.1805, d_4^{20} 1.190, n_D^{20} 1.422. Purified by repeated fractional distillation and stored in sealed ampoules in the dark. [*J Am Chem Soc* 75 1583 1953.] Also fractionally distd through a 15 plate column (0.8 x 32cm packed with 1/16in single turn helices from Pt-Ir wire). [*J Am Chem Soc* 68 1161 1946; 70 433 1948.]

tert-Butyldiphenylchlorosilane (TBDPSCI, tert-butylchlorodiphenylsilane) [58479-61-1] M 274.9, b 90°/0.015mm, d 1.057, n 1.568. Purified by repeated fractional distn. It is soluble in DMF and pentane [Hanessian and Lavalee *Can J Chem* 53 2975 1975; Robl et al. *J Med Chem* 34 2804 1991].

n-Butylmercuric chloride [543-63-5] M 293.1, m 130°. Crystd from EtOH.

n-Butylphenyl n-butylphosphonate [36411-99-1] M 270.3. Crystd three times from hexane as its compound with uranyl nitrate. See *tri-n-butyl phosphate* below.

tert-Butyldimethylsilyl chloride (TBDMSCl) [18162-48-6] M 150.7, m 87-89°, 92.5°, b 125°/760mm. Fractionally distd at atmospheric pressure. [*J Am Chem Soc* 76 1030 1954; 94 6190 1972.]

p-tert-Butylphenyl diphenyl phosphate [981-40-8] M 382.4, b 261°/6mm, n^{25} 1.5522. Purified by vacuum distn, and percolation through an alumina column, followed by passage through a packed column maintained at 150° to remove residual traces of volatile materials in a counter-current stream of N₂ at reduced pressure [Dobry and Keller *J Phys Chem* 61 1448 1957].

n-Butylstannic acid [PhSn(OH)₃, trihydroxy-n-butylstannane] [22719-01-3] M 208.8. Purified by adding excess KOH in CHCl₃ to remove *n*-BuSn(OH)Cl₂ and *n*-BuSn(OH)₂Cl, and isolated by acidification [Holmes et al. *J Am Chem Soc* 109 1408 1987].

Cacodylic acid (dimethylarsinic acid) [75-60-5] M 138.0, m 195-196°, pK^{25} 6.15 [Me₂As(O)OH]. Crystd from warm EtOH (3mL/g) by cooling and filtering. Dried in vacuum desiccator over CaCl₂. Has also been twice recrystd from propan-2-ol. [Koller and Hawkrigde *J Am Chem Soc* 107 7412 1985.]

[Previous Page](#)

IR(endo,anti)-3-Bromocamphor-8-sulfonic acid ammonium salt [55870-50-3] M 328.2, m 284-285°(dec), $[\alpha]_D^{25} +84.8^\circ$ (c 4, H₂O). Passage of a hot aqueous soln through an alumina column removed water-soluble coloured impurities which remained on the column when the ammonium salt was eluted with hot water. The salt was crystd from water and dried over CaCl₂ [Craddock and Jones *J Am Chem Soc* 84 1098 1962; Kauffmann *J Prakt Chem* 33 295 1966].

Bromopyrogallol Red [16574-43-9] M 576.2, ϵ 5.45 x 10⁴ at 538nm (water pH 5.6-7.5). See Bromopyrogallol Red (5,5'-dibromopyrogallolsulfonephthalein) on p. 141 in Chapter 4.

Bromosulfalein (phenoltetrabromophthalein 3',3'-disulfonic acid disodium salt) [71-67-0] M 838.0. Purified by TLC on silica Gel G (Merck 250 μ particle size) in two solvent systems (BuOH-AcOH-H₂O 30:7.5:12.5 v/v; and BuOH-propionic acid-H₂O 30:20:7.5 v/v). When the solvent reached a height of 10cm the plate was removed, dried in air and developed with NH₃ vapour giving blue coloured spots. Also the dye was chromatographed on MN Silica Gel with *t*-BuOH-H₂O-*n*-BuOH (32:10:5 v/v and visualised with a dilute KOH (or NaOH if the Na salt is required) spray. The product corresponding to bromosulfalein was scraped off and eluted with H₂O, filtered and evap to dryness in a vacuum. It was dissolved in H₂O and filtered through Sephadex G-25 and evaporated to dryness. [UV and IR identification: *J Pharm Sci* 57 819 1968; NMR: *Chem Pharm Bull Jpn* 20 581 1972; *Anal Biochem* 83 75 1977.]

Bromtrimethylsilane (trimethylbromosilane, trimethylsilyl bromide) [2857-97-8] M 153.1, m -43.5° to -43.2°; b 40.5°/200mm, 77.3°/735mm, 79°/744mm, 79.8-79.9°/754mm, d_4^{20} 1.1805, d_4^{20} 1.190, n_D^{20} 1.422. Purified by repeated fractional distillation and stored in sealed ampoules in the dark. [*J Am Chem Soc* 75 1583 1953.] Also fractionally distd through a 15 plate column (0.8 x 32cm packed with 1/16in single turn helices from Pt-Ir wire). [*J Am Chem Soc* 68 1161 1946; 70 433 1948.]

***tert*-Butyldiphenylchlorosilane (TBDPSCI, *tert*-butylchlorodiphenylsilane)** [58479-61-1] M 274.9, b 90°/0.015mm, d 1.057, n 1.568. Purified by repeated fractional distn. It is soluble in DMF and pentane [Hanessian and Lavalee *Can J Chem* 53 2975 1975; Robl et al. *J Med Chem* 34 2804 1991].

***n*-Butylmercuric chloride** [543-63-5] M 293.1, m 130°. Crystd from EtOH.

***n*-Butylphenyl *n*-butylphosphonate** [36411-99-1] M 270.3. Crystd three times from hexane as its compound with uranyl nitrate. See *tri-n-butyl phosphate* below.

***tert*-Butyldimethylsilyl chloride (TBDMSCl)** [18162-48-6] M 150.7, m 87-89°, 92.5°, b 125°/760mm. Fractionally distd at atmospheric pressure. [*J Am Chem Soc* 76 1030 1954; 94 6190 1972.]

***p-tert*-Butylphenyl diphenyl phosphate** [981-40-8] M 382.4, b 261°/6mm, n^{25} 1.5522. Purified by vacuum distn, and percolation through an alumina column, followed by passage through a packed column maintained at 150° to remove residual traces of volatile materials in a counter-current stream of N₂ at reduced pressure [Dobry and Keller *J Phys Chem* 61 1448 1957].

***n*-Butylstannic acid [PhSn(OH)₃, trihydroxy-*n*-butylstannane]** [22719-01-3] M 208.8. Purified by adding excess KOH in CHCl₃ to remove *n*-BuSn(OH)Cl₂ and *n*-BuSn(OH)₂Cl, and isolated by acidification [Holmes et al. *J Am Chem Soc* 109 1408 1987].

Cacodylic acid (dimethylarsinic acid) [75-60-5] M 138.0, m 195-196°, pK²⁵ 6.15 [Me₂As(O)OH]. Crystd from warm EtOH (3mL/g) by cooling and filtering. Dried in vacuum desiccator over CaCl₂. Has also been twice recrystd from propan-2-ol. [Koller and Hawkrigde *J Am Chem Soc* 107 7412 1985.]

Cadion [1-(4-nitrophenyl)-3-(4-phenylazophenyl)-triazene] [5392-67-6] M 346, m 198°. Commercial cadion is purified by recrystn from 95% EtOH and dried. It is stable in 0.2 N KOH (in 20% aqueous EtOH) at 25°. It is a sensitive reagent for Cd, and the Cd complex has λ_{\max} (EtOH) 475nm. [*Aust Chem Inst J Proc* 4 26 1937 ; *Anal Chim Acta* 19 377 1958.]

Cadmium [7440-43-9] M 112.4, m 321.1°, b 767°. Oxide has been removed by filtering the molten metal, under vacuum through quartz wool.

Cadmium acetate (2H₂O) [5743-04-4] M 230.5, m 255°(anhydr), d 2.01 (hydr), 2.34 (anhydr), pK_1^{25} 9.7, pK_2^{25} ~11.0 (for Cd²⁺). Crystd twice from anhydrous acetic acid and dried under vacuum for 24h at 100°.

Cadmium bromide (4H₂O) [13464-92-1 (4H₂O); 7789-42-6 (anhydr)] M 344.2, m 566°, b 963, d 5.19°. Crystd from water (0.6mL/g) between 100° and 0°, and dried at 110°. Forms monohydrate below 36° and the 4H₂O above 36°.

Cadmium chloride [10108-64-2] M 183.3, m 568°, b 960°, d 4.06. Crystd from water (1mL/g) by addition of EtOH and cooling.

Cadmium fluoride [7790-79-6] M 150.4, m >1000°, b 1748°, d 6.35. Crystd by dissolving in water at room temperature (25mL/g) and heating to 60°.

Cadmium iodide [7790-80-9] M 366.2, m 388°, b 787°, d 5.66. Crystd from ethanol (2mL/g) by partial evaporation.

Cadmium ionophore I [*N,N,N',N'*-tetramethyl-3,6-dioxooctanedi-(thioamide)] [73487-00-0] M 432.7, m 35-36°. Wash well with pet ether, then several times with 2N HCl (if it has a slight odour of pyridine) then H₂O and dry in a vacuum over H₂SO₄. It is a polar selectrophore for Cd. [*Helv Chim Acta* 63 217 1980.]

Cadmium lactate [16039-55-7] M 290.6. Crystd from water (10mL/g) by partial evapn in a desiccator.

Cadmium nitrate (4H₂O) [10022-68-1] M 308.5, m 59.5°. Crystd from water (0.5mL/g) by cooling in ice-salt.

Cadmium potassium iodide [13601-63-3] M 532.2. Crystd from ethanol by partial evapn.

Cadmium salicylate [19010-79-8] M 248.5, 242°(dec). Crystd from distd H₂O by evapn in a desiccator.

Cadmium sulfate [7790-84-3 (for 3CdSO₄ 8H₂O); 10124-36-4 (anhydr)] M 208.4 (anhydr), 769.5 (hydr). Crystd from distd water by partial evapn in a desiccator. On heating gives monohydrate at 80°.

Calcein sodium salt [2',7'-bis-{*N,N*-di(carboxymethyl)aminomethyl}fluorescein Na salt, **Fluorexon**, **Fluorescein Complexon**] [1461-15-0] M 666.5, $pK_{\text{Est}(1)}$ ~ 1.9, $pK_{\text{Est}(2)}$ ~ 2.5, $pK_{\text{Est}(3)}$ ~ 8.0, $pK_{\text{Est}(4)}$ ~ 10.5 (all for N-CH₂COOH), and $pK_{\text{Est}(5)}$ ~ 3.5 (for benzoic COOH). Dissolve in distilled H₂O and acidify with dilute HCl to pH 3.5. Filter off the solid acid and wash well with H₂O. Redissolve ca 10g in 300mL H₂O containing 12g of NaOAc. Ppte again by adding HCl, filter and wash with H₂O. Add the solid to 200mL of EtOH stir for 1h and filter. Repeat the EtOH wash and dry the bright yellow solid in a vacuum. This acid decomposes on heating at ca 180°. See below for the prepn of the Na salt. [*Anal Chem* 28 882 1956].

Dissolve in H₂O and acidify with 3N HCl to pH 3.5. Collect the solid and wash with H₂O. The air-dried ppte is extracted with 70% aqueous EtOH, filtered hot and cooled slowly. Fine yellow needles of the acid crystallise out, are filtered and dissolved in the minimum quantity of 0.01N NaOH and reppted with N HCl to pH 3.5. It is then recrystd from 70% aqueous EtOH (3x). The final product (acid) is dried at 80° in a vacuum for 24h, m >300°dec. It contains one mol of water per mol of acid (C₃₀H₃₆N₄O₁₃·H₂O). The product is pure as revealed

by electrophoresis at pH 5.6 and 8.6, and by TLC in *i*-BuOH-*i*-PrOH-AcOH-H₂O (60:60:5:5 by vol) or *i*-PrOH or pH 8.0 borate buffer. [Wallach et al. *Anal Chem* 31 456 1959.]

The Na salt is prepared by dissolving the in H₂O containing 2 mols of NaOH per mol of acid reagent and lyophilising. It complexes with Ca and Mg ions.

Calcium [7440-70-2] **M 40.1, m 845°**. Cleaned by washing with ether to remove adhering paraffin, filed in an argon-filled glove box, and washed with ethanol containing 2% of conc HCl. Then washed with dry ethanol, dried in a vac and stored under pure argon [Addison, Coltreay and Halsstead, *J Chem Soc* 3868 1962].

Calcium acetate monohydrate [5743-26-0 (H₂O), 62-54-4 (xH₂O)] **M 176.2 (H₂O), m 150° (loses H₂O), pK²⁵ 12.7 (for Ca₂⁺)**. Crystd from water (3mL/g) by partial evapn in a desiccator.

Calcium benzoate (3H₂O) [2090-05-3] **M 336.4**. Crystd from water (10m/g) between 90° and 0°.

Calcium bromide (H₂O) [62648-72-0; 71626-99-8 (xH₂O); 7789-41-5 (anhydr)] **M 217.9, d 3.35**. Crystd from EtOH or Me₂CO. It loses H₂O on heating and is anhydrous at 750° then it loses Br. Deliquescent.

Calcium butyrate [5743-36-2] **M 248.2**. Crystd from water (5mL/g) by partial evapn in a desiccator.

Calcium carbamate [543-88-4] **M 160.1**. Crystd from aqueous ethanol.

Calcium chloride (anhydrous) [10043-52-4] **M 111.0, m 772°, b >1600°, d₄¹⁵ 2.15**. Available as fused granules or cubic crystals. It is very *hygroscopic*. Very soluble in H₂O (exothermic), and EtOH. Store in a tightly closed container.

Calcium chloride (2H₂O) [10035-04-8] **M 147.0, m 175°(dehydr), 772°(dec)**. Crystd from ethanol, and is hygroscopic. Loses H₂O at 200° so it can be dried at high temperatures to dehydrate. *Hexahydrate* [7774-34-7] has **m 30°** and **d 1.67**.

Calcium dithionite [13812-88-9] **M 168.2, m dec on heating**. Crystd from water, or water followed by acetone and dried in air at room temperature.

Calcium D-gluconate monohydrate [299-28-5] **M 448.4, m dec on heating, [α]₅₄₆²⁰ +11.0°, [α]_D²⁰ +9.0° (c 1.2, H₂O)**. It is sol in H₂O (3.5g in 100g at 25°). Dissolve in H₂O, filter and ppt by adding MeOH. Filter off solid and dry in a vacuum at 85°. Alternatively, dissolve in H₂O, filter (from insol inorganic Ca) and evaporate to dryness under vacuum at 85°. [*J Am Pharm Assoc* 41 366 1952.]

Calcium D-heptagluconate dihydrate [17140-60-2] **M 526.4, [α]₅₄₆²⁰ +5.2°, [α]_D²⁰ +4.4° (c 5, H₂O)**. Purified same as calcium D-gluconate.

Calcium formate [544-17-2] **M 130.1, m dec on heating, d 2.01**. Crystd from water (5mL/g) by partial evaporation in a desiccator.

Calcium hexacyanoferrate (II) (11H₂O) [13821-08-4] **M 490.3**. Recrystd three times from conductivity H₂O and air dried to constant weight over partially dehydrated salt. [*Trans Faraday Soc* 45 855 1949.] Alternatively the Ca salt can be purified by pptn with absolute EtOH in the cold (to avoid oxidation) from an air-free saturated aqueous soln. The pure lemon yellow crystals are centrifuged, dried in a vacuum desiccator first over dry charcoal for 24h, then over partly dehydrated salt and stored in a dark glass stoppered bottle. No deterioration occurred after 18 months. No trace of Na, K or NH₄ ions could be detected in the salt from the residue after decomposition of the salt with conc H₂SO₄. Analyses indicate 11mols of H₂O per mol of salt. The solubility in H₂O is 36.45g (24.9°) and 64.7g (44.7°) per 100g of solution. [*J Chem Soc* 50 1926.]

Calcium hydroxide [1305-62-0] **M 74.1, m loses H₂O on heating, pK²⁵ 12.7 (for Ca²⁺)**. Heat analytical grade calcium carbonate at 1000° during 1h. Allow the resulting oxide to cool and add slowly to

water. Heat the suspension to boiling, cool and filter through a sintered glass funnel of medium porosity (to remove soluble alkaline impurities). Dry the solid at 110° and crush to a uniformly fine powder.

Calcium iodate [7789-80-2 (H_2O)] **M 389.9, m >540°, pK²⁵ 0.79 (for HIO₃)**. Crystd from water (100mL/g).

Calcium iodide (xH_2O) [71626-98-7 (xH_2O); 10102-68-8 (*anhydr*)] **M 293.9 (for 4H₂O), m 740°, b 1100°**. Dissolved in acetone, which was then diluted and evaporated. This drying process was repeated twice, then the CaI₂ was crystd from acetone-diethyl ether and stored over P₂O₅. Very *hygroscopic* when anhydrous and is light sensitive [Cremllyn et al. *J Chem Soc* 528 1958]. *Hexahydrate* has m 42°.

Calcium ionophore I (ETH 1001) [58801-34-6] **M 685.0**. This is a neutral Ca selectophore. It can be purified by thick layer (2mm) chromatography (Kieselgel F₂₄₅) and eluted with Me₂CO-CHCl₃ (2:1). [*Helv Chim Acta* 56 1780 1973.]

Calcium ionophore II (ETH 129) [74267-27-9] **M 460.7, m 153-154°**. Recrystd from Me₂CO. It forms 1:2 and 1:3 metal/ligand complexes with Mg²⁺ and Ca²⁺ ions respectively, and induces selectivity in membranes for Ca²⁺ over Mg²⁺ by a factor of *ca* 10⁴. [*Helv Chim Acta* 63 191 1980.]

Calcium ionophore III [A23187 calcimycin] [52665-69-7] **M 523.6, m 181-182°, [α]_D²⁵ -56.0° (c 1, CHCl₃)**. Recrystallises from Me₂CO as colourless needles. Protect from light and moisture, store in a refrigerator. Soluble in Me₂SO or EtOH and can be stored for 3 months without loss of activity. Mg and Ca salts are soluble in organic solvents and cross biological membranes. It has a pK_a of 6.9 in 90% Me₂SO. The Ca complex cryst from 50% EtOH as colourless prisms. *Highly TOXIC* [*Ann Rev Biochem* 45 501 1976; *J Am Chem Soc* 96 1932 1974, *J Antibiotics* 29 424 1976.]

Calcium isobutyrate [533-90-4] **M 248.2**. Crystd from water (3mL/g) by partial evapn in a desiccator.

Calcium lactate (5H₂O) [814-80-2] **M 308.3, m anhydr at 120°**. Crystd from warm water (10mL/g) by cooling to 0°.

Calcium nitrate (4H₂O) [13477-34-4] **M 236.1, m 45°(dehydr), 560°(anhydr)**. Crystd four times from water (0.4mL/g) by cooling in a CaCl₂-ice freezing mixture. The tetrahydrate was dried over conc H₂SO₄ and stored over P₂O₅, to give the anhydrous salt. It is deliquescent. After 3 recrystns of ACS grade it had Co, Fe, Mg, Sr and Zn at 0.2, 1. 0, 0.02, 10 and 0.02 ppm resp.

Calcium nitrite (2H₂O) [13780-06-8 (30%w/w aq soln)] **M 150.1(hydr), m dec on heating, d 2.22**. Crystd from hot water (1.4mL/g) by adding ethanol and cooling to give the hydrate. It is deliquescent.

(+)-Calcium pantothenate (H₂O) (D(+)- 137-08-6; 63409-48-3] **M 476.5**. See R(+)-pantothenic acid calcium salt on p. 555 in Chapter 6.

Calcium permanganate (4H₂O) [10118-76-0 (*anhydr*)] **M 350.0 (for 4H₂O)**. Crystd from water (3.3mL/g) by partial evapn in a desiccator. It is deliquescent.

Calcium propionate [4075-81-4] **M 186.2, m dec on heating**. Crystd from water (2mL/g) by partial evapn in a desiccator.

Calcium salicylate (2H₂O) [824-35-1] **M 350.4**. Crystd from water (3mL/g) between 90° and 0°.

Calcium sulfate dihydrate [10101-41-4] **M 172.1, m 150(dec), d 2.32**. Loses only part of its H₂O at 100-150° (see below). Soluble in H₂O and very slowly soluble in glycerol. Insoluble in most organic solvents.

Calcium sulfate hemihydrate [10034-76-1] **M 145.2**. Sol in H₂O (0.2 parts/100 at 18.75°). Completely dehydrated >650°. Dry below 300° to give a solid with estimated pore size *ca* 38% of vol. Anhydrous CaSO₄ has high affinity for H₂O and will absorb 6.6% of its weight of H₂O to form the hemihydrate (gypsum). It sets to a hard mass with H₂O, hence should be kept in a tightly sealed container.

Calcium thiosulfate [10124-41-1] **M 152.2, m 43-49°, pK₁^{2.5} 0.6, pK₂^{2.5} 1.74 (for H₂S₂O₃)**. Recrystd from water below 60° in a N₂ atmosphere, followed by drying with EtOH and Et₂O. Stored in a refrigerator. [Pethybridge and Taba *J Chem Soc, Faraday Trans 1* **78** 1331 1982.]

(4-Carbamylphenylarsylenedithio)diacetic acid [531-72-6] **M 345.1, pK_{Est}~3.5**. Crystd from MeOH or EtOH.

Carbonate ionophore I [ETH 6010] (heptyl 4-trifluoroacetylbenzoate) [129476-47-7] **M 316.3, b 170°/0.02 Torr, d 0.909**. Purified by flash chromatography (2g of reagent with 30g of Silica Gel 60) and eluted with EtOAc/hexane (1:19). The fractions that absorbed at 260nm were pooled, evapd and dried at room temp (10.3 Torr). The oily residue was distd in a bubbled-tube apparatus (170°/0.02 Torr). Its IR (CHCl₃) had peaks at 1720, 1280, 940cm⁻¹ and its sol in tetrahydrofuran is 50mg/0.5mL. It is a lipophilic neutral ionophore selective for carbonate as well as being an optical humidity sensor. [*Anal Chim Acta* **233** 41 1990.]

Carbon dioxide [124-38-9] **M 44.0, sublimates at -78.5°, pK₁^{2.5} 6.35, pK₂^{2.5} 10.33 (for H₂CO₃)**. Passed over CuO wire at 800° to oxidise CO and other reducing impurities (such as H₂), then over copper dispersed on Kieselguhr at 180° to remove oxygen. Drying at -78° removed water vapour. Final purification was by vacuum distn at liquid nitrogen temperature to remove non-condensable gases [Anderson, Best and Dominey *J Chem Soc* 3498 1962]. Sulfur dioxide can be removed at 450° using silver wool combined with a plug of platinised quartz wool. Halogens are removed by using Mg, Zn or Cu, heated to 450°.

Carbon disulfide, see entry on p. 156 in Chapter 4.

Carbon monoxide [630-08-0] **M 28.0, b -191.5°**. Iron carbonyl is a likely impurity in CO stored under pressure in steel tanks. It can be decomposed by passage of the gas through a hot porcelain tube at 350-400°. Passage through alkaline pyrogallol soln removes oxygen (and CO₂). Removal of CO₂ and water are effected by passage through soda-lime followed by Mg(ClO₄)₂. Carbon monoxide can be condensed and distd at -195°. **HIGHLY POISONOUS gas.**

Carbonyl bromide [593-95-3] **M 187.8, b 64.5°/760mm**. Purified by distn from Hg and from powdered Sb to remove free bromine, then vacuum distd to remove volatile SO₂ (the major impurity) [Carpenter et al. *J Chem Soc, Faraday Trans 2* 384 1977]. **TOXIC**

Carbonyl sulfide [463-58-1] **M 60.1, m -138°, b -47.5°, -50°**. Purified by scrubbing through three consecutive fritted washing flasks containing conc NaOH at 0° (to remove HCN), and then through conc H₂SO₄ (to remove CS₂) followed by a mixture of NaN₃ and NaOH solution; or passed through traps containing satd aq lead acetate, then through a column of anhydrous CaSO₄. Then freeze-pumped repeatedly and distd through a trap packed with glass wool and cooled to -130° (using an *n*-pentane slurry). It liquefies at 0°/12.5mm. The gas is stored over conc H₂SO₄. **TOXIC**

Catecholborane (1,3,2-Benzodioxaborole) [274-07-7] **M 119.2, b 50°/50mm, 66°/80mm, 76-77°/100mm, 88°/165mm, d 1.125, n 1.507 (also available as a 1.0M soln in THF)**. A moisture sensitive flammable liquid which is purified by distn in a vacuum under a N₂ atmosphere and stored under N₂ at 0-4°. It liberates H₂ when added to H₂O or MeOH. A soln in THF after 25h at 25° has residual hydride of 95%(under N₂) and 80% (under air) [Brown and Gupta *J Am Chem Soc* **97** 5249 1975].

Celite 545 (diatomaceous earth) [12003-10-0]. Stood overnight in conc HCl after stirring well, then washed with distilled water until neutral and free of chloride ions. Washed with methanol and dried at 50°.

Ceric ammonium nitrate [16774-21-3] M 548.2, pK_1^{25} -1.15, pK_2^{25} -0.72, pK_3^{25} 1.68, pK_4^{25} 2.29 (for aquo Ce^{4+}). Ceric ammonium nitrate (125g) is warmed with 100mL of dilute HNO_3 (1:3 v/v) and 40g of NH_4NO_3 until dissolved, and filtered off on a sintered-glass funnel. The solid which separates on cooling in ice is filtered off on a sintered funnel (at the pump) and air is sucked through the solid for 1-2 h to remove most of the nitric acid. Finally, the solid is dried at 80-85°.

Cerous acetate [537-00-8] M 317.3, pK_1^{25} 8.1 (9.29), pK_2^{25} 16.3, pK_3^{25} 26.0 (for Ce^{3+}). Crystd twice from anhydrous acetic acid, then pumped dry under vacuum at 100° for 8h.

Cesium bromide [7787-69-1] M 212.8, m 636°, b ca 1300°, d 4.44. Very soluble in H_2O , soluble in EtOH but insoluble in Me_2CO . Dissolve in the minimum volume of H_2O , filter and ppte by adding Me_2CO . Filter solid and dry at 100°. Also recrystd from water (0.8mL/g) by partial evaporation in a desiccator.

Cesium carbonate [534-17-8] M 325.8, m 792°(at red heat). Crystd from ethanol (10mL/g) by partial evaporation.

Cesium chloride [7647-17-8] M 168.4, m 645°, b 1303°, d 3.99. Soluble in H_2O but can be purified by crystn from H_2O [sol in g per cent: 162.3(0.7°), 182.2(16.2°) and 290(at bp 119.4°)] and dried in high vac. Sol in EtOH and is deliquescent, keep in a tightly closed container. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 951 1963.] For further purification of $CsCl$, a conc aqueous soln of the practically pure reagent is treated with an equivalent weight of I_2 and Cl_2 bubbled into the soln until pptn of $CsCl_2I$ ceased. Recrystn yields a salt which is free from other alkali metals. It is then decomposed to pure $CsCl$ on heating. [*J Am Chem Soc* 52 3886 1930.] Also recrystd from acetone-water, or from water (0.5mL/g) by cooling in $CaCl_2$ /ice. Dried at 78° under vacuum.

Cesium chromate [56320-90-2] M 381.8, pK_1^{25} 0.74, pK_2^{25} 6.49 (for H_2CrO_4). Crystd from water (1.4mL/g) by partial evapn in a desiccator.

Cesium fluoride [13400-13-0] M 151.9, m 703°. Crystd from aqueous soln by adding ethanol.

Cesium iodide [7789-17-5] M 259.8, m 621°, b-1280°, d 4.5. Crystd from warm water (1mL/g) by cooling to -5°.

Cesium nitrate [7789-18-6] M 194.9, m 414°(dec), d 3.65. Crystd from water (0.6mL/g) between 100° and 0°. After 1 crystn of 99.9% grade it had K, Na and Se at 0.8, 0.4 and 0.2 ppm resp.

Cesium oleate [31642-12-3] M 414.4. Crystd from EtOAc, dried in an oven at 40° and stored over P_2O_5 .

Cesium perchlorate [13454-84-7] M 232.4, pK^{25} -2.4 to -3.1 (for $HClO_4$). Crystd from water (4mL/g) between 100° and 0°.

Cesium perfluoro-octanoate [17125-60-9] M 546.0. Recrystd from a butanol-petroleum ether mixture, dried in an oven at 40° and stored over P_2O_5 under vacuum.

Cesium sulfate [10294-54-9] M 361.9, m 1005°, d 4.24. Crystd from water (0.5mL/g) by adding ethanol and cooling.

Chloramine-T (*N*-chloro-*p*-toluenesulfonamide sodium salt) $3H_2O$ [7080-50-4] M 281.7, m 168-170°(dec). Crystd from hot water (2mL/g). Dried in a desiccator over $CaCl_2$ where it loses water. Protect from sunlight. Used for detection of bromate and halogens, and Co, Cr, Fe, Hg, Mn, Ni and Sb ions.

Chlorazol Sky Blue FF {6,6'-[(3,3'-dimethoxy[1,1'-biphenyl]-4,4'-diyl)bis(azo)bis(4-amino-5-hydroxy-1,3-naphthylenedisulfonic acid) tetra-Na salt [2610-05-1] M 996.9, m

>300°(dec). Freed from other electrolytes by adding aqueous sodium acetate to a boiling soln of the dye in distd water. After standing, the salted-out dye was filtered on a Büchner funnel, the process being repeated several times. Finally, the pptd dye was boiled several times with absolute EtOH to wash out any sodium acetate, then dried (as the sodium salt) at 105°. [McGregor, Peters and Petropolous *Trans Faraday Soc* **58** 1045 1962.]

Chlorine [7782-50-5] **M 70.9, m -101.5°, b -34.0°, d 2.898.** Passed in succession through aqueous KMnO₄, dilute H₂SO₄, conc H₂SO₄, and a drying tower containing Mg(ClO₄)₂. Or, bubbled through with water, dried over P₂O₅ and distd from bulb to bulb in a vac line. **HIGHLY TOXIC.**

Chlorine trifluoride [7790-91-2] **M 92.5, b 12.1°.** Impurities include chloryl fluoride, chlorine dioxide and hydrogen fluoride. Passed first through two U-tubes containing NaF to remove HF, then through a series of traps in which the liquid is fractionally distd. Can be purified *via* the KF complex, KClF₄, formed by adding excess ClF₃ to solid KF in a stainless steel cylinder in a dry-box and shaking overnight. After pumping out the volatile materials, pure ClF₃ is obtained by heating the bomb to 100-150° and condensing the evolved gas in a -196° trap [Schack, Dubb and Quaglino *Chem Ind (London)* 545 1967]. **HIGHLY TOXIC.**

Chlorodiphenylphosphine (diphenylphosphinous chloride) [1079-66-9] **M 220.6, m 15-16°, b 124-126°/0.6mm, 174°/5mm, 320°/atm, d 1.229, n 1.636.** Air sensitive, pale yellow lachrymatory liquid which is purified by careful fractional distn and discarding the lower boiling fraction which contains the main impurity PhPCl₂ (b 48-51°/0.7mm); and checking for impurities by NMR. [Weinberg *J Org Chem* **40** 3586 1975; Honer et al. *Chem Ber* **94** 2122 1961.]

4-(Chloromercuri)benzenesulfonic acid monosodium salt [14110-97-5] **M 415.2, dec on heating.** The free acid is obtained by acidifying an aq soln, filtering off the acid, washing it with H₂O and recrystallising from hot H₂O to give a colourless solid which is dried in a vacuum over P₂O₅ and should give negative Cl⁻ ions. The Na salt is made by dissolving in one equivalent of aqueous NaOH and evaporate to dryness. [*Chem Ber* **67** 130 1934; *J Am Chem Soc* **76** 4331 1954.] **HIGHLY TOXIC.**

4-Chloromercuribenzoic acid [59-85-8] **M 357.2, m >300°.** Its suspension in water is stirred with enough 1M NaOH to dissolve most of it: a small amount of insoluble matter is removed by centrifugation. The chloromercuribenzoic acid is then pptd by adding 1M HCl and centrifuged off. The pptn is repeated twice. Finally, the ppte is washed three times with distilled water (by centrifuging), then dried in a thin layer under vacuum over P₂O₅ [Boyer *J Am Chem Soc* **76** 4331 1954].

Chloromethylphosphonic acid dichloride [1983-26-2] **M 167.4, b 50°/0.5mm, 52-53(59)°/2mm, 63-65°/3mm, 78-79°/10mm, 87-88°/15mm, 102-103°/30mm, d₄²⁰ 1.638, n_D²⁰ 1.4971.** Fractionally distd using a short Claisen column and redistd. The *aniline salt* has m 199-201°. The ³¹P NMR has a line at -38±2 ppm from 85% H₃PO₄. [Kinnear and Perren *J Chem Soc* 3437 1952; NMR: *J Am Chem Soc* **78** 5715 1956; *J Org Chem* **22** 462 1957.]

2-Chloro-2-oxo-1,3,2-dioxaphospholane [6609-64-9] **M 142.5, m 12-14°, b 89-91°/0.8mm, d₄²⁰ 1.549, n_D²⁰ 1.448.** Should be distd at high vacuum as some polymerisation occurs on distn. It has IR bands at 3012, 2933, 1477, 1366, 1325, 1040, 924 and 858 cm⁻¹. In H₂O at 100° it is hydrolysed to HOCH₂CH₂OPO₃H₂ in 30min [IR: Cox and Westheimer *J Am Chem Soc* **80** 5441 1958].

2-Chlorophenyl diphenyl phosphate [115-85-5] **M 360.7, b 236°/4mm, n_D²⁵ 1.5707.** Purified by vacuum distn, percolated through a column of alumina, then passed through a packed column maintained by a countercurrent stream of nitrogen at reduced pressure [Dobry and Keller *J Phys Chem* **61** 1448 1957].

Chlorosulfonic (chlorosulfuric) acid [7790-94-5] **M 116.5, b 151-152°/750mm, d₄ 1.753, n 1.4929, pK -5.9 (aq H₂SO₄).** Distd in an all-glass apparatus, taking the fraction boiling at 156-158°. Reacts **EXPLOSIVELY** with water [Cremlyn *Chlorosulfonic acid: A Versatile Reagent*, Royal Society of Chemistry UK, 2002, 308 pp, ISBN 0854044981].

Chloro-(2,2':6',2'-terpyridine)platinum (II) chloride (2H₂O) [60819-00-3] **M 535.3**. Recrystd from hot dilute HCl and cooling to give the red dihydrate. The trihydrate crystals slowly from a cold aq soln and is air dried. The red dihydrate can be obtained from the trihydrate by desiccation over conc H₂SO₄, by washing with EtOH or by precipitating from a warm aq soln with HCl. The dihydrate is also formed by decomposing the black trihydrate form by heating in water (slowly), or more rapidly with hot 2N HCl. [*J Chem Soc* 1498 1934.]

Chloro-tri-isopropyl titanium [20717-86-6] **M 260.6, m 45-50°, b 61-65°/0.1mm**. Distd under vacuum and sets slowly to a solid on standing. Stock reagents are made by dissolving the warm liquid in pentane, toluene, Et₂O, THF, CH₂Cl₂, and can be stored in pure state or in soln under dry N₂ for several months. The reagent is *hygroscopic* and is hydrolysed by H₂O. [*Chem Ber* 118 1421 1985.]

Chlorotriphenylsilane (triphenylchlorosilane) [76-86-8] **M 294.9, m 90-92°, 91-93°, 94-95°, 97-99°, b 156°/1mm, 161°/0.6mm**. Likely impurities are tetraphenylsilane, small amounts of hexaphenyldisiloxane and traces of triphenylsilanol. Purified by distn at 2mm, then crystd from EtOH-free CHCl₃, and from pet ether (b 30-60°) or hexane by cooling in a Dry-ice/acetone bath. [*J Chem Soc* 3671 1957; *J Am Chem Soc* 72 4471 1958, 77 6395 1955, 79 1843 1957.]

Chlorotris(triphenylphosphine) rhodium I (Wilkinson's catalyst) [14694-95-2] **M 925.2, m 138°(dec), 140°(dec), 157-158°(dec)**. Forms dark burgundy crystals from hot EtOH after refluxing for 30min. When the soln is heated for only 5min orange crystals are formed, Heating the orange crystals in EtOH yields red crystals. Crystn from Me₂CO gives the orange crystals. The two forms have similar IR spectra but X-rays are slightly different. [Osborne et al. *J Chem Soc (A)* 1711 1966; Osborne and Wilkinson *Inorg Synth* 10 67 1967; Bennett and Donaldson *Inorg Chem* 16 655 1977.] Sol in CH₂Cl₂ ~2% (25°), in toluene 0.2% (25°), and less sol in Me₂CO, MeOH, BuOH and AcOH, but insol in pet ethers and cyclohexane. It reacts with donor solvents such as pyridine, DMSO and MeCN.

Chromazurol S [1667-99-8] **M 539.3, λ_{max} 540nm, ε 7.80 x 10⁴ (10M HCl), CI 43825, pK₁^{2.5} <0, pK₂^{2.5} 2.25, pK₃^{2.5} 4.88, pK₄^{2.5} 11.75**. Crude *phenolic triphenylmethanecarboxysulfonic acid triNa salt* (40g) is dissolved in water (250mL) and filtered. Then added conc HCl (50mL) to filtrate, with stirring. Ppte is filtered off, washed with HCl (2M) and dried. Redissolved in water (250mL) and pptn repeated twice more in water bath at 70°. Then dried under vacuum over solid KOH (first) then P₂O₅ [Martynov et al. *Zh Analyt Khim* 32 519 1977]. It has also been purified by paper chromatography using *n*-butanol, acetic acid and water (7:3:1). First and second spots were extracted. It chelates Al and Be. Used for estimating fluoride.

Chromic chloride (anhydrous) [10025-73-7] **M 158.4, m 1152°, pK₁^{2.5} 3.95, pK₂^{2.5} 5.55, pK₃^{2.5} 10.5 (for Cr³⁺)**. Sublimed in a stream of dry HCl. Alternatively, the impure chromic chloride (100g) was added to 1L of 10% aq K₂Cr₂O₇ and several millilitres of conc HCl, and the mixture was brought to a gentle boil with constant stirring for 10 min. (This removed a reducing impurity.) The solid was separated, and washed by boiling with successive 1L lots of distilled water until the wash water no longer gave a test for chloride ion, then dried at 110° [Poulsen and Garner *J Am Chem Soc* 81 2615 1959].

Chromium (III) acetylacetonate [21679-31-2] **M 349.3, m 212-216°, 216°, pK₁^{2.5} 4.0 (see chromic chloride)**. Purified by dissolving 6g in hot *C₆H₆ (20mL) and adding 75mL of pet ether slowly. Cool to room temp then chill on ice, filter off and dry in air to give 2.9g. Also crystallises from EtOH. Sol in heptane, *C₆H₆, toluene and pentane-2,4-dione at 20-40°. It forms a 1:2 complex with CHCl₃. [*Inorg Synth* 5 130 1957; *J Am Chem Soc* 80 1839 1958.]

Chromium ammonium sulfate (12H₂O) [34275-72-4 (*hydr*); 13548-43-1 (*anhydr*)] **M 478.4, m 94° loses 9H₂O then dehydr at 300°, d 1.72**. Crystd from a saturated aqueous soln at 55° by cooling slowly with rapid mechanical stirring. The resulting fine crystals were filtered on a Büchner funnel, partly dried on a porous plate, then equilibrated for several months in a vacuum desiccator over crude chromium ammonium sulfate (partially dehydrated by heating at 100° for several hours before use) [Johnson, Hu and Horton *J Am Chem Soc* 75 3922 1953].

Chromium (II) chloride (anhydrous) [10049-05-5] M 122.9, m 824°, d₄¹⁴ 2.75. Obtained from the dihydrate by heating *in vacuo* at 180°. It is a very *hygroscopic* white powder which dissolves in H₂O to give a sky blue solution. Stable in dry air but oxidises rapidly in moist air and should be stored in air tight containers. It sublimates at 800° in a current of HCl gas and cooled in the presence of HCl gas. Alternatively it can be washed with air-free Et₂O and dried at 110-120°. [*Inorg Synth* 3 150 1950.]

Chromium hexacarbonyl [13007-92-6] M 220.1, m 130°(dec), d 1.77. Wash with cold EtOH then Et₂O and allow to dry in air. Alternatively recrystallise from dry Et₂O. This is best accomplished by placing the hexacarbonyl in a Soxhlet extractor and extracting exhaustively with dry Et₂O. Pure Cr(CO)₆ is filtered off and dried in air. Completely colourless refracting crystals are obtained by sublimation at 40-50°/ <0.5mm in an apparatus where the collecting finger is cooled by Dry Ice and in which there is a wide short bore between the hot and cold sections to prevent clogging by the crystals. Loss of product in the crystn and sublimation is slight. It is important not to overdo the drying as the solid is appreciably volatile and **TOXIC** [vapour pressure is 0.04(8°), 1.0(48°) and 66.5(100°) mm]. Also do not allow the Et₂O solns to stand too long as a brown deposit is formed which is sensitive to light, and to avoid the possibility of violent decomposition. It sinters at 90°, dec at 130°, and **EXPLODES** at 210°. [*Inorg Synth* 3 156 1950; *J Am Chem Soc* 83 2057 1961.]

Chromium potassium sulfate (12H₂O) [7788-99-0] M 499.4, pK₁²⁵ 0.74, pK₂²⁵ 6.49 (for H₂CrO₄, chromic acid). Crystd from hot water (2mL/g) by cooling.

Chromium trioxide (chromic anhydride) [1333-82-0] M 100.0, m 197°, dec at 250° to Cr₂O₃, d 2.70 (pK₁²⁵ 0.74, pK₂²⁵ 6.49, for H₂CrO₄, chromic acid). Red crystals from water (0.5mL/g) between 100° and -5°, or from water/conc HNO₃ (1:5). It separates when potassium or sodium dichromate are dissolved in conc H₂SO₄. Dried in a vacuum desiccator over NaOH pellets; *hygroscopic*, powerful oxidant, can ignite with organic compounds. It is a skin and pulmonary **IRRITANT**.

§ Commercially available on polymer support. **CANCER SUSPECT**.

Chromium (III) tris-2,4-pentanedionate [21679-31-2] M 349.3, m 216°, pK²⁵ 4.0 (see **chromic chloride**). See chromium (III) acetylacetonate on p. 412.

Chromoionophore I [ETH 5294] [9-diethylamino-5-octadecanoyl-imino-5-*H*-benzo[*a*]-phenoxazine] [125829-24-5] M 583.9. Purified by flash chromatography (Silica Gel) and eluted with EtOAc. The coloured fractions are pooled, evaporated and recrystd from EtOAc. It is a lipophilic chromoionophore and is a selectophore for K and Ca ions. [*Anal Chem* 62 738 1990.]

Chromotropic acid (4,5-dihydroxynaphthalene-2,7-disulfonic acid di-Na salt) [5808-22-0 (2H₂O)] M 400.3, m >300°, pK₁ 0.61(SO₃⁻), pK₂ 0.7(SO₃⁻), pK₃ 5.45(OH), pK₄ 15.5(OH). See disodium 4,5(1,8)-dihydroxynaphthalene-2,7(3,6)-disulfonate (2H₂O) on p. 421.

Chromyl chloride [14977-61-8] M 154.9, b 115.7°, d 1.911. Purified by distn under reduced pressure. **TOXIC**.

Claisen alkali (alkali Claisen). Prepared from KOH (35g) in H₂O (25mL) and diluted to 100mL with MeOH. **STRONGLY CAUSTIC**.

Cobalt (II) meso-5,10,15,20-tetraphenylporphine complex [14172-90-8] M 671.7. Brown crystals from Et₂O or CHCl₃-MeOH (*cf iron chloride complex*). Recrystd by extraction (Soxhlet) with *C₆H₆. Sol in most organic solvents except MeOH and pet ether. [UV, IR: *J Am Chem Soc* 70 1808 1948; 81 5111 1959.]

Cobaltous acetate (4H₂O) [6147-53-1] M 249.1, pK₁²⁵ 9.85 (for Co²⁺). Crystd several times as the tetrahydrate from 50% aqueous acetic acid. Converted to the anhydrous salt by drying at 80°/1mm for 60h.

Cobaltous acetylacetonate [14024-48-7] M 257.2, m 172°. Crystd from acetone.

Cobaltous ammonium sulfate (6H₂O) [13596-46-8] M 395.5, d 1.90. Crystd from boiling water (2mL/g) by cooling. Washed with ethanol.

Cobaltous bromide (6H₂O) [85017-77-2 (*x*H₂O); 7789-43-7 (*anhydr*)] M 326.9 (6H₂O), m 47°(dec), b 100°(dec), d 4.9. Crystd from water (1mL/g) by partial evaporation in a desiccator.

Cobaltous chloride (6H₂O) [7791-13-1 (6H₂O); 7646-79-9 (*anhydr*)] M 237.9, m 87°(dec), d 1.92. A saturated aqueous soln at room temperature was fractionally crystd by standing overnight. The first half of the material that crystd in this way was used in the next crystn. The process was repeated several times, water being removed in a dry-box using air filtered through glass wool and dried over CaCl₂ [Hutchinson *J Am Chem Soc* 76 1022 1954]. Has also been crystd from dilute aq HCl.

Cobaltous nitrate (6H₂O) [10026-22-9] M 291.0, m ~55(6H₂O), 100-105°(dec), d 1.88. Crystd from water (1mL/g), or ethanol (1mL/g), by partial evapn. After 3 crystns (H₂O) it contains: metal(ppm) As (8), Fe (1.2), K (1), Mg (4), Mn (4), Mo (4), Na (0.6), Ni (18), Zn (1.6).

Cobaltous perchlorate (6H₂O) [13478-33-6] M 365.9, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from warm water (0.7mL/g) by cooling.

Cobaltous potassium sulfate [13596-22-0] M 329.4. Crystd from water (1mL/g) between 50° and 0°, and dried in a vacuum desiccator over conc H₂SO₄.

Cobaltous sulfate (7H₂O) [10026-24-1 (7H₂O); 60459-08-7 (*x*H₂O); 10124-43-3 (*anhydr*)] M 281.1, m 41°(dec), d 2.03. Crystd three times from conductivity water (1.3mL/g) between 100° and 0°.

Copper (I) thiophenolate [1192-40-1] M 172.7, m *ca* 280°, pK₁²⁵ 6.62 (for PhS⁻). The Cu salt can be extracted from a thimble (Soxhlet) with boiling MeOH. It is a green-brown powder which gives a yellow-green soln in pyridine. Wash with EtOH and dry in a vacuum. It can be pted from a pyridine soln by addition of H₂O, collect ppte, wash with EtOH and dry in a vacuum. [*Synthesis* 662 1974; *J Am Chem Soc* 79 170 1957; *Chem Ber* 90 425 1957.]

12-Crown-4 (lithium ionophore V, 1,4,7,10-tetraoxacyclododecane) [294-93-9] M 176.2, m 17°. The distilled crude product had to be crystd from pentane at -20° to remove acyclic material. It is then dried over P₂O₅. [*Acta Chem Scand* 27 3395 1973.]

Cupferon ammonium salt (*N*-nitroso-*N*-phenylhydroxylamine ammonium salt) [135-20-6] M 155.2, m 150-155°(dec), 162.5-163.5°, 163-164°, pK²⁵ 4.16 (free base). Recrystd twice from EtOH after treatment with Norite and finally once with EtOH. The crystals are washed with diethyl ether and air dried then stored in the dark over solid ammonium carbonate. A standard soln (*ca* 0.05M prepared in air-free H₂O) is prepared daily from this material for analytical work and is essentially 100% pure. [*Anal Chem* 26 1747 1954.] It can also be washed with Et₂O, dried and stored as stated. In a sealed, dark container it can be stored for at least 12 months without deterioration. λ_{max} 260nm (CHCl₃). [*Org Synth Coll Vol I* 177 1948; *J Am Chem Soc* 78 4206 1956.] Possible **CARCINOGEN**.

Cupric acetate (H₂O) [6046-93-1 (H₂O); 142-71-2 (*anhydr*)] M 199.7, m 115°, 240°(dec), d 1.88, pK₁²⁵ 8.0, pK₂²⁵ 13.1 (for Cu²⁺). Crystd twice from warm dilute acetic acid solns (5mL/g) by cooling.

Cupric ammonium chloride (2H₂O) [10534-87-9 (*hydr*); 15610-76-1 (*anhydr*)] M 277.5, m 110-120°(anhydr) then dec at higher temp, d 2.0. Crystd from weak aqueous HCl (1mL/g).

Cupric benzoate [533-01-7] M 305.8. Crystd from hot water.

Cupric bromide [7789-45-9] **M 223.4, m 498°, b 900°, d 4.7.** Crystd twice by dissolving in water (140mL/g), filtering to remove any Cu_2Br_2 , and concentrating under vac at 30° until crystals appeared. The cupric bromide was then allowed to crystallise by leaving the soln in a vac desiccator containing P_2O_5 [Hope, Otter and Prue *J Chem Soc* 5226 1960].

Cupric chloride [7447-39-4] **M 134.4, m 498°, 630°(dec).** Crystd from hot dilute aq HCl (0.6mL/g) by cooling in a CaCl_2 -ice bath. Dehydrated by heating on a steam-bath under vacuum. It is deliquescent in moist air but efflorescent in dry air.

Cupric lactate (H_2O) [814-81-3] **M 295.7.** The monohydrate crystals from hot H_2O (3mL/g) on cooling.

Cupric nitrate ($3\text{H}_2\text{O}$) [10031-43-3 ($3\text{H}_2\text{O}$); 3251-23-8 (anhydr)] **M 241.6, m 114°, b 170°(dec), d 2.0.** Crystd from weak aqueous HNO_3 (0.5mL/g) by cooling from room temperature. The anhydrous salt can be prepared by dissolving copper metal in a 1:1 mixture of liquid NO_2 and ethyl acetate and purified by sublimation [Evans et al. *J Chem Soc, Faraday Trans 1* 75 1023 1979]. The *hexahydrate* dehydr to trihydrate at 26°, and the *anhydrous salt* sublimates between 150 and 225°, but melts at 255-256° and is deliquescent.

Cupric oleate [1120-44-1] **M 626.5.** Crystd from diethyl ether.

Cupric perchlorate ($6\text{H}_2\text{O}$) [10294-46-9 (hydr); 13770-18-8] **M 370.5, m 230-240°, pK^{25} -2.4 to -3.1 (for HClO_4).** Crystd from distilled water. The *anhydrous salt* is hygroscopic.

Cupric phthalocyanine [147-14-8] **M 576.1.** Precipitated twice from conc H_2SO_4 by slow dilution with water. Also purified by two or three sublimations at 580° in an argon flow at 300-400Pa.

Cupric sulfate [7758-98-7] **M 159.6, m >560°.** After adding 0.02g of KOH to a litre of nearly saturated aq soln, it was left for two weeks, then the ppte was filtered on to a fibreglass filter with pore diameter of 5-15 microns. The filtrate was heated to 90° and allowed to evaporate until some $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ had crystd. The soln was then filtered hot and cooled rapidly to give crystals which were freed from mother liquor by filtering under suction [Geballe and Giauque *J Am Chem Soc* 74 3513 1952]. Alternatively crystd from water (0.6mL/g) between 100° and 0°.

Cupric trifluoromethylsulfonate (copper II triflate) [34946-82-2] **M 361.7, pK^{25} <-3.0 (for triflic acid).** Dissolve in MeCN, add dry Et_2O until cloudy and cool at -20° in a freezer. The light blue ppte is collected and dried in a vacuum oven at 130°/20mm for 8h. It has λ_{max} 737nm (ϵ 22.4M⁻¹cm⁻¹) in AcOH. [*J Am Chem Soc* 95 330 1973]. It has also been dried in a vessel at 0.1Torr by heating with a Fischer burner [*J Org Chem* 43 3422 1978]. It has been dried at 110-120°/5mm for 1h before use and forms a *benzene complex which should be handled in a dry box because it is air sensitive [*Chem Pharm Bull Jpn* 28 262 1980; *J Am Chem Soc* 95 330 1973].

Cuprous bromide [7787-70-4] **M 143.4, m 497°, b 1345°, d 4.72.** Purified as for cuprous iodide but using aqueous NaBr.

Cuprous bromide dimethylsulfide complex [54678-23-8] **M 205.6, m ca 135°(dec).** Purified by recrystn in the presence of Me_2S . A soln of the complex (1.02g) in Me_2S (5mL) is slowly diluted with hexane (20mL) and the pure colourless prisms of the complex (0.96g) separate and are collected and dried, **m** 124-129°dec. The complex is insoluble in hexane, Et_2O , Me_2CO , CHCl_3 and CCl_4 . It dissolves in DMF and DMSO but the soln becomes hot and green indicating dec. It dissolves in * C_6H_6 , Et_2O , MeOH and CHCl_3 if excess of Me_2S is added a colourless soln is obtained. [*J Org Chem* 40 1460 1975.] Prior to use, the complex was dissolved in Me_2S and evaporated to dryness in the weighed reaction flask [*J Organomet Chem* 228 321 1983].

Cuprous chloride [7758-89-6] **M 99.0, m 430°, b~1400°.** Dissolved in strong HCl, pted by dilution with water and filtered off. Washed with ethanol and diethyl ether, then dried and stored in a vacuum desiccator [Österlöf *Acta Chem Scand* 4 375 1950]. Alternatively, to an aq. soln of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ was added, with stirring,

an aqueous soln of anhydrous sodium sulfite. The colourless product was dried at 80° for 30min and stored under N₂. CuCl₂ can be purified by zone-refining [Hall et al. *J Chem Soc, Faraday Trans 1* **79** 243 1983].

Cuprous cyanide [544-92-3] **M 89.6, m 474°**. Wash thoroughly with boiling H₂O, then with EtOH. Dry at 100° to a fine soft powder. [*J Chem Soc* **79** 1943.]

Cuprous iodide [7681-65-4] **M 190.5, m 605°, b 1336°, d₄²⁵ 5.63**. It can be freshly prepared by dissolving an appropriate quantity of CuI in boiling saturated aqueous NaI over 30min. Pure CuI is obtained by cooling and diluting the soln with water, followed by filtering and washing sequentially with H₂O, EtOH, EtOAc and Et₂O, pentane, then drying *in vacuo* for 24h [Dieter, *J Am Chem Soc* **107** 4679 1985]. Alternatively wash with H₂O then EtOH and finally with Et₂O containing a little iodine. Traces of H₂O are best removed first by heating at 110° and then at 400°. Excess of I₂ is removed completely at 400°. It dissolves in Et₂O if an amine is present to form the amine complex. [*Chem Ind (London)* 1180 1957.]

Cuprous iodide trimethylphosphite [34836-53-8] **M 314.5, m 175-177°, 192-193°**. Cuprous iodide dissolves in a *C₆H₆ soln containing trimethylphosphite to form the complex. The complex crystallises from *C₆H₆ or pet ether. [*Chem Ber* **38** 1171 1905; *Bull Chem Soc Jpn* **34** 1177 1961.]

Cuprous thiocyanate [18223-42-2] **M 121.6, pK²⁵ -1.85 (for HSCN)**. Purified as for cuprous iodide but using aq NaSCN.

Cyanamide [420-04-2] **M 42.0, m 43°, 45°, 46°, b 85-87°/0.5mm, pK₁²⁰ -0.36 (1.1 at 29°), pK₂²⁰ 10.27**. Purified by placing *ca* 15g in a Soxhlet thimble and extracting exhaustively (2-3h) with two successive portions of Et₂O (400mL, saturated with H₂O by shaking before use) containing two drops of 1N acetic acid. Two successive portions of Et₂O are used so that the NH₂CN is not heated for too long. Each extract is dried over Na₂SO₄(30g), then combined and evaporated under reduced pressure. The NH₂CN may be stored unchanged at 0° in Et₂O soln in the presence of a trace of AcOH. Extracts from several runs may be combined and evaporated together. The residue from evaporation of an Et₂O soln is a colourless viscous oil which sets to a solid, and can be recrystd from a mixture of 2 parts of *C₆H₆ and 1 part of Et₂O. Concentrating an aqueous soln of NH₂CN at high temps causes **EXPLOSIVE** polymerisation. [*Org Synth Coll Vol IV* 645 1963; *Inorg Synth* **3** 39 1950; *J Org Chem* **23** 613 1958.] *Hygroscopic*.

Cyanogen bromide [506-68-3] **M 105.9, m 49-51°, b 60-62°/atm**. *All operations with this substance should be performed in a very efficient fume cupboard - it is very POISONOUS and should be handled in small amounts. Fresh commercial material is satisfactory for nearly all purposes and does not need to be purified. It is a white crystalline solid with a strong cyanide odour. If it is reddish in colour and partly liquid or paste-like then it is too far gone to be purified, and fresh material should be sought.* It can be purified by distn using small amounts at a time, and using a short wide-bore condenser because it readily solidifies to a crystalline white solid and may clog the condenser. *An appropriate gas mask should be used when transferring the molten solid from one container to another and the operation should be done in an efficient fume cupboard.* The melting point (m 49-51°) should be measured in a sealed tube. [*Org Synth Coll Vol II* 150 1948.]

Cyanogen iodide [506-78-5] **M 152.9, m 146-147°**. *This compound is POISONOUS and the precautions for cyanogen bromide (above) apply here.* The reagent (*ca* 5.9g) is dissolved in boiling CHCl₃ (15mL), filtered through a plug of glass wool into a 25mL Erlenmeyer flask. Cool to room temperature for 15min, then place in an ice-salt bath and cool to -10°. This cooling causes a small aqueous layer to separate as ice. The ice is filtered with the CNI, but melts on the filter and is also removed with the CHCl₃ used as washing liquid. The CNI which is collected on a sintered glass funnel is washed 3x with CHCl₃ (1.5mL at 0°) and freed from last traces of solvent by being placed on a watch glass and exposed to the atmosphere in a good fume cupboard at room temp for 1h to give colourless needles (*ca* 4.5g), m 146-147° (sealed capillary totally immersed in the oil bath). The yield depends slightly on the rapidity of the operation, in this way loss by sublimation can be minimised. If desired, it can be sublimed under reduced pressure at temps at which CNI is only slowly decomposed into I₂ and (CN)₂. The vacuum will need to be renewed constantly due to the volatility of CNI. [*Org Synth* **32** 29 1952.]

Decaborane [17702-41-9] M 122.2, m 99.7-100°. Purified by vacuum sublimation at 80°/0.1mm, followed by crystn from methylcyclohexane, CH₂Cl₂, or dry olefin-free-*n*-pentane, the solvent being subsequently removed by storing the crystals in a vacuum desiccator containing CaCl₂.

Deuterium [7782-39-0] M 4. Passed over activated charcoal at -195° [MacIver and Tobin *J Phys Chem* 64 451 1960]. Purified by diffusion through nickel [Pratt and Rogers, *J Chem Soc, Faraday Trans I* 92 1589 1976]. Always check deuterium for radioactivity to find out the amount of tritium in it (see D₂O below).

Deuterium oxide [7789-20-0] M 20, f 3.8°/760mm, b 101.4°/760mm, d 1.105. Distd from alkaline KMnO₄ [de Giovanni and Zamenhof *Biochem J* 92 79 1963]. NOTE that D₂O invariably contains tritiated water and will therefore be RADIOACTIVE; always check the radioactivity of D₂O in a scintillation counter before using.

cis-Diamminedichloroplatinum(II) (Cisplatin) [15663-27-1] M 300.1, m 270°(dec). Recrystd from dimethylformamide and the purity checked by IR and UV-VIS spectroscopy. [Raudaschl et al. *Inorg Chim Acta* 78 143 1983.] HIGHLY TOXIC, SUSPECTED CARCINOGEN.

Diammonium hydrogen orthophosphate [7783-28-0] M 132.1. Crystd from water (1mL/g) between 70° and 0°. After one crystn of ACS grade had Fe, Mo, Na, Se and Ti at 1, 0.2, 1.4, 0.2 and 0.8ppm resp.

Di-*n*-amyl *n*-amylphosphonate [6418-56-0] M 292.4, b 150-151°/2mm, n 1.4378. Purified by three crystns of its uranyl nitrate complex from hexane (see *tributyl phosphate*). Extracts Zr²⁺ from NaCl solns.

6,6-Dibenzyl-14-crown-4 (lithium ionophore VI; 6,6-dibenzyl-1,4,8,11-tetra-oxa-cyclo-tetradecane) [106868-21-7] M 384.5, m 102-103°. Dissolve in CHCl₃, wash with saturated aqueous NaCl, dry with MgSO₄, evaporate and purify by chromatography on silica gel and gradient elution with *C₆H₆-MeOH followed by preparative reverse phase HPLC on an octadecyl silanised silica (ODS) column and eluting with MeOH. It can be crystd from MeOH (ν_{CBr} 1120 cm⁻¹, C-O-C). [*J Chem Soc Perkin Trans I* 1945 1986.]

Di-*n*-butyl boron triflate (di-*n*-butylboryl trifluoromethanesulfonate) [60669-69-4] M 274.1, b 37°/0.12mm, 60°/2mm, pK²⁵ <-3.0 (for triflic acid). Distil in vacuum under argon and store under argon. Should be used within 2 weeks of purchase or after redistn. Use a short path distn system. It has IR bands in CCl₄ at ν 1405, 1380, 1320, 1200 and 1550cm⁻¹; and ¹³C NMR (CDCl₃) with δ at 118.1, 25.1, 21.5 and 13.6. [*Org Synth* 68 83 1990; *J Am Chem Soc* 103, 3099 1981.] TOXIC

Di-*n*-butyl cyclohexylphosphonate [1085-92-3] M 245.4. The compound with uranyl nitrate was crystd three times from hexane. For method see *tributyl phosphate*.

Di-*tert*-butyl dichlorosilane (DTBCl₂) [18395-90-9] M 213.2, m -15°, b 190°/729mm, 195-197°/atm, d 1.01. Purified by fractional distn. It is a colourless liquid with a pleasant odour and does not fume in moist air, but does not titrate quantitatively with excess of dil alkali. [*J Am Chem Soc* 70 2877 1948.]

Di-*n*-butyl *n*-butylphosphonate [78-46-6] M 250.3, b 150-151°/10mm, 160-162°/20mm, n²⁵ 1.4302. Purified by three recrystallisations of its compound with uranyl nitrate, from hexane. For method, see *tributyl phosphate*.

Di-*tert*-butyl silyl bis(trifluoromethanesulfonate) [85272-31-7] M 440.5, b 73.5-74.5°/0.35mm, d 1.36 (see pK for triflic acid). Purified by fractional distillation. It is a pale yellow liquid which should be stored under argon. It is less reactive than the diisopropyl analogue. The presence of the intermediate monochloro compound can be detected by ¹H NMR, (CHCl₃): *tert*-Bu₂Si(OTf)₂ [δ 1.25s]; but

impurities have δ 1.12s for *tert*-Bu₂Si(H)OTf and δ 1.19s for *tert*-Bu₂HSi(Cl)OTf. [*Tetrahedron Lett* **23** 487 1982.] **TOXIC.**

Di-*n*-butyltin oxide [818-08-6] **M 248.9, m >300°.** It is prepd by hydrolysis of di-*n*-butyltin dichloride with KOH. Hence wash with a little aq M KOH then H₂O and dry at -80°/10mm until the IR is free from OH bands. [Cummings *Aust J Chem* **18** 98 1965.]

Dicarbonyl(cyclopentadienyl)Co (I) [1207-25-0] **M 180.1, b 75°/22mm, b 139-140°(dec)/710mm.** Best distd in an atmosphere of CO in a vac. The red brown liquid decomposes slightly on distn even in a vac to liberate some CO. Operations should be performed in an efficient fume cupboard. It is sol in organic solvents and stable in air but decomposes slowly in sunlight and rapidly under UV. [Piper et al. *J Inorg Nucl Chem* **1** 165 1955.] **TOXIC.**

Dichlorodimethylsilane see dimethylchlorosilane p. 419.

2,6-Dichlorophenol-indophenol sodium salt (2H₂O) [620-45-1] **M 326.1, ϵ 2.1 x 10⁴ at 600nm and pH 8, pK³⁰ 5.7 (oxidised form), pK₁³⁰ 7.0, pK₂³⁰ 10.1 (reduced form).** Dissolved in 0.001M phosphate buffer, pH 7.5 (alternatively, about 2g of the dye was dissolved in 80mL of M HCl), and extracted into diethyl ether. The extract was washed with water, extracted with aqueous 2% NaHCO₃, and the sodium salt of the dye was ppted by adding NaCl (30g/100mL of NaHCO₃ soln), then filtered off, washed with dilute NaCl soln and dried.

Dicobalt octacarbonyl [10210-68-1] **M 341.9, m 51°.** Orange-brown crystals by recrystn from *n*-hexane under a carbon monoxide atmosphere [Ojima et al. *J Am Chem Soc* **109** 7714 1987; see also Hileman in *Preparative Inorganic Reactions*, Jolly Ed. Vol 1 101 1987].

Diethyl aluminium chloride [96-10-6] **M 120.6, m -75.5°, b 106.5-108°/24.5mm, d 0.96.** Distd from excess dry NaCl (to remove ethyl aluminium dichloride) in a 50-cm column containing a heated nichrome spiral.

***O,O*-Diethyl-*S*-2-diethylaminoethyl phosphorothiolate** [78-53-5] **M 269.3, m 98-99°.** Crystd from isopropanol/diethyl ether.

Di-(2-ethylhexyl)phosphoric acid ('diisooctyl' phosphate) [27215-10-7; 298-07-7] **M 322.4, b 209°/10mm, d 0.965, pK_{Est} ~1.7.** Contaminants of commercial samples include the monoester, polyphosphates, pyrophosphate, 2-ethylhexanol and metal impurities. Dissolved in *n*-hexane to give an 0.8M soln. Washed with an equal volume of M HNO₃, then with saturated (NH₄)₂CO₃ soln, with 3M HNO₃, and twice with water [Petrov and Allen *Anal Chem* **33** 1303 1961]. Similarly, the impure sodium salt, after scrubbing with pet ether, was acidified with HCl and the free organic acid was extracted into pet ether and purified as above. [Peppard et al. *J Inorg Nucl Chem* **7** 231 1958] or Stewart and Crandall [*J Am Chem Soc* **73** 1377 1951]. Purified also *via* the copper salt [McDowell et al. *J Inorg Nucl Chem* **38** 2127 1976].

Diethylmethylsilane [760-32-7] **M 102.3, b 78.4°/760mm, 77.2-77.6°/atm, d 0.71.** Fractionally distilled through a *ca* 20 plate column and the fraction boiling within a range of less than 0.5° is collected. [*Izv Akad Nauk SSSR Otd Khim* 1416 1957; *J Am Chem Soc* **69** 2600 1947.]

Diethyl trimethylsilyl phosphite [13716-45-5] **M 210.3, b 61°/10mm, 66°/15mm, d 0.9476, n 1.4113.** Fractionated under reduced pressure and has δ_p -128 ± 0.5 relative to H₃PO₄. [*J Org Chem* **46** 2097 1981; *J Gen Chem USSR (Engl Transl)* **45** 231 1975.]

***N,N*-Diethyltrimethylsilylamine** [996-50-9] **M 145.3, b 33°/26mm, 126.8-127.1°/738mm, 126.1-126.4°, d 0.763, n 1.411.** Fractionated through a 2ft vac-jacketed column containing Helipak packing with a reflux ratio of 10:1. [*J Am Chem Soc* **68** 241 1946; *J Org Chem* **23** 50 1958; *J Prakt Chem* **9** 315 1959.]

***N,N'*-Diheptyl-*N,N'*-5,5-tetramethyl-3,7-dioxanonanediamide [lithium ionophore I (ETH 149)]** [58821-96-8] **M 442.7**. Purified by chromatography on Kieselgel using CHCl_3 as eluent (IR ν 1640cm^{-1}). [*Helv Chim Acta* **60** 2326 1977.]

Dihexadecyl phosphate [2197-63-9] **M 546.9**, **m 75°** , **$\text{pK}_{\text{Est}} \sim 1.2$** . Crystd from MeOH [Lukac *J Am Chem Soc* **106** 4387 1984].

1,2-Dihydroxybenzene-3,5-disulfonic acid, di-Na salt (TIRON) [149-45-1] **M 332.2**, ϵ **6.9×10^4** at **260nm**, **pH 10.8**, **pK_1 and $\text{pK}_2 < 2$** (for SO_3^-), **pK_3 7.7**, **pK_4 12.6** (for OHs of disulfonate dianion). Recrystd from water [Hamaguchi et al. *Anal Chim Acta* **9** 563 1962]. Indicator color reagent for Fe, Mn, Ti and Mo ions and complexes with Al, Cd, Co, Co, Fe (III), Mn, Pd, UO_2^{2+} , VO^{2+} and Zn.

Diisooctyl phenylphosphonate [49637-59-4] **M 378.5**, **n^{25} 1.4780**. Vacuum distilled, percolated through a column of alumina, then passed through a packed column maintained at 150° to remove residual traces of volatile materials in a countercurrent stream of N_2 at reduced pressure [Dobry and Keller *J Phys Chem* **61** 1448 1957].

Diisopropyl chlorosilane (chlorodiisopropylsilane) [2227-29-4] **M 150.7**, **b $59^\circ/8\text{mm}$** , **$80^\circ/10\text{mm}$** , **$200^\circ/738\text{mm}$** , **d 0.9008**, **n 1.4518**. Impurities can be readily detected by ^1H NMR. Purified by fractional distn [*J Am Chem Soc* **69** 1499 1947; *J Chem Soc* 3668 1957; *J Organometal Chem* **282** 175 1985].

Dilongifolyl borane [77882-24-7] **M 422.6**, **m $169-172^\circ$** . Wash with dry Et_2O and dry in a vacuum under N_2 . It has **m $160-161^\circ$** in a sealed evacuated capillary. It is sparingly soluble in pentane, tetrahydrofuran, carbon tetrachloride, dichloromethane, and chloroform, but the suspended material is capable of causing asymmetric hydroboration. Disappearance of solid indicates that the reaction has proceeded. [*J Org Chem* **46** 2988 1981.]

Dimethyl carbonate [616-38-6] **M 90.1**, **b $89.5^\circ/755\text{mm}$** , **$90.2^\circ/\text{atm}$** , **d 1.0446**, **n 1.3687**. If the reagent has broad intense bands at 3300cm^{-1} and above (i.e. OH stretching) then it should be purified further. Wash successively with 10% Na_2CO_3 soln, saturated CaCl_2 , H_2O and dried by shaking mechanically for 1h with anhydrous CaCl_2 , and fractionated. [*J Chem Soc* **78** 1939, 1847 1948.]

Dimethyl dicarbonate (dimethyl pyrocarbonate) [4525-33-1] **M 134.1**, **m 15.2°** , **b $45-46^\circ/5\text{mm}$** , **d 1.2585**, **n 1.3950**. Dissolve in Et_2O , shake with a small vol of 0.1N HCl, dry Et_2O with Na_2SO_4 and distil in vac below 100° to give a clear liquid. It dec to CO_2 and dimethyl carbonate on heating at $123-149^\circ$. It is readily hydrolysed by H_2O and is an **IRRITANT**. [*J Gen Chem USSR* **22** 1546 1952; see also *Chem Ber* **71** 1797 1938.]

Dimethyldichlorosilane [75-78-5] **M 129.1**, **m -75.5°** , **b $68.5-68.7^\circ/750\text{mm}$** , **$70.5^\circ/760\text{mm}$** , **d 1.0885**, **n 1.4108**. Other impurities are chlorinated silanes and methylsilanes. Fractionated through a 3/8in diameter 7ft Stedman column rated at 100 theoretical plates at almost total reflux (see p. 441). See purification of MeSiCl_2 . [*J Am Chem Soc* **70** 3590 1948.]

2,6-Dimethyl-1,10-phenanthrolinedisulfonic acid, di-Na salt (H_2O) (bathocuproine-disulfonic acid di-Na salt) [52698-84-7] **M 564.5**, **$\text{pK}_{\text{Est}} \sim 0$** (for free acid). Inorganic salts and some coloured species can be removed by dissolving the crude material in the minimum volume of water and precipitating by adding EtOH. Purified reagent can be obtained by careful evapn of the filtrate.

Dimethylphenylsilyl chloride (chlorodimethylphenylsilane) [768-33-2] **M 170.7**, **b $85-87^\circ/32\text{mm}$** , **$196^\circ/\text{atm}$** , **d 1.017**, **n 1.509**. See phenyl methyl chlorosilane on p. 449.

Dinitrogen tetroxide (nitrogen dioxide, N_2O_4) [10544-72-6] **M 92.0 m -11.2°, b 21.1°**. Purified by oxidation at 0° in a stream of oxygen until the blue colour changed to red-brown. Distd from P_2O_5 , then solidified on cooling in a deep-freeze (giving nearly colourless crystals). Oxygen can be removed by alternate freezing and melting. **TOXIC VAPOUR.**

Diocetyl phenylphosphonate [1754-47-8] **M 378.8, d 1.485, n^{25} 1.4780**. Purified as described under diisooctyl phenylphosphonate.

(1,3-Dioxalan-2-ylmethyl)triphenylphosphonium bromide [52509-14-5] **M 429.3, m 191.5-193°, 193-195°**. Wash the crystals with Et_2O , dry in a vac and recryst from CH_2Cl_2 -dry Et_2O to give prisms **m 172-174°**, which is raised to 191.5-193°, on drying at 56°/0.5mm. [Cresp et al. *J Chem Soc, Perkin Trans 1* 37 1974.]

Diphenyldiselenide [1666-13-3] **M 312.1, m 62-64**. Crystd twice from hexane [Kice and Purkiss *J Org Chem* 52 3448 1987].

Diphenyl hydrogen phosphate [838-85-7] **M 250.2, m 99.5°, pK^{20} 0.26**. Crystd from $CHCl_3$ /pet ether.

Diphenylmercury [587-85-9] **M 354.8, m 125.5-126°**. Sublimed, then crystd from nitromethane or ethanol. If phenylmercuric halides are present they can be converted to phenylmercuric hydroxide which, being much more soluble, remains in the alcohol or *benzene used for crystn. Thus, crude material (10g) is dissolved in warm ethanol (ca 150mL) and shaken with moist Ag_2O (ca 10g) for 30min, then heated under reflux for 30min and filtered hot. Concentration of the filtrate by evaporation gives diphenylmercury, which is recrystd from *benzene [Blair, Bryce-Smith and Pengilly *J Chem Soc* 3174 1959]. **TOXIC.**

4,7-Diphenyl-1,10-phenanthroline-disulfonic acid, di-Na salt $3H_2O$ (bathophenanthroline-disulfonic acid di-Na salt) [52746-49-3] **M 590.6, m 300°, $pK_{Est} \sim 0$ (for free acid)**. Dissolve crude sample in the minimum volume of water and add $EtOH$ to ppt the contaminants. Carefully evaporate the filtrate to obtain pure material.

It forms a dark red complex with Fe^{2+} with λ_{max} 535nm (ϵ 2.23 x $10^4 mol^{-1} cm^{-1}$) [*Anal Chim Acta* 115 407 1980]. Prepared by sulfonating bathophenanthroline with $ClSO_3H$: to 100g of bathophenanthroline was added 0.5mL of Fe free $ClSO_3H$ and heated over a flame for 30sec. Cool and carefully add 10mL of pure distd H_2O and warm on a water bath with stirring till all solid dissolved. A stock soln is made by diluting 3mL of this reagent to 100mL with 45% aq $NaOAc$, filter off the solid and store in a dark bottle. In this way it is stable for several months. [*Am J Clinical Pathology* 29 590 1958.]

Diphenylphosphinic acid [1707-03-5] **M 218.2, m 194-195°, pK^{20} 1.72**. Recrystd from 95% $EtOH$ and dried under vacuum at room temperature. [see Kosolapoff *Organophosphorus Compounds* J Wiley, NY, 1950; Kosolapoff and Maier *Organic Phosphorus Compounds* Wiley-Interscience, NY, 1972-1976.]

Diphenylsilane [775-12-2] **M 184.3, b 75-76°/0.5mm, 113-114°/9mm, 124-126°/11mm, 134-135°/16mm, d 1.0027, n 1.5802, 1.5756**. Dissolve in Et_2O , mix slowly with ice-cold 10% $AcOH$. The Et_2O layer is then shaken with H_2O until the washings are neutral to litmus. Dry over Na_2SO_4 , evaporate the Et_2O and distil the residual oil under reduced pressure using a Claisen flask with the take-off head modified into a short column. Ph_2SiH_2 boils at 257°/760mm but it cannot be distd at this temp because exposure to air leads to flashing, decomposition and formation of silica. It is a colourless, odourless oil, miscible with organic solvents but not H_2O . A possible impurity is Ph_3SiH which has **m 43-45°** and would be found in the residue. [*J Org Chem* 18 303 1953 ; *J Am Chem Soc* 74 6481952, 81 5925 1959.]

Diphenylsilanediol [947-42-2] **M 216.3, m 148°(dec)**. Crystd from $CHCl_3$ -methyl ethyl ketone.

Diphenyl tolyl phosphate [26444-49-5] **M 340.3, n^{25} 1.5758**. Vac distd, then percolated through a column of alumina. Finally, passed through a packed column maintained at 150° to remove traces of volatile

impurities in a countercurrent stream of nitrogen at reduced pressure. [Dobry and Keller *J Phys Chem* **61** 1448 1947.]

Disodium calcium ethylenediaminetetraacetate [39208-14-5] **M 374.3**, (see pKs for EDTA in entry below). Dissolved in a small amount of water, filtered and ppted with excess EtOH. Dried at 80°.

Disodium dihydrogen ethylenediaminetetraacetic acid (2H₂O) [6381-92-6] **M 372.2**, **m 248°(dec)**, **pK₁^{2.5} 0.26**, **pK₂^{2.5} 0.96**, **pK₃^{2.5} 2.60**, **pK₄^{2.5} 2.67**, **pK₅^{2.5} 6.16**, **pK₆^{2.5} 10.26** (see EDTA Cha 4). Analytical reagent grade material can be used as primary standard after drying at 80°. Commercial grade material can be purified by crystn from water or by preparing a 10% aqueous soln at room temperature, then adding ethanol slowly until a slight permanent ppt is formed, filtering, and adding an equal volume of ethanol. The ppt is filtered off on a sintered-glass funnel, is washed with acetone, followed by diethyl ether, and dried in air overnight to give the dihydrate. Drying at 80° for at least 24h converts it to the anhydrous form.

Disodium 4,5(1,8)-dihydroxynaphthalene-2,7(3,6)-disulfonate (2H₂O) [5808-22-0] **M 400.3**, **m >300°**, **pK₁ 0.61(SO₃⁻)**, **pK₂ 0.7(SO₃⁻)**, **pK₃ 5.45(OH)**, **pK₄ 15.5(OH)**. Crystd from H₂O or H₂O by addition of EtOH. Complexes with Ag, ClO₃⁻, Cr, Hg, NO₂⁻, NO₃⁻ and Ti. [cf Chromotropic acid p. 413.]

Disodium ethylenebis[dithiocarbamate] [142-59-6] **M 436.5**, **pK_{Est} ~ 3.0**. Crystd (as hexahydrate) from aqueous ethanol.

Disodium-β-glycerophosphate [819-83-0 (4H₂O)] **M 216.0**, **m 102-104°**, **pK₂^{2.5} 6.66** (free acid). Crystd from water.

Disodium hydrogen orthophosphate (anhydrous) [7558-79-4] **M 142.0**, (see pK of H₃PO₄). Crystd twice from warm water, by cooling. Air dried, then oven dried overnight at 130°. *Hygroscopic*: should be dried before use.

Disodium magnesium ethylenediaminetetraacetate [14402-88-1] **M 358.5**, **pK₁^{2.5} 0.26**, **pK₂^{2.5} 0.96**, **pK₃^{2.5} 2.60**, **pK₄^{2.5} 2.67**, **pK₅^{2.5} 6.16**, **pK₆^{2.5} 10.26** (see EDTA on p. 237 in Chapter 4). Dissolved in a small amount of water, filtered and ppted with an excess of MeOH. Dried at 80°.

Disodium naphthalene-1,5-disulfonate [1655-29-4] **M 332.3**, **pK_{Est} ~0**. Recrystd from aqueous acetone [Okahata et al. *J Am Chem Soc* **108** 2863 1986].

Disodium 4-nitrophenylphosphate (6H₂O) [4264-83-9] **M 371.1** Dissolve in hot aqueous MeOH, filter and ppt by adding Me₂CO. Wash the solid with Me₂CO and repeat the purification. Aq MeOH and Et₂O can also be used as solvents. The white fibrous crystals contain less than 1% of free 4-nitrophenol [assay: *J Biol Chem* **167** 57 1947].

Disodium phenylphosphate (2H₂O) [3279-54-7] **M 254.1**, **pK₁^{2.5} 1.46**, **pK₂^{2.5} 6.29** [for PhPO(OH)₂]. Dissolved in a minimum amount of methanol, filtering off an insoluble residue of inorganic phosphate, then ppted by adding an equal volume of diethyl ether. Washed with diethyl ether and dried [Tsuboi *Biochim Biophys Acta* **8** 173 1952].

Disodium succinate [150-90-3] **M 162.1**. Crystd twice from water (1.2mL/g) and dried at 125°. Freed from other metal ions by passage of a 0.1M soln through a column of Dowex resin A-1 (Na form).

Di-p-tolylmercury [50696-65-6] **M 382.8**, **m 244-246°**. Crystd from xylene.

Di-p-tolyl phenylphosphonate [94548-75-1] **M 388.3**, **n²⁵ 1.5758**. Purified as described under diisooctyl phenylphosphonate.

1,3-Divinyl-1,1,3,3-tetramethyldisiloxane [2627-95-4] M 186.4, m -99.7°; b 128-129°/atm, 139°/760mm, d 0.811, n 1.4122. Dissolve in Et₂O, wash with H₂O, dry over CaCl₂ and distil. [*J Am Chem Soc* 77 1685 1955; *Collect Czech Chem Comm* 24 3758 1959.]

Eosin B (Bluish, Eosin Scarlet, 4',5'-dibromo-2',7'-dinitrofluorescein disodium salt) [548-24-3] M 624.1, λ_{\max} 514nm, CI 45400. Freed from inorganic halides by repeated crystn from butan-1-ol.

Eosin Y (as di-Na salt) [17372-87-1] M 691.9. Dissolved in water and ppted by addition of dilute HCl. The ppt was washed with water, crystd from ethanol, then dissolved in the minimum amount of dilute NaOH soln and evaporated to dryness on a water-bath. The purified disodium salt was then crystd twice from ethanol [Parker and Hatchard *Trans Faraday Soc* 57 1894 1961].

Eosin YS (Eosin Yellowish, 2',4',5'7'-tetrabromofluorescein di-Na salt) [17372-87-1] M 691.9, CI 45380. Dissolve in the minimum vol of H₂O (1g/mL), filter and add EtOH until separation of salt is complete. Filter off, wash with abs EtOH, then Et₂O and dry first in air, then at 100°. Used for staining blood cells and for estimating traces of Ag. [Selsted and Becker *Anal Biochem* 155 270 1986; El-Ghamry and Frei *Anal Chem* 40 1986 1968.]

Eriochrome Black T [1787-61-7] M 416.4, $A_{1\text{cm}}^{\%}$ (λ_{\max}) 656(620nm) at pH 10, using the dimethylammonium salt, pK_2^{25} 5.81, pK_3^{25} 11.55. The sodium salt (200g) was converted to the free acid by stirring with 500mL of 1.5M HCl, and, after several minutes, the slurry was filtered on a sintered-glass funnel. The process was repeated and the material was air dried after washing with acid. It was extracted with *benzene for 12h in a Soxhlet extractor, then the *benzene was evaptd and the residue was air dried. A further desalting with 1.5M HCl (1L) was followed by crystn from dimethylformamide (in which it is very soluble) by forming a saturated soln at the boiling point, and allowing to cool slowly. The crystalline dimethylammonium salt so obtained was washed with *benzene and treated repeatedly with dilute HCl to give the insoluble free acid which, after air drying, was dissolved in alcohol, filtered and evaporated. The final material was air dried, then dried in a vacuum desiccator over Mg(ClO₄)₂ [Diehl and Lindstrom, *Anal Chem* 31 414 1959]. Indicator for complexometry of alkaline earth metals.

Eriochrome Blue Black R (Palatine Chrome Black 6BN, Calcon, 3-hydroxy-4-(2-hydroxy-1-naphthylazo)naphthalene-1-sulfonic acid Na salt) [2538-85-4] M 416.4, pK_2^{25} 7.0, pK_3^{25} 13.5. Freed from metallic impurities by three pptns from aqueous soln by addn of HCl. The ppted dye was dried at 60° under vacuum. Indicator for complexometry of Al, Fe and Zr.

Ethoxycarbonylmethylene triphenylphosphonium bromide [1530-45-6] M 429.3, m 155-155.5°, 158°(dec). Wash with pet ether (b 40-50°) and recryst from CHCl₃/Et₂O and dry in high vac at 65°. [Isler et al. *Helv Chim Acta* 40 1242 1957; Wittig and Haag *Chem Ber* 88 1654, 1664 1955.]

(Ethoxycarbonylmethylene)triphenylphosphorane [ethyl (triphenylphosphoranylidene)-acetate] [1099-45-2] M 348.4, m 116-117°, 128-130°. Cryst by dissolving in AcOH and adding pet ether (b 40-50°) to give colorless plates. UV λ_{\max} ($A_{1\text{mm}}^1$): 222nm (865) and 268nm (116) [Isler et al. *Helv Chim Acta* 40 1242 1957].

Ethylarsonic acid [507-32-4] M 154.0, m 99.5°, pK_1 4.72 (As(OH)O⁻), pK_2 8.00 [AsO₂²⁻]. Crystd from ethanol.

2-Ethyl-1,2-benzisoxazolium tetrafluoroborate [4611-62-5] M 235.0, m 107-109°, 109.5-110.2°. Recrystd from MeCN-EtOAc to give magnificent crystals. It is not hygroscopic but on long exposure to moisture it etches glass. It is light-sensitive and should be stored in brown glass bottles. UV (H₂O), λ_{\max} 258nm (ϵ 13 100) and λ_{\max} 297nm (ϵ 2 900); IR (CH₂Cl₂): 1613 (C=N) and 1111-1000 (BF₄⁻) [UV, IR, NMR: Kemp and Woodward *Tetrahedron* 21 3019 1965].

[Previous Page](#)

1,3-Divinyl-1,1,3,3-tetramethyldisiloxane [2627-95-4] M 186.4, m -99.7°; b 128-129°/atm, 139°/760mm, d 0.811, n 1.4122. Dissolve in Et₂O, wash with H₂O, dry over CaCl₂ and distil. [*J Am Chem Soc* 77 1685 1955; *Collect Czech Chem Comm* 24 3758 1959.]

Eosin B (Bluish, Eosin Scarlet, 4',5'-dibromo-2',7'-dinitrofluorescein disodium salt) [548-24-3] M 624.1, λ_{\max} 514nm, CI 45400. Freed from inorganic halides by repeated crystn from butan-1-ol.

Eosin Y (as di-Na salt) [17372-87-1] M 691.9. Dissolved in water and ppted by addition of dilute HCl. The ppte was washed with water, crystd from ethanol, then dissolved in the minimum amount of dilute NaOH soln and evaporated to dryness on a water-bath. The purified disodium salt was then crystd twice from ethanol [Parker and Hatchard *Trans Faraday Soc* 57 1894 1961].

Eosin YS (Eosin Yellowish, 2',4',5',7'-tetrabromofluorescein di-Na salt) [17372-87-1] M 691.9, CI 45380. Dissolve in the minimum vol of H₂O (1g/mL), filter and add EtOH until separation of salt is complete. Filter off, wash with abs EtOH, then Et₂O and dry first in air, then at 100°. Used for staining blood cells and for estimating traces of Ag. [Selsted and Becker *Anal Biochem* 155 270 1986; El-Ghamry and Frei *Anal Chem* 40 1986 1968.]

Eriochrome Black T [1787-61-7] M 416.4, $A_{1\text{cm}}^{\%}$ (λ_{\max}) 656(620nm) at pH 10, using the dimethylammonium salt, $\text{pK}_2^{2.5}$ 5.81, $\text{pK}_3^{2.5}$ 11.55. The sodium salt (200g) was converted to the free acid by stirring with 500mL of 1.5M HCl, and, after several minutes, the slurry was filtered on a sintered-glass funnel. The process was repeated and the material was air dried after washing with acid. It was extracted with *benzene for 12h in a Soxhlet extractor, then the *benzene was evaptd and the residue was air dried. A further desalting with 1.5M HCl (1L) was followed by crystn from dimethylformamide (in which it is very soluble) by forming a saturated soln at the boiling point, and allowing to cool slowly. The crystalline dimethylammonium salt so obtained was washed with *benzene and treated repeatedly with dilute HCl to give the insoluble free acid which, after air drying, was dissolved in alcohol, filtered and evaporated. The final material was air dried, then dried in a vacuum desiccator over Mg(ClO₄)₂ [Diehl and Lindstrom, *Anal Chem* 31 414 1959]. Indicator for complexometry of alkaline earth metals.

Eriochrome Blue Black R (Palatine Chrome Black 6BN, Calcon, 3-hydroxy-4-(2-hydroxy-1-naphthylazo)naphthalene-1-sulfonic acid Na salt) [2538-85-4] M 416.4, $\text{pK}_2^{2.5}$ 7.0, $\text{pK}_3^{2.5}$ 13.5. Freed from metallic impurities by three pptns from aqueous soln by addn of HCl. The ppted dye was dried at 60° under vacuum. Indicator for complexometry of Al, Fe and Zr.

Ethoxycarbonylmethylene triphenylphosphonium bromide [1530-45-6] M 429.3, m 155-155.5°, 158°(dec). Wash with pet ether (b 40-50°) and recryst from CHCl₃/Et₂O and dry in high vac at 65°. [Isler et al. *Helv Chim Acta* 40 1242 1957; Wittig and Haag *Chem Ber* 88 1654, 1664 1955.]

(Ethoxycarbonylmethylene)triphenylphosphorane [ethyl (triphenylphosphoranylidene)-acetate] [1099-45-2] M 348.4, m 116-117°, 128-130°. Cryst by dissolving in AcOH and adding pet ether (b 40-50°) to give colorless plates. UV λ_{\max} ($A_{1\text{mm}}^{\%}$): 222nm (865) and 268nm (116) [Isler et al. *Helv Chim Acta* 40 1242 1957].

Ethylarsonic acid [507-32-4] M 154.0, m 99.5°, pK_1 4.72 (As(OH)O⁻), pK_2 8.00 [AsO₂²⁻]. Crystd from ethanol.

2-Ethyl-1,2-benzisoxazolium tetrafluoroborate [4611-62-5] M 235.0, m 107-109°, 109.5-110.2°. Recrystd from MeCN-EtOAc to give magnificent crystals. It is not hygroscopic but on long exposure to moisture it etches glass. It is light-sensitive and should be stored in brown glass bottles. UV (H₂O), λ_{\max} 258nm (ϵ 13 100) and λ_{\max} 297nm (ϵ 2 900); IR (CH₂Cl₂): 1613 (C=N) and 1111-1000 (BF₄⁻) [UV, IR, NMR: Kemp and Woodward *Tetrahedron* 21 3019 1965].

Ethylene bis(diphenylphosphine) [1,2-bis(diphenylphosphino)ethane] [1663-45-2] **M 398.4, m 139-140°**. See 1,2-bis-(diphenylphosphino)ethane (DIPHOS) on p. 402.

Ethylmercuric chloride [107-27-7] **M 265.1, m 193-194°**. Mercuric chloride can be removed by suspending ethylmercuric chloride in hot distilled water, filtering with suction in a sintered-glass crucible and drying. Then crystd from ethanol and sublimed under reduced pressure. It can also be crystd from water.

Ethylmercuric iodide [2440-42-8] **M 356.6, m 186°**. Crystd once from water (50mL/g).

Ethyl Orange (sodium 4,4'-diethylaminophenylazobenzenesulfonate) [62758-12-7] **M 355.4, pK_{Est} ~ 3.8**. Recrystd twice from water.

Ethyl trimethylsilylacetate [4071-88-9] **M 160.3, b 74.5°/41mm, 75.5°/42 mm, 157°/730mm, d 0.8762, n 1.4149**. Purified by distilling *ca* 10g of reagent through a 15cm, Vigreux column and then redistilling through a 21cm glass helices-packed column [J Am Chem Soc 75 994 1953]. Alternatively, dissolve in Et₂O, wash with H₂O, dilute Na₂CO₃, dry over Na₂CO₃, evaporate Et₂O, and distil through a column of 15 theoretical plates [J Am Chem Soc 70 2874 1948].

Ethyl 3-(trimethylsilyl)propionate [17728-88-0] **M 174.3, b 93°/40mm, 178°-180°/atm, d 0.8763, n 1.4198**. Dissolve in Et₂O, wash with H₂O, dilute Na₂CO₃, dry over Na₂SO₄, evaporate Et₂O and fractionally distil. [J Am Chem Soc 72 1935 1950.]

Ethynyl tributylstannane [994-89-8] **M 315.1, b 76°/0.2mm, 130-135°/0.7mm, 200°/2mm, d 1.1113, n 1.4770**. Purified by dissolving the reagent (*ca* 50g) in heptane (250mL), washing with H₂O (100mL), drying (MgSO₄), evaporating and distilling in a vacuum. It has IR ν 3280 (\equiv C-H), 2950, 2850, 2005 (C \equiv C), 1455, 1065 and 865cm⁻¹. [J Org Chem 46 5221 1981; J Am Chem Soc 109 2138 1987; J Gen Chem USSR (Engl Edn) 37 1469 1967.]

Ethynyl trimethylsilane [1066-54-2] **M 98.2, b 53°/atm, 52.5°/atm, d 0.71, n 1.3871**. Distil through an efficient column. The IR has bands at 2041 (C \equiv C) and 3289 (\equiv C-H) cm⁻¹. [Chem Ber 92 30 1959.]

Ethyl triphenylphosphonium bromide [1530-32-1] **M 371.3, m 203-205°**. Recrystd from H₂O and dried in high vacuum at 100°. IR has bands at 1449, 1431 and 997cm⁻¹. [Justus Liebigs Ann Chem 606 1 1957; J Org Chem 23 1245 1958.]

Europium (III) acetate (2H₂O) [62667-64-5] **M 383.1, pK₁^{2.5} 8.31 (for aquo Eu³⁺)**. Recrystd several times from water [Ganapathy et al. J Am Chem Soc 108 3159 1986]. For europium shift reagents see lanthanide shift reagents in Chapter 4.

Ferric acetylacetonate [14024-18-1] **M 353.2, m 181.3-182.3°**. Recrystd twice from *benzene-pet ether **m 181.3-182.3°** corr [J Chem Soc 1256 1938]. Recrystd from EtOH or Et₂O, **m 179°** [Justus Liebigs Ann Chem 323 13 1902]. Recrystd from absolute EtOH, **m 159.5°** [Chem Ber 67 286 1934]. Dry for 1hr at 120°.

Ferric Bromide [10031-26-2] **M 395.6, m >130°(dec)**. Sublimed in a sealed tube with Br₂ at 120°-200°. [Lux in Handbook of Preparative Inorganic Chemistry (Ed. Brauer) Vol II, p 1494 1963.]

Ferric chloride (anhydrous) [7705-08-0] **M 162.2, m >300°(dec)**. Sublimed at 200° in an atmosphere of chlorine. Stored in a weighing bottle inside a desiccator.

Ferric chloride (6H₂O) [10025-77-1] **M 270.3, m 37°(dec), pK₁²⁵ 2.83, pK₂²⁵ 4.59** (for hydrolysis of Fe³⁺). An aqueous soln, saturated at room temperature, was cooled to -20° for several hours. Pptn was slow, even with scratching and seeding, and it was generally necessary to stir overnight. The presence of free HCl retards the pptn [Linke *J Phys Chem* **60** 91 1956].

Ferric nitrate (9H₂O) [7782-61-8] **M 404.0, m 47°(dec)**. Cryst from aqueous solutions of moderately strong HNO₃ as the violet nonahydrate. With more concentrated aqueous solns (containing some HNO₃), the hexahydrate crystals out. The anhydrous salt is slightly deliquescent and decomposes at 47°.

Ferric perchlorate (9H₂O) [13537-24-1] **M 516.3, pK²⁵ -2.4 to -3.1 (for HClO₄)**. Crystd twice from conc HClO₄, the first time in the presence of a small amount of H₂O₂ to ensure that the iron is fully oxidised [Sullivan *J Am Chem Soc* **84** 4256 1962]. Extreme care should be taken with this preparation because it is potentially **DANGEROUS**.

Ferric sulfate (xH₂O) [10028-22-5] **M 399.9 + xH₂O**. Dissolve in the minimum volume of dilute aqueous H₂SO₄ and allow to evaporate at room temp until crystals start to form. Do not concentrate by boiling off the H₂O as basic salts will be formed. Various *hydrates* are formed the—common ones are the *dodeca* and *nona hydrates* which are violet in colour. The anhydrous salt is colourless and very *hygroscopic* but dissolves in H₂O slowly unless ferrous sulfate is added.

Ferrocene [102-54-5] **M 186.0, m 173-174°**. Purified by crystn from pentane or cyclohexane (also *C₆H₆ or MeOH can be used). Moderately soluble in Et₂O. Sublimes readily above 100°. Crystallisation from EtOH gave **m 172.5-173°**. [*Org Synth Coll Vol IV* 473 1963; *J Chem Soc* 632 1952.] Also crystd from methanol and sublimed *in vacuo*. [Saltiel et al. *J Am Chem Soc* **109** 1209 1987.]

Ferrocene carboxaldehyde [12093-10-6] **M 214.1, m 117-120°, 118-120°, 121°, 124.5°**. Red crystals from EtOH or pet ether and sublimed at 70°/1mm. *Semicarbazone* **m 217-219°(dec)** cryst from aqueous EtOH. *O-Acetyloxime* **m 80-81°** cryst from hexane [*J Org Chem* **22** 355 1957]. *2,4-Dinitrophenylhydrazone* **m 248°(dec)**. [*Beilstein* **16 4th Suppl** 1798; *J Am Chem Soc* **79** 3416 1957; *J Chem Soc* 650 1958.]

Ferrocene carboxylic acid [1271-42-7] **M 230.1, m 210°(dec), 225-230°(dec)**. Yellow crystals from pet ether. Also crystd from aqueous ethanol. [Matsue et al. *J Am Chem Soc* **107** 3411 1985.] *Acid chloride* **m 49°** crystallises from pentane, λ_{max} 458nm [*J Org Chem* **24** 280 1959]. *Methyl ester* crystallises from aq MeOH **m 70-71°**. *Anhydride* **m 143-145°** from pet ether [*J Org Chem* **24** 1487 1959]. *Amide* **m 168-170°** from CHCl₃-Et₂O or **m 167-169°** from *C₆H₆-MeOH. [*J Am Chem Soc* **77** 6295 1955; **76** 4025 1954.]

Ferrocene-1,1'-dicarboxylic acid [1293-87-4] **M 274.1, m >250°(dec), >300°**. Orange-yellow crystals from AcOH. Sublimes above 230°. *Monomethyl ester* **m 147-149°** [*Dokl Acad Nauk USSSR* **115**, 518 1957]. *Dimethyl ester* **m 114-115°** [*J Am Chem Soc* **74**, 3458 1958]. *Diacid chloride* **m 92-93°** from pet ether. [*Dokl Acad Nauk SSSR* **120** 1267 1958; **127** 333 1959.]

Ferrocene-1,1,-dimethanol [1291-48-1] **M 246.1, m 107-108°**. Obtained from the diacid with LiAlH₄ reduction and recrystd from Et₂O-pet ether. [*J Am Chem Soc* **82** 4111 1960.]

Ferrous bromide [20049-65-4] **M 215.7 + xH₂O, m 684°, d²⁵ 4.63**. Crystn from air-free H₂O provides the *hexahydrate* as pale green to bluish-green rhombic prisms. On heating at 49° H₂O is lost and the *tetrahydrate* is formed. Further heating at 83° more H₂O is lost and the *dihydrate* is formed as a light yellow to dark brown *hygroscopic* powder. The ferrous iron in the aqueous solns of these salts readily oxidises to ferric iron. The salts should be stored over H₂SO₄ under N₂ in tightly closed containers. They have some solubility in EtOH. [*Chem Ber* **38** 236 1904.]

Ferrous chloride (4H₂O) [13478-10-9] **M 198.8, m 105°(dec), pK₁²⁵ 6.7, pK₂²⁵ 9.3** (for aquo Fe²⁺). A 550mL round-bottomed Pyrex flask was connected, *via* a glass tube fitted with a medium porosity

sintered-glass disc, to a similar flask. To 240g of $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ in the first flask was added conductivity water (200mL), 38% HCl (10mL), and pure electrolytic iron (8-10g). A stream of purified N_2 was passed through the assembly, escaping through a mercury trap. The salt was dissolved by heating which was continued until complete reduction had occurred. By inverting the apparatus and filtering (under N_2 pressure) through the sintered glass disc, unreacted iron was removed. After cooling and crystn, the unit was again inverted and the crystals of ferrous chloride were filtered free from mother liquor by applied N_2 pressure. Partial drying by overnight evacuation at room temperature gave a mixed hydrate which, on further evacuation on a water bath at 80° , lost water of hydration and its absorbed HCl (with vigorous effervescence) to give a white powder, $\text{FeCl}_2 \cdot 2\text{H}_2\text{O}$ [Gayer and Wootner *J Am Chem Soc* **78** 3944 1956].

Ferrous chloride [7758-94-3] **M 126.8, m 674° , b 1023° , d^{25} 3.16.** Sublimes in a stream of HCl at *ca* 700° , or in H_2 below 300° . Its vapour pressure at 700° is 12mm. Anhydrous FeBr_2 can be obtained by carefully dehydrating the *tetrahydrate* in a stream of HBr and N_2 , and it can be sublimed under N_2 . White *hygroscopic* rhombohedral crystals with a green tint. They oxidise in air to $\text{FeCl}_3 + \text{Fe}_2\text{O}_3$. Sol in H_2O , EtOH Me_2CO but insol in Et_2O . The *tetrahydrate* is pale green to pale blue in colour and loses $2\text{H}_2\text{O}$ at 105 - 115° . The *dihydrate* loses H_2O at 120° . The ferrous iron in the aqueous solns of these salts readily oxidises to ferric iron. [*Inorg Synth* **6** 172 1960; *Handbook of Preparative Inorganic Chemistry* (Ed Brauer) Vol II 1491 1965.]

Ferrous perchlorate ($6\text{H}_2\text{O}$) [13933-23-8] **M 362.9, pK^{25} -2.4 to -3.1 (for HClO_4).** Crystd from HClO_4 .

Ferrous sulfate ($7\text{H}_2\text{O}$) [7782-63-0] **M 278.0, m $\sim 60^\circ$ (dec).** Crystd from 0.4M H_2SO_4 .

Fluorine [7782-41-4] **M 38.0, b -129.2° .** Passed through a bed of NaF at 100° to remove HF and SiF_4 . [For description of stills used in fractional distn, see Greenberg et al. *J Phys Chem* **65** 1168 1961; Stein, Rudzitis and Settle *Purification of Fluorine by Distillation*, Argonne National Laboratory, ANL-6364 1961 (from Office of Technical Services, US Dept of Commerce, Washington 25).] **HIGHLY TOXIC.**

Fluoroboric acid [16872-11-0] **M 87.8, pK -4.9.** Crystd several times from conductivity water.

Fluorotrimethylsilane (trimethylsilyl fluoride, TMSF) [420-56-4] **M 92.2, m -74° , b $16^\circ/760\text{mm}$, $19^\circ/730\text{mm}$, d^0 0.793.** It is a **FLAMMABLE** gas which is purified by fractional distn through a column at low temperature and with the exclusion of air [Booth and Suttle *J Am Chem Soc* **68** 2658 1946; Reid and Wilkins *J Chem Soc* 4029 1955].

Gallium [7440-55-3] **M 69.7, m 29.8° .** Dissolved in dilute HCl and extracted into Et_2O . Pptn with H_2S removed many metals, and a second extraction with Et_2O freed Ga more completely, except for Mo, Th(III) and Fe which were largely removed by pptn with NaOH. The soln was then electrolysed in 10% NaOH with a Pt anode and cathode (2-5A at 4-5V) to deposit Ga, In, Zn and Pb, from which Ga was obtained by fractional crystn of the melt [Hoffman *J Res Nat Bur Stand* **13** 665 1934]. Also purified by heating to boiling in 0.5-1M HCl, then heating to 40° in water and pouring the molten Ga with water under vacuum onto a glass filter (30-50 μ pore size), to remove any unmelted metals or oxide film. The Ga was then fractionally crystd from the melt under water.

Gallium (III) Chloride [13450-90-3] **M 176.1, m 77.8° , b $133^\circ/100\text{mm}$, $197.7^\circ/700\text{mm}$, d 2.47, pK_1^{25} 2.91, pK_2^{25} 3.70, pK_3^{20} 4.42 (for Ga^{3+}).** Pure compound can be obtained by redistn in a stream of Cl_2 or Cl_2/N_2 followed by vacuum sublimation or zone refining. Colourless needles which give *gallium dichloride* [$\text{Ga}(\text{GaCl}_4)$, m 172.4°] on heating. Dissolves in H_2O with liberation of heat. Soluble in Et_2O . [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 846 1963.]

Gallium (III) nitrate ($9\text{H}_2\text{O}$) [63462-65-7] **M 417.9, m *ca* 65° .** Recrystd from H_2O (sol: 295g/100mL at 20°). White deliquescent colourless powder soluble in H_2O , absolute EtOH and Et_2O . Loses

HNO₃ upon heating at 40°. Addition of Et₂O to a warm ethanolic soln (40-50°) of Ga(NO₃)₃·9H₂O precipitates Ga(OH)₂NO₃·Ga(OH)₃·2H₂O. If the salt has partly hydrolysed, dissolve in conc HNO₃, reflux, dilute with H₂O and concentrate on a sand bath. Wash several times by adding H₂O and evaporate until there is no odour of acid. Dilute the residue to a Ga concentration of 26g/100mL. At this concentration, spongy Ga(NO₃)₃·xH₂O separates from the viscous soln. After standing for several days the crystals are collected and dried in a stream of dry air first at room temp then at 40°. Dehydration is complete after 2 days. Recrystallise from H₂O and dry on a water pump at room temperature. [*Z Naturforsch* **20B** 71 1965; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 856/1963.]

Gallium (III) sulfate [13494-91-2 (*anhydr*); 13780-42-2 (*hydr*)] **M 427.6**. Recrystn from H₂O gives the 16-18H₂O hydrate (sol at 20° is 170g/100mL). Alternatively dissolve in 50% H₂SO₄ and evaporate (60-70°), cool and ppt by adding EtOH/Et₂O. On heating at 165° it provides the *anhydrous* salt which is a white *hygroscopic* solid. [*Z Naturforsch* **20B** 71 1965.]

Germanium [7440-56-4] **M 72.6, m 937°, 925-975°, b 2700°, d 5.3**. Copper contamination on the surface and in the bulk of single crystals of Ge can be removed by immersion in molten alkali cyanide under N₂. The Ge was placed in dry cyanide powder in a graphite holder in a quartz or porcelain boat. The boat was then inserted into a heated furnace which, after a suitable time, was left to cool to room temperature. At 750°, a 1mm thickness requires about 1min, whereas 0.5cm needs about half hour. The boat was removed and the samples were taken out with plastic-coated tweezers, carefully rinsed in hot water and dried in air [Wang *J Phys Chem* **60** 45 1956].

Germanium (IV) oxide [1310-53-8] **M 104.6, m 1080°(soluble form), d²⁵ 6.239; m 1116°(insoluble form) d²⁵ 4.228, pK₁²⁵ 9.02, pK₂²⁵ 12.82 (for germanic acid H₂GeO₃)**. The oxide (GeO₂) is usually prepared by hydrolysing redistd GeCl₄ and igniting in order to remove H₂O and chloride. It can be further purified by dissolving in hot H₂O (sol: 4g/L cold) evaporating and drying the residual crystalline solid. When the *soluble* form (which is produced in H₂O at 355°) is heated for 100h it is converted to the *insoluble* form. This form is stable at temperatures up to 1033°, and fusion at 1080° for 4h causes complete devitrification and it reverts to the *soluble* form. [*J Am Chem Soc* **46** 2358 1924, **47** 1945 1925, **54** 2303 1032.]

Germanium tetrachloride [10038-98-9] **M 214.4, m -49.5° (α), -52.0° (β), b 83.1°/760mm, 86.5°/760mm corr, d₄²⁰ 1.84**. Traces of Cl₂ and HCl can be removed from the liquid by blowing dry air through it for a few hours at room temperature or shake it with Hg or Hg₂Cl₂ and then fractionally distil in a vacuum. It decomposes on heating at 950°. It has a sharp penetrating odour and fumes in moist air to give a chalky coat of GeO₂. It is slowly hydrolysed by H₂O to give GeO₂. [*J Am Chem Soc* **44** 306 1922.]

Germanium tetraethoxide [14165-55-0] **M 252.8, m -72°; b 54.5°/5mm, 71-72°/11mm, 188-190°/722mm, d²⁵ 1.1288**. Distil through a 10cm Vigreux column under reduced pressure. Alternatively distil through a Fenske glass helices column fitted with a total condensation variable take-off stillhead. Fractionate under reduced pressure using a reflux ratio of 10:1. [*J Am Chem Soc* **75** 718 1953; *J Chem Soc* 4916 1956.]

Glass powder (100-300 mesh). Washed with 10% HNO₃, water and dried.

Gold (III) bromide (gold tribromide) [10294-28-7] **M 436.7, m 150°(dec)**. Purified by adding pure Br₂ to the dark powder, securely stopper the container, warm a little and shake while keeping away from light for ca 48h. Remove the stopper and place over NaOH until free Br₂ is no longer in the apparatus (48-60h). The bright yellow needles of the tribromide are stable over NaOH in the dark. It is sol in H₂O and in EtOH where it is slowly reduced. Keep in a cooled closed container and protect from light as decomposition causes gold to be formed. *Aurobromic acid* can be obtained by adding the calculated amount of conc HBr to AuBr₃ (actually Au₂Br₆) until all dissolves, whereby the acid crystallises out as H₂AuBr₄·5H₂O, deliquescent solid soluble in EtOH with **m ca 27°**, and store as above. [*J Chem Soc* 2410 1931, 217, 219 1935.]

Gold (III) chloride (hydrate) [16903-35-8] M 339.8 + xH₂O, m 229°, b 354°(dec), d 3.9. Obtained as a dark red crystalline mass by dissolving Au in aqua regia and evaporating. When sublimed at 180° the crystals are ruby red. The anhydrous salt is *hygroscopic* sol in H₂O but sparingly soluble in EtOH and Et₂O. *Aurochloric acid* is formed when AuCl₃ is dissolved in HCl. [*J Am Chem Soc* 35 553 1913; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1056 1965.]

Gold (I) cyanide [506-65-0] M 223.0, m dec on heating. The lemon yellow powder is sparingly soluble in H₂O and EtOH but soluble in aqueous NH₃. It is obtained by heating H[Au(CN)₂] at 110°. Wash well with H₂O and EtOH and dry at 110°. It has an IR band at ν 2239cm⁻¹ typical for C≡N stretching vibration. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1064 1965.] CARE: may evolve HCN.

Gold (I) iodide [10294-31-2] M 323.9, m 120°(dec), d 8.25. It has been prepared by heating gold and iodine in a tube at 120° for 4 months. Since it decomposes to Au and I₂ in the presence of UV light and heat then the main impurity is Au. The salt is therefore purified by heating at 120° with I₂ for several weeks. The crystals should be kept dry and in a cool place in the dark. [*Z Naturforsch* 11B 604 1956.]

Gold (III) oxide hydrate [1303-58-8] M 441.9 + xH₂O, evolves O₂ at 110°, pK₁²⁵ < 11.7, pK₂²⁵ 13.36, pK₃²⁵ > 15.3 [for Au(OH)₃]. Most probable impurities are Cl⁻ ions. Dissolve in strong boiling KOH soln (ca 5M) and precipitate (care) with excess of 3N H₂SO₄. Then shake and centrifuge, resuspend in H₂O and repeat wash several times until free from SO₄ and Cl ions. This gives a *wet* oxide which is dried in air, and dec to Au in sunlight. It is best to keep it wet as it decomposes on drying (analyse wet sample). Store away from light in the presence of H₂O vapour. It evolves O₂ at 110°. It is insoluble in H₂O but soluble in HCl and conc HNO₃. [*J Am Chem Soc* 49 1221 1927.]

Graphite [7782-42-5]. Treated with hot 1:1 HCl. Filtered, washed, dried, powdered and heated in an evacuated quartz tube at 1000° until a high vacuum was obtained. Cooled and stored in an atmosphere of helium [Craig, Van Voorhis and Bartell *J Phys Chem* 60 1225 1956].

Haematoporphyrin IX [8,13-bis(1-hydroxyethyl)-3,7,12,17-tetramethyl-21H-23H-porphin-2,18-dipropionic acid [14459-29-1] M 598.7, pK_{Est} ~ 4.8 (-CH₂CH₂COOH). See haematoporphyrin on p. 541 in Chapter 6.

Helium [7440-59-7] M 4.0. Dried by passage through a column of Linde 5A molecular sieves and CaSO₄, then passed through an activated-charcoal trap cooled in liquid N₂, to adsorb N₂, argon, xenon and krypton. Passed over CuO pellets at 300° to remove hydrogen and hydrocarbons, over Ca chips at 600° to remove oxygen, and then over titanium chips at 700° to remove N₂ [Arnold and Smith *J Chem Soc, Faraday Trans 2* 77 861 1981].

Hexabutyl-distannane [hexabutyl-ditin, bis(tributyl)tin] [813-19-4] M 580.4, b 160-162°/0.3mm, d 1.148, n 1.512. Purified by distn in a vacuum and stored in the dark. [Shirai et al. *Yakugaku Zasshi* 90 59 1970, *Chem Abstr* 72 90593 1970.]

Hexachlorocyclotriphosphazene [940-71-6] M 347.7, m 113-114°, 113-115°. See phosphonitric chloride trimer on p. 450.

Hexachloroplatinic acid hydrate (H₂PtCl₆, chloroplatinic acid, platinum IV chloride solution) [16941-12-1] M 409.8 + H₂O, m 60° (deliquescent solid). If it is to be purified, or regenerated from Pt recovered from catalytic hydrogenations, it was dissolved in aqua regia followed by evaporation to dryness and dissolution in the minimum vol of H₂O. Then the aqueous solution was treated with saturated ammonium chloride until all the ammonium hexachloroplatinate separated. The (NH₄)₂PtCl₆ was filtered off and dried at 100°. Ignite the salt to give Pt sponge, dissolve the Pt sponge in aqua regia, boil to

dryness, dissolve in concentrated HCl, boil to dryness again and repeat the process. Protect from light. [Hickers *J Am Chem Soc* **43** 1268 1921; *Org Synth Coll Vol I* 463, 466 1941; Bruce *J Am Chem Soc* **58** 687 1936.]

Hexaethyldisiloxane [924-49-0] **M 246.5, b 114-115°/16mm, 235.5°/760mm, d 0.8443, n 1.4330.** Distil in a vacuum, but can be distilled at atmospheric pressure without decomposition. It is characterised by completely dissolving in conc H₂SO₄. [*J Chem Soc* 3077 1950.]

2,2,4,4,6,6-Hexamethylcyclotrisiloxane [1009-93-4] **M 219.5, m -10°, b 81-82°/19mm, 111-112°/85mm, 188°/756mm, d 0.9196, n 1.448.** Purified by fractional distillation at atmospheric pressure until the temperature reaches 200°. The residue in the flask is mostly octamethylcyclotetrasilazane. [*J Am Chem Soc* **70** 3888 1948.]

Hexamethyldisilane [1450-14-2] **M 164.4, m 9-12°, b 113.1°/750mm, d 0.7272, n 1.4229.** Most likely impurity is trimethylchlorosilane (*cf* boiling point). Wash with H₂O, cold conc H₂SO₄, H₂O again then aqueous NaHCO₃, dry over CaSO₄ and fractionate at atmospheric pressure. [*J Chem Soc* 2811 1958.] Grossly impure sample (25% impurities) was purified by repeated spinning band distn. This lowered the impurity level to 500 ppm. The main impurity was identified as 1-hydroxypentamethyldisilane.

Hexamethyldisilazane (HMDS) [999-97-3] **M 161.4, b 125-125.6°/atm, 126°/760mm, d 0.7747, n 1.407.** Possible impurity is Me₃SiCl. Wash well with pet ether and fractionate through a vacuum jacketed column packed with Helipac using a reflux ratio of 10:1. [*J Org Chem* **23** 50 1958.]

Hexamethyldisiloxane [107-46-0] **M 162.4, b 99.4°/760mm, 100.4°/764mm, d 0.7633, n 1.3777.** Fractionally distilled through a column packed with glass helices with *ca* 15 theoretical plates. [*J Am Chem Soc* **76** 2672 1954; *J Gen Chem USSR (Engl ed)* **25** 469 1955.]

Hexamethylditin (hexamethyldistannane) [661-69-8] **M 327.6, m 23.5°, b 85-88°/45mm, 182°/756mm, d²⁵ 1.57.** Wash with H₂O and extract with *C₆H₆, dry by filtering through powdered Na₂SO₄, remove *C₆H₆ on a rotary evaporator and fractionally dist the oily residue under vacuum (**b 85-88°/45mm**). *It boils at ca 182° at atmospheric press but it cannot be distilled in air because the hot vapours flash in the condenser.* [*J Am Chem Soc* **47** 2361 1925, **63** 2509 1941; *Trans Faraday Soc* **53** 1612 1957.]

Hexamethylphosphoric triamide (HMPA) [680-31-9] **M 179.2, f 7.2°, b 68-70°/1mm, 235°/760mm, d 1.024, n 1.460.** The industrial synthesis is usually by treatment of POCl₃ with excess of dimethylamine in isopropyl ether. Impurities are water, dimethylamine and its hydrochloride. It is purified by refluxing over BaO or CaO at about 4mm pressure in an atmosphere of nitrogen for several hours, then distd from sodium at the same pressure. The middle fraction (**b ca 90°**) is collected, refluxed over sodium under reduced pressure under nitrogen and distd. It is kept in the dark under nitrogen, and stored in solid CO₂. Can also be stored over 4A molecular sieves.

Alternatively, it is distd under vacuum from CaH₂ at 60° and crystd twice in a cold room at 0°, seeding the liquid with crystals obtained by cooling in liquid nitrogen. After about two-thirds frozen, the remaining liquid is drained off [Fujinaga, Izutsu and Sakara *Pure Appl Chem* **44** 117 1975]. For tests of purity see Fujinaga et al. in *Purification of Solvents*, Coetzee Ed., Pergamon Press, Oxford, 1982. For efficiency of desiccants in drying HMPT see Burfield and Smithers [*J Org Chem* **43** 3966 1978; Sammes et al. *J Chem Soc, Faraday Trans 1* 281 1986]. **CARCINOGEN.**

Hexamethylphosphorous triamide (HMPT) [1608-26-0] **M 163.2, m 7.2°, b 49-51°/12mm, 162-164°/12mm, d 0.989, n 1.466.** It may contain more than 1% of phosphoric triamide. The yellow oil is first distd at atm press then under reduced press and stored under N₂. It is air sensitive, **TOXIC**, should not be inhaled and is absorbed through the skin. [Mark *Org Synth Coll Vol V* 602 1973.]

Hexamminecobalt(III) chloride [10534-89-1] **M 267.5.** Crystd from warm water (8mL/g) by cooling. [Bjerrum and McReynolds *Inorg Synth* **2** 217 1946.]

Hexammineruthenium(III) chloride [14282-91-8] **M 309.6.** Crystd twice from 1M HCl.

Hexarhodium hexadecacarbonyl [28407-51-4] **M 1065.6, m 220°(dec, in air), d 2.87.** Slowly loses CO when heated in air; may be regenerated by heating at 80-200° in the presence of CO at 200atm pressure for 15h, preferably in the presence of Cu. Forms black crystals which are insoluble in hexane. It has bands at 2073, 2026 and 1800cm⁻¹ in the IR. [*Z Anorg Allg Chem* **251** 96 1963; *J Am Chem Soc* **85** 1202 1963; *Tetrahedron Lett* **22** 1783 1981.]

Hydrazine (anhydrous) [302-01-2] **M 32.1, f 1.5-2.0°, b 47°/26mm, 56°/71mm, 113-113.5°/atm, n 1.470, d 1.91, pK₁²⁵ -0.88, pK₂²⁵ 8.11.** Hydrazine hydrate is dried by refluxing with an equal weight of KOH pellets for 3h, then distilled from fresh NaOH or BaO in a current of dry N₂.

Hydrazine dihydrochloride [5341-61-7] **M 105.0, m 198°, d 1.42.** Crystd from aqueous EtOH and dried under vacuum over CaSO₄.

Hydrazine monohydrochloride [2644-70-4] **M 68.5, m 89°.** Prepared by dropwise addition of cold conc HCl to cold liquid hydrazine in equimolar amounts. The crystals were harvested from water and were twice recrystd from absolute MeOH and dried under vacuum. [Kovack et al. *J Am Chem Soc* **107** 7360 1985.]

Hydriodic acid [10034-85-2] **M 127.9, b 127°(aq azeotrope), d 1.701, pK²⁵ -8.56.** Iodine can be removed from aqueous HI, probably as the amine hydrogen triiodide, by three successive extractions using a 4% soln of Amberlite LA-2 (a long-chain aliphatic amine) in CCl₄, toluene or pet ether (10mL per 100mL of acid). [Davidson and Jameson *Chem Ind (London)* 1686 1963.] Extraction with tributyl phosphate in CHCl₃ or other organic solvents is also suitable. Alternatively, a De-acidite FF anion-exchange resin column in the OH⁻-form using 2M NaOH, then into its I⁻-form by passing dilute KI soln, can be used. Passage of an HI solution under CO₂ through such a column removes polyiodide. The column can be regenerated with NaOH. [Irving and Wilson *Chem Ind (London)* 653 1964.] The earlier method was to reflux with red phosphorus and distil in a stream of N₂. The colourless product was stored in ampoules in the dark [Bradbury *J Am Chem Soc* **74** 2709 1952; *Inorg Synth* **1** 157 1939]. Fumes in moist air. **HARMFUL VAPOURS.**

Hydrobromic acid [10035-10-6] **M 80.9, b 125°(aq azeotrope, 47.5% HBr), d 1.38 (34% HBr), pK²⁵ -8.69.** A soln of aqueous HBr ca 48% (w/w, constant boiling) was distilled twice with a little red phosphorus, and the middle half of the distillate was taken. (The azeotrope at 760mm contains 47.8% (w/w) HBr.) [Hetzer, Robinson and Bates *J Phys Chem* **66** 1423 1962]. Free bromine can be removed by Irvine and Wilson's method for HI (see above), except that the column is regenerated by washing with an ethanolic solution of aniline or styrene. Hydrobromic acid can also be purified by aerating with H₂S, distilling and collecting the fraction boiling at 125-127°. [*Inorg Synth* **1** 155 1939.] **HARMFUL VAPOURS.**

Hydrochloric acid [7647-01-0] **M 36.5, b 108.6°(aq azeotrope, 20.2% HCl), d 1.09(20%), pK²⁵ -6.1.** Readily purified by fractional distillation as constant boiling point acid, following dilution with H₂O. The constant-boiling fraction contains 1 mole of HCl in the following weights of distillate at the stated pressures: 179.555g (730mm), 179.766g (740mm), 179.979 (750mm), 180.193 (760mm), 180.407 (770mm) [Foulk and Hollingsworth *J Am Chem Soc* **45** 1220 1923..] **HARMFUL VAPOURS.**

Hydrofluoric acid [7664-39-3] **M 20.0, b 112.2°(aq azeotrope, 38.2% HF), d 1.15 (47-53% HF), pK²⁵ 3.21.** Freed from lead (Pb ca 0.002ppm) by co-precipitation with SrF₂, by addition of 10mL of 10% SrCl₂ soln per kilogram of the conc acid. After the ppte has settled, the supernatant is decanted through a filter in a hard-rubber or paraffin lined-glass vessel [Rosenqvist *Am J Sci* **240** 358 1942. Pure aqueous HF solutions (up to 25M) can be prepared by isothermal distn in polyethylene, polypropylene or platinum apparatus [Kwestroo and Visser *Analyst* **90** 297 1965]. **HIGHLY TOXIC.**

Hydrogen [1333-74-0] **M 2.0, m -259.1°, -252.9°.** Usually purified by passage through suitable absorption train. Carbon dioxide is removed with KOH pellets, soda-lime or NaOH pellets. Oxygen is removed with a "De-oxo" unit or by passage over Cu heated to 450-500°, Cu on Kieselguhr at 250°. Passage over a mixture of MnO₂ and CuO (Hopcalite) oxidises any CO to CO₂ (which is removed as above). Hydrogen can be dried by passage through dried silica-alumina at -195°, through a dry-ice trap followed by a liquid-N₂ trap

packed with glass wool, through CaCl_2 tubes, or through $\text{Mg}(\text{ClO}_4)_2$ or P_2O_5 . Other purification steps include passage through a hot palladium thimble [Masson *J Am Chem Soc* **74** 4731 1952], through an activated-charcoal trap at -195° , and through non-absorbent cotton-wool filter or small glass spheres coated with a thin layer of silicone grease. *Potentially VERY EXPLOSIVE in air.*

Hydrogen bromide (anhydrous) [10035-10-6] **M 80.9**. Dried by passage through $\text{Mg}(\text{ClO}_4)_2$ towers. This procedure is **hazardous**, see Stoss and Zimmermann [*Ind Eng Chem* **17** 70 1939]. Shaken with mercury, distd through a -78° trap and condensed at $-195^\circ/10^{-5}\text{mm}$. Fumes in moist air. **HARMFUL VAPOURS**.

Hydrogen chloride [7647-01-0] **M 36.5**. Passed through conc H_2SO_4 , then over activated charcoal and silica gel. Fumes in moist air. Hydrogen chloride in gas cylinder include ethylene, 1,1-dichloroethane and ethyl chloride. The latter two may be removed by fractionating the HCl through a trap cooled to -112° . Ethylene is difficult to remove. Fumes in moist air. **HARMFUL VAPOURS**.

Hydrogen cyanide (anhydrous) [74-90-8] **M 27.0, b 25.7°, pK²⁵ 9.21 (aq acid)**. Prepared from NaCN and H_2SO_4 , and dried by passage through H_2SO_4 and over CaCl_2 , then distilled in a vacuum system and degassed at 77°K before use [Arnold and Smith *J Chem Soc, Faraday Trans 2* **77** 861 1981]. Cylinder HCN may contain stabilisers against explosive polymerisation, together with small amounts of H_3PO_4 , H_2SO_4 , SO_2 , and water. It can be purified by distn over P_2O_5 , then frozen in Pyrex bottles at Dry-ice temperature for storage. **EXTREMELY POISONOUS**.

Hydrogen fluoride (anhydrous) [7664-39-3] **M 20.0, b 19.4°**. Can be purified by trap-to-trap distn, followed by drying over CoF_2 at room temperature and further distn. Alternatively, it can be absorbed on NaF to form NaHF_2 which is then heated under vacuum at 150° to remove volatile impurities. The HF is regenerated by heating at 300° and stored with CoF_3 in a nickel vessel, being distilled as required. (Water content *ca* 0.01%.) To avoid contact with base metal, use can be made of nickel, polychlorotrifluoroethylene and gold-lined fittings [Hyman, Kilpatrick and Katz *J Am Chem Soc* **79** 3668 1957]. **HIGHLY TOXIC**.

Hydrogen iodide (anhydrous) [10034-85-2] **M 127.9, b -35.5°**. After removal of free iodine from aqueous HI, the solution is frozen, then covered with P_2O_5 and allowed to melt under vacuum. The gas evolved is dried by passage through P_2O_5 on glass wool. It can be freed from iodine contamination by repeated fractional distillation at low temperatures. Fumes in moist air. **HARMFUL VAPOURS**.

Hydrogen ionophore II (ETH 1907) (4-nonadecylpyridine - Proton ionophore) [70268-36-9] **M 345.6, b 180°/0.07mm, pK_{Est}~ 6.0**. Dissolve the waxy solid (*ca* 60g) in CHCl_3 (200mL), wash with H_2O (3 x 200mL), dry and evaporate to dryness then distil in vacuum. A waxy solid is formed on cooling the distillate. UV: 257nm (ϵ $1.86 \times 10^3 \text{ M}^{-1}\text{cm}^{-1}$), 308nm (ϵ $1.7 \times 10^2 \text{ M}^{-1}\text{cm}^{-1}$). [IR, NMR UV: *Inorg Chem* **18** 2160 1979.]

Hydrogen ionophore III (N,N-dioctadecyl methylamine) [4088-22-6] **M 536.0, m 40°, 44-46°, 48-49°, b 252-259°, pK_{Est}~ 10**. It can be distd at high vacuum; but dissolving in $^*\text{C}_6\text{H}_6$, filtering and evaporating gives a waxy solid suitable for electrode use. It recrystallises from Me_2CO . [*Chem Ber* **69** 60 1936; *Talanta* **34** 435 1987.]

Hydrogen ionophore IV ETH 1778 (octadecyl isonicotinate) [103225-02-1] **M 375.6, m 57.5°, pK_{Est}~ 3.5**. Dissolve in Et_2O and wash 3 times with H_2O . Dry, evaporate, and recrystallise the residue from EtOAc /hexane (4:1). [*Anal Chem* **58** 2285 1986.]

Hydrogen peroxide [7722-84-1] **M 34.0, d 1.110, pK²⁵ 11.65**. The 30% material has been steam distilled using distilled water. Gross and Taylor [*J Am Chem Soc* **72** 2075 1950] made 90% H_2O_2 approximately 0.001M in NaOH and then distilled under its own vapour pressure, keeping the temperature below 40° , the receiver being cooled with a Dry-ice/isopropyl alcohol mush. The 98% material has been rendered anhydrous by repeated fractional crystn in all-quartz vessels. **EXPLOSIVE IN CONTACT WITH ORGANIC MATERIAL**.

Hydrogen sulfide [7783-06-4] **M 34.1, b -59.6°, pK₁²⁵ 7.05, pK₂²⁵ 12.89.** Washed, then passed through a train of flasks containing saturated Ba(OH)₂ (2x), water (2x), and dilute HCl [Goates et al. *J Am Chem Soc* 73 707 1951]. **HIGHLY POISONOUS.**

Hydroquinone-2-sulfonic acid K salt [21799-87-1] **M 228.3, m 250°(dec), pK_{Est(1)}~1, pK_{Est(2)}~8.5, pK_{Est(3)}~11.** Recrystd from water.

Hydroxylamine [7803-49-8] **M 33.0, m 33.1°, b 56.5°/22mm, d 1.226, pK²⁰ 5.96.** Crystd from *n*-butanol at -10°, collected by vacuum filtration and washed with cold diethyl ether. **Harmful vapours.**

Hydroxylamine hydrochloride [5470-11-1] **M 69.5, m 151°.** Crystallised from aqueous 75% ethanol or boiling methanol, and dried under vacuum over CaSO₄ or P₂O₅. Has also been dissolved in a minimum of water and saturated with HCl; after three such crystns it was dried under vacuum over CaCl₂ and NaOH.

Hydroxylamine sulfate [10039-54-0] **M 164.1, m 170°(dec).** Crystallised from boiling water (1.6mL/g) by cooling to 0°.

Hydroxylamine-O-sulfonic acid [2950-43-8] **M 113.1, m 210-211°, 215°(dec), pK⁴⁵ 1.48.** Stir the solid vigorously with anhydrous Et₂O and filter off using large volumes of dry Et₂O. Drain dry at the pump for 5min and then for 12-14h in a vacuum. Store in a vacuum desiccator/conc H₂SO₄. Determine the purity by oxidation of iodide to I₂. Must be stored in a dry atmosphere at 0-4°. It decompose slowly in H₂O at 25° and more rapidly above this temperature. [*Inorg Synth* 5 122 1957.]

Hydroxynaphthol Blue tri-Na salt [63451-35-4] **M 620.5, m dec on heating, pK_{Est} <0.** Crude material was treated with hot EtOH to remove soluble impurities, then dissolved in 20% aqueous MeOH and chromatographed on a cellulose powder column with propanol:EtOH:water (5:5:4) as eluent. The upper of three zones was eluted to give the pure dye which was ppted as the monosodium salt trihydrate by adding conc HCl to the concentrated eluate [Ito and Ueno *Analyst* 95 583 1970].

4-Hydroxy-3-nitrobenzenearsonic acid [121-19-7] **M 263.0.** See 2-nitrophenol-4-arsonic acid on p. 446.

Hydroxyurea [127-07-1] **M 76.1, m 70-72° (unstable form), m 133-136°, 141° (stable form), pK 10.6.** Recrystallise from absolute EtOH (10g in 150mL). Note that the rate of solution in boiling EtOH is slow (15-30 min). It should be stored in a cool dry place but some decomposition could occur after several weeks. [*Org Synth Coll Vol V* 645 1973.] It is very soluble in H₂O and can be crystd from Et₂O. [*Acta Chem Scand* 10 256 1956.]

Hypophosphorous acid (Phosphinic acid) [6303-21-5] **M 66.0, m 26.5°, d₄³⁰ 1.217, 1.13 and 1.04 for 50, 30-32, and 10% aq solns resp, pK²⁵ 1.31 (H₃PO₂).** Phosphorous acid is a common contaminant of commercial 50% hypophosphorous acid. Jenkins and Jones [*J Am Chem Soc* 74 1353 1952] purified this material by evaporating about 600mL in a 1L flask at 40°, under reduced pressure (in N₂), to a volume of about 300mL. After the soln was cooled, it was transferred to a wide-mouthed Erlenmeyer flask which was stoppered and left in a Dry-ice/acetone bath for several hours to freeze (if necessary, with scratching of the wall). When the flask was then left at ca 5° for 12h, about 30-40% of it liquefied, and again filtered. This process was repeated, then the solid was stored over Mg(ClO₄)₂ in a vacuum desiccator in the cold. Subsequent crystns from *n*-butanol by dissolving it at room temperature and then cooling in an ice-salt bath at -20° did not appear to purify it further. The free acid forms deliquescent crystals **m 26.5°**, and is soluble in H₂O and EtOH. The NaH₂PO₂ salt can be purified through an anion exchange resin [*Z Anorg Allg Chem* 260 267 1949.]

Indigocarmine (2[1,3-dihydro-3-oxo-5-sulfo-2H-indol-2-ylidene]-2,3-dihydro-3-oxo-1H-indole-5-sulfonic acid di-Na salt) [860-22-0] **M 466.4, pK₁²⁰ 2.8, pK₂²⁰ 12.3.** Its

solubility in H₂O is 1g/100mL at 25°. Could be purified by dissolving in H₂O, filtering and adding EtOH to cause the salt to separate. Wash the solid with EtOH, Et₂O and dry *in vacuo*. [Vörländer and Schubert *Chem Ber* **34** 1860 1901; UV: Smit et al. *Anal Chem* **27** 1159 1955; Preisler et al. *J Am Chem Soc* **81** 1991 1959.]

Indium [7440-74-6] **M 114.8, m 156.6°, b 2000°, d 7.31**. Before use, the metal surface can be cleaned with dilute HNO₃, followed by a thorough washing with water and an alcohol rinse.

Indium (III) chloride [10025-82-8] **M 211.2, m 586°, d 4.0, pK₁²⁵ 3.54, pK₂²⁵ 4.28, pK₃²⁵ 5.16 (for aquo In³⁺)**. The anhydrous salt forms yellow deliquescent crystals which can be sublimed at 600° in the presence of Cl₂/N₂ (1:1) {does not melt}. It is resublimed in the presence of Cl₂/N₂ (1:10) and finally heated to 150° to expel excess Cl₂. It is soluble in H₂O and should be stored in a tightly closed container. [*J Am Chem Soc* **55** 1943 1933.]

Indium (III) oxide [1312-43-2] **M 277.6, d 7.18, m sublimes at 850°**. Wash with H₂O and dry below 850°. Volatilises at 850° and dissolves in hot mineral acids to form salts. Store away from light because it darkens due to formation of In.

Indium sulfate [13464-82-9] **M 517.8**. Crystd from dilute aqueous H₂SO₄.

Indium (III) sulfate (5H₂O) [17069-79-3] **M 607.9, d 3.44**. Dissolve in strong H₂SO₄ and slowly evaporate at *ca* 50°. Wash crystals with glacial AcOH and then heat in a furnace at a temperature of 450-500° for 6h. Sol in H₂O is 5%. The pentahydrate is converted to an anhydrous *hygroscopic* powder on heating at 500° for 6h; but heating above this temperature over N₂ yields the oxide sulfate. Evaporation of neutral aqueous solutions provides basic sulfates. [*J Am Chem Soc* **55** 1943 1933, **58** 2126 1936.]

Iodic acid [7782-68-5] **M 175.9, m 118°(dec), d 4.628, pK²⁵ 0.79**. Dissolve in the minimum volume of hot dilute HNO₃, filter and evaporate in a vacuum desiccator until crystals are formed. Collect crystals and wash with a little cold H₂O and dry in air in the dark. Soluble in H₂O: 269g/100mL at 20° and 295g/100mL at 40°. Soluble in dilute EtOH and darkens on exposure to light. It is converted to HIO₃.I₂O₅ on heating at 70°, but at 220° complete conversion to HIO₃ occurs. [*J Am Chem Soc* **42** 1636 1920, **53** 44 1931.]

Iodine [7553-56-2] **M 253.8, m 113.6°**. Usually purified by vacuum sublimation. Preliminary purifications include grinding with 25% by weight of KI, blending with 10% BaO and subliming; subliming with CaO; grinding to a powder and treating with successive portions of H₂O to remove dissolved salts, then drying; and crystn from *benzene. Barrer and Wasilewski [*Trans Faraday Soc* **57** 1140 1961] dissolved I₂ in conc KI and distilled it, then steam distilled three times, washing with distilled H₂O. Organic material was removed by sublimation in a current of O₂ over platinum at about 700°, the iodine being finally sublimed under vacuum. **HARMFUL VAPOURS.**

Iodine monobromide [7789-33-5] **M 206.8, m 42°**. Purified by repeated fractional crystallisation from its melt.

Iodine monochloride [7790-99-0] **M 162.4, m 27.2°**. Purified by repeated fractional crystallisation from its melt.

Iodine pentafluoride [7783-66-6] **M 221.9, m -8.0°, b 97°**. Rogers et al. [*J Am Chem Soc* **76** 4843 1954] removed dissolved iodine from IF₅ by agitating with a mixture of dry air and ClF₃ in a Fluorothene beaker using a magnetic stirrer. The mixture was transferred to a still and the more volatile impurities were pumped off as the pressure was reduced below 40mm. The still was gradually heated (kept at 40mm) to remove the ClF₃ before IF₅ distilled. Stevens [*J Org Chem* **26** 3451 1961] pumped IF₅ under vacuum from its cylinder, trapping it at -78°, then allowing it to melt in a stream of dry N₂. **HARMFUL VAPOURS.**

Iodine trichloride [865-44-1] **M 233.3, m 33°, b 77°(dec)**. Purified by sublimation at room temperature.

Iodomethyl trimethylsilane [4206-67-1] **M 214.1, b 139.5°/744mm, d 1.44, n_D^{25} 1.4917.** If slightly violet in colour wash with aqueous 1% sodium metabisulfite, H₂O, dry over Na₂SO₄ and fractionally distil at atmospheric pressure. [*J Am Chem Soc* **68** 481 1946.]

Iodotrimethylsilane (trimethylsilyl iodide, TMSI) [16029-98-4] **M 200.1, b 106.8°/742mm, 107.5°/760mm, d 1.470.** Add a little antimony powder and fractionate with this powder in the still. Stabilise with 1% wt of Cu powder. [*J Chem Soc* 3077 1950.]

Iridium [7439-88-5] **M 192.2, m 2450°, b ~4500°, d 22.65.** It is a silver white hard solid which oxidises on the surface in air. Scrape the outer tarnished layer until silver clear and store under paraffin. Stable to acids but dissolves in aqua regia. [*Chem Rev* **32** 277 1943.]

Iridium (IV) chloride hydrate (hexachloroiridic acid) [16941-92-7 (6H₂O); 207399-11-9 (xH₂O)] **M 334.0+H₂O.** If it contains nitrogen then repeatedly concentrate a conc HCl solution until free from nitrogen, and dry free from HCl in a vac over CaO until crystals are formed. The solid is very *hygroscopic*. [*J Am Chem Soc* **53** 884 1931; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1592 1965.]

Iron (wire) [7439-89-6] **M 55.9, m 1535°.** Cleaned in conc HCl, rinsed in de-ionised water, then reagent grade acetone and dried under vacuum.

Iron enneacarbonyl (di-iron nonacarbonyl) [15321-51-4] **M 363.7, m 100°(dec).** Wash with EtOH and Et₂O and dry in air. Sublimes at 35° at high vacuum. Dark yellow plates stable for several days when kept in small amounts. Large amounts, especially when placed in a desiccator spontaneously *ignite* in a period of one day. It decomposes in moist air. It is insoluble in hydrocarbon solvents but forms complexes with several organic compounds. [*J Am Chem Soc* **72** 1107 1950; *Chem Ber* **60** 1424 1927.]

Iron (III) meso-5,10,15,20-tetraphenylporphine chloride complex [16456-81-8] **M 704.0.** Crystallise by extraction from a thimble (Soxhlet) with CHCl₃. Concentrate the extract to ca 10mL and add ca 80mL of hot MeOH. Dark blue crystals separate on cooling. It can be recrystallised several times from CHCl₃-MeOH. Avoid prolonged heating. It is quite soluble in organic solvents but insoluble in pet ether. [*J Am Chem Soc* **70** 1808 1948; UV: **73** 4315 1951.]

Iron pentacarbonyl (pentacarbonyl iron) [13463-40-6] **M 195.9, m -20°, b 102.8°/749mm, 103°/760mm, n 1.520, d 1.490.** It is a pale yellow viscous liq which is PYROPHORIC and readily absorbed by the skin. **HIGHLY TOXIC (protect from light and air).** It should be purified in a vacuum line by distilling and collecting in a trap at -96° (toluene-Dry ice slush). It has been distd at atm pressure (use a very efficient fume cupboard). At 180°/atm it decomp to give Fe and CO. In UV light in pet ether it forms Fe₂(CO)₉. [Hagen et al. *Inorg Chem* **17** 1369 1978; Ewens et al. *Trans Faraday Soc* **35** 6811 1939.]

Isopropyldimethyl chlorosilane [3634-56-8] **M 140.7, b 109.8-110.0°/738mm, d 0.88, n 1.4158.** Probable impurity is Me₃SiCl (b 56.9°/783mm) which can be removed by efficient fractional distillation. [*J Am Chem Soc* **76** 801 1954.]

(2,3-O-Isopropylidene)-2,3-dihydroxy-1,4-bis-(diphenylphosphino)butane (DIOP) [4R,5S(-)-32305-98-9; 4S,5R(+)-37002-48-5] **M 498.5, m 88-90°, $[\alpha]_D^{19}$ (-) and (+) 26° (c 2.3, CHCl₃), pK_{Est} ~ 4.5.** It has been recrystd from *C₆H₆-pet ether. After 2 recrystns from EtOH it was pure by TLC on silica gel using Me₂CO-hexane as solvent. [Kagan and Dang *J Am Chem Soc* **94** 6429 1972.]

Lanthanide shift reagents see p. 277 in Chapter 4.

Lanthanum [7439-91-0] **M 138.9, m 920°, b 3470°, d 6.16.** White metal that slowly tarnishes in air due to oxidation. Slowly decomposed by H₂O in the cold and more rapidly on heating to form the

hydroxide. The metal is cleaned by scraping off the tarnished areas until the shiny metal is revealed and stored under oil or paraffin. It burns in air at 450°.

Lanthanum triacetate [917-70-4] **M 316.0, pK₁²⁵ 9.06 (for aquo La³⁺)**. Boil with redistilled Ac₂O for 10min (does not dissolve and is a white solid). Cool, filter, wash with Ac₂O and keep in a vacuum desiccator (NaOH) till free from solvent. [*J Indian Chem Soc* **33** 877 1956.]

***N*-Lauroyl-*N*-methyltaurine sodium salt (sodium *N*-decanoyl-*N*-methyl-2-aminoethane sulfonate)** [4337-75-1] **M 343.5, pK_{Est} ~1.5**. Prepared from methyldecanoate (at 180° under N₂) or decanoyl chloride and sodium *N*-methylthane sulfonate and purified by dissolving in H₂O and precipitating by addition of Et₂O. Decomposes on heating. [Desseigne and Mathian *Mém Services Chim Etat Paris* **31** 359 1944, *Chem Abstr* **41** 705 1947.]

Lead II acetate [301-04-2 (anhydr); 6080-56-4 (3H₂O)] **M 325.3, m 280°, pK₁²⁵ 7.1 (for Pb²⁺), pK₂²⁵ 10.1 (HPbO₂⁻), pK₃²⁵ 10.8 (PbO₂²⁻)**. Crystallised twice from anhydrous acetic acid and dried under vacuum for 24h at 100°.

Lead (bis-cyclopentadienyl) [1294-74-2] **M 337.4**. Purified by vacuum sublimation. Handled and stored under N₂.

Lead (II) bromide [10031-22-8] **M 367.0, m 373°**. Crystallised from water containing a few drops of HBr (25mL of water per gram PbBr₂) between 100° and 0°. A neutral solution was evaporated at 110° and the crystals that separated were collected by rapid filtration at 70°, and dried at 105° (to give the *monohydrate*). To prepare the anhydrous bromide, the hydrate is heated for several hours at 170° and then in a Pt boat at 200° in a stream of HBr and H₂. Finally fused [Clayton et al. *J Chem Soc, Faraday Trans 1* **76** 2362 1980].

Lead (II) chloride [7758-95-4] **M 278.1, m 501°**. Crystallised from distilled water at 100° (33mL/g) after filtering through sintered-glass and adding a few drops of HCl, by cooling. After three crystals the solid was dried under vacuum or under anhydrous HCl vapour by heating slowly to 400°.

Lead diethyldithiocarbamate [17549-30-3] **M 503.7, pK₁²⁵ 3.36 (for *N,N*-diethyldithiocarbamate)**. Wash with H₂O and dry at 60-70°, or dissolve in the min vol of CHCl₃ and add the same vol of EtOH. Collect the solid that separates and dry as before. Alternatively, recryst by slow evaporation of a CHCl₃ soln at 70-80°. Filter the crystals, wash with H₂O until all Pb²⁺ ions are eluted (check by adding chromate) and then dry at 60-70° for at least 10h. [*Justus Liebigs Ann Chem* **49** 1146 1977.]

Lead (II) formate [811-54-4] **M 297.3, m 190°**. Crystd from aqueous formic acid.

Lead (II) iodide [10101-63-0] **M 461.0, m 402°**. Crystd from a large volume of water.

Lead monoxide [1317-36-8] **M 223.2, m 886°**. Higher oxides were removed by heating under vacuum at 550° with subsequent cooling under vacuum. [Ray and Ogg *J Am Chem Soc* **78** 5994 1956.]

Lead nitrate [10099-74-8] **M 331.2, m 470°**. Ppted twice from hot (60°) conc aqueous soln by adding HNO₃. The ppte was sucked dry in a sintered-glass funnel, then transferred to a crystallising dish which was covered by a clock glass and left in an electric oven at 110° for several hours [Beck, Singh and Wynne-Jones *Trans Faraday Soc* **55** 331 1959]. After 2 recrystns of ACS grade no metals above 0.001 ppm were detected.

Lead tetraacetate [546-67-8] **M 443.4, m 175-180°, pK₁ 1.8, pK₂ 3.2, pK₃ 5.2, pK₄ 6.7**. Dissolved in hot glacial acetic acid, any lead oxide being removed by filtration. White crystals of lead tetraacetate separated on cooling. Stored in a vacuum desiccator over P₂O₅ and KOH for 24h before use.

Lissamine Green B {1-[bis-(4,4'-dimethylaminophenyl)methyl]-2-hydroxynaphthalene-3,6-disulfonic acid sodium salt, Acid Green 50} [3087-16-9] **M 576.6, m >200°(dec), CI 44090, λ_{max} 633nm**. Crystd from EtOH/water (1:1, v/v).

Lissapol C (mainly sodium salt of 9-octadecene-1- sulfate) [2425-51-6]. Refluxed with 95% EtOH, then filtered to remove insoluble inorganic electrolytes. The alcohol solution was then concentrated and the residue was poured into dry acetone. The ppt was filtered off, washed in acetone and dried under vacuum. [Biswas and Mukerji *J Phys Chem* **64** 1 1960].

Lissapol LS (mainly sodium salt of anisidine sulfate) [28903-20-0]. Refluxed with 95% EtOH, then filtered to remove insoluble inorganic electrolytes. The alcohol solution was then concentrated and the residue was poured into dry acetone. The ppt was filtered off, washed in acetone and dried under vacuum. [Biswas and Mukerji *J Phys Chem* **64** 1 1960].

Lithium (metal) [7439-93-2] **M 6.9, m 180.5°, b 1342°, d 0.534**. After washing with pet ether to remove storage oil, lithium was fused at 400° and then forced through a 10-micron stainless-steel filter with argon pressure. It was again melted in a dry-box, skimmed, and poured into an iron distillation pot. After heating under vacuum to 500°, cooling and returning to the dry-box for a further cleaning of its surface, the lithium was distd at 600° using an all-iron distn apparatus [Gunn and Green *J Am Chem Soc* **80** 4782 1958].

Lithium acetate (2H₂O) [546-89-4] **M 102.0, m 54-56°**. Crystallised from EtOH (5mL/g) by partial evaporation.

Lithium aluminium hydride [16853-85-3] **M 37.9, m 125°(dec)**. Extracted with Et₂O, and, after filtering, the solvent was removed under vacuum. The residue was dried at 60° for 3h, under high vacuum [Ruff *J Am Chem Soc* **83** 1788 1961]. **IGNITES in the presence of a small amount of water and reacts EXPLOSIVELY.**

Lithium amide [7782-89-0] **M 23.0, m 380-400°, d^{17.5} 1.178**. Purified by heating at 400° while NH₃ is passed over it in the upper of two crucibles (the upper crucible is perforated). The LiNH₂ will drip into the lower crucible through the holes in the upper crucible. The product is cooled in a stream of NH₃. Protect it from air and moisture, store under N₂ in a clear glass bottle sealed with paraffin. Store small quantities so that all material is used once the bottle is opened. If the colour of the amide is yellow it should be destroyed as it is likely to have oxidised and to **EXPLODE**. On heating above 450° it is decomposed to Li₂NH which is stable up to 750-800°. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer) Vol I* 463 1963; *Inorg Synth* **2** 135 1953.]

Lithium benzoate [553-54-8] **M 128.1**. from EtOH (13mL/g) by partial evaporation.

Lithium borohydride [16949-15-8] **M 21.8, mCrystd 268°, b 380°(dec), d 0.66**. Crystd from Et₂O, and pumped free of ether at 90-100° during 2h [Schaeffer, Roscoe and Stewart *J Am Chem Soc* **78** 729 1956].

Lithium bromide [7550-35-8] **M 86.8, m 550°**. Crystd several times from water or EtOH, then dried under high vacuum for 2 days at room temperature, followed by drying at 100°.

Lithium carbonate [554-13-2] **M 73.9, m 552°, 618°**. Crystd from water. Its solubility decreases as the temperature is raised.

Lithium chloride [7447-41-8] **M 42.4, m 600°, 723°**. Crystd from water (1mL/g) or MeOH and dried for several hours at 130°. Other metal ions can be removed by preliminary crystallisation from hot aqueous 0.01M disodium EDTA. Has also been crystallised from conc HCl, fused in an atmosphere of dry HCl gas, cooled under dry N₂ and pulverised in a dry-box. Kolthoff and Bruckenstein [*J Am Chem Soc* **74** 2529 1952] pptd with ammonium carbonate, washed with Li₂CO₃ five times by decantation and finally with suction, then dissolved in HCl. The LiCl solution was evaporated slowly with continuous stirring in a large evaporating dish, the dry powder being stored (while still hot) in a desiccator over CaCl₂.

Lithium diisopropylamide [4111-54-0] **M 107.1, b 82-84°/atm, 84°/atm, d²² 0.722, flash point -6°**. It is purified by refluxing over Na wire or NaH for 30min and then distilled into a receiver under

N_2 . Because of the low boiling point of the amine a dispersion of NaH in mineral oil can be used directly in this purification without prior removal of the oil. It is **HIGHLY FLAMMABLE**, and is decomposed by air and moisture. [*Org Synth* 50 67 1970.]

Lithium dodecylsulfate [2044-56-6] M 272.3. Recrystd twice from absolute EtOH and dried under vacuum.

Lithium fluoride [7789-24-4] M 25.9, m 842°, 848°, b 1676°, 1681°, d 2.640. Possible impurities are LiCO_3 , H_2O and HF. These can be removed by calcining at red heat, then pulverised with a Pt pestle and stored in a paraffin bottle. Solubility in H_2O is 0.27% at 18°. It volatilises between 1100-1200°. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 235 1963].

Lithium formate (H_2O) [6108-23-2 (H_2O); 556-63-8 (anhydr)] M 70.0, d 1.46. Crystd from hot water (0.5mL/g) by chilling.

Lithium hydride. [7580-67-8] M 7.95, m 680°, d 0.76-0.77. It should be a white powder otherwise replace it. It darkens rapidly on exposure to air and is decomposed by H_2O to give H_2 and LiOH, and reacts with lower alcohols. One gram in H_2O liberates 2.8L of H_2 (Could be explosive).

Lithium hydroxide (H_2O) [1310-66-3 (H_2O); 1310-65-2 (anhydr)] M 42.0, m 471°, d 1.51, pK^{25} 13.82. Crystd from hot water (3mL/g) as the monohydrate. Dehydrated at 150° in a stream of CO_2 -free air.

Lithium iodate [13765-03-2] M 181.9. Crystd from water and dried in a vacuum oven at 60°.

Lithium iodide [10377-51-2] M 133.8, m 469°, b 1171°, d 4.06. Crystd from hot water (0.5mL/g) by cooling in CaCl_2 -ice, or from acetone. Dried under vacuum over P_2O_5 for 1h at 60° and then at 120°.

Lithium methylate (lithium methoxide) [865-34-9] M 38.0. Most probable impurity is LiOH due to hydrolysis by moisture. It is important to keep the sample dry. It can be dried by keeping in a vacuum at 60-80° under dry N_2 using an oil pump for a few hours. Store under N_2 in the cold. It should not have bands above 3000cm^{-1} ; IR has ν_{KBr} 1078, 2790, 2840 and 2930cm^{-1} . [*J Org Chem* 21 156 1956.]

Lithium nitrate [7790-69-4] M 68.9, m 253°, d 2.38. Crystd from water or EtOH. Dried at 180° for several days by repeated melting under vacuum. If it is crystallised from water keeping the temperature above 70°, formation of trihydrate is avoided. The anhydrous salt is dried at 120° and stored in a vac desiccator over CaSO_4 . After 99% salt was recrystd 3 times it contained: metal (ppm) Ca (1.6), K (1.1), Mo (0.4), Na (2.2).

Lithium nitrite (H_2O) [13568-33-7] M 71.0. Crystd from water by cooling from room temperature.

Lithium picrate [18390-55-1] M 221.0. Recrystd three times from EtOH and dried under vacuum at 45° for 48h [D'Aprano and Sesta *J Phys Chem* 91 2415 1987]. The necessary precautions should be taken in case of **EXPLOSION**.

Lithium perchlorate [7791-03-9] M 106.4, pK^{25} -2.4 to -3.1 (for HClO_4). Crystd from water or 50% aq MeOH. Rendered anhydrous by heating the trihydrate at 170-180° in an air oven. It can then be recrystd twice from acetonitrile and again dried under vacuum [Mohammad and Kosower *J Am Chem Soc* 93 2713 1971].

Lithium salicylate [552-38-5] M 144.1. Crystd from EtOH (2mL/g) by partial evaporation.

Lithium sulfate (anhydrous) [10377-48-7] M 109.9, loses H_2O at 130° and m 859°, d 2.21. Crystd from H_2O (4mL/g) by partial evaporation.

Lithium tetrafluoroborate [14283-07-9] M 93.7, pK^{25} 13.82 (Li^+), pK^{25} -4.9 (for HBF_4). Dissolve in THF just below its solubility, filter from insol material and evap to dryness in a vacuum below

50°. Wash the residue with dry Et₂O, and pass dry N₂ gas over the solid and finally heat in an oven at 80-90°. Solubility in Et₂O: 1.9 (1.3)g in 100mL at 25°, in THF: 71g in 100mL at 25°. It is *hygroscopic* and is an **IRRITANT**. [*J Am Chem Soc* 74 5211 1952, 75 1753 1953.]

Lithium thiocyanate (lithium rhodanide) [556-65-0] **M 65.0, pK²⁵ -1.85 (for HSCN)**. It crystallises from H₂O as the dihydrate but on drying at 38-42° it gives the monohydrate. It can be purified by allowing an aqueous soln to crystallise in a vac over P₂O₅. The crystals are collected, dried out in vacuum at 80°/P₂O₅ in a stream of pure N₂ at 110°. [*J Chem Soc* 1245 1936.]

Lithium trimethylsilanolate (trimethylsilanol Li salt) [2004-14-0] **M 96.1, m 120°(dec in air)**. Wash with Et₂O and pet ether. Sublimes at 180°/1mm as fine transparent needles. [*J Org Chem* 17 1555 1952.]

Magnesium [7439-95-4] **M 24.3, m 651°, b 1100°, d 1.739**. Slowly oxidises in moist air and tarnishes. If dark in colour do not use. Shiny solid should be degreased by washing with dry Et₂O, dry and keep in a N₂ atmosphere. It can be activated by adding a crystal of I₂ in the Et₂O before drying and storing.

Magnesium acetate [142-72-3 (*anhydr*); 16674-78-5 (*4H₂O*)] **M 214.5, m 80°**. Crystd from anhydrous acetic acid, then dried under vacuum for 24h at 100°

Magnesium benzoate (3H₂O) [553-70-8] **M 320.6, m ~200°**. Crystd from water (6mL/g) between 100° and 0°.

Magnesium bromide (anhydrous) [7789-48-2] **M 184.1, m 711°, d 3.72**. Crystd from EtOH.

Magnesium chloride (6H₂O) [7791-18-6] **M 203.3, m ~100°(dec), pK₁²⁵ 10.3, pK₂²⁵ 12.2 (for Mg²⁺ hydrolysis)**. Crystd from hot water (0.3mL/g) by cooling.

Magnesium dodecylsulfate [3097-08-3] **M 555.1**. Recrystd three times from EtOH and dried in a vacuum.

Magnesium ethylate (magnesium ethoxide) [2414-98-4] **M 114.4**. Dissolve *ca* 1g of solid in 12.8mL of absolute EtOH and 20mL of dry xylene and reflux in a dry atmosphere (use CaCl₂ in a drying tube at the top of the condenser). Add 10mL of absolute EtOH and cool. Filter solid under dry N₂ and dry in a vacuum. Alternatively dissolve in absolute EtOH and pass through molecular sieves (40 mesh) under N₂, evap under N₂, and store in a tightly stoppered container. [*J Am Chem Soc* 68 889 1964.]

Magnesium D-gluconate [3632-91-5] **M 414.6, [α]₅₄₆²⁰ +13.5°, [α]_D²⁰ +11.3° (c 1, H₂O)**. Cryst from dilute EtOH to give *ca* trihydrate, and then dry at 98° in high vacuum. Insol in EtOH and solubility in H₂O is 16% at 25°.

Magnesium iodate (4H₂O) [7790-32-1] **M 446.2**. Crystd from water (5mL/g) between 100° and 0°.

Magnesium iodide [10377-58-9] **M 278.1, m 634°**. Crystd from water (1.2mL/g) by partial evapn in a desiccator.

Magnesium ionophore I (ETH 1117), (N,N'-diheptyl-N,N'-dimethyl-1,4-butanediamide) [75513-72-3] **M 340.6**. Purified by flash chromatography (at 40 kPa) on silica and eluting with EtOH-hexane (4:1). IR has ν(CHCl₃) 1630cm⁻¹. [*Helv Chim Acta* 63 2271 1980.] It is a good magnesium selectophore compared with Na, K and Ca [*Anal Chem* 52 2400 1980].

Magnesium ionophore II (ETH 5214), [*N,N''*-octamethylene-bis(*N'*-heptyl-*N''*-methyl methylmalonamide)] [119110-37-1] M 538.8. Reagent (ca 700mg) can be purified by flash chromatography on Silica Gel 60 (30g) and eluting with CH₂Cl₂-Me₂CO (4:1). [*Anal Chem* 61 574 1989.]

Magnesium lactate [18917-37-1] M 113.4. Crystd from water (6mL/g) between 100° to 0°.

Magnesium nitrate (6H₂O) [13446-18-9] M 256.4, m ~95°(dec). Crystd from water (2.5mL/g) by partial evapn in a desiccator. After 2 recrystns ACS grade has: metal (ppm) Ca (6.2), Fe (8.4), K (2), Mo (0.6), Na (0.8), Se (0.02).

Magnesium perchlorate (Anhydron, Dehydrite) [10034-81-8 (anhydr)] M 259.2, m >250°, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from water to give the *hexahydrate* M 331.3 [13346-19-0]. Coll, Nauman and West [*J Am Chem Soc* 81 1284 1959] removed traces of unspecified contaminants by washing with small portions of Et₂O and drying in a vac (CARE). The anhydrous salt is commercially available as an ACS reagent, and is as efficient a dehydrating agent as P₂O₅ and is known as "Dehydrite" or "Anhydron". [Smith et al. *J Am Chem Soc* 44 2255 1922 and *Ind Eng Chem* 16 20 1924.] Hygroscopic, Keep in a tightly closed container. **EXPLOSIVE** in contact with organic materials, and is a **SKIN IRRITANT**.

Magnesium succinate [556-32-1] M 141.4. Crystd from water (0.5mL/g) between 100° and 0°.

Magnesium sulfate (anhydrous) [7487-88-9] M 120.4, m 1127°. Crystd from warm H₂O (1mL/g) by cooling. Dry heptahydrate at ~250° until it loses 25% of its wt. Store in a sealed container.

Magnesium trifluoromethanesulfonate [60871-83-2] M 322.4, m >300°. Wash with CH₂Cl₂ and dry at 125°/2h and 3mmHg. [*Tetrahedron Lett* 24 169 1983.]

Magon [3-hydroxy-4-(hydroxyphenylazo)-2-naphthoyl-2,4-dimethylanilide; Xylidyl Blue II] [523-67-1] M 411.5, m 246-247°. Suspend in H₂O and add aqueous NaOH until it dissolves, filter and acidify with dil HCl. Collect the dye, dissolve in hot EtOH (sol is 100mg/L at ca 25°) concentrate to a small volume and allow to cool. Sol in H₂O of the Na salt is 0.4mg/mL. [*Anal Chim Acta* 16 155 1957; *Anal Chem* 28 202 1956.]

Manganese (III) acetate (2H₂O) [19513-05-4] M 268.1, pK²⁵ 0.06 (for Mn³⁺ hydrolysis). Wash the acetate with AcOH then thoroughly with Et₂O and dry in air to obtain the dihydrate. The *anhydrous* salt can be made by stirring vigorously a mixt of the hydrated acetate (ca 6g) and Ac₂O (22.5mL) and heat carefully (if necessary) until the mixture is clear. It is set aside overnight for the material to crystallise. Filter the solid, wash with Ac₂O and dry over P₂O₅. The dihydrate can also be obtained from the *di-* and *tetra-* hydrate mixture of the divalent acetate by adding 500mL of Ac₂O and 48g of the hydrated acetate and refluxing for 20min, then add slowly 8.0g of KMnO₄. After refluxing for an additional 30min, the mixture was cooled to room temperature and 85mL of H₂O added. It should be noted that larger amounts of H₂O change the yield and nature of the manganese acetate and the yields of reactions that use this reagent, e.g. formation of lactones from olefines. The Mn(OAc)₃.2H₂O is then filtered off after 16h, washed with cold AcOH and air dried. [*J Am Chem Soc* 90 5903, 5905 1968, Heiba et al. 91 138 1969.]

Alternatively dissolve the salt (30g) in glacial acetic acid (200mL) by heating and filter. If crystals do not appear, the glass container should be rubbed with a glass rod to induce crystn which occurs within 1h. If not, allow to stand for a few days. Filter the cinnamon brown crystals which have a silky lustre and dry over CaO. Keep away from moisture as it is decomposed by cold H₂O. [Lux in *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II, p 1469 1963; Williams and Hunter *Can J Chem* 54 3830 1976.]

Manganese (II) acetylacetonate [14024-58-9] M 253.2, m ~250°. Purify by stirring 16g of reagent for a few min with 100mL absolute EtOH and filter by suction as rapidly as possible through coarse filter paper. Sufficient EtOH is added to the filtrate to make up for the loss of EtOH and to redissolve any solid that separates. Water (15mL) is added to the filtrate and the solution is evaporated with a stream of N₂ until reduced to half its vol. Cool for a few min and filter off the yellow crystals, dry under a stream of N₂, then in a

vacuum at room temp for 6-8h. These conditions are important for obtaining the *dihydrate*. A vacuum to several mm of Hg or much lower pressure for several days produces the anhydrous complex. The degree of hydration can be established by determining the loss in weight of 100g of sample after heating for 4h at 100° and <20mmHg. The theoretical loss in weight for 2H₂O is 12.5%. Material sublimates at 200°/2mm. It is soluble in heptane, MeOH, EtOH or *C₆H₆ at 30°. [*Inorg Synth* 6 164 1960, 5 105 1957.]

Manganese decacarbonyl Mn₂(CO)₁₀ [10170-69-1] M 390.0, m 151-152°, 154-155°(sealed tube), d²⁵ 1.75. Golden yellow crystals which in the absence of CO begin to decompose at 110°, and on further heating yield a metallic mirror. In the presence of 3000psi of CO it does not decompose on heating to 250°. It is soluble in common organic solvents, insoluble in H₂O, not very stable in air, to heat or UV light. Dissolves in a lot of *C₆H₆ and can be crystallised from it. It distils with steam at 92-100°. It can be purified by sublimation under reduced pressure (<0.5mm) at room temperature to give well formed golden yellow crystals. If the sample is orange coloured this sublimation leads to a mixture of golden-yellow and dark red crystals of the carbonyl and carbonyl iodide respectively which can be separated by hand picking under a microscope. Separate resublimations provide the pure compounds. **POISONOUS** [*J Am Chem Soc* 76 3831 1954, 80 6167 1958, 82 1325 1960].

Manganous acetate (4H₂O) [6156-78-1 (4H₂O); 638-38-0 (anhydr)] M 245.1, m 80°, d 1.59, pK²⁵ 10.59 (for Mn²⁺ hydrolysis). Crystd from water acidified with acetic acid.

Manganous bromide (anhydrous) [13446-03-2] M 214.8, m 695°; 4H₂O [10031-20-6] M 286.8, m 64°(dec). Rose-red deliquescent crystals soluble in EtOH. The H₂O is removed by heating at 100° then in HBr gas at 725° or dry in an atmosphere of N₂ at 200°.

Manganous chloride (4H₂O) [13446-34-9; 7773-01-5 (anhydr)] M 197.9, m 58°, 87.5°, d 2.01. Crystd from water (0.3mL/g) by cooling.

Manganous ethylenebis(dithiocarbamate) [12427-38-2] M 265.3, pK_{Est} ~ 3.0 (for —NCSSH). Crystd from EtOH.

Manganous lactate (3H₂O) [51877-53-3] M 287.1. Crystd from water.

Manganous sulfate (H₂O) [10034-96-5 (H₂O); 15244-36-7 (xH₂O)] M 169.0, d 2.75. Crystd from water (0.9mL/g) at 54-55° by evaporating about two-thirds of the water. Dehydr at >400°.

2-Mercaptopyridine N-oxide sodium salt (pyridinethione or pyrithione sodium salt) [3811-73-2] M 149.1, m ~250°(dec), pK₁ -1.95, pK₂ 4.65. When recrystd from water it assayed as 98.7% based on AgNO₃ titration [Krivis et al. *Anal Chem* 35 966 1963, see also Krivis et al. *Anal Chem* 48 1001 1976; and Barton and Crich *J Chem Soc. Perkin Trans 1* 1603, 1613 1986].

Mercuric acetate [1600-27-7] M 318.7, pK₁²⁵ 2.47, pK₂²⁵ 3.49 (for Hg²⁺ hydrolysis). Crystd from glacial acetic acid. **POISONOUS**.

Mercuric bromide [7789-47-1] M 360.4, m 238.1°. Crystd from hot saturated ethanolic soln, dried and kept at 100° for several hours under vacuum, then sublimed. **POISONOUS**.

Mercuric chloride [7487-94-7] M 271.5, m 276°, b 304°, d 5.6. Crystd twice from distilled water, dried at 70° and sublimed under high vacuum. **POISONOUS**.

Mercuric cyanide [592-04-1] M 252.6, m 320°(dec), d 4.00. Crystd from water. **POISONOUS**.

Mercuric iodide (red) [7774-29-0] M 454.4, m 259°(yellow >130°), b ~350°(subl), d 6.3. Crystd from MeOH or EtOH, and washed repeatedly with distilled water. Has also been mixed thoroughly with excess 0.001M iodine solution, filtered, washed with cold distilled water, rinsed with EtOH and Et₂O, and dried in air. **POISONOUS**.

Mercuric oxide (yellow) [21908-53-2] M 216.6, m 500°(dec). Dissolved in HClO₄ and ppted with NaOH soln.

Mercuric thiocyanate [592-85-8] M 316.8, m 165°(dec), pK²⁵ -1.85 (for HSCN). Recryst from H₂O, and can form various crystal forms depending on conditions. Solubility in H₂O is 0.069% at 25°, but is more soluble at higher temps. Decomposes to Hg above 165°. **Poisonous.** [*J Phys Chem* 35 1128 1931; *Chem Ber* 68 919 1935.]

Mercurous nitrate (2H₂O) [7782-86-7 (2H₂O); 7783-34-8 (H₂O); 10415-75-5 (anhydr)] M 561.2, m 70°(dec), d 4.78, pK²⁵ 2.68 (for Hg₂²⁺ hydrolysis). Solubility in H₂O containing 1% HNO₃ is 7.7%. Recrystd from a warm saturated soln of dilute HNO₃ and cool to room temp slowly to give elongated prisms. Rapid cooling gives plates. Colourless crystals to be stored in the dark. **POISONOUS.** [*J Chem Soc* 1312 1956.]

Mercurous sulfate [7783-36-0] M 497.3, d 7.56. Recrystallise from dilute H₂SO₄, and dry in a vacuum under N₂ and store in the dark. Solubility in H₂O is 0.6% at 25°. **POISONOUS.**

Mercury [7439-97-6] M 200.6, m -38.9°, b 126°/1mm, 184°/10mm, 261°/100mm, 356.9°/atm, d 13.534. After air had been bubbled through mercury for several hours to oxidise metallic impurities, it was filtered to remove coarser particles of oxide and dirt, then sprayed through a 4-ft column containing 10% HNO₃. It was washed with distilled water, dried with filter paper and distilled under vacuum.

Mercury(II) bis(cyclopentadienyl) [18263-08-6] M 330.8. Purified by low-temp recrystn from Et₂O.

Mercury dibromofluorescein {mercurochrome, merobromin, [2',7'-dibromo-4'-(hydroxy-mercurio)-fluorescein di-Na salt]} [129-16-8] M 804.8, m>300°. The Na salt is dissolved in the minimum vol of H₂O, or the free acid suspended in H₂O and dilute NaOH added to cause it to dissolve, filter and acidify with dilute HCl. Collect the ppte wash with H₂O by centrifugation and dry in vacuum. The di Na salt can be purified by dissolving in the minimum volume of H₂O and ppted by adding EtOH, filter, wash with EtOH or Me₂CO and dry in a vacuum. Solubility in 95% EtOH is 2% and in MeOH it is 16%. [*J Am Chem Soc* 42 2355 1920.]

Mercury orange [1-(4-chloromercuriophenylazo)-2-naphthol] [3076-91-3] M 483.3, m 291.5-293°(corr) with bleaching. Wash several times with boiling 50% EtOH and recrystallise from 1-butanol (0.9g/L of boiling alcohol). Fine needles insoluble in H₂O but slightly soluble in cold alcohols, CHCl₃ and soluble in aqueous alkalis. [*J Am Chem Soc* 70 3522 1948.]

Mercury(II) trifluoroacetate [13257-51-7] M 426.6. Recrystd from trifluoroacetic anhydride/trifluoroacetic acid [Lan and Kochi *J Am Chem Soc* 108 6720 1986]. Very **TOXIC** and *hygroscopic*.

Metanil Yellow (3[{4-phenylamino}phenylazo]-benzenesulfonic acid) [587-98-4] M 375.4, pK_{Est} <0. Salted out from water three times with sodium acetate, then repeatedly extracted with EtOH [McGrew and Schneider, *J Am Chem Soc* 72 2547 1950].

(Methoxycarbonylmethyl)triphenylphosphorane [methyl (triphenylphosphoranylidene)-acetate] [2605-67-6] M 334.4, m 162-163°, 169-171°. Cryst by dissolving in AcOH and adding pet ether (b 40-50°) to give colorless plates. UV λ_{max} (A_{1mm}^{1%}): 222nm (865) and 268nm (116) [Isler et al. *Helv Chim Acta* 40 1242 1957].

Methoxycarbonylmethyltriphenylphosphonium bromide [1779-58-4] M 415.3, m 163°, 165-170°(dec). Wash with pet ether (b 40-50°) and recryst from CHCl₃/Et₂O and dry in high vac at 65°. [Isler et al. *Helv Chim Acta* 40 1242 1957; Wittig and Haag *Chem Ber* 88 1654, 1664 1955.]

Methoxymethyl trimethylsilane (trimethylsilylmethyl methyl ether) [14704-14-4] M 118.3, b 83°/740mm, d_4^{25} 0.758, n_D^{25} 1.3878. Forms an azeotrope with MeOH (b 60°). If it contains MeOH (check IR for bands above 3000cm⁻¹) then wash with H₂O and fractionate. A possible impurity could be chloromethyl trimethylsilane (b 97°/740mm). [*J Am Chem Soc* 70 4142 1948.]

1-Methoxy-2-methyl-1-trimethylsiloxypropene (dimethyl ketene methyl trimethylsilyl acetal) [31469-15-5] M 174.3, b 121-122°/0.35mm, 125-126°/0.4mm, 148-150°/atm, d 0.86. Add Et₂O, wash with cold H₂O, dry (Na₂SO₄), filter, evaporate Et₂O, and distil oily residue in a vacuum. [*J Organometal Chem* 46 59 1972.]

trans-1-Methoxy-3-(trimethylsilyloxy)-1,3-butadiene (Danishefsky's diene) [54125-02-9] M 172.3, b 68-69°/14mm, 70-72°/16mm, d 0.885, n 1.4540. It may contain up to 1% of the precursor 4-methoxybut-4-ene-2-one. Easily distd through a Vigreux column in a vac and taking the middle fraction. [Danishefsky and Kitihara *J Am Chem Soc* 96 7807 1974; Danishefsky *Acc Chem Res* 14 400 1981.]

Methylarsonic acid [124-58-3] M 137.9, m 161°, pK_1^{25} 1.54, pK_2^{25} 6.31 [As(OH)₂]. Crystd from abs EtOH.

Methyl dichlorosilane (dichloro methylsilane) [75-54-7] M 115.0, m -92.5°, b 41°/748mm, 40.9°/760mm, 40-45°/atm, d_{27}^{27} 1.105. Impurities are generally other chloromethyl silanes. Distilled through a conventional Stedman column of 20 theoretical plates or more. [Stedman column. A plain tube containing a series of wire-gauze discs stamped into flat, truncated cones and welded together, alternatively base-to-base and edge-to-edge, with a flat disc across each base. Each cone has a hole, alternately arranged, near its base, vapour and liquid being brought into intimate contact on the gauze surfaces (Stedman *Can J Res B* 15 383 1937)]. It should be protected from H₂O by storing over P₂O₅. [*Chem Ber* 52 695 1919; *J Am Chem Soc* 68 9621946.]

Methylmercuric chloride [115-09-3] M 251.1, m 167°. Crystd from absolute EtOH (20mL/g).

Methyl Orange (sodium 4,4'-dimethylaminophenylazobenzenesulfonate) [547-58-0] M 327.3, pK 3.47. Crystd twice from hot water, then washed with a little EtOH followed by diethyl ether. Indicator: pH 3.1 (red) and pH 4.4 (yellow).

Methylphenyl dichlorosilane (dichloro methyl phenylsilane) [149-74-6] M 191.1, b 114-115°/50mm, 202-205°/atm, d 1.17. Purified by fractionation using an efficient column. It hydrolyses *ca* ten times more slowly than methyltrichlorosilane and *ca* sixty times more slowly than phenyltrichlorosilane. [*J Phys Chem* 61 1591 1957].

Methylphosphonic acid [993-13-5] M 96.0, m 104-106°, 105-107°, 108°, pK_1^{25} 2.12, pK_2^{25} 7.29. If it tests for Cl⁻, add H₂O and evaporate to dryness; repeat several times till free from Cl⁻. The residue solidifies to a wax-like solid. Alternatively, dissolve the acid in the minimum volume of H₂O, add charcoal, warm, filter and evaporate to dryness in a vacuum over P₂O₅. [*J Am Chem Soc* 75 3379 1953.] The *di-Na salt* is prepared from 24g of acid in 50mL of dry EtOH and a solution of 23g Na dissolved in 400mL EtOH is added. A white ppt is formed but the mixture is refluxed for 30min to complete the reaction. Filter off and recrystallise from 50% EtOH. Dry crystals in a vacuum desiccator. [*J Chem Soc* 3292 1952.]

Methylphosphonic dichloride [676-97-1] M 132.9, m 33°, 33-37°, b 53-54°/10mm, 64-67°/20.5mm, 86°/44mm, 162°/760mm, d_4^{40} 1.4382. Fractionally redistd until the purity as checked by hydrolysis and acidimetry for Cl⁻ is correct and should solidify on cooling. [*J Chem Soc* 3437 1952; *J Am Chem Soc* 75 3379 1952; for IR see *Can J Chem* 34 1611 1956.]

Methyl Thymol Blue, sodium salt [1945-77-3] M 844.8, ϵ 1.89 x 10⁴ at 435nm, pH 5.5. Starting material for synthesis is Thymol Blue. Purified as for Xylenol Orange on p. 387 in Chapter 4.

Methyl trichlorosilane [75-79-6] **M 149.5, b 13.7°/101mm, 64.3°/710.8mm, 65.5°/745mm, 66.1°/atm, d 1.263, n 1.4110.** If very pure distil before use. Purity checked by ^{29}Si nmr, δ in MeCN is 13.14 with respect to Me_4Si . Possible contaminants are other silanes which can be removed by fractional distillation through a Stedman column of >72 theoretical plates with total reflux and 0.35% take-off (see p. 441). The apparatus is under N_2 at a rate of 12 bubbles/min fed into the line using an Hg manometer to control the pressure. Sensitive to H_2O . [*J Am Chem Soc* 73 4252 1951; *J Org Chem* 48 3667 1983.]

Methyl triethoxysilane [2031-67-6] **M 178.31, b 142-144.5°/742mm, 141°/765mm, 141.5°/775mm, d 0.8911, n 1.3820.** Repeated fractionation in a stream of N_2 through a 3' Heligrad packed Todd column (see p. 174). Hydrolysed by H_2O and yields cyclic polysiloxanes on hydrolysis in the presence of acid in $^*\text{C}_6\text{H}_6$. [*J Am Chem Soc* 77 1292, 3990 1955.]

Methyl trimethoxysilane [1185-55-3] **M 136.2, b 102°/760mm, d 1.3687, n 1.3711.** Likely impurities are 1,3-dimethyltetramethoxy disiloxane (b 31°/1mm) and cyclic polysiloxanes, see methyl triethoxysilane. [*J Org Chem* 16 1400 1952, 20 250 1955.]

N-Methyl-N-trimethylsilylacetamide [7449-74-3] **M 145.3, b 48-49°/11mm, 84°/13mm, 105-107°/35mm (solid at room temp), d 0.90, n 1.4379.** Likely impurity is $\text{Et}_3\text{N.HCl}$ which can be detected by its odour. If it is completely soluble in $^*\text{C}_6\text{H}_6$, then redistil, otherwise dissolve in this solvent, filter and evaporate first in a vacuum at 12mm then fractionate, all operations should be carried out in a dry N_2 atmosphere. [*J Am Chem Soc* 88 3390 1966; *Chem Ber* 96 1473 1963.]

Methyl trimethylsilylacetate [2916-76-9] **M 146.3, b 65-68°/50mm, d 0.89.** Dissolved in Et_2O , shaken with 1M HCl, washed with H_2O , aqueous saturated NaHCO_3 , H_2O again, and dried (a ppt may be formed in the NaHCO_3 soln and should be drawn off and discarded). The solvent is distd off and the residue is fractionated through a good column. IR (CHCl_3) ν 1728 cm^{-1} . [*J Org Chem* 32 3535 1967, 45 237 1980.]

Methyl 2-(trimethylsilyl)propionate [55453-09-3] **M 160.3, b 155-157°/atm, d 0.89.** Dissolve in Et_2O , wash with aqueous NaHCO_3 , H_2O , 0.1M HCl, H_2O again, dry (MgSO_4), evaporated and distil. [*J Chem Soc Perkin Trans 1* 541 1985; *Tetrahedron* 39 3695 1983.]

Methyl triphenoxyphosphonium iodide [17579-99-6] **M 452.2, m 146°.** Gently heat the impure iodide with good grade Me_2CO . The saturated solution obtained is decanted rapidly from undissolved salt and treated with an equal volume of dry Et_2O . The iodide separates as beautiful flat needles which are collected by centrifugation, washed several times with dry Et_2O , and dried in a vacuum over P_2O_5 . For this recrystn it is essential to minimise the time of contact with Me_2CO and to work rapidly and with rigorous exclusion of moisture. If the crude material is to be used, it should be stored under dry Et_2O , and dried and weighed *in vacuo* immediately before use. [*J Chem Soc Perkin Trans 1* 982 1974; *J Chem Soc* 224 1953.]

Methyl triphenylphosphonium bromide [1779-49-3] **M 357.3, m 229-230°(corr), 227-229°, 230-233°.** If the solid is sticky, wash with $^*\text{C}_6\text{H}_6$ and dry in a vacuum over P_2O_5 . [Marvel and Gall *J Org Chem* 24 1494 1959; *Chem Ber* 87 1318 1954; Milas and Priesing *J Am Chem Soc* 79 6295 1957; Wittig and Schöllkopf *Org Synth* 40 66 1960..] The iodide crystd from H_2O has m 187.5-188.5° [*J Chem Soc* 1130 1953; *Justus Liebigs Ann Chem* 580 44 1953].

N-Methyl-N-trimethylsilyl trifluoroacetamide [24589-78-4] **M 199.3, b 78-79°/130mm.** Fractionate through a 40mm Vigreux column. Usually it contains *ca* 1% of methyl trifluoroacetamide and 1% of other impurities which can be removed by gas chromatography or fractionating using a spinning band column. [*J Chromatogr* 42 103 1969, 103 91 1975.]

Methyl vinyl dichlorosilane (dichloro methyl vinyl silane) [124-70-9] **M 141.1, b 43-45.5°/11-11.5mm, 91°/742mm, 92.5°/743.2mm, 92.5-93°/atm, d 1.0917, n 1.444.** Likely impurities are dichloromethylsilane, butadienyl-dichloromethylsilane. Fractionate through a column packed

with metal filings (20 theoretical plates) at atmospheric pressure. [*Izv Akad Nauk SSSR Ser Khim* 1474 1957 and 767 1958.]

Milling Red SWB {1-[4-[4-[4-toluenesulfonyloxy]phenylazo](3,3'-dimethyl-1,1'-biphenyl)-4'-azo]-2-hydroxynaphthalene-6,8-disulfonic acid di-Na salt, Acid Red 114} [6459-94-5] M 830.8, m dec >250°, CI 23635, λ_{\max} ~514nm. Salted out three times with sodium acetate, then repeatedly extracted with EtOH. [McGrew and Schneider *J Am Chem Soc* 72 2547 1950.] See Solochrome Violet R on p. 352 in Chapter 4.

Milling Yellow G [51569-18-7]. Salted out three times with sodium acetate, then repeatedly extracted with EtOH. [McGrew and Schneider *J Am Chem Soc* 72 2547 1950.] See Solochrome Violet R on p. 352 in Chapter 4.

Molybdenum hexacarbonyl [13939-06-5] M 264.0, m 150°(dec), b 156°. Sublimed in a vacuum before use [Connor et al. *J Chem Soc, Dalton Trans* 511 1986].

Molybdenum hexafluoride [7783-77-9] M 209.9, b 35°/760mm. Purified by low-temperature trap-to-trap distillation over predried NaF. [Anderson and Winfield *J Chem Soc, Dalton Trans* 337 1986.]
Poisonous vapours.

Molybdenum trichloride [13478-18-7] M 202.3, m 1027°, d 3.74. Boiled with 12M HCl, washed with absolute EtOH and dried in a vacuum desiccator.

Molybdenum trioxide (molybdenum IV oxide, MoO₃) [1313-27-5] M 143.9, m 795°, d 4.5. Crystd from water (50mL/g) between 70° and 0°.

Monocalcium phosphate (2H₂O) (monobasic) [7789-77-7 (2H₂O); 7757-93-9 (anhydr)] M 154.1, m 200°(dec, loses H₂O at 100°), d 2.2. Crystd from a near-saturated soln in 50% aqueous reagent grade phosphoric acid at 100° by filtering through fritted glass and cooling to room temperature. The crystals were filtered off and this process was repeated three times using fresh acid. For the final crystn the solution was cooled slowly with constant stirring to give thin plate crystals that were filtered off on fritted glass, washed free of acid with anhydrous acetone and dried in a vacuum desiccator [Egan, Wakefield and Elmore, *J Am Chem Soc* 78 1811 1956].

Monoperoxyphthalic acid magnesium salt 6H₂O. (MMPP) [84665-66-7] M 494.7, m ~93°(dec). MMPP is a safer reagent than *m*-chloroperbenzoic acid because it is not as explosive and has advantages of solubility. It is sol in H₂O, low mol wt alcohols, *i*-PrOH and DMF. The product of reaction, Mg phthalate, is sol in H₂O. It has been used in aq phase to oxidise compds in e.g. CHCl₃ and using a phase transfer catalyst e.g. methyltrioctylammonium chloride [Brougham *Synthesis* 1015 1987]. The oxidising activity can be checked (as for perbenzoic acid in Silbert et al. *Org Synth Coll Vol V* 906 1973), and if found to be low it would be best to prepare afresh from phthalic anhydride (1mol), H₂O₂ (1mol) and MgO at 20-25° to give MMPP. [Hignett, European Pat Appl 27 693 1981, *Chem Abstr* 95 168810 1981.]

Naphthalene Scarlet Red 4R [1-(4-sulfonaphthalene-1-azo)-2-hydroxynaphthalene-6,8-disulfonic acid tri-Na salt, New Coccine, Acid Red 18] [2611-82-7] M 604.5, m >250°(dec), CI 16255, λ_{\max} 506nm. Dissolved in the minimum quantity of boiling water, filtered and enough EtOH was added to ppt ca 80% of the dye. This process was repeated until a soln of the dye in aqueous 20% pyridine had a constant extinction coefficient.

Naphthol Yellow S (citronin A, flavianic acid sodium salt, 8-hydroxy-5,7-dinitro-2-naphthalene sulfonic acid disodium salt) [846-70-8] M 358.2, m dec on heating. Greenish yellow powder soluble in H₂O. The *free sulfonic acid* can be recrystd from dil HCl (m 150°) or AcOH-EtOAc (m 148-149.5°). The disodium salt is then obtained by dissolving the acid in two equivalents of aqueous NaOH

[Previous Page](#)

with metal filings (20 theoretical plates) at atmospheric pressure. [*Izv Akad Nauk SSSR Ser Khim* 1474 1957 and 767 1958.]

Milling Red SWB {1-[4-[4-[4-toluenesulfonyloxy]phenylazo](3,3'-dimethyl-1,1'-biphenyl)-4'-azo]-2-hydroxynaphthalene-6,8-disulfonic acid di-Na salt, Acid Red 114} [6459-94-5] M 830.8, m dec >250°, CI 23635, λ_{\max} ~514nm. Salted out three times with sodium acetate, then repeatedly extracted with EtOH. [McGrew and Schneider *J Am Chem Soc* 72 2547 1950.] See Solochrome Violet R on p. 352 in Chapter 4.

Milling Yellow G [51569-18-7]. Salted out three times with sodium acetate, then repeatedly extracted with EtOH. [McGrew and Schneider *J Am Chem Soc* 72 2547 1950.] See Solochrome Violet R on p. 352 in Chapter 4.

Molybdenum hexacarbonyl [13939-06-5] M 264.0, m 150°(dec), b 156°. Sublimed in a vacuum before use [Connor et al. *J Chem Soc, Dalton Trans* 511 1986].

Molybdenum hexafluoride [7783-77-9] M 209.9, b 35°/760mm. Purified by low-temperature trap-to-trap distillation over predried NaF. [Anderson and Winfield *J Chem Soc, Dalton Trans* 337 1986.]
Poisonous vapours.

Molybdenum trichloride [13478-18-7] M 202.3, m 1027°, d 3.74. Boiled with 12M HCl, washed with absolute EtOH and dried in a vacuum desiccator.

Molybdenum trioxide (molybdenum IV oxide, MoO₃) [1313-27-5] M 143.9, m 795°, d 4.5. Crystd from water (50mL/g) between 70° and 0°.

Monocalcium phosphate (2H₂O) (monobasic) [7789-77-7 (2H₂O); 7757-93-9 (anhydr)] M 154.1, m 200°(dec, loses H₂O at 100°), d 2.2. Crystd from a near-saturated soln in 50% aqueous reagent grade phosphoric acid at 100° by filtering through fritted glass and cooling to room temperature. The crystals were filtered off and this process was repeated three times using fresh acid. For the final crystn the solution was cooled slowly with constant stirring to give thin plate crystals that were filtered off on fritted glass, washed free of acid with anhydrous acetone and dried in a vacuum desiccator [Egan, Wakefield and Elmore, *J Am Chem Soc* 78 1811 1956].

Monoperoxyphthalic acid magnesium salt 6H₂O. (MMPP) [84665-66-7] M 494.7, m ~93°(dec). MMPP is a safer reagent than *m*-chloroperbenzoic acid because it is not as explosive and has advantages of solubility. It is sol in H₂O, low mol wt alcohols, *i*-PrOH and DMF. The product of reaction, Mg phthalate, is sol in H₂O. It has been used in aq phase to oxidise compds in e.g. CHCl₃ and using a phase transfer catalyst e.g. methyltrioctylammonium chloride [Brougham *Synthesis* 1015 1987]. The oxidising activity can be checked (as for perbenzoic acid in Silbert et al. *Org Synth Coll Vol V* 906 1973), and if found to be low it would be best to prepare afresh from phthalic anhydride (1mol), H₂O₂ (1mol) and MgO at 20-25° to give MMPP. [Hignett, European Pat Appl 27 693 1981, *Chem Abstr* 95 168810 1981.]

Naphthalene Scarlet Red 4R [1-(4-sulfonaphthalene-1-azo)-2-hydroxynaphthalene-6,8-disulfonic acid tri-Na salt, New Coccine, Acid Red 18] [2611-82-7] M 604.5, m >250°(dec), CI 16255, λ_{\max} 506nm. Dissolved in the minimum quantity of boiling water, filtered and enough EtOH was added to ppt ca 80% of the dye. This process was repeated until a soln of the dye in aqueous 20% pyridine had a constant extinction coefficient.

Naphthol Yellow S (citronin A, flavianic acid sodium salt, 8-hydroxy-5,7-dinitro-2-naphthalene sulfonic acid disodium salt) [846-70-8] M 358.2, m dec on heating. Greenish yellow powder soluble in H₂O. The *free sulfonic acid* can be recrystd from dil HCl (m 150°) or AcOH-EtOAc (m 148-149.5°). The disodium salt is then obtained by dissolving the acid in two equivalents of aqueous NaOH

and evaporating to dryness and drying the residue in a vacuum desiccator. The sodium salt can be recrystd from the minimum volume of H₂O or from EtOH [Dermer and Dermer *J Am Chem Soc* **61** 3302 1939].

1,2-Naphthoquinone-4-sulfonic acid sodium salt (3,4-dihydro-3,4-dioxo-1-naphthlene sulfonic acid sodium salt) [521-24-4] **M 260.2, pK_{Est} <0**. Yellow crystals from aqueous EtOH and dry at 80° *in vacuo*. Solubility in H₂O is 5% [Org Synth Coll Vol III 633 1955; Danielson *J Biol Chem* **101** 507 1933; UV: Rosenblatt et al. *Anal Chem* **27** 1290 1955].

1-Naphthyl phosphate disodium salt [2183-17-7] **M 268.1, pK₁²⁵ 0.97, pK₂²⁵ 5.85 (for free acid)**. The free acid has **m 157-158°** (from Me₂CO/*C₆H₆). The *free acid* is crystd several times by adding 20 parts of boiling *C₆H₆ to a hot solution of 1 part of *free acid* and 1.2 parts of Me₂CO. It has **m 157-158°**. [*J Am Chem Soc* **77** 4002 1955.] The *monosodium salt* was pptd from a soln of the acid phosphate in MeOH by addition of an equivalent of MeONa in MeOH. [*J Am Chem Soc* **72** 624 1950.] See entry on p.550 in Chapter 6.

2-Naphthyl phosphate monosodium salt [14463-68-4] **M 246.1, m 203-205°, pK₁²⁵ 1.25, pK₂²⁵ 5.83 (for free acid)**. Recrystd from H₂O (10mL) containing NaCl (0.4g). The salt is collected by centrifugation and dried in a vac desiccator, **m 203-205°** (partially resolidifies and melts at 244°). Crystd from MeOH (**m 222-223°**). The free acid is recrystd several times by addn of 2.5 parts of hot CHCl₃ to a hot soln of the free acid (1 part) in Me₂CO (1.3 parts), **m 177-178°**. [*J Am Chem Soc.* **73** 5292 1951, **77** 4002 1955.] See entry on p. 551 in Chapter 6.

Neodymium chloride 6H₂O [13477-89-9] **M 358.7, m 124°, pK₁²⁵ 8.43 (for Nd³⁺ hydrolysis)**. Forms large purple prisms from conc solns of dilute HCl. Soluble in H₂O (2.46 parts in 1 part of H₂O) and EtOH.

Neodymium nitrate (6H₂O) [16454-60-7] **M 438.4, m 70-72°**. Crystallises with 5 and 6 molecules of H₂O from conc solutions in dilute HNO₃ by slow evaporation; 1 part is soluble in 10 parts of H₂O.

Neodymium oxide [1313-97-9] **M 336.5, m 2320°**. Dissolved in HClO₄, pptd as the oxalate with doubly recrystd oxalic acid, washed free of soluble impurities, dried at room temperature and ignited in a platinum crucible at higher than 850° in a stream of oxygen [Tobias and Garrett *J Am Chem Soc* **80** 3532 1958].

Neon [7440-01-9] **M 20.2**. Passed through a copper coil packed with 60/80 mesh 13X molecular sieves which is cooled in liquid N₂, or through a column of Ascarite (NaOH-coated silica adsorbent).

Neopentoxylithium [3710-27-8] **M 94.1**. Recrystd from hexane [Kress and Osborn *J Am Chem Soc* **109** 3953 1987].

New Methylene Blue N (2,8-dimethyl-3,6-bis(ethylamino)phenothiazinium chloride 0.5 ZnCl₂) [6586-05-6] **M 416.1, m >200°(dec), pK₁ 3.54, pK₂ 4.82**. Crystd from *benzene/MeOH (3:1).

Nickel (II) acetate (4H₂O) [6018-89-9] **M 248.9, d 1.744, pK₁²⁵ 8.94 (from Ni²⁺ hydrolysis)**. Recryst from aqueous AcOH as the green tetrahydrate. Soluble in 6 parts of H₂O. It forms lower hydrates and should be kept in a well closed container. [*Z Anorg Allg Chem* **343** 92 1966.]

Nickel (II) acetylacetonate [3264-82-2] **M 256.9, m 229-230°, b 220-235°/11mm, d¹⁷ 1.455**. Wash the green solid with H₂O, dry in a vacuum desiccator and recrystallise from MeOH. [*J Phys Chem* **62** 440 1958.] The complex can be conveniently dehydrated by azeotropic distn with toluene and the crystals may be isolated by concentrating the toluene solution. [*J Am Chem Soc* **76** 1970 1954.]

Nickel bromide [13462-88-9] **M 218.5, m 963°(loses H₂O at ~ 200°)**. Crystd from dilute HBr (0.5mL/g) by partial evaporation in a desiccator.

Nickel chloride (6H₂O) [7791-20-0 (6H₂O); 69098-15-3 (xH₂O); 7718-54-9 (anhydr)] **M 237.7**. Crystd from dilute HCl.

Nickel nitrate (6H₂O) [13478-00-7] **M 290.8, m 57°**. Crystd from water (0.3mL/g) by partial evaporation in a desiccator.

Nickelocene [bis-(cyclopentadienyl)nickel II] [1271-28-9] **M 188.9, m 173-174°(under N₂)**. Dissolve in Et₂O, filter and evaporate in a vacuum. Purify rapidly by recrystn from pet ether using a solid CO₂-Me₂CO bath, **m 171-173°(in an evacuated tube)**. Also purified by vacuum sublimation. [*J Am Chem Soc* **76** 1970 1954; *J Inorg Nucl Chem* **295**, 110 1956.]

Nickel (II) phthalocyanine [14055-02-8] **M 571.3, m >300°**. Wash well with H₂O and boiling EtOH and sublime at high vacuum in a slight stream of CO₂. A special apparatus is used (see reference) with the phthalocyanine being heated to red heat. The sublimate is made of needles with an extremely bright red lustre. The powder is dull greenish blue in colour. [*J Chem Soc* 1719 1936.]

Nickel sulfate (7H₂O) [1010-98-1] **M 280.9, m loses 5H₂O at 100°, anhydr m at ~280°**. Crystd from warm water (0.25mL/g) by cooling.

Nickel 5,10,15,20-tetraphenylporphyrin [14172-92-0] **M 671.4, λ_{max} 414(525)nm**. Purified by chromatography on neutral (Grade I) alumina, followed by recrystn from CH₂Cl₂/MeOH [*Yamashita J Phys Chem* **91** 3055 1987].

Niobium (V) chloride [10026-12-7] **M 270.2, m 204.7-209.5°, b ~250°(begins to sublime at 125°), d 2.75**. Yellow very deliquescent crystals which decompose in moist air to give HCl. Should be kept in a dry box flushed with N₂ in the presence of P₂O₅. Wash with CCl₄ and dry over P₂O₅. The yellow crystals usually contain a few small dirty white pellets among the yellow needles. These should be easily picked out. Upon grinding in a dry box, however, they turn yellow. NbCl₅ has been sublimed and fractionated in an electric furnace. [*Inorg Synth* **7** 163 1963; *J Chem Soc* suppl 233 1949.]

Nitric acid [7697-37-2] **M 63.0, m -42°, b 83°, d₂₅ 1.5027, [Constant boiling acid has composition 68% HNO₃ + 32% H₂O, b 120.5°, d 1.41], pK²⁵ -1.27 (1.19)**. Obtained colourless (approx. 92%) by direct distn of fuming HNO₃ under reduced pressure at 40-50° with an air leak at the head of the fractionating column. Stored in a desiccator kept in a refrigerator. Nitrite-free HNO₃ can be obtained by vac distn from urea.

Nitric oxide [10102-43-9] **M 30.0, b -151.8°**. Bubbling through 10M NaOH removes NO₂. It can also be freed from NO₂ by passage through a column of Ascarite followed by a column of silica gel held at -197°K. The gas is dried with solid NaOH pellets or by passing through silica gel cooled at -78°, followed by fractional distillation from a liquid N₂ trap. This purification does not eliminate nitrous oxide. Other gas scrubbers sometimes used include one containing conc H₂SO₄ and another containing mercury. It is freed from traces of N₂ by a freeze and thaw method. **TOXIC**.

p-Nitrobenzenediazonium fluoroborate [456-27-9] **M 236.9**. Crystd from water. **Can be EXPLOSIVE when dry**.

Nitrogen [7727-37-9] **M 28.0, b -195.8°**. Cylinder N₂ can be freed from oxygen by passage through Fieser's soln [which comprises 2g sodium anthraquinone-2-sulfonate and 15g sodium hydrosulfite dissolved in 100mL of 20% KOH [Fieser, *J Am Chem Soc* **46** 2639 1924] followed by scrubbing with saturated lead acetate soln (to remove any H₂S generated by the Fieser soln), conc H₂SO₄ (to remove moisture), then soda-lime (to remove any H₂SO₄ and CO₂). Alternatively, after passage through Fieser's solution, N₂ can be dried by washing with a soln of the metal ketyl from benzophenone and Na wire in absolute diethyl ether. [If ether vapour in N₂ is undesirable, the ketyl from liquid Na-K alloy under xylene can be used.]

Another method for removing O₂ is to pass the nitrogen through a long tightly packed column of Cu turnings, the surface of which is constantly renewed by scrubbing it with ammonia (sg 0.880) soln. The gas is then

passed through a column packed with glass beads moistened with conc H_2SO_4 (to remove ammonia), through a column of packed KOH pellets (to remove H_2SO_4 and to dry the N_2), and finally through a glass trap packed with chemically clean glass wool immersed in liquid N_2 . Nitrogen has also been purified by passage over Cu wool at 723°K and Cu(II) oxide [prepared by heating $\text{Cu}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ at 903°K for 24h] and then into a cold trap at 77°K .

A typical dry purification method consists of a mercury bubbler (as trap), followed by a small column of silver and gold turnings to remove any mercury vapour, towers containing anhydrous CaSO_4 , dry molecular sieves or $\text{Mg}(\text{ClO}_4)_2$, a tube filled with fine Cu turnings and heated to 400° by an electric furnace, a tower containing soda-lime, and finally a plug of glass wool as filter. Variations include tubes of silica gel, traps containing activated charcoal cooled in a Dry-ice bath, copper on Kieselguhr heated to 250° , and Cu and Fe filings at 400° .

2-Nitrophenol-4-arsonic acid (4-hydroxy-3-nitrophenylarsonic acid) [121-19-7] M 263.0, $\text{pK}_{\text{Est}(1)} \sim 4.4$ $\text{As}(\text{O})-(\text{OH})-(\text{O}^-)$, $\text{pK}_{\text{Est}(2)} \sim 7.4$ (phenolic OH), $\text{pK}_{\text{Est}(3)} \sim 7.7$ ($\text{As}(\text{O})-2(\text{O}^-)$). Crystd from water.

1-Nitroso-2-naphthol-3,6-disulfonic acid, di-Na salt, hydrate (Nitroso-R-salt) [525-05-3] M 377.3, m $>300^\circ$, $\text{pK}_{\text{Est}(1)} < 0$ (SO_3^-), $\text{pK}_{\text{Est}(2)} \sim 7$ (OH). Purified by dissolution in aqueous alkali and precipitation by addition of HCl.

Nitrosyl chloride [2696-92-6] M 65.5, b -5.5° . Fractionally distilled at atmospheric pressure in an all-glass, low temperature still, taking the fraction boiling at -4° and storing it in sealed tubes.

Nitrous oxide [10024-97-2] M 44.0, b -88.5° . Washed with conc alkaline pyrogallol solution, to remove O_2 , CO_2 , and NO_2 , then dried by passage through columns of P_2O_5 or Drierite, and collected in a dry trap cooled in liquid N_2 . Further purified by freeze-pump-thaw and distn cycles under vacuum [Ryan and Freeman *J Phys Chem* 81 1455 1977].

Nuclear Fast Red (1-amino-2,4-dihydroxy-5,10-anthraquinone-3-sulfonic acid Na Salt) [6409-77-4] M 357.3, m $>290^\circ(\text{dec})$, λ_{max} 518nm. A soln of 5g of the dye in 250mL of warm 50% EtOH was cooled to 15° for 36h, then filtered on a Büchner funnel, washed with EtOH until the washings were colourless, then with 100mL of diethyl ether and dried over P_2O_5 . [Kingsley and Robnett *Anal Chem* 33 552 1961.]

Octadecyl isonicotinate see hydrogen ionophore IV, ETH 1778 on p. 430.

Octadecyl trichlorosilane [112-04-9] M 387.9, b $159-162^\circ/13\text{mm}$, $185-199^\circ/2-3\text{mm}$, d_4^{30} 0.98. Purified by fractional distillation. [*J Am Chem Soc* 69 2916 1947.]

Octadecyl trimethylammonium bromide [1120-02-1] M 392.5, m $\sim 250^\circ\text{dec}$, $230-240^\circ(\text{dec})$. Cryst from EtOH or H_2O (sol 1 in 1000parts). Very soluble in Me_2CO . [*J Am Chem Soc* 68 714 1946.]

Octamethyl cyclotetrasiloxane [556-67-2] M 296.6, m $17-19^\circ$, 17.58° , 18.5° , b $74^\circ/20\text{mm}$, $176.4^\circ/760\text{mm}$, $d_4^{29.3}$ 0.9451, n_D^{30} 1.3968. Solid has two forms, m 16.30° and 17.65° . Dry over CaH_2 and distil. Further fractionation can be effected by repeated partial freezing and discarding the liquid phase. [*J Am Chem Soc* 76 399 1954, 75 6313 1954.]

Octamethyl trisiloxane [107-51-7] M 236.5, m -80° , b $151.7^\circ/747\text{mm}$, $153^\circ/760\text{mm}$. Distil twice, the middle fraction from the first distillation is again distilled, and the middle fraction of the second distillation is used. [*J Am Chem Soc* 68 358, 691 1946, *J Chem Soc* 1908 1953.]

Octaphenyl cyclotetrasiloxane [546-56-5] M 793.2, m $201-202^\circ$, $203-204^\circ$, b $330-340^\circ/1\text{mm}$. Recryst from AcOH, EtOAc, $^*\text{C}_6\text{H}_6$ or $^*\text{C}_6\text{H}_6/\text{EtOH}$. It forms two stable polymorphs and both

forms as well as the mixture melt at 200-201°. There is a metastable form which melts at 187-189°. [*J Am Chem Soc* 67 2173 1945, 69 488 1947.]

Octyl trichlorosilane [5283-66-9] M 247.7, b 96.5°/10mm, 112°/15mm. 119°/28mm, 229°/760mm, d 1.0744, n 1.4453. Purified by repeated fractionation using a 15-20 theoretical plates glass column packed with glass helices. This can be done more efficiently using a spinning band column. The purity can be checked by analysing for Cl [ca 0.5-1g of sample is dissolved in 25mL of MeOH, diluted with H₂O and titrated with standard alkali. [*J Am Chem Soc* 68 475 1946, 80 1737 1958.]

Orange I [tropaeolin 000 Nr1, 4-(4-hydroxy-1-naphthylazo)benzenesulfonic acid sodium salt] [523-44-4] M 350.3, m >260°(dec). Purified by dissolving in the minimum volume of H₂O, adding, with stirring, a large excess of EtOH. The salt separates as orange needles. It is collected by centrifugation or filtration, washed with absolute EtOH (3 x) and Et₂O (2x) in the same way and dried in a vacuum desiccator over KOH. The free acid can be recrystallised from EtOH. [*Chem Ber* 64 86 1931.] The purity can be checked by titration with titanium chloride [*J Am Chem Soc* 68 2299 1946].

Orange II [tropaeolin 000 Nr2, 4-(2-hydroxy-1-naphthylazo)benzenesulfonic acid sodium salt] [633-96-5] M 350.3. Purification is as for Orange I. The solubility in H₂O is 40g/L at 25°. [*Helv Chim Acta* 35 2579 1952.] Also purified by extracting with a small volume of water, then crystd by dissolving in boiling water, cooling to ca 80°, adding two volumes of EtOH and cooling. When cold, the ppte is filtered off, washed with a little EtOH and dried in air. It can be salted out from aqueous solution with sodium acetate, then repeatedly extracted with EtOH. Meggy and Sims [*J Chem Soc* 2940 1956], after crystallising the sodium salt twice from water, dissolved it in cold water (11mL/g) and conc HCl added to ppte the dye acid which was separated by centrifugation, redissolved and again ppted with acid. After washing the ppte three times with 0.5M acid it was dried over NaOH, recrystd twice from absolute EtOH, washed with a little Et₂O, dried over NaOH and stored over conc H₂SO₄ in the dark.

Orange G (1-phenylazo-2-naphthol-6,8-disulfonic acid di-Na salt) [1936-15-8] M 452.4, pK_{Est} ~ 9. Recryst from 75% EtOH, dry for 3h at 110° and keep in a vacuum desiccator over H₂SO₄. The free acid crystallises from EtOH or conc HCl in deep red needles with a green reflex. [*J Am Chem Soc* 48 2483 1923, *J Chem Soc* 292 1938.]

Orange RO {acid orange 8, 1,8-[bis(4-n-propyl-3-sulfophenyl-1-amino)]anthra-9,10-quinone di-Na salt} [5850-86-2] M 364.4, CI 15575, λ_{max} 490nm. Salted out three times with sodium acetate, then repeatedly extracted with EtOH.

Osmium tetroxide (osmic acid) [20816-12-0] M 524.2, m 40.6°, b 59.4°/60mm, 71.5°/100mm, 109.3°/400mm, 130°/760mm, d 5.10, pK₁^{2.5} 7.2, pK₂^{2.5} 12.2, pK₃^{2.5} 13.95, pK₄^{2.5} 14.17 (H₄OsO₆). It is VERY TOXIC and should be manipulated in a very efficient fume cupboard. It attacks the eyes severely (use also face protection) and is a good oxidising agent. It is volatile and has a high vapour pressure (11mm) at room temp. It sublimes and dists well below its boiling point. It is sol in *C₆H₆, H₂O (7.24% at 25°), CCl₄ (375% at 25°), EtOH and Et₂O. It is estimated by dissolving a sample in a glass stoppered flask containing 25mL of a solution of KI (previously saturated with CO₂) and acidified with 0.35M HCl. After gentle shaking in the dark for 30min, the solution is diluted to 200mL with distilled H₂O satd with CO₂ and titrated with standard thiosulfate using Starch indicator. This method is not as good as the gravimetric method. Hydrazine hydrochloride (0.1 to 0.3g) is dissolved in 3M HCl (10mL) in a glass stoppered bottle. After warming to 55-65°, a weighed sample of OsO₄ solution is introduced, and the mixture is digested on a water bath for 1h. The mixture is transferred to a weighed glazed crucible and evaporated to dryness on a hot plate. A stream of H₂ is started through the crucible and the crucible is heated over a burner for 20-30 min. The stream of H₂ is continued until the crucible is cooled to room temperature, and then the H₂ is displaced by CO₂ in order to avoid rapid combustion of H₂. Finally the crucible is weighed. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1603 1965; *J Am Chem Soc* 60 1822 1938.]

§ Available commercially on a polymer support.

Oxygen [7782-44-7] M 32.00, m -218.4°, b -182.96°, d-183 1.149, d-252.5 1.426. Purified by passage over finely divided platinum at 673°K and Cu(II) oxide (see under nitrogen) at 973°, then condensed in liquid N₂-cooled trap. **HIGHLY EXPLOSIVE in contact with organic matter.**

Palladium (II) acetate [3375-31-3] M 244.5, m 205°dec, pK₁²⁵ 1.0, pK₂²⁵ 1.2 (for Pd₂²⁺). Recrystd from CHCl₃ as purple crystals. It can be washed with AcOH and H₂O and dried in air. Large crystals can be obtained by dissolving in *C₆H₆ and allowing to evaporate slowly at room temp. It forms green adducts with nitrogen donors, dissolved in KI soln but is insoluble in aqueous saturated NaCl, and NaOAc. Soluble in HCl to form PdCl₄²⁻. [*Chem Ind (London)* 544 1964; *J Chem Soc* 658 1970.]

Palladium (II) acetyl acetone [14024-61-4] M 304.6. Recrystd from *C₆H₆-pet ether and sublimed *in vacuo*. It is soluble in heptane, *C₆H₆ (1.2% at 20°, 2.2 at 40°), toluene (0.56% at 20°, 1.4% at 40°) and acetylacetone (1.2% at 20°, 0.05% at 40°). [*J Inorg Nucl Chem* 5 295 1957/8; *Inorg Synth* 5 105 1957.]

Palladium (II) chloride [7647-10-1] M 177.3, m 678-680°. The anhydrous salt is insoluble in H₂O and dissolves in HCl with difficulty. The dihydrate forms red *hygroscopic* crystals that are readily reduced to Pd. Dissolve in conc HCl through which dry Cl₂ was bubbled. Filter this solution which contains H₂PdCl₄ and H₂PdCl₆ and on evaporation yields a residue of pure PdCl₂. [*Handbook of Preparative Inorganic Chemistry (Ed Brauer)* Vol 2 1582 1965; *Org Synth Coll Vol III* 685 1955.]

Palladium (II) cyanide [2035-66-7] M 158.1. A yellow solid, wash well with H₂O and dry in air. [*Inorg Chem* 2 245 1946]. **POISONOUS.**

Palladium (II) trifluoroacetate [42196-31-6] M 332.4, m 210°(dec). Suspend in trifluoroacetic acid and evaporate on a steam bath a couple of times. The residue is then dried in vacuum (40-80°) to a brown powder. [*J Chem Soc* 3632 1965; *J Am Chem Soc* 102 3572 1980.]

Pentafluorophenyl dimethylchlorosilane (Flophemesyl chloride) [20082-71-7] M 260.7, b 89-90°/10mm, d₄³⁰ 1.403, n_D³⁰ 1.447. If goes turbid on cooling due to separation of some LiCl, then dissolve in Et₂O, filter and fractionate. [*J Chromatogr* 89 225 1974, 132 548 1977.]

Perchloric acid [7601-90-3] M 100.5, d 1.665, pK²⁵ -2.4 to -3.1 (HClO₄). The 72% acid has been purified by double distn from silver oxide under vacuum: this frees the acid from metal contamination. Anhydrous acid can be obtained by adding gradually 400-500mL of oleum (20% fuming H₂SO₄) to 100-120mL of 72% HClO₄ in a reaction flask cooled in an ice-bath. The pressure is reduced to 1mm (or less), with the reaction mixture at 20-25°. The temperature is gradually raised during 2h to 85°, the distillate being collected in a receiver cooled in Dry-ice. For further details of the distillation apparatus [see Smith *J Am Chem Soc* 75 184 1953]. **HIGHLY EXPLOSIVE, a strong protective screen should be used at all times.**

Phenylarsonic acid (benzenearsonic acid) [98-05-5] M 202.2, m 155-158°(dec), pK₁²⁵ 3.65, pK₂²⁵ 8.77. Crystd from H₂O (3mL/g) between 90° and 0°.

Phenylboric acid (benzeneboronic acid) [98-80-6] M 121.9, m ~43°, 215-216° (anhydride), 217-220°, pK₁²⁵ 8.83. It recrystallises from H₂O, but can convert spontaneously to benzeneboronic anhydride or phenylboroxide on standing in dry air. Possible impurity is dibenzeneborinic acid which can be removed by washing with pet ether. Heating in an oven at 110°/760mm 1h converts it to the *anhydride* m 214-216°. Its solubility in H₂O is 1.1% at 0° and 2.5% at 25° and in EtOH it is 10% (w/v). [Gilman and Moore *J Am Chem Soc* 80 3609 1958.] If the acid is required, not the anhydride, the acid (from recrystallisation in H₂O) is dried in a slow stream of air saturated with H₂O. The anhydride is converted to the acid by recrystallisation from H₂O. The acid gradually dehydrates to the anhydride if left in air at room temperature with 30-40% relative humidity. The melting point is usually that of the anhydride because the acid dehydrates before it melts [Washburn et al. *Org Synth Coll Vol IV* 68 1963].

Phenyl dimethyl chlorosilane (DMPCl, chlorodimethylphenylsilane) [768-33-2] M 170.7, b 79°/15mm, 85-87°/32mm, 196°/760mm, d 1.017, n 1.509. Fractionate through a 1.5 x 18 inch column packed with stainless steel helices; or a spinning band column. [*J Am Chem Soc* 74 386 1952; 70 1115 1948; *J Chem Soc* 494 1953.] Used for standardising MeLi or MeMgBr which form Me₃PhSi, estimated by GC [Maienthal et al. *J Am Chem Soc* 76 6392 1954; House and Respass *J Organomet Chem* 4 95 1965.] **TOXIC** and **MOISTURE SENSITIVE**.

1,2-Phenylene phosphorochloridate (2-chloro-1,3,2-benzodioxaphosphole-2-oxide) [1499-17-8] M 190.5, m 52°, 58-59°, 59-61°, b 80-81°/1-2mm, 118°/10mm, 122°/12mm, 125°/16mm, 155°/33mm. Distil in a vacuum, sets to a colourless solid. It is soluble in pet ether, *benzene and slightly soluble in Et₂O. [*J Chem Soc (C)* 2092 1970; *Justus Liebigs Ann Chem* 454 109 1927.]

Phenylmercuric hydroxide [100-57-2] M 294.7, m 195-203°. Crystd from dilute aqueous NaOH.

Phenylmercuric nitrate [8003-05-2] M 634.4, m 178-188°. Crystd from water.

Phenylphosphinic acid [benzenephosphinic acid, PhPH(O)(OH)] [1779-48-2] M 142.1, m 70°, 71°, 83-85°, 86°, pK₁²⁵ 1.75. Cryst from H₂O (sol 7.7% at 25°). Purified by placing the solid in a flask covered with dry Et₂O, and allowed to stand for 1 day with intermittent shaking. Et₂O was decanted off and the process repeated. After filtration, excess Et₂O was removed in vacuum. [*Justus Liebigs Ann Chem* 181 265 1876; *Anal Chem* 29 109 1957; NMR: *J Am Chem Soc* 78 5715 1956.]

Phenylphosphonic acid [1571-33-1] M 158.1, m 164.5-166°, pK₂²⁵ 7.43 (7.07). Best recryst from H₂O by concentrating an aqueous soln to a small volume and allowing to crystallise. Wash the crystals with ice cold H₂O and dry in a vacuum desiccator over H₂SO₄. [*J Am Chem Soc* 78 1045 1954.] pK²⁵ values in H₂O are 7.07, and in 50% EtOH 8.26. [*J Am Chem Soc* 75 2209 1953.] [IR: *Anal Chem* 23 853 1951.]

Phenylphosphonic dichloride (P,P-dichlorophenyl phosphine oxide) [824-72-6] M 195.0, b 83-84°/1mm, 135-136°/23mm, d₄³⁰ 1.977, n_D³⁰ 1.5578. Fractionally distilled using a spinning band column. [*J Am Chem Soc* 76 1045 1954; NMR: *J Am Chem Soc* 78 3557, 5715 1956; IR: *Anal Chem* 23 853 1951.]

Phenylphosphonous acid [PhP(OH)₂] [121-70-0] M 141.1, m 71°, pK_{Est} <0, pK¹⁷ 2.1. Crystd from hot H₂O.

Phenylphosphonous dichloride (P,P-dichloro phenyl phosphine) [644-97-3] M 179.0, b 68-70°/1mm, 224-226°/atm, d₄³⁰ 1.9317, n_D³⁵ 1.5962. Vacuum distilled by fractionating through a 20cm column packed with glass helices (better use a spinning band column) [*J Am Chem Soc* 73 755 1951; NMR: *J Am Chem Soc* 78 3557 1956; IR: *Anal Chem* 23 853 1951]. It forms a yellow Ni complex: Ni(C₆H₅Cl₂P)₄ (m 91-92°, from H₂O) [*J Am Chem Soc* 79 3681 1957] and a yellow complex with molybdenum carbonyl: Mo(CO)₃.(C₆H₅Cl₂P)₃ (m 106-110°dec) [*J Chem Soc* 2323 1959].

Phenyl phosphoro chloridate (diphenyl phosphoryl chloride) [2524-64-3] M 268.6, b 141°/1mm, 194°/13mm, 314-316°/272mm, d₄³⁰ 1.2960, n_D³⁵ 1.5490. Fractionally distd in a good vac, better use a spinning band column. [*J Am Chem Soc* 81 3023 1959; IR: *J Chem Soc* 475, 481 1952.]

Phenyl phosphoryl dichloride [770-12-7] M 211.0, m -1°, b 103-104°/2mm, 110-111°/10mm, 130-134°/21mm, 241-243°/atm, d₄³⁰ 1.4160, n_D³⁰ 1.5216. Fractionally distilled under as good a vacuum as possible using an efficient fractionating column or a spinning band column. It should be redistilled if the IR is not very good [IR: *J Chem Soc* 475, 481 1952; *J Am Chem Soc* 60 750 1938, 80 727 1958]. **HARMFUL VAPOURS**.

Phenylthio trimethylsilane (trimethyl phenylthio silane) [4551-15-9] M 182.4, b 95-99°/12mm, d₄³⁰ 0.97. Purification is as for phenyl trimethyl silylmethyl sulfide on p. 450.

Phenyl trimethoxysilane (trimethoxysilyl benzene) [2996-92-1] **M 198.3, b 103°/20mm, 130.5-131°/45mm, d_4^{35} 1.022, n_D^{35} 1.4698.** Fractionate through an efficient column but note that it forms an azeotrope with MeOH which is a likely impurity. [*J Am Chem Soc* **75** 2712 1953; *J Gen Chem USSR (Engl Transl)* **25** 1079 1955.]

Phenyl trimethylsilane (trimethylphenylsilane) [768-32-1] **M 150.3, b 67.3°/20mm, 98-99°/80mm, 170.6°/738mm, d_4^{25} 0.8646.** See trimethylphenylsilane on p. 489.

Phenyl trimethylsilylmethyl sulfide [(phenylthiomethyl)trimethylsilane] [17873-08-4] **M 196.4, b 48°/0.04mm, 113-115°/12mm, 158.5°/52mm, d_4^{30} 0.9671, n_D^{30} 1.5380.** If the sample is suspect then add H₂O, wash with 10% aqueous NaOH, H₂O again, dry (anhydrous CaCl₂) and fractionally distil through a 2ft column packed with glass helices. [*J Am Chem Soc* **76** 3713 1954.]

Phosgene [75-44-5] **M 98.9, b 8.2°/756mm.** Dried with Linde 4A molecular sieves, degassed and distilled under vacuum. This should be done in a closed system such as a vacuum line. **HIGHLY TOXIC, should not be inhaled. If it is inhaled operator should lie still and made to breath ammonia vapour which reacts with phosgene to give urea.**

Phosphine [7803-51-2] **M 34.0, m -133°, b -87.7°, pK -14, pK_b 28.** Best purified in a gas line (in a vacuum) in an efficient fume cupboard. It is spontaneously flammable, has a strong odour of decayed fish and is **POISONOUS**. The gas is distd through solid KOH towers (two), through a Dry ice-acetone trap (-78°, to remove H₂O, and P₂H₄ which causes spontaneous ignition with O₂), then through two liquid N₂ traps (-196°), followed by distn into a -126° trap (Dry ice-methylcyclohexane slush), allowed to warm in the gas line and then seal in ampoules preferably under N₂. IR: ν 2327 (m), 1121 (m) and 900 (m) cm⁻¹. [Klement in *Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I, pp. 525-530 1963; Gokhale and Jolly *Inorg Synth* **9** 56 1967.] PH₃ has also been absorbed into a soln of cuprous chloride in hydrochloric acid (when CuCl.PH₃ is formed). PH₃ gas is released when the soln is heated and the gas is purified by passage through KOH pellets and over then P₂O₅. The solubility is 0.26mL/1 mL of H₂O at 20° and a crystalline hydrate is formed on releasing the pressure on an aq soln.

Phosphonitrilic chloride (tetramer) [1832-07-1] **M 463.9.** Purified by zone melting, then crystd from pet ether (b 40-60°) or *n*-hexane. [van der Huizen et al. *J Chem Soc, Dalton Trans* 1317 1986.]

Phosphonitrilic chloride (trimer) (hexachlorocyclotriphosphazine) [940-71-6] **M 347.7, m 112.8°, 113-114°.** Purified by zone melting, by crystallisation from pet.ether, *n*-hexane or *benzene, and by sublimation. [van der Huizen et al. *J Chem Soc, Dalton Trans* 1311 1986; Meirovitch et al. *J Phys Chem* **88** 1522 1984; Alcock et al. *J Am Chem Soc* **106** 5561 1984; Winter and van de Grampel *J Chem Soc, Dalton Trans* 1269 1986.]

Phosphoric acid [7664-38-2] **M 98.0, m 42.3°, pK₁²⁵ 2.15, pK₂²⁵ 7.21, pK₃²⁵ 12.33.** Pyrophosphate can be removed from phosphoric acid by diluting with distilled H₂O and refluxing overnight. By cooling to 11° and seeding with crystals obtained by cooling a few millilitres in a Dry-ice/acetone bath, 85% orthophosphoric acid crystallises as H₃PO₄.H₂O. The crystals are separated using a sintered glass filter. It has pK_a²⁵ values of 2.15, 7.20 and 12.37 in H₂O.

Phosphorus (red) [7723-14-0] **M 31.0, m 590°/43atm, ignites at 200°, d 2.34.** Boiled for 15min with distilled H₂O, allowed to settle and washed several times with boiling H₂O. Transferred to a Büchner funnel, washed with hot H₂O until the washings are neutral, then dried at 100° and stored in a desiccator.

Phosphorus (white) [7723-14-0] **M 31.0, m 590, d 1.82.** Purified by melting under dilute H₂SO₄⁻ dichromate (possible **carcinogen**) mixture and allowed to stand for several days in the dark at room temperature. It remains liquid, and the initial milky appearance due to insoluble, oxidisable material gradually disappears. The phosphorus can then be distilled under vacuum in the dark [Holmes *Trans Faraday Soc* **58** 1916]

1962]. Other methods include extraction with dry CS_2 followed by evaporation of the solvent, or washing with 6M HNO_3 , then H_2O , and drying under vacuum. **POISONOUS.**

Phosphorus oxychloride [10025-87-3] **M 153.3, b 105.5°, n 1.461, d 1.675.** Distilled under reduced pressure to separate from the bulk of the HCl and the phosphoric acid, the middle fraction being distilled into ampoules containing a little purified mercury. These ampoules are sealed and stored in the dark for 4-6 weeks with occasional shaking to facilitate reaction of any free chloride with the mercury. The POCl_3 is then again fractionally distd and stored in sealed ampoules in the dark until used [Herber *J Am Chem Soc* **82** 792 1960]. Lewis and Sowerby [*J Chem Soc* 336 1957] refluxed their distilled POCl_3 with Na wire for 4h, then removed the Na and again distilled. *Use Na only with almost pure POCl_3 to avoid explosions.* **HARMFUL VAPOURS.**

Phosphorus pentabromide [7789-69-7] **M 430.6, m <100°, b 106°(dec).** Dissolved in pure nitrobenzene at 60°, filtering off any insoluble residue on to sintered glass, then crystallised by cooling. Washed with dry Et_2O and removed the ether in a current of dry N_2 . (All manipulations should be performed in a dry-box.) [Harris and Payne *J Chem Soc* 3732 1958]. Fumes in moist air because of hydrolysis. **HARMFUL VAPOURS.**

Phosphorus pentachloride [10026-13-8] **M 208.2, m 179-180°(sublimes).** Sublimed at 160-170° in an atmosphere of chlorine. The excess chlorine was then displaced by dry N_2 gas. All subsequent manipulations were performed in a dry-box [Downs and Johnson *J Am Chem Soc* **77** 2098 1955]. Fumes in moist air. **HARMFUL VAPOURS.**

Phosphorus pentasulfide [1314-80-3] **M 444.5, m 277-283°.** Purified by extraction and crystallisation with CS_2 , using a Soxhlet extractor. Liberates H_2S in moist air. **HARMFUL VAPOURS.**

Phosphorus pentoxide [1314-56-3] **M 141.9, m 562°, b 605°.** Sublimed at 250° under vacuum into glass ampoules. Fumes in moist air and reacts violently with water. **HARMFUL VAPOURS and attacks skin.**

Phosphorus sesquisulfide P_4S_3 [1314-85-8] **M 220.1, m 172°.** Extracted with CS_2 , filtered and evapd to dryness. Placed in H_2O , and steam was passed through for an hour. The H_2O was then removed, the solid was dried, followed by crystallisation from CS_2 [Rogers and Gross *J Am Chem Soc* **74** 5294 1952].

Phosphorus sulfochloride (phosphorus thiochloride) [3982-91-0] **M 169.4, m -35°, b 122-124°, 125°(corr), d_4^{30} 1.64, n_D^{30} 1.556.** Possible impurities are PCl_5 , H_3PO_4 , HCl and AlCl_3 . Gently mix with H_2O to avoid a heavy emulsion, the product decolorises immediately and settles to the bottom layer. **HARMFUL VAPOURS.**

Phosphorus tribromide [7789-60-8] **M 270.7, m -41.5°, b 168-170°/725mm, 171-173°/atm, 172.9°/760mm(corr), d_4^{30} 2.852.** It is decomposed by moisture, should be kept dry and is *corrosive*. Purified by distillation through an efficient fractionating column (see Whitmore and Lux *J Am Chem Soc* **54** 3451) in a slow stream of dry N_2 , i.e. under strictly dry conditions. [*Inorg Synth* **2** 147 1946; *Org Synth Col Vol II* 358 1943.] Dissolve in CCl_4 , dry over CaCl_2 , filter and distil. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) vol I 532 1963.] Store in sealed ampoules under N_2 and kept away from light. **HARMFUL VAPOURS.**

Phosphorus trichloride [7719-12-2] **M 137.3, b 76°, n 1.515, d 1.575.** Heated under reflux to expel dissolved HCl, then distilled. It has been further purified by vacuum fractionation several times through a -45° trap into a receiver at -78°. **HARMFUL VAPOURS.**

Phosphorus triiodide [13455-01-1] **M 411.7, m 61°.** Decomposes in moist air and must be kept in a desiccator over CaCl_2 . It is crystallised from sulfur-free CS_2 otherwise the **m** decreases to *ca* 55°. It is best prepared freshly. [*J Am Chem Soc* **49** 307 1927; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) vol I 541 1963.] **HARMFUL VAPOURS.**

12-Phosphotungstic acid [12501-23-4] M 2880.2, m -96° . A few drops of conc HNO_3 were added to 100g of phosphotungstic acid dissolved in 75mL of water, in a separating funnel, and the soln was extracted with diethyl ether. The lowest of the three layers, which contained a phosphotungstic acid-ether complex, was separated, washed several times with 2M HCl, then with water and again extracted with ether. Evaporation of the ether, under vacuum with mild heating on a water bath gave crystals which were dried under vacuum and ground [Matijevic and Kerker, *J Am Chem Soc* **81** 1307 1959].

Phthalocyanine [574-93-6] M 514.6. Purified by sublimation (two to three times) in an argon flow at 300-400Pa. Similarly for the Cu(II), Ni(II), Pb(II), VO(II) and Zn(II) phthalocyanine complexes.

Platinum (II) acetylacetonate [15170-57-7] M 393.3, m 249-252 $^{\circ}$. Recrystd from $^*\text{C}_6\text{H}_6$ as yellow crystals and dried in air or in a vacuum desiccator. [*Chem Ber* **34** 2584 1901.]

Platinum (II) chloride [10025-65-7] M 266.0, d 5.87. It is purified by heating at 450 $^{\circ}$ in a stream of Cl_2 for 2h. Some sublimation occurs because the PtCl_2 sublimes completely at 560 $^{\circ}$ as red (almost black) needles. This sublimate can be combined to the bulk chloride and while still at ca 450 $^{\circ}$ it should be transferred to a container and cooled in a desiccator. A probable impurity is PtCl_4 . To test for this add a few drops of H_2O (in which PtCl_4 is soluble) to the salt, filter and add an equal volume of saturated aqueous NH_4Cl to the filtrate. If no ppt is formed within 1 min then the product is pure. If a ppt appears then the whole material should be washed with small volumes of H_2O until the soluble PtCl_4 is removed. The purified PtCl_2 is partly dried by suction and then dried in a vacuum desiccator over P_2O_5 . It is insoluble in H_2O but soluble in HCl to form chloroplatinic acid (H_2PtCl_4) by disproportionation. [*Inorg Synth* **6** 209 1960.]

Polystyrenesulfonic acid sodium salt (-CH₂CH(C₆H₄SO₃Na)-) [25704-18-1]. Purified by repeated pptn of the sodium salt from aqueous soln by MeOH, with subsequent conversion to the free acid by passage through an Amberlite IR-120 ion-exchange resin. [Kotin and Nagasawa *J Am Chem Soc* **83** 1026 1961.] Recrystd from EtOH. Also purified by passage through cation and anion exchange resins in series (Rexyn 101 cation exchange resin and Rexyn 203 anion exchange resin), then titrated with NaOH to pH 7. The sodium form of polystyrenesulfonic acid ppted by addition of 2-propanol. Dried in a vac oven at 80 $^{\circ}$ for 24h, finally increasing to 120 $^{\circ}$ prior to use. [Kowblansky and Ander *J Phys Chem* **80** 297 1976.]

Pontacyl Carmine 2G (Acid Red 1, Amido Naphthol Red G, Azophloxine, 1-acetamido-8-hydroxy-7-phenylazonaphthalene-3,7-disulfonic acid di-Na salt) [3734-67-6] M 510.4, CI 18050, λ_{max} 532nm. Salted out three times with sodium acetate, then repeatedly extracted with EtOH. See Solochrome Violet R on p. 352 in Chapter 4. [McGrew and Schneider *J Am Chem Soc* **72** 2547 1950.]

Pontacyl Light Yellow GX [Acid Yellow 17, 1-(2,5-dichloro-4-sulfophenyl)-3-methyl-4-(4-sulfophenylazo)-5-hydroxypyrazole di-Na Salt] [6359-98-4] M 551.3, CI 18965, λ_{max} 400nm. Purification as for Pontacyl Carmine 2G above.

Potassium (metal) [7440-09-7] M 39.1, m 62.3 $^{\circ}$, d 0.89. Oil was removed from the surface of the metal by immersion in *n*-hexane and pure Et_2O for long periods. The surface oxide was next removed by scraping under ether, and the potassium was melted under vacuum. It was then allowed to flow through metal constrictions into tubes that could be sealed, followed by distillation under vacuum in the absence of mercury vapour (see Sodium). **EXPLOSIVE IN WATER.**

Potassium acetate [127-08-2] M 98.2, m 292 $^{\circ}$, d 1.57, pK^{25} 16 (for aquo K^+). Crystd three times from water-ethanol (1:1) dried to constant weight in a vacuum oven, or crystd from anhydrous acetic acid and pumped dry under vacuum for 30h at 100 $^{\circ}$.

Potassium 4-aminobenzoate [138-84-1] M 175.2. Crystd from EtOH.

Potassium antimonyltartrate (H₂O) [28300-74-5] M 333.9, $[\alpha]_{\text{D}}$ +141 $^{\circ}$ (c 2, H₂O). Crystd from water (3mL/g) between 100 $^{\circ}$ and 0 $^{\circ}$. Dried at 100 $^{\circ}$.

Potassium benzoate [582-25-2] M 160.2. Crystd from water (1mL/g) between 100° and 0°.

Potassium bicarbonate [298-14-6] M 100.1. Crystd from water at 65-70° (1.25mL/g) by filtering, then cooling to 15°. During all operations, CO₂ is passed through the stirred mixture. The crystals, sucked dry at the pump, are washed with distilled water, dried in air and then over H₂SO₄ in an atmosphere of CO₂.

Potassium biiodate [13455-24-8] M 389.9. Crystd three times from hot water (3mL/g), stirred continuously during each cooling. After drying at 100° for several hours, the crystals are suitable for use in volumetric analysis.

Potassium bisulfate [7646-93-7] M 136.2, m 214°. Crystd from H₂O(1mL/g) between 100° and 0°.

Potassium borohydride [13762-51-1] M 53.9, m ~500°(dec). Crystd from liquid ammonia.

Potassium bromate [7758-01-2] M 167.0, m 350°(dec at 370°), d 3.27. Crystd from distilled H₂O(2mL/g) between 100° and 0°. To remove bromide contamination, a 5% soln in distilled H₂O, cooled to 10°, has been bubbled with gaseous chlorine for 2h, then filtered and extracted with reagent grade CCl₄ until colourless and odourless. After evaporating the aqueous phase to about half its volume, it was cooled again slowly to about 10°. The crystalline KBrO₃ was separated, washed with 95% EtOH and vacuum dried [Boyd, Cobble and Wexler *J Am Chem Soc* 74 237 1952]. Another way to remove Br⁻ ions was by stirring several times in MeOH and then dried at 150° [Field and Boyd *J Phys Chem* 89 3767 1985].

Potassium bromide [7758-02-3] M 119.0, m 734°, d 2.75. Crystd from distilled water (1mL/g) between 100° and 0°. Washed with 95% EtOH, followed by Et₂O. Dried in air, then heated at 115° for 1h, pulverised and heated in a vacuum oven at 130° for 4h. Has also been crystd from aqueous 30% EtOH, or EtOH, and dried over P₂O₅ under vacuum before heating in an oven.

Potassium tert-butoxide [865-47-4] M 112.2. It sublimes at 220°/1 Torr. Last traces of *tert*-BuOH are removed by heating at 150-160°/2mm for 1h. It is best prepared fresh; likely impurities are *tert*-BuOH, KOH and K₂CO₃ depending on exposure to air. Its solubility at 25°-26° in hexane, toluene, Et₂O, and THF is 0.27%, 2.27%, 4.34% and 25.0% respectively. [*J Am Chem Soc* 78 5938, 4364 1956.]

Potassium carbonate [584-08-7] M 138.2, m 898°, d 2.3. Crystd from water between 100° and 0°. After 2 recrystns tech grade had B, Li and Fe at 1.0, 0.04 and 0.01 ppm resp.

Potassium chlorate [3811-04-9] M 122.6, m 368°. Crystd from water (1.8mL/g) between 100° and 0°, and the crystals are filtered onto sintered glass.

Potassium chloride [7447-40-7] M 74.6, m 771°, d 1.98. Dissolved in conductivity water, filtered, and saturated with chlorine (generated from conc HCl and KMnO₄). Excess chlorine was boiled off, and the KCl was pptd by HCl (generated by dropping conc HCl into conc H₂SO₄). The ppte was washed with water, dissolved in conductivity water at 90-95°, and crystd by cooling to about -5°. The crystals were drained at the centrifuge, dried in a vacuum desiccator at room temperature, then fused in a platinum dish under N₂, cooled and stored in desiccator. Potassium chloride has also been sublimed in a stream of prepurified N₂ gas and collected by electrostatic discharge [Craig and McIntosh *Can J Chem* 30 448 1952].

Potassium chromate [7789-00-6] M 194.2, m 975°, d 2.72, pK₁²⁵ 0.74, pK₂²⁵ 6.49 (for H₂CrO₄). Crystd from conductivity water (0.6g/mL at 20°), and dried between 135° and 170°.

Potassium cobalticyanide [13963-58-1] M 332.4, m dec on heating, d 1.91. Crystd from water to remove traces of HCN.

Potassium cyanate [590-28-3] M 81.1, d 2.05, pK²⁵ 3.46 (for HCNO). Common impurities include ammonia and bicarbonate ion (from hydrolysis). Purified by preparing a saturated aqueous solution at

50°, neutralising with acetic acid, filtering, adding two volumes of EtOH and keeping for 3-4h in an ice bath. (More EtOH can lead to co-precipitation of KHCO_3 .) Filtered, washed with EtOH and dried rapidly in a vacuum desiccator (P_2O_5). The process is repeated [Vanderzee and Meyers *J Chem Soc* **65** 153 1961].

Potassium cyanide [151-50-8] **M 65.1, m 634°, d 1.52**. A saturated solution in H_2O -ethanol (1:3) at 60° was filtered and cooled to room temperature. Absolute EtOH was added, with stirring, until crystallisation ceased. The solution was again allowed to cool to room temperature (during 2-3h) then the crystals were filtered off, washed with absolute EtOH, and dried, first at 70-80° for 2-3h, then at 105° for 2h [Brown, Adisesh and Taylor *J Phys Chem* **66** 2426 1962]. Also purified by vacuum melting and zone refining. **HIGHLY POISONOUS**.

Potassium dichromate [7778-50-9] **M 294.2, m 398°(dec), d 2.68**. Crystd from water (1mL/g) between 100° and 0° and dried under vacuum at 156°. (Possible **CARCINOGEN**).

Potassium dihydrogen citrate [866-83-1] **M 230.2**. Crystd from water. Dried at 80°, or in a vacuum desiccator over Sicapent.

Potassium dihydrogen phosphate [7778-77-0] **M 136.1**. Dissolved in boiling distilled water (2mL/g), kept on a boiling water-bath for several hours, then filtered through paper pulp to remove any turbidity. Cooled rapidly with constant stirring, and the crystals were separated on to hardened filter paper, using suction, washed twice with ice-cold water, once with 50% EtOH, and dried at 105°. Alternative crystns are from water, then 50% EtOH, and again water, or from conc aqueous solution by addition of EtOH. Freed from traces of Cu by extracting its aqueous solution with diphenylthiocarbazon in CCl_4 , followed by repeated extraction with CCl_4 to remove traces of diphenylthiocarbazon.

Potassium dithionate [13455-20-4] **M 238.3, pK_{Est(1)} -3.4, pK₂^{2.5} 0.49 (for dithionic acid)**. Crystd from water (1.5mL/g) between 100° and 0°.

Potassium ethylxanthate [140-89-6] **M 160.3, m > 215°(dec)**. Crystd from absolute EtOH, ligroin-ethanol or acetone by addition of Et_2O . Washed with ether, then dried in a desiccator.

Potassium ferricyanide [13746-66-2] **M 329.3, pK²⁵ <1 (for ferricyanide)**. Crystd repeatedly from hot water (1.3mL/g). Dried under vacuum in a desiccator.

Potassium ferrocyanide (3H₂O) [14459-95-1] **M 422.4, pK₃^{2.5} 2.57, pK₄^{2.5} 4.35 (for ferrocyanide)** Crystd repeatedly from distilled water, never heating above 60°. Prepared anhydrous by drying at 110° over P_2O_5 in a vacuum desiccator. To obtain the trihydrate, it is necessary to equilibrate in a desiccator over saturated aqueous soln of sucrose and NaCl. Can also be ppted from a saturated solution at 0° by adding an equal volume of cold 95% EtOH, standing for several hours, then centrifuging and washing with cold 95% EtOH. Finally sucked air dry with a water-pump. The anhydrous salt can be obtained by drying in a platinum boat at 90° in a slow stream of N_2 [Loftfield and Swift *J Am Chem Soc* **60** 3083 1938].

Potassium fluoroborate [14075-53-7] **M 125.9, m 530°**. See potassium tetrafluoroborate on p. 458.

Potassium fluorosilicate [16871-90-2] **M 220.3, d 2.3, pK 1.92 (for H₂SiF₆)**. Crystd several times from conductivity water (100mL/g) between 100° and 0°.

Potassium hexachloroiridate (IV) [16920-56-2] **M 483.1**. Crystd from hot aqueous solution containing a few drops of HNO_3 .

Potassium hexachloroosmate (IV) [16871-60-6] **M 481.1**. Crystd from hot dilute aqueous HCl.

Potassium hexachloroplatinate (IV) [16921-30-5] **M 486.0, m 250°(dec)**. Crystd from water (20mL/g) between 100° and 0°.

Potassium hexacyanochromate (III) (3H₂O) [13601-11-1] M 418.5. Crystd from water.

Potassium hexafluorophosphate [17084-13-8] M 184.1, pK₁^{2.5} ~ 0.5, pK₂⁵ 5.12 (for fluorophosphoric acid H₂PO₃F). Crystd from alkaline aqueous solution, using polyethylene vessels, or from 95% EtOH, and dried in a vacuum desiccator over KOH.

Potassium hexafluorozirconate (K₂ZrF₆) [16923-95-8] M 283.4, d 3.48. Recrystd from hot water (solubility is 0.78% at 2° and 25% at 100°).

Potassium hydrogen fluoride [7789-29-9] M 78.1. Crystd from water.

Potassium hydrogen D-glucarate [18404-47-2] M 248.2, m 188°(dec). Crystd from water.

Potassium hydrogen malate [4675-64-3] M 172.2. A saturated aqueous solution at 60° was decolorised with activated charcoal, and filtered. The filtrate was cooled in water-ice bath and the salt was ppted by addition of EtOH. After being crystallised five times from ethanol-water mixtures, it was dried overnight at 130° in air [Eden and Bates *J Res Nat Bur Stand* 62 161 1959].

Potassium hydrogen oxalate (H₂O) [127-95-7] M 137.1. Crystd from water by dissolving 20g in 100mL water at 60° containing 4g of potassium oxalate, filtering and allowing to cool to 25°. The crystals, after washing three or four times with water, are allowed to dry in air.

Potassium hydrogen phthalate [877-24-7] M 204.2. Crystd first from a dilute aqueous solution of K₂CO₃, then H₂O(3mL/g) between 100° and 0°. Before being used as a standard in volumetric analysis, analytical grade potassium hydrogen phthalate should be dried at 120° for 2h, then allowed to cool in a desiccator.

Potassium hydrogen d-tartrate [868-14-4] M 188.2, [α]₅₄₆²⁰ +37.5° (c 10, M NaOH). Crystd from water (17mL/g) between 100° and 0°. Dried at 110°.

Potassium hydroxide (solution) [1310-58-3] M 56.1, pK²⁵ 16 (for aquo K⁺). Its carbonate content can be reduced by rinsing KOH sticks rapidly with water prior to dissolving them in boiled out distilled water. Alternatively, a slight excess of saturated BaCl₂ or Ba(OH)₂ can be added to the soln which, after shaking well, is left so that the BaCO₃ ppte will separate out. Davies and Nancollas [Nature 165 237 1950] rendered KOH solutions carbonate free by ion exchange using a column of Amberlite IR-100 in the OH⁻ form.

Potassium iodate [7758-05-6] M 214.0, pK²⁵ 0.80 (for HIO₃). Crystd twice from distilled water (3mL/g) between 100° and 0°, dried for 2h at 140° and cooled in a desiccator. Analytical reagent grade material dried in this way is suitable for use as an analytical standard.

Potassium iodide [7681-11-0] M 166.0, pK²⁵ -8.56 (for HI). Crystd from distilled water (0.5mL/g) by filtering the near-boiling soln and cooling. To minimise oxidation to iodine, the crystn can be carried out under N₂ and the salt is dried under vacuum over P₂O₅ at 70-100°. Before drying, the crystals can be washed with EtOH or with acetone followed by pet ether. Has also been recrystallised from water/ethanol. After 2 recrystns ACS/USP grade had Li and Sb at <0.02 and <0.01 ppm resp.

Potassium ionophore I (valinomycin) [2001-95-8] M 111.3, m 186-187°, 190°, [α]_D²⁰ +31.0° (c 1.6, *C₆H₆). See valinomycin on p. 573 in Chapter 6.

Potassium isoamyl xanthate [61792-26-5] M 202.4, pK 1.82 (pK⁰ 2.8 free acid). Crystd twice from acetone-diethyl ether. Dried in a desiccator for two days and stored under refrigeration.

Potassium laurate [10124-65-9] M 338.4. Recrystd three times from EtOH [Neto and Helene *J Phys Chem* 91 1466 1987].

Potassium nickel sulfate (6H₂O) [13842-46-1] **M 437.1**. Crystd from H₂O (1.7mL/g) between 75° and 0°.

Potassium nitrate [7757-79-1] **M 101.1, m 334°**. Crystd from hot H₂O (0.5mL/g) by cooling (*cf* KNO₂ below). Dried for 12h under vacuum at 70°. After 2 recrystns tech grade had < 0.001 ppm of metals.

Potassium nitrite [7758-09-0] **M 85.1, m 350°(dec), pK²⁰ 3.20 (for HNO₂)**. A saturated solution at 0° can be warmed and partially evaporated under vacuum, the crystals so obtained being filtered from the warm solution. (This procedure is designed to reduce the level of nitrate impurity and is based on the effects of temperature on solubility. The solubility of KNO₃ in water is 13g/100mL at 0°, 247g/100mL at 100°; for KNO₂ the corresponding figures are 280g/100mL and 413g/100mL.)

Potassium nitrosodisulfonate (Fremy's Salt) [14293-70-0] **M 268.3**. Yellow needles (dimeric) which dissolve in H₂O to give the violet monomeric free radical. Purified by dissolving (~12g) in 2M KOH (600mL) at 45°, filtering the blue soln and keeping it in a refrigerator overnight. The golden yellow crystals (10g) are filt'd off, washed with MeOH (3x), then Et₂O and stored in a glass container in a vac over KOH. It is stable indefinitely when dry. [Cram and Reeves *J Org Chem* **80** 3094 1958; Schenk *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I p. 505 1963.]

Potassium nonafluorobutane sulfonate [29420-49-3] **M 338.2**. Wash with H₂O and dry in vacuum. The K salt when distilled with 100% H₂SO₄ gives the free acid which can be distilled (**b** 105°/22mm, 210-212°/760mm) and then converted to the K salt. [*J Chem Soc* 2640 1957.]

Potassium oleate [143-18-0] **M 320.6**. Crystd from EtOH (1mL/g).

Potassium osmate (VI) dihydrate [19718-36-6] **M 368.4**. *Hygroscopic POISONOUS* crystals which are soluble in H₂O but insol in EtOH and Et₂O. It decomposes slowly in H₂O to form the tetroxide which attacks the eyes. The solid should be kept dry and in this form it is relatively safe. [*Synthesis* 610 1972.]

Potassium oxalate [6487-48-5] **M 184.2, m 160°(dec), d 2.13**. Crystd from hot water.

Potassium perchlorate [7778-74-7] **M 138.6, m 400(dec), d 2.52, pK²⁵ -2.4 to -3.1 (for HClO₄)**. Crystd from boiling water (5mL/g) by cooling. Dried under vacuum at 105°.

Potassium periodate (potassium metaperiodate) [7790-21-8] **M 230.0, m 582°, d 3.62**. Crystd from distilled water.

Potassium permanganate [7722-64-7] **M 158.0, m 240(dec), d 2.7, pK²⁵ -2.25 (for HMnO₄)**. Crystd from hot water (4mL/g at 65°), then dried in a vacuum desiccator over CaSO₄. Phillips and Taylor [*J Chem Soc* 4242 1962] cooled an aqueous solution of KMnO₄, saturated at 60°, to room temperature in the dark, and filtered through a No.4 porosity sintered-glass filter funnel. The solution was allowed to evaporate in air in the dark for 12h, and the supernatant liquid was decanted from the crystals, which were dried as quickly as possible with filter paper.

Potassium peroxydisulfate (potassium persulfate) [7727-21-1] **M 270.3**. Crystd twice from distilled water (10mL/g) and dried at 50° in a vacuum desiccator.

Potassium peroxymonosulfate (Oxone, potassium monopersulfate triple salt; 2KHSO₅·KHSO₄·K₂SO₄), [37222-66-5, triple salt] [70693-62-8] **M 614.8**. This is a stable form of Caro's acid and should contain >4.7% of active oxygen. It can be used in EtOH/H₂O and EtOH/AcOH/H₂O solutions. If active oxygen is too low it is best prepared afresh from 1mole of KHSO₅, 0.5moles of KHSO₄ and 0.5moles of K₂SO₄. [Kennedy and Stock *J Org Chem* **25** 1901 1960; Stephenson US Patent 2,802,722 1957.] A rapid prepn of *Caro's acid* is made by stirring finely powdered potassium persulfate (M 270.3) into ice cold conc H₂SO₄ (7mL) and when homogeneous add ice (40-50g). It is stable for several days if kept cold.

Keep away from organic matter as it is a **STRONG OXIDANT**. A detailed prepn of **Caro's acid** (*hypersulfuric acid*, H_2SO_5 , [7722-86-3]) in crystalline form $m \sim 45^\circ$ from H_2O_2 and chlorosulfonic acid was described by Fehér in *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I p. 388 1963.

Potassium perrhenate (KReO_4) [10466-65-6] **M 289.3**, $\text{pK}^{25} -1.25$ (for HReO_4). Crystd from water (7mL/g), then fused in a platinum crucible in air at 750° .

Potassium phenol-4-sulfonate (**4-hydroxybenzene-1-sulfonic acid K salt**) [30145-40-5] **M 212.3**. Crystd several times from distilled water at 90° , after treatment with charcoal, by cooling to *ca* 10° . Dried at 90 - 100° .

Potassium phthalimide (**phthalimide K salt**) [1074-82-4] **M 185.2**, $m >300^\circ$. The solid may contain phthalimide and K_2CO_3 from hydrolysis. If too much hydrolysis has occurred (this can be checked by extraction with cold Me_2CO in which the salt is insoluble, evaporation of the Me_2CO and weighing the residue) it would be better to prepare it afresh. If little hydrolysis had occurred then recryst from a large volume of EtOH, and wash solid with a little Me_2CO and dry in a continuous vacuum to constant weight. [Salzerg and Supriawski *Org Synth Coll Vol I* 119 1941; Raman and IR: Hase *J Mol Struct* **48** 33 1978; Dykman *Chem Ind (London)* 40 1972; IR, NMR: Assef et al. *Bull Soc Chim Fr* II-167 1979.]

Potassium picrate [573-83-1] **M 267.2**. Crystd from water or 95% EtOH, and dried at room temperature in vacuum. It is soluble in 200 parts of cold water and 4 parts of boiling water. **THE DRY SOLID EXPLODES WHEN STRUCK OR HEATED.**

Potassium propionate [327-62-8] **M 112.2**. Crystd from water (30mL/g) or 95% EtOH.

Potassium reineckate [34430-73-4] **M 357.5**. Crystd from KNO_3 soln, then from warm water [Adamson *J Am Chem Soc* **80** 3183 1958].

Potassium (VI) ruthenate [31111-21-4] **M 243.3**. Dissolve in H_2O and evaporate until crystals are formed. The crystals are iridescent green prisms which appear red as thin films. Possible impurity is RuO_4 ; in this case wash with CCl_4 (which dissolves RuO_4). The concn of an aqueous solution of RuO_4^{2-} (orange colour) can be estimated from the absorbance at 385nm (ϵ 1030 $\text{M}^{-1} \text{cm}^{-1}$), or at 460nm (ϵ 1820 $\text{M}^{-1} \text{cm}^{-1}$). [Can J Chem **50** 3741 1972; *J Am Chem Soc* **74** 5012 1952; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1600 1965].

Potassium selenocyanate [3425-46-5] **M 144.1**. Dissolved in acetone, filtered and ppted by adding Et_2O .

Potassium sodium tartrate ($4\text{H}_2\text{O}$) [6381-59-5 ($4\text{H}_2\text{O}$); 304-59-6 (*R,R*)] **M 282.3**. Crystd from distilled water (1.5mL/g) by cooling to 0° .

Potassium sulfate [7778-80-5] **M 174.3**, m 1069 $^\circ$, d 2.67 Crystd from distilled water (4mL/g at 20° ; 8mL/g at 100°) between 100° and 0° .

Potassium *d*-tartrate (H_2O) [921-53-9 , 6381-59-5] **M 235.3**, m loses H_2O at 150° , d 1.98. Crystd from distilled water (solubility: 0.4mL/g at 100° ; 0.7mL/g at 14°).

Potassium tetrachloroplatinate(II) [10025-99-7] **M 415.1**, m 500 $^\circ$ (dec). Crystd from aqueous 0.75M HCl (20mL/g) between 100° and 0° . Washed with ice-cold water and dried.

Potassium tetracyanopalladate (II) $3\text{H}_2\text{O}$ [10025-98-6] **M 377.4**. All operations should be carried out in an efficient fume cupboard - **Cyanide is very POISONOUS** Dissolve the complex (*ca* 5g) in a solution of KCN (4g) in H_2O (75mL) with warming and stirring and evaporate hot till crystals appear. Cool, filter off the crystals and wash with a few drops of cold H_2O . Further concentration of the mother liquors provides more crystals. The complex is recrystallised from H_2O as the colourless *trihydrate*. It effloresces in

dry air and dehydrates at 100° to the *monohydrate*. The *anhydrous salt* is obtained by heating at 200°, but at higher temperatures it decomposes to $(\text{CN})_2$, Pd and KCN. [*Inorg Synth* 2 245 1946.]

Potassium tetrafluoroborate (potassium borofluoride) [14075-53-7] M 125.9, m 530°, d_4^{30} 2.505, pK^{25} -4.9 (for HBF_4). Cryst from H_2O (sol % (temp): 0.3 (3°), 0.45 (20°), 1.4 (40°), 6.27 (100°), and dry under vacuum. Non-hygroscopic salt. A 10% solution is transparent blue at 100°, green at 90° and yellow at 60°. [*Chem Ber* 65 535 1932; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol 1 223 1963.]

Potassium tetraoxalate ($2\text{H}_2\text{O}$) [oxalic acid hemipotassium salt] [6100-20-5 ($2\text{H}_2\text{O}$); 127-96-8 (*anhydr*)] M 254.2. Crystd from water below 50°. Dried below 60° at atmospheric pressure.

Potassium tetraphenylborate [3244-41-5] M 358.3. Ppted from a soln of KCl acidified with dilute HCl, then crystallised twice from acetone, washed thoroughly with water and dried at 110° [Findeis and de Vries *Anal Chem* 28 1899 1956]. It has also been recrystd several times from conductivity water.

Potassium thiocyanate [333-20-0] M 97.2, m 172°, pK^{25} -1.85 (for HSCN). Crystd from H_2O if much chloride ion is present in the salt, otherwise from EtOH or MeOH (optionally by addition of Et_2O). Filtered on a Büchner funnel without paper, and dried in a desiccator at room temperature before being heated for 1h at 150°, with a final 10-20min at 200° to remove the last traces of solvent [Kolthoff and Lingane *J Am Chem Soc* 57 126 1935]. Stored in the dark.

Potassium thiosulfate hydrate [13446-67-8; 10294-66-3 (75% aq soln)] M 190.3, pK_1^{25} 0.6, pK_2^{25} 1.74 (for $\text{H}_2\text{S}_2\text{O}_3$) Crystd from warm water (0.5mL/g) by cooling in an ice-salt mixture.

Potassium thiotosylate [28519-50-8] M 226.4. Recrystallise from absolute EtOH and dry at 130°. In wet EtOH the *monohydrate* can be obtained. [*J Gen Chem USSR (Engl Transl)* 28 1345 1958.]

Potassium trifluoroacetate [2923-16-2] M 152.1, m 140-142°, pK^{25} 0.52 (for $\text{CF}_3\text{CO}_2\text{H}$). To purify dissolve the salt in trifluoroacetic acid with ca 2% of trifluoroacetic anhydride, filter and evaporate carefully to dryness (avoid over heating), and finally dry in a vacuum at 100°. It can be recrystallised from trifluoroacetic acid (solubility in the acid is ca 50.1%). [*J Am Chem Soc* 74 4746 1952, 76 4285 1954; *J Inorg Nucl Chem* 9 166 1959.]

Potassium trimethylsilanolate (trimethylsilanol K salt) [10519-96-7] M 128.3, m 131-135° (cubic form), d^{25} 1.11, 125°dec (orthorhombic form). Recryst from H_2O and dried at 100°/1-2mm. [*J Am Chem Soc* 75 5615 1953; IR: *J Org Chem* 17 1555 1952.]

Potassium tungstate (ortho $2\text{H}_2\text{O}$) [37349-36-3; 7790-60-5] M 362.1, m 921°, d 3.12, pK_1^{25} 2.20, pK_2^{25} 3.70 (for H_2WO_4). Crystd from hot water (0.7mL/g).

Praseodymium acetate [6192-12-7] M 318.1. Recrystd several times from water [Ganapathy *J Am Chem Soc* 109 3159 1986].

Praseodymium trichloride ($6\text{H}_2\text{O}$) [10361-79-2] M 355.4, pK_1^{25} 8.55 (for Pr^{3+} hydrol). Its 1M soln in 6M HCl was passed twice through a Dowex-1 anion-exchange column. The eluate was evaporated in a vac desiccator to about half its vol and allowed to crystallise [Katzin and Gulyas *J Phys Chem* 66 494 1962].

Praseodymium oxide (Pr_6O_{11}) [12037-29-5] M 1021.4. Dissolved in acid, ppted as the oxalate and ignited at 650°.

Propargyl triphenyl phosphonium bromide [2091-46-5] M 381.4, m 179°. Recrystallises from 2-propanol as white plates. Also crystallises from EtOH, m 156-158°. IR has ν 1440, 1110 cm^{-1} (P-C str). [*Justus Liebigs Ann Chem* 682 62 1965; *J Org Chem* 42 200 1977].

Propenyloxy trimethylsilane [1833-53-0] M 130.3, b 93-95°/atm, d 0.786. Purified by fractional distillation using a very efficient column at atmospheric pressure. Usually contains 5% of hexamethyldisiloxalane which boils at 99-101°, but is generally non-reactive and need not be removed. [*J Am Chem Soc* 71 5091 1952.] It has been distilled under N₂ through a 15cm column filled with glass helices. Fraction b 99-104° is further purified by gas chromatography through a Carbowax column (Autoprep A 700) at a column temperature of 87°, retention time is 9.5min. [*J Organometal Chem* 1 476 1963-4.]

1-Propenyltrimethylsilane (cis and trans mixture) [17680-01-2] M 114.3, b 85-88°, n_D²⁰ 1.4121. Dissolve (~ 20g) in THF (200mL), shake with H₂O (2x 300 mL), dry (Na₂SO₄) and fractionate. This is a mixture of *cis* and *trans* isomers which can be separated by gas chromatography on an AgNO₃ column (for prep: see Seyferth and Vaughan *J Organomet Chem* 1 138 1963) at 25° with He as carrier gas at 9psi. The *cis*-isomer has n_D²⁵ 1.4105 and the *trans*-isomer has n_D²⁵ 1.4062. [Seyferth et al. *Pure Appl Chem* 13 159 1966.]

Pyridinium chlorochromate [26299-14-9] M 215.6, m 205°(dec). Dry in a vacuum for 1h. It is not hygroscopic and can be stored for extended periods at room temp without change. If very suspect it can be readily prepared. [*Tetrahedron Lett* 2647 1975; *Synthesis* 245 1982.]

§ Available commercially on a polymer support.

Pyridinium dichromate [20039-37-6] M 376.2, m 145-148°, 152-153°. Dissolve in the minimum volume of H₂O and add 5 volumes of cold Me₂CO and cool to -20°. After 3h the orange crystals are collected, washed with a little cold Me₂CO and dried in a vacuum. It is soluble in dimethylformamide (0.9g/mL at 25°), and in H₂O, and has a characteristic IR with ν 930, 875, 765 and 730cm⁻¹. [*Tetrahedron Lett* 399 1979; *Chem Ind (London)* 1594 1969.] (Possible carcinogen).

§ Available commercially on a polymer support.

3-(2-Pyridyl)-5,6-diphenyl-1,2,4-triazine-*p,p'*-disulfonic acid, monosodium salt (H₂O) [63451-29-6] M 510.5. Purified by recrystn from water or by dissolving in the minimum volume of water, followed by addition of EtOH to ppt the pure salt.

Pyrocatechol Violet (tetraphenolictriphenylmethanesulfonic acid Na salt) [115-41-3] M 386.4, ϵ 1.4 x 10⁴ at 445nm in acetate buffer pH 5.2-5.4, pK_{Est(1)}>0 (SO₃H), pK_{Est(2)}~ 9.4, pK_{Est(3)}~ 13. It was recrystd from glacial acetic acid. Very *hygroscopic*. Indicator standard for metal complex titrations. [Mustafin et al. *Zh Anal Khim* 22 1808 1967.]

Pyrogallol Red (tetraphenolic xanthyliumphenylsulfonate) [32638-88-3] M 418.4, m >300°(dec), ϵ 4.3 x 10⁴ at 542nm, pH 7.9-8.6, pK as above. Recrystd from aqueous alkaline solution (Na₂CO₃ or NaOH) by precipitation on acidification [Suk *Collect Czech Chem Commun* 31 3127 1966].

Pyronin B [di-(3,6-bis(diethylamino)xanthylium chloride) diFeCl₃ complex] [2150-48-3] M 358.9 (Fe free), m 176-178° (diFe complex), CI 45010, λ_{max} 555nm, pK²⁵ 7.7. Crystd from EtOH. Forms Fe stain.

Quinolinium chlorochromate [108703-35-1] M 265.6, m 127-130°. A yellow-brown solid which is stable in air for long periods. If it has deteriorated or been kept for too long, it is best to prepare it freshly. Add freshly distilled quinoline (13mL) to a mixture of chromic acid (CrO₃) (10g) and ~ 5M HCl (11mL of conc HCl and 10mL of H₂O) at 0°. A yellow-brown solid separates, it is filtered off on a sintered glass funnel, dried for 1h in a vacuum, and can be stored for extended periods without serious loss in activity. It is a good oxidant for primary alcohol in CH₂Cl₂. [Singh et al. *Chem Ind (London)* 751 1986; method of Corey and Suggs *Tetrahedron Lett* 2647 1975.]

Reinecke salt see ammonium reineckate on p. 394.

Resorufin (7-hydroxy-3H-phenoxazine-3-one Na salt) [635-78-9] **M 213.2**, **pK₁³⁰ 6.93**, **pK₂³⁰ 9.26**, **pK₃³⁰ 10.0**. Washed with water and recrystd from EtOH several times.

Rhodium (II) acetate dimer (2H₂O) [15956-28-2] **M 478.0**. Dissolve 5g in boiling MeOH (ca 600mL) and filter. Concentrate to 400mL and chill overnight at ca 0° to give dark green crystals of the MeOH adduct. Conc'n of the mother liquors gives a further crop of [Rh(OAc)₂]₂·2MeOH. The adduct is then heated at 45° in vacuum for 2h (all MeOH is lost) to leave the emerald green crystals of the acetate. [*J Chem Soc (A)* 3322 1970.] Alternatively dissolve in glacial AcOH and reflux for a few hs to give an emerald green soln. Evaporate most of the AcOH on a steam bath then heat the residue at 120°/1h. Extract the residue with boiling Me₂CO. Filter, concentrate to half its volume and keep at 0°/18h. Collect the crystals, wash with ice cold Me₂CO and dry at 110°. It is soluble in most organic solvents with which it forms adducts including NMe₃ and Me₂S and give solutions with different colours varying from green to orange and red. [UV: *Inorg Synth* 2 960 1963.]

Rhodium (III) chloride [10049-07-7] **M 209.3**, **m >100°(dec)**, **b 717°**. Probable impurities are KCl and HCl. Wash solid well with small volumes of H₂O to remove excess KCl and KOH and dissolve in the minimum volume of conc HCl. Evaporate to dryness on a steam bath to give wine-red coloured RhCl₃·3H₂O. Leave on the steam bath until odour of HCl is lost - do not try to dry further as it begins to decompose above 100° to the oxide and HCl. It is not soluble in H₂O but soluble in alkalis or CN solns and forms double salts with alkali chlorides. [*Inorg Synth* 7 214 1063.]

Rhodizonic acid sodium salt (5,6-dihydroxycyclohex-5-ene-1,2,3,4-tetraone di-Na salt) [523-21-7] **M 214.0**, **pK₁³⁰ 4.1 (4.25)**, **pK₂³⁰ 4.5 (4.72)**. The free acid, obtained by acidifying and extracting with Et₂O, drying (MgSO₄), filtering, evaporating and distilling in vacuum (**b** 155-160°/14mm). The *free acid* solidifies on cooling and the colourless crystals can be recrystd from tetrahydrofuran-pet ether or *C₆H₆. It forms a *dihydrate* **m** 130-140°. The pure di Na salt is formed by dissolving in 2 equivs of NaOH and evaporating in a vacuum. It forms violet crystals which give an orange soln in H₂O that is unstable for extended periods even at 0°, and should be prepared freshly before use. Salts of rhodizonic acid cannot be purified by recrystn without great loss due to conversion to croconate, so that the original material must be prepared pure. It can be washed with NaOAc soln then EtOH to remove excess NaOAc dried under vacuum and stored in the dark. [UV and tautomerism: Schwarzenbach and Suter *Helv Chim Acta* 24 617 1941; Polarography: Preisler and Berger *J Am Chem Soc* 64 67 1942; Souchay and Taibouet *J Chim Phys* 49 C108 1952.]

Rose Bengal [Acid Red 94, 4,5,6,7-tetrachloro-2'.4',5',7'-tetraiodofluorescein di-Na or di-K salt] [*di-Na salt* 632-69-9] **M 1017.7 (di-Na salt)** [*di-K salt* 11121-48-5] **M 1049.8 (di-K salt)**. This biological stain can be purified by chromatography on silica TLC using a 35:65 mix of EtOH/acetone as eluent.

Rubidium bromide [7789-39-1] **M 165.4**, **m 682°**, **b 1340°**, **d 3.35**. A white crystalline powder which crystallises from H₂O (solubility: 50% in cold and 67% in boiling H₂O to give a neutral soln). Also crystd from near-boiling water (0.5mL/g) by cooling to 0°.

Rubidium chlorate [13446-71-4] **M 168.9**, **d 3.19**. Crystd from water (1.6mL/g) by cooling from 100°.

Rubidium chloride [7791-11-9] **M 120.9**, **m 715°**, **d 2.80**. Crystd from water (0.7mL/g) by cooling to 0° from 100°.

Rubidium nitrate [13126-12-0] **M 147.5**, **m 305°**, **d 3.11**. Crystd from hot water (0.25mL/g) by cooling to room temperature.

Rubidium perchlorate [13510-42-4] M 184.9, d 2.80, pK_1^{25} -2.4 to -3.1 (for $HClO_4$). Crystd from hot water (1.6mL/g) by cooling to 0°.

Rubidium sulfate [7488-54-2] M 267.0, m 1050°, d 6.31. Crystd from water (1.2mL/g) between 100° and 0°.

Ruthenium (III) acetylacetonate [14284-93-6] M 398.4, m 240°(dec). Purified by recrystn from *benzene. [*J Am Chem Soc* 74 6146 1952.]

Ruthenium (III) chloride (2H₂O) (β-form) [14898-67-0] M 207.4 + H₂O, m >500°(dec), d 3.11, pK_1^{25} 3.40 (for aquo Rh^{3+} hydrolysis). Dissolve in H₂O, filter and concentrate to crystallise in the absence of air to avoid oxidation. Evaporate the solution in a stream of HCl gas while being heated just below its boiling point until a syrup is formed and finally to dryness at 80-100° and dried in a vacuum over H₂SO₄. When heated at 700° in the presence of Cl₂ the insoluble α-form is obtained [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1598 1965; *J Org Chem* 46 3936 1981].

Ruthenium (IV) oxide [12036-10-1] M 133.1, d 6.97. Freed from nitrates by boiling in distilled water and filtering. A more complete purification is based on fusion in a KOH-KNO₃ mix to form the soluble ruthenate and perruthenate salts. The melt is dissolved in water, and filtered, then acetone is added to reduce the ruthenates to the insoluble hydrate oxide which, after making a slurry with paper pulp, is filtered and ignited in air to form the anhydrous oxide [*Campbell, Ortner and Anderson Anal Chem* 33 58 1961].

Ruthenocene [bis-(cyclopentadienyl)ruthenium] [1287-13-4] M 231.2, m 195.5°, 199-210°. Sublime in high vacuum at 120°. Yellow crystals which can be recrystallised from CCl₄ as transparent plates. [*J Am Chem Soc* 74 6146 1952].

Samarium (II) iodide [32248-43-4] M 404.2, m 520°, b 1580. Possible impurity is SmI₃ from which it is made. If present, grind solid to a powder and heat in a stream of pure H₂. The temperature (~ 500-600°) should be below the m (~ 628°) of SmI₃, since the molten compounds react very slowly. [*Wetzel in Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II pp. 1149, 1150 1965.]

Selenious acid [7783-00-8] M 129.0, m 70°(dec), d 3.0, pK_1^{25} 2.62, pK_2^{25} 8.32 (H₂SeO₃). Crystd from water. On heating it loses water and SeO₂ sublimes.

Selenium [7782-49-2] M 79.0, m 217.4°, d 4.81. Dissolved in small portions in hot conc HNO₃ (2mL/g) filtered and evaporated to dryness to give selenious acid which was then dissolved in conc HCl. Passage of SO₂ into the solution pptd selenium (but not tellurium) which was filtered off and washed with conc HCl. This purification process was repeated. The selenium was then converted twice to the selenocyanate by treating with a 10% excess of 3M aqueous KCN, heating for half an hour on a sand-bath and filtering. Addition of an equal weight of crushed ice to the cold solution, followed by an excess of cold, conc HCl, with stirring (in a well ventilated fume hood because HCN is evolved) pptd selenium powder, which, after washing with water until colourless, and then with MeOH, was heated in an oven at 105°, then by fusion for 2h under vacuum. It was cooled, crushed and stored in a desiccator [*Tideswell and McCullough J Am Chem Soc* 78 3036 1956].

Selenium dioxide [7446-08-4] M 111.0, m 340°. Purified by sublimation, or by solution in HNO₃, pptn of selenium which, after standing for several hours or boiling, is filtered off, then re-oxidised by HNO₃ and cautiously evaporated to dryness below 200°. The dioxide is dissolved in water and again evaporated to dryness.

Selenopyronine [85051-91-8] M 365.8, λ_{max} 571nm (ϵ 81,000). Purified as the hydrochloride from hydrochloric acid [*Fanghanel et al. J Phys Chem* 91 3700 1987].

[Previous Page](#)

Rubidium perchlorate [13510-42-4] M 184.9, d 2.80, pK_1^{25} -2.4 to -3.1 (for $HClO_4$). Crystd from hot water (1.6mL/g) by cooling to 0°.

Rubidium sulfate [7488-54-2] M 267.0, m 1050°, d 6.31. Crystd from water (1.2mL/g) between 100° and 0°.

Ruthenium (III) acetylacetonate [14284-93-6] M 398.4, m 240°(dec). Purified by recrystn from *benzene. [*J Am Chem Soc* 74 6146 1952.]

Ruthenium (III) chloride (2H₂O) (β-form) [14898-67-0] M 207.4 + H₂O, m >500°(dec), d 3.11, pK_1^{25} 3.40 (for aquo Rh^{3+} hydrolysis). Dissolve in H₂O, filter and concentrate to crystallise in the absence of air to avoid oxidation. Evaporate the solution in a stream of HCl gas while being heated just below its boiling point until a syrup is formed and finally to dryness at 80-100° and dried in a vacuum over H₂SO₄. When heated at 700° in the presence of Cl₂ the insoluble α-form is obtained [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1598 1965; *J Org Chem* 46 3936 1981].

Ruthenium (IV) oxide [12036-10-1] M 133.1, d 6.97. Freed from nitrates by boiling in distilled water and filtering. A more complete purification is based on fusion in a KOH-KNO₃ mix to form the soluble ruthenate and perruthenate salts. The melt is dissolved in water, and filtered, then acetone is added to reduce the ruthenates to the insoluble hydrate oxide which, after making a slurry with paper pulp, is filtered and ignited in air to form the anhydrous oxide [Campbell, Ortnier and Anderson *Anal Chem* 33 58 1961].

Ruthenocene [bis-(cyclopentadienyl)ruthenium] [1287-13-4] M 231.2, m 195.5°, 199-210°. Sublime in high vacuum at 120°. Yellow crystals which can be recrystallised from CCl₄ as transparent plates. [*J Am Chem Soc* 74 6146 1952].

Samarium (II) iodide [32248-43-4] M 404.2, m 520°, b 1580. Possible impurity is SmI₃ from which it is made. If present, grind solid to a powder and heat in a stream of pure H₂. The temperature (~ 500-600°) should be below the m (~ 628°) of SmI₃, since the molten compounds react very slowly. [Wetzel in *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II pp. 1149, 1150 1965.]

Selenious acid [7783-00-8] M 129.0, m 70°(dec), d 3.0, pK_1^{25} 2.62, pK_2^{25} 8.32 (H₂SeO₃). Crystd from water. On heating it loses water and SeO₂ sublimes.

Selenium [7782-49-2] M 79.0, m 217.4°, d 4.81. Dissolved in small portions in hot conc HNO₃ (2mL/g) filtered and evaporated to dryness to give selenious acid which was then dissolved in conc HCl. Passage of SO₂ into the solution pptd selenium (but not tellurium) which was filtered off and washed with conc HCl. This purification process was repeated. The selenium was then converted twice to the selenocyanate by treating with a 10% excess of 3M aqueous KCN, heating for half an hour on a sand-bath and filtering. Addition of an equal weight of crushed ice to the cold solution, followed by an excess of cold, conc HCl, with stirring (in a well ventilated fume hood because HCN is evolved) pptd selenium powder, which, after washing with water until colourless, and then with MeOH, was heated in an oven at 105°, then by fusion for 2h under vacuum. It was cooled, crushed and stored in a desiccator [Tideswell and McCullough *J Am Chem Soc* 78 3036 1956].

Selenium dioxide [7446-08-4] M 111.0, m 340°. Purified by sublimation, or by solution in HNO₃, pptn of selenium which, after standing for several hours or boiling, is filtered off, then re-oxidised by HNO₃ and cautiously evaporated to dryness below 200°. The dioxide is dissolved in water and again evaporated to dryness.

Selenopyronine [85051-91-8] M 365.8, λ_{max} 571nm (ϵ 81,000). Purified as the hydrochloride from hydrochloric acid [Fanghanel et al. *J Phys Chem* 91 3700 1987].

Selenourea [630-10-4] M 123.0, m 214-215°(dec). Recrystd from water under nitrogen.

Silica [7631-86-9 (*colloidal*); 112945-52-5 (*fumed*)]. Purification of silica for high technology applications uses isopiestic vapour distillation from conc volatile acids and is absorbed in high purity water. The impurities remain behind. Preliminary cleaning to remove surface contaminants uses dip etching in HF or a mixture of HCl, H₂O₂ and deionised water [Phelan and Powell *Analyst* 109 1299 1984].

Silica gel [63231-67-4; 112926-00-8]. Before use as a drying agent, silica gel is heated in an oven, then cooled in a desiccator. Conditions in the literature range from heating at 110° for 15h to 250° for 2-3h. Silica gel has been purified by washing with hot acid (in one case successively with aqua regia, conc HNO₃, then conc HCl; in another case digested overnight with hot conc H₂SO₄), followed by exhaustive washing with distilled water (one week in a Soxhlet apparatus has also been used), and prolonged oven drying. Alternatively, silica gel has been extracted with acetone until all soluble material was removed, then dried in a current of air, washed with distilled water and oven dried. Silica gel has also been washed successively with water, M HCl, water, and acetone, then activated at 110° for 15h.

Silicon monoxide [10097-28-6] M 44.1, m > 1700°, d 2.18. Purified by sublimation in a porcelain tube in a furnace at 1250° (4h) in a high vacuum (10⁻⁴mm) in a stream of N₂. It is obtained as brownish black scales. [*Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 696 1963.]

Silicon tetraacetate [562-90-3] M 264.3, m 110-111°, b 148°/5-6mm, pK₁²⁵ 9.7, pK₂²⁵ 11.9 (for H₄SiO₄ free acid). It can be crystallised from mixtures of CCl₄ and pet ether or Et₂O, or from acetic anhydride and then dried in a vacuum desiccator over KOH. Ac₂O adheres to the crystals and is removed first by drying at room temp then at 100° for several hours. It is soluble in Me₂CO, is very *hygroscopic* and effervesces with H₂O. It decomposes at 160-170°. [*Z Obshch Khim (Engl Transl)* 27 985 1957; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 701 1963.]

Silicon tetrachloride [10026-04-7] M 169.9, m -70°, b 57.6°, d 1.483. Distd under vacuum and stored in sealed ampoules under N₂. Very sensitive to moisture.

12-Silicotungstic acid (tungstosilicic acid; H₄SiW₁₂O₄₀) [12027-43-9] M 2914.5. Extracted with diethyl ether from a solution acidified with HCl. The diethyl ether was evaporated under vacuum, and the free acid was crystallised twice [Matijevic and Kerker *J Phys Chem* 62 1271 1958].

Silver (metal) [7440-22-4] M 107.9, m 961.9°, b 2212°, d 10.5. For purification by electrolysis, see Craig et al. [*J Res Nat Bur Stand* 64A 381 1960].

Silver acetate [563-63-3] M 166.9, pK²⁵ >11.1 (for aquo Ag⁺ hydrolysis). Shaken with acetic acid for three days, the process being repeated with fresh acid, the solid then being dried in a vacuum oven at 40° for 48h. Has also been recrystallised from water containing a trace of acetic acid, and dried in air.

Silver bromate [7783-89-3] M 235.8, m dec on heating, d 5.21. Crystd from hot water (80mL/g).

Silver bromide [7785-23-1] M 187.8, m 432°, d 6.47. Purified from Fe, Mn, Ni and Zn by zone melting in a quartz vessel under vacuum.

Silver chlorate [7783-92-8] M 191.3, m 230°, b 270°(dec), d 4.43. Recrystd three times from water (10mL/g at 15°; 2mL/g at 80°).

Silver chloride [7783-90-6] M 143.3, m 455°, b 1550°, d 5.56. Recrystd from conc NH₃ solution.

Silver chromate [7784-01-2] M 331.8, d²⁵ 5.625, pK₁²⁵ 0.74, pK₂²⁵ 6.49 (for H₂CrO₄). Wash the red-brown powder with H₂O, dry in a vacuum, then powder well and dry again in a vacuum at 90°/5h. Solubility in H₂O is 0.0014% at 10°. [*J Org Chem* 42 4268 1977.]

Silver cyanide [506-64-9] M 133.9, m dec at 320°, d 3.95. **POISONOUS** white or grayish white powder. Stir thoroughly with H₂O, filter, wash well with EtOH and dry in air in the dark. It is very insoluble in H₂O (0.000023g in 100mL H₂O) but is soluble in HCN or aqueous KCN to form the soluble Ag(CN)₂⁻ complex. [*Chem Ber* 72 299 1939; *J Am Chem Soc* 52 184 1930.]

Silver diethyldithiocarbamate [1470-61-7] M 512.3, pK₁²⁵ 3.36 (for *N,N*-diethyldithiocarbamate). Purified by recrystn from pyridine. Stored in a desiccator in a cool and dark place.

Silver difluoride [7783-95-1] M 145.9, m 690°, d 4.7. Highly **TOXIC** because it liberates HF and F₂. Very *hygroscopic* and reacts violently with H₂O. It is a powerful oxidising agent and liberates O₃ from dilute acids, and I₂ from I⁻ soln. Store in quartz or iron ampoules. White when pure, otherwise it is brown-tinted. Thermally stable up to 700°. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I 241 1963.]

Silver fluoride [7775-41-9] M 126.9, m 435°, b ca 1150°, d 5.852. *Hygroscopic* solid with a solubility of 135g/100mL of H₂O at 15°, and forms an insoluble basic fluoride in moist air. Purified by washing with AcOH and dry *C₆H₆, then kept in a vacuum desiccator at room temperature to remove *benzene and stored in opaque glass bottles. Flaky *hygroscopic* crystals which darken on exposure to light. It *attacks* bone and teeth. [*J Chem Soc* 4538 1952; *Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I 240 1963.]

Silver iodate [7783-97-3] M 282.8, m >200°, d 5.53. Washed with warm dilute HNO₃, then H₂O and dried at 100°, or recrystd from NH₃ soln by adding HNO₃, filtering, washing with H₂O and drying at 100°.

Silver lactate [128-00-7] M 196.9, m ~ 100°. Recrystd from H₂O by adding EtOH. The solid was collected washed with EtOH then Et₂O and dried at 80° to give the dihydrate. White powder soluble in 15 parts of H₂O but only slightly soluble in EtOH. [*Justus Liebigs Ann Chem* 63 89 1847; *Helv Chim Acta* 2 251 1919.]

Silver nitrate [7761-88-8] M 169.9, m 212°, b 444°(dec), d 4.35. Purified by recrystn from hot water (solubility of AgNO₃ in water is 992g/100mL at 100° and 122g/100mL at 0°). It has also been purified by crystn from hot conductivity water by slow addition of freshly distilled EtOH.

CAUTION: avoid using EtOH for washing the ppte; and avoid concentrating the filtrate to obtain further crops of AgNO₃ owing to the risk of EXPLOSION (as has been reported to us) caused by the presence of silver fulminate. When using EtOH in the purification the apparatus should be enveloped in a strong protective shield. [Tully, *News Ed (Am Chem Soc)* 19 3092 1941; Garin and Henderson *J Chem Educ* 47 741 1970; Bretherick, *Handbook of Reactive Chemical Hazards* 4th edn, Butterworths, London, 1985, pp 13-14.] Before being used as a standard in volumetric analysis, analytical reagent grade AgNO₃ should be finely powdered, dried at 120° for 2h, then cooled in a desiccator.

Recovery of silver residues as AgNO₃ [use protective shield during the whole of this procedure] can be achieved by washing with hot water and adding 16M HNO₃ to dissolve the solid. Filter through glass wool and concentrate the filtrate on a steam bath until precipitation commences. Cool the solution in an ice-bath and filter the precipitated AgNO₃. Dry at 120° for 2h, then cool in a desiccator in a vacuum. Store over P₂O₅ in a vacuum in the dark. *AVOID contact with hands due to formation of black stains.*

Silver nitrite [7783-99-5] M 153.9, m 141°(dec), d 4.45. Crystd from hot conductivity water (70mL/g) in the dark. Dried in the dark under vacuum.

Silver(I) oxide [20667-12-3] M 231.7, m ~200°(dec), d 7.13. Leached with hot water in a Soxhlet apparatus for several hours to remove any entrained electrolytes.

Silver (II) oxide [1301-96-8] M 123.9, m >100°(dec), d²⁵ 7.22. Soluble in 40,000 parts of H₂O, and should be protected from light. Stir with an alkaline solution of potassium peroxydisulfate (K₂S₂O₈) at 85-

90°. The black AgO is collected, washed free from sulfate with H₂O made slightly alkaline and dried in air in the dark. [*Inorg Synth* 4 12 1953.]

Silver perchlorate (H₂O) [14242-05-8 (H₂O); 7783-93-9 (anhydr)] M 207.3, pK²⁵ -2.4 to -3.1 (for HClO₄). Refluxed with *benzene (6mL/g) in a flask fitted with a Dean and Stark trap until all the water was removed azeotropically (ca 4h). The soln was cooled and diluted with dry pentane (4mL/g of AgClO₄). The pptd AgClO₄ was filtered off and dried in a desiccator over P₂O₅ at 1mm for 24h [Radell, Connolly and Raymond *J Am Chem Soc* 83 3958 1961]. It has also been recrystallised from perchloric acid. [Caution due to **EXPLOSIVE** nature in the presence of organic matter.]

Silver permanganate [7783-98-4] M 226.8, d 4.49. Violet crystals which can be crystallised from hot H₂O (sol is 9g/L at 20°). Store in the dark. Oxidising agent, decomposed by light.

Silver sulfate [10294-26-5] M 311.8, m 652°, b 1085°(dec), d 5.45. Crystd form hot conc H₂SO₄ contg a trace of HNO₃, cooled and diluted with H₂O. The ppte was filt'd off, washed and dried at 120°.

Silver thiocyanate [1701-93-5] M 165.9, m 265°(dec), d 3.746, pK²⁵ -1.85 (for HSCN). Digest the solid salt with aqueous NH₄NCS, wash thoroughly with H₂O and dry at 110° in the dark. Soluble in dilute aqueous NH₃. Dissolve in strong aqueous NH₄NCS solution, filter and dilute with large volume of H₂O when the Ag salt separates. The solid is washed with H₂O by decantation until free from NCS⁻ ions, collected, washed with H₂O, EtOH and dried in an air oven at 120°. Alternatively dissolve in dilute aqueous NH₃ and single crystals are formed by free evaporation of the solution in air. [*J Chem Soc* 836, 2405 1932; IR and Raman: *Acta Chem Scand* 13 1607 1957; *Acta Cryst* 10 29 1957.]

Silver tosylate [16836-95-6] M 279.1. The anhydrous salt is obtained by recrystn from H₂O. [*Chem Ber* 12 1851 1879.]

Silver trifluoroacetate [2966-50-9] M 220.9, m 251-255°. Extract the salt (Soxhlet) with Et₂O. The extract is filtered and evaporated to dryness, then the powdered residue is completely dried in a vacuum desiccator over silica gel. Solubility in Et₂O is 33.5g in 750mL. It can be recrystd from *C₆H₆ (sol: 1.9g in 30mL of *C₆H₆; and 33.5g will dissolve in 750mL of anhydrous Et₂O). [*J Org Chem* 23 1545 1958; *J Chem Soc* 584 1951.] It is also soluble in trifluoroacetic acid (15.2% at 30°), toluene, *o*-xylene and dioxane [*J Am Chem Soc* 76 4285 1954].

Silver trifluoromethanesulfonate [2923-28-6] M 256.9. Recrystd twice from hot CCl₄ [Alo et al. *J Chem Soc, Perkin Trans 1* 805 1986].

Sodium (metal) [7440-23-5] M 23.0, m 97.5°, d 0.97. The metal was placed on a coarse grade of sintered-glass filter, melted under vacuum and forced through the filter using argon. The Pyrex apparatus was then re-evacuated and sealed off below the filter, so that the sodium could be distilled at 460° through a side arm and condenser into a receiver bulb which was then sealed off [Gunn and Green *J Am Chem Soc* 80 4782 1958]. **EXPLODES and IGNITES in water.**

Sodium acetate [127-09-3] M 82.0, m 324°, d 1.53. Crystd from acetic acid and pumped under vacuum for 10h at 120°. Alternatively, crystd from aqueous EtOH, as the trihydrate. This material can be converted to the anhydrous salt by heating slowly in a porcelain, nickel or iron dish, so that the salt liquefies. Steam is evolved and the mass again solidifies. Heating is now increased so that the salt melts again. (NB: if it is heated too strongly, the salt chars.) After several minutes, the salt is allowed to solidify and cooled to a convenient temperature before being powdered and bottled (water content should now less than 0.02%).

Sodium acetylide [1066-26-8] M 48.0. It disproportionates at ca 180° to sodium carbide. It sometimes contains diluents, e.g. xylene, butyl ether or dioxane which can be removed by filtration followed by a vacuum at 65-60°/5mm. Alternatively the acetylide is purged with HC≡CH at 100-125° to remove diluent. NaC₂H adsorbs 2.2x, 2.0x and 1.6x its wt of xylene, butyl ether and dioxane respectively. Powdered NaC₂H is yellow or yellow-gray in colour and is relatively stable. It can be heated to ca 300° in the absence of air. Although no

explosion or evolution of gas occurs, it turns brown due to disproportionation. At 170-190° in air it ignites slowly and burns smoothly. At 215-235° in air it flash-ignites and burns quickly. It can be dropped into a *slight* excess of H₂O without flashing or burning but vigorous evolution of HC≡CH (**HIGHLY FLAMMABLE IN AIR**) occurs. The sample had been stored in the absence of air for one year without deterioration. Due to the high flammability of HC≡CH the salt should be stored dry, and treated with care. After long storage, NaC≡CH can be redissolved in liquid NH₃ and used for the same purposes as the fresh material. However it may be slightly turbid due to the presence of moisture. [*J Org Chem* **22** 649 1957; *J Am Chem Soc* **77** 5013 1955; *Inorg Synth* **2** 76, 81 1946; *Org Synth* **30** 15 1950.] See p. 89, Chapter 4 for preparation.

Sodium alginate [9005-38-3]. Freed from heavy metal impurities by treatment with ion-exchange resins (Na⁺-form), or with a dilute solution of the sodium salt of EDTA. Also dissolved in 0.1M NaCl, centrifuged and fractionally pptd by gradual addition of EtOH or 4M NaCl. The resulting gels were centrifuged off, washed with aq EtOH or acetone, and dried under vacuum. [Büchner, Cooper and Wassermann *J Chem Soc* 3974 1961.]

Sodium *n*-alkylsulfates. Crystd from EtOH/Me₂CO [Hashimoto and Thomas *J Am Chem Soc* **107** 4655 1985].

Sodium amide [7782-92-5] **M 39.0, m 210°.** It reacts *violently* with H₂O and is soluble in liquid NH₃ (1% at 20°). It should be stored in wax-sealed container in small batches. It is very *hygroscopic* and absorbs CO₂ and H₂O. If the solid is discoloured by being yellow or brown in colour then it should be destroyed as it can be highly **EXPLOSIVE**. It should be replaced if discoloured. It is best destroyed by covering with much toluene and slowly adding dilute EtOH with stirring until all the ammonia is liberated (FUME CUPBOARD). [*Inorg Synth* **1** 74 1939; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol I 465 1963; *Org Synth Coll Vol III*, 778 1955.]

Sodium 4-aminobenzoate [555-06-6] **M 159.1.** Crystd from water.

Sodium 4-aminosalicylate (2H₂O) [6018-19-5] **M 175.1.** Crystd from water at room temperature (2mL/g) by adding acetone and cooling.

Sodium ammonium hydrogen phosphate [13011-54-6] **M 209.1, m 79°(dec), d 1.55.** Crystd from hot water (1mL/g).

Sodium amylpenicillin [575-47-3] **M 350.4.** Crystd from moist acetone or moist ethyl acetate.

Sodium anthraquinone-1,5-disulfonate (H₂O) [853-35-0] **M 412.3.** Separated from insoluble impurities by continuous extraction with water. Crystd twice from hot water and dried under vacuum.

Sodium anthraquinone-1-sulfonate (H₂O) [107439-61-2] **M 328.3.** Crystd from hot water (4mL/g) after treatment with active charcoal, or from water by addition of EtOH. Dried under vacuum over CaCl₂, or in an oven at 70°. Stored in the dark.

Sodium anthraquinone-2-sulfonate (H₂O) [131-08-8] **M 328.3.** See 9,10-anthraquinone-2-sulfonic acid disodium salt on p. 395.

Sodium antimonyl tartrate [34521-09-0] **M 308.8.** Crystd from water.

Sodium arsenate (7H₂O) [10048-95-0] **M 312.0, m 50 (loses 5H₂O), m 130°, d 1.88 pK₁^{2.5} 2.22, pK₂^{2.5} 6.98 (for H₃AsO₄).** Crystd from water (2mL/g).

Sodium azide [26628-22-8] **M 65.0, m 300°(dec, explosive), pK^{2.5} 4.72 (for HN₃).** Crystd from hot water or from water by the addition of absolute EtOH or acetone. Also purified by repeated crystn from an aqueous solution saturated at 90° by cooling it to 10°, and adding an equal volume of EtOH. The crystals were washed with acetone and the azide dried at room temperature under vacuum for several hours in an

Abderhalden pistol. [Das et al. *J Chem Soc, Faraday Trans 1* 78 3485 1982.] **HIGHLY POISONOUS and potentially explosive.**

Sodium barbitone (sodium 5,5-diethylbarbiturate) [144-02-5] M 150.1, pK_1^{25} 3.99, pK_2^{25} 12.5 (barbituric acid). Crystd from water (3mL/g) by adding an equal volume of EtOH and cooling to 5°. Dried under vacuum over P_2O_5 .

Sodium benzenesulfinate [873-55-2] M 164.2, m >300°. See benzenesulfonic acid sodium salt on p. 400.

Sodium benzenesulfonate [515-42-4] M 150.1, pK_1^{25} 0.70 (2.55) (for PhSO_3H_2). Crystd from EtOH or aqueous 70-100% MeOH, and dried under vacuum at 80-100°.

Sodium benzoate [532-32-1] M 144.1. Crystd from EtOH (12mL/g).

Sodium benzylpenicillin see *N*-benzylpenicillin sodium salt on p. 514 in Chapter 6.

Sodium bicarbonate [144-55-8] M 84.0, m ~50°(dec, $-\text{CO}_2$). Crystd from hot water (6mL/g). The solid should not be heated above 40° due to the formation of carbonate.

Sodium bis(trimethylsilyl)amide (hexamethyl disilazane sodium salt) [1070-89-9] M 183.4, m 165-167°(sintering at 140°). It can be sublimed at 170°/2 Torr (bath temp 220-250°) onto a cold finger, and can be recrystd from $^*\text{C}_6\text{H}_6$ (sol: 10g in 100mL at 60°). It is slightly soluble in Et_2O and is decomposed by H_2O . [*Chem Ber* 94 1540 1961.]

Sodium bisulfite [7631-90-5] M 104.1, d 1.48. Crystd from hot H_2O (1mL/g). Dried at 100° under vac for 4h.

Sodium borate (borax) [1330-43-4] M 201.2, m 741°, d 2.37. Most of the water of hydration was removed from the decahydrate by evacuation at 25° for three days, followed by heating to 100° and evacuation with a high-speed diffusion pump. The dried sample was then heated gradually to fusion (above 966°), allowed to cool gradually to 200°, then transferred to a desiccator containing P_2O_5 [Grenier and Westrum *J Am Chem Soc* 78 6226 1956].

Sodium borate (decahydrate, hydrated borax) [1303-96-4] M 381.2, m 75°(loses $5\text{H}_2\text{O}$ at 60°), d 1.73. Crystd from water (3.3mL/g) keeping below 55° to avoid formation of the pentahydrate. Filtered at the pump, washed with water and equilibrated for several days in a desiccator containing an aqueous solution saturated with respect to sucrose and NaCl. Borax can be prepared more quickly (but its water content is somewhat variable) by washing the recrystd material at the pump with water, followed by 95% EtOH, then Et_2O , and air dried at room temperature for 12-18h on a clock glass.

Sodium borohydride [16940-66-2] M 37.8, m ~400°(dec), d 1.07. After adding NaBH_4 (10g) to freshly distilled diglyme (120mL) in a dry three-necked flask fitted with a stirrer, nitrogen inlet and outlet, the mixture was stirred for 30min at 50° until almost all of the solid had dissolved. Stirring was stopped, and, after the solid had settled, the supernatant liquid was forced under N_2 pressure through a sintered-glass filter into a dry flask. [The residue was centrifuged to obtain more of the solution which was added to the bulk.] The solution was cooled slowly to 0° and then decanted from the white needles that separated. The crystals were dried by pumping for 4h to give anhydrous NaBH_4 . Alternatively, after the filtration at 50° the solution was heated at 80° for 2h to give a white ppte of substantially anhydrous NaBH_4 which was collected on a sintered-glass filter under N_2 , then pumped at 60° for 2h [Brown, Mead and Subba Rao *J Am Chem Soc* 77 6209 1955].

NaBH_4 has also been crystd from isopropylamine by dissolving it in the solvent at reflux, cooling, filtering and allowing the solution to stand in a filter flask connected to a Dry-ice/acetone trap. After most of the solvent was passed over into the cold trap, crystals were removed with forceps, washed with dry diethyl ether and dried under vacuum. [Kim and Itoh *J Phys Chem* 91 126 1987.] Somewhat less pure crystals were obtained more rapidly by using Soxhlet extraction with only a small amount of solvent and extracting for about 8h. The

crystals that formed in the flask were filtered off, then washed and dried as before. [Stockmayer, Rice and Stephenson *J Am Chem Soc* **77** 1980 1955.] Other solvents used for crystallisation include water and liquid ammonia.

Sodium bromate [7789-38-0] **M 150.9, m 381°**, **d 3.3**. Crystd from hot water (1.1mL/g) to decrease contamination by NaBr, bromine and hypobromite. [Noszticzius et al. *J Am Chem Soc* **107** 2314 1985.]

Sodium bromide [7647-15-6] **M 102.9, m 747°**, **b 1390°**, **d 3.2**. Crystd from water (0.86mL/g) between 50° and 0°, and dried at 140° under vacuum (this purification may not eliminate chloride ion).

Sodium 4-bromobenzenesulfonate [5015-75-8] **M 258.7**. Crystd from MeOH, EtOH or distd water.

Sodium tert-butoxide [865-48-5] **M 96.1**. It sublimes at 180°/1 Torr. Its solubility in *tert*-BuOH is 0.208M at 30.2° and 0.382M at 60°, and is quite soluble in tetrahydrofuran (32g/100g). It should not be used if it has a brown colour. [*J Am Chem Soc* **78** 4364, 3614 1956, *Inorg Synth* **1** 87 1939; IR: *J Org Chem* **21** 156 1956.]

Sodium butyrate [156-54-7] **M 110.1**. Prepared by neutralisation of the acid and recrystn from EtOH.

Sodium cacodylate (3H₂O) [124-65-2] **M 214.0, m 60°**. Crystd from aqueous EtOH.

Sodium carbonate [497-19-8] **M 106.0, m 858°**, **d 2.5**. Crystd from water as the decahydrate which was redissolved in water to give a near-saturated soln. By bubbling CO₂, NaHCO₃ was pptd. It was filtered, washed and ignited for 2h at 280° [MacLaren and Swinehart *J Am Chem Soc* **73** 1822 1951]. Before being used as a volumetric standard, analytical grade material should be dried by heating at 260-270° for 0.5h and allowed to cool in a desiccator. For preparation of primary standard sodium carbonate, see *Pure Appl Chem* **25** 459 1969. After 3 recrystns tech grade had Cr, Mg, K, P, Al, W, Sc and Ti at 32, 9.4, 6.6, 3.6, 2.4, 0.6, 0.2 and 0.2 ppm resp; another technical source had Cr, Mg, Mo, P, Si, Sn and Ti at 2.6, 0.4, 4.2, 13.4, 32, 0.6, 0.8 ppm resp.

Sodium carboxymethylcellulose [9004-32-4]. Dialysed for 48h against distilled water.

Sodium cetyl sulfate [1120-01-0] **M 344.5**. See sodium hexadecylsulfate on p. 471.

Sodium chlorate [7775-09-9] **M 106.4, m 248°**, **b >300°(dec)**, **d 2.5**. Crystd from hot water (0.5mL/g).

Sodium chloride [7647-14-5] **M 58.4, m 800.7°**, **b 1413°**, **d 2.17**. Crystd from saturated aqueous solution (2.7mL/g) by passing in HCl gas, or by adding EtOH or acetone. Can be freed from bromide and iodide impurities by adding chlorine water to an aqueous solution and boiling for some time to expel free bromine and iodine. Traces of iron can be removed by prolonged boiling of solid NaCl in 6M HCl, the crystals then being washed with EtOH and dried at *ca* 100°. Sodium chloride has been purified by sublimation in a stream of pre-purified N₂ and collected by electrostatic discharge [Ross and Winkler *J Am Chem Soc* **76** 2637 1954]. For use as a primary analytical standard, analytical reagent grade NaCl should be finely ground, dried in an electric furnace at 500-600° in a platinum crucible, and allowed to cool in a desiccator. For most purposes, however, drying at 110-120° is satisfactory.

Sodium chlorite [7758-19-2] **M 90.4, m ~180°(dec)**. Crystd from hot water and stored in a cool place. Has also been crystd from MeOH by counter-current extraction with liquid ammonia [Curti and Locchi *Anal Chem* **29** 534 1957]. Major impurity is chloride ion; can be recrystallised from 0.001M NaOH.

Sodium 4-chlorobenzenesulfonate [5138-90-9] **M 214.6, pK_{Est} <0 (for SO₃H)**. Crystd twice from MeOH and dried under vacuum.

Sodium 3-chloro-5-methylbenzenesulfonate [5138-92-1] M 228.7, $pK_{\text{Est}} < 0$ (for SO_3H). Crystd twice from MeOH and dried under vacuum.

Sodium chromate ($4\text{H}_2\text{O}$) [10034-82-9] M 234.0, m $\sim 20^\circ$ (for $10\text{H}_2\text{O}$), d, 2.7, $pK_1^{25} 0.74$, $pK_2^{25} 6.49$ (for H_2CrO_4). Crystd from hot water (0.8mL/g).

dl-Sodium creatine phosphate ($4\text{H}_2\text{O}$) [922-32-7] M 327.1. See creatine phosphate di-Na salt on p. 523 in Chapter 6.

Sodium cyanate [917-61-3] M 65.0, m 550° , $d_4^{20} 1.893$, $pK^{25} 3.47$ (for HCNO). Colourless needles from EtOH. Solubility in EtOH is 0.22g/100g at 0°C . Soluble in H_2O but can be recrystallised from small volumes of it.

Sodium cyanoborohydride [25895-60-7] M 62.8, m $240\text{-}242^\circ(\text{dec})$, $d^{28} 1.20$. Very hygroscopic solid, soluble in H_2O (212% at 29° , 121% at 88°), tetrahydrofuran (37% at 28° , 42.2% at 62°), very soluble in EtOH but insoluble in Et_2O , $^*\text{C}_6\text{H}_6$ and hexane. It is stable to acid up to pH 3 but is hydrolysed in 12N HCl. The rate of hydrolysis at pH 3 is 10^{-8} that of NaBH_4 . The fresh commercially available material is usually sufficiently pure. If very pure material is required one of the following procedures must be used [*Synthesis* 135 1975]: (a) The NaBH_3CN is dissolved in tetrahydrofuran (20% w/v), filtered and the filtrate is treated with a fourfold volume of CH_2Cl_2 . The solid is collected and dried in a vacuum [*Inorg Chem.* 9 2146 1970].

Dissolve the NaBH_3CN in dry MeNO_2 , filter, and pour the filtrate into a 10-fold volume of CCl_4 with vigorous stirring. The white ppt is collected, washed several times with CCl_4 and dried in a vacuum [*Inorg Chem* 9 624 1970]. (b) When the above procedures fail to give a clean product then dissolve the NaBH_3CN (10g) in tetrahydrofuran (80mL) and add N MeOH/HCl until the pH is 9. Pour the solution with stirring into dioxane (250mL). The solution is filtered, and heated to reflux. A further volume of dioxane (150mL) is added slowly with swirling. The solution is cooled slowly to room temp then chilled in ice and the crystalline dioxane complex is collected, dried in a vacuum for 4h at 25° , then 4h at 80° to yield the amorphous dioxane-free powder is 6.7g with purity $>98\%$ [*J Am Chem Soc* 93 2897 1971]. The purity can be checked by iodometric titration [*Anal Chem* 91 4329 1969].

Sodium p-cyrenesulfonate [77060-21-0] M 236.3. Dissolved in water, filtered and evaporated to dryness. Crystd twice from absolute EtOH and dried at 110° .

Sodium decanoate (sodium caproate) [1002-62-6] M 194.2. Neutralised by adding a slight excess of the free acid, recovering the excess acid by Et_2O extraction. The salt is crystd from solution by adding pure acetone, repeating the steps several times, then dried in an oven at $ca 110^\circ$ [Chaudhury and Awuwallia *Trans Faraday Soc* 77 3119 1981].

Sodium 1-decanesulfonate [13419-61-9] M 244.33. Recrystd from absolute EtOH and dried over silica gel.

Sodium n-decylsulfate [142-87-0] M 239.3. Rigorously purified by continuous Et_2O extraction of a 1% aqueous solution for two weeks.

Sodium deoxycholate (H_2O) [302-95-4] M 432.6, $[\alpha]_{\text{D}}^{20} +48^\circ$ (c 1, EtOH). Crystd from EtOH and dried in an oven at 100° . The solution is freed from soluble components by repeated extraction with acid-washed charcoal.

Sodium dibenzylthiocarbamate [55310-46-8] M 295.4, m $230^\circ(\text{dec})$, $pK^{20} 3.13$ (for monobenzylthiocarbamic acid). The free acid when recrystd twice from dry Et_2O has m $80\text{-}82^\circ$. The Na salt is repped from aqueous EtOH or EtOH by addition of Et_2O or Me_2CO [*Anal Chem* 50 896 1978]. The NH_4 salt has m $130\text{-}133^\circ$; Cu salt (yellow crystals) has m $284\text{-}286^\circ$ and the Ti salt has m $64\text{-}70^\circ$.

Sodium 2,5-dichlorobenzenesulfonate [5138-93-2] M 249.0, $pK_{\text{Est}} < 0$ (for SO_3H). Crystd from MeOH, and dried under vacuum.

Sodium dichromate [7789-12-0] M 298.0, m 84.6° (2H₂O), 356° (anhydr); b 400°(dec), d₄²⁵ 2.348. Crystd from small volumes of H₂O by evaporation to crystallisation. Solubility in H₂O is 238% at 0° and 508% at boiling. Red dihydrate is slowly dehydrated by heating at 100° for long periods. It is deliquescent, a powerful oxidising agent-*do not place in contact with skin- wash immediately as it is caustic*. (Possible carcinogen).

Sodium diethyldithiocarbamate (3H₂O) [20624-25-3] M 225.3, m 94-96°(anhydr), pK²⁰ 3.65 (diethyldithiocarbamic acid). Recrystd from water.

Sodium di(ethylhexyl)sulfosuccinate (Aerosol-OT) [577-11-7] M 444.6. Dissolved in MeOH and inorganic salts which ppted were filtered off. Water was added and the solution was extracted several times with hexane. The residue was evaporated to one fifth its original volume, *benzene was added and azeotropic distillation was continued until no water remained. Solvent was then evaporated. The white solid was crushed and dried in vacuum over P₂O₅ for 48h [El Seoud and Fendler *J Chem Soc, Faraday Trans 1* 71 452 1975].

Sodium diethyloxaloacetate [63277-17-8] M 210.2. Extracted several times with boiling Et₂O (until the solvent remained colourless) and then the residue was dried in air.

Sodium diformylamide [18197-26-7] M 95.0. Grind under dry tetrahydrofuran (fumehood), filter and wash with this solvent then dry in vacuum. [IR and prepn: *Synthesis* 122 1990; *Chem Ber* 100 355 1967, 102 4089 1969.]

Sodium dihydrogen orthophosphate (2H₂O) [13472-35-0 (2H₂O); 10049-21-5 (H₂O); 7558-80-7 (anhydr)] M 156.0, m 60°(dec), d 1.91. Crystd from warm water (0.5mL/g) by chilling.

Sodium 2,2'-dihydroxy-1-naphthaleneazobenzene-5'-sulfonate [2092-55-9] M 354.3. See Solochrome Violet R on p. 352 in Chapter 4.

Sodium 2,4-dihydroxyphenylazobenzene-4'-sulfonate [547-57-9] M 304.2. Crystd from absolute EtOH.

Sodium *p*-(*p*-dimethylaminobenzeneazo)-benzenesulfonate [23398-40-5] M 327.3. Crystd from water.

Sodium *p*-dimethylaminoazobenzene-*o*'-carboxylate [845-10-3] M 219.2. Ppted from aqueous soln as the free acid which was recrystallised from 95% EtOH, then reconverted to the sodium salt.

Sodium *p*-dimethylaminoazobenzene-*p*'-carboxylate [845-46-5] M 219.2. Ppted from aqueous soln as the free acid which was recrystallised from 95% EtOH, then reconverted to the sodium salt.

Sodium 2,4-dimethylbenzenesulfonate [827-21-4] M 208.2. Crystd from MeOH and dried under vacuum.

Sodium 2,5-dimethylbenzenesulfonate [827-19-0] M 208.2. Dissolved in distilled water, filtered, then evaporated to dryness. Crystd twice from absolute EtOH or MeOH and dried at 110° under vacuum.

Sodium dimethyldithiocarbamate hydrate [128-04-1] M 143.2, m 106-108°, 120-122°, pK²⁵ 3.36 (diethyldithiocarbamic acid). Crystallise from a small volume of H₂O, or dissolve in minimum volume of H₂O and add cold Me₂CO and dry in air. The solution in Me₂CO is ~50g/400mL. The dihydrate loses H₂O on heating at 115° to give the hemi hydrate which decomposes on further heating [IR: *Can J Chem.* 34 1096 1956].

Sodium *N,N*-dimethylsulfanilate [2244-40-8] M 223.2, m >300°. Crystd from water.

Sodium dithionite (2H₂O) [7631-94-9] M 242.1, m 110°(loses 2H₂O), 267°(dec), d 2.19, pK_{Est(1)} -3.4, pK₂²⁵ 0.49 (for dithionic acid). Crystd from hot water (1.1mL/g) by cooling.

Sodium dodecanoate (sodium laurate) [629-25-4] M 222.3, pK²⁰ 5.3 (-COOH). Neutralised by adding a slight excess of dodecanoic acid, removing it by ether extraction. The salt is recrystd from aq soln by adding pure Me₂CO and repeating the process (see sodium decanoate on p. 468). Also recrystd from MeOH.

Sodium 1-dodecanesulfonate [2386-53-0] M 272.4. Twice recrystd from EtOH.

Sodium dodecylbenzenesulfonate [25155-30-0] M 348.5. Recrystd from propan-2-ol.

Sodium dodecylsulfate (SDS, sodium laurylsulfate) [151-21-3] M 288.4, m 204-207°. Purified by Soxhlet extraction with pet ether for 24h, followed by dissolution in acetone:MeOH:H₂O 90:5:5(v/v) and recrystn [Politi et al. *J Phys Chem* **89** 2345 1985]. Also purified by two recrystns from absolute EtOH, aqueous 95% EtOH, MeOH, isopropanol or a 1:1 mixture of EtOH:isopropanol to remove dodecanol, and dried under vacuum [Ramesh and Labes *J Am Chem Soc* **109** 3228 1987]. Also purified by foaming [see Cockbain and McMullen *Trans Faraday Soc* **47** 322 1951] or by liquid-liquid extraction [see Harrold *J Colloid Sci* **15** 280 1960]. Dried over silica gel. For DNA work it should be dissolved in excess MeOH passed through an activated charcoal column and evaporated until it crystallises out. Also purified by dissolving in hot 95% EtOH (14mL/g), filtering and cooling, then drying in a vacuum desiccator. Alternatively, it was crystd from H₂O, vacuum dried, washed with anhydrous Et₂O, vacuum dried. These operations were repeated five times [Maritato *J Phys Chem* **89** 1341 1985; Lennox and McClelland *J Am Chem Soc* **108** 3771 1986; Dressik *J Am Chem Soc* **108** 7567 1986].

Sodium ethoxide [141-52-6] M 68.1. *Hygroscopic* powder which should be stored under N₂ in a cool place. Likely impurity is EtOH which can be removed by warming at 60-80° under high vacuum. Hydrolysed by H₂O to yield NaOH and EtOH. Other impurities, if kept in air for long periods are NaOH and Na₂CO₃. In this case the powder cannot be used if these impurities affect the reactivity and a fresh sample should be acquired [IR: *J Org Chem* **21** 156 1956].

Sodium ethylmercurithiosalicylate [54-64-8] M 404.8. Crystd from ethanol-diethyl ether

Sodium ethylsulfate [546-74-7] M 166.1. Recrystd three times from MeOH-Et₂O and vacuum dried.

Sodium ferricyanide (H₂O) [14217-21-1; 13601-19-9 (anhydr)] M 298.9, pK²⁵ <1 (for ferricyanide). Crystd from hot water (1.5mL/g) or by precipitation from 95% EtOH.

Sodium ferrocyanide (10H₂O) [13601-19-9] M 484.1, m 50-80° (loses 10H₂O), 435°(dec), d 1.46, pK₃²⁵ 2.57, pK₄²⁵ 4.35 (for ferrocyanide). Crystd from hot water (0.7mL/g), until free of ferricyanide as shown by absence of Prussian Blue formation with ferrous sulfate soln.

Sodium fluoride [7681-49-4] M 42.0, m 996°, b 1695°, d 2.56. Crystd from water by partial evaporation in a vacuum desiccator, or dissolved in water, and ca half of it pptd by addition of EtOH. Ppte was dried in an air oven at 130° for one day, and then stored in a desiccator over KOH.

Sodium fluoroacetate (mono) [62-74-8] M 100.0, m 200-205°(dec). A free flowing white TOXIC powder which is purified by dissolving in ca 4 parts of H₂O and the pH is checked. If it is alkaline, add a few drops of FCH₂CO₂H to make the solution just acidic. Evaporate (fumehood) on a steam bath until crystals start to separate, cool and filter the solid off. More solid can be obtained by adding EtOH to the filtrate. Dry at 100° in vacuum. [*J Chem Soc* 1778 1948.]

Sodium fluoroborate [13755-29-8] M 109.8, m 384°, d 2.47, pK -4.9 (for fluoroboric acid H₃O⁺BF₄⁻). Crystd from hot water (50mL/g) by cooling to 0°. Alternatively, purified from insoluble material by dissolving in a minimum amount of water, then fluoride ion was removed by adding conc lanthanum nitrate in excess. After removing lanthanum fluoride by centrifugation, the supernatant was passed

through a cation-exchange column (Dowex 50, Na⁺-form) to remove any remaining lanthanum [Anbar and Guttman *J Phys Chem* **64** 1896 1960]. Also recrystd from anhydrous MeOH and dried in a vacuum at 70° for 16h. It is affected by moisture. [Delville et al. *J Am Chem Soc* **109** 7293 1987.]

Sodium fluorosilicate [16893-85-9] **M 188.1**. Crystd from hot water (40mL/g) by cooling.

Sodium formaldehyde sulfoxylate dihydrate (sodium hydroxymethylsulfinate, Rongalite) [149-44-0] **M 134.1, m 63-64° (dihydrate)**. Crystallises from H₂O as the dihydrate, decomposes at higher temperatures. Store in a closed container in a cool place. It is insoluble in EtOH and Et₂O and is a good reducing agent. [X-ray structure: *J Chem Soc* 3064 1955.] Note that this compound {HOCH₂SO₂Na} should not be confused with formaldehyde sodium bisulfite adduct {HOCH₂SO₃Na} from which it is prepared by reduction with Zn.

Sodium formate (anhydrous) [141-53-7] **M 68.0, m 253°, d 1.92**. A saturated aqueous solution at 90° (0.8mL water/g) was filtered and allowed to cool slowly. (The final temperature was above 30° to prevent formation of the hydrate.) After two such crystns the crystals were dried in an oven at 130°, then under high vacuum. [Westrum, Chang and Levitin *J Phys Chem* **64** 1553 1960; Roecker and Meyer *J Am Chem Soc* **108** 4066 1986.] The salt has also been recrystd twice from 1mM DTPA (diethylenetriaminepentaacetic acid which was recrystd 4x from MilliQ water and dried in a vac), then twice from water [Bielski and Thomas *J Am Chem Soc* **109** 7761 1987].

Sodium D-gluconate [527-07-1] **M 218.1, m 200-205°dec, [α]_D²⁰ +14°, [α]_D²⁰ +12 (c 20, H₂O)**. Crystallise from a small volume of H₂O (sol 59g/100mL at 25°), or dissolve in H₂O and add EtOH since it is sparingly soluble in EtOH. Insoluble in Et₂O. It forms a Cu complex in alkaline soln and a complex with Fe in neutral solution. [*J Am Chem Soc* **81** 5302 1959.]

Sodium glycochenodeoxycholate [16564-43-5] **M 472.6**. Dissolved in EtOH, filtered and concentrated to crystallisation, and recrystallised from a little EtOH.

Sodium glycocholate [863-57-0] **M 488.6**. Dissolved in EtOH, filtered and concentrated to crystallisation, and recrystallised from a little EtOH.

Sodium glycolate (2H₂O) [2836-32-0] **M 98.0**. Ppted from aqueous solution by EtOH, and air dried.

Sodium hexadecylsulfate [1120-01-0] **M 344.5**. Recrystd from absolute EtOH or MeOH and dried in vac [Abu Hamdiyyah and Rahman *J Phys Chem* **91** 1531 1987].

Sodium hexafluorophosphate [21324-39-0] **M 167.9, pK₁^{2.5}~ 0.5, pK₂^{2.5} 5.12 (for fluorophosphoric acid H₂PO₃F)**. Recrystd from acetonitrile and vacuum dried for 2 days at room temperature. It is an irritant and is *hygroscopic*. [Delville et al. *J Am Chem Soc* **109** 7293 1987.]

Sodium hexanitrocobaltate III (Na₃[Co(NO)₆]) [13600-98-1] **M 403.9**. Dissolve (ca 60g) in H₂O (300mL), filter to obtain a clear solution, add 96% EtOH (250mL) with vigorous stirring. Allow the ppte to settle for 2h, filter, wash with EtOH (4 x 25mL), twice with Et₂O and dry in air [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer) Vol II* 1541 1965]. Yellow to brown yellow crystals which are very soluble in H₂O, are decomposed by acid and form an insoluble K salt. Used for estimating K.

Sodium hydrogen diglycollate [50795-24-9] **M 156.1**. Crystd from hot water (7.5mL/g) by cooling to 0° with constant stirring, the crystals being filtered off on to a sintered-glass funnel and dried at 110° overnight.

Sodium hydrogen oxalate (2H₂O) [1186-49-8] **M 130.0, m 100°(loses 2H₂O), b 200°(dec)**. Crystd from hot water (5mL/g) by cooling.

Sodium hydrogen succinate [2922-54-5] **M 140.0**. Crystd from water and dried at 110°.

Sodium hydrogen *d*-tartrate [526-94-3] **M 190.1, m 100°(loses H₂O), b 234°, [α]₅₄₆ +26° (c 1, H₂O).** Crystd from warm water (10mL/g) by cooling to 0°.

Sodium hydroxide (anhydrous) [1310-73-2] **M 40.0, m 323°, b 1390°, d 2.13.** Common impurities are water and sodium carbonate. Sodium hydroxide can be purified by dissolving 100g in 1L of pure EtOH, filtering the solution under vacuum through a fine sintered-glass disc to remove insoluble carbonates and halides. (This and subsequent operations should be performed in a dry, CO₂-free box.) The soln is concentrated under vacuum, using mild heating, to give a thick slurry of the mono-alcoholate which is transferred to a coarse sintered-glass disc and pumped free of mother liquor. After washing the crystals several times with purified alcohol to remove traces of water, they are vacuum dried, with mild heating, for about 30h to decompose the alcoholate, leaving a fine white crystalline powder [Kelly and Snyder *J Am Chem Soc* 73 4114 1951].

Sodium hydroxide solutions (caustic), pK²⁵ 14.77. Carbonate ion can be removed by passage through an anion-exchange column (such as Amberlite IRA-400; OH⁻-form). The column should be freshly prepared from the chloride form by slow prior passage of sodium hydroxide soln until the effluent gives no test for chloride ions. After use, the column can be regenerated by washing with dilute HCl, then water. Similarly, other metal ions are removed when a 1M (or more dilute) NaOH soln is passed through a column of Dowex ion-exchange A-1 resin in its Na⁺-form.

Alternatively, carbonate contamination can be reduced by rinsing sticks of NaOH (analytical reagent quality) rapidly with H₂O, then dissolving in distilled H₂O, or by preparing a concentrated aqueous soln of NaOH and drawing off the clear supernatant liquid. (Insoluble Na₂CO₃ is left behind.) Carbonate contamination can be reduced by adding a slight excess of conc BaCl₂ or Ba(OH)₂ to a NaOH soln, shaking well and allowing the BaCO₃ ppte to settle. If the presence of Ba in the soln is unacceptable, an electrolytic purification can be used. For example, sodium amalgam is prepared by the electrolysis of 3L of 30% NaOH with 500mL of pure mercury for cathode, and a platinum anode, passing 15 Faradays at 4Amps, in a thick-walled polyethylene bottle. The bottle is then fitted with inlet and outlet tubes, the spent soln being flushed out by CO₂-free N₂. The amalgam is then washed thoroughly with a large volume of deionised water (with the electrolysis current switched on to minimize loss of Na). Finally, a clean steel rod is placed in contact in the solution with the amalgam (to facilitate hydrogen evolution), reaction being allowed to proceed until a suitable concentration is reached, before being transferred to a storage vessel and diluted as required [Marsh and Stokes *Aust J Chem* 17 740 1964].

Sodium 2-hydroxy-4-methoxybenzophenone-5-sulfonate [6628-37-1] **M 330.3.** Crystd from MeOH and dried under vacuum.

Sodium *p*-hydroxyphenylazobenzene-*p*'-sulfonate [2623-36-1] **M 288.2.** Crystd from 95% EtOH.

Sodium hypophosphite monohydrate [10039-56-2] **M 106.0 (see pK of hypophosphorous acid).** Dissolve in boiling EtOH, cool and add dry Et₂O till all the salt separates. Collect and dry in vacuum. It is soluble in 1 part of H₂O. It liberates PH₃ on heating and can *ignite* spontaneously when heated. The anhydrous salt is soluble in ethylene glycol (33% w/w) and propylene glycol (9.7%) at 25°.

Sodium iodate [7681-55-2] **M 197.9, m dec on heating, d 4.28.** Crystd from water (3mL/g) by cooling.

Sodium iodide [7681-82-5] **M 149.9, m 660°, b 1304°, d 3.67.** Crystd from water/ethanol soln and dried for 12h under vacuum, at 70°. Alternatively, dissolved in acetone, filtered and cooled to -20°, the resulting yellow crystals being filtered off and heated in a vacuum oven at 70° for 6h to remove acetone. The NaI was then crystd from very dilute NaOH, dried under vacuum, and stored in a vacuum desiccator [Verdin *Trans Faraday Soc* 57 484 1961].

Sodium ionophore I (ETH 227) (*N,N,N*'-triheptyl-*N,N,N*'-trimethyl-4,4',4''-propylidynetris(3-oxabutynamide) [61183-76-4] **M 642.0.** It is purified (*ca* 200mg) by TLC on Kieselgel F₂₅₄ with CHCl₃/Me₂CO (1:1) as solvent, followed by HPLC (50mg) with an octadecyltrimethylsilane modified column (Mercksorb SI 100, 10µm) [IR, NMR, MS: *Helv Chim Acta* 59 2417 1976].

Sodium ionophore V (ETH 4120) [4-octadecanoyloxymethyl-*N,N,N',N'*-tetracyclohexyl-1,2-phenylenedioxydiacetamide] [129880-73-5] M 849.3. Purified by recrystn from EtOAc. [Preparation and properties: *Anal Chim Acta* 233 295 1990].

Sodium ionophore VI {bis[(12-crown-4)methyl]dodecyl methyl malonate} [80403-59-4] M 662.9. Purified by gel permeation or column chromatography. [Preparation and NMR data: *J Electroanal Chem* 132 99 1982.]

Sodium isopropylxanthate [140-93-2] M 158.2, pK 2.16 (for -S⁻). Crystd from ligroin/ethanol.

Sodium laurate [629-25-4] M 222.3. See sodium dodecanoate on p. 470.

Sodium *RS*-mandelate [114-21-6] M 174.1. Crystd from 95% EtOH.

Sodium 2-mercaptoethanesulfonate (MESNA) [19767-45-4] M 164.2, pK₁²⁰ < 0 (SO₃⁻), pK₂²⁰ 9.53 (SH). It can be recrystd from H₂O and does not melt below 250°. It can be purified further by converting to the free acid by passing a 2M soln through an ion exchange (Amberlite IR-120) column in the acid form, evaporating the eluate in a vacuum to give the acid as a viscous oil (readily dec) which can be checked by acid and SH titration. It is then dissolved in H₂O, carefully neutralised with aqueous NaOH, evaporated and recrystd from H₂O [*J Am Chem Soc* 77 6231 1955].

Sodium metanilate [1126-34-7] M 195.2. Crystd from hot water.

Sodium metaperiodate (NaIO₄) [7790-28-5] M 213.9, m ~300°(dec), d 4.17. Crystd from hot water.

Sodium metasilicate (5H₂O) [6834-92-0] M 212.1, m 1088°, d 2.4. Crystd from aqueous 5% NaOH solution.

Sodium methanethiolate [sodium methylmercaptide] [5188-07-8] M 70.1, pK²⁵ 10.33 (MeS⁻). Dissolve the salt (10g) in EtOH (10mL) and add Et₂O (100mL). Cool and collect the ppte, wash it with Et₂O and dry it in vacuum. It is a white powder which is very soluble in EtOH and H₂O. [*Bull Soc Chim Fr* 3 2318 1936.]

Sodium methoxide [124-41-4] M 54.0. It behaves the same as sodium ethoxide. It is *hygroscopic* and is hydrolysed by moist air to NaOH and EtOH. Material that has been kept under N₂ should be used. If erratic results are obtained, even with recently purchased NaOMe it should be freshly prepared thus: Clean Na (37g) cut in 1-3g pieces is added in small portions to stirred MeOH (800mL) in a 2L three necked flask equipped with a stirrer and a condenser with a drying tube. After all the Na has dissolved the MeOH is removed by distillation under vacuum and the residual NaOMe is dried by heating at 150° under vacuum and kept under dry N₂ [*Org Synth* 39 51 1959].

Sodium 3-methyl-1-butanesulfonate [5343-41-9] M 174.1. Crystd from 90% MeOH.

Sodium molybdate (2H₂O) [10102-40-6] M 241.9, m 100°(loses 2H₂O), 687°, d 3.28, pK²⁵ 4.08 (for H₂MoO₄). Crystd from hot water (1mL/g) by cooling to 0°.

Sodium monensin [22373-78-0] M 693.8. Recrystd from EtOH-H₂O [Cox et al. *J Am Chem Soc* 107 4297 1985].

Sodium 1-naphthalenesulfonate [130-14-3] M 230.2. Recrystd from water or aqueous acetone [Okadata et al. *J Am Chem Soc* 108 2863 1986].

Sodium 2-naphthalenesulfonate [532-02-5] M 230.2. Crystd from hot 10% aqueous NaOH or water, and dried in a steam oven.

Sodium 2-naphthylamine-5,7-disulfonate [79004-97-0] M 235.4. Crystd from water (charcoal) and dried in a steam oven.

Sodium nitrate [7631-99-4] M 85.0, m 307°, b 380°, d 2.26. Crystd from hot water (0.6mL/g) by cooling to 0°, or from concentrated aqueous solution by addition of MeOH. Dried under vacuum at 140°. After 2 recrystns tech grade had K, Mg, B, Fe Al, and Li at 100, 29, 0.6, 0.4, 0.2 and 0.2 ppm resp.

Sodium nitrite [7632-00-0] M 69.0, m 271°, b 320°, d 2.17. Crystd from hot water (0.7mL/g) by cooling to 0°, or from its own melt. Dried over P₂O₅.

Sodium 1-octanesulfonate [5324-84-5] M 216.2. Recrystd from absolute EtOH.

Sodium oleate [143-19-1] M 304.4, m 233-235°. Crystd from EtOH and dried in an oven at 100°.

Sodium oxalate [62-76-0] M 134.0, m 250-270°(dec), d 2.34. Crystd from hot water (16mL/g) by cooling to 0°. Before use as a volumetric standard, analytical grade quality sodium oxalate should be dried for 2h at 120° and allowed to cool in a desiccator.

Sodium palmitate [408-35-5] M 278.4, m , 270°, 285-201°. Crystd from EtOH, dried in an oven.

Sodium perchlorate (anhydrous) [7601-89-0] M 122.4, m 130°(for monohydrate), d 2.02, pK²⁵ -2.4 to -3.1 (for HClO₄). Because its solubility in water is high (2.1g/mL at 15°) and it has a rather low temperature coefficient of solubility, sodium perchlorate is usually crystd from acetone, MeOH, water-ethanol or dioxane-water (33g dissolved in 36mL of water and 200mL of dioxane). After filtering and crystallising, the solid is dried under vacuum at 140-150° to remove solvent of crystn. Basic impurities can be removed by crystn from hot acetic acid, followed by heating at 150°. If NaClO₄ is ppted from distilled water by adding HClO₄ to the chilled solution, the ppte contains some free acid. **EXPLOSIVE**

Sodium phenol-4-sulfonate (2H₂O) (4-hydroxybenzenesulfonic acid Na salt) [825-90-1] M 232.2. Crystd from hot water (1mL/g) by cooling to 0°, or from MeOH, and dried in vacuum.

Sodium phenoxide [139-02-6] M 116.1, m 61-64°. Washed with Et₂O, then heated under vacuum to 200° to remove any free phenol.

Sodium phenylacetate [114-70-5] M 158.1. Its aqueous solution was evaporated to crystallisation on a steam bath, the crystals being washed with absolute EtOH and dried under vacuum at 80°.

Sodium o-phenylphenolate (4H₂O) [132-27-4] M 264.3. Crystd from acetone and dried under vacuum at room temperature.

Sodium phosphamidate [3076-34-4] M 119.0. Dissolved in water below 10°, and acetic acid added dropwise to pH 4.0 so that the monosodium salt was ppted. The ppte was washed with water and Et₂O, then air dried. Addition of one equivalent of NaOH to the solution gave the sodium salt, the solution being adjusted to pH 6.0 before use [Rose and Heald *Biochem J* 81 339 1961].

Sodium phytate (H₂O) [*myo*-inositolhexakis(H₂PO₄) Na salt] [14306-25-3] M 857.9. Crystd from water.

Sodium piperazine-*N,N'*-bis(2-ethanesulfonate) H₂O (PIPES-Na salt) [76836-02-7] M 364.3. Crystd from water and EtOH.

Sodium polyacrylate (NaPAA) [9003-04-7]. Commercial polyacrylamide was neutralised with an aqueous solution of NaOH and the polymer ppted with acetone. The ppte was redissolved in a small amount of water and freeze-dried. The polymer was repeatedly washed with EtOH and water to remove traces of low

molecular weight material, and finally dried in vacuum at 60° [Vink *J Chem Soc, Faraday Trans 1* **75** 1207 1979]. Also dialysed overnight against distilled water, then freeze-dried.

Sodium poly(α -L-glutamate). It was washed with acetone, dried, dissolved in water and pptd with isopropanol at 5°. Impurities and low molecular weight fractions were removed by dialysis of the aqueous solution for 50h, followed by ultrafiltration through a filter impermeable to polymers of molecular weights greater than 10^4 . The polymer was recovered by freeze-drying. [Mori et al. *J Chem Soc, Faraday Trans 1* **2583** 1978.]

Sodium propionate [137-40-6] **M 96.1, m 287-289°.** Recrystd from H₂O (solubility 10%), and dried by heating at 100° for 4h. Solubility of anhydrous salt in MeOH is 13% at 15° and 13.77% at 68°. It is insoluble in *C₆H₆ and Me₂CO. [*J Chem Soc* 1341 1934.]

Sodium pyrophosphate (10H₂O) [13472-36-1] **M 446.1, d 1.82, pK₁²⁵ 1.52, pK₂²⁵ 2.36, pK₃²⁵ 6.60, pK₄²⁵ 9.25 (for pyrophosphoric acid, H₄P₂O₇).** Crystd from hot H₂O and air dried at room temp.

Sodium selenate [13410-01-0] **M 188.9, pK₁²⁵ ~0, pK₂²⁵ 1.66 (for selenic acid, H₂SeO₄).** Crystd from water.

Sodium selenite [10102-18-8] **M 172.9, m >350°, pK₁²⁵ 2.62, pK₂²⁵ 8.32 (for H₂SeO₃).** Crystd from water.

Sodium silicate solution [1344-09-8] **pK₁²⁵ 9.51, pK₂²⁵ 11.77 (for silicic acid, H₄SiO₄)** Purified by contact filtration with activated charcoal.

Sodium succinate [150-90-3] **M 162.1.** See disodium succinate on p. 421.

Sodium sulfanilate [515-74-2] **M 195.2.** Crystd from water.

Sodium sulfate (10H₂O) [7727-73-3 (10H₂O); 7757-82-6 (anhydr)] **M 322.2, m 32°(dec), 884° (anhydr), d 2.68 (anhydr).** Crystd from water at 30° (1.1mL/g) by cooling to 0°. Sodium sulfate becomes anhydrous at 32°.

Sodium sulfide (9H₂O) [1313-84-4 (9H₂O); 1313-82-2 (anhydr)] **M 240.2, m ~50(loses H₂O), 950(anhydr), d 1.43 (10H₂O), 1.86 (anhydr).** Some purification of the hydrated salt can be achieved by selecting large crystals and removing the surface layer (contaminated with oxidation products) by washing with distilled water. Other metal ions can be removed from Na₂S solutions by passage through a column of Dowex ion-exchange A-1 resin, Na⁺-form. The hydrated salt can be rendered anhydrous by heating in a stream of H₂ or N₂ until water is no longer evolved. (The resulting cake should not be heated to fusion because it is readily oxidised.) Recrystd from distilled water [Anderson and Azowlay *J Chem Soc, Dalton Trans* 469 1986].

Sodium sulfite [7757-83-7] **M 126.0, d 2.63.** Crystd from warm water (0.5mL/g) by cooling to 0°. Purified by repeated crystns from deoxygenated water inside a glove-box, finally drying under vacuum. [Rhee and Dasgupta *J Phys Chem* **89** 1799 1985.]

Sodium R-tartrate (2H₂O) [6106-24-7] **M 230.1, m 120°(loses H₂O), d 1.82.** Crystd from warm dilute aqueous NaOH by cooling.

Sodium taurocholate [145-42-6] **M 555.7.** Purified by recrystn and gel chromatography using Sephadex LH-20.

Sodium tetradecylsulfate [1191-50-0] **M 316.4.** Recrystd from absolute EtOH [Abu Hamdiyyah and Rahman *J Phys Chem* **91** 1531 1987].

Sodium tetrafluoroborate [13755-29-8] M 109.8, d 2.47, pK^{25} -4.9 (for HBF_4). See Sodium fluoroborate on p. 470.

Sodium tetrametaphosphate [13396-41-3] M 429.9, pK_1^{25} 2.60, pK_2^{25} 6.4, pK_3^{25} 8.22, pK_4^{25} 11.4 (tetrametaphosphoric acid, $\text{H}_4\text{P}_4\text{O}_{12}$). Crystd twice from water at room temperature by adding EtOH (300g of $\text{Na}_4\text{P}_4\text{O}_{12}\cdot\text{H}_2\text{O}$, 2L of water, and 1L of EtOH), washed first with 20% EtOH then with 50% EtOH and air dried [Quimby *J Phys Chem* 58 603 1954].

Sodium tetraphenylborate [tetraphenyl boron Na] [143-66-8] M 342.2. Dissolve in dry MeOH and add dry Et_2O . Collect the solid and dry in a vacuum at $80^\circ/2\text{mm}$ for 4h. Also can be extracted (Soxhlet) using CHCl_3 and crystallises from CHCl_3 as snow white needles. It is freely sol in H_2O , Me_2CO but insol in pet ether and Et_2O . An aqueous soln has pH ~ 5 and can be stored for days at 25° or lower, and for 5 days at 45° without deterioration. Its solubility in polar solvents increases with decrease in temp [Justus Liebigs Ann Chem 574 195 1950]. The salt can also be recrystd from acetone-hexane or CHCl_3 , or from Et_2O -cyclohexane (3:2) by warming the soln to ppt the compound. Dried in a vacuum at 80° . Dissolved in Me_2CO and added to an excess of toluene. After a slight milkiness developed on standing, the mixture was filtered. The clear filtrate was evaporated at room temperature to a small bulk and again filtered. The filtrate was then warmed to $50\text{-}60^\circ$, giving clear dissolution of crystals. After standing at this temperature for 10min the mixture was filtered rapidly through a pre-heated Büchner funnel, and the crystals were collected and dried in a vacuum desiccator at room temperature for 3 days [Abraham et al. *J Chem Soc, Faraday Trans 1* 80 489 1984]. If the product gives a turbid aq solution, the turbidity can be removed by treating with freshly prepared alumina gel.

Sodium thioantimonate ($\text{Na}_3\text{SbS}_4\cdot 9\text{H}_2\text{O}$, Schlippe's salt) [13776-84-6] M 481.1, m 87° , b 234° , d 1.81. Crystd from warm water (2mL/g) by cooling to 0° .

Sodium thiocyanate [540-72-7] M 81.1, m 300° , pK^{25} -1.85 (for HSCN). It is recrystd from EtOH or Me_2CO and the mother liquor is removed from the crystals by centrifugation. It is very deliquescent and should be kept in an oven at 130° before use. It can be dried in vacuum at $120^\circ/\text{P}_2\text{O}_5$ [*Trans Faraday Soc* 30 1104 1934]. Its solubility in H_2O is 113% at 10° , 178% at 46° , 225.6% at 101.4° ; in MeOH 35% at 15.8° , 51% at 48° , 53.5% at 52.3° ; in EtOH 18.4% at 18.8° , 24.4% at 70.9° ; and in Me_2CO 6.85% at 18.8° and 21.4% at 56° [*J Chem Soc* 2282 1929].

Sodium thiocyanate has also been recrystd from water, acetonitrile or from MeOH using Et_2O for washing, then dried at 130° , or dried under vacuum at 60° for 2 days. [Strasser et al. *J Am Chem Soc* 107 789 1985; Szezygiel et al. *J Am Chem Soc* 91 1252 1987.] (The latter purification removes material reacting with iodine.) Sodium thiocyanate solns can be freed from traces of iron by repeated batch extractions with Et_2O .

Sodium thioglycolate [367-51-1] M 114.1. Crystd from 60% EtOH (charcoal). Hygroscopic.

Sodium thiosulfate ($5\text{H}_2\text{O}$) [10102-17-7 (hydr); 7772-98-7 (anhydr)] M 248.2(anhydr), m 48(rapid heat), d 1.69, pK_1^{25} 0.6, pK_2^{25} 1.74 (for $\text{H}_2\text{S}_2\text{O}_3$) Crystd from EtOH- H_2O solns or from water (0.3mL/g) below 60° by cooling to 0° , and dried at 35° over P_2O_5 under vacuum.

Sodium *p*-toluenesulfinate [824-79-3] M 178.2, pK^{25} 2.80 (1.99)(for $-\text{SO}_2^-$). Crystd from water (to constant UV spectrum), and dried under vacuum or extracted with hot *benzene, then dissolved in EtOH- H_2O and heated with decolorising charcoal. The solution was filtered and cooled to give crystals of the dihydrate.

Sodium *p*-toluenesulfonate [657-84-1] M 194.2, pK^{25} -1.34 (for $-\text{SO}_3^-$). Dissolved in distilled water, filtered to remove insoluble impurities and evaporated to dryness. Then crystd from MeOH or EtOH, and dried at 110° . Its solubility in EtOH is not high (maximum 2.5%) so that Soxhlet extraction with EtOH may be preferable. Sodium *p*-toluenesulfonate has also been crystd from Et_2O and dried under vacuum at 50° .

Sodium trifluoroacetate [2923-18-4] M 136.0, m $206\text{-}210^\circ(\text{dec})$, pK^{25} 0.52 (for CF_3CO_2^-). A possible contaminant is NaCl. The solid is treated with $\text{CF}_3\text{CO}_2\text{H}$ and evaporated twice. Its solubility in $\text{CF}_3\text{CO}_2\text{H}$ is 13.1% at 29.8° . The residue is crystd from dil EtOH and the solid dried in vacuum at 100° . [*J*

Am Chem Soc 76 4285 1954.] It can be pptd from EtOH by adding dioxane, then crystd several times from hot absolute EtOH. Dried at 120-130°/1mm.

Sodium 2,2',4-trihydroxyazobenzene-5'-sulfonate [3564-26-9] M 295.3. Purified by precipitating the free acid from aqueous solution using concentrated HCl, then washing and extracting with EtOH in a Soxhlet extractor. Evaporation of the EtOH left the purified acid.

Sodium trimetaphosphate (6H₂O) [7785-84-4] M 320.2, m 53°, d 1.79, pK₂^{2.5} 1.64, pK₃^{2.5} 2.07 (for trimetaphosphoric acid, H₃P₃O₉). Pptd from an aq soln at 40° by adding EtOH. Air dried.

Sodium 2,4,6-trimethylbenzenesulfonate [6148-75-0] M 222.1. Crystd twice from MeOH and dried under vacuum.

Sodium trimethylsilanolate (sodium trimethylsilanol) [18027-10-6] M 112.2, m 230°(dec). It is very soluble in Et₂O and *C₆H₆ but moderately soluble in pet ether. It is purified by sublimation at 130-150° in a high vacuum. [IR: *J Am Chem Soc* 75 5615 1953; *J Org Chem* 17 1555 1952.]

Sodium tripolyphosphate [7758-29-4] M 367.9, pK₁^{2.5} ~ 1, pK₂^{2.5} 2.0, pK₃^{2.5} 2.13, pK₄^{2.5} 5.78, pK₅^{2.5} 8.56 (for tripolyphosphoric acid, H₅P₃O₁₀). Purified by repeated pptn from aqueous solution by slow addition of MeOH and air dried. Also a solution of anhydrous sodium tripolyphosphate (840g) in water (3.8L) was filtered, MeOH (1.4L) was added with vigorous stirring to ppte Na₅P₃O₁₀·6H₂O. The ppte was collected on a filter, air dried by suction, then left to dry in air overnight. It was crystd twice more in this way, using a 13% aqueous solution (w/w), and leaching the crystals with 200mL portions of water [Watters, Loughran and Lambert *J Am Chem Soc* 78 4855 1956]. Similarly, EtOH can be added to ppte the salt from a filtered 12-15% aqueous solution, the final solution containing ca 25% EtOH (v/v). Air drying should be at a relative humidity of 40-60%. Heat and vac drying should be avoided. [Quimby *J Phys Chem* 58 603 1954.]

Sodium tungstate (2H₂O) [10213-10-2] M 329.9, m 698°, d 4.18, pK₁^{2.5} 2.20, pK₂^{2.5} 3.70 (for tungstic acid, H₂WO₄). Crystd from hot water (0.8mL/g) by cooling to 0°.

Sodium m-xylenesulfonate [30587-85-0] M 208.2. Dissolved in distilled water, filtered, then evaporated to dryness. Crystd twice from absolute EtOH and dried at 110°.

Sodium p-xylenesulfonate [827-19-0] M 208.2. See sodium 2,5-dimethylbenzenesulfonate on p. 469.

Stannic chloride (tin IV chloride, stannic tetrachloride) [7646-78-8] M 260.5, m -33°, -30.2°, b 114°/760mm, d 2.23, pK^{2.5} 14.15 (for aquo Sn⁴⁺ hydrolysis). Fumes in moist air due to hydrate formation. Fractionate in a ground glass still and store in the absence of air. Possible impurities are SO₂ and HCl [Baudler *Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I p. 729 1963]. It forms a solid pentahydrate [10026-06-9] which smells of HCl and is formed when the anhydrous salt is dissolved in a small vol of H₂O. Also refluxed with clean mercury or P₂O₅ for several hours, then distd under (reduced) N₂ pressure into a receiver containing P₂O₅. Finally redistd. Alternatively, distd from Sn metal under vacuum in an all-glass system and sealed off in large ampoules. Fumes in moist air. SnCl₄ is available commercially as 1M solns in CH₂Cl₂ or hexane. **HARMFUL VAPOURS.**

Stannic iodide (SnI₄) [7790-47-8] M 626.3, m 144°, b 340, d 4.46. Crystd from anhydrous CHCl₃, dried under vacuum and stored in a vacuum desiccator. Sublimes at 180°.

Stannic oxide (SnO₂) [18282-10-5] M 150.7, m 1630°, d 6.95. Refluxed repeatedly with fresh HCl until the acid showed no tinge of yellow. The oxide was then dried at 110°.

Stannous bis-cyclopentadienyl [26078-96-6] M 248.9. Purified by vacuum sublimation. Handled and stored under dry N₂. The related thallium and indium compounds are similarly prepared.

Stannous chloride (anhydrous) [7772-99-8] M 189.6, m 247°, b 606°, d 3.95, pK_1^{25} 1.7, pK_2^{25} 3.7 (for aquo Sn^{2+} hydrolysis). Analytical reagent grade stannous chloride dihydrate is dehydrated by adding slowly to vigorously stirred, redistilled acetic anhydride (120g salt per 100g of anhydride). (In a fume cupboard.) After *ca* an hour, the anhydrous SnCl_2 is filtered on to a sintered-glass or Büchner funnel, washed free from acetic acid with dry Et_2O (2 x 30mL), and dried under vacuum. It is stored in a sealed container. [Stephen *J Chem Soc* 2786 1930].

Strontium acetate [543-94-2] M 205.7, d 2.1, pK^{25} 13.0 (for aquo Sr^{2+} hydrolysis). Crystd from AcOH, then dried under vacuum for 24h at 100°.

Strontium bromide [10476-81-0] M 247.4, m 643°, d 4.22. Crystd from water (0.5mL/g).

Strontium chloride (6H₂O) [1025-70-4] M 266.6, m 61°(rapid heating), 114-150°(loses 5H₂O), 868°(anhydr). Crystd from warm water (0.5mL/g) by cooling to 0°.

Strontium chromate [7789-06-2] M 203.6, d 3.9, pK_1^{25} 0.74, pK_2^{25} 6.49 (for H_2CrO_4). Crystd from water (40mL/g) by cooling.

Strontium hydroxide (8H₂O) [1311-10-0 (8H₂O); 18480-07-4 (anhydr)] M 265.8, m 100°(loses H₂O), d 1.90, m 375(anhydr), d 3.63 (anhydr). Crystd from hot water (2.2mL/g) by cooling to 0°.

Strontium lactate (3H₂O) [29870-99-3] M 319.8, m 120°(loses 3H₂O). Crystd from aq EtOH.

Strontium nitrate [10042-76-9] M 211.6, m 570°, b 645°, d 2.99. Crystd from hot water (0.5mL/g) by cooling to 0°.

Strontium oxalate (H₂O) [814-95-9] M 193.6, m 150°. Crystd from hot water (20mL/g) by cooling.

Strontium salicylate [526-26-1] M 224.7. Crystd from hot water (4mL/g) or EtOH.

Strontium tartrate [868-19-9] M 237.7. Crystd from hot water.

Strontium thiosalicylate (5H₂O) [15123-90-7] M 289.8. Crystd from hot water (2mL/g) by cooling to 0°.

Sulfamic acid [5329-14-6] M 97.1, m 205°(dec), pK^{25} 0.99 ($\text{NH}_2\text{SO}_3\text{H}$). Crystd from water at 70° (300mL per 25g), after filtering, by cooling a little and discarding the first batch of crystals (about 25g) before standing in an ice-salt mixture for 20min. The crystals were filtered by suction, washed with a small quantity of ice water, then twice with cold EtOH and finally with Et_2O . Air dried for 1h, then stored in a desiccator over $\text{Mg}(\text{ClO}_4)_2$ [Butler, Smith and Audrieth *Ind Eng Chem (Anal Ed)* 10 690 1938]. For preparation of primary standard material see *Pure Appl Chem* 25 459 1969.

Sulfamide [7803-58-9] M 96.1, m 91.5°. Crystd from absolute EtOH.

Sulfur [7704-34-9] M 32.1, m between 112.8° and 120°, depending on form. Murphy, Clabaugh and Gilchrist [*J Res Nat Bur Stand* 64A 355 1960] have obtained sulfur of about 99.999% purity by the following procedure: Roll sulfur was melted and filtered through a coarse-porosity glass filter funnel into a 2L round-bottomed Pyrex flask with two necks. Conc H_2SO_4 (300mL) was added to the sulfur (2.5Kg), and the mixture was heated to 150°, stirring continuously for 2h. Over the next 6h, conc HNO_3 was added in about 2mL portions at 10-15min intervals to the heated mixture. It was then allowed to cool to room temperature and the acid was poured off. The sulfur was rinsed several times with distilled water, then remelted, cooled, and rinsed several times with distd water again, this process being repeated four or five times to remove most of the acid entrapped in the sulfur. An air-cooled reflux tube (*ca* 40cm long) was attached to one of the necks of the flask, and a gas delivery tube (the lower end about 1in above the bottom of the flask) was inserted into the other. While the sulfur was boiled under reflux, a stream of helium or N_2 was passed through to remove any

water, HNO_3 or H_2SO_4 , as vapour. After 4h, the sulfur was cooled so that the reflux tube could be replaced by a bent air-cooled condenser. The sulfur was then distilled, rejecting the first and the final 100mL portions, and transferred in 200mL portions to 400mL glass cylinder ampoules (which were placed on their sides during solidification). After adding about 80mL of water, displacing the air with N_2 , and sealing the ampoule was cooled, and the water was titrated with 0.02M NaOH, the process being repeated until the acid content was negligible. Finally, entrapped water was removed by alternate evacuation to 10mm Hg and refilling with N_2 while the sulfur was kept molten. Other purifications include crystn from CS_2 (which is less satisfactory because the sulfur retains appreciable amounts of organic material), *benzene or *benzene/acetone, followed by melting and degassing. Has also been boiled with 1% MgO, then decanted, and dried under vacuum at 40° for 2 days over P_2O_5 . [For purification of S_6 , "recryst. S_8 " and "Bacon-Fanelli sulfur" see Bartlett, Cox and Davis *J Am Chem Soc* **83** 103, 109 1961.]

Sulfur dichloride [10545-99-0] M 103.0, m -78° , b $59^\circ/760\text{mm}(\text{dec})$, d 1.621. Twice distilled in the presence of a small amount of PCl_3 through a 12in Vigreux column, the fraction boiling between $55-61^\circ$ being redistd (in the presence of PCl_3), and the fraction distilling between $58-61^\circ$ retained. (The PCl_3 is added to inhibit the decomposition of SCl_2 into S_2Cl_2 and Cl_2). The SCl_2 must be used as quickly as possible after distn, within 1h at room temperature, The sample contains 4% S_2Cl_2 . On long standing this reaches 16-18%. **HARMFUL VAPOURS.**

Sulfur dioxide [7446-09-5] M 64.1, b -10° . Dried by bubbling through concentrated H_2SO_4 and by passage over P_2O_5 , then passed through a glass-wool plug. Frozen with liquid air and pumped to a high vacuum to remove dissolved gases. **HARMFUL VAPOURS.**

Sulfuric acid [7664-93-9] M 98.1, d 1.83, $\text{pK}_1^{25} \sim -8.3$, $\text{pK}_2^{25} 1.99$. Sulfuric acid, and also 30% fuming H_2SO_4 , can be distilled in an all-Pyrex system, optionally from potassium persulfate. Also purified by fractional crystn of the monohydrate from the liquid. Dehydrates and attacks skin—wash immediately with H_2O .

Sulfur monochloride (sulfur monochloride) [10025-67-9] M 135.0, m -77° ; b 19.1° , $29-30^\circ/12\text{mm}$, $72^\circ/100\text{mm}$, $138^\circ/760\text{mm}$, $d^{20} 1.677$, $n_D^{20} 1.67$. Pungent, irritating golden yellow liquid. When impure its colour is orange to red due to SCl_2 formed. It fumes in moist air and liberates HCl, SO_2 and H_2S in the presence of H_2O . Distil and collect the fraction boiling above 137° at atmospheric pressure. Fractionate this fraction over sulfur at *ca* 12mm using ground glass apparatus (b $29-30^\circ$). Alternatively purify by distn below 60° from a mixture containing sulfur (2%) and activated charcoal (1%), under reduced pressure (e.g. 50mm). It is soluble in EtOH, * C_6H_6 , Et_2O , CS_2 and CCl_4 . Store in a closed container in the dark in a refrigerator. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer) Vol I 371 1963.*] **HARMFUL VAPOURS.**

Sulfur trioxide pyridine complex [26412-87-3] M 159.2, m $155-165^\circ$, 175° . Wash the solid with a little CCl_4 , then H_2O to remove traces of pyridine sulfate, and dry over P_2O_5 [*Chem Ber* **59** 1166 1926; *Synthesis* **59** 1979].

Sulfuryl chloride [7791-25-2] M 135.0, m -54.1° , b $69.3^\circ/760\text{mm}$, $d_4^{20} 1.67$, $n_D^{30} 1.44$. Pungent, irritating colourless liquid. It becomes yellow with time due to decomposition to SO_2 and HCl. Distil and collect fraction boiling below $75^\circ/\text{atm}$ which is mainly SO_2Cl_2 . To remove HSO_3Cl and H_2SO_4 impurities, the distillate is poured into a separating funnel filled with crushed ice and *briefly* shaken. The lower cloudy layer is removed, dried for some time in a desiccator over P_2O_5 and finally fractionated at atmospheric pressure. The middle fraction boils at $69-70^\circ$ and is pure SO_2Cl_2 . It decomposes gradually in H_2O to H_2SO_4 and HCl. Reacts **violently** with EtOH and MeOH and is soluble in * C_6H_6 , toluene Et_2O and acetic acid. [*Handbook of Preparative Inorganic Chemistry (Ed Brauer) Vol I 383 1963; Inorg Synth* **1** 114 1939]. **HARMFUL VAPOURS.**

Tantalum (V) chloride (tantalum pentachloride) [7721-01-9] M 358.2, m 216.2° , $216.5-220^\circ$; b $239^\circ/\text{atm.}$, d 3.68. Purified by sublimation in a stream of Cl_2 . Colourless

[Previous Page](#)

water, HNO_3 or H_2SO_4 , as vapour. After 4h, the sulfur was cooled so that the reflux tube could be replaced by a bent air-cooled condenser. The sulfur was then distilled, rejecting the first and the final 100mL portions, and transferred in 200mL portions to 400mL glass cylinder ampoules (which were placed on their sides during solidification). After adding about 80mL of water, displacing the air with N_2 , and sealing the ampoule was cooled, and the water was titrated with 0.02M NaOH, the process being repeated until the acid content was negligible. Finally, entrapped water was removed by alternate evacuation to 10mm Hg and refilling with N_2 while the sulfur was kept molten. Other purifications include crystn from CS_2 (which is less satisfactory because the sulfur retains appreciable amounts of organic material), *benzene or *benzene/acetone, followed by melting and degassing. Has also been boiled with 1% MgO, then decanted, and dried under vacuum at 40° for 2 days over P_2O_5 . [For purification of S_6 , "recryst. S_8 " and "Bacon-Fanelli sulfur" see Bartlett, Cox and Davis *J Am Chem Soc* **83** 103, 109 1961.]

Sulfur dichloride [10545-99-0] M 103.0, m -78° , b $59^\circ/760\text{mm}(\text{dec})$, d 1.621. Twice distilled in the presence of a small amount of PCl_3 through a 12in Vigreux column, the fraction boiling between $55-61^\circ$ being redistd (in the presence of PCl_3), and the fraction distilling between $58-61^\circ$ retained. (The PCl_3 is added to inhibit the decomposition of SCl_2 into S_2Cl_2 and Cl_2). The SCl_2 must be used as quickly as possible after distn, within 1h at room temperature, The sample contains 4% S_2Cl_2 . On long standing this reaches 16-18%. **HARMFUL VAPOURS.**

Sulfur dioxide [7446-09-5] M 64.1, b -10° . Dried by bubbling through concentrated H_2SO_4 and by passage over P_2O_5 , then passed through a glass-wool plug. Frozen with liquid air and pumped to a high vacuum to remove dissolved gases. **HARMFUL VAPOURS.**

Sulfuric acid [7664-93-9] M 98.1, d 1.83, $\text{pK}_1^{25} \sim -8.3$, $\text{pK}_2^{25} 1.99$. Sulfuric acid, and also 30% fuming H_2SO_4 , can be distilled in an all-Pyrex system, optionally from potassium persulfate. Also purified by fractional crystn of the monohydrate from the liquid. Dehydrates and attacks skin—wash immediately with H_2O .

Sulfur monochloride (sulfur monochloride) [10025-67-9] M 135.0, m -77° ; b 19.1° , $29-30^\circ/12\text{mm}$, $72^\circ/100\text{mm}$, $138^\circ/760\text{mm}$, $d^{20} 1.677$, $n_D^{20} 1.67$. Pungent, irritating golden yellow liquid. When impure its colour is orange to red due to SCl_2 formed. It fumes in moist air and liberates HCl, SO_2 and H_2S in the presence of H_2O . Distil and collect the fraction boiling above 137° at atmospheric pressure. Fractionate this fraction over sulfur at ca 12mm using ground glass apparatus (b $29-30^\circ$). Alternatively purify by distn below 60° from a mixture containing sulfur (2%) and activated charcoal (1%), under reduced pressure (e.g. 50mm). It is soluble in EtOH, * C_6H_6 , Et_2O , CS_2 and CCl_4 . Store in a closed container in the dark in a refrigerator. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I 371 1963.] **HARMFUL VAPOURS.**

Sulfur trioxide pyridine complex [26412-87-3] M 159.2, m $155-165^\circ$, 175° . Wash the solid with a little CCl_4 , then H_2O to remove traces of pyridine sulfate, and dry over P_2O_5 [*Chem Ber* **59** 1166 1926; *Synthesis* 59 1979].

Sulfuryl chloride [7791-25-2] M 135.0, m -54.1° , b $69.3^\circ/760\text{mm}$, $d_4^{20} 1.67$, $n_D^{30} 1.44$. Pungent, irritating colourless liquid. It becomes yellow with time due to decomposition to SO_2 and HCl. Distil and collect fraction boiling below $75^\circ/\text{atm}$ which is mainly SO_2Cl_2 . To remove HSO_3Cl and H_2SO_4 impurities, the distillate is poured into a separating funnel filled with crushed ice and briefly shaken. The lower cloudy layer is removed, dried for some time in a desiccator over P_2O_5 and finally fractionated at atmospheric pressure. The middle fraction boils at $69-70^\circ$ and is pure SO_2Cl_2 . It decomposes gradually in H_2O to H_2SO_4 and HCl. Reacts violently with EtOH and MeOH and is soluble in * C_6H_6 , toluene Et_2O and acetic acid. [*Handbook of Preparative Inorganic Chemistry (Ed Brauer)* Vol I 383 1963; *Inorg Synth* **1** 114 1939]. **HARMFUL VAPOURS.**

Tantalum (V) chloride (tantalum pentachloride) [7721-01-9] M 358.2, m 216.2° , $216.5-220^\circ$; b $239^\circ/\text{atm.}$, d 3.68. Purified by sublimation in a stream of Cl_2 . Colourless

needles when pure (yellow when contaminated with even less than 1% of NbCl_5). Sensitive to H_2O , even in conc HCl it decomposes to tantallic acid. Sol in EtOH . [*J Am Chem Soc* 80 2952 1958; *Handbook of Preparative Inorganic Chemistry* (Ed Brauer) Vol II 1302 1965.]

Tantalium pentaethoxide [6074-84-6] M 406.3, b 147°/0.2mm, 202°/10mm, pK_1^{25} 9.6, pK_2^{25} 13.0 (for tantallic acid). Purified by distillation. It aggregates in $^*\text{C}_6\text{H}_6$, EtOH , MeCN , pyridine and diisopropyl ether. [*J Chem Soc* 726 1955, 5 1956.]

Telluric acid [7803-68-1] M 229.6, pK_1^{25} 7.70, pK_2^{25} 11.04 (H_6TaO_6). Crystd once from nitric acid, then repeatedly from hot water (0.4mL/g).

Tellurium [13494-80-9] M 127.6, m 450°. Purified by zone refining and repeated sublimation to an impurity of less than 1 part in 10^8 (except for surface contamination by TeO_2). [Machol and Westrum *J Am Chem Soc* 80 2950 1958.] Tellurium is volatile at 500°/0.2mm. Also purified by electrode deposition [Mathers and Turner *Trans Amer Electrochem Soc* 54 293 1928].

Tellurium dioxide [7446-07-3] M 159.6, m 733°, d 6.04. Dissolved in 5M NaOH , filtered and ppted by adding 10M HNO_3 to the filtrate until the soln was acid to phenolphthalein. After decanting the supernatant, the ppt was washed five times with distilled water, then dried for 24h at 110° [Horner and Leonhard *J Am Chem Soc* 74 3694 1952].

Terbium oxide [12037-01-3] M 747.7, pK^{25} 8.16 (for Tb^{3+} hydrolysis). Dissolved in acid, ppted as its oxalate and ignited at 650°.

Tetraallyltin (tetraallylstannane) [7393-43-3] M 283.0, b 52°/0.2mm, 69-70°/15mm, d 1.179, n 1.536. Possible contaminants are allyl chloride and allyltin chloride. Check ^1H NMR and IR [Fishwick and Wallbridge *J Organometal Chem* 25 69 1970], and if impure, dissolve in Et_2O and shake with a 5% aq soln of NaF which ppts allyltin fluoride. Separate the Et_2O layer, dry (MgSO_4) and dist at ~ 0.2mm. It decomposes slightly on repeated distn. [O'Brien et al. *Inorg Synth* 13 75 1972; Fishwick et al. *J Chem Soc (A)* 57 1971.]

Tetrabutylammonium borohydride [33725-74-5] M 257.3, m 128-129°. Purified by recrystn from EtOAc followed by careful drying under vacuum at 50-60°. Samples purified in this way showed no signs of loss of active H after storage at room temperature for more than 1 year. Nevertheless samples should be stored at ca 6° in tightly stoppered bottles if kept for long periods. It is soluble in CH_2Cl_2 . [*J Org Chem* 41 690 1976; *Tetrahedron Lett* 3173 1972.]

Tetrabutylammonium chlorochromate [54712-57-1] M 377.9, m 184-185°. Recrystd from EtOAc -hexane. IR ν 920 cm^{-1} in CHCl_3 [*Synthesis* 749 1983]. Powerful oxidant.

Tetrabutylammonium tetrafluoroborate [429-42-5] M 329.3, m 161.8°, 161-163°, pK^{25} -4.9 (for HBF_4). Recryst from H_2O , aq EtOH or from EtOAc by cooling in Dry ice. Also recrystd from ethyl acetate/pentane or dry acetonitrile. Dried at 80° under vacuum. [Detty and Jones *J Am Chem Soc* 109 5666 1987; Hartley and Faulkner *J Am Chem Soc* 107 3436 1985.] Acetate m 118±2° (from BuCl); bromide m 118° (from EtOAc) and nitrate m 120° (from $^*\text{C}_6\text{H}_6$). [*J Am Chem Soc* 69 2472 1947, 77 2024 1955.]

Tetrabutyl orthotitanate monomer (titanium tetrabutoxide) [5593-70-4] M 340.4, b 142°/0.1mm, 134-136°/0.5mm, 160°/0.8mm, 174°/6mm, 189°/13mm, d_4^{35} 0.993, n_D^{25} 1.49. Dissolve in $^*\text{C}_6\text{H}_6$, filter if solid is present, evaporate and vacuum fractionate through a Widmer 24inch column. The ester hydrolyses when exposed to air to give hydrated ortho-titanic acid. Titanium content can be determined thus: weigh a sample (ca 0.25g) into a weighed crucible and cover with 10mL of H_2O and a few drops of conc HNO_3 . Heat (hot plate) carefully till most of the H_2O has evaporated. Cool and add more H_2O (10mL) and conc HNO_3 (2mL) and evaporate carefully (no spillage) to dryness and ignite residue at 600-650°/1h. Weigh the residual TiO_2 . [*J Chem Soc* 2773 1952; *J Org Chem* 14 655 1949.]

Tetrabutyl tin (tin tetrabutyl) [1461-25-2] M 347.2, b 94.5-96°/0.28mm, 145°/11mm, 245-247°/atm, d_4^{20} 1.05, n_D^{24} 1.473. Dissolve in Et₂O, dry over MgSO₄, filter, evaporate and distil under reduced pressure. Although it does not crystallise easily, once the melt has crystallised then it will recrystallise more easily. It is soluble in Et₂O, Me₂CO, EtOAc and EtOH but insoluble in MeOH and H₂O and shows no apparent reaction with H₂O. [J Org Chem 19 74 1954, J Chem Soc 1992 1954.]

Tetraethoxysilane (tetraethyl orthosilicate) [78-10-4] M 208.3, m -77°, b 165-166°/atm, d_4^{20} 0.933, n_D^{25} 1.382. Fractionate through an 80cm Podbielniak type column (see p. 141) with heated jacket and partial take-off head. Slowly decomposed by H₂O, soluble in EtOH. It is *flammable* - irritates the eyes and mucous membranes. [J Am Chem Soc 78 5573 1956, cf J Chem Soc 5020 1952.]

Tetraethylammonium hexafluorophosphate [429-07-2] M 275.2, m >300°, 331°(dec), pK_1^{25} ~0.5, pK_2^{25} 5.12 (for fluorophosphoric acid H₂PO₃F). Dissolve salt (0.8g) in hot H₂O (3.3mL) and cool to crystallise. Yield of prisms is 0.5g. Solubility in H₂O is 8.1g/L at 19° [Chem Ber 63 1067 1930].

Tetraethylammonium tetrafluoroborate [429-06-1] M 217.1, m 235°, 356-367°, pK^{25} -4.9 (for HBF₄). Dissolve in hot MeOH, filter and add Et₂O. It is soluble in ethylene chloride [J Am Chem Soc 69 1016 1947, 77 2025 1955]. See entry on p. 359 in Chapter 4.

Tetraethyl lead [78-00-2] M 323.5, b 200°, 227.7°(dec), d 1.653, n 1.5198. Its more volatile contaminants can be removed by exposure to a low pressure (by continuous pumping) for 1h at 0°. Purified by stirring with an equal volume of H₂SO₄ (d 1.40), keeping the temperature below 30°, repeating this process until the acid layer is colourless. It is then washed with dilute Na₂CO₃ and distilled water, dried with CaCl₂ and fractionally distd at low pressure under H₂ or N₂ [Calingaert Chem Rev 2 43 1926]. **VERY POISONOUS.**

Tetraethylsilane [631-36-7] M 144.3, b 153.8°/760mm, d_4^{30} 0.77, n_D^{30} 1.427. Fractionate through a 3ft vacuum jacketed column packed with 1/4" stainless steel saddles. The material is finally percolated through a 2ft column packed with alumina and maintained in an inert atmosphere. [J Chem Soc 1992 1954; J Am Chem Soc 77 272 1955.]

1.1.3.3-Tetraisopropyldisiloxane [18043-71-5] M 246.5, b 129-130°/6mm, d_4^{30} 0.89, n_D^{30} 1.47. Fractionate under reduced pressure in a N₂ atm. [J Am Chem Soc 69 1500 1947.]

Tetraisopropyl orthotitanate (titanium tetraisopropyl) [546-68-9] M 284.3, m 18.5°; b 80°/2mm, 78°/12mm, 228-229°/755mm. Dissolve in dry *C₆H₆, filter if a solid separates, evap and fractionate. It is hydrolysed by H₂O to give solid Ti₂O(*iso*-OPr)₂ m ca 48°. [J Chem Soc 2027, 1952, 469 1957.]

Tetrakis(diethylamino) titanium [(titanium tetrakis(diethylamide))] [4419-47-0] M 336.4, b 85-90°/0.1mm, 112°/0.1mm, d_4^{30} 0.93, n_D^{30} 1.54. Dissolve in *C₆H₆, filter if a solid separates, evaporate under reduced pressure and distil. Orange liquid which reacts violently with alcohols. [J Chem Soc 3857 1960.]

Tetrakis(hydroxymethyl)phosphonium chloride [124-64-1] M 190.6, m 151°. Crystd from AcOH and dried at 100° in a vacuum. An 80% w/v aqueous solution has d_4^{20} 1.33 [J Am Chem Soc 77 3923 1955].

Tetrakis(triphenylphosphine) palladium [14221-01-3] M 1155.58, m 100-105°(dec). Yellow crystals from EtOH. It is stable in air only for a short time, and prolonged exposure turns its colour to orange. Store in an inert atmosphere below room temp in the dark. [J Chem Soc 1186 1957.]

Tetrakis(triphenylphosphine) platinum [14221-02-4] M 1244.3, m 118°. Recrystd by adding hexane to a cold saturated solution in *C₆H₆. It is soluble in *C₆H₆ and CHCl₃ but insoluble in EtOH and hexane. A less pure product is obtained if crystd by adding hexane to a CHCl₃ soln. Stable in air for several hours and completely stable under N₂. [J Am Chem Soc 2323 1958.]

Tetramethoxysilane (tetramethyl orthosilicate) [681-84-5] M 152.2, m 4.5°, b 122°/760mm. Purification as for tetraethoxysilane. It has a vapour pressure of 2.5mm at 0°. [IR: *J Am Chem Soc* 81 5109 1959.]

Tetramethylammonium borohydride [16883-45-7] M 89.0. Recrystn from H₂O three times yields ca 94% pure compound. Dry in high vacuum at 100° for 3h. The solubility in H₂O is 48% (20°), 61% (40°); and in EtOH 0.5% (25°) and MeCN 0.4% (25°). It decompose slowly in a vacuum at 150°, but rapidly at 250°. The rate of hydrolysis of Me₄N.BH₄ (5.8M) in H₂O at 40° is constant over a period of 100h at 0.04% of original wt/h. The rate decreases to 0.02%/h in the presence of Me₄NOH (5% of the wt of Me₄N.BH₄). [*J Am Chem Soc* 74 2346 1952.]

Tetramethylammonium hexafluorophosphate [558-32-7] M 219.1, m >300°, d₄²⁵ 1.617, pK₁²⁵ ~ 0.5, pK₂²⁵ 5.12 (for fluorophosphoric acid H₂PO₃F). The salt (0.63g) is recrystd from boiling H₂O (76mL), yielding pure (0.45) Me₄N.PF₆ after drying at 100°. It is a good supporting electrolyte. [*Chem Ber* 63 1067 1930.]

Tetramethylammonium perchlorate [2537-36-2] M 123.6, m>300°, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystallise twice from H₂O and dry at 100° in an oven. Insol in most organic solvents. [*J Chem Soc* 1210 1933.]

Tetramethylammonium triacetoxyborohydride [109704-53-2] M 263.1, m 93-98°, 96.5-98°. If impure, wash with freshly distd Et₂O and dry overnight in a vac to give a free flowing powder. Check ¹H NMR, and if still suspect prepare freshly from Me₄NBH₄ and AcOH in *C₆H₆ and store away from moisture [Banus et al. *J Am Chem Soc* 74 2346 1952; Evans and Chipman *Tetrahedron Lett* 27 5939 1986]. It is an IRRITANT and MOISTURE SENSITIVE.

Tetramethylammonium triphenylborofluoride [437-11-6] M 392.2. Crystd from acetone or acetone/ethanol.

2,4,6,8-Tetramethylcyclotetrasiloxane [2370-88-9] M 240.5, m -69°, b 134°/750mm, 134.5-134.9°/755mm, d₄²⁰ 0.99, n_D²⁰ 1.3872. It is purified by repeated redistillation, and fractions with the required ¹H NMR data are collected. [*J Gen Chem USSR (Engl Transl)* 29 262 1959; *J Am Chem Soc* 68 962 1946].

1,1,3,3-Tetramethyldisiloxane [3277-26-7] M 134.3, b 70.5-71°/731mm, 71-72°/atm, d₄³⁰ 0.75, n_D²⁵ 11.367. Possible impurity is 1,1,5,5-tetramethyl-3-trimethylsiloxytrisiloxane b 154-155°/733mm. Fractionate, collect fractions boiling below 80° and refractionate. Purity can be analysed by alkaline hydrolysis and measuring the volume of H₂ liberated followed by gravimetric estimation of silica in the hydrolysate. It is unchanged when stored in glass containers in the absence of moisture for 2-3 weeks. Small amounts of H₂ are liberated on long storage. *Care should be taken when opening a container due to pressure developed.* [*J Am Chem Soc* 79 974 1958; *J Chem Soc* 609 1958; IR: *Z Anorg Chem* 299 78 1959.]

N,N,N'-Tetramethylphosphonic diamide (methylphosphonic bis-dimethylamide) [2511-17-3] M 150.2, b 60.5°/0.6mm, 138°/32mm, 230-230°/atm, d₄³⁰ 1.0157, n_D³⁰ 1.4539. Dissolve in heptane or ethylbenzene shake with 30% aqueous NaOH, stir for 1h, separate the organic layer and fractionate. [*J Org Chem* 21 413 1956]. IR has v 1480, 1460, 1300, 1184, 1065 and 988-970cm⁻¹ [*Can J Chem* 33 1552 1955].

Tetramethylsilane (TMS) [75-76-3] M 88.2, b 26.3°, n 1.359, d 0.639. Distilled from conc H₂SO₄ (after shaking with it) or LiAlH₄, through a 5ft vacuum-jacketted column packed with glass helices into an ice-cooled condenser, then percolated through silica gel to remove traces of halide.

2,4,6,8-Tetramethyl tetravinyl cyclotetrasiloxane [2554-06-5] M 344.7, m -43.5°, b 111-112°/10mm, 145-146°/13mm, 224-224.5°/758mm, d₄³⁰ 0.98, n_D³⁰ 1.434. A 7mL sample was

distilled in a small Vigreux column at atmospheric pressure without polymerisation or decomposition. It is soluble in cyclohexane. [*J Am Chem Soc* 77 1685 1955.]

Tetraphenylarsonium chloride hydrate [507-28-8] M 418.8, m 261-263°. A neutralised aqueous soln was evaporated to dryness. The residue was extracted into absolute EtOH, evaporated to a small volume and ppted by addition of absolute Et₂O. It was again dissolved in a small volume of absolute EtOH or ethyl acetate and reprecipitated with Et₂O. Alternatively purified by adding conc HCl to ppt the chloride dihydrate. Redissolved in water, neutralised with Na₂CO₃ and evaporated to dryness. The residue was extracted with CHCl₃ and finally crystallised from CH₂Cl₂ or EtOH by adding Et₂O. If the aqueous layer is somewhat turbid treat with Celite and filter through filter paper. **POISONOUS.**

Tetraphenylarsonium iodide [7422-32-4] M 510.2. Crystd from MeOH. **POISONOUS.**

Tetraphenylarsonium perchlorate [3084-10-4] M 482.8, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from MeOH. **POISONOUS.**

Tetraphenylboron potassium [3244-41-5] M 358.2. See potassium tetraphenylborate on p. 458.

Tetraphenylphosphonium chloride [2001-45-8] M 374.9, m 273-275°. Crystd from acetone. Dried at 70° under vacuum. Also recrystd from a mixture of 1:1 or 1:2 dichloromethane/pet ether, the solvents having been dried over anhydrous K₂CO₃. The purified salt was dried at room temperature under vacuum for 3 days, and at 170° for a further 3 days. *Extremely hygroscopic.*

Tetraphenylsilane [1048-08-4] M 336.4, m 231-233°. Crystd from *benzene.

Tetraphenyltin [595-90-4] M 427.1, m 224-225°, 226°. Yellow crystals from CHCl₃, pet ether (b 77-120°), xylene or *benzene/cyclohexane, and dried at 75°/20mm. [*J Am Chem Soc* 74 531 1952.]

Tetrapropylammonium perchlorate [15780-02-6] M 285.8, m 238-240°, pK²⁵ -2.4 to -3.1 (for HClO₄). Purified by recrystns from H₂O or MeCN/H₂O (1:4.v/v), and dried in an oven at 60° for several days, or in vacuum over P₂O₅ at 100°. [*Z Phys Chem* 165A 245 1933, 144 281 1929, 140 97 1929.]

Tetra-*n*-propylammonium perruthenate (TPAP, tetrapropyl tetraoxoruthenate) [114615-82-6] M 351.4, m 160°(dec). It is a strong oxidant and may explode on heating. It can be washed with aq *n*-propanol, then H₂O and dried over KOH in a vac. It is stable at room temp but best stored in a refrigerator. It is sol in CH₂Cl₂ and MeCN. [Dengel et al. *Transition Met Chem* 10 98 1985; Griffith et al. *J Chem Soc, Chem Commun* 1625 1987.] § Polymer supported reagent is available commercially.

Tetrasodium pyrene-1,3,6,8-tetrasulfonate [59572-10-0] M 610.5. Recrystd from aqueous acetone [Okahata et al. *J Am Chem Soc* 108 2863 1986].

Thallium (I) acetate [563-68-8] M 263.4, m 126-128°, 127°, pK²⁵ 13.2 (for Tl⁺). Likely impurity is H₂O because the white solid is deliquescent. Dry in a vacuum over P₂O₅ or for several days in a desiccator, and store in a well closed container. 7.5g dissolve in 100g of liquid SO₂ at 0°, and ca 2mol% in AcOH at 25°. **POISONOUS.** [*Trans Faraday Soc* 32 1660 1936; *J Am Chem Soc* 52 516.]

Thallos bromide [7789-40-4] M 284.3, m 460°. Thallos bromide (20g) was refluxed for 2-3h with water (200mL) containing 3mL of 47% HBr. It was then washed until acid-free, heated to 300° for 2-3h and stored in brown bottles. **POISONOUS.**

Thallos carbonate [6533-73-9] M 468.7, m 268-270°. Crystd from hot water (4mL/g) by cooling. **POISONOUS.**

Thallos chlorate [13453-30-0] M 287.8, d 5.05. Crystd from hot water (2mL/g) by cooling. **POISONOUS.**

Thallos chloride [7791-12-0] M 239.8, m 429.9°, d 7.0. Crystd from 1% HCl and washed until acid-free, or crystd from hot water (50mL/g), then dried at 140° and stored in brown bottles. Also purified by subliming in vacuum, followed by treatment with dry HCl gas and filtering while molten. (Soluble in 260 parts of cold water and 70 parts of boiling water). **POISONOUS**.

Thallos hydroxide [12026-06-1] M 221.4, m 139°(dec), pK²⁵ 13.2 (for TI⁺). Crystd from hot water (0.6mL/g) by cooling. **POISONOUS**.

Thallos iodide [7790-30-9] M 331.3, m 441.8°, b 824°, d 7.1. Refluxed for 2-3h with water containing HI, then washed until acid-free, and dried at 120°. Stored in brown bottles. **POISONOUS**.

Thallos nitrate [10102-45-1] M 266.4, m 206°, b 450°(dec), d 5.55. Crystd from warm water (1mL/g) by cooling to 0°. **POISONOUS**.

Thallos perchlorate [13453-40-2] M 303.8, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from hot water (0.6mL/g) by cooling. Dried under vacuum for 12h at 100° (protect from possible **EXPLOSION**).

Thallos sulfate [7446-18-6] M 504.8, m 633°, d 6.77. Crystd from hot water (7mL/g) by cooling, then dried under vacuum over P₂O₅. **POISONOUS**.

Thexyl dimethyl chlorosilane (dimethyl-[2,3-dimethyl-2-butyl] chlorosilane) [67373-56-2] M 178.8, b 55-56°/10mm, 158-159°/720mm, d₄²⁰ 0.970, n_D²⁰ 1.428. Purified by fractional distillation and stored in small aliquots in sealed ampoules. It is very sensitive to moisture and is estimated by dissolving an aliquot in excess of 0.1M NaOH and titrating with 0.1M HCl using methyl red as indicator. [*Helv Chim Acta* 67 2128 1984].

N-(Thexyl dimethylsilyl)dimethylamine (N-[2,3-dimethyl-2-butyl]dimethylsilyl dimethylamine) [81484-86-8] M 187.4, b 156-160°/720mm. Dissolve in hexane, filter, evaporate and distil. Colourless oil extremely sensitive to humidity. It is best to store small quantities in sealed ampoules after distillation. For estimation of purity crush an ampoule in excess 0.1N HCl and titrate the excess acid with 0.1M NaOH using methyl red as indicator. [*Helv Chim Acta* 67 2128 1984.]

Thionyl chloride [7719-09-7] M 119.0, b 77°, d 1.636. Crude SOCl₂ can be freed from sulfuryl chloride, sulfur monochloride and sulfur dichloride by refluxing with sulfur and then fractionally distilling twice. [The SOCl₂ is converted to SO₂ and sulfur chlorides. The S₂Cl₂ (b 135.6°) is left in the residue, whereas SCl₂ (b 59°) passes over in the forerun]. The usual purification is to distil from quinoline (50g SOCl₂ to 10g quinoline) to remove acid impurities, followed by distillation from boiled linseed oil (50g SOCl₂ to 20g of oil). Precautions must be taken to exclude moisture.

Thionyl chloride for use in organic syntheses can be prepared by distillation of technical SOCl₂ in the presence of diterpene (12g/250mL SOCl₂), avoiding overheating. Further purification is achieved by redistillation from linseed oil (1-2%) [*Rigby Chem Ind (London)* 1508 1969]. Gas chromatographically pure material is obtained by distillation from 10% (w/w) triphenyl phosphite [*Friedman and Wetter J Chem Soc (A)* 36 1967; *Larsen et al. J Am Chem Soc* 108 6950 1986]. **Harmful vapours**.

Thorium chloride [10026-08-1] M 373.8, pK₁²⁵ 10.45, pK₂²⁵ 10.62, pK₃²⁵ 10.80, pK₄²⁵ 11.64 (for aquo Th⁴⁺). Freed from anionic impurities by passing a 2M soln of ThCl₄ in 3M HCl through a Dowex-1 anion-resin column. The eluate was partially evaporated to give crystals which were filtered off, washed with Et₂O and stored in a desiccator over H₂SO₄ to dry. Alternatively, a saturated solution of ThCl₄ in 6M HCl was filtered through quartz wool and extracted twice with ethyl, or isopropyl, ether (to remove iron), then evaporated to a small volume on a hot plate. (Excess silica ppted, and was filtered off. The filtrate was cooled to 0° and saturated with dry HCl gas.) It was shaken with an equal volume of Et₂O, agitating with HCl gas, until the mixture becomes homogeneous. On standing, ThCl₄·8H₂O ppted and was filtered off, washed with Et₂O and dried [*Kremer J Am Chem Soc* 64 1009 1942].

Thorium sulfate (4H₂O) [10381-37-0] M 496.2, m 42°(loses H₂O), d 2.8. Crystd from water.

Tin (powder) [7440-31-5] M 118.7. The powder was added to about twice its weight of 10% aqueous NaOH and shaken vigorously for 10min. (This removed oxide film and stearic acid or similar material sometimes added for pulverisation.) It was then filtered, washed with water until the washings were no longer alkaline to litmus, rinsed with MeOH and air dried. [Sisido, Takeda and Kinugama *J Am Chem Soc* 83 538 1961.]

Tin tetramethyl [594-27-4] M 178.8, m 16.5°, b 78.3°/740mm. It is purified by fractionation using a Todd column of 35-40 plates at atmospheric pressure (p. 177). The purity of the fractions can be followed by IR [*J Am Chem Soc* 77 6486 1955]. It readily dissolves stopcock silicone greases which give bands in the 8-10 μ region. [*J Am Chem Soc* 76 1169 1954.]

Titanium tetrabromide [7789-68-6] M 367.5, m 28.3°, b 233.5°, d 3.3. Purified by distn. Distillate forms light orange hygroscopic crystals. Store in the dark under N₂ preferably in sealed brown glass ampules. [Olsen and Ryan *J Am Chem Soc* 54 2215 1932.]

Titanium tetrachloride [7550-45-0] M 189.7, b 136.4°, 154°, d 1.730, pK₁²⁵ 0.3, pK₂²⁵ 1.8, pK₃²⁵ 2.1, pK₄²⁵ 2.4 (for aquo Ti⁴⁺ hydrolysis). Refluxed with mercury or a small amount of pure copper turnings to remove the last traces of light colour [due to FeCl₃ and VCl₄], then distilled under N₂ in an all-glass system, taking precautions to exclude moisture. Clabaugh, Leslie and Gilchrist [*J Res Nat Bur Stand* 55 261 1955] removed organic material by adding aluminium chloride hexahydrate as a slurry with an equal amount of water (the slurry being *ca* one-fiftieth the weight of TiCl₄), refluxing for 2-6h while bubbling in chlorine, which was subsequently removed by passing a stream of clean dry air. The TiCl₄ was then distilled, refluxed with copper and again distilled, taking precautions to exclude moisture. Volatile impurities were then removed using a technique of freezing, pumping and melting. [Baxter and Fertig *J Am Chem Soc* 45 1228 1923; Baxter and Butler *J Am Chem Soc* 48 3117 1926.] **HARMFUL VAPOURS.**

Titanium trichloride [7705-07-9] M 154.3, m >500°, pK₁²⁵ 2.55 (for hydrolysis of Ti³⁺ to TiOH²⁺). Brown purple powder that is very reactive with H₂O and pyrophoric when dry. It should be manipulated in a dry box. It is soluble in CH₂Cl₂ and tetrahydrofuran and is used as a M solution in these solvents in the ratio of 2:1, and stored under N₂. It is a powerful reducing agent. [*Inorg Synth* 6 52 1960; *Synthesis* 833 1989.]

Titanocene dichloride [1271-19-8] M 248.9, m 260-280°(dec), 289.2°, 298-291°, d 1.60. Bright red crystals from toluene or xylene-CHCl₃ (1:1) and sublimes at 190°/2mm. It is moderately soluble in EtOH and insoluble in Et₂O, *C₆H₆, CS₂, CCl₄, pet ether and H₂O. [IR: *J Am Chem Soc* 76 4281 1954; NMR and X-ray: *Can J Chem* 51 2609 1973, 53 1622 1975.]

Titanyl sulfate (TiOSO₄.2H₂O) [13825-74-6] M 160.0. Dissolved in water, filtered and crystd three times from boiling 45% H₂SO₄, washing with EtOH to remove excess acid, then with Et₂O. Air dried for several hours, then oven dried at 105-110°. [Hixson and Fredrickson *Ind Eng Chem* 37 678 1945.]

Tribenzyl chlorosilane [18740-59-5] M 336.9, m 139-142°, 141-142°, b 300-360°/100mm. It is recrystd three times from pet ether; slender colourless needles, m 141°, sparingly soluble in pet ether and soluble in Et₂O. Does not fume in air but is decomposed by H₂O to give *tribenzyl silanol* m 106° (from pet ether). [*J Chem Soc* 93 439 1908; *J Org Chem* 15 556 1950.]

Tribenzyl phosphine [76650-89-7] M 304.4, m 96-101°, b 203-210°/0.5mm, pK_{Est} ~8.8. Dissolve in Et₂O, dry over Na₂SO₄, evap and distil in an inert atmosphere. Distillate solidifies on cooling and is sublimed at 140°/0.001mm. This has m 92-95°(evacuated capillary). When air is bubbled through an Et₂O solution, it is oxidised to *tribenzyl phosphine oxide*, m 209-212° (evacuated capillary) (from Me₂CO). [*J Chem Soc* 2835 1959.]

Tri-*n*-butyl borate [688-74-4] M 230.2, b 232.4°, n 1.4092, d 0.857. The chief impurities are *n*-butyl alcohol and boric acid (from hydrolysis). It must be handled in a dry-box, and can readily be purified by fractional distillation, under reduced pressure.

Tri-*n*-butyl chlorosilane [995-45-9] M 234.9, b 93-94°/4.5mm, 134-139°/16mm, 250-252°/atm, 142-144°/29mm, d_4^{20} 0.88, n_D^{20} 1.447. Fractionate and store in small aliquots in sealed ampoules. [*J Am Chem Soc* 74 1361 1952; *J Org Chem* 24 219 1959.]

Tri-*n*-butyl phosphate (butyl phosphate) [126-73-8] M 266.3, m -80°, b 47°/0.45mm, 98°/0.1mm, 121-124°/3mm, 136-137°/5.5mm, 166-167°/17mm, 177-178°/27mm, 289°/760atm (some dec), d_4^{20} 0.980, n_D^{20} 1.44249. The main contaminants in commercial samples are organic pyrophosphates, mono- and di- butyl phosphates and butanol. It is purified by washing successively with 0.2M HNO₃ (three times), 0.2M NaOH (three times) and water (three times), then fractionally distilled under vacuum. [Yoshida *J Inorg Nucl Chem* 24 1257 1962.] It has also been purified *via* its uranyl nitrate addition compound, obtained by saturating the crude phosphate with uranyl nitrate. This compound was crystd three times with *n*-hexane by cooling to -40°, and then decomposed by washing with Na₂CO₃ and water. Hexane was removed by steam distn and the water was then evaporated under reduced pressure and the residue was distilled under reduced pressure. [Siddall and Dukes *J Am Chem Soc* 81 790 1959.]

Alternatively wash with water, then with 1% NaOH or 5% Na₂CO₃ for several hours, then finally with water. Dry under reduced pressure and fractionate carefully under vacuum. Stable colourless oil, sparingly soluble in H₂O (1mL dissolves in 165mL of H₂O), but freely miscible in organic solvents. [*J Am Chem Soc* 74 4953 1952, 80 5441 1958; ³¹P NMR: *J Am Chem Soc* 78 5715 1956; *J Chem Soc* 1488 1957.]

Tri-*n*-butyl phosphine [998-40-3] M 202.3, b 109-110°/10mm, 115-116°/12mm, 149.5°/50mm, 240.4-242.2°/atm, d_4^{20} 0.822, n_D^{20} 1.4463, pK_{Est}~7.6. Fractionally distilled under reduced pressure in an inert atm (N₂) through an 8" gauze packed column (b 110-111°/10mm) and redistilled in a vacuum and sealed in thin glass ampoules. It is easily oxidised by air to *tri-*n*-butylphosphine oxide*, b 293-296°/745mm. It has a characteristic odour, it is soluble in EtOH, Et₂O, and *C₆H₆ but insoluble in H₂O and is less easily oxidised by air than the lower molecular weight phosphines. It forms complexes, e.g. with CS₂ (1:1) m 65.5° (from EtOH). [*J Chem Soc* 33 1929, 1401 1956.]

Tri-*n*-butyl phosphite [102-85-2] M 250.3, b 114-115°/5mm, 122°/12mm, 130°/17mm, 137°/26mm, d_4^{20} 0.926, n_D^{20} 1.4924. Fractionate with an efficient column. Stable in air but is slowly hydrolysed by H₂O. [*J Chem Soc* 1464 1940, 1488 1957; *J Am Chem Soc* 80 2358, 2999 1958.]

Tri-*n*-butyl tin chloride [1461-22-9] M 325.5, b 98-100°/0.4mm, 140-152°/10mm, 172°/25mm, d_4^{20} 1.21, n_D^{20} 1.492. Fractionate in an inert atmosphere, and seal in small aliquots in glass ampoules. Sensitive to moisture. [*J Chem Soc* 1446 1947; *J Appl Chem* 6 93 1956.]

Tributyl tin hydride [688-73-3] M 291.1, b 76°/0.7mm, 81°/0.9mm, d_4^{20} 1.098, n_D^{20} 1.473. Dissolve in Et₂O, add quinol (500mg for 300mL), dry over Na₂SO₄, filter, evaporate and distil under dry N₂. It is a clear liquid if dry and decompose very slowly. In the presence of H₂O traces of tributyl tin hydroxide are formed in a few days. Store in sealed glass ampoules in small aliquots. It is estimated by reaction with aq NaOH when H₂ is liberated. CARE: stored samples may be under pressure due to liberated H₂. [*J Appl Chem* 7 366 1957.]

B-Trichloroborazine [933-18-6] M 183.1, m 87°, b 88-92°/21mm. Purified by distillation from mineral oil.

Trichloromethyl trimethylsilane (trimethylsilyl trichloromethane) [5936-98-1] M 191.6, m 130-132°, b 146-156°/749mm. It distils at atmospheric pressure without decomposition and readily sublimates at 70°/10mm. It has one peak in the ¹H NMR spectrum (CH₂Cl₂) δ: 0.38. [*Synthesis* 626 1980.]

Tricyclohexylphosphine [2622-14-2] M 280.4, m 82-83°, pK_{Est}~9.5. Recrystd from EtOH [Boert et al. *J Am Chem Soc* 109 7781 1987].

Triethoxysilane [998-30-1] M 164.3, m -170°, b 131.2-131.8°/atm, 131.5°/760mm, d_4^{20} 0.98753, n_D^{20} 1.4377. Fractionated using a column packed with glass helices of *ca* 15 theoretical plates in

an inert atmosphere. Store in aliquots in sealed ampoules because it is sensitive to moisture. [*J Am Chem Soc* 72 1377, 2032 1950; *J Org Chem* 13 280 1948.]

Triethylborane [97-94-9] M 146.0, b 118.6°, n 1.378, d 0.678. Distilled at 56-57°/220mm.

Triethyl borate [150-46-9] M 146.0, b 118°, n 1.378, d 0.864. Dried with sodium, then distilled.

Triethyl phosphate [78-40-0] M 182.2, b 40-42°/0.25-0.3mm, 98-98.5°/8-10mm, 90°/10mm, 130°/55mm, 204°/680mm, 215-216°/760mm, d_4^{25} 1.608, n_D^{20} 1.4053. Dried by refluxing with solid BaO and fractionally distilled under reduced pressure. It is kept with Na and distilled. Stored in the receiver protected from light and moisture. Alternatively it is dried over Na₂SO₄ and distilled under reduced pressure. The middle fraction is stirred for several weeks over anhydrous Na₂SO₄ and again fractionated under reduced pressure until the specific conductance reached a constant low value of κ^{25} 1.19 x 10⁸, κ^{40} 1.68 x 10⁸, and κ^{55} 2.89 x 10⁸ ohm⁻¹ cm⁻¹. It has also been fractionated carefully under reduced pressure through a glass helices packed column. It is soluble in EtOH, Et₂O and H₂O (dec). [*J Am Chem Soc* 77 4767 1955, 78 6413, 3557 (P NMR) 1956; *J Chem Soc* 3582 1959, IR: *J Chem Soc* 475 1952 and *Can J Chem* 36 820 1958; Kosolapoff *Organophosphorus Compounds*, Wiley p. 258 1950.]

Triethylphosphine [554-70-1] M 118.2, b 100°/7mm, 127-128°/744mm, d_4^{15} 0.812, n_D^{18} 1.457, pK²⁵ 8.69 (also available as a 1.0M soln in THF). All operations should be carried out in an efficient fume cupboard because it is flammable, toxic and has a foul odour. Purified by fractional distn at atm pressure in a stream of dry N₂, as it is oxidised by air to the oxide. In 300% excess of CS₂ it forms Et₃PCS₂ (m 118-120° cryst from MeOH) which decomposes in CCl₄ to give Et₃PS as a white solid m 94° when recryst from EtOH. [Sorettas and Isbell *J Org Chem* 27 273 1962; *J Am Chem Soc* 82 5791 1960; pK: Henderson and Streuli *J Am Chem Soc* 82 5791 1960; see also trimethylphosphine.] Store in a sealed vial under N₂.

Alternatively, dissolve in Et₂O and shake with a solution of AgI and KI to form the insoluble complex. Filter off the complex, dry over P₂O₅ and the Et₃P is regenerated by heating the complex in a tube attached to a vacuum system. [*J Chem Soc* 530 1953, 1828 1937; *J Org Chem* 27 2573 1962; Kosolapoff *Organophosphorus Compounds*, Wiley p. 31 1950.]

Triethyl phosphite [122-52-1] M 166.2, b 48-49°/11mm, b 52°/12mm, 57.5°/19mm, 157.9°/757mm, d_4^{20} 0.9687, n_D^{20} 1.4135. Treat with Na (to remove water and any dialkyl phosphonate), then decant and distil under reduced pressure, with protection against moisture or distil in vacuum through an efficient Vigreux column or a column packed with Penn State 0.16 x 0.16 in protruded nickel packing and a variable volume take-off head. [*Org Synth Coll Vol IV* 955 1963; *J Am Chem Soc* 78 5817 1956, 80 2999 1958; Kosolapoff *Organophosphorus Compounds*, Wiley p. 203 1950.]

Triethyl phosphonoacetate (triethyl carboxymethyl phosphonate) [867-13-0] M 224.2, b 83-84°/0.5mm, 103°/1.2mm, 143-144°/11mm, 260-262°/atm, d_4^{20} 1.1215, n_D^{20} 1.4310. Purified by fractional distn, preferably *in vacuo*. ³¹P NMR has P resonance at 19.5 relative to orthophosphate. [Kosolapoff and Powell *J Am Chem Soc* 68 1103 1946; 72 4198 1950; Speziale and Freeman *J Org Chem* 23 1586 1958.]

Triethyl phosphonoforate [1474-78-8] M 210.2, b 70-72°/0.1mm, 122.5-123°/8mm, 130-131°/10mm, 138.2°/12.5mm, d_4^{20} 1.22, n_D^{20} 1.423. Dissolve in Et₂O, shake with H₂O (to remove any trace of NaCl impurity), dry (Na₂SO₄), evaporate and distil using an efficient fractionating column. [*Chem Ber* 57 1035 1924.]

Triethyl 2-phosphonopropionate [3699-66-9] M 238.2, b 76-77°/0.2mm, 137-138.5°/17mm, d_0^{20} 1.096, n_D^{20} 1.432. Purified by fractional distillation with high reflux ratio, preferably using a spinning band column. [*J Am Chem Soc* 4198 1950.]

Triethylsilane [617-86-7] **M 116.3, b 105-107°, 107-108°, d 0.734, n 1.414.** Refluxed over molecular sieves, then distilled. It was passed through neutral alumina before use [Randolph and Wrighton *J Am Chem Soc* **108** 3366 1986].

Triethylsilyl-1,4-pentadiene (1,4-pentadien-3-yloxy-trimethylsilane) [62418-65-9] **M 198.4, b 72-74°/12mm, d_4^{20} 0.842, n_D^{20} 1.439.** Dissolve in pentane, wash with H₂O, dry (Na₂SO₄), evaporate, and distil under vacuum. R_F values on Kieselgel 60 are 0.15 (pentane) and 0.60 (*C₆H₆). [IR, NMR, MS: *Helv Chim Acta* **64** 2002 1981.]

Triethyltin hydroxide [994-32-1] **M 222.9.** Treated with HCl, followed by KOH, and filtered to remove diethyltin oxide [Prince *J Chem Soc* 1783 1959].

Tri-*n*-hexylborane [1188-92-7] **M 265.3.** Treated with hex-1-ene and 10% anhydrous Et₂O for 6h at gentle reflux under N₂, then vacuum distilled through an 18in glass helices-packed column under N₂ taking the fraction **b 130°/2.1mm to 137°/1.5mm.** The distillate still contained some di-*n*-hexylborane [Mirviss *J Am Chem Soc* **83** 3051 1961].

Triiron dodecacarbonyl [17685-52-8] **M 503.7, m 140°(dec).** It usually contains 10% by weight of MeOH as stabiliser. This can be removed by keeping in a vacuum at 0.5mm for at least 5h. It can be sublimed slowly at high vacuum and is soluble in organic solvents. [*J Org Chem* **37** 930 1972, *J Chem Soc* 4632 1960; *Inorg Synth* **7** 193 1963.]

Triisoamyl phosphate [919-62-0] **M 308.4, b 143°/3mm.** Purified by repeated crystallisation, from hexane, of its addition compound with uranyl nitrate. (see *tributyl phosphate*.) [Siddall *J Am Chem Soc* **81** 4176 1959].

Triisobutyl phosphate [126-71-6] **M 266.3, b 119-129°/8-12mm, 192°/760mm, d 0.962, n 1.421.** Purified by repeated crystallisation, from hexane, of its addition compound with uranyl nitrate. (see *tributyl phosphate*.) [Siddall *J Am Chem Soc* **81** 4176 1959.]

Triisooctyl thiophosphate [30108-39-5] **M 450.6.** Purified by passage of its solution in CCl₄ through a column of activated alumina.

Triisopropyl phosphite [116-17-6] **M 208.2, b 58-59°/7mm, n_D^{25} 1.4082.** Distilled from sodium, under vacuum, through a column with glass helices. (This removes any dialkyl phosphonate).

Trimesitylphosphine [23897-15-6] **M 388.5, m 205-206°, pK_{Est} ~8.0.** Recrystd from EtOH [Boert et al. *J Am Chem Soc* **109** 7781 1987].

Trimethylalyl phosphate [14019-81-9] **M 260.3, b 134.5-140°/5mm, n_D^{25} 1.4454.** Purified as for triisoamyl phosphate.

Trimethoxysilane [2487-90-3] **M 122.2, m -114.8°, 81.1°/760mm, 84°/atm, d_4^{20} 0.957, n_D^{20} 1.359.** Likely impurities are Si(OMe)₄ and H₂Si(OMe)₂. Efficient fractionation is essential for removing these impurities [IR: *J Am Chem Soc* **81** 5109 1959].

Trimethyl borate (methylborate, trimethoxyboron) [121-43-7] **M 103.9, b 67-68°/742mm, d_4^{20} 0.928, n_D^{20} 1.3610.** Carefully fractionated through a gauze-packed column. Redistil and collect in weighed glass vials and seal. Keep away from moisture. It undergoes alkyl exchange with alcohols and forms azeotropes, e.g. with MeOH the azeotrope consists of 70% (MeO)₃B and 30% MeOH with **b 52-54°/atm, d 0.87.** [*J Chem Soc* 2288 1952; *Chem Ind (London)* 53 1952; *J Am Chem Soc* **75** 213 1953.] Also dried with Na, then distilled.

Trimethyl boroxine [823-96-1] **M 125.5, b 80°/742mm, 79.3°/755mm, d_4^{20} 0.902.** Possible impurity is methylboronic acid. If present then add a few drops of conc H₂SO₄ and distil immediately, then

fractionate through an efficient column. [*J Am Chem Soc* **79** 5179 1957; IR: *Z Anorg Chem* **272** 303 1953.]

Trimethyl chlorosilane (chlorotrimethylsilane) [75-77-4] **M 108.6, b 56-57°/atm, 58°/760mm, d 0.86, n 1.388.** Likely impurities are other chlorinated methylsilanes, and tetrachlorosilane (**b 57.6°**), some of which can form azeotropes. To avoid the latter very efficient fractional distillation is required. It has been fractionated through a 12 plate glass helices packed column with only the heart-cut material used. It has also been fractionated through a 90cm, 19mm diameter Stedman column (see p. 441). Also purified by redistilling from CaH_2 before use. [*J Am Chem Soc* **70**, 4254, 4258 1948; *J Org Chem* **23** 50 1958.] **FLAMMABLE and CORROSIVE.**

Trimethyloxonium tetrafluoroborate [420-37-1] **M 147.9, m 141-143°(sinters, open capillary), 179.6-180.0°(dec), 210-220°(dec).** The salt must be a white crystalline solid **m ~ 179.6-180.0°** (dec, sealed tube). Under a N_2 atmosphere (e.g. Dry Box), wash twice with CH_2Cl_2 then twice with Na-dried Et_2O , and dry by passing dry N_2 over the salt until free from Et_2O [Curphey *Org Synth Coll Vol VI* 1019 1988]. The oxonium salt purified in this way can be handled in air for short periods. The sample kept in a desiccator (Drierite) for 1 month at -20° had the same **m**, and samples stored in this way for >1 year are satisfactory for alkylations. $^1\text{H NMR}$ in liq SO_2 in a sealed tube had a single peak at δ 4.54 (impurities have δ at 3.39). [Meerwein *Org Synth Coll Vol V* 1096 1973.] If the sample looks good, dry in a vac desiccator for 2h ($25^\circ/1\text{mm}$) and stored under N_2 at -20° . Melting point depends on heating rate.

Trimethylphenylsilane (phenyltrimethylsilane) [768-32-1] **M 150.3, b 67.3°/20mm, 170.6°/738mm, d_4^{25} 0.8646, n 1.491.** Fractionally distd at atm or reduced pressure (Podbielniak column, p.141) and estimated by GC with a column packed with Silicone Fluid No 710 on Chromosorb P support. [Gilman et al. *J Org Chem* **18** 1743 1953; Maienthal et al. *J Am Chem Soc* **76** 6392 1954; House and Respass *J Organomet Chem* **4** 95 1965; *J Am Chem Soc* **71** 2923 1949, **73** 4770 1951, **75** 2821 1953]

Trimethyl phosphate [512-56-1] **M 140.1, b 77°/12mm, 94°/22mm, 110°/60mm, 197.2°/atm, d_4^{20} 1.0213, n_D^{20} 1.3961.** Purified by fractionation through and efficient column at high reflux ratio. It is quite soluble in H_2O , solubility is 1:1 at 25° . [*J Am Chem Soc* **74** 2923 1952; IR: *J Chem Soc* 847 1952; *Can J Chem* **36** 820 1958; Kosolapoff *Organophosphorus Compounds*, Wiley p. 258 1950.]

Trimethylphosphine [594-02-2] **M 76.1, m -86° , b 38-39°/atm, pK^{25} 8.65, (also available as a 1.0M soln in THF or toluene).** All operations should be carried out in an efficient fume cupboard because it is flammable, toxic and has a foul odor. Distd at atm pressure in a stream of dry N_2 (apparatus should be held together with springs to avoid loss of gas from increased pressure in the system) and the distillate run into a soln of AgI in aq KI whereby the complex $[\text{Me}_3\text{PAgI}]_4$ separates steadily. Filter off the complex, wash with satd aq KI soln, then H_2O and dry in a vac desiccator over P_2O_5 . The dry complex is heated in a flask (in a stream of dry N_2) in an oil bath at 140° , when pure Me_3P distils off (bath temp can be raised up to 260°). The vapour pressure of Me_3P at 20° is 466mm and the **b** is 37.8° [Thomas and Eriks *Inorg Synth* **9** 59 1967]. Alternatively, freshly distilled Me_3P (6g) is shaken with a solution of AgI (13.2g, 1.1mol) in saturated aqueous KI solution (50mL) for 2h. A white solid, not wetted with H_2O , separates rapidly. It is collected, washed with the KI solution, H_2O , and dried [*J Chem Soc* 1829 1937]. The silver complex is stable if kept dry in the dark in which state it can be kept indefinitely. Me_3P can be generated from the complex when required. Store under N_2 in a sealed container. It has been distd in a vacuum line at -78° *in vacuo* and condensed at -96° [IR and NMR: Crosbie and Sheldrick *J Inorg Nucl Chem* **31** 3684 1969]. The pK^{22} by NMR was 8.80 [Silver and Lutz *J Am Chem Soc* **83** 786 1961; pK 8.65: Henderson and Strueuli *J Am Chem Soc* **82** 5791 1960].

$[\text{Me}_2\text{PAgI}]_4$ [12389-34-3] complex is a flammable solid which has **m 140-142°**. It is decomposed by heating gently in one arm of an inverted U tube. The other arm is kept in a freezing mixture. The complex dissociates and pure Me_3P collects in the cold arm and is used at once. It should not be allowed to come in contact with air [*J Chem Soc* 708 1938]. The CS_2 complex has **m 119°** (cryst from 95% EtOH) and decomposes in CCl_4 to give Me_3PS **m 154°** (from EtOH) [Soretas and Isbell *J Org Chem* **27** 273 1962].

Triphenylphosphine hydrochloride is unstable and volatilises at 75°/0.4mm (120°/14mm). [*J Am Chem Soc* 67 503 1945; IR: *Trans Faraday Soc* 40 41 1944; Kosolapoff *Organophosphorus Compounds*, Wiley p. 31 1950.]

Trimethyl phosphite [121-45-9] M 124.1, b 22°/23mm, 86-86.5°/351mm, 111-112°/760mm, 111°/atm, d_4^{20} 1.0495, n_D^{20} 1.408. Treated with Na (to remove water and any dialkyl phosphonate), then decanted and distilled with protection against moisture. It has also been treated with sodium wire for 24h, then distilled in an inert atmosphere onto activated molecular sieves [Connor et al. *J Chem Soc, Dalton Trans* 511 1986]. It has also been fractionally distilled using a spinning band column at high reflux ratio. It is a colourless liquid which is slowly hydrolysed by H₂O. [*J Am Chem Soc* 80 2999 1958; IR: *J Chem Soc* 255 1950, P NMR: *J Am Chem Soc* 79 2719 1957; Kosolapoff *Organophosphorus Compounds*, Wiley p. 203 1950.]

Trimethylsilyl acetamide [13435-12-6] M 131.3, m 38-43°, 52-54°, b 84°/13mm, 185-186°/atm. Repeated distillation in an inert atmosphere, all operations to be performed under anhydrous atmosphere. In the presence of moisture trimethylsilanol (b 31-34°/26mm) is formed and is a likely impurity (check by NMR). [*Chem Ber* 96 1473 1963.]

Trimethylsilyl acetonitrile (TMSAN) [18293-53-3] M 113.2, b 49-51°/10mm, 65-70°/20mm, d_4^{20} 0.8729, n_D^{20} 1.4420. Check if NMR and IR spectra are correct, if not dissolve in *C₆H₆ (10vols), wash with buffer (AcOH-AcONa pH ca 7) several times, dry (CaCl₂), evaporate and distil. IR: ν (CCl₄) 2215 (CN) cm⁻¹; NMR δ (CCl₄): 0.23 (s, 9H, SiMe₃), and 1.53 (s, 2H, CH₂CN). [*J Chem Soc Perkin Trans 1* 26 1979.]

Trimethylsilyl azide [4648-54-8] M 115.2, b 92-95°/atm, 95-99°/atm, d_4^{20} 0.878, n_D^{20} 1.441. Distil through a Vigreux column in a N₂ atmosphere maintaining the oil bath temperature thermostated at 135-140°. Check the purity by ¹H NMR [CHCl₃, δ : single peak at 13cps from Me₄Si. Likely impurities are siloxane hydrolysis products. The azide is thermally stable even at 200° when it decomposes slowly without explosive violence. All the same it is advisable to carry out the distillation behind a thick safety screen in a fumehood because unforeseen **EXPLOSIVE** azides may be formed on long standing. [Birkofer and Wagner *Org Synth Coll Vol VI* 1030 1988.]

Trimethylsilyl chloroacetate [18293-71-5] M 166.7, m -20°, b 57-58°/14mm, 70-71°/30mm, 159°/760mm, d_4^{20} 1.057, n_D^{20} 1.4231. Purified by repeated fractionation and taking the fractions with clean NMR spectra. [*J Am Chem Soc* 2371 1952.]

Trimethylsilyl cyanide [7677-24-9] M 99.2, m 8-11°, 10.5-11.5°, 11-12°, 12-12.5°; b 54-55°/87mm, 67-71°/168mm, 114-117°/760mm, 118-119°/760mm, d_4^{20} 0.79, n_D^{20} 1.43916. Material should have only one sharp signal in the ¹H NMR (in CCl₄ with CHCl₃ as internal standard) δ : 0.4 and IR with ν at 2210cm⁻¹ [*J Am Chem Soc* 74 5247 1952, 77 3224 1955]; otherwise purify by fractionating through an 18 x 1/4 in column. [*J Am Chem Soc* 81 4493 1959.] It has also been carefully distilled using a 60cm vac jacketed column. If volume of sample is small the cyanide can be chased (in the distillation) with xylene that had be previously distilled over P₂O₅. [*J Org Chem* 39 914 1974.]

2-Trimethylsilyl-1,3-dithiane [13411-42-2] M 192.2, b 54.5°/0.17mm, 100°/8mm, d_4^{20} 1.04, n_D^{20} 1.533. Fractionally distil through an efficient column and collect the fractions that have the correct NMR and IR spectra. ¹H NMR (CCl₄) τ 6.36 (SiMe₃), 9.87 (SCHS) and dithiane H at 7 and 8 (ratio 1:9:4:2) from Me₄Si; UV λ_{max} 244nm (ϵ 711); sh 227nm (ϵ 800). [*J Am Chem Soc* 89 434 1967.]

Trimethylsilyl ethanol [2916-68-9] M 118.3, b 53-55°/11mm, 75°/41mm, 95°/100mm, d_4^{25} 0.8254, n_D^{25} 1.4220. If the NMR spectrum is not clean then dissolve in Et₂O, wash with aqueous NH₄Cl solution, dry (Na₂SO₄), evaporate and distil. The 3,4-dinitrobenzoyl deriv has m 66° (from EtOH). [NMR: *J Am Chem Soc* 79 974 1957; *Z Naturforsch* 14b 137 1959.]

2-(Trimethylsilyl)ethoxymethyl chloride (SEMCl) [76513-69-4] **M 166.7, b 57-59°/8mm, d 0.942, n 1.4350**. Dissolve in pentane, dry (MgSO₄), evaporate and dist residual oil in a vac. Stabilise with 10ppm of diisopropylamine. Store under N₂ in a sealed container in a refrigerator. [Lipshutz and Pegram *Tetrahedron Lett* **21** 3343 1980.]

2-(Trimethylsilyl)ethoxymethyltrimethylphosphonium chloride [82495-75-8] **M 429.0, m 140-142°**. Wash the solid with AcOH and recryst from CH₂Cl₂-EtOAc. Dry in a vacuum desiccator. *Hygroscopic*. ¹H NMR (CDCl₃) δ: -0.2 (s, Me₃Si), 0.8 (t, 8Hz, CH₂Si), 3.83 (t, 8Hz, OCH₂), 5.77 (d, *J*_{PH} 4Hz, P⁺-CH₂O) and 7.70 (m, aromatic H). [Justus Liebigs *Ann Chem* 1031 1983.]

Trimethylsilylethyl phenylsulfone (phenyl-2-trimethylsilylethylsulfone) [73476-18-3] **M 242.4, m 52°**. Dissolve in Et₂O, wash with saturated HCO₃⁻, saturated NaCl, H₂O and dried (MgSO₄). Evaporation leaves residual crystals **m 52°**. [Tetrahedron Lett **23** 1963 1982, *J Org Chem* **53** 2688 1985.]

1-Trimethylsilyloxy-1,3-butadiene [6651-43-0] **M 142.3, b 131°/760mm (mixt of isomers), 49.5°/25mm (E-isomer), d 0.8237, n 1.447**. Purified by fractional distn and collecting the fractions with the required ¹H NMR. Store under N₂ — it is a flammable and moisture sensitive liquid. [Caseau et al. *Bull Soc Chim Fr* 16658 1972; Belge Patent 670,769, *Chem Abstr* **65** 5487d 1966.]

1-(Trimethylsilyloxy)cyclopentene [19980-43-9] **M 156.3, b 45°/11mm, 75-80°/20-21mm, d₄²⁰ 0.878, n_D²⁰ 1.441**. If too impure as seen by the NMR spectrum then dissolve in 10 vols of pentane, shake with cold NaHCO₃ (3 x 500mL), then 1.5M HCl (200mL) and aqueous NaHCO₃ (200mL) again, dry (Na₂SO₄), filter, evaporate and distil through a short Vigreux column. ¹H NMR: (CDCl₃) δ: 0.21 (s, 9H), 1.55 (m, 2H), 1.69 (m, 2H), 2.05 (br d, 4H) and 4.88 (br s, 1H). GLPC in a 6ft x 1/8in with 3% SP2100 on 100-120 mesh Supelcoport column should give one peak. [Org Synth Coll Vol VIII 460 1993.]

2-(Trimethylsilyloxy)furan [61550-02-5] **M 156.3, b 34-35°/9-10mm, 42-50°/17mm, 40-42°/25mm, d₄²⁰ 0.950, n_D²⁰ 1.436**. Fractionally distilled using a short path column. ¹H NMR in CCl₄ has δ: 4.90 (dd, *J* 1.3Hz, 3H), 6.00 (t, *J* 3Hz, 4H) and 6.60 (m, 5H). [Heterocycles **4** 1663 1976.]

4-Trimethylsilyloxy-3-penten-2-one (cis) (acetylacetone enol trimethylsilyl ether) [13257-81-3] **M 172.3, b 66-68°/4mm, 61-63°/5mm, d₄²⁰ 0.917, n_D²⁰ 1.452**. Fractionally distilled and stored in glass ampoules which are sealed under N₂. It hydrolyses readily in contact with moisture giving, as likely impurities, hexamethyldisiloxane and 2,4-pentanedione. [*J Am Chem Soc* **80** 3246 1958.]

Trimethylsilyl isocyanate [1118-02-1] **M 115.2, b 90-92°/atm, b 91.3-91.6°/atm, d₄²⁰ 0.850, n_D²⁰ 1.43943**. Purified by repeated fractionation as for the isothiocyanate. [*J Chem Soc* 3077 1950.]

Trimethylsilyl isothiocyanate [2290-65-5] **M 131.3, m -33°, b 142.6-143.1°/759mm, 143.8°/760mm, n_D²⁰ 1.4809**. The ¹H NMR spectrum should have only one peak, if not purify by repeated fractionation in an all glass system using a 50cm (4mm internal diameter) column without packing. [*J Am Chem Soc* **69** 3049 1947; *Chem Ber* **90** 1934 1957; *Synthesis* 51 1975.]

(Trimethylsilyl)methanol [3219-63-4] **M 104.2, b 120-122°/754mm, 122-123°/768mm, d₄²⁰ 0.83, n_D²⁰ 1.4176**. If the NMR indicates impurities (should have only two signals) then dissolve in Et₂O, shake with aqueous 5N NaOH, M H₂SO₄, saturated aqueous NaCl, dry (MgSO₄) and distil using an efficient column at atmospheric pressure. The 3,5-dinitrobenzoate has **m 70-70.5°** (from 95% EtOH). [Huang and Wang *Acta Chem Sin* **23** 291 1957, *Chem Abs* **52** 19911 1958; Speier et al. *J Am Chem Soc* **81** 1844 1959 and *J Am Chem Soc* **70** 1117 1949.]

(Trimethylsilyl)methylamine (aminomethyl trimethylsilane) [18166-02-4] **M 103.2, b 101.6°/735mm, d₄²⁰ 0.77, n_D²⁰ 1.416**. A possible contaminant is hexamethyldisiloxane. Should have two ¹H NMR signals in CDCl₃, if not, dissolve in ^{*}C₆H₆, shake with 15% aq KOH, separate, dry (Na₂SO₄), filter, evaporate and distil using a still of ca 10 theoretical plates. The water azeotrope has **b 83°/735mm**,

hence it is important to dry the extract well. The *hydrochloride* has *m* 198/199° (from MeOH or Me₂CO). [*J Am Chem Soc* 73 3867 1951; NMR, IR: *J Organometal Chem* 44 279 1972.]

Trimethylsilylmethyl phenylsulfone (phenyltrimethylsilylmethylsulfone) [17872-92-3] *M* 228.4, *m* 28-32°, *b* 121°/0.01mm, 160°/6mm, *n*_D²⁰ 1.5250. Fractionate at high vacuum and recrystallise from pentane at -80°. If too impure (*cf* IR) dissolve in CH₂Cl₂ (*ca* 800mL for 100g), wash with 2M aqueous NaOH (2 x 200mL), brine, dry, evaporate and distil. [*J Chem Soc, Perkin Trans 1* 1949 1985; IR and NMR: *J Am Chem Soc* 76 3713 1954.]

1-(Trimethylsilyl)-2-phenylacetylene (1-phenyl-2-trimethylsilylacetylene) [78905-09-6] *M* 174.3, *b* 45-46°/0.1mm, 67°/5mm, 87.5°/9mm, *d* 0.8961 *n* 1.5284. Dissolve in Et₂O, wash with H₂O, dry and fractionate through a Todd column (see p. 174). [*J Am Chem Soc* 80 5298 1958.]

3-(Trimethylsilyl)propyne [13361-64-3] *M* 112.3, *b* 99-100°/760mm, *d* 0.7581, *n* 1.4091. Fractionally distilled and 0.5% of 2,6-di-*tert*-butyl-*p*-cresol added to stabilise it. [*Doklady Acad Nauk USSR* 93 293 1953; *Chem Abs* 48 13616 1954.]

1-Trimethylsilyl-1,2,4-triazole [18293-54-4] *M* 141.3, *b* 74°/12mm, *d*₄²⁰ 0.99, *n*_D²⁰ 1.4604. Fractionally distilled at atmospheric pressure in an inert atmosphere because it is moisture sensitive. [*Chem Ber* 93 2804 1960.]

Trimethylsilyl trifluoromethane (trifluoromethyl trimethylsilane) [81290-20-2] *M* 142.2, *b* 54-55°, 55-55.5°, *d*₄²⁰ 0.962, *n*_D²⁰ 1.332. Purified by distilling from trap to trap in a vacuum of 20mm using a bath at 45° and Dry ice-Me₂CO bath for the trap. The liquid in the trap is then washed with ice cold H₂O (3x), the top layer is collected, dried (Na₂SO₄), the liquid was decanted and fractionated through a helices packed column at atmospheric pressure. ¹H, ¹³C, ¹⁹F, and ²⁹Si NMR can be used for assessing the purity of fractions. [*Tetrahedron Lett* 25 2195 1984; *J Org Chem* 56 984 1991.]

Trimethyl vinyl silane [754-05-2] *M* 100.2, *b* 54.4°/744mm, 55.5°/767mm, *d*_{S(25,4)} 0.6865, *n*_D²⁵ 1.3880. If the ¹H NMR spectrum shows impurities then dissolve in Et₂O, wash with aq NH₄Cl soln, dry over CaCl₂, filter, evaporate and distil at atmospheric pressure in an inert atmosphere. It is used as a copolymer and may polymerise in the presence of free radicals. It is soluble in CH₂Cl₂. [*J Org Chem* 17 1379 1952.]

Trineopentyl phosphate [14540-59-1] *M* 320.4. Crystd from hexane.

Tri-(4-nitrophenyl)phosphate [3871-20-3] *M* 461.3, *m* 155-156°, 156°, 156-158°, 157-159°. It has been recrystd from AcOH, dioxane, AcOEt and Me₂CO and dried in vacuum over P₂O₅. [*J Am Chem Soc* 72 5777 1950, 79 3741 1957.]

Tri-*n*-octylphosphine oxide [78-50-2] *M* 386.7, *m* 59.5-60°, *pK*_{Est} <0. Mason, McCarty and Peppard [*J Inorg Nuclear Chem* 24 967 1962] stirred an 0.1M solution in *benzene with an equal volume of 6M HCl at 40° in a sealed flask for 48h, then washed the *benzene solution successively with water (twice), 5% aq Na₂CO₃ (three times) and water (six times). The *benzene and water were then evaporated under reduced pressure at room temperature. Zingaro and White [*J Inorg Nucl Chem* 12 315 1960] treated a pet ether solution with aqueous KMnO₄ (to oxidise any phosphinous acids to phosphinic acids), then with sodium oxalate, H₂SO₄ and HCl (to remove any manganese compounds). The pet ether solution was slurried with activated alumina (to remove phosphinic acids) and recrystd from pet ether or cyclohexane at -20°. It can also be crystd from EtOH.

Triphenylantimony [603-36-1] *M* 353.1, *m* 52-54°. Recrystd from acetonitrile [Hayes et al. *J Am Chem Soc* 107 1346 1985].

Triphenylarsine [603-32-7] *M* 306.2, *m* 60-62°. Recrystd from EtOH or aqueous EtOH [Dahlinger et al. *J Chem Soc, Dalton Trans* 2145 1986; Boert et al. *J Am Chem Soc* 109 7781 1987].

Triphenylbismuth (bismuth triphenyl) [603-33-8] M 440.3, m 75-76°, 77-78°, 78.5°, d_4^{25} 1.6427(melt). Dissolve in EtOH, ppt with H₂O, extract with Et₂O, dry and evaporate when the residue crystallises. It has been recrystd from EtOH and Et₂O-EtOH and is a stable compound. [*J Chem Soc suppl* p121 1949; *Chem Ber* 37 4620 1904; *J Am Chem Soc* 62 665 1940; UV: *J Chem Phys* 22 1430 1954.]

Triphenyl borane (borane triphenyl) [960-71-4] M 242.1, m 134-140°, 137°, 139-141°, 142-142.5°, 147.5-148°, 151°, b 203°/15mm. Recryst three times from Et₂O or *C₆H₆ under N₂ and dry at 130°. It can be distilled in a high vacuum at 300-350°, and has been distilled (b 195-215°) in vacuum using a bath temp of 240-330°. N₂ was introduced into the apparatus before dismantling. It forms complexes with amines. [*Chem Ind (London)* 1069 1957; *Justus Liebigs Ann Chem* 563 110 1949; *J Am Chem Soc* 57 1259 1935.]

Triphenylchlorostannane (triphenyltin chloride) [639-58-7] M 385.5, m 103-106°(dec). See triphenyltin chloride on p. 494.

Triphenyl phosphate [115-86-6] M 326.3, m 49.5-50°, b 245°/0.1mm. Crystd from EtOH.

Triphenylphosphine [603-35-0] M 262.3, m 77-78°, 79°, 79-81°, 80.5°, 80-81°, b >360°(in inert gas), d_4^{25} 1.194, d_4^{80} 1.075 (liq), pK 2.73. Crystd from hexane, MeOH, diethyl ether, CH₂Cl₂/hexane or 95% EtOH. Dried at 65°/1mm over CaSO₄ or P₂O₅. Chromatographed through alumina using (4:1) *benzene/CHCl₃ as eluent. [Blau and Espenson et al. *J Am Chem Soc* 108 1962 1986; Buchanan et al. *J Am Chem Soc* 108 1537 1986; Randolph and Wrighton *J Am Chem Soc* 108 3366 1986; Asali et al. *J Am Chem Soc* 109 5386 1987.] It has also been crystd twice from pet ether and 5 times from Et₂O-EtOH to give m 80.5°. Alternatively, dissolve in conc HCl, and upon dilution with H₂O it separates because it is weakly basic; it is then crystallised from EtOH-Et₂O. It recrystallises unchanged from AcOH. [*J Chem Soc Suppl.* p121 1949; *J Am Chem Soc* 78 3557 1956.] 3Ph₃P .4HCl crystallises when HCl gas is bubbled through an Et₂O solution; it has m 70-73°, but recrystallises very slowly and is deliquescent. The HI, made by adding Ph₃P to hydriodic acid is not hygroscopic and decomps at ~100°. The chlorate (1:1) salt has m 165-167°, but decomposes slowly at 100°. All salts hydrolyse in H₂O to give Ph₃P [IR, UV: *J Am Chem Soc* 80 2117 1958; pK: *J Am Chem Soc* 82 5791 1960; Kosolapoff, *Organophosphorus Compounds*, Wiley 1950].
§ Available commercially on a polystyrene or polyethyleneglycol support.

Triphenyl phosphine dibromide [1034-39-5] M 422.1, m 235°, 245-255°(dec). Recrystd from MeCN-Et₂O. Although it has been recrystd from EtOH, this is not recommended as it converts alcohols to alkyl bromides. It deteriorates on keeping and it is best to prepare it afresh. [Anderson and Frenor *J Am Chem Soc* 86 5037 1964; Horner et al. *Justus Liebigs Ann Chem* 626 26 1959.]

Triphenylphosphine oxide [791-28-6] M 278.3, m 152.0°, pK_{Est} ~ -2.10 (aq H₂SO₄). Crystd from absolute EtOH. Dried *in vacuo*.

Triphenyl phosphite [101-02-0] M 310.3, b 181-189°/1mm, d 1.183. Its ethereal soln was washed successively with aqueous 5% NaOH, distilled water and saturated aqueous NaCl, then dried with Na₂SO₄ and distilled under vacuum after evaporating the diethyl ether.

Triphenylphosphorylidene acetaldehyde (formylmethylenetriphenylphosphorane) [2136-75-6] M 304.3, m 185-187°, 186-187°(dec). Recryst from Me₂CO, or dissolve in *C₆H₆, wash with N NaOH, dry (MgSO₄), evap, and cryst residue from Me₂CO. It can be prepd from its precursor, formylmethyltriphenylphosphonium chloride (crystd from CHCl₃/EtOAc), by treatment with Et₃N and extraction with *C₆H₆. [Trippett and Walker *Chem Soc* 1266 1961.]

Triphenyl silane [789-25-3] M 260.4, m 45°, b 148-151°/1mm. Purified by recrystn from MeOH. [*J Am Chem Soc* 81 5925 1959; *Acta Chem Scand* 9 947 1955; IR: *J Am Chem Soc* 76 5880 1954.]

Triphenylsilanol (hydroxytriphenylsilane) [791-31-1] **M 276.4, m 150-153°, 151-153°, 154-155°, 156°**. It can be purified by dissolving in pet ether, passing through an Al_2O_3 column, eluting thoroughly with CCl_4 to remove impurities and then eluting the silanol with MeOH. Evaporation gives crystals **m 153-155°**. It can be recrystallised from pet ether, CCl_4 or from *benzene or Et_2O -pet ether (1:1). It has also been recrystallised by partial freezing from the melt to constant melting point. [*J Am Chem Soc* **81** 3288 1959; IR: *J Org Chem* **17** 1555 1952 and *J Chem Soc* 124 1949.]

Triphenyltin chloride (chlorotriphenylstannane) [639-58-7] **M 385.5, m 103-106°(dec), 108°(dec), b 240°/13.5mm**. Purify by distillation, followed by recrystn from MeOH by adding pet ether (b 30-60°), **m 105-106°** [*Chem Ber* **67** 1348 1934], or by crystn from Et_2O [Krause *Chem Ber* **51** 914 1918]. It sublimes in a vacuum. **HIGHLY TOXIC**.

Triphenyltin hydroxide [76-87-9] **M 367.0, m 122-123.5°, 124-126°**. West, Baney and Powell [*J Am Chem Soc* **82** 6269 1960] purified a sample which was grossly contaminated with tetraphenyltin and diphenyltin oxide by dissolving it in EtOH, most of the impurities remaining behind as an insoluble residue. Evaporation of the EtOH gave the crude hydroxide which was converted to triphenyltin chloride by grinding in a mortar under 12M HCl, then evaporating the acid soln. The chloride, after crystallisation from EtOH, had **m 104-105°**. It was dissolved in Et_2O and converted to the hydroxide by stirring with excess aqueous ammonia. The ether layer was separated, dried, and evaporated to give triphenyltin hydroxide which, after crystn from EtOH and drying under vacuum, was in the form of white crystals (**m 119-120°**), which retained some cloudiness in the melt above 120°. The hydroxide retains water (0.1-0.5 moles of water per mole) tenaciously.

Triphenyl vinyl silane [18666-68-7] **M 286.5, m 58-59°, 57-59.5°, 67-68°, b 190-210°/3mm**. It has been recrystallised from EtOH, 95% EtOH, $\text{EtOH} \cdot \text{C}_6\text{H}_6$, pet ether (b 30-60°) and Et_2O , and has been distilled under reduced pressure. [*J Am Chem Soc* **74** 4582 1952; *J Org Chem* **17** 1379 1952.]

Tri-*n*-propyl borate [688-71-1] **M 188.1, b 175-177°, d 0.857, n 1.395**. Dried with sodium and then distilled.

Triquinol-8-yl phosphate [52429-99-9] **M 479.4, m 193-197°, 202-203°**. Purified by recrystn from dimethylformamide. Purity was checked by paper chromatography, R_F 0.90 [*i*-PrOH, saturated $(\text{NH}_4)_2\text{SO}_4$, H_2O ; 2:79:19 as eluent]; IR (KBr) ν 1620-1570 (C=C, C=N) and 1253 (P=O). [*Bull Chem Soc Jpn* **47** 779 1974.]

Tri-ruthenium dodecacarbonyl [15243-33-1] **M 639.1, m 154-155°**. Recryst from * C_6H_6 or cyclohexane as orange-red crystals, and sublime at 80-100°/0.1mm. It has ν_{CO} 2062 and 2032. [*J Chem Soc, Chem Commun* 684 1966; *J Chem Soc (A)* 1238 1967; IR,UV: *Angew Chem Int Ed Engl* **7** 427 1968.]

Tris-(2-biphenyl) phosphate [132-28-5] **M 554.6, m 115.5-117.5°**. Crystd from MeOH containing a little acetone.

Tris(2,2'-bipyridine)ruthenium(II) dichloride (6H₂O) [50525-27-4] **M 748.6**. Recrystd from water then from MeOH [Ikezawa et al. *J Am Chem Soc* **108** 1589 1986].

Tris-(1,2-dioxyphenyl)cyclotriphosphazine {trisp[1,3,5,2,4,6-triazatriphosphorine]-2,2':-2,4''':2,6'''-tris(1,3,2)benzodioxaphosphole} [311-03-5] **M 459.0, m 244-245°, 245°, 245-246°**. Recrystd from * C_6H_6 or chlorobenzene, then triple sublimed (175°/0.1mm, 200°/0.1mm, 230°/0.05mm). UV has λ_{max} nm (log ϵ): 276 (3.72), 271 (3.79) 266sh (3.68) and 209 (4.38) in MeCN. IR (ν): 1270 (O-Ph), 1220 (P=N), 835 (P-O-Ph) and 745 (Ph) cm^{-1} . [Alcock *J Am Chem Soc* **86** 2591 1964; Alcock et al. *J Am Chem Soc* **98** 5120 1976; Meirovitch *J Phys Chem* **88** 1522 1984.]

(±)-Tris-(2-ethylhexyl)phosphate (TEHP, tri-isooctylphosphate, "trioctyl" phosphate, [78-42-2; 25103-23-5] M 434.6, b 186°/1mm, 219°/5mm, d^{25} 0.92042, n 1.44464. TEHP, in an equal volume of diethyl ether, was shaken with aqueous 5% HCl and the organic phase was filtered to remove

traces of pyridine (used as a solvent during manufacture) as its hydrochloride. This layer was shaken with aqueous Na_2CO_3 , then water, and the ether was distilled off at room temperature. The ester was filtered, dried for 12h at $100^\circ/15\text{mm}$, and again filtered, then shaken intermittently for 2 days with activated alumina (100g/L). It was decanted through a fine sintered-glass disc (with exclusion of moisture), and distd under vacuum. [French and Muggleton *J Chem Soc* 5064 1957.] *Benzene can be used as a solvent (to give 0.4M soln) instead of ether. IR: 1702, 1701, 481 and 478cm^{-1} [Bellamy and Becker *J Chem Soc* 475 1952]. The *uranyl nitrate* salt was purified by partial crystallisation from hexane [Siddall *J Am Chem Soc* 81 4176 1959].

Trisodium citrate ($2\text{H}_2\text{O}$) [68-04-2] M 294.1, m 150° (loses H_2O). Crystd from warm water by cooling to 0° .

Trisodium 8-hydroxy-1,3,6-pyrenetrisulfonate [6358-69-6] M 488.8, m >300 (dec). Purified by chromatography with an alumina column, and eluted with *n*-propanol-water (3:1, v/v). Recrystd from aqueous acetone (5:95, v/v) using decolorising charcoal.

Trisodium 1,3,6-naphthalenetrisulfonate [5182-30-9] M 434.2. The free acid was obtained by passage through an ion-exchange column and converted to the lanthanum salt by treatment with La_2O_3 . This salt was crystallised twice from hot water. [The much lower solubility of $\text{La}_2(\text{SO}_4)_3$ and its retrograde temperature dependence allows a good separation from sulfate impurity]. The lanthanum salt was then passed through an appropriate ion-exchange column to obtain the free acid, the sodium or potassium salt. (The sodium salt is *hygroscopic*). [Atkinson, Yokoi and Hallada *J Am Chem Soc* 83 1570 1961.] Also recrystd from aqueous acetone [Okahata et al. *J Am Chem Soc* 108 2863 1986].

Trisodium orthophosphate ($12\text{H}_2\text{O}$) [10101-89-0] M 380.1, pK_1^{25} 2.15, pK_2^{25} 7.21, pK_3^{25} 12.33 (for H_3PO_4). Crystd from warm dilute aqueous NaOH (1mL/g) by cooling to 0° .

Tris(2,4-pentandionate)aluminium [13963-57-0] M 324.3. See aluminum acetylacetonate on p. 390.

Tris-(trimethylsilyl)silane (TTMSS) [1873-77-4] M 248.7, b $73^\circ/5\text{mm}$, d 0.808, n 1.49. Purified by fractional distn and taking the middle cut. Store under N_2 or Ar as it is an IRRITANT and PYROPHORIC. [Chatgialiloglu *Acc Chem Res* 25 188 1992; NMR: Gilman et al. *J Organomet Chem* 4 163 1965.]

Tritium [10028-17-8] M 6.0. Purified from hydrocarbons and ^3He by diffusion through the wall of a hot nickel tube [Landecker and Gray *Rev Sci Instrum* 25 1151 1954]. RADIOACTIVE.

Tri-*p*-tolyl phosphate [20756-92-7; 1330-78-5 (isomeric tritolyl phosphate mixture)] M 368.4, b $232-234^\circ$, d^{25} 1.16484, n 1.56703. Dried with CaCl_2 , then distd under vacuum and percolated through a column of alumina. Passage through a packed column at 150° , with a counter-current stream of nitrogen, under reduced pressure, removed residual traces of volatile impurities.

Tri-*o*-tolylphosphine [6163-58-2] M 304.4, m $129-130^\circ$, $\text{pK}_{\text{Est}} < 0$. Crystd from EtOH [Boert et al. *J Am Chem Soc* 109 7781 1987].

Tungsten (rod) [7440-33-7] M 183.6. m 3410° , b 5900° , d 19.0. Cleaned with conc NaOH solution, rubbed with very fine emery paper until its surface was bright, washed with previously boiled and cooled conductivity water and dried with filter paper.

Tungsten hexacarbonyl [14040-11-0] M 351.9, d 2.650. Sublimed *in vacuo* before use [Connoe et al. *J Chem Soc, Dalton Trans* 511 1986].

Tungsten (VI) trichloride [13283-01-7] M 396.6, m 265° (dec), 275° , b 346° , d_4^{25} 3.520, pK_1^{25} 2.20, pK_2^{25} 3.70 (for tungstic acid, H_2WO_4). Sublimed in a stream of Cl_2 in a high temperature furnace and collected in a receiver cooled in a Dry Ice-acetone bath in an inert atmosphere because it is sensitive to moisture. It is soluble in CS_2 , CCl_4 , CHCl_3 , POCl_3 , $^*\text{C}_6\text{H}_6$, pet ether and Me_2CO . Solns decompose on

standing. Good crystals can be obtained by heating WCl_6 in CCl_4 to 100° in a sealed tube, followed by slow cooling (tablets of four-sided prisms). Store in a desiccator over H_2SO_4 in the dark. [*Inorg Synth* 3 163 1950, 9 133 1967; *Handbook of Preparative Inorganic Chemistry* (Ed. Brauer) Vol II 1417 1965.]

Uranium hexafluoride [7783-81-5] M 352.0, b $0^\circ/17.4\text{mm}$, $56.2^\circ/765\text{mm}$, m 64.8° , pK^{25} 1.68 (for hydrolysis of U^{4+} to UOH^{3+}). Purified by fractional distillation to remove HF. Also purified by low temperature trap-to-trap distillation over pre-dried NaF [Anderson and Winfield *J Chem Soc, Dalton Trans* 337 1986].

Uranium trioxide [1344-58-7] M 286.0, d 7.29. The oxide was dissolved in HClO_4 (to give a uranium content of 5%), and the solution was adjusted to pH 2 by addition of dilute ammonia. Dropwise addition of 30% H_2O_2 , with rapid stirring, ppted U(VI) peroxide, the pH being held constant during the pptn, by addition of small amounts of the ammonia soln. (The H_2O_2 was added until further quantities caused no change in pH.) After stirring for 1h, the slurry was filtered through coarse filter paper in a Büchner funnel, washed with 1% H_2O_2 acidified to pH 2 with HClO_4 , then heated at 350° for three days in a large platinum dish [Baes *J Phys Chem* 60 878 1956].

Uranyl nitrate ($6\text{H}_2\text{O}$) [13520-83-7] M 502.1, m 60.2° , b 118° , pK^{25} 5.82 (for aquo UO_2^{2+}). Crystd from water by cooling to -5° , taking only the middle fraction of the solid which separated. Dried as the hexahydrate over 35-40% H_2SO_4 in a vacuum desiccator.

Vanadium (metal) [7440-62-2] M 50.9, m 1910° , d 6.0. Cleaned by rapid exposure consecutively to HNO_3 , HCl, HF, de-ionised water and reagent grade acetone, then dried in a vacuum desiccator.

Vanadium (III) acetylacetonate [13476-99-8] M 348.3, m $181-184^\circ$, $185-190^\circ$, pK_1^{25} 2.92, pK_2^{25} 3.5 (for aquo V^{3+} hydrolysis). Crystd from acetylacetone as brown plates. It can be distilled in small quantities without decomposition. It is soluble in CHCl_3 and $^*\text{C}_6\text{H}_6$ and evaporation of a CHCl_3 solution yields brown crystals which are washed with cold EtOH and dried in vacuum or at 100° in a CO_2 atmosphere. Under moist conditions it readily oxidises [$\text{V}(\text{AcAc})_3$ to $\text{V}(\text{AcAc})_2\text{O}$]. [*J Chem Soc* 103 78 1913, *Inorg Synth* 5 105 1957; *Anal Chem* 30 526 1958; UV: *J Am Chem Soc* 80 5686 1958.]

Vanadyl acetylacetonate [3153-26-2] M 265.2, m $256-259^\circ$. Crystd from acetone.

Vanadyl trichloride (VOCl_3) [7727-18-6] M 173.3, m -79.5° , b $124.5-125.5^\circ/744\text{mm}$, $127.16^\circ/760\text{mm}$, d_4^{20} 1.854, d_4^{32} 1.811. Should be lemon yellow in colour. If red it may contain VCl_4 and Cl_2 . Fractionally distil and then redistil over metallic Na but be careful to leave some residue because the residue can become **EXPLOSIVE** in the presence of the metal **USE A SAFETY SHIELD and avoid contact with moisture**. It readily hydrolyses to vanadic acid and HCl. Store in a tightly closed container or in sealed ampoules under N_2 . [*Inorg Synth* 1 106 1939, 4 80 1953.]

Vinyl chlorosilane [75-94-5] M 161.5, b $17.7^\circ/46.3\text{mm}$, $82.9^\circ/599.4\text{mm}$, $92^\circ/742\text{mm}$, $91-91.5^\circ/\text{atm}$, d_4^{20} 0.1.2717, n_D^{20} 1.435. Fractionally distil at atmospheric pressure. It is H_2O sensitive and is stored in the dark and is likely to polymerise. [*Chem Ber* 91 1805 1958, 92 1012 1959; *Anal Chem* 24 1827 1952]

Vinylferrocene (ferrocenylethene) [1271-51-8] M 212.1, m $51-52.5^\circ$, b $80-85^\circ/0.2\text{mm}$. Dissolve in Et_2O , wash with H_2O and brine, dry (Na_2SO_4), evap to a small vol. Purify through an Al_2O_3 (Spence grade H) column by eluting the yellow band with pet ether (b $40-60^\circ$). The low melting orange crystals which can be sublimed. The *tetracyanoethylene adduct* [49716-63-4] crysts from $^*\text{C}_6\text{H}_6$ -pentane and has m 137-

139°(dec). [Horspool and Sutherland *Can J Chem* **46** 3453 1968; Berger et al. *J Org Chem* **39** 377 1974; Rauch and Siegel *J Organomet Chem* **11** 317 1968.]

Vinyltributylstannane (vinyltributyltin) [7486-35-3] **M 317.1, b 104-106°/3.5mm, d 1.081, n 1.4751**. Fractionate under reduced pressure and taking the middle fraction to remove impurities such as (*n*-Bu)₃SnCl. [Seyferth and Stone *J Am Chem Soc* **79** 515 1957.]

Water [7732-18-5] **M 18.0, m 0°, b 100°, pK²⁵ 14.00**. Conductivity water (specific conductance *ca* 10⁻⁷ mho) can be obtained by distilling water in a steam-heated tin-lined still, then, after adding 0.25% of solid NaOH and 0.05% of KMnO₄, distilling once more from an electrically heated Barnstead-type still, taking the middle fraction into a Jena glass bottle. During these operations suitable traps must be used to protect against entry of CO₂ and NH₃. Water only a little less satisfactory for conductivity measurements (but containing traces of organic material) can be obtained by passing ordinary distilled water through a mixed bed ion-exchange column containing, for example, Amberlite resins IR 120 (cation exchange) and IRA 400 (anion exchange), or Amberlite MB-1. This treatment is also a convenient one for removing traces of heavy metals. (The metals Cu, Zn, Pb, Cd and Hg can be tested for by adding pure concentrated ammonia to 10mL of sample and shaking vigorously with 1.2mL 0.001% dithizone in CCl₄. Less than 0.1μg of metal ion will impart a faint colour to the CCl₄ layer.) For almost all laboratory purposes, simple distillation yields water of adequate purity, and most of the volatile contaminants such as ammonia and CO₂ are removed if the first fraction of distillate is discarded.

Xylenol Orange (sodium salt) [3618-43-7]. See entry on p. 387 in Chapter 4.

Zinc (dust) [7440-66-6] **M 65.4**. Commercial zinc dust (1.2Kg) was stirred with 2% HCl (3L) for 1min, (then the acid was removed by filtration), and washed in a 4L beaker with a 3L portion of 2% HCl, three 1L portions of distilled water, two 2L portions of 95% EtOH, and finally with 2L of absolute Et₂O. (The wash solutions were removed each time by filtration.) The material was then dried thoroughly and if necessary, any lumps were broken up in a mortar.

Zinc (metal) [7440-66-6] **M 65.4, m 420°, d 7.141**. Fused under vacuum, cooled, then washed with acid to remove the oxide.

Zinc acetate (2H₂O) [5970-45-6] **M 219.5, m 100°(loses 2H₂O), 237°, d 1.74, pK²⁵ 8.96 (for hydrolysis of Zn²⁺ to ZnOH⁺)**. Crystd (in poor yield) from hot water or, better, from EtOH.

Zinc acetylacetonate [14024-63-6] **M 263.6, m 138°**. Crystd from hot 95% EtOH.

Zinc bromide [7699-45-8] **M 225.2, m 384, b 697**. Heated to 300° under vacuum (2 x 10⁻²mm) for 1h, then sublimed.

Zinc caprylate [557-09-5] **M 351.8**. Crystd from EtOH.

Zinc chloride [7646-85-7] **M 136.3, m 283°, 290°**. The anhydrous material can be sublimed under a stream of dry HCl, followed by heating to 400° in a stream of dry N₂. Also purified by refluxing (50g) in dioxane (400mL) with 5g zinc dust, filtering hot and cooling to ppt ZnCl₂. Crystd from dioxane and stored in a desiccator over P₂O₅. It has also been dried by refluxing in thionyl chloride. [Weberg et al. *J Am Chem Soc* **108** 6242 1986.] *Hygroscopic: minimal exposure to the atmosphere is necessary.*

Zinc cyanide [557-21-1] M 117.4, m 800°(dec), d 1.852. It is a **POISONOUS** white powder which becomes black on standing if Mg(OH)₂ and carbonate are not removed in the preparation. Thus wash well with H₂O, then well with EtOH, Et₂O and dry in air at 50°. Analyse by titrating the cyanide with standard AgNO₃. Other likely impurities are ZnCl₂, MgCl₂ and traces of basic zinc cyanide; the first two salts can be washed out. It is soluble in aq KCN solns. However, if purified in this way Zn(CN)₂ is not reactive in the Gattermann synthesis. For this the salt should contain at least 0.33 mols of KCl or NaCl which will allow the reaction to proceed faster. [*J Am Chem Soc* 45 2375 1923, 60 1699 1938; *Org Synth Coll Vol III* 549 1955.]

Zinc diethyldithiocarbamate [14324-55-1] M 561.7, pK²⁵ 3.04 (for Et₂NCS₂⁻). Crystd several times from hot toluene or from hot CHCl₃ by addition of EtOH.

Zinc dimethyldithiocarbamate [137-30-4] M 305.8, m 248-250°, pK²⁵ 3.36 (for Me₂NCS₂⁻). Crystd several times from hot toluene or from hot CHCl₃ by addition of EtOH.

Zinc ethylenebis[dithiocarbamate] [12122-67-7] M 249.7. Crystd several times from hot toluene or from hot CHCl₃ by addition of EtOH.

Zinc fluoride [7783-49-5] M 103.4, m 872°, b 1500°, d²⁵ 5.00. Possible impurity is H₂O which can be removed by heating at 100° or by heating to 800° in a dry atmosphere. Heating in the presence of NH₄F produces larger crystals. It is sparingly sol in H₂O (1.51g/100mL) but more sol in HCl, HNO₃ and NH₄OH. It can be stored in glass bottles. [*Handbook of Preparative Inorganic Chemistry (Ed. Brauer)* Vol I 242 1963.]

Zinc formate (2H₂O) [557-41-5] M 191.4, m 140°(loses H₂O), d 2.21. Crystd from water (3mL/g).

Zinc iodide [10139-47-6] M 319.2, m 446, b 624°(dec), d 4.74. Heated to 300° under vacuum (2 x 10⁻²mm) for 1h, then sublimed.

Zinc RS-lactate (3H₂O) [554-05-2; 16039-53-5 (L)] M 297.5. Crystd from water (6mL/g).

Zincon (o-[1-(2-hydroxy-5-sulfo)-3-phenyl-5-formazono]-benzoic acid) [135-52-4] M 459.4. Main impurities are inorganic salts which can be removed by treatment with dilute acetic acid. Organic contaminants are removed by refluxing with ether. It can be recrystd from dilute H₂SO₄. [Fichter and Schiess *Chem Ber* 33 751 1900.]

Zincon disodium salt (o-[1-(2-hydroxy-5-sulfo)-3-phenyl-5-formazono]-benzoic acid di-Na salt) [135-52-4; 56484-13-0] M 484.4, m ~250-260° (dec). Zincon soln is prepared by dissolving 0.13g of the powder in aqueous N NaOH (2mL diluted to 100mL with H₂O). This gives a deep red colour which is stable for one week. It is a good reagent for zinc ions but also forms stable complexes with transition metal ions. [UV-VIS: Bush and Yoe *Anal Chem* 26 1345 1954; Hunter and Roberts *J Chem Soc* 820 1941; Platte and Marcy *Anal Chem* 31 1226 1959] The free acid has been recrystd from dilute H₂SO₄. [Fichter and Scheiss *Chem Ber* 33 751 1900.]

Zinc perchlorate (6H₂O) [13637-61-1] M 372.4, m 105-107°, pK²⁵ -2.4 to -3.1 (for HClO₄). Crystd from water.

Zinc phenol-o-sulfonate (8H₂O) [127-82-2] M 555.8. Crystd from warm water by cooling to 0°.

Zinc phthalocyanine [14320-04-8] M 580.9. Sublimed repeatedly in a flow of oxygen-free N₂.

Zinc sulfate (7H₂O) [7446-20-0] M 287.5, m 100°(dec), 280°(loses all 7H₂O), >500°(anhydr), d 1.97. Crystd from aqueous EtOH.

Zinc 5,10,15,20-tetraphenylporphyrin [14074-80-7] M 678.1, λ_{\max} 418(556)nm. Purified by chromatography on neutral (Grade I) alumina, followed by recrystallisation from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ [Yamashita et al. *J Phys Chem* 91 3055 1987].

Zinc trifluoromethanesulfonate [54010-75-2] M 363.5, m $>300^\circ$. It should be dried at 125° for 2h at 3mm. It is soluble in CH_2Cl_2 but insoluble in pet ether. [*Tetrahedron Lett* 24 169 1983.]

Zirconium (IV) propoxide [23519-77-9] M 327.6, b $198^\circ/0.03\text{mm}$, $208^\circ/0.1\text{mm}$, d_4^{20} 1.06, n_D^{20} 1.454. Although it was stated that it could not be crystallised or sublimed even at $150^\circ/10^{-4}\text{mm}$ [*J Chem Soc* 280 1951], the propoxide has, when properly prepared, been purified by distn in a high vacuum [*J Chem Soc* 2025 1953].

Zirconium tetrachloride [10026-11-6] M 233.0, m 300° (sublimes), pK_1^{25} -0.32, pK_2^{25} 0.06, pK_3^{25} 0.35, pK_4^{25} 0.46 (for hydrolysis of aquo Zr^{4+}). Crystd repeatedly from conc HCl.

Zirconocene chloride hydride (bis[cyclopentadienyl]zirconium hydride chloride) (Schwartz reagent) [37342-97-5] M 257.9. It is a moisture and light sensitive compound. Its purity can be determined by reaction with a slight excess of Me_2CO whereby the active H reacts to produce $\text{Cp}_2\text{ZrClOPr}^i$ and the integrals of the residual Me_2CO in the ^1H NMR will show how pure the sample is. The presence of Cp_2ZrH_2 can be determined because it forms $\text{Cp}_2\text{Zr(OPr}^i)_2$. For very active compound it is best to prepare freshly from the dichloride by reduction with Vitride [$\text{LiAl(OCH}_2\text{CH}_2\text{OH)}_2\text{H}_2$], the white ppt is filtered off, washed with tetrahydrofuran, $^* \text{C}_6\text{H}_6$, Et_2O , dried in vacuum and stored under anhydrous conditions and in the dark. [IR: *J Chem Soc, Chem Commun* 1105 1969; *J Am Chem Soc* 96 8115 1974, 101 3521 1979; *Synthesis* 1 1988.]

Zirconocene dichloride (bis[cyclopentadienyl]zirconium dichloride) [1291-32-3] M 292.3, m $242\text{-}245^\circ$, 248° . Purified by recrystn from CHCl_3 or xylene, and dried in vacuum. ^1H NMR (CDCl_3) δ : 6.52 from Me_4Si . Store in the dark under N_2 as it is moisture sensitive. [IR, NMR, MS: *Aust J Chem* 18 173 1965; method of *J Am Chem Soc* 81 1364 1959; and references in the previous entry.]

Zirconyl chloride ($6\text{H}_2\text{O}$) [7699-43-6] M 286.2, m 150° (loses $6\text{H}_2\text{O}$). Crystd repeatedly from 8M HCl as $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$. On drying $\text{ZrOCl}_2 \cdot 6\text{H}_2\text{O}$ m 150° . The product was not free from hafnium.

Zirconyl chloride ($8\text{H}_2\text{O}$) [13520-92-8] M 322.3, m 150° (loses $6\text{H}_2\text{O}$), 210° (loses all H_2O). 400° (anhydr dec), d 1.91. Recrystd several times from water [Ferragina et al. *J Chem Soc, Dalton Trans* 265 1986]. Recrystn from 8M HCl gives the octahydrate as white needles on concentrating. It is also formed by hydrolysing ZrCl_4 with water. After one recryst from H_2O , 99+% grade had Ag, Al, As, Cd, Cu, Hf, Mg, Na, Sc and V at 20, 1.8, 0.6, 0.6, 0.4, 8.4, 0.4, 2.4, 80 and 3 ppm resp.

CHAPTER 6

PURIFICATION OF BIOCHEMICALS AND RELATED PRODUCTS

Biochemicals are chemical substances produced by living organisms. They range widely in size, from simple molecules such as formic acid and glucose to macromolecules such as proteins and nucleic acids. Their *in vitro* synthesis is often impossibly difficult and in such cases they are available (if at all) only as commercial tissue extracts which have been subjected to purification procedures of widely varying stringency. The desired chemical may be, initially, only a minor constituent of the source tissue which may vary considerably in its composition and complexity. Recent advances in molecular biology have made it possible to produce substantial amounts of biological materials, which are present in nature in extremely small amounts, by recombinant DNA technology and expression in bacteria, yeast, insect and mammalian cells. The genes for these substances can be engineered such that the gene products, e.g. polypeptides or proteins, can be readily obtained in very high states of purity. However, many such products which are still obtained from the original natural sources are available commercially and may require further purification.

As a preliminary step the tissue might be separated into phases [e.g. whole egg into white and yolk, blood into plasma (or serum) and red cells], and the desired phase may be homogenised. Subsequent treatment usually comprises filtration, solvent extraction, salt fractionation, ultracentrifugation, chromatographic purification, gel filtration and dialysis. Fractional precipitation with ammonium sulfate gives crude protein species. Purification is finally judged by the formation of a single band of macromolecule (e.g. protein) on electrophoresis and/or analytical ultracentrifugation. Although these generally provide good evidence of high purity, none-the-less it does not follow that one band under one set of experimental conditions is an absolute indication of homogeneity.

During the past 20 or 30 years a wide range of methods for purifying substances of biological origin have become available. For small molecules (including many sugars and amino acids) reference should be made to Chapters 1 and 2. The more important methods used for large molecules, polypeptides and proteins in particular, comprise:

1. *Centrifugation.* In addition to centrifugation for sedimenting proteins after ammonium sulfate precipitation in dilute aqueous buffer, this technique has been used for fractionation of large molecules in a denser medium or a medium of varying density. By layering sugar solutions of increasing densities in a centrifuge tube, proteins can be separated in a sugar-density gradient by centrifugation. Smaller DNA molecules (e.g. plasmid DNA) can be separated from RNA or nuclear DNA by centrifugation in aqueous cesium chloride (*ca* 0.975g/mL of buffer) for a long time (e.g. 40h at 40,000 x g). The plasmid DNA band appears at about the middle of the centrifuge tube, and is revealed by the fluorescent pink band formed by the binding of DNA to ethidium bromide which is added to the CsCl buffer. *Microfuges* are routinely used for centrifugation in Eppendorf tubes (1.2-2mL) and can run up to speeds of 12,000 x g. *Analytical centrifugation*, which is performed under specific conditions in an analytical ultracentrifuge is very useful for determining purity, aggregation of protein subunits and the molecular weight of macromolecules. [D.Rickwood, T.C.Ford and J.Steensgaard *Centrifugation: Essential Data Series*, J Wiley & Sons, NY, 1994].
2. *Gel filtration* with polyacrylamide (mol wt exclusion limit from 3000 to 300,000) and agarose gel (mol wt exclusion limit 0.5 to 150 x 10⁶) is useful for separating macromolecules. In this technique high-molecular weight substances are too large to fit into the gel microapertures and pass rapidly through the matrix (with the void volume), whereas low molecular weight species enter these apertures and are held there for longer periods of time, being retarded by the column material in the equilibria, relative to the larger molecules. This method is also used for desalting solutions of macromolecules. *Dry gels* and *crushed beads* are also

useful in the gel filtration process. Selective retention of water and inorganic salts by the gels or beads (e.g. Sephadex G-25) results in increased concentration and purity of the protein fraction which moves with the void volume. (See also Chapter 1, pp 23, 41).

3. *Ion exchange matrices* are microreticular polymers containing carboxylic acid (e.g. Bio-Rad 70) or phosphoric acid (Pharmacia, Amersham Biosciences, Mono-P) exchange functional groups for weak acidic cation exchangers, sulfonic acid groups (Dowex 50W) for strong acidic cation exchangers, diethylaminoethyl (DEAE) groups for weakly basic anion exchangers and quaternary ammonium (QEAE) groups for strong anion exchangers. The old cellulose matrices for ion exchanges have been replaced by Sephadex, Sepharose or Fractogel which have more even particle sizes with faster and more reproducible flow rates. Some can be obtained in fine, medium or coarse grades depending on particle size. These have been used extensively for the fractionation of peptides, proteins and enzymes. The use of pH buffers controls the strength with which the large molecules are bound to the support in the chromatographic process. Careful standardisation of experimental conditions and similarly the very uniform size distribution of Mono beads has led to high resolution in the purification of protein solutions. MonoQ (Pharmacia, Amersham Biosciences) is a useful strong anion exchanger, and MonoS (Pharmacia, Amersham Biosciences) is a useful strong cation exchanger whereas MonoP is a weak cation exchanger. These have been successful with medium pressure column chromatography (FPLC, see below in 8). Chelex 100 binds strongly and removes metal ions from macromolecules. [See also Chapter 1, pp. 22-24.]
4. *Hydroxylapatite* is used for the later stages of purification of enzymes. It consists essentially of hydrated calcium phosphate which has been precipitated in a specific manner. It combines the characteristics of gel and ionic chromatography. Crystalline hydroxylapatite is a structurally organised, highly polar material which, in aqueous solution (in buffers) strongly adsorbs macromolecules such as proteins and nucleic acids, permitting their separation by virtue of the interaction with charged phosphate groups and calcium ions, as well as by physical adsorption. The procedure therefore is not entirely ion-exchange in nature. Chromatographic separations of singly and doubly stranded DNA are readily achievable whereas there is negligible adsorption of low molecular weight species.
5. *Affinity chromatography* is a chromatographic technique whereby the adsorbant has a particular and specific affinity for one of the components of the mixture to be purified. For example the adsorbant can be prepared by chemically binding an inhibitor of a specific enzyme (which is present in the crude complex mixture) to a matrix (e.g. Sepharose). When the mixture of impure enzyme is passed through the column containing the adsorbant, only the specific enzyme binds to the column. After adequate washing, the pure enzyme can be released from the column by either increasing the salt concentration (e.g. NaCl) in the eluting buffer or adding the inhibitor to the eluting buffer. The salt or inhibitor can then be removed by dialysis, gel filtration (above) or ultrafiltration (see below). [See W.H.Scouten, *Affinity Chromatography: Bioselective Adsorption on Inert Matrices*, J.Wiley & Sons, NY, 1981, ISBN 0471026492; H.Schott, *Affinity Chromatography: Template Chromatography of Nucleic Acids and Proteins*, Marcel Dekker, NY, 1984, ISBN 0824771117; P.Matejtschuk ed. *Affinity Separations* Oxford University Press 1997 ISBN 01996335501 (paperback); M.A.Vijayalakshmi, *Biochromatography, Theory and Practice*, Taylor & Francis Publ, 2002, ISBN 0415269032; and Chapter 1, p. 25.]
6. In the *Isoelectric focusing* of large charged molecules on polyacrylamide or agarose gels, slabs of these are prepared in buffer mixtures (e.g. ampholines) which have various pH ranges. When a voltage is applied for some time the buffers arrange themselves on the slabs in respective areas according to their pH ranges (prefocusing). Then the macromolecules are applied near the middle of the slab and allowed to migrate in the electric field until they reach the pH area similar to their isoelectric points and focus at that position. This technique can also be used in a chromatographic mode, *chromatofocusing*, whereby a gel in a column is run (also under HPLC conditions) in the presence of ampholines (narrow or wide pH ranges as required) and the macromolecules are then run through in a buffer. *Capillary electrophoresis* systems in which a current is applied to set the gradient are now available in which the columns are fine capillaries and are used for qualitative and quantitative purposes [See R.Kuhn and S.Hoffstetter-Kuhn, *Capillary Electrophoresis: Principles and Practice*, Springer-Verlag Inc, NY, 1993; P.Camilleri ed. *Capillary Electrophoresis - Theory and Practice*, CRC Press, Boca Raton, Florida, 1993; D.R.Baker, *Capillary Electrophoresis*, J Wiley & Sons, NY, 1995; P.G.Righetti, A.Stoyanov and M.Zhukov, *The Proteome Revisited, Isoelectric Focusing: J.Chromatography Library Vol 63* 2001, Elsevier, ISBN 0444505261.] The bands are eluted according to their isoelectric points. Isoelectric focusing standards are available which can be used in a preliminary run in order to calibrate the effluent from the column, or alternatively the pH of the effluent is recorded using a glass electrode designed for the purpose. Several efficient commercially available apparatus are available for separating proteins on a preparative and semi-preparative scale.
7. *High performance liquid chromatography (HPLC)* is liquid chromatography in which the eluting liquid is sent through the column containing the packing (materials as in 2-6 above, which can withstand higher than atmospheric pressures) under pressure. On a routine basis this has been found useful for purifying

proteins (including enzymes) and polypeptides after enzymic digestion of proteins or chemical cleavage (e.g. with CNBr) prior to sequencing (using reverse-phase columns such as μ -Bondapak C18). Moderate pressures (50-300psi) have been found most satisfactory for large molecules (FPLC). [See *Scopes Anal Biochem* **114** 8 1981; *High Performance Liquid Chromatography and Its Application to Protein Chemistry*, Hearn in *Advances in Chromatography*, **20** 7 1982; B. A. Bidlingmeyer *Practical HPLC Methodology and Applications*, J Wiley & Sons, NY 1991; L.R.Snyder, J.L.Glajch and J.J.Kirkland *Practical HPLC Method Development*, J Wiley & Sons, NY 1988; ISBN 0471627828; R.W.A.Oliver, *HPLC of Macromolecules: A Practical Approach*, 2nd Edn, Oxford University Press, 1998, T.Hanai, *HPLC: A Practical Guide*, Royal Society of Chemistry (UK), 1999, ISBN 084045155; P.Millner *High Resolution Chromatography*, Oxford University Press, 1999 ISBN 0199636486; see also Chapter 1, bibliography.]

8. *Ultrafiltration* using a filter (e.g. Millipore) can remove water and low-molecular weight substances without the application of heat. Filters with a variety of molecular weight exclusion limits not only allow the concentration of a particular macromolecule to be determined, but also the removal (by washing during filtration) of smaller molecular weight contaminants (e.g. salts, inhibitors or cofactors). This procedure has been useful for changing the buffer in which the macromolecule is present (e.g. from Tris-Cl to ammonium carbonate), and for desalting. Ultrafiltration can be carried out in a stirrer cell (Amicon) in which the buffer containing the macromolecule (particularly protein) is pressed through the filter, with stirring, under argon or nitrogen pressure (e.g. 20-60psi). During this filtration process the buffer can be changed. This is rapid (e.g. 2L of solution can be concentrated to a few mLs in 1 to 2h depending on pressure and filter). A similar application uses a filter in a specially designed tube (Centricon tubes, Amicon) and the filtration occurs under centrifugal force in a centrifuge (4-6000rpm at 0°/40min). The macromolecule (usually DNA) then rests on the filter and can be washed on the filter by centrifugation. The macromolecule is recovered by inverting the filter, placing a conical receiver tube on the same side where the macromolecule rests, filling the other side of the filter tube with eluting solution (usually a very small volume e.g. 100 μ L), and during further centrifugation this solution passes through the filter and collects the macromolecule from the underside into the conical receiver tube.
9. *Partial precipitation* of a protein in solution can often be achieved by controlled addition of a strong salt solution, e.g. ammonium sulfate. This is commonly the first step in the purification process. Its simplicity is offset by possible denaturation of the desired protein and the (sometimes gross) contamination with other proteins. It should therefore be carried out by careful addition of small aliquots of the powdered salt or concentrated solution (below 4°, with gentle stirring) and allowing the salt to be evenly distributed in the solution before adding another small aliquot. Under carefully controlled conditions and using almost pure protein it is sometimes possible to obtain the protein in crystalline form suitable for X-ray analysis (see below).
10. *Dialysis*. This is a process by which small molecules, e.g. ammonium sulfate, sodium chloride, are removed from a solution containing the protein or DNA using a membrane which is porous to small molecules. The solution (e.g. 10mL) is placed in a dialysis bag or tube tied at both ends, and stirred in a large excess of dialysing solution (e.g. 1.5 to 2 L), usually a weak buffer at ca 4°. The dialysing buffer is replaced with fresh buffer several times, e.g. four times in 24h. This procedure is similar to ultrafiltration (above) and allows the replacement of buffer in which the protein, or DNA, is dissolved. It is also possible to concentrate the solutions by placing the dialysis tube or bag in Sephadex G25 which allows the passage of water and salts from the inside of the bag thus concentrating the protein (or DNA) solution. Dialysis tubing is available from various distributors but "Spectra/por" tubing (from Spectrum Medical Industries, Inc, LA) is particularly effective because it retains macromolecules and allows small molecules to dialyse out very rapidly thus reducing dialysing time considerably. This procedure is used when the buffer has to be changed so as to be compatible with the next purification or storage step, e.g. when the protein (or DNA) needs to be stored frozen in a particular buffer for extended periods.
11. *Gel Electrophoresis*. This is becoming a more commonly used procedure for purifying proteins, nucleic acids, nucleoproteins, polysaccharides and carbohydrates. The gels can be electroblotted onto membranes and the modern procedures of identifying, sequencing (proteins and nucleic acids) and amplifying (nucleic acids) on sub-micro scales have made this technique of separation a very important one. See below for polyacrylamide gel electrophoresis (PAGE), [D.Patel *Gel Electrophoresis*, J.Wiley-Liss, Inc., 1994; P.Jones and D.Rickwood, *Gel Electrophoresis: Nucleic Acids*, J.Wiley and Sons, 1999 (paperback) ISBN 0471960438; D.M.Gersten and D.Gersten, *Gel Electrophoresis: Proteins*, J.Wiley and Sons Inc, 1996 ISBN 0471962651; R.Westermeier *Electrophoresis in Practice*, 3rd Edn, Wiley-VCH, NY, 2001, ISBN 3527303006].
12. *Crystallisation*. The ultimate in purification of proteins or nucleic acids is crystallisation. This involves very specialised procedures and techniques and is best left to the experts in the field of X-ray crystallography who provide a complete picture of the structure of these large molecules. [A. Ducruix and R. Giegé eds, *Crystallisation of Nucleic Acids and Proteins: A Practical Approach*, 2nd Edition, 2000,

Oxford University Press, ISBN 0199636788 (paperback); T.L.Blundell and L.N.Johnson *Protein Crystallisation*, Academic Press, NY, 1976; A.McPherson *Preparation and Analysis of Protein Crystals*, J.Wiley & Sons, NY, 1982; A.McPherson, *Crystallisation of Biological Macromolecules*, Cold Spring Harbour Laboratory Press, 2001 ISBN 0879696176.]

Other details of the above will be found in Chapters 1 and 2 which also contain relevant references.

Several illustrations of the usefulness of the above methods are given in the *Methods Enzymol* series (Academic Press) in which 1000-fold purifications or more, have been readily achieved. In applying these sensitive methods to macromolecules, reagent purity is essential. It is disconcerting, therefore, to find that some commercial samples of the widely used affinity chromatography ligand Cibacron Blue F3GA contained this dye only as a minor constituent. The major component appeared to be the dichlorotriazinyl precursor of this dye. Commercial samples of Procion Blue and Procion Blue MX-R were also highly heterogeneous [Hanggi and Cadd *Anal Biochem* **149** 91 1985]. Variations in composition of sample dyes can well account for differences in results reported by different workers. The purity of substances of biological origin should therefore be checked by one or more of the methods given above. Water of high purity should be used in all operations. Double glass distilled water or water purified by a MilliQ filtration system (see Chapter 2) is most satisfactory.

Brief general procedures for the purification of polypeptides and proteins. Polypeptides of up to *ca* 1-2000 (10-20 amino acid residues) are best purified by reverse phase HPLC. The desired fractions that are collected are either precipitated from solution with EtOH or lyophilised. The purity can be checked by HPLC and identified by microsequencing (1-30 picomoles) to ascertain that the correct polypeptide was in hand. Polypeptides larger than these are sometimes classified as proteins, and are purified by one or more of the procedures described above. The purification of enzymes and functional proteins which can be identified by specific interactions is generally easier to follow because enzyme activities or specific protein interactions can be checked (by assaying) after each purification step. The commonly used procedures for purifying soluble proteins involve the isolation of an aqueous extract from homogenised tissues or extracts from ruptured cells from microorganisms or specifically cultured cells, for example, by sonication, freeze shocking or passage through a small orifice under pressure. Contaminating nucleic acids are removed by precipitation with a basic protein, e.g. protamine sulfate. The soluble supernatant is then subjected to fractionation with increasing concentrations of ammonium sulfate. The required fractions are then further purified by the procedures described in sections 2-9 above. If an affinity adsorbant has been identified then affinity chromatography can provide an almost pure protein in one step sometimes even from the crude extract. The rule of thumb is that a solution with a protein concentration of 1mg/mL has an absorbance A_{1cm} at 280nm of 1.0 units. Membrane-bound proteins are usually insoluble in water or dilute aqueous buffer and are obtained from the insoluble fractions, e.g. the microsomal fractions from the $>100,000 \times g$ ultracentrifugation supernatant. These are solubilised in appropriate detergents, e.g. Mega-10 (nonionic), Triton X-100 (ionic) detergents, and purified by methods 2 to 8 (previous section) in the presence of detergent in the buffer used. They are assayed also in the presence of detergent or membrane lipids.

The purity of proteins is best checked by *polyacrylamide gel electrophoresis* (PAGE). The gels are either made or purchased as pre-cast gels and can be with uniform or gradient gel composition. Proteins are applied onto the gels *via* wells set into the gels or by means of a comb, and travel along the gel surface by means of the current applied to the gel. When the buffer used contains sodium dodecylsulfate (SDS) the proteins are denatured and the denatured proteins (e.g. as protein subunits) separate on the gels mainly according to their molecular sizes. These can be identified by running marker proteins, with a range of molecular weights, simultaneously on a track alongside the proteins under study. The protein bands are visualised by fixing the gel (20% acetic acid) and staining with Coomassie blue followed by silver staining if higher sensitivity is required. An Amersham-Pharmacia "Phast Gel Electrophoresis" apparatus is very useful for rapid analysis of proteins. It uses small pre-cast polyacrylamide gels (two gels can be run simultaneously) with various uniform or gradient polyacrylamide concentrations as well as gels for isoelectric focusing. The gels are usually run for 0.5-1h and can be stained and developed (1-1.5h) in the same apparatus. The equipment can be used to electroblot the protein bands onto a membrane from which the proteins can be isolated and sequenced or subjected to antibody or other identification procedures. It should be noted that all purification procedures are almost always carried out at *ca* 4° in order to avoid denaturation or inactivation of the protein being investigated. Anyone contemplating the purification of a protein is referred to: Professor R.K.Scopes's monograph *Protein Purification*, 3rd edn, Springer-Verlag, New

York, 1994, ISBN 0387940723; M.L.Ladisch ed. *Protein Purification - from Molecular Mechanisms to Large-scale Processes*, American Chemical Society, Washington DC, 1990; E.L.V.Harris and S.Angal, *Protein Purification Applications - A Practical Approach*, IRL Press, Oxford, 1990; J.C.Janson and L.Rydén, *Protein Purification - Principles, High Resolution Methods and Applications*, VCH Publ. Inc., 1989; ISBN 0895731223 R.Burgess, *Protein Purification - Micro to Macro*, A.R.Liss, Inc., NY, 1987; S.M.Wheelwright, *Protein Purification: Design and Scale up of Downstream Processing*, J Wiley & Sons, NY, 1994, references in the bibliography in Chapter 1, and selected volumes of *Methods Enzymol*, e.g. M.P.Deutscher ed. Guide to Protein Purification, *Methods Enzymol*, Academic Press, NY, Vol 182 1990, ISBN 0121820831; T.Palzkil, *Proteomics*, Kluwer Academic Publ, 2001, ISBN 0792375653; M.A.Vijayalakshmi, *Biochromatography, Theory and Practice*, Taylot & Francis Publ, 2002, ISBN 0415269032; J.S.Davies, *Amino Acids, Peptides and Proteins Vol 32 2001*, A Specialist Periodical Report, Royal Society of Chemistry, ISBN 0854042326; S.Roe, *Protein Purification Techniques: A Practical Approach*, 2nd Edn, Oxford University Press, 2001, ISBN 0199636737; T.Palmer, *Enzymes, Biochemistry, Biotechnology, Clinical Chemistry*, Horwood Publishing, 2001, ISBN 1898563780.

Brief general procedures for purifying DNA. Oligo-deoxyribonucleotides (up to *ca* 60-mers) are conveniently purified by HPLC (e.g. using a Bio-Rad MA7Q anion exchange column and a Rainin Instrument Co, Madison, Dynamax-300A C₈ matrix column) and used for a variety of molecular biology experiments. Plasmid and chromosomal DNA can be isolated by centrifugation in caesium chloride buffer (see section 1. centrifugation above), and then re-precipitated with 70% ethanol at -70° (18h), collected by centrifugation (microfuge) and dried in air before dissolving in TE (10mM TrisHCl, 1mM EDTA pH 8.0). The DNA is identified on an Agarose gel slab (0.5 to 1.0% DNA grade in 45mM Tris-borate + 1mM EDTA or 40mM Tris-acetate + 1mM EDTA pH 8.0 buffers) containing ethidium bromide which binds to the DNA and under UV light causes it be visualised as pink fluorescent bands. Marker DNA (from λ phage DNA cut with the restriction enzymes Hind III and/or EcoRI) are in a parallel track in order to estimate the size of the unknown DNA. The DNA can be isolated from their band on the gel by transfer onto a nitro-acetate paper (e.g. NA 45) electrophoretically, by binding to silica or an ion exchange resin, extracted from these adsorbents and precipitated with ethanol. The DNA pellet is then dissolved in TE buffer and its concentration determined. A solution of duplex DNA (or RNA) of 50 μ g/mL gives an absorbance of 1.0units at 260nm/1cm cuvette (single stranded DNA or RNA gives a value of 1.3 absorbance units). DNA obtained in this way is suitable for molecular cloning. For experimental details on the isolation, purification and manipulation of DNA and RNA the reader is referred to: J.Sambrook, E.F.Fritsch and T.Maniatis, *Molecular Cloning - A Laboratory Manual* , 2nd edn, (3 volumes), Cold Spring Harbor Laboratory Press, NY, 1989, ISBN 0879693096 (paperback); P.D.Darbre, *Basic Molecular Biology: Essential Techniques*, J.Wiley and Sons, 1998, ISBN 0471977055; J.Sambrook and D.W.Russell, *Molecular Cloning - A Laboratory Manual* , 3rd edn, (3 volumes), Cold Spring Harbor Laboratory Press, NY, 2001, ISBN 0079695773 (paperback), ISBN 0079695765 (cloth bound), also available on line; M.A.Vijayalakshmi, *Biochromatography, Theory and Practice*, Taylot & Francis Publ, 2002, ISBN 0415269032; A.Travers and M.Buckle, *DNA-Protein Interactions: A Practical Approach*, Oxford University Press, 2000, ISBN 0199636915 (paperback); R.Rapley and D.L.Manning eds *RNA: Isolation and Characterisation Protocols*, Humana Press 1998 ISBN 086034941; R.Rapley, *The Nucleic Acid Protocols Handbook*, Humana Press 2000 ISBN 0896038416 (paperback).

This chapter lists some representative examples of biochemicals and their origins, a brief indication of key techniques used in their purification, and literature references where further details may be found. Simpler low molecular weight compounds, particularly those that may have been prepared by chemical syntheses, e.g. acetic acid, glycine, will be found in Chapter 4. Only a small number of enzymes and proteins are included because of space limitations. The purification of some of the ones that have been included has been described only briefly. The reader is referred to comprehensive texts such as the *Methods Enzymol* (Academic Press) series which currently runs to more than 344 volumes and *The Enzymes* (3rd Edn, Academic Press) which runs to 22 volumes for methods of preparation and purification of proteins and enzymes. Leading references on proteins will be found in *Advances in Protein Chemistry* (59 volumes, Academic Press) and on enzymes will be found in *Advances in Enzymology* (72 volumes, then became *Advances in Enzymology and Related Area of Molecular Biology*, J Wiley & Sons). The *Annual Review of Biochemistry* (Annual Review Inc. Patlo Alto California) also is an excellent source of key references to the up-to-date information on known and new natural compounds, from small molecules, e.g. enzyme cofactors to proteins and nucleic acids.

Abbreviations of titles of periodical are defined as in the Chemical Abstracts Service Source Index (CASSI).

Ionisation constants of ionisable compounds are given as **pK** values (published from the literature) and refer to the **pKa** values at room temperature (~ 15°C to 25°C). The values at other temperatures are given as superscripts, e.g. **pK²⁵** for 25°C. Estimated values are entered as **pK_{Est(1)}**~ (see Chapter 1, p 6 for further information).

Benzene, which has been used as a solvent successfully and extensively in the past for reactions and purification by chromatography and crystallisation is now considered a **very dangerous substance** so it has to be used with extreme care. We emphasise that an alternative solvent to benzene (e.g. toluene, toluene-petroleum ether, or a petroleum ether to name a few) should be used first. However, if benzene has to be used then all operations have to be performed in a well ventilated fumehood and precautions taken to avoid inhalation and contact with skin and eyes. Whenever benzene is mentioned in the text and asterisk e.g. *C₆H₆ or *benzene, is inserted to remind the user that special precaution should be adopted.

Amino acids, carbohydrates and steroids not found below are in Chapter 4 (see also CAS Registry Numbers Index and General Index).

Abrin A and Abrin B [1393-62-0] **M_r 63,000-67,000**. Toxic proteins from seeds of *Abras precatorius*. Purified by successive chromatography on DEAE-Sephadex A-50, carboxymethylcellulose, and DEAE-cellulose. [Wei et al. *J Biol Chem* **249** 3061 1974.]

Acetoacetyl coenzyme A trisodium salt trihydrate [102029-52-7] **M 955.6, pK₁ 4.0 (NH₂), pK₂ 6.4 (PO₄⁻)**. The pH of solution (0.05g/mL H₂O) is adjusted to 5 with 2N NaOH. This solution can be stored frozen for several weeks. Further purification can be carried out on a DEAE-cellulose formate column, then through a Dowex 50 (H⁺) column to remove Na ions, concentrated by lyophilisation and redissolved in H₂O. Available as a soln of 0.05g/mL of H₂O. The concn of acetoacetylcoenzyme A is determined by the method of Stern et al. *J Biol Chem* **221** 15 1956. It is stable at pH 7-7.5 for several hours at 0° (half life *ca* 1-2h). At room temperature it is hydrolysed in *ca* 1-2h at pH 7-7.5. At pH 1.0/20° it is more stable than at neutrality. It is stable at pH 2-3/-17° for at least 6 months. [*J Biol Chem* **159** 1961 1964; **242** 3468 1967; Clikensbeard et al. *J Biol Chem* **250** 3108 1975; *J Am Chem Soc* **75** 2520 1953, **81** 1265 1959; see Simon and Shemin *J Am Chem Soc* **75** 2520 1953; Salem et al. *Biochem J* **258** 563 1989.]

Acetobromo- α -D-galactose [3068-32-4] **M 411.2, m 87°, [α]₅₄₆²⁰ +255°, [α]_D²⁰ +210° (c 3, CHCl₃)**. Purified as for the glucose analogue (see next entry). If the compound melts lower than 87° or is highly coloured then dissolve in CHCl₃ (*ca* 3 vols) and extract with H₂O (2 vols), 5% aqueous NaHCO₃, and again with H₂O and dry over Na₂SO₄. Filter and evaporate in a vacuum. The partially crystalline solid or syrup is dissolved in dry Et₂O (must be very dry) and recrystd by adding pet ether (b 40-60°) to give a white product. [McKellan and Horecker *Biochem Prep* **11** 111 1960.]

Acetobromo- α -D-glucose [572-09-8] **M 411.2, m 87-88°, 88-89°, [α]₅₄₆²⁰ +230°, [α]_D²⁰ +195° (c 3, CHCl₃)**. If nicely crystalline recryst from Et₂O-pentane. Alternatively dissolve in diisopropyl ether (dried over CaCl₂ for 24hours, then over P₂O₅ for 24hours) by shaking and warming (for as short a period as possible), filter warm. Cool to *ca* 45° then slowly to room temperature and finally at 5° for more than 2hours. Collect the solid, wash with cold dry diisopropyl ether and dry in a vacuum over Ca(OH)₂ and NaOH. Store dry in a desiccator in the dark. Solutions can be stabilised with 2% CaCO₃. [Redemann and Niemann *Org Synth* **65** 236 1987, Coll Vol III 11 1955.]

Acetoin dehydrogenase [from beef liver; acetoin NAD oxidoreductase] [9028-49-3] **M_r 76000, [EC 1.1.1.5]**. Purified *via* the acetone cake then Ca-phosphate gel filtration (unabsorbed), lyophilised and then fractionated through a DEAE-22 cellulose column. The Km for diacetyl in 40 μ M and for

NADH it is 100 μ M in phosphate buffer at pH 6.1. [Burgos and Martin *Biochim Biophys Acta* **268** 261 1972; **289** 13 1972.]

(-)-3- β -Acetoxy-5-etiolic acid [**3- β -acetoxy-5-etiocholenic acid, androst-5-ene-17- β -carboxylic acid**] [51424-66-9] **M 306.5, m 238-240 $^{\circ}$, 241-242 $^{\circ}$, 243-245 $^{\circ}$, 246-247 $^{\circ}$, $[\alpha]_{\text{D}}^{20}$ -19.9 $^{\circ}$ (c 1, Me₂CO), -36 $^{\circ}$ (c 1, Dioxane), -33.5 $^{\circ}$ (CHCl₃), pK_{Est} ~ 4.7.** It is purified by recrystn from Me₂CO, Et₂O-pentane, or AcOH, and dried in a vacuum oven (105 $^{\circ}$ /20mm) and sublimed at high vacuum. [Staunton and Eisenbram *Org Synth* **42** 4 1962; Steiger and Reichstein *Helv Chim Acta* **20** 1404 1937.]

Acetylcarnitine chloride (2-acetoxy-3-carboxy-*N,N,N*-trimethylpropanamine HCl) [*S*(D+)-5080-50-2; *R*(L-)-5061-35-8; *RS* 2504-11-2] **M 239.7, m 181 $^{\circ}$, 197 $^{\circ}$ (dec), $[\alpha]_{\text{D}}^{25}$ -28 $^{\circ}$ (c 2, H₂O) for *S*-isomer, pK²⁵ 3.6.** Recrystd from isopropanol. Dried over P₂O₅ under high vacuum.

Acetylcholine bromide [66-23-9] **M 226.1, m 143 $^{\circ}$, 146 $^{\circ}$.** *Hygroscopic* solid but less than the hydrochloride salt. It crystd from EtOH as prisms. Some hydrolysis occurs in boiling EtOH particularly if it contains some H₂O. It can also be recryst from EtOH or MeOH by adding dry Et₂O. [*Acta Chem Scand* **12** 1492, 1497, 1502 1958.]

Acetylcholine chloride [60-31-1] **M 181.7, m 148-150 $^{\circ}$, 151 $^{\circ}$.** It is very sol in H₂O (> 10%), and is very *hygroscopic*. If pasty, dry in a vacuum desiccator over H₂SO₄ until a solid residue is obtained. Dissolve in abs EtOH, filter and add dry Et₂O and the hydrochloride separates. Collect by filtration and store under very dry conditions. [*J Am Chem Soc* **52** 310 1930.] The *chloroplatinate* crystallises from hot H₂O in yellow needles and can be recrystd from 50% EtOH, m 242-244 $^{\circ}$ [*Biochem J* **23** 1069 1929], other m given is 256-257 $^{\circ}$. The *perchlorate* crystallises from EtOH as prisms m 116-117 $^{\circ}$. [*J Am Pharm Assocn* **36** 272 1947.]

***N*⁴-Acetylcytosine** [14631-20-0] **M 153.1, m >300 $^{\circ}$, 326-328 $^{\circ}$, pK_{Est(1)} ~1.7, pK_{Est(2)} ~10.0.** If TLC or paper chromatography show that it contains unacetylated cytosine then reflux in Ac₂O for 4h, cool at 3-4 $^{\circ}$ for a few days, collect the crystals, wash with cold H₂O, then EtOH and dry at 100 $^{\circ}$. It is insoluble in EtOH and difficulty soluble in H₂O but crystallises in prisms from hot H₂O. It is hydrolysed by 80% aq AcOH at 100 $^{\circ}$ /1h. [*Am Chem J* **29** 500 1903; UV: *J Chem Soc* 2384 1956; *J Am Chem Soc* **80** 5164 1958.] It forms an Hg salt [*J Am Chem Soc* **79** 5060 1957].

β -D-*N*-Acetylglucosaminidase [from *M sexta* insects] [9012-33-3] **M_r ~61,000, [EC 3.2.1.52].** Purified by chromatography on DEAD-Biogel, hydroxylapatite chromatography and gel filtration through Sephacryl S200. Two isoforms: a hexosaminidase EI with Km 177 μ M (*V*_{max} 328 sec⁻¹) and EII a chitinase with Km 160 μ M (*V*_{max} 103 sec⁻¹) with 4-nitrophenyl- β -acetylglucosamine as substrate. [Dziadil-Turner *Arch Biochem Biophys* **212** 546 1981.]

β -D-*N*-Acetylhexosaminidase A and B (from human placenta) [9012-33-3] **M_r ~61,000, [EC 3.2.1.52].** Purified by Sephadex G-200 filtration and DEAE-cellulose column chromatography. Hexosaminidase A was further purified by DEAE-cellulose column chromatography, followed by an ECTEOLA-cellulose column, Sephadex-200 filtration, electrofocusing and Sephadex G-200 filtration. Hexosaminidase B was purified by a CM-cellulose column, electrofocusing and Sephadex G-200 filtration. [Srivastava et al. *J Biol Chem* **249** 2034 1974.]

***N*-Acetyl-D-lactosamine** [2-acetyl-amino-*O*- β -D-lactopyranosyl-2-deoxy-D-glucose] [32181-59-2] **M 383.4, m 169-171 $^{\circ}$, 170-171 $^{\circ}$, $[\alpha]_{\text{D}}^{18}$ +51.5 $^{\circ}$ \rightarrow +28.8 $^{\circ}$ (in 3h, c 1, H₂O).** Purified by recrystn from MeOH (with 1 mol of MeOH) or from H₂O. It is available as a soln of 0.5g /mL of H₂O. [Zilliken *J Biol Chem* **271** 181 1955.]

***O*-Acetyl- β -methylcholine chloride** [Methacholine chloride, Amechol, Provocholine, 2-acetoxypropyl-ammonium chloride] [62-51-1] **M 195.7, m 170-173 $^{\circ}$, 172-173 $^{\circ}$.** It forms white *hygroscopic* needles from Et₂O and is soluble in H₂O, EtOH and CHCl₃. It decomposes readily in alkaline solns and slowly in H₂O. It should be handled and stored in a dry atmosphere. The *bromide* is less

hygroscopic and the *picrate* has *m* 129.5-131° (from EtOH). [racemate: Annis and Ely *Biochem J* 53 34 1953; IR of iodide: Hansen *Acta Chem Scand* 13 155 1959.]

***N*-Acetyl muramic acid** [NAMA, *R*-2-(acetylamino)-3-*O*-(1-carboxyethyl)-2-deoxy-D-glucose] [10597-89-4] *M* 292.3, *m* ~125°(dec), $[\alpha]_{\text{D}}^{20} +41.2^\circ$ (c 1.5, H₂O, after 24h), *pK*_{Est} ~ 3.6. See muramic acid below.

***N*-Acetyl neuraminic acid (NANA, *O*-Sialic acid, 5-acetamido-3,5-dideoxy-D-glycero-D-glacto-2-nonulosonic acid, lactaminic acid)** [131-48-6] *M* 309.3, *m* 159°(dec), 181-183°(dec), 185-187°(dec), $[\alpha]_{\text{D}}^{25} -33^\circ$ (c 2, H₂O, l 2), *pK* 2.6. A Dowex-1x8 (200-400 mesh) in the formate form was used, and was prepd by washing with 0.1M NaOH, then 2N sodium formate, excess formate was removed by washing with H₂O. *N*-Acetyl neuraminic acid in H₂O is applied to this column, washed with H₂O, then eluted with 2N formic acid at a flow rate of 1mL/min. Fractions (20mL) were collected and tested (Bial's orcinol reagent, *cf Biochem Prep* 7 1 1959). NANA eluted at formic acid molarity of 0.38 and the Bial positive fractions are collected and lyophilised. The residue is recrystd from aqueous AcOH: Suspend 1.35g of residue in AcOH, heat rapidly to boiling, add H₂O dropwise until the suspension dissolves (do not add excess H₂O, filter hot and then keep at +5° for several hours until crystn is complete. Collect and dry in a vacuum over P₂O₅. Alternatively dissolve 1.35g of NANA in 14mL of H₂O, filter, add 160mL of MeOH followed by 360mL of Et₂O. Then add pet ether (b 40-60°) until heavy turbidity. Cool at 20° overnight. Yield of NANA is *ca* 1.3g. Dry over P₂O₅ at 1mm vacuum and 100° to constant weight. It mutarotates in Me₂SO: $[\alpha]_{\text{D}}^{20} -115^\circ$ (after 7min) to -32° (after 24h). It is available as aqueous soln (0.01g/mL). [IR and synthesis: Cornforth et al. *Biochem J* 68 57 1958; Zillikin and O'Brien *Biochem Prep* 7 1 1960; ¹³C NMR and 1-¹³C synthesis: Nguyen, Perry *J Org Chem* 43 551 1978; Danishevski, DeNinno *J Org Chem* 51 2615 1986; Gottschalk, *The Chemistry and Biology of Sialic Acids and Related Substances*, Cambridge University Press, London, 1960.]

***N*-Acetyl neuraminic acid aldolase [from *Clostridium perfringens*, *N*-acetylneuraminic acid pyruvate lyase]** [9027-60-5] *M*_r 32,000 [EC 4.1.3.3]. Purified by extraction with H₂O, protamine pptn, (NH₄)₂SO₄ pptn, Me₂CO pptn, acid treatment at pH 5.7 and pptn at pH 4.5. The equilibrium constant for pyruvate + *n*-acetyl-D-mannosamine \rightleftharpoons *N*-acetylneuraminidate at 37° is 0.64. The *K*_m for *N*-acetylneuraminic acid is 3.9mM in phosphate at pH 7.2 and 37°. [Comb and Roseman *Methods Enzymol* 5 391 1962.] The enzyme from Hogg kidney (cortex) has been purified 1700 fold by extraction with H₂O, protamine sulfate pptn, (NH₄)₂SO₄ pptn, heating between 60-80°, a second (NH₄)₂SO₄ pptn and starch gel electrophoresis. The *K*_m for *N*-acetylneuraminic acid is 1.5mM. [Brunetti et al. *J Biol Chem* 237 2447 1962.]

***N*-Acetyl penicillamine** [D- 15537-71-0, DL-59-53-0] *M* 191.3, *m* 183°, 186-187° (DL-form), 189-190° (D-form), D-form $[\alpha]_{\text{D}}^{25} +18^\circ$ (c 1, 50% EtOH), *pK*_{Est(1)}~3.0 (CO₂H), *pK*_{Est(2)}~ 8.0 (SH). Both forms are recrystd from hot H₂O. A pure sample of the D-form was obtained after five recrystns. [Crooks in *The Chemistry of Penicillin* Clarke, Johnson and Robinson eds, Princeton University Press, 470 1949.]

***p*-Acetylphenyl sulfate potassium salt**, [38533-41-4] *M* 254.3, *m* dec on heating, *pK*_{Est} ~2,1. Purified by dissolving in the minimum vol of hot water (60°) and adding EtOH, with stirring, then left at 0° for 1h. Crystals were filt'd off and recrystd from H₂O until free of Cl⁻ and SO₄²⁻ ions. Dried in a vac over P₂O₅ at room temperature. It is a specific substrate for arylsulfatases which hydrolyse it to *p*-acetylphenol [λ_{max} 327nm (ϵ 21700 M⁻¹cm⁻¹)] [Milsom et al. *Biochem J* 128 331 1972].

***S*-Acetylthiocholine bromide** [25025-59-6] *M* 242.2, *m* 217-223°(dec). It is a *hygroscopic* solid which can be recrystd from ligroin-EtOH (1:1), dried and kept in a vacuum desiccator. Crystn from *C₆H₆-EtOH gave *m* 227° or from propan-1-ol the *m* was 213°. [*Acta Chem Scand* 11 537 1957, 12 1481 1958.]

***S*-Acetylthiocholine chloride** [6050-81-3] *M* 197.7, *m* 172-173° The chloride can be purified in the same way as the bromide, and it can be prepared from the iodide. A few milligrams dissolved in H₂O can be purified by applying onto a Dowex-1 Cl⁻ resin column (prepared by washing with N HCl followed by CO₃²⁻—free H₂O until the pH is 5.8). After equilibration for 10min elution is started with CO₃²⁻—free distilled H₂O and

3mL fractions are collected and their OD at 229nm measured. The fractions with appreciable absorption are pooled and lyophilised at 0-5°. Note that at higher temps decomposition of the ester is appreciable; hydrolysis is appreciable at pH >10.5/20°. The residue is dried *in vacuo* over P₂O₅, checked for traces of iodine (conc H₂SO₄ and heat, violet vapours are released), and recrystd from propan-1-ol. [*Clin Chim Acta* 2 316 1957.]

S-Acetylthiocholine iodide [1866-15-5] **M 289.2, m 203-204°, 204°, 204-205°**. Recrystd from propan-1-ol (or *iso*-PrOH, or EtOH/Et₂O) until almost colourless and dried in a vacuum desiccator over P₂O₅. Solubility in H₂O is 1% w/v. A 0.075M (21.7mg/mL) solution in 0.1M phosphate buffer pH 8.0 is stable for 10-15 days if kept refrigerated. Store away from light. It is available as a 1% soln in H₂O. [*Biochemical Pharmacology* 7, 88 1961; IR: Hansen *Acta Chem Scand* 13 151 1959, 11 537 1957; *Clin Chim Acta* 2 316 1957; *Zh Obshch Khim* 22 267 1952.]

Actinomycin C (Cactinomycin) [8052-16-2] **M ~1255**. (A commercial mixture of Actinomycin C₁ ~5%, C₂ ~30% and C₃ ~65%). *Actinomycin C₁ (native)* crystals from EtOAc as red crystals, is sol in CHCl₃, *C₆H₆ and Me₂CO and has **m 246-247°(dec)**, $[\alpha]_D^{20}$ -328° (0.22, MeOH) and λ_{\max} 443nm (ϵ 25,000) and 240nm (ϵ 34,000). *Actinomycin C₂ (native)* crystals as red needles from EtOAc and has **m 244-246°(dec)**, $[\alpha]_D^{20}$ -325° (c 0.2, MeOH), λ_{\max} 443nm (ϵ 25,300) and (ϵ 33,400). *Actinomycin C₃ (native)* recryst from cyclohexane, or *C₆H₆/MeOH/cyclohexane as red needles **m 238-241° (dec)**, $[\alpha]_D^{20}$ -321° (c 0.2, MeOH), λ_{\max} 443nm (ϵ 25,000) and 240nm (ϵ 33,300). [Brockman and Lackner, *Chem Ber* 101 1312 1968.] It is *light sensitive*.

Actinomycin D (Dactinomycin) [50-76-0] **M 1255.5, m 241-243°(dec)**, $[\alpha]_D^{22}$ -296° (c 0.22, MeOH). Crystallises as bright red rhombic crystals from absolute EtOH or from MeOH-EtOH (1:3). It will also crystallise from EtOAc-cyclohexane (**m 246-247° dec**), CHCl₃-pet ether (**m 245-246° dec**), and EtOAc-MeOH-*C₆H₆ (**m 241-243° dec**). Its solubility in MeCN is 1mg/mL. $[\alpha]_D^{20}$ varies from -296° to -327° (c 0.2, MeOH). λ_{\max} (MeOH) 445, 240nm (log ϵ 4.43, 4.49), λ_{\max} (MeOH, 10N HCl, 1:1) 477nm (log ϵ 4.21) and λ_{\max} (MeOH, 0.1N NaOH) 458, 344, 285 (log ϵ 3.05, 4.28, 4.13). It is **HIGHLY TOXIC**, light sensitive and antineoplastic. [Bullock and Johnson, *J Chem Soc* 3280 1957.]

Acyl-coenzyme A Synthase [from beef liver] [9013-18-7] **M_r 57,000, [EC 6.2.1.2]**. Purified by extraction with sucrose-HCO₃ buffer, protamine sulfate pptn, (NH₄)₂SO₄ (66-65%) pptn at pH 4.35 and a second (NH₄)₂SO₄ (35-60%) pptn at pH 4.35. It has Km 0.15mM (*V_{rel}* 1.0) for octanoate; 0.41mM (*V_{rel}* 2.37) for heptanoate and 1.59mM (*V_{rel}* 0.63). Km for ATP is 0.5mM all at pH 9.0 in ethylene glycol buffer at 38°. [Jencks et al. *J Biol Chem* 204 453 1953; *Methods Enzymol* 5 467 1962.]

Acyl-coenzyme A Synthase (from yeast) [9012-31-1] [EC 6.2.1.1]. This enzyme has been purified by extraction into phosphate buffer pH 6.8-7.0 containing 2-mercaptoethanol and EDTA, protamine sulfate pptn, polyethylene glycol fractionation, Alumina γ gel filtration, concentration by (NH₄)₂SO₄ pptn, Bio-Gel A-0.5m chromatography and DEAE-cellulose gradient chromatography. It has M_r ~151,000, Km (apparent) 0.24mM (for acetate) and 0.035mM (for CoA); 1.2 mM (for ATP) and Mg²⁺ 4.0mM. [Frenkel and Kitchens *Methods Enzymol* 71 317 1981.]

Adenosine-5'-diphosphate [adenosine-5'-pyrophosphate, ADP] [58-64-0] **M 427.2, $[\alpha]_D^{25}$ -25.7° (c 2, H₂O), pK₁²⁵ <2 (PO₄H), pK₂²⁵ <2 (PO₄H), pK₃²⁵ 3.95 (NH₂), pK₄²⁵ 6.26 (PO₄H)**. Characterised by conversion to the *acridine salt* by addition of alcoholic acridine (1.1g in 50mL), filtering off the yellow salt and recrystallising from H₂O. The salt has **m 215°(dec)**, λ_{\max} 259nm (ϵ 15,400) in H₂O. [Baddiley and Todd *J Chem Soc* 648 1947, 582 1949, cf LePage *Biochem Prep* 1 1 1949; Martell and Schwarzenbach *Helv Chim Acta* 39 653 1956].

Adenosine-3'-monophosphoric acid hydrate [3'-adenylic acid, 3'-AMP] [84-21-9] **M 347.3, m 197°(dec, as 2H₂O), 210°(dec), m 210°(dec), $[\alpha]_{546}$ -50° (c 0.5, 0.5M Na₂HPO₄), pK₁²⁵ 3.65, pK₂²⁵ 6.05**. It crystallises from large volumes of H₂O in needles as the monohydrate, but is not very soluble in boiling H₂O. Under acidic conditions it forms an equilibrium mixture of 2' and 3' adenylic acids *via* the 2',3'-cyclic phosphate. When heated with 20% HCl it gives a quantitative yield of furfural after 3hours, unlike 5'-adenylic acid which only gives traces of furfural. The yellow *monoacridine salt* has **m 175°(dec)** and

the *diacridine salt* has m 177° (225°)(dec). [Brown and Todd *J Chem Soc* 44 1952; Takaku et al. *Chem Pharm Bull Jpn* 21 1844 1973; NMR: Ts'O et al. *Biochemistry* 8 997 1969.]

Adenosine-5'-monophosphoric acid monohydrate [5'-adenylic acid, 5'-AMP] [18422-05-4] M 365.2, m 178°, 196-200°, 200° (sintering at 181°), $[\alpha]_D^{20}$ -47.5°, $[\alpha]_{546}$ -56° (c 2, in 2% NaOH), -26.0° (c 2, 10% HCl), -38° (c 1, 0.5M Na₂HPO₄), pK_1^{25} 3.89, pK_2^{25} 6.14, pK_3^{25} 13.1. It has been recrystd from H₂O (fine needles) and is freely soluble in boiling H₂O. Crysts also from H₂O by addition of acetone. Purified by chromatography on Dowex 1 (in formate form), eluting with 0.25M formic acid. It was then adsorbed onto charcoal (which had been boiled for 15min with M HCl, washed free of chloride and dried at 100°), and recovered by stirring three times with isoamyl alcohol/H₂O (1:9 v/v). The aqueous layer from the combined extracts was evaporated to dryness under reduced pressure, and the product was crystallised twice from hot H₂O. [Morrison and Doherty *Biochem J* 79 433 1961]. It has λ_{max} 259nm (ϵ 15,400) in H₂O at pH 7.0. [Alberty et al. *J Biol Chem* 193 425 1951; Martell and Schwarzenbach *Helv Chim Acta* 39 653 1956]. The *acridinium salt* has m 208° [Baddiley and Todd *J Chem Soc* 648 1947; Pettit *Synthetic Nucleotides*, van Nostrand-Reinhold, NY, Vol 1 252 1972; NMR: Sarma et al. *J Am Chem Soc* 96 7337 1974; Norton et al. *J Am Chem Soc* 98 1007 1976; IR of *diNa salt*: Miles *Biochem Biophys Acta* 27 324 1958].

Adenosine 5''-[β -thio]diphosphate tri-lithium salt [73536-95-5] M 461.1. Purified by ion-exchange chromatography on DEAE-Sephadex A-25 using gradient elution with 0.1-0.5M triethylammonium bicarbonate. [*Biochem Biophys Acta* 276 155 1972.]

Adenosine 5''-[α -thio]monophosphate di-lithium salt [19341-57-2] M 375.2. Purified as for the diNa salt [Murray and Atkinson *Biochemistry* 7 4023 1968]. Dissolve 0.3g in dry MeOH (7mL) and M LiI (6mL) in dry Me₂CO containing 1% of mercaptoethanol and the Li salt is ppted by adding Me₂CO (75mL). The residue is washed with Me₂CO (4 x 30mL) and dried at 55°/25mm. λ_{max} (HCl, pH 1.2) 257nm (ϵ 14,800); (0.015M NaOAc, pH 4.8) 259nm (ϵ 14,800); and (0.015M NH₄OH, pH 10.1) 259nm (ϵ 15,300).

Adenosine-5'-triphosphate (ATP) [56-65-5] M 507.2, $[\alpha]_{546}$ -35.5 (c 1, 0.5 M Na₂HPO₄), pK_1^{25} 4.00, pK_2^{25} 6.48. Ppted as its barium salt when excess barium acetate soln was added to a 5% soln of ATP in water. After filtering off, the ppte was washed with distd water, redissolved in 0.2M HNO₃, and again pptd with barium acetate. The ppte, after several washings with distd water, was dissolved in 0.2M HNO₃ and slightly more 0.2M H₂SO₄ than was needed to ppt all the barium as BaSO₄, was added. After filtering off the BaSO₄, the ATP was ppted by addition of a large excess of 95% ethanol, filtered off, washed several times with 100% EtOH and finally with dry diethyl ether. [Kashiwagi and Rabinovitch *J Phys Chem* 59 498 1955.]

S-(5'-Adenosyl)-L-homocysteine [979-92-0] M 384.4, m 202°(dec), 204°(dec), 205-207°(dec), $[\alpha]_D^{25}$ +93° (c 1, 0.2N HCl), $[\alpha]_D^{23}$ +44° (c 0.1, 0.05N HCl), (pK see SAM hydrochloride below). It has been recrystd several times from aqueous EtOH or H₂O to give small prisms and has λ_{max} 260nm in H₂O. The *picrate* has m 170°(dec) from H₂O. [Baddiley and Jameison *J Chem Soc* 1085 1955; de la Haba and Cantoni *J Biol Chem* 234 603 1959; Borchardt et al. *J Org Chem* 41 565 1976; NMR: Follmann et al. *Eur J Biochem* 47 187 1974.]

(-)-S-Adenosyl-L-methionine chloride (SAM hydrochloride) [24346-00-7] M 439.9, $pK_{Est(1)} \sim 2.13$, $pK_{Est(2)} \sim 4.12$, $pK_{Est(3)} \sim 9.28$. Purified by ion exchange on Amberlite IRC-150, and eluting with 0.1-4M HCl. [Stolowitz and Minch *J Am Chem Soc* 103 6015 1981.] It has been isolated as the tri-reineckate salt by adding 2 volumes of 1% solution of ammonium reineckate in 2% perchloric acid. The reineckate salt separates at once but is kept at 2° overnight. The salt is collected on a sintered glass funnel, washed with 0.5% of ammonium reineckate, dried (all operations at 2°) and stored at 2°. To obtain adenosylmethionine, the reineckate is dissolved in a small volume of methyl ethyl ketone and centrifuged at room temp to remove a small amount of solid. The clear dark red supernatant is extracted (in a separating funnel) with a slight excess of 0.1 N H₂SO₄. The aqueous phase is re-extracted with fresh methyl ethyl ketone until it is colourless. [Note that reineckates have UV absorption at 305nm (ϵ 15,000), and the optical density at 305nm is used to detect the presence of reineckate ions.] Methyl ethyl ketone is removed from the aqueous layer containing adenosylmethionine sulfate, the pH is adjusted to 5.6-6.0 and extracted with two volumes of Et₂O.

The *sulfate* is obtained by evaporating the aqueous layer in *vacuo*. The *hydrochloride* can be obtained in the same way but using HCl instead of H₂SO₄. SAM-HCl has a solubility of 10% in H₂O. The salts are stable in the cold at pH 4-6 but decompose in alkaline media. [Cantoni *Biochem Prep* 5 58 1957.] The purity of SAM can be determined by paper chromatography [Cantoni *J Biol Chem* 204 403 1953; *Methods Enzymol* 3 601 1957], and electrophoretic methods or enzymic analysis [Cantoni and Vignos *J Biol Chem* 209 647 1954].

L-Adrenaline [L-epinephrine, L-(-)-(3,4-dihydroxyphenyl)-2-methylaminoethanol] [51-43-4] **M** 183.2, **m** 210°(dec), 211°(dec), 211-212°(dec), 215°(dec), $[\alpha]_{\text{D}}^{25} -52^\circ$ (c 2, 5% HCl), **pK₁²⁵ 8.88**, **pK₂²⁵ 9.90**, **pK₃²⁵ 12.0**. It has been recrystd from EtOH + AcOH + NH₃ [Jensen *J Am Chem Soc* 57 1765 1935]. It is sparingly soluble in H₂O, readily in acidic or basic solns but insoluble in aqueous NH₃, alkali carbonate solns, EtOH, CHCl₃, Et₂O or Me₂CO. It is readily oxidised in air and turns brown on exposure to light and air. Store in the dark under N₂. Its pKa values in H₂O are 8.88 and 9.90 [Lewis *Br J Pharmacol Chemother* 9 488 1954]. The *hydrogen oxalate salt* has **m** 191-192°(dec, *evac capillary*) after recrystn from H₂O or EtOH [Pickholz *J Chem Soc* 928 1945].

Adrenolone hydrochloride [3',4'-dihydroxy-2-methylaminoacetophenone hydrochloride] [62-13-5] **M** 217.7, **m** 244-249°(dec), 248°(dec), 256°(dec), **pK** 5.5. It was purified by recrystn from EtOH or aqueous EtOH. [Gero *J Org Chem* 16 1222 1951; Kindler and Peschke *Arch Pharm* 269 581, 603 1931.]

ADP-Ribosyl transferase (from human placenta) [9026-30-6]. Purified by making an affinity absorbent for ADP-ribosyltransferase by coupling 3-aminobenzamide to Sepharose 4B. [Burtscher et al. *Anal Biochem* 152 285 1986.]

Agglutinin (from peanuts) [*Arachis hypogaea*] [1393-62-0] **M_r** 134,900 (tetramer). Purified by affinity chromatography on Sepharose- ζ -aminocaproyl- β -D-galactopyranosylamine. [Lotan et al. *J Biol Chem* 250 8518 1974.]

Alamethicin (from *Trichoderma viridae*). [27061-78-5] **M** 1964.3, **m** 259-260°, 275-270°, $[\alpha]_{\text{D}}^{22} -45^\circ$ (c 1.2, EtOH), **pK** 6.04 (aq EtOH). Recrystd from MeOH. [Panday et al. *J Am Chem Soc* 99 8469 1977.] The *acetate* [64918-47-4] has **m** 195-180° from MeOH/Et₂O and the *acetate-methyl ester* [64936-53-4] has **m** 145-140° from aq MeOH.

Albumin (bovine and human serum) [9048-46-8 (bovine); 70024-90-7 (human)] **M_r** ~67,000 (bovine), 69 000 (human), UV: A $\frac{1\%}{280\text{nm}}$ 6.6 (bovine) and 5.3 (human) in H₂O, $[\alpha]_{546}^{25} -78.2^\circ$ (H₂O). Purified by soln in conductivity water and passage at 2-4° through two ion-exchange columns, each containing a 2:1 mixture of anionic and cationic resins (Amberlite IR-120, H-form; Amberlite IRA-400, OH-form). This treatment removed ions and lipid impurities. Care was taken to exclude CO₂, and the soln was stored at -15°. [Möller, van Os and Overbeek *Trans Faraday Soc* 57 312 1961.] More complete lipid removal was achieved by lyophilising the de-ionised soln, covering the dried albumin (human serum) with a mixture of 5% glacial acetic acid (v/v) in iso-octane (previously dried with Na₂SO₄) and allowing to stand at 0° (without agitation) for upwards of 6h before decanting and discarding the extraction mixture, washing with iso-octane, re-extracting, and finally washing twice with iso-octane. The purified albumin was dried under vacuum for several hours, then dialyzed against water for 12-24h at room temperature, lyophilised, and stored at -10°C [Goodman *Science* 125 1296 1957]. It has been recrystd in high (35%) and in low (22%) EtOH solutions from Cohn's Fraction V.

The **high EtOH recrystn** was as follows: To 1 Kg of Fraction V albumin paste at -5° was added 300mL of 0.4 M pH (pH 5.5) acetate buffer in 35% EtOH pre-cooled to -10° and 430 mL of 0.1 M NaOAc in 25% EtOH also at -10°. Best results were obtained by adding all of the buffer and about half of the NaOAc and stirring slowly for 1hour. The rest of the NaOAc was added when all the lumps had disintegrated. The mixture was set aside at -5° for several days to crystallise. 35% EtOH (1 L) was then added to dilute the crystalline suspension and lower the ionic strength prior to centrifugation at -5° (yield 80%). The crystals were further dissolved in 1.5 volumes of 15% EtOH-0.02M NaCl at -5° and clarified by filtration through washed, calcined diatomaceous earth. This soln may be recrystd by re-adjusting to the conditions in the first crystallisation, or it may be recrystd at 22% EtOH with the aid of a very small amount of decanol (enough to give a final concn of 0.02%).

Note that crystn from lower EtOH gave better purification (i.e. by removing globulins and carbohydrates) and producing a more stable product.

The **low EtOH recrystn** was as follows: To 1 Kg of Fraction V at -10° to -15° was added 500mL of 15% EtOH at -5° , stirred slowly until a uniform suspension was formed. 15% EtOH (500mL) and sufficient 0.2M NaHCO_3 soln at 0° to bring the pH (1:10 diln) to 5.3. This required 125- 150mL. Some temp rise occurs and care must be taken to keep the temp $< -5^{\circ}$. If the albumin is incompletely dissolved a small amount of H_2O was added (100mL at a time at 0° , allowing 15min between additions). Undissolved albumin can be easily distinguished from small amounts of undissolved globulins, or as the last albumin dissolves, the appearance of the soln changes from milky white to hazy grey-green in colour. Keep the soln at -5° for 12h and filter by suspending in 15g of washed fine calcined diatomaceous earth, and thus filtering using a Büchner funnel precoated with coarser diatomaceous earth. The filtrate may require two or more similar filtrations to give a clear soln. To crystallise the filtrate add through a capillary pipette, and with careful stirring, 1/100volume of a soln containing 10% decanol and 60% EtOH (at -10°), and seeded with the needle-type albumin crystals. After 2-3 days crystn is complete. The crystals are centrifuged off. These are suspended with gentle mechanical stirring in one third their weight of 0.005 M NaCl pre-cooled to 0° . With careful stirring, H_2O (at 0°) is added slowly in an amount equal to 1.7 times the weight of the crystals. At this stage there is about 7% EtOH and the temp cannot be made lower than -2.5° to -1° . Clarify and collect as above. [Cohn et al. *J Am Chem Soc* **69** 1753 1947.]

Human serum albumin has been purified similarly with 25% EtOH and 0.2% decanol. The isoelectric points of bovine and human serum albumins are 5.1 and 4.9.

Amethopterin (Methotrexate, 4-amino-4-deoxy-*N*¹⁰-methylpteroyl-L-glutamic acid) [59-05-2] **M 454.4, m 185-204^o(dec)**, $[\alpha]_D^{20} +19^{\circ}$ (c 2, 0.1N aq NaOH), **pK₁ <0.5 (pyrimidine²⁺)**, **pK₂ 2.5 (N⁵-Me⁺)**, **pK₃ 3.49 (α -CO₂H)**, **pK₄ 4.99 (γ -CO₂H)**, **pK₅ 5.50 (pyrimidine⁺)**. Commonest impurities are 10-methyl pteroylglutamic acid, 4-amino-10-methylpteroylglutamic acid, aminopterin and pteroylglutamic acid. Purified by chromatography on Dowex-1 acetate, followed by filtration through a mixture of cellulose and charcoal. It has been recrystd from aqueous HCl or by dissolution in the minimum volume of N NaOH and acidified until pptn is complete, filter or collect by centrifugation, wash with H_2O (also by centrifugation) and dry at $100^{\circ}/3\text{mm}$. It has UV λ_{max} at 244 and 307nm (ϵ 17300 and 19700) in H_2O at pH 1; 257, 302 and 370nm (ϵ 23000, 22000 and 7100) in H_2O at pH 13. [Momle *Biochem Prep* **8** 20 1961; Seeger et al. *J Am Chem Soc* **71** 1753 1949.] It is a potent inhibitor of dihydrofolate reductase and used in cancer chemotherapy. [Blakley *The Biochemistry of Folic Acid and Related Pteridines* (North-Holland Publ Co., Amsterdam, NY) pp157-163 1969.] It is **CARCINOGENIC, HANDLE WITH EXTREME CARE**.

α -Amino acids. All the α -amino acids 'natural' configuration [*S* (L), except for cysteine which is *R*(L)] at the α - carbon atom are available commercially in a very high state of purity. Many of the 'non-natural' α -amino acids with the [*R*(D)] configuration as well as racemic mixtures are also available and generally none require further purification before use unless they are of 'Technical Grade'. The *R* or *S* enantiomers are optically active except for glycine which has two hydrogen atoms on the α - carbon atom, but these are *pro*-chiral and enzymes or proteins do distinguish between them, e.g. serine hydroxymethyltransferase successfully replaces the *pro*- α - hydrogen atom of glycine with CH_2OH (from formaldehyde) to make *S*-serine. The twenty common natural α -amino acids are: **amino acid**, three letter abbreviation, **one letter abbreviation**, pK ($-\text{COOH}$) and pK ($-\text{NH}_3^+$): **Alanine**, Ala, **A**, 2.34, 9.69; **Arginine**, Arg, **R**, 2.17, 9.04; **Asparagine**, Asn, **N**, 2.01, 8.80; **Aspartic acid**, Asp, **D**, 1.89, 9.60; **Cysteine**, Cys, **C**, 1.96, 8.18; **Glutamine**, Gln, **Q**, 2.17, 9.13; **Glutamic acid**, Glu, **E**, 2.19, 9.67; **Glycine**, Gly, **G**, 2.34, 9.60; **Histidine**, His, **H**, 1.8, 9.17; **Isoleucine**, Ile, **I**, 2.35, 9.68; **Leucine**, Leu, **L**, 2.36, 9.60; **Lysine**, Lys, **K**, 2.18, 8.95; **Methionine**, Met, **M**, 2.28, 9.20; **Phenylalanine**, Phe, **F**, 1.83, 9.12; **Proline**, Pro, **P**, 1.99, 10.96; **Serine**, Ser, **S**, 2.21, 9.15; **Threonine**, Thr, **T**, 2.11, 9.62; **Tryptophan**, Trp, **W**, 2.38, 9.39; **Tyrosine**, Tyr, **Y**, 2.2, 9.11, **Valine**, Val, **V**, 2.32, 9.61 repectively. Technical grade amino acids can be purified on ion exchange resins (e.g. Dowex 50W and eluting with a gradient of HCl or AcOH) and the purity is checked by TLC in two dimensions and stained with ninhydrin. (J.P.Greenstein and M.Winitz, *Chemistry of the Amino Acids* (3 Volumes), J.Wiley & Sons, NY, 1961; C.Cooper, N.Packer and K.Williams, *Amino Acid Analysis Protocols*, Humana Press, 2001, ISBN 0896036561). Recently codons for a further two amino acids have been discovered which are involved in ribosome-mediated protein synthesis giving proteins containing these amino

acids. The amino acids are *R(L)*-selenocysteine [Stadtman *Ann Rev Biochem* **65** 83 1996] and pyrrolysine [(4*R*, 5*R*)-4-substituted (with Me, NH₂ or OH) pyrroline-5-carboxylic acid] [Krzycki and Chan et al. *Science* **296** 1459 and 1462 2002.] They are, however, rare at present and only found in a few microorganisms.

9-Aminoacridine hydrochloride monohydrate (Acramine yellow, Monacrin) [52417-22-8] **M 248.7, m >355°**, **pK₁²⁰ 4.7, pK₂²⁰ 9.99**. Recrystd from boiling H₂O (charcoal; 1g in 300 mL) to give pale yellow crystals with a neutral reaction. It is one of the most fluorescent substances known. At 1:1000 dilution in H₂O it is pale yellow with only a faint fluorescence but at 1:100,000 dilution it is colourless with an intense blue fluorescence. [Albert and Ritchie *Org Synth Coll Vol III* 53 1955; Falk and Thomas *Pharm J* **153** 158 1944.] See entry in Chapter 4 for the free base.

Aminopterin (4-amino-4-deoxypteroyl-L-glutamic acid) [54-62-6] **M 440.4, m 231-235°(dec)**, **[α]_D²⁰ +18°** (c 2, 0.1N aq NaOH), **pK₁ <0.5 (pyrimidine²⁺)**, **pK₂ 2.5 (N5-Me⁺)**, **pK₃ 3.49 (α-CO₂H)**, **pK₄ 4.65 (γ-CO₂H)**, **pK₅ 5.50 (pyrimidine⁺)**. Purified by recrystn from H₂O, and has properties similar to those of methotrexate. It has UV at λ_{max} 244, 290 and 355nm (ε 18600, 21300 and 12000) in H₂O at pH 1; 260, 284 and 370nm (ε 28500, 26400 and 8600) in H₂O at pH 13. [Seeger et al. *J Am Chem Soc* **71** 1753 1949; Angier and Curran *J Am Chem Soc* **81** 2814 1959; Blakley *The Biochemistry of Folic Acid and Related Pteridines* (North-Holland Publ Co., Amsterdam, NY) pp157-163 1969.] For small quantities chromatography on DEAE cellulose with a linear gradient of ammonium bicarbonate pH 8 and increasing the molarity from 0.1 to 0.4 and followed by UV is best. For larger quantities a near boiling solution of aminopterin (5g) in H₂O (400mL) was slowly treated with small portions of MgO powder (~0.7g) calcined magnesnia) with vigorous stirring until a small amount of MgO remained undissolved and the pH rises from 3-4 to 7-8. Charcoal (1g) is added to the hot solution, filtered at once through a large sintered glass funnel of medium porosity and lined with a hot wet pad of Celite (~2-3 mm thick). The filtrate is cooled in ice and the crystals of the Mg salt are collected by filtration and recrystd from boiling H₂O (200mL) and the crystals washed with EtOH. The Mg salt is redissolved in boiling H₂O (200mL) and carefully acidified with vigorous agitation with AcOH (2mL). Pure aminopterin (3g) separates in fine yellow needles (dihydrate) which are easily fildt. The filtrate is washed with cold H₂O, then Me₂CO and dried in vac. If a trace of impurity is still present as shown by DEAE cellulose chromatography, then repetition of the process will remove it, see UV above. [Loo *J Med Chem* **8** 139 1965.] **CARCINOGENIC**

3-Aminopyridine adenine dinucleotide [21106-96-7] **M 635.4** (see NAD for pK) Purified by ion exchange chromatography [Fisher et al. *J Biol Chem* **248** 4293 1973; Anderson and Fisher *Methods Enzymol* **66** 81 1980].

α-Amino-thiophene-2-acetic acid 2-(2-thienyl)glycine [*R*(+)- 65058-23-3; *S*(-)- 4052-59-9; (-)- 43189-45-3; *RS*(±)- 21124-40-3] **M 57.2, m 236-237°** (*R*), **189-191°**, **235-236°** (*S*), **208-210°**, **223-224°** (dec)(*RS*), **[α]_D²⁰ (+) and (-) 84°** (c 1, 1% aq HCl), **[α]_D²⁵ (+) and (-) 71°** (c 1 H₂O), **pK_{Est(1)}~ 1.5, pK_{Est(2)}~ 8.0**. Recrystd by dissolving in H₂O (1g in 3 mL), adjusting the pH to 5.5 with aq NH₃, diluting with MeOH (20 mL), stirring, adjusting the pH to 5.5 and cooling to 0°. Also recrystd from small vols of H₂O. [*R*-isomer: Nishimura et al. *Nippon Kagaku Zasshi* **82** 1688 1961; *S*-isomer: Johnson and Panetta *Chem Abstr* **63** 14869h 1965; Johnson and Hardcastle *Chem Abstr* **66** 10930m 1967; *RS*-isomer: LiBassi et al. *Gazz Chim Ital* **107** 253 1977.] The (±) *N*-acetyl derivative has **m 191°** (from H₂O) [Schouteenten et al. *Bull Soc Chim Fr* II-248, II-252 1978].

4(6)-Aminouracil (4-amino-2,6-dihydroxypyrimidine) [873-83-6] **M 127.1, m >350°**, **pK₁²⁰ 0.00 (basic)**, **pK₂²⁰ 8.69 (acidic)**, **pK₃²⁰ 15.32 (acidic)**. Purified by dissolving in 3M aq NH₃, filter hot, and add 3M formic acid until pptn is complete. Cool, filter off (or centrifuge), wash well with cold H₂O, then EtOH and dry in air. Dry further in a vac at ~80°. [Barlin and Pfeiderer *J Chem Soc (B)* 1424 1971.]

Amylose [9005-82-7] (C₆H₁₀O₅)_n (for use in iodine complex formation). Amylopectin was removed from impure amylose by dispersing in aqueous 15% pyridine at 80-90° (concn 0.6-0.7%) and leaving the soln stand at 44-45° for 7 days. The ppte was re-dispersed and recrystd during 5 days. After a further dispersion in 15% pyridine, it was cooled to 45°, allowed to stand at this temperature for 12hours, then cooled

to 25° and left for a further 10 hours. The combined ppt was dispersed in warm water, ppted with EtOH, washed with absolute EtOH, and vacuum dried [Foster and Paschall *J Am Chem Soc* **75** 1181 1953].

Angiotensin (from rat brain) [70937-97-2] **M 1524.8**. Purified using extraction, affinity chromatography and HPLC [Hermann et al. *Anal Biochem* **159** 295 1986].

Angiotensinogen (from human blood serum) [64315-16-8]. Purified by chromatography on Blue Sepharose, Phenyl-Sepharose, hydroxylapatite and immobilised 5-hydroxytryptamine [Campbell et al. *Biochem J* **243** 121 1987].

Anion exchange resins. Should be conditioned before use by successive washing with water, EtOH and water, and taken through two OH⁻—H⁺—OH⁺ cycles by successive treatment with N NaOH, water, N HCl, water and N NaOH, then washed with water until neutral to give the OH⁻ form. (See commercial catalogues on ion exchange resins).

β-Apo-4'-carotenal [12676-20-9] **M 414.7, m 139°, A_{1cm}^{1%} 2640 at 461nm** Recrystd from CHCl₃/EtOH mixture or *n*-hexane. [Bobrowski and Das *J Org Chem* **91** 1210 1987.]

β-Apo-8'-carotenal [1107-26-2] **M 414.7, m 136-139°**. Recrystd from CHCl₃/EtOH mixture or *n*-hexane. [Bobrowski and Das *J Org Chem* **91** 1210 1987.]

β-Apo-8'-carotenoic acid ethyl ester [1109-11-1] **M 526.8, m 134-138°, A_{1cm}^{1%} 2550 at 449nm**. Crystd from pet ether or pet ether/ethyl acetate. Stored in the dark in an inert atmosphere at -20°.

β-Apo-8'-carotenoic acid methyl ester [16266-99-2] **M 512.7, m 136-137°, A_{1cm}^{1%} 2575 at 446nm and 2160 at 471nm, in pet ether**. Crystd from pet ether or pet ether/ethyl acetate. Stored in the dark in an inert atmosphere at -20°.

Apocodeine [641-36-1] **M 281.3, m 124°, pK_{Est(1)}~ 7.0, pK_{Est(2)}~ 8.2**. Crystd from MeOH and dried at 80°/2mm.

Apomorphine [58-00-4] **M 267.3, m 195°(dec), pK₁¹⁵ 7.20 (NH₂), pK₂¹⁵ 8.91 (phenolic OH)**. Crystd from CHCl₃ and pet ether, also from Et₂O with 1 mol of Et₂O which it loses at 100°. It is white but turns green in moist air or in alkaline soln. **NARCOTIC**

Apomorphine hydrochloride [41372-20-7] **M 312.8, m 285-287°(dec), [α]_D²⁰ -48° (c 1 H₂O)**. Cryst from H₂O and EtOH. Crystals turn green on exposure to light. **NARCOTIC**

Aureomycin (7-chlorotetracycline) [57-62-5] **M 478.5, m 172-174°(dec), [α]_D²³ -275° (MeOH), pK₁ 3.3, pK₂ 7.44, pK₃ 9.27**. Dehydrated by azeotropic distn of its soln with toluene. On cooling anhydrous material crystallises out and is recrystd from *C₆H₆, then dried under vacuum at 100° over paraffin wax. (If it is crystd from MeOH, it contains MeOH which is not removed on drying.) [Stephens et al. *J Am Chem Soc* **76** 3568 1954; *Biochem Biophys Res Commun* **14** 137 1964].

Aureomycin hydrochloride (7-chlorotetracycline hydrochloride) [64-72-2] **M 514.0, m 234-236°(dec), [α]_D²⁵ -23.5° (H₂O)**. Purified by dissolving 1g rapidly in 20mL of hot water, cooling rapidly to 40°, treating with 0.1mL of 2M HCl, and chilling in an ice-bath. The process is repeated twice. Also recrystd from Me₂NCHO + Me₂CO. [Stephens et al. *J Am Chem Soc* **76** 3568 1954 ; UV: McCormick et al. *J Am Chem Soc* **79** 2849 1975.]

Avidin (from egg white) [1405-69-2] **M_r ~70,000**. Purified by chromatography of an ammonium acetate soln on CM-cellulose [Green *Biochem J* **101** 774 1966]. Also purified by affinity chromatography on 2-iminobiotin-6-aminoethyl-Sepharose 4B [Orr *J Biol Chem* **256** 761 1981]. It is a biotin binding protein.

Azurin (from *Pseudomonas aeruginosa*) [12284-43-4] M_r 30,000. Material with $A_{625}/A_{280} = 0.56$ was purified by gel chromatography on G-25 Sephadex with 5mM phosphate pH 7 buffer as eluent [Cho et al. *J Phys Chem* **91** 3690 1987]. It is a blue Cu protein used in biological electron transport and its reduced form is obtained by adding a slight excess of $Na_2S_2O_4$. [See *Structure and Bonding* Springer Verlag, Berlin **23** 1 1975.]

Bacitracin (Altracin, Topitracin) [1405-87-4] M 1422.7, $[\alpha]_D^{23} + 5^\circ$ (H_2O). It has been purified by carrier displacement using *n*-heptanol, *n*-octanol and *n*-nonanol as carriers and 50% EtOH in 0.1 N HCl. The pure material gives one spot with $R_F \sim 0.5$ on paper chromatography using AcOH:*n*-BuOH: H_2O (4:1:5). [Porath *Acta Chem Scand* **6** 1237 1952.] It has also been purified by ion-exchange chromatography. It is a white powder soluble in H_2O and EtOH but insoluble in Et_2O , $CHCl_3$ and Me_2CO . It is stable in acidic soln but unstable in base. (Abraham and Bewton *Biochem J* **47** 257 1950; Synthesis: Munekata et al. *Bull Chem Soc Jpn* **46** 3187, 3835 1973.]

***N*⁶-Benzyladenine** [1214-39-7] M 225.3, m 231-232°, 232.5°(dec), $pK_{Est(1)} \sim 4.2$, $pK_{Est(2)} \sim 10.1$. Purified by recrystn from aqueous EtOH. It has λ_{max} at 207 and 270nm (H_2O), 268 nm (pH 6), 274nm (0.1 N HCl) and 275nm (0.1 N NaOH). [Daly *J Org Chem* **21** 1553 1956; Bullock et al. *J Am Chem Soc* **78** 3693 1956.]

***N*⁶-Benzyladenosine** [4294-16-0] M 357.4, m 177-179°, 185-187°, $[\alpha]_D^{25} - 68.6^\circ$ (c 0.6, EtOH)(see above entry for pK). Purified by recrystn from EtOH. It has λ_{max} 266nm (aq EtOH-HCl) and 269 nm (aqueous EtOH-NaOH). [Kissman and Weiss *J Org Chem* **21** 1053 1956.]

***N*-Benzylcinchoninium chloride (9*S*-benzyl-9-hydroxycinchoninium chloride)** [69221-14-3] M 421.0, $[\alpha]_D^{20} + 169^\circ$ (c 0.4, H_2O), $pK_{Est} \sim 5$. Recrystd from isoPrOH, toluene or small volumes of H_2O . Good chiral phase transfer catalyst [Julia et al. *J Chem Soc Perkin Trans 1* 574 1981; Hughes et al. *J Am Chem Soc* **106** 446 1984; Hughes et al. *J Org Chem* **52** 4745 1987]. See cinchonine below.

***R*-(-)-*N*-Benzylcinchonidinium chloride** [69257-04-1] M 421.0, m 212-213° (dec), $[\alpha]_D^{20} - 175.4^\circ$, -183° (c 5, 0.4, H_2O), $pK_{Est} \sim 5$. Dissolve in minimum volume of H_2O and add absolute Me_2CO . Filter off and dry in a vacuum. Also recrystd from hot EtOH or EtOH- Et_2O . (A good chiral phase transfer catalyst - see above) [Colonna et al. *J Chem Soc Perkin Trans 1* 547 1981, Imperali and Fisher *J Org Chem* **57** 757 1992]. See cinchonidine below.

***N*-Benzylpenicillin sodium salt** [69-57-8] M 356.37, m 215° (charring and dec), 225° (dec), $[\alpha]_D^{20} + 269^\circ$ (c 0.7, MeOH), $[\alpha]_D^{25} + 305^\circ$ (c 1, H_2O), $pK^{25} 2.76$ (4.84 in 80% aq EtOH)(for free acid). Purified by dissolving in a small volume of MeOH (in which it is more soluble than EtOH) and treating gradually with ~ 5 volumes of EtOAc. This gives an almost colourless crystalline solid (rosettes of clear-cut needles) and recrystallising twice more if slightly yellow in colour. The salt has also been conveniently recrystd from the minimum amount of 90% Me_2CO and adding an excess of absolute Me_2CO . A similar procedure can be used with wet *n*-BuOH. If yellow in colour then dissolve (~ 3.8 g) in the minimum volume of H_2O (3mL), add *n*-BuOH and filter through a bed of charcoal. The salt forms long white needles on standing in a refrigerator overnight. More crystals can be obtained on concentrating the mother liquors *in vacuo* at 40°. A further recrystn (without charcoal) yields practically pure salt. A good preparation has ~ 600 Units/mg. The presence of H_2O in the solvents increases the solubility considerably. The solubility in mg/100mL at 0° is 6.0 (Me_2CO), 15.0 ($Me_2CO + 0.5\% H_2O$), 31.0 ($Me_2CO + 1.0\% H_2O$), 2.4 (methyl ethyl ketone), 81.0 (*n*-butanol) and 15.0 (dioxane at 14°). Alternatively it is dissolved in H_2O (solubility is 10%), filtered if necessary and ppted by addition of EtOH and dried in a vacuum over P_2O_5 . A sample can be kept for 24h at 100° without loss of physiological activity. [IR: *Anal Chem* **19** 620 1947; *The Chemistry of Penicillin* [Clarke, Johnson and Robinson eds.] Princeton University Press, Princeton NJ, Chapter V 85 1949.]

Other salts, e.g. the **potassium salt** can be prepared from the Na salt by dissolving it (147mg) ice-cold in H_2O acidified to pH 2, extracting with Et_2O (~ 50 mL), wash once with H_2O , and extract with 2mL portions of 0.3% $KHCO_3$ until the pH of the extract rose to ~ 6.5 (~ 7 extracts). The combined aqueous extracts are

lyophilised and the white residue is dissolved in *n*-BuOH (1mL, absolute) with the addition of enough H₂O to effect soln. Remove insoluble material by centrifugation and add absolute *n*-BuOH to the supernatant. Crystals should separate on scratching, and after 2.5h in a refrigerator they are collected, washed with absolute *n*-BuOH and EtOAc and dried (yield 51.4mg). The *potassium salt* has *m* 214-217° (dec) (block preincubated at 200°; heating rate of 3°/min) and $[\alpha]_D^{22} +285^\circ$ (c 0.748, H₂O). The *free acid* has *m* 186-187° (MeOH-Me₂CO), 190-191° (H₂O) $[\alpha]_D^{25} +522^\circ$.

(+)-Bicuculine [*R*-6(5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolon-5-yl)-furo[3,4-*c*]-1,3-benzodioxolo-8(6*H*)-one] {485-49-4} *M* 367.4, *m* 177°, 193-195°, 193-197°, 215°, $[\alpha]_D^{20} +126^\circ$ (c 1, CHCl₃), $[\alpha]_{546}^{20} +159^\circ$ (c 1, CHCl₃), *pK* 4.84. Recrystallises from CHCl₃-MeOH as plates. Crystals melt at 177° then solidify and re-melt at 193-195° [Manske *Canad J Research* 21B 13 1943]. It is soluble in CHCl₃, *C₆H₆, EtOAc but sparingly soluble in EtOH, MeOH and Et₂O. [Stereochem: Blaha et al. *Collect Czech Chem Commun* 29 2328 1964; Snatzke et al. *Tetrahedron* 25 5059 1969; Pharmcol: Curtis et al. *Nature* 266 1222 1970].

L-erythro-Biopterin (2-amino-4-hydroxy-6-[[1*R*,2*S*]-1,2-dihydroxypropyl]pteridine) [22150-76-1] *M* 237.2, *m* >300°(dec), $[\alpha]_{546}^{20} -65^\circ$ (c 2.0, M HCl), *pK*_{1⁵} 2.23(2.45), *pK*_{2⁵} 7.89(8.05). Purified by chromatography on Florisil washed thoroughly with 2M HCl, and eluted with 2M HCl. The fractions with the UV-fluorescent band are evapd *in vacuo* and the residue recrystd. Biopterin is best recrystd (90% recovery) by dissolving in 1% aq NH₃ (ca 100 parts), and adding this soln dropwise to an equal vol of M aq formic acid at 100° and allowing to cool at 4° overnight. It is dried at 20° to 50°/01mm in the presence of P₂O₅. [Schircks, Bieri and Viscontini *Helv Chim Acta* 60 211 1977; Armarego, Waring and Paal *Aust J Chem* 35 785 1982.] Also crystd from ca 50 parts of water or 100 parts of hot 3M aq HCl by adding hot 3M aq NH₃ and cooling. It has UV: λ_{\max} at 212, 248 and 321nm (log ϵ 4.21, 4.09 and 3.94) in H₂O at pH 0.0; 223infl, 235.5, 274.5 and 345nm (log ϵ 4.07inflexion, 4.10, 4.18 and 3.82) in H₂O at pH 5.0; 221.5, 254.5 and 364nm (log ϵ 3.92, 4.38 and 3.84) in H₂O at pH 10.0 [Sugimoto and Matsuura *Bull Chem Soc Jpn* 48 3767 1875].

D-(+)-Biotin (vitamin H, hexahydro-2-oxo-1*H*-thieno[3,4-*d*]imidazole-4-pentanoic acid) [58-85-5] *M* 244.3, *m* 229-231°, 230.2°(dec), 230-231°, 232-234°(dec), $[\alpha]_{546}^{20} +108^\circ$, $[\alpha]_D^{20} +91.3^\circ$ (c 1, 0.1N NaOH), *pK*_{Est} ~ 4.8. Crystd from hot water in fine long needles with a solubility of 22 mg/100mL at 25°. Its solubility in 95% EtOH is 80 mg/100 mL at 25°. Its isoelectric point is at pH 3.5. Store solid and solutions under sterile conditions because it is susceptible to mould growth. [Confalone *J Am Chem Soc* 97 5936 1975; Wolf et al. *J Am Chem Soc* 67 2100 1945; Synthesis: Ohuri and Emoto *Tetrahedron Lett* 2765 1975; Harris et al. *J Am Chem Soc* 66 1756 1944.] The (+)-*methyl ester* has *m* 166-167° (from MeOH-Et₂O), $[\alpha]_D^{22} +57^\circ$ (c 1, CHCl₃) [du Vigneaud et al. *J Biol Chem* 140 643, 763 1941]; the (+)-*S-oxide* has *m* 200-203°, $[\alpha]_D^{20} +130^\circ$ (c 1.2, 0.1N NaOH) [Melville *J Biol Chem* 208 495 1954]; the *SS-dioxide* has *m* 274-275°(dec, 268-270°) and the *SS-dioxide methyl ester* has *m* 239-241° (from MeOH-Et₂O) [Hofmann et al. *J Biol Chem* 141 207, 213 1941.]

D-(+)-Biotin hydrazide [66640-86-6] *M* 258.4, *m* 238-240°, 245-247°, $[\alpha]_D^{20} +66^\circ$ (c 1, Me₂NCHO). Wash the material with H₂O, dry, wash with MeOH then Et₂O, dry, and recrystallise from hot H₂O (clusters of prisms) [Hofmann et al. *J Biol Chem* 144 513 1942].

D-(+)-Biotin *N*-hydroxysuccinimide ester (+-biotin *N*-succinimidyl ester) [35013-72-0] *M* 342.4, *m* 210°, 212-214°, $[\alpha]_D^{20} +53^\circ$ (c 1, Me₂NCHO). Recrystd from refluxing isoPrOH and dried in a vacuum over P₂O₅ + KOH. [Jasiewicz et al. *Exp Cell Biol* 100 213 1976.]

D-(+)-Biotin 4-nitrophenyl ester [33755-53-2] *M* 365.4, *m* 160-163°, 163-165°, $[\alpha]_D^{25} +47^\circ$ (c 2, Me₂NCHO containing 1% AcOH). It has been recrystd by dissolving 2g in 95% EtOH (30mL), heated to dissolve, then cooled in an ice-water bath. The crystals are collected, washed with ice-cold 95% EtOH (5mL) and dried over P₂O₅. The *R_F* on silica plates (CHCl₃:MeOH-19:1) is 0.19 [Bodanszky and Fagan *J Am Chem Soc* 99 235 1977].

***N*-(+)-Biotinyl-4-aminobenzoic acid** [6929-40-4] **M 363.4, m 295-297°, 295-300°, $[\alpha]_D^{25} +56.55^\circ$ (c 0.5, 0.1N NaOH), $pK_{Est} \sim 4.0$** . Dissolve in NaHCO₃ soln, cool and ppt by adding N HCl. Collect the solid, dry at 100° and recrystallise from MeOH. Note that it is hydrolysed by aq 3M, 1M and 0.2M HCl at 120°, but can be stored in 5% aq NaHCO₃ at -20° without appreciable hydrolysis [Knappe et al. *Biochem Zeitschrift* **338** 599 1963; *J Am Chem Soc* **73** 4142 1951; Bayer and Wilchek *Methods Enzymol* **26** 1 1980]

***N*-Biotinyl-6-aminocaproic *N*-succinimidyl ester** [72040-63-2] **M 454.5, m 149-152°**. Dissolve ~400mg in dry propan-2-ol (~25mL) with gentle heating. Reduce the volume to ~10mL by gentle boiling and allow the soln to cool. Decant the supernatant carefully from the white crystals, dry the crystals in a vacuum over P₂O₅ at 60° overnight. Material gives one spot on TLC. [Costello et al. *Clin Chem* **25** 1572 1979; Kincaid et al. *Methods Enzymol* **159** 619 1988.]

***N*-(+)-Biotinyl-6-aminocaproyl hydrazide (biotin-6-aminohexanoic hydrazide)** [109276-34-8] **M 371.5, m 189-191°, 210°, $[\alpha]_D^{20} +23^\circ$ (c 1, Me₂NCHO)**. Suspend in ice-water (100mg/mL), allow to stand overnight at 4°, filter and dry the solid in a vacuum. Recrystd from isoPrOH. R_F 0.26 on SiO₂ plate using CHCl₃-MeOH (7:3) as eluent. [O'Shannessy et al. *Anal Biochem* **163** 204 1987.]

***N*-(+)-Biotinyl-L-lysine (Biocytin)** [576-19-2] **M 372.5, m 228.5°, 228-230° (dec), 241-243°, 245-252° (dec, sintering at 227°), $[\alpha]_D^{25} +53^\circ$ (c 1.05, 0.1 N NaOH)**. Recrystd rapidly from dilute MeOH or Me₂CO. Also recrystd from H₂O by slow evaporation or by dissolving in the minimum volume of H₂O and adding Me₂CO until solid separates. It is freely soluble in H₂O and AcOH but insoluble in Me₂CO. [Wolf et al. *J Am Chem Soc* **76** 2002 1952, **72** 1048 1050.] It has been purified by chromatography on superfiltrol-Celite, Al₂O₃ and by countercurrent distribution and then recrystd [IR: Peck et al. *J Am Chem Soc* **74** 1991 1952]. The *hydrochloride* can be recrystd from aqueous Me₂CO + HCl and has m 227° (dec).

2-(4-Biphenyl)-5-phenyl-1,3,4-oxadiazole [852-38-0] **M 298.4, m 166-167°, 167-170°**. Recrystd from toluene. It is a good scintillation material [Brown et al. *Discussion Faraday Soc* **27** 43 1959].

2,5-Bis(4-biphenyl)-1,3,4-oxadiazole (BBOD) [2043-06-3] **M 374.5, m 229-230°, 235-238°**. Recrystd from heptane or toluene. It is a good scintillant. [Hayes et al. *J Am Chem Soc* **77** 1850 1955.]

4,4-Bis(4-hydroxyphenyl)valeric acid [diphenolic acid] [126-00-1] **M 286.3, m 168-171°, 171-172°, $pK_{Est(1)} \sim 4.8$ (CO₂H), $pK_{Est(2)} \sim 7.55$ (OH), $pK_{Est(3)} \sim 9.0$ (OH)**. When recrystd from *C₆H₆ the crystals have 0.5 mol of *C₆H₆ (m 120-122°) and when recrystd from toluene the crystals have 0.5 mol of toluene. Purified by recrystn from hot H₂O. It is sol in Me₂CO, AcOH, EtOH, propan-2-ol, methyl ethyl ketone. It is also recrystallised from AcOH, heptane-Et₂O or Me₂CO + *C₆H₆. It has λ_{max} 225 and 279nm in EtOH. The *methyl ester* has m 87-89° (aqueous MeOH to give the trihydrate). [Bader and Kantowicz *J Am Chem Soc* **76** 4465 1954.]

Bis(2-mercaptoethyl)sulfone (BMS) [145626-87-5] **M 186.3, m 57-58°, $pK_1^{25} 7.9, pK_2^{25} 9.0$** . Recrystd from hexane as white fluffy crystals. Large amounts are best recrystd from de-oxygenated H₂O (charcoal). It is a good alternative reducing agent to dithiothreitol. Its IR (film) has ν 2995, 2657, 1306, 1248, 1124 and 729 cm⁻¹. The synthetic intermediate *thioacetate* has m 82-83° (white crystals from CCl₄). The *disulfide* was purified by flash chromatography on SiO₂ and elution with 50% EtOAc-hexane and recrystd from hexane, m 137-139°. [Lamoureux and Whitesides *J Org Chem* **58** 633 1993.]

Bombesin (2-L-glutamin-3-6-L-asparaginealytesin) [31362-50-2] **M 619.9**. Purified by gel filtration on a small column of Sephadex G-10 and eluted with 0.01 M AcOH. This procedure removes lower molecular weight contaminants which are retarded on the column. The procedure should be repeated twice and the material should now be homogeneous on electrophoresis, and on chromatography gives a single active spot which is negative to ninhydrin but positive to Cl₂ and iodoplatinate reagents. R_F on paper chromatography (*n*-BuOH-pyridine-AcOH-H₂O (37.5: 25:7.5: 30)) is 0.55 for Bombesin and 0.65 for Alytin. [Bernardi *Experientia* **B** **27** 872 1971; **A** **27** 166 1971.] The *hydrochloride* has m 185°(dec) (from EtOH) $[\alpha]_D^{24} -20.6^\circ$ [c 0.65, Me₂NCHO-(Me₂N)₃PO (8:2)].

Bradykinin [ArgProProGlyPheSerProPheArg] [5979-11-3] M_r 1,240.4. Purified by ion-exchange chromatography on CMC (*O*-carboxymethyl cellulose) and partition chromatography on Sephadex G-25. Purity was checked by paper chromatography using BuOH:AcOH:H₂O (4:1:5) as eluent. [Park et al. *Can J Biochem* 56 92 1978; ORD and CD: Bodanszky et al. *Experientia* 26 948 1970; activity: Regoli and Barabé *Pharmacol Rev* 32 1 1980.]

Brefeldin A [1-*R*-2*c*,15*c*-dihydroxy-7*t*-methyl-(1*r*,13*t*)-6-oxa-bicyclo[11.3.0]hexadeca-3*t*,11*t*-dien-5-one, Decumbin] [20350-15-6] M 280.4, m 200-202°, 204°, 204-205°, $[\alpha]_D^{22} +95^\circ$ (c 0.81, MeOH). Isolated from *Penicillium brefeldianum* and recrystd from aqueous MeOH-EtOAc or MeOH. Solubility in H₂O is 0.6mg/mL, 10mg/mL in MeOH and 24.9mg/mL in EtOH. The *O*-acetate recrystallises from Et₂O-pentane and has m 130-131°, $[\alpha]_D^{22} +17^\circ$ (c 0.95, MeOH). [Sigg *Helv Chim Acta* 47 1401 1964; UV and IR: Härrri et al. *Helv Chim Acta* 46 1235 1963; total synthesis: Kitahara et al. *Tetrahedron* 3021 1979; X-ray : Weber et al. *Helv Chim Acta* 54 2763 1971.]

Bromelain (anti-inflammatory Ananase from pineapple) [37189-34-7] M_r ~33 000, [EC 3.4.33.4]. This protease has been purified *via* the acetone powder, G-75 Sephadex gel filtration and Bio-Rex 70 ion-exchange chromatography and has $A_{1cm}^{1\%}$ 20.1 at 280nm. The protease from pineapple hydrolyses benzoyl glycine ethyl ester with a K_m (app) of 210mM and k_{cat} of 0.36 sec⁻¹. [Murachi *Methods Enzymol* 19 273 1970; Balls et al. *Ind Eng Chem* 33 950 1941.]

5-Bromo-2'-deoxyuridine [59-14-3] M 307.1, m 193-197°(dec), 217-218°, $[\alpha]_D^{25} -41^\circ$ (c 0.1, H₂O), $pK^{25} \sim 8.1$. Recrystd from EtOH or 96% EtOH. It has λ_{max} 279 nm at pH 7.0, and 279 nm (log ϵ 3.95) at pH 1.9. Its R_F values are 0.49, 0.46 and 0.53 in *n*-BuOH-AcOH-H₂O (4:1:1), *n*-BuOH-EtOH-H₂O (40:11:19) and *i*-PrOH-25% aq NH₃-H₂O (7:1:1) respectively. [*Nature* 209 230 1966; *Collect Czech Chem Comm* 29 2956 1964.]

6-Bromo-2-naphthyl- α -D-galactopyranoside [25997-59-5] M 385.2, m 178-180°, 224-226°, 225°, $[\alpha]_D^{28} +60^\circ$ (c 1.2, pyridine). It was prepared from penta-*O*-acetyl-D-galactoside and 6-bromo-2-naphthol and ZnCl₂. The resulting tetra-acetate (2g) was hydrolysed by dissolving in 0.3N KOH (100mL) and heated until the soln was clear, filtered and cooled to give colourless crystals of the α -isomer which are collected and recrystd twice from hot MeOH. The high specific rotation is characteristic of the α -isomer. The *tetra-acetate* has m 155-156° $[\alpha]_D^{20} +60^\circ$ (c 1, CHCl₃) [Dey and Pridham *Biochem J* 115 47 1969] [reported m 75-85°, $[\alpha]_D^{24} +94^\circ$ (c 1.3, dioxane), Monis et al. *J Histochem Cytochem* 11 653 1963].

5-Bromouridine [957-75-5] M 323.1, m 215-217°, 217-218°, $[\alpha]_D^{25} -4.1^\circ$ (c 0.1, H₂O), $pK^{25} 8.1$. Recrystd from 96% EtOH. UV λ_{max} 279nm (log ϵ 3.95) in H₂O pH 1.9. R_F in *n*-BuOH:AcOH:H₂O (4:4:1) is 0.49; in *n*-BuOH:EtOH:H₂O (40:11:9) is 0.46 and in isoPrOH:25%NH₃:H₂O (7:1:2) is 0.53 using Whatman No 1 paper. [Prystas and Sorm *Collect Czech Chem Commun* 29 2956 1964.]

Brucine [357-57-3 (anhydr), 5892-11-5 (4H₂O)] M 430.5, m 178-179°, $[\alpha]_{546}^{20} -149.9^\circ$ (anhydrous; c 1, in CHCl₃), $pK_1^{15} 2.50$, $pK_2^{15} 8.16$ ($pK_2^{25} 8.28$). Crystd once from water or aq Me₂CO, as tetrahydrate, then suspended in CHCl₃ and shaken with anhydrous Na₂SO₄ (to dehydrate the brucine, which then dissolves). Pptd by pouring the soln into a large bulk of dry pet ether (b 40-60°), filtered and heated to 120° in a high vacuum [Turner *J Chem Soc* 842 1951]. **VERY POISONOUS**

α -Brucine sulfate (hydrate) [4845-99-2] M 887.0, m 180°(dec). Crystd from water.

Butyryl choline iodide [(2-butiryloxyethyl)trimethyl ammonium iodide] [2494-56-6] M 301.7, m 85-89°, 87°, 93-94°. Recrystd from isoPrOH or Et₂O. [Tammelin *Acta Chem Scand* 10 145 1956.] The *perchlorate* has m 72° (from isoPrOH). [Aldridge *Biochem J* 53 62 1953.]

S-Butyryl thiocholine iodide [(2-butirylycerthioethyl)trimethyl ammonium iodide] [1866-16-6] M 317.2, m 173°, 173-176°. Recrystd from propan-1-ol and dried *in vacuo*; store in the dark under N₂. The *bromide* has m 150° (from Me₂CO) or m 140-143° (from butan-1-ol). [Gillis *Chem and Ind (London)* 111 1957; Hansen *Acta Chem Scand* 11 537 1957.]

L-Canavanine sulfate (from jackbean, *O*-guanidino-L-homoserine) [2219-31-0] M_r 274.3, m 160-165°(dec), 172°(dec), $[\alpha]_D^{18.5} +19.8^\circ$ (c 7, H₂O), $pK_1^{2.5} 7.40$ (CO₂H), $pK_2^{2.5} 9.25$ (α -NH₂), $pK_3^{2.5} 11.5$ (guanidinoxy). Recrystd by dissolving (~1g) in H₂O (10mL), and adding with stirring 0.5 to 1.0 vols of 95% EtOH whereby crystals separate. These are collected, washed with Me₂CO-EtOH (1:1) and dried over P₂O₅ in a vacuum. [Hunt and Thompson *Biochem Prep* 13 416 1971; Feacon and Bell *Biochem J* 59 221 1955.]

Carbonic anhydrase (carbinic acid hydrolase) [9001-03-0] M_r 31,000 [EC 4.2.1.1]. Purified by hydroxylapatite and DEAE-cellulose chromatography [Tiselius et al. *Arch Biochem Biophys* 65 132 1956, *Biochim Biophys Acta* 39 218 1960], and is then dialysed for crystn. A 0.5 to 1% soln of the enzyme in 0.05 M Tris-HCl pH 8.5 was dialysed against 1.75M soln of (NH₄)₂SO₄ in the same buffer, and this salt soln was slowly increased in salt concn by periodic removal of small amounts of dialysate and replacement with an equal volume of 3.5M (NH₄)₂SO₄. The final salt concn in which the DEAE-cellulose fractions which gave beautiful birefringent suspensions of crystals ranged from 2.4 to 2.7M, and appeared first as fine crystals then underwent transition to thin fragile plates. Carbonic anhydrase is a Zn enzyme which exists as several isoenzymes of varying degrees of activity [*J Biol Chem* 243 6474 1968; crystal structure: *Nature, New Biology* 235 131 1972; see also P.D. Boyer Ed. *The Enzymes* Academic Press NY, pp 587-665 1971].

Carboxypeptidase A (from bovine pancreas, peptidyl-L-aminoacid lyase) [11075-17-5] M_r 34,600 [EC 3.4.17.1]. Purified by DEAE-cellulose chromatography, activation with trypsin and dialysed against 0.1M NaCl, yielding crystals. It is recrystd by dissolving in 20 mL of M NaCl and dialysed for 24hours each against the following salts present in 500mL of 0.02M sodium veronal pH 8.0: .05M NaCl, 0.2M NaCl and 0.15M NaCl. The last dialysate usually induces crystn. If it does not crystallise then dialyse the last soln against 0.02M sodium veronal containing 0.10M NaCl. Only 2 or 3 recrystns are required to attain maximum activity. [Cox et al. *Biochemistry* 3 44 1964.] Enzyme activity is measured by hydrolysing hippuryl-L-phenylalanine (or phenylacetic acid) and observing the rate of change of optical density at 254nm (reaction extinction coefficient is ~0.592 cm²/μmole at pH 7.5 [Bergmyer *Methods in Enzymatic Analysis* (Academic Press) 1 436 1974].

Carminic acid (7- α -D-glucopyranosyl-9,10-dihydro-3,5,6,8-tetrahydroxy-1-methyl-9,10-dioxo-2-anthracene carboxylic acid, Neutral Red 4: CI 75470) [1260-17-9] M_r 492.4, m 120°(dec), $[\alpha]_{654}^{15} +51.6^\circ$ (H₂O), (several phenolic pKs). Forms red prisms from EtOH. It gives a red colour in Ac₂O and yellow to violet in acidic solution. UV: λ_{max} (H₂O) 500nm (ϵ 6,800); (0.02N HCl) 490-500nm (ϵ 5,800) and (0.0001N NaOH) 540nm (ϵ 3,450). IR: ν_{max} (Nujol) 1708s, 1693s, 1677m, 1648m, 1632m, 1606s, 1566s, 1509 cm⁻¹. Periodate oxidation is complete after 4h at 0° with the consumption of 6.2 mols. The *tetra-O-methyl carminate* has m 186-188° (yellow needles from *C₆H₆ + pet ether). [IR: Ali and Haynes *J Chem Soc* 1033 1959; Bhatia and Venkataraman *Indian J Chem* 3 (2) 92 1965; Synthesis: Davis and Smith *Biochem Prep* 4 38 1955.]

Carnitine (α -hydroxy- β -N,N,N-trimethylaminopropionic acid) [R(+)- 541-14-0 ; S(L)- 541-15-1; RS 461-06-3] M_r 161.2, m R or S isomer 197-198°(dec), 210-212°(dec), RS isomer 195-197°, $[\alpha]_{546}^{20}$ (+) and (-) 36° (c 10, H₂O), $pK^{2.5} 3.6$. The S(L) isomer is levocarnitine, Vitamin B₇. The R or S isomers crystallise from EtOH + Me₂CO (hygroscopic). The R or S hydrochlorides crystallise from hot EtOH and have m 142°(dec). The RS isomer crystallises from hot EtOH (hygroscopic). The RS hydrochloride crystallises in needles from hot EtOH and has m 196°(dec).

L-Carnosine (β -alanyl-L-histidine) [305-84-0] M_r 226.2, m 258-260°(dec), 260°(capillary tube), 262°(dec), $[\alpha]_D^{25} +20.5^\circ$ (c 1.5, H₂O), $pK_1^{2.5} 2.64$, $pK_2^{2.5} 6.83$, $pK_3^{2.5} 9.51$. Likely impurities: histidine, β -alanine. Crystd from water by adding EtOH in excess. Recrystd from aqueous EtOH by slow addition of EtOH to a strong aqueous soln of the dipeptide. Its solubility in H₂O is 33.3% at 25°. [Vinick and Jung *J Org Chem* 48 392 1983; Turner *J Am Chem Soc* 75 2388 1953; Sifford and du Vigneaud *J Biol Chem* 108 753 1935.]

α -Carotene [7488-99-5] M 536.9, m 184-188°, $[\alpha]_{643}^{20} +385^\circ$ (c 0.08, *C₆H₆) λ_{\max} 422, 446, 474 nm, in hexane, $A_{1\text{cm}}^{1\%}$ 2725 (at 446nm), 2490 (at 474nm). Purified by chromatography on columns of calcium hydroxide, alumina or magnesia. Crystd from CS₂/MeOH, toluene/MeOH, diethyl ether/pet ether, or acetone/pet ether. Stored in the dark, under inert atmosphere at -20°.

all-trans- β -Carotene [7235-40-7] M 536.9, m 178-179°, 179-180°, 180°, 181°, 183° (evac capillary), $\epsilon_{1\text{cm}}^{1\%}$ 2590 (450nm), 2280 (478nm), in hexane. It forms purple prisms when crystd from *C₆H₆-MeOH and red rhombs from pet ether. Its solubility in hexane is 0.1% at 0°. It is oxygen sensitive and should be stored under N₂ at -20° in the dark. It gives a deep blue colour with SbCl₃ in CHCl₃. UV: (*C₆H₆) 429nm, λ_{\max} 454 and 484nm. The principal peak at 454nm has $A_{1\text{cm}}^{1\%}$ 2000. [Synthesis: Surmatis and Ofner *J Org Chem* 26 1171 1961; Milas et al. *J Am Chem Soc* 72 4844 1950.] β -Carotene was also purified by chromatography (Al₂O₃ activity I-II) - it was dissolved in pet ether-*C₆H₆ (10:1), applied to the column and eluted with pet ether-EtOH, the desired fraction was evaporated and the residue recrystd from *C₆H₆-MeOH as violet-red plates. [UV: Inhoffen et al. *Justus Liebigs Ann Chem* 570 54,68 1950; Review: Fleming *Selected Organic Synthesis* (J Wiley, Lond) pp. 70-74 1973.] Alternatively it can be purified by chromatography on a magnesia column, thin layer of Kieselguhr or magnesia. Crystd from CS₂/MeOH, Et₂O/pet ether, acetone/pet ether or toluene/MeOH. Stored in the dark, under inert atmosphere, at -20°. Recrystd from 1:1 EtOH/CHCl₃ [Bobrowski and Das *J Phys Chem* 89 5079 1985; Johnston and Scaiano *J Am Chem Soc* 108 2349 1986].

γ -Carotene [472-93-5, 10593-83-6] M 536.9, $A_{1\text{cm}}^{1\%}$ (λ_{\max}) 2055 (437nm), 3100 (462nm), 2720 (494nm) in hexane. Purified by chromatography on alumina or magnesia columns. Crystd from *C₆H₆/MeOH (2:1). Stored in the dark, under inert atmosphere, at 0°.

ξ -Carotene [38894-81-4] M 536.9, m 38-42°, λ_{\max} 378, 400, 425nm, $A_{1\text{cm}}^{1\%}$ (λ_{\max}) 2270 (400nm), in pet ether. Purified by chromatography on 50% magnesia-HyfloSupercel, developing with hexane and eluting with 10% EtOH in hexane. It was crystd from toluene/MeOH. [Gorman et al. *J Am Chem Soc* 107 4404 1985.] Stored in the dark under inert atmosphere at -20°.

λ -Carrageenan [9064-57-7, 9000-07-1 (κ + little of λ)]. This D-galactose-anhydro-D or L-galactoside polysaccharide is pptd from 4g of Carrageenan in 600mL of water containing 12g of KOAc by addn of EtOH. The fraction taken, pptd between 30 and 45% (v/v) EtOH. [Pal and Schubert *J Am Chem Soc* 84 4384 1962.]

Cation exchange resins. Should be conditioned before use by successive washing with water, EtOH and water, and taken through two H⁺-OH⁻-H⁺ cycles by successive treatment with M HCl, water, M NaOH, water and M HCl, then washed with water until neutral to give the H⁺ form. (See commercial catalogues on ion exchange resins).

Cathepsin B (from human liver) [9047-22-7] M_r 27,500 [EC 3.4.22.1]. Purified by affinity chromatography on the semicarbazone of Gly-Phe-glycinal-linked to Sepharose 4B, with elution by 2,2'-dipyridyl disulfide [Rich et al. *Biochem J* 235 731 1986; *Methods Enzymol* 80 551 1981].

Cathepsin D (from bovine spleen) [9025-26-7] M_r 56,000, [EC 3.4.23.5]. Purified on a CM column after ammonium sulfate fractionation and dialysis, then starch-gel electrophoresis and by ultracentrifugal analysis. Finally chromatographed on a DEAE column [Press et al. *Biochem J* 74 501 1960].

Cephalosporin C potassium salt [28240-09-7] M 453.5, $[\alpha]_{\text{D}}^{20} +103^\circ$ (H₂O), pK₁ <2.6, pK₂ 3.1, pK₃ 9.8. Purified by dissolving in the minimum volume of H₂O (filter) and adding EtOH until separation of solid is complete. A soln is stable in the pH range 2.5-8. It has UV λ_{\max} is 260nm (log ϵ 3.95) in H₂O. The Ba salt has $[\alpha]_{\text{D}}^{20} +80^\circ$ (c 0.57, H₂O) [Woodward et al. *J Am Chem Soc* 88 852 1966; Abraham and Newton *Biochem J* 79 377 1961; Hodgkin and Maslen *Biochem J* 79 402 1961; see also *Quart Reviews Chem Soc* London 21 231 1967].

Ceruloplasmin (from human blood plasma) [9031-37-2] M_r 134,000. This principle Cu transporter (90-90% of circulating Cu) is purified by precipitation with polyethylene glycol 4000, batchwise adsorption and elution from QAE-Sephadex, and gradient elution from DEAE-Sepharose CL-6B. Ceruloplasmin

was purified 1640-fold. Homogeneous on anionic polyacrylamide gel electrophoresis (PAGE), SDS-PAGE, isoelectric focusing and low speed equilibrium centrifugation. [Oestnuizen *Anal Biochem* **146** 1 1985; Cohn et al. *J Am Chem Soc* **68** 459 1946.]

Chemokines. These are small proteins formed from longer precursors and are chemoattractants for lymphocytes and lymphoid organs. They are characterised by having cysteine groups in specific relative positions. The two largest families are the α and β families that have four cysteine residues arranged (C-X-C) and (C-C) respectively. The mature chemokines have ~70 amino acids with internal cys S-S bonds and attract myeloid type cells *in vitro*. The γ -family (Lymphotactin) has only two cys residues. The δ -family (Neurotactin, Fractalkine) has the C-C-X-X-X-C sequence (*ca* 387 amino acids), binds to membrane promoting adhesion of lymphocytes. The soluble domain of human Fractalkine chemoattracts monocytes and T cells. Several chemokines are available commercially (some prepared by recombinant DNA techniques) including 6Ckine/exodus/SLC which belongs to the β -family with 6 cysteines (110 amino acids mature protein), as the name implies (C-C-C-C-X.....X-C-C) and homes lymphocytes to secondary lymphoid organs with lymphocyte adhesion antitumor properties. Other chemokines available are C10 (β CC) and Biotaxin. Several chemokine receptors and antibodies are available commercially and can generally be used without further purification. [Murphy 'Molecular biology of lymphocyte chemoattractant receptors' in *Ann Rev Immunol* **12** 593 1994.]

Chirazymes. These are commercially available enzymes e.g. lipases, esterases, that can be used for the preparation of a variety of optically active carboxylic acids, alcohols and amines. They can cause regio and stereospecific hydrolysis and do not require cofactors. Some can be used also for esterification or transesterification in neat organic solvents. The proteases, amidases and oxidases are obtained from bacteria or fungi, whereas esterases are from pig liver and thermophilic bacteria. For preparative work the enzymes are covalently bound to a carrier and do not therefore contaminate the reaction products. Chirazymes are available from Roche Molecular Biochemicals and are used without further purification.

Chlorambucil [4-{bis(2-chloroethyl)amino}benzene)butyric acid] [305-03-3] M 304.2, m 64-66°, pK₁ 5.8 (6.0 at 66°, 50% aq Me₂CO), pK₂ 8.0. It is recrystd from pet ether (flat needles) and has a solubility at 20° of 66% in EtOH, 40% in CHCl₃, 50% in Me₂CO but is insoluble in H₂O [Everett et al. *J Chem Soc* 2386 1953]. **CARCINOGEN.**

Chloramphenicol [Amphicol, 1R,2R-(-)-2-{2,2-dichloroacetyl-amino}-1-{4-nitrophenyl}-propan-1,3-diol] [56-75-7] M 323.1, m 149-151°, 150-151°, 151-152°, [α]_D²⁰ +20.5° (c 3, EtOH), [α]_D²⁵ -25.5° (EtOAc). Purified by recrystn from H₂O (sol 2.5mg/mL at 25°) or ethylene dichloride as needles or long plates and by sublimation at high vacuum. It has A_{1cm}^{1%} 298 at λ_{\max} 278nm and it is slightly soluble in H₂O (0.25%) and propylene glycol (1.50%) at 25° but is freely soluble in MeOH, EtOH, BuOH, EtOAc and Me₂CO. [Relstock et al. *J Am Chem Soc* **71** 2458 1949; Confroulis et al. *J Am Chem Soc* **71** 2463 1949; Long and Troutman *J Am Chem Soc* **71** 2469, 2473 1949, Ehrhart et al. *Chem Ber* **90** 2088 1957.]

Chloramphenicol palmitate [530-43-8] M 561.5, m 90°, [α]_D²⁶ +24.6° (c 5, EtOH). Crystd from *benzene.

2-Chloroadenosine [146-77-0] M 301.7, m 145-146°(dec), 147-149°(dec), pK_{Est(1)} ~ 0.5, pK_{Est(2)} ~ 7.6. Purified by recrystn from H₂O (~1% in cold) and has λ_{\max} at 264 nm (pH 1 and 7) and 265 nm (pH 13) in H₂O. [Brown and Weliky *J Org Chem* **23** 125 1958; Schaeffer and Thomas *J Am Chem Soc* **80** 3738 1958; IR: Davoll and Lewy *J Am Chem Soc* **74** 1563 1952.]

Chlorophylls a and b see entries on p. 167 in Chapter 4.

6-Chloropurine riboside (6-chloro-9- β -D-ribofuranosyl-9H-purine) [2004-06-0] M 286.7, m 158-162°(dec), 165-166°(sintering At 155°), 168-170°(dec), [α]_D²⁶ -45° (c 0.8, H₂O). Purified by suspending the dry solid (~12 g) in hot MeOH (130 mL) and then adding enough hot H₂O (~560 mL) to cause solution, filter and set aside at 5° overnight. The colourless crystals of the riboside are filtered off, washed with Me₂CO, Et₂O and dried at 60°/0.1mm. More material can be obtained from the filtrate by evapn to

dryness and recrystn of the residue from MeOH-H₂O (2:1) (15mL/g). It has λ_{\max} 264nm (ϵ 9140) in H₂O. [Robins *Biochem Prep* 10 145 1963; Baker et al. *J Org Chem* 22 954 1957.]

Chromomycin A₃ [7059-24-7] M 1183.3, m 185°dec, $[\alpha]_{\text{D}}^{23}$ -57° (c 1, EtOH). Dissolve reagent (10g) in EtOAc and add to a column of Silica Gel (Merck 0.05-0.2microns, 4x70cm) in EtOAc containing 1% oxalic acid. Elute with EtOAc+1% oxalic acid and check fractions by TLC. Pool fractions, wash with H₂O thoroughly, dry and evaporate. Recryst from EtOAc. The *hepta-acetate* has m 214°, $[\alpha]_{\text{D}}^{23}$ -20° (c 1, EtOH). [*Tetrahedron* 23 421 1967; *J Am Chem Soc* 91 5896 1969.]

α -Chymotrypsin [9004-07-3] M_r ~25000 [EC 3.4.21.1]. Crystd twice from four-tenths saturated ammonium sulfate soln, then dissolved in 1mM HCl and dialysed against 1mM HCl at 2-4°. The soln was stored at 2° [Lang, Frieden and Grunwald *J Am Chem Soc* 80 4923 1958].

Cinchonidine [485-71-2] M 294.4, m 210.5°, $[\alpha]_{\text{D}}^{20}$ -127.5° (c 0.5, EtOH), pK₁¹⁵ 4.17, pK₂¹⁵ 8.4. Crystd from aqueous EtOH. For *N-benzyl chloride* see entry in Chapter 6.

Cinchonine [118-10-5] M 294.4, m 265°, $[\alpha]_{\text{D}}^{20}$ +268° (c 0.5, EtOH), pK₁¹⁵ 4.28, pK₂¹⁵ 8.35. Crystd from EtOH or diethyl ether. For *N-benzyl chloride* see entry in Chapter 6.

Citranaxanthin [3604-90-8] M 456.7, m 155-156°, A_{1cm}^{1%} (λ_{\max}) 410 (349nm), 275 (466nm) in hexane. Purified by chromatography on a column of 1:1 magnesium oxide and HyfloSupercel (diatomaceous filter aid). Crystd from pet ether. Stored in the dark, under inert atmosphere, at 0°.

Citric acid cycle components (from rat heart mitochondria). Resolved by anion-exchange chromatography [LaNoue et al. *J Biol Chem* 245 102 1970].

Clonidine hydrochloride [Catapres, 2-(2,6-dichloroanilino)-2-imidazoline hydrochloride] [4205-91-8] M 266.6, m 305°, pK 5.88 (free base). It is recrystd from EtOH-Et₂O and dried in a vacuum (solubility in H₂O is 5%). It has a pKa of 5.88. The *free base* has m 124-125° and is recrystallised from hexane. [Jen et al. *J Med Chem* 18 90 1975; NMR: Jackman and Jen *J Am Chem Soc* 97 2811 1975.]

Clostripain [9028-00-6] [EC 3.4.22.8] M_r ~55,000. Isolated from *Clostridium histolyticum* callogenase by extraction in pH 6.7 buffer, followed by hydroxylapatite chromatography with a 0.1-0.2 M phosphate gradient, then Sephadex G-75 gel filtration with 0.05M phosphate pH 6.7, dialysis and a second hydroxylapatite chromatography (gradient elution with 0.1M → 0.3M phosphate, pH 6.7). It has proteinase and esterase activity and is assayed by hydrolysing *n*-benzoyl-L-arginine methyl ester. [Mitchell and Harrington *J Biol Chem* 243 4683 1968, *Methods Enzymol* 19 635 1970.]

Cloxacillin sodium salt (sodium 3-*o*-chlorophenyl-5-methyl-4-isoxazolyl penicillin monohydrate) [642-78-4] M 457.9, m 170°, $[\alpha]_{\text{D}}^{20}$ +163° (H₂O pH 6.0-7.5), pK_{Est} ~ 2.8 (COOH). Purified by dissolving in isoPrOH containing 20% of H₂O, and diluting with isoPrOH to a water content of 5% and chilled, and recrystd again in this manner. The sodium salt is collected and dried at 40° in air to give the colourless monohydrate. It is soluble in H₂O (5%), MeOH, EtOH, pyridine and ethylene glycol. [Doyle et al. *J Chem Soc* 5838 1963; Naylor et al. *Nature* 195 1264 1962.]

β -Cocaine {2 β -carbomethoxy-3- β -benzoxypyrone, methyl [1*R*-(*exo,exo*)]-3-(benzoyloxy)-2-methyl-8-azabicyclo[3.2.1]octane-2-carboxylate} [50-36-2] M 303.4, m 98°, b 187-188°/0.1mm, $[\alpha]_{\text{D}}^{20}$ -15.8° (c 4, CHCl₃), pK²⁴ 8.39. Crystallises from EtOH and sublimes below 90° in a vacuum in a non-crystalline form.

Coccarboxylase tetrahydrate (aneurine pyrophosphoric acid tetrahydrate, thiamine pyrophosphoric acid tetrahydrate) [136-09-4] M 496.4, m 220-222°(sinters at 130-140°), 213-214°, pK_{Est(1)}~2, pK_{Est(2)}~6, pK_{Est(3)}~9. Recrystd from aqueous Me₂CO. [Wenz et al. *Justus Liebigs Ann Chem* 618 210 1958; UV: Melnick *J Biol Chem* 131 615 1939; X-ray: Carlisle and Cook *Acta Cryst (B)* 25 1359 1969.] The *hydrochloride salt* has m 242-244°(dec), 241-243°(dec) or 239-240°(dec) and is

recrystd from aqueous HCl + EtOH, EtOH containing HCl or HCl + Me₂CO. [Weijlard *J Am Chem Soc* **63** 1160 1941; Synthesis: Weijlard and Tauber *J Am Chem Soc* **60** 2263 1938.]

Codeine [76-57-3] M 299.4, m 154-156°, $[\alpha]_D^{20}$ -138° (in EtOH), pK²⁵ 8.21. Crystd from water or aqueous EtOH. Dried at 80°.

Coenzyme A trihydrate [85-61-0] M 821.6, pK₁ 4.0 (adenine NH₂), pK₂ 6.5 (PO₄H), pK₃ 9.6 (SH). White powder best stored in an inert atmosphere in the dark in sealed ampoules after drying *in vacuo* over P₂O₅ at 34°. It has UV: λ_{\max} 259 nm (ϵ 16,800) in H₂O. [Buyske et al. *J Am Chem Soc* **76** 3575 1954.] It is sol in H₂O but insol in EtOH, Et₂O and M₂CO. It is readily oxidised in air and is best kept as the more stable *trilithium salt* [Moffat and Khorana *J Am Chem Soc* **83** 663 1961; see also Beinert et al. *J Biol Chem* **200** 384 1953; De Vries et al. *J Am Chem Soc* **72** 4838 1950; Gregory et al. *J Am Chem Soc* **74** 854 1952 and Baddiley *Adv Enzymol* **16** 1 1955].

Coenzyme Q₀ (2,3-Dimethoxy-5-methyl-1,4-benzoquinone, 3,4-dimethoxy-2,5-toluquinone, fumigatin methyl ether) [605-94-7] M 182.2, m 56-58°, 58-60°, 59°. It crystallises in red needles from pet ether (b 40-60°) and can be sublimed in high vacuum with a bath temperature of 46-48° [Ashley, Anslow and Raistrick *J Chem Soc* 441 1938; UV in EtOH: Vischer *J Chem Soc* 815 1953; UV in cyclohexane: Morton et al. *Helv Chim Acta* **41** 2343 1858; Aghoramurthy et al. *Chem Ind (London)* 1327 1954].

Coenzyme Q₄ (Ubiquinone-4, 2,3-dimethoxy-5-methyl-6-[3,7,11,15-tetramethyl-hexadeca-2*t*,6*t*,10*t*,14-tetraenyl]-[1,4]benzoquinone [4370-62-1] M 454.7, m 30°, 33-45°, A^{1%}_{1cm} (275nm) 185. A red oil purified by TLC chromatography on SiO₂ and eluted with Et₂O-hexane. Purity can be checked by HPLC (silica column using 7% Et₂O-hexane). It has λ_{\max} 270 nm (ϵ 14,800) in pet ether. [NMR and MS: Naruta *J Org Chem* **45** 4097 1980; cf Morton *Biochemical Spectroscopy* (Adam Hilger, London, 1975) p 491]. It has also been dissolved in MeOH/EtOH (1:1 v/v) and kept at 5° until crystals appear [Lester and Crane *Biochim Biophys Acta* **32** 497 1958].

Coenzyme Q₉ (Ubiquinone-9, 2,3-dimethoxy-5-methyl-6-[3,7,11,15,19,23,27,31,35-nonamethyl-hexatriaconta-2*t*,6*t*,10*t*,14*t*,18*t*,22*t*,26*t*,30*t*,34-nonaenyl]-1,4-benzoquinone) [303-97-9] M 795.3, m 40.5-42.5°, 44-45°, 45°. Yellow crystals purified by recrystn from pet ether and by TLC chromatography on SiO₂ and eluted with Et₂O-hexane. Purity can be checked by HPLC (silica column using 7% Et₂O-hexane). It has λ_{\max} 270nm (ϵ 14,850) in pet ether. [NMR and MS: Naruta *J Org Chem* **45** 4097 1980; Le et al. *Biochem Biophys Acta* **32** 497 1958; cf Morton *Biochemical Spectroscopy* (Adam Hilger, London, 1975) p 491; IR: Lester et al. *Biochim Biophys Acta* **33** 169 1959; UV: Ruegg et al. *Helv Chim Acta* **42** 2616 1959; Shunk *J Am Chem Soc* **81** 5000 1959.]

Coenzyme Q₁₀ (Ubiquinone-10, 2,3-dimethoxy-5-methyl-6-[3,7,11,15,19,23,27,31,35,-39-decamethyl-tetraconta-2*t*,6*t*,10*t*,14*t*,18*t*,22*t*,26*t*,30*t*,34*t*,38-decaenyl]-1,4-benzoquinone) [303-98-0] M 795.3, m 48-49°, 49°, 49.5-50.5°, 50°. Purified by recrystn from EtOH, EtOH + Me₂CO or Et₂O-EtOH and by chromatography on silica gel using isoPrOH-Et₂O (3:1) to give orange crystals. It has λ_{\max} 270nm (ϵ 15,170) in pet ether. [Terao et al. *J Org Chem* **44** 868 1979; NMR and MS: Naruta et al. *J Org Chem* **45** 4097 1980; IR: Lester et al. *Biochem Biophys Acta* **42** 1278 1959, NMR: Planta et al. *Helv Chim Acta* **42** 1278 1959; cf Morton *Biochemical Spectroscopy* (Adam Hilger, London, 1975) p 491].

Colcemide (Demecocine) [477-30-5] M 371.4, m 182-185°, 183-185°, 186°, $[\alpha]_D^{20}$ -129° (c 1, CHCl₃). It has been purified by chromatography on silica and eluting with CHCl₃-MeOH (9:1) and recrystn from EtOAc-Et₂O and forms yellow prisms. UV in EtOH has λ_{\max} 243nm (ϵ 30,200) and 350nm (ϵ 16,3000). [Synthesis, IR, NMR, MS: Capraro and Brossi *Helv Chim Acta* **62** 965 1979.]

Colchicine [64-86-8] M 399.5, m 155-157°(dec), $[\alpha]_{546}^{20}$ -570° (c 1, H₂O), pK²⁰ 1.85. Commercial material contains up to 4% desmethylcolchicine. Purified by chromatography on alumina, eluting with CHCl₃ [Ashley and Harris *J Chem Soc* 677 1944]. Alternatively, an acetone solution on alkali-free alumina has been used, and eluting with acetone [Nicholls and Tarbell *J Am Chem Soc* **75** 1104 1953].

Colchicoside [477-29-2] M 547.5, m 216-218°. Crystd from EtOH.

Colicin E (from *E.coli*) [11032-88-5]. Purified by salt extraction of extracellular-bound colicin followed by salt fractionation and ion-exchange chromatography on a DEAE-Sephadex column, and then by CM-Sephadex column chromatography [Schwartz and Helinski *J Biol Chem* **246** 6318 1971].

Collagenase (from human polymorphonuclear leukocytes) [9001-12-1] M_r 68,000-125,000 [EC 3.4.24.3]. Purified by using *N*-ethylmaleimide to activate the enzyme, and wheat germ agglutinin-agarose affinity chromatography [Callaway et al. *Biochemistry* **25** 4757 1986].

Compactin [73573-88-3] M 390.5, m 151-153°, 152°, $[\alpha]_D^{22} +283^\circ$ (c 0.48, acetone). Purified by recrystn from aqueous EtOH. UV: λ_{max} 230, 237 and 246nm (log ϵ 4.28, 4.30 and 4.11); IR (KBr): ν 3520, 1750 (lactone CO) and 1710 (CO ester) cm^{-1} . [Clive et al. *J Am Chem Soc* **110** 6914 1988; Synthesis review: Rosen and Heathcock *Tetrahedron* **42** 4909 1986; IR, NMR, MS: Brown et al. *J Chem Soc Perkin Trans 1* 1165 1976.]

Convallatoxin (α cardenolide mannoside) [508-75-8] M 550.6, m 238-239°, $[\alpha]_D^{25} 9.4^\circ$ (c 0.7, dioxane). Crystd from EtOAc. *Tetra-acetate* has m 238-242° (MeOH/Et₂O), $[\alpha]_D^{25} -5^\circ$ (CHCl₃).

Copper-zinc-superoxide dismutase (from blood cell haemolysis) [9054-89-1] M_r ~32,000 [EC 1.15.1.1]. Purified by DEAE-Sepharose and copper chelate affinity chromatography. The preparation was homogeneous by SDS-PAGE, analytical gel filtration chromatography and by isoelectric focusing [Weselake et al. *Anal Biochem* **155** 193 1986; Fridovich *J Biol Chem* **244** 6049 1969].

Coproporphyrin I [531-14-6] M 654.7, λ_{max} 591, 548, 401nm in 10% HCl. Crystd from pyridine/glacial acetic acid. The *dihydrochloride* [69477-27-6] has M 727.7 and λ_{max} 395nm in water.

Corticosterone (11 β , 21-dihydroxypregn-4-en-3,20-dione) [50-22-6] M 346.5, m 180-181°, 180-182°, 181-184°, $[\alpha]_D^{15} +223^\circ$ (c 1.1, EtOH), $[\alpha]_D^{23-25} +194^\circ$ (c 0.1, dioxane). Purified by recrystn from Me₂CO (trigonal plates), EtOH or isoPrOH. UV λ_{max} at 240nm, and gives an orange-yellow soln with strong fluorescence on treatment with concentrated H₂SO₄. Insoluble in H₂O but soluble in organic solvents. [Reichstein and Euw *Helv Chim Acta* **21** 1197 1938, **27** 1287 1944; Mason et al. *J Biol Chem* **114** 613 1936; ORD: Foltz et al. *J Am Chem Soc* **77** 4359 1955; NMR: Shoolery and Rogers *J Am Chem Soc* **80** 5121 1958.] The *21-O-benzoyl* derivative has m 201-202° [Reichstein *Helv Chim Acta* **20** 953 1937].

Corticotropin [92307-52-3] **polypeptide** M_r ~4697. Extract separated by ion-exchange on CM-cellulose, desalted, evapd and lyophilised. Then run on gel filtration (Sephadex G-50) [Lande et al. *Biochemical Preparations* **13** 45 1971; Esch et al. *Biochem Biophys Res Commun* **122** 899 1984].

Cortisol see hydrocortisone on p. 541.

Cortisone [53-06-5] M 360.5, m 230-231°, $[\alpha]_{546}^{20} +225^\circ$ (c 1, in EtOH). Crystd from 95% EtOH or acetone.

Cortisone-21-acetate [50-04-4] M 402.5, m 242-243°, $[\alpha]_{546}^{20} +227^\circ$ (c 1, in CHCl₃). Crystd from acetone.

Creatine (H₂O) (*N*-guanidino-*N*-methylglycine) [6020-87-7] M 131.1, m 303°, $pK_1^{25} 2.63$, $pK_2^{25} 14.3$. Likely impurities are creatinine and other guanidino compounds. Crystd from water as monohydrate. Dried under vacuum over P₂O₅ to give anhydrous material.

Creatine phosphate di Na, 4H₂O salt (phosphocreatine) [922-32-7] M 327.1, $pK_1^{27} 2.7$, $pK_2^{27} 4.58$, $pK_3^{27} \sim 12$. To 3-4g of salt in H₂O (220mL) is added 4 vols of EtOH with thorough stirring and allowed to stand at 20° for 12hrs (this temp is critical as crystals did not readily form at 23° or 25°). The salt first appears as oily droplets which slowly settle and crystallise. After 12hrs the supernatant is clear. Stirring

and scratching the flask containing the filtrate brings out additional (0.3-1g) crystals if the salt is kept at 20° for 12hrs. Filter at room temp, wash with 3 x 5mL of ice-cold 90% EtOH then 5mL of abs EtOH and dry in a vac desiccator (Drierite or CaCl₂) for 16-30hrs. The hexahydrate (plates) is converted to the tetrahydrate salt (needles) in vac at -10°. [Ennor and Stocken *Biochem Prep* 5 9 1957; *Biochem J* 43 190 1958.]

Creatinine (2-imino-1-methyl-4-oxoimidazolidine) [60-27-5] M 113.1, m 260°(dec), pK₁²⁵ 4.80, pK₂²⁵ 9.2. Likely impurities are creatine and ammonium chloride. Dissolved in dilute HCl, then neutralised by adding ammonia. Recrystd from water by adding excess of acetone.

Crotaline (monocrotaline, 12,13-dihydroxy-(13β-14βH)-14,19-dihydro-20-norcrotalanan-11,15-dione) [315-22-0] M 325.4, m 196-197°(dec), 197-198°(dec), 203°(dec), [α]_D²⁰ -55° (c 1, EtOH). It forms prisms from absolute EtOH and recrystallises also from CHCl₃. UV in 96% EtOH has λ_{max} 217nm (log ε 3.32). [Adams et al. *J Am Chem Soc* 74 5612 1952; Culvenor and Smith *Aust J Chem* 10 474 1957.] The *hydrochloride* has m 212-214° (from MeOH-Et₂O) and [α]_D²⁸ -38.4° (c 5, H₂O) [Adams and Gianturco *J Am Chem Soc* 78 1922 1956]. The *picrate* has m 230-231.5°(dec) [Adams et al. *J Am Chem Soc* 74 5614 1952].

α-Cyclodextrin (H₂O) [10016-20-3] M 972.9, m >280°(dec), [α]₅₄₆ +175° (c 10, H₂O). Recrystd from 60% aq EtOH, then twice from water, and dried for 12hours in a vacuum at 80°. Also purified by pptn from water with 1,1,2-trichloroethylene. The ppte was isolated, washed and resuspended in water. This was boiled to steam distil the trichloroethylene. The soln was freeze-dried to recover the cyclodextrin. [Armstrong et al. *J Am Chem Soc* 108 1418 1986].

β-Cyclodextrin (H₂O) [7585-39-9] M 1135.0, m >300°(dec), [α]₅₄₆ +170° (c 10, H₂O). Recrystd from water and dried for 12hours in a vacuum at 110°, or 24hours in a vacuum at 70°. The purity was assessed by TLC on cellulose with a fluorescent indicator. [Taguchi, *J Am Chem Soc* 108 2705 1986; Tabushi et al. *J Am Chem Soc* 108 4514 1986; Orstam and Ross *J Phys Chem* 91 2739 1987.]

D-(R-natural) and L-(S-non-natural) Cycloserine (2-amino-3-isoxazolidone) [R- 68-41-7 and S- 339-72-0] M 102.1, m 145-150° (dec), 154-155°, 155-156° (dec), 156° (dec), [α]_D²⁵ (+) and (-) 137° (c 5, 2N NaOH), pK₁¹⁰ 4.5, pK₂¹⁰ 7.74, pK₁²⁵ 4.50, pK₂²⁵ 7.43, pK₁⁵⁰ 4.44, pK₂⁵⁰ 7.20. Purified by recrystn from aqueous EtOH or MeOH or aqueous NH₃ + EtOH or isoPrOH. Also recrystd from aqueous ammoniacal soln at pH 10.5 (100mg/mL) by diluting with 5 volumes of isopropanol and then adjusting to pH 6 with acetic acid. An aqueous soln buffered to pH 10 with Na₂CO₃ can be stored in a refrigerator for 1 week without decomposition. UV: λ_{max} 226nm (A_{1cm}^{1%} 4.02). The *tartrate salt* has m 165-166° (dec), 166-168° (dec), and [α]_D²⁴ -41° (c 0.7, H₂O). [Stammer et al. *J Am Chem Soc* 79 3236 1959; UV: Kuehl *J Am Chem Soc* 77 2344 1955.]

Cystamine dihydrochloride [2,2'-diaminodiethylene disulfide dihydrochloride, 2,3'-dithio-bis(ethylamine) dihydrochloride] [56-17-7] M 225.2, m 219-220°(dec), pK₁³⁰ 8.82, pK₂³⁰ 9.58. Recrystd by dissolving in EtOH containing a few drops of dry EtOH-HCl, filtering and adding dry Et₂O. The solid is dried in a vacuum and stored in dry and dark atmosphere. It has been recrystd from EtOH (solubility: 1g in 60mL of boiling EtOH) or MeOH (plates). The *free base* has b 90-100°/0.001mm, 106-108°/5mm and 135-136°/atm, d₄²⁰ 1.1559, n_D²⁰ 1.5720. [Verly and Koch *Biochem J* 58 663 1954; Gonick et al. *J Am Chem Soc* 76 4671 1954; Jackson and Block *J Biol Chem* 113 137 1936]. The *dihydrobromide* has m 238-239° (from EtOH-Et₂O) [Viscontini *Helv Chim Acta* 36 835 1953].

S,S-(L,L)-Cystathionine (S-2-amino-2-carboxyethyl-L-homocysteine, L-2-amino-4[(2-amino-2-carboxyethyl)thio]butyric acid) [56-88-2] M 222.3, m >300°, dec at 312° with darkening at 270°, [α]_D²⁰ +23.9° (c 1, M HCl). Could be converted to the HCl salt by dissolving in 20% HCl and carefully basifying with aqueous NH₃ until separation is complete. Filter off and dry in a vacuum. It forms prisms from H₂O. The *dibenzoyl* derivative has m 229° (from EtOH). [IR: Greenstein and Winitz *Chemistry of the Amino Acids (J Wiley)* Vol 3 2690 1961 and Tallan et al. *J Biol Chem* 230 707 1958; Synthesis: du Vigneaud et al. *J Biol Chem* 143 59 1942; Anslow et al. *J Biol Chem* 166 39 1946.]

[Prepn: Weiss and Stekol *J Am Chem Soc* **73** 2497 1951; see also du Vigneaud et al. *J Biol Chem* **143** 60 1942; Biological synthesis: Greenberg *Methods Enzymol* **5** 943 1962.]

Cysteamine (2-aminoethanethiol, 2-mercaptoethylamine) [60-23-1] **M 77.2, m 97-98.5°, 98-99°, 99-100°, pK₁⁰ 9.15, pK₂⁰ 11.93, pK₁^{3.0} 8.42, pK₂^{3.0} 10.83.** Soluble in H₂O giving an alkaline reaction and it has a disagreeable odour. Likely impurity is the disulfide, cystamine which is not soluble in alkaline solution. Under a N₂ atmosphere dissolve in EtOH, evaporate to dryness and wash the white residue with dry pet ether, then sublime at 0.1mm and store under N₂ (out of contact with air) at 0-10° in the dark. Its HgCl₂ (2:3) complex has **m 181-182°** (from H₂O), and its picrate has **m 125-126°**. [Mills and Bogert *J Am Chem Soc* **57** 2328 1935, **62** 1173 1940; Baddiley and Thain *J Chem Soc* 800 1952; Shirley *Preparation of Organic Intermediates (J. Wiley)* Vol 3 189 1951; Barkowski and Hedberg *J Am Chem Soc* **109** 6989 1987.]

Cysteamine hydrochloride [156-57-0] **M 113.6, m 70.2-70.7°, 70-72°.** Purified by recrystn from EtOH. It is freely soluble in H₂O and should be stored in a dry atmosphere. [Mills and Bogert *J Am Chem Soc* **62** 1177 1940.] The picrate has **m 125-126°**, see previous entry for *free base*.

(±)-Cysteic acid (3-sulfoalanine, 1-amino-3-sulfopropionic acid) [13100-82-8, 3024-83-7] **M 169.2, m 260°(dec).** Likely impurities are cystine and oxides of cysteine. Crystd from water by adding 2 volumes of EtOH. When recrystd from aqueous MeOH it has **m 264-266°**, and the anhydrous acid has **m ~260°(dec)**. [Chapeville and Formageot *Biochim Biophys Acta* **26** 538 1957; *J Biol Chem* **72** 435 1927.]

R(L)-Cysteic acid (H₂O) [23537-25-9] **M 187.2, m 275-280° (dec), 289°, [α]_D²⁰ +8.66°** (c 7.4, H₂O, pH 1) and **+1.54°** (H₂O, pH 13), **pK₁^{2.5} 1.9** (SO₃H), **pK₂^{2.5} 8.7** (CO₂H), **pK₃^{2.5} 12.7** (NH₂). Likely impurities are cystine and oxides of cysteine. Crystd from water by adding 2 volumes of EtOH. When recrystd from aqueous MeOH it has **m 264-266°**, and the anhydrous acid has **m ~260°(dec)**. [Chapeville and Formageot *Biochim Biophys Acta* **26** 538 1957; *J Biol Chem* **72** 435 1927.]

D-(S)- and L-(R)- Cysteine (S- and R-2-amino-3-mercaptopropionic acid) [*S*(+)-: 921-01-7 and *R*(-)-: 52-90-4] **M 121.2, m 230°, 240° (dec), [α]_D²⁰ (+) and (-) 7.6°** (c 2, M HCl) and **(+) and (-) 10.1°** (c 2, H₂O, pH 10), **pK₁^{2.5} 1.92** (CO₂), **pK₂^{2.5} 8.35** (NH₂), **pK₃^{2.5} 10.46** (SH). Purified by recrystn from H₂O (free from metal ions) and dried in a vacuum. It is soluble in H₂O, EtOH, Me₂CO, EtOAc, AcOH, *C₆H₆ and CS₂. Acidic solns can be stored under N₂ for a few days without deterioration. [For synthesis and spectra see Greenstein and Winitz *Chemistry of the Amino Acids (J. Wiley)* Vol 3 p1879 1961.]

L-Cysteine hydrochloride (H₂O) [52-89-1] **M 175.6, m 175-178° (dec), [α]_D²⁵ +6.53°** (5M HCl). Likely impurities are cystine and tyrosine. Crystd from MeOH by adding diethyl ether, or from hot 20% HCl. Dried under vacuum over P₂O₅. *Hygroscopic*.

(±)-Cysteine hydrochloride [10318-18-0] **M 157.6.** Crystd from hot 20% HCl; dried under vacuum over P₂O₅.

L-Cystine [56-89-3] **M 240.3, [α]_D^{18.5} -229°** (c 0.92 in M HCl), **pK₁^{2.5} 1.04** (1.65), **pK₂^{2.5} 2.05** (2.76), **pK₃^{2.5} 8.00** (7.85), **pK₄^{2.5} 10.25** (8.7, 9.85). Cystine disulfoxide was removed by treating an aqueous suspension with H₂S. The cystine was filtered off, washed with distilled water and dried at 100° under vacuum over P₂O₅. Crystd by dissolving in 1.5M HCl, then adjusting to neutral pH with ammonia. Likely impurities are D-cystine, meso-cystine and tyrosine.

Cytidine [65-46-3] **M 243.2, m 210-220°(dec), 230° (dec), 251-252° (dec), [α]_D²⁰ +37°** (c 9, H₂O), **[α]_D²⁰ +29°** (c 9, H₂O), **pK 3.85.** Crystd from 90% aqueous EtOH. Also has been converted to the *sulfate* by dissolving (~200mg) in a soln of EtOH (10mL) containing H₂SO₄ (50mg), whereby the salt crystallises out. It is collected, washed with EtOH and dried for 5hours at 120°/0.1mm. The *sulfate* has **m 225°**. The *free base* can be obtained by shaking with a weak ion-exchange resin, filtering, evaporating and recrystallising the residue from EtOH as before. [Fox and Goodman *J Am Chem Soc* **73** 3256 1956; Fox and

Shugar *Biochim Biophys Acta* **9** 369 1952; see Prytsas and Sorm in *Synthetic Procedures in Nucleic Acid Chemistry* (Zorbach and Tipson Eds) **Vol 1** 404 1973.]

Cytisine see entry in Chapter 4.

Cytochalasin B (from dehydrated mould matter) [14930-96-2] **M 479.6**. Purified by MeOH extraction, reverse phase C18 silica gel batch extraction, selective elution with 1:1 v/v hexane/tetrahydrofuran, crystn, subjected to TLC and recrystallised [Lipski et al. *Anal Biochem* **161** 332 1987].

Cytochrome c₁ (from horse, beef or fishes' heart, or pigeon breast muscle) [9007-43-6] **M ~ 13,000**. Purified by chromatography on CM-cellulose (CM-52 Whatman) [Brautigan et al. *Methods Enzymol* **53D** 131 1978]. It has a high PI (isoelectric point) and has been purified further by adsorption onto an acidic cation exchanger, e.g. Amberlite IRC-50 (polycarboxylic) or in ground form Amberlite XE-40 (100-200 mesh) or Decalso-F (aluminium silicate), where the non-cytochrome protein is not adsorbed and is readily removed. The cytochrome is eluted using a soln containing 0.25g ions/L of a univalent cation at pH 4.7 adsorbed onto the NH₄⁺ salt of Amberlite IRC-50 at pH 7, washed with H₂O and then with 0.12M NH₄OAc to remove non-cytochrome protein. When the cytochrome begins to appear in the eluate then the NH₄OAc concn is increased to 0.25 M. The fractions with *ca* Fe = 0.465—0.467 are collected, dialysed against H₂O and adsorbed onto a small IRC-50 column and eluted with 0.5M NH₄OH, then dialysed and lyophilised. (A second fraction (II) can be eluted from the first resin with 0.5M NH₄OH but is discarded). [Keilin and Hartree *Biochem Prep* **1** 1 1952; Margoliash *Biochem Prep* **8** 33 1957.]

Cytochrome c has been recrystd as follows: The above eluate (*ca* 100mL) is dialysed against H₂O (10 vols) at 4° for 24 h (no more), then passed through an XE-40 column (2 x 1 cm above) which is equilibrated with 0.1M NH₄OAc pH 7.0. The column is washed with 0.1% (NH₄)₂SO₄ pH 8.0 and the dark red resin in the upper part of the column is collected and in 0.1% (NH₄)₂SO₄ pH 8.0 transferred to another column (7 mm diameter) and the cytochrome c is eluted with 5% (NH₄)₂SO₄ pH 8.0. More than 98% of the red colour is collected in a volume of *ca* 4mL in a weighed centrifuge tube. Add a drop of octanol, 0.43g of (NH₄)₂SO₄/g of soln. When the salt has dissolved ascorbic acid (5mg), add a few drops of 30% NH₃ and keep the soln at 10° for 10min (turns lighter colour due to reduction). Then add finely powdered (NH₄)₂SO₄ in small portions (stir with a glass rod) until the soln becomes turbid. Stopper the tube tightly, and set aside at 15-25° for 2 days while the cytochrome c separates as fine needles or rosettes. Further (NH₄)₂SO₄ (20mg) are added per mL of suspension and kept in the cold for a few days to complete the crystallisation. The crystals are collected by centrifugation (5000xg), suspended in saturated (NH₄)₂SO₄ (pH 8.0 at 10°) then centrifuged again. For recrystn the crystals are dissolved in the least volume of H₂O, one drop of ammonia and 1 mg of ascorbic acid are added and the above process is repeated. The yield of twice recrystd cytochrome c from 2Kg of muscle is *ca* 200 mg but this varies with the source and freshness of the muscle used. The crystals are stored as a solid after dialysis against 0.08M NaCl or 0.1M sodium buffer and lyophilising, or as a suspension in saturated (NH₄)₂SO₄ at 0°. [Hagihara et al. *Biochem Prep* **6** 1 1958.] *Purity of cytochrome c:* This is checked by the ratio of the absorbance at 500nm (reduced form) to 280nm (oxidised form), i.e. $\epsilon_{500}/\epsilon_{280}$ should be between 1.1 and 1.28, although values of up to 1.4 have been obtained for pure preparations.

For the preparation of the *reduced form* see Margoliash *Biochem Prep* **5** 33 1957 and Yonetani *Biochem Prep* **11** 19 1966.

Cytochrome from *Rhodospirillum rubrum*. ($\epsilon_{270}/\epsilon_{551}$ 0.967). Purified by chromatography on a column of CM-Whatman cellulose [Paleus and Tuppy *Acta Chem Scand* **13** 641 1959].

Cytochrome c oxidase (from bovine heart mitochondria). [9001-16-5] **M_r 100,000/haeme**, [EC 1.9.3.1]. Purified by selective solubilisation with Triton X-100 and subsequently with lauryl maltoside; finally by sucrose gradient centrifugation [Li et al. *Biochem J* **242** 417 1978].

Also purified by extraction in 0.02 M phosphate buffer (pH 7.4) containing 2% of cholic acid (an inhibitor which stabilises as well as solubilises the enzyme) and fractionated with (NH₄)₂SO₄ collecting the 26-33% saturation cut and refractionating again and collecting the 26-33% saturation fraction. The pellet collected at 10,000xg appears as an oily paste. The cholate needs to be removed to activate the enzyme as follows: The ppte is dissolved in 10mL of 0.1M phosphate buffer pH 7.4, containing 1% of Tween-80 and dialysed against 1L of 0.01 M PO₄ buffer (pH 7.4) containing 1% of Tween-80 for 10 h at 0° and aliquoted. The enzyme is

stable at 0° for 2 weeks and at -15° for several months. It is assayed for purity (see reference) by oxidation of reduced cytochrome c (Km 10µM). [Yonetani *Biochem Prep* 11 14 1966; *J Biol Chem* 236 1680 1961.]

Cytokines see chemokines, interferons, interleukins.

Cytosine [71-30-7] M 111.1, m 320-325° (dec), pK₁²⁵ 4.6, pK₂²⁵ 12.1. Crystd from water.

Cytosine-1-β-O-arabinofuranoside (Cytarabin) [147-94-4] M 243.2, m ~220°(dec), 212-213.5°, [α]_D²⁰ +155° (c 1, H₂O), pK²⁵ 4.3. Purified by recrystn from aqueous EtOH. It has λ_{max} 212 and 279nm at pH 2 and 272nm at pH 12. [Walwick et al. *Proc Chem Soc (London)* 84 1959.]

N-Decanoyl-N-methylglucamine (Mega-10, N-D-glucidyl-N-methyl deconamide) [85261-20-7] M 349.5, m 91-93°, 92°. Possible impurities are decanoic acid and N-methylglycamine. The former is removed by grinding the solid with Et₂O and then with pet ether and dried over P₂O₅. Twice recrystd from MeOH-Et₂O by dissolving in the minimum volume of MeOH and adding Et₂O and drying in a vacuum. To remove the glycamine the solid (800mg) is dissolved in hot H₂O (10mL) and set aside. Mega-10 crystallises in colourless needles. These are filtered off and dried in a vacuum to constant weight. It is a good non-ionic non-hygroscopic detergent with a critical micelle concentration (CMC) of 7.4mM (0.26%) in 0.1M Tris-HCl pH 7.4 at 25°. [Hildreth *Biochem J* 207 363 1982.]

Demeclocycline hydrochloride (7-chloro-6-demethyltetracycline hydrochloride, Clortetrin) [64-73-3] M 501.3, m 174-178°(dec, for sesquihydrate), [α]_D²⁵ -258° (c 0.5, 0.1N H₂SO₄), pK 4.45 [H₂O-Me₂NCHO (1:1)]. Crystd from EtOH-Et₂O or H₂O and dried in air [McCormick et al. *J Am Chem Soc* 79 4561 1957; Dobrynin et al. *Tetrahedron Lett* 901 1962].

2'-Deoxyadenosine (adenine 2'-deoxyriboside) [16373-93-6] M 269.3, m 187-189°, 189-191°, [α]_D²⁰ -25° (c 0.5, H₂O), [α]₅₈₉²⁵ -26°, [α]₃₁₀²⁵ -206° (c 0.5, H₂O), pK²⁰ 3.79. Purified by recrystn from H₂O (as hydrated crystals; solubility of mono-hydrate is 1.1% in H₂O at 20°). It has λ_{max} 258nm (pH 1), 260nm (pH 7) and 261nm (pH 13). [Ness and Fletcher *J Am Chem Soc* 81 4752 1959; Walker and Butler *Can J Chem* 34 1168 1956.] The 3',5'-O-diacetyl derivative has m 151-152° (recrystd from EtOAc-pet ether).

3'-Deoxyadenosine (Cordycepin, adenine 3'-deoxyriboside) [73-03-0] M 251.2, m 225-226°, 225-229°, [α]_D²⁰ -47° (H₂O), pK_{Est} ~ 4.8. It forms needles from EtOH, n-BuOH and n-PrOH, and from H₂O as the mono-hydrate. It has λ_{max} 260nm (ε 14,600) in EtOH. The picrate has m 195°(dec, yellow crystals from H₂O). [Kaczka et al. *Biochim Biophys Acta* 14 456 1964; Todd and Ulbricht *J Chem Soc* 3275 1960; Lee et al. *J Am Chem Soc* 83 1906 1961; Walton et al. *J Am Chem Soc* 86 2952 1964.]

11-Deoxycorticosterone acetate (21-acetoxy-4-pregnen-3,20-dione) [56-47-3] M 372.5, m 154-159°, 154-160°, 155-157°, 155-161°, [α]_D²⁰ +174° (c 1, dioxane), [α]_D²²⁻²⁴ +196° (c 1, CHCl₃). Recrystallises from EtOH as needles or Me₂CO-hexane, and sublimes at high vacuum. Partly soluble in MeOH, Me₂CO, Et₂O and dioxane but insoluble in H₂O. [Romo et al. *J Am Chem Soc* 79 5034 1957; NMR: Shoolery and Rogers *J Am Chem Soc* 80 5121 1959.]

2'-Deoxycytidine monohydrate [951-77-9] M 245.2, m 119-200°, 207-209°, 213-215°, [α]_D²⁵ +78° (c 0.4, N NaOH), [α]_D²³ +57.6° (c 2, H₂O), pK 4.25. Purified by recrystn from MeOH-Et₂O or EtOH and dried in air. [NMR: Miles *J Am Chem Soc* 85 1007 1963; UV: Fox and Shugar *Biochim Biophys Acta* 9 369 1952.] The hydrochloride crystallises from H₂O-EtOH and has m 174°(dec, 169-173°) [Walker and Butler *Can J Chem* 34 1168 1956.] The picrate has m 208°(dec). [Fox et al. *J Am Chem Soc* 83 4066 1961.]

2'-Deoxycytidine 5'-monophosphoric acid (deoxycytidylic acid) [1032-65-1] M 307.2, m 170-172°(dec), 183-184°(dec), 183-187°(dec), [α]_D²¹ +35° (c 0.2, H₂O), pK₁ 4.6, pK₂ 6.6.

Recrystd from H₂O or aqueous EtOH and dried in a vacuum. [Volkin et al. *J Am Chem Soc* **73** 1533 1951; UV: Fox et al. *J Am Chem Soc* **75** 4315 1953; IR: Michelson and Todd *J Chem Soc* 3438 1954.]

2'-Deoxyguanosine monohydrate (9-[2-deoxy-β-D-ribofuranosyl]guanidine) [961-07-9] **M 285.3, m ca 200°(dec)**, $[\alpha]_D^{20} +37.5^\circ$ (c 2, H₂O), $[\alpha]_D^{14} -47.7^\circ$ (c 0.9, N NaOH), **pK_{Est(1)}~ 3.3, pK_{Est(2)}~ 9.2**. Recrystd from H₂O as the monohydrate. [Brown and Lythgoe *J Chem Soc* 1990 1950; Levene and London *J Biol Chem* **81** 711 1929, **83** 793 1929]; UV: Hotchkiss *J Biol Chem* **175** 315 1948; ORD: Levendahl and James *Biochim Biophys Acta* **26** 89 1957.] The 3',5'-di-O-acetyl derivative crystd from aqueous EtOH has **m 222°(dec)**, $[\alpha]_D^{18} -38^\circ$ (c 0.3, 10% aq EtOH) [Hayes et al. *J Chem Soc* 808, 813 1955].

2'-Deoxyinosine [890-38-0] **M 252.2, m 206°(dec), 218-220°(dec)**, $[\alpha]_D^{25} -2.1^\circ$ (c 2, N NaOH), $[\alpha]_D^{21.5}$ (c 1, H₂O), **pK_{Est(1)}~ 8.9, pK_{Est(2)}~ 12.4**. Purified by recrystn from H₂O. [Brown and Lythgoe *J Chem Soc* 1990 1950; UV: : MacNutt *Biochem J* **50** 384 1952.]

5-Deoxy-5-(methylthio)adenosine [2457-80-9] **M 297.3, m 210-213°(dec), 211°, 212°, 213-214°**, $[\alpha]_D^{20} -23.7^\circ$ (c 0.02, pyridine), $[\alpha]_D^{20} -8^\circ$ (c 1, 5% aq NaOH), $[\alpha]_D^{25} +15^\circ$ (c 0.4-1.0, 0.3N aq AcOH), **pK_{Est}~3.5**. It has been recrystd from H₂O and sublimed at 200°/0.004mm. [v.Euler and Myrbäck *Z physiol Chem* **177** 237 1928; Weygand and Trauth *Chem Ber* **84** 633 1951; Baddiley et al. *J Chem Soc* 2662 1953.] The hydrochloride has **m 161-162°** [Kuhn and Henkel *Hoppe Seyler's Z Physiol Chem* **269** 41 1941]. The picrate has **m 183°(dec)** (from H₂O).

Deoxyribonucleic acid (from plasmids). Purified by two buoyant density ultracentrifugations using ethidium bromide-CsCl. The ethidium bromide was extracted with Et₂O and the DNA was dialysed against buffered EDTA and lyophilised. [Marmur and Doty *J Mol Biol* **5** 109 1962; Guerry et al. *J Bacteriol* **116** 1064 1973.] See p. 504.

3'-Deoxythymidine {2',3'-dideoxythymidine, 1-[(2r)-5c-hydroxymethyltetrahydro(2r)-furyl]-5-methylpyrimidine-2,4-dione} [3416-05-5] **M 226.2, m 145°, 149-151°**, $[\alpha]_D^{26} +18^\circ$ (c 1, H₂O), **pK_{Est}~ 9.2**. Recrystd from Me₂CO + MeOH. [Michelson and Todd *J Chem Soc* 816 1955.]

2'-Deoxyuridine [1-(β-D-erythro-2-deoxypentofuranosyl)-1H-pyrimidine-2,4-dione] [951-78-0] **M 228.2, m 163°, 163-163.5°, 165-167° 167°**, $[\alpha]_D^{26} +30^\circ$ (c 2, H₂O), $[\alpha]_D^{22} +50^\circ$ (c 1, N NaOH), **pK₂₅ 9.3**. Forms needles from absolute EtOH or 95% EtOH. [Dekker and Todd *Nature* **166** 557 1950; Brown et al. *J Chem Soc* 3035 1958; NMR Jardetzky *J Am Chem Soc* **83** 2919 1961; Fox and Shugar *Biochim Biophys Acta* **9** 369 1952; UV: MacNutt *Biochem J* **50** 384 1952.]

3'-Deoxyuridine {1-[(2R)-5c-hydroxymethyltetrahydro(2r)furyl]-5-methylpyrimidin-2,4-dione, 2',3'-dideoxythymidine} [7057-27-4, 3416-05-5] **M 226.2, m 149-151°**, $[\alpha]_D^{20} +18^\circ$ (c 1, H₂O), **pK_{Est}~ 9.3**. Recrystd from Me₂CO + MeOH and dried in a vacuum. [Michelson and Todd *J Chem Soc* 816 1955.]

Dermatan sulfate (chondroitin sulfate B from pig skin) [54328-33-5 (Na salt)]. Purified by digestion with papain and hyaluronidase, and fractionation using aqueous EtOH. [Gifonelli and Roden *Biochem Prep* **12** 1 1968.]

Desthiobiotin [533-48-2] **M 214.3, m 156-158°**, $[\alpha]_D^{20} +10.5^\circ$ (c 2, H₂O), **pK_{Est}~2.8**. Crystd from H₂O or 95% EtOH.

Dextran [9004-54-0] **M_r 6,000-220,000**. Solutions keeps indefinitely at room temperature if 0.2mL of Roccal (10% alkylidimethylbenzylammonium chloride) or 2mg phenyl mercuric acetate are added per 100mL solution. [Scott and Melvin *Anal Biochem* **25** 1656 1953.]

Diacetone-D-Glucose (1,2:5,6-di-O-isopropylidene-α-D-glucofuranoside) [582-52-5] **M 260.3, m 107-110°, 110.5°, 111-113°, 112°**, $[\alpha]_D^{15} -18.4^\circ$ (c 1, H₂O). It crystallises from Et₂O, (needles), pet ether or *C₆H₆ and sublimes *in vacuo*. It is sol in 7 vols of H₂O and 200 vols of pet ether at their

boiling points. The solubility in H₂O at 17.5° is 4.3%. It pptes from aq solns on basification with NaOH. [Schmid and Karrer *Helv Chim Acta* **32** 1371 1949; Fischer and Rund *Chem Ber* **49** 90, 93 1916; IR: Kuhn *Anal Chem* **22** 276 1950.]

***N,N'*-Diacetylchitobiose** (2-acetyl-*O*⁴-[2-acetyl-amino-2-deoxy-β-D-glucopyranosyl]-2-deoxy-D-glucose) [35061-50-8] *M* 424.4, *m* 245-247°(dec), 251.5-252.5°, 260-262°, [α]_D²⁵ +39.5° (extrapolated) → +18.5° (after 60 min, c 1, H₂O). Recrystd from aqueous MeOH or aqueous EtOH + 1,2-dimethoxyethane. [Zilliken et al. *J Chem Soc* **77** 1296 1955.]

1,8-Diazafluorenone (cyclopenta[1.2-*b*:4,3-*b'*]dipyridin-9-one) [54078-29-4] *M* 182.2, *m* 205°, 229-231°, *pK*_{Est} ~ 2.6. Recrystd from Me₂CO. The *oxime* has *m* 119-200°. [Druey and Schmid *Helv Chim Acta* **33** 1080 1950.]

Di- and tri-carboxylic acids. Resolution by anion-exchange chromatography. [Bengtsson and Samuelson *Anal Chim Acta* **44** 217b 1969.]

Digitonin [11024-24-1] *M* 1229.3, *m* >270°(dec), [α]₅₄₆²⁰ -63° (c 3, MeOH). Crystd from aqueous 85% EtOH or MeOH/diethyl ether.

Digitoxigenin [143-62-4] *M* 374.5, *m* 253°, [α]₅₄₆²⁰ +21° (c 1, MeOH). Crystd from aqueous 40% EtOH.

D(+)-Digitoxose [527-52-6] *M* 148.2, *m* 112°, [α]₅₄₆²⁰ +57° (c 1, H₂O). Crystd from MeOH/diethyl ether, or ethyl acetate.

Dihydrofolate reductase (from *Mycobacterium phlei*) [9002-03-3] *M_r* ~18,000 [EC 1.5.1.3]. Purified by ammonium sulfate pptn, then fractionated on Sephadex G-75 column, applied to a Blue Sepharose column and eluted with 1mM dihydrofolate. [Al Rubeai and Dole *Biochem J* **235** 301 1986.]

7,8-Dihydrofolic acid (7,8-dihydropteroyl-L-glutamic acid, DHFA) [4033-27-6] *M* 443.4, *pK*₁ 2.0 (basic 10-NH), *pK*₂ 2.89 (2-NH₂), *pK*₃ 3.45 (α-CO₂H), *pK*₄ 4.0 (basic 5N), *pK*₅ 4.8 (γ-CO₂H), *pK*₆ 9.54 (acidic 3NH). Best purified by suspending (1g mostly dissolved) in ice-cold sodium ascorbate (300mL of 10% at pH 6.0 [prepared by adjusting the pH of 30g of sodium ascorbate in 150mL of H₂O by adding 1N NaOH dropwise using a glass electrode till the pH is 6.0]). This gave a clear solution with pH ~5. While stirring at 0° add N HCl dropwise slowly (0.1mL/min) until the pH drops to 2.8 when white birefringent crystals separate. These are collected by centrifugation (1000xg for 5min), washed 3x with 0.001N HCl by centrifugation and decantation. The residue is then dried in a vacuum (0.02mm) over P₂O₅ (change the P₂O₅ frequently at first) and KOH at 25° in the dark. After 24hours the solid reaches constant weight.

For the assay of *dihydrofolate reductase* (see below): suspend ~66.5mg of DHFA in 10mL of 0.001M HCl containing 10mM dithiothreitol (DTT stock made from 154mg in 10mL H₂O making 0.1M), shake well and freeze in 400μL aliquots. Before use mix 400μL of this suspension with 0.1M DTT (200μL, also made in frozen aliquots), and the mixture is diluted with 200μL of 1.5M Tris-HCl pH 7.0 and 1.2mL of H₂O (making a total volume of ~2mL) to give a clear solution. To estimate the concentration of DHFA in this solution, dilute 20μL of this solution to 1mL with 0.1M Tris-HCl pH 7.0 and read the OD at 282nm in a 1cm pathlength cuvette. ε at 282nm is 28,000M⁻¹cm⁻¹. [Reyes and Rathod *Methods Enzymol* **122** 360 1986.]

Dihydropteridine reductase (from sheep liver) [9074-11-7] *M_r* 52,000 [EC 1.6.99.7]. Purified by fractionation with ammonium sulfate, dialysed *versus* tris buffer, adsorbed and eluted from hydroxylapatite gel. Then run through a DEAE-cellulose column and also subjected to Sephadex G-100 filtration. [Craine et al. *J Biol Chem* **247** 6082 1972.]

Dihydropteridine reductase (from human liver) [9074-11-7] *M_r* 52,000 [EC 1.6.99.7]. Purified to homogeneity on a naphthoquinone affinity adsorbent, followed by DEAE-Sephadex and CM-Sephadex

chromatography. [Firgaira, Cotton and Danks, *Biochem J* **197** 31 1981.] [For other dihydropteridine reductases see Armarego et al. *Med Res Rev* **4**(3) 267 1984.]

DL-erythro-Dihydrospingosine (dl-erythro-2-aminooctadecan-1,3-diol) [3102-56-5] **M 301.5, m 85-86°, 85-87°, pK_{Est} ~ 8.8.** Purified by recrystn from pet ether-EtOAc or CHCl₃. The (±)-*N*-dichloroacetyl derivative has **m 142-144° (from MeOH).** [Shapiro et al. *J Am Chem Soc* **80** 2170 1958; Shapiro and Sheradsky *J Org Chem* **28** 2157 1963.] The D-isomer crystallises from pet ether-Et₂O and has **m 78.5-79°, [α]₅₄₆²⁸ +6° (CHCl₃ + MeOH, 10:1).** [Grob and Jenny *Helv Chim Acta* **35** 2106 1953, Jenny and Grob *Helv Chim Acta* **36** 1454 1953.]

Dihydrostreptomycin sesquisulfate [5490-27-7] **M 461.4, m 250°(dec), 255-265°(dec), [α]_D²⁰ -92.4° (c 1, H₂O), pK_{Est(1)} ~ 9.5 (NMe), pK_{Est(2,3)} ~ 13.4 (guanidino).** It crystallises from H₂O with MeOH, *n*-BuOH or methyl ethyl ketone. The crystals are not hygroscopic like the amorphous powder, however both forms are soluble in H₂O but the amorphous solid is about 10 times more soluble than the crystals. The *free base* also crystallises from H₂O-Me₂CO and has [α]_D²⁶ -92° (aqueous solution pH 7.0). [Solomons and Regina *Science* **109** 515 1949; Wolf et al. *Science* **109** 515 1949; McGilveray and Rinehart *J Am Chem Soc* **87** 4003 1956].

3-(3,4-Dihydroxyphenyl)-L-alanine (DOPA, EUODOPA) [59-92-7, 5796-17-8] **M 197.2, m 275°(dec), 267-268°(dec), 284-286°(dec), ~295°(dec), [α]_D¹³ -13.1° (c 5.12, N HCl), pK₁^{2.5} 2.32 (CO₂H), pK₂^{2.5} 8.72 (NH₂), pK₃^{2.5} 9.96 (OH), pK₄^{2.5} 11.79 (OH).** Likely impurities are vanillin, hippuric acid, 3-methoxytyrosine and 3-aminotyrosine. Recryst from large vols of H₂O as colourless white needles; solubility in H₂O is 0.165%, but it is insoluble in EtOH, *C₆H₆, CHCl₃, and EtOAc. Also crystd by dissolving in dilute HCl and adding dilute ammonia to give pH 5, under N₂. Alternatively, crystd from dil aqueous EtOH. It is rapidly oxidised in air when moist, and darkens; particularly in alkaline solution. Dry in a vacuum at 70° in the dark, and store in a dark container preferably under N₂. λ_{max} 220.5nm (log ε 3.79) and 280nm (log ε 3.42) in 0.001N HCl. [Yamada et al. *Chem Pharm Bull Jpn* **10** 693 1962; Bretschneider et al. *Helv Chim Acta* **56** 2857 1973; NMR: Jardetzky and Jardetzky *J Biol Chem* **233** 383 1958.]

3,4-Dihydroxyphenylalanine-containing proteins. Boronate affinity chromatography is used in the selective binding of proteins containing 3,4-dihydroxyphenylalanine to a *m*-phenylboronate agarose column and eluting with 1M NH₄OAc at pH 10. [Hankus et al. *Anal Biochem* **150** 187 1986.]

3-(3,4-Dihydroxyphenyl)-2-methyl-L-alanine [methyldopa, 2-amino-3-(3,4-dihydroxyphenyl)-2-methylpropionic acid] [555-30-6] **M 238.2, m >300°, 300-301°(dec), pK₁^{2.5} 2.2, pK₂^{2.5} 9.2, pK₃^{2.5} 10.6, pK₄^{2.5} 12.0.** Recrystd from H₂O. [Reinhold et al. *J Org Chem* **33** 1209 1968.] The *L*-isomer forms a sesquihydrate from H₂O **m 302-304° (dec),** and the anhydrous crystals are *hygroscopic*, [α]_D²³ -4.0° (c 1, 0.1N HCl), [α]₅₄₆ +154.5° (c 5, CuSO₄ solution). It has λ_{max} 281nm (ε 2780). Solubility in H₂O at 25° is ~10mg/mL and the pH of an aqueous solution is ~5.0. It is almost insoluble in most organic solvents. [Stein et al. *J Am Chem Soc* **77** 700 1955.]

(±)-7-(2,3-Dihydroxypropyl)theophylline (Diprophylline, Dyphylline) [479-18-5] **M 254.3, m 158°, 160-164°, 161°, 161-164°, pK_{Est} ~ 8.7.** Recrystd from EtOH or H₂O. Solubility in H₂O is 33% at 25°, in EtOH it is 2% and in CHCl₃ it is 1%. λ_{max} (H₂O) 273nm (ε 8,855). [Roth *Arch Pharm* **292** 234 1959.] The 4-nitrobenzoyl derivative has **m 178°** [Oshay *J Chem Soc* 3975 1956].

3,5-Diiodo-L-thyronine (3,5-diiodo-4-[4-hydroxyphenoxy]-1-phenylalanine) [1041-01-6] **M 525.1, m 255°(dec), 255-257°(dec), [α]_D²² +26° [2N HCl-EtOH (1:2)], pK₁^{2.0} 3.25, pK₂^{2.0} 5.32, pK₃^{2.0} 9.48.** Recrystd from EtOH. [Chambers et al. *J Chem Soc* 3424 1949.]

3,5-Diiodo-L-tyrosine dihydrate [300-39-0] **M 469.0, m 199-210°, 202°(dec), [α]_D²⁰ +2.89° (c 4.9, 4% HCl), pK₁^{2.5} 2.12, pK₂^{2.5} 6.48, pK₃^{2.5} 7.82.** It forms crystals from H₂O [solubility (g/L): 0.204 at 0°, 1.86 at 50°, 5.6 at 75° and 17.0 at 100°]. Also recrystd from 50% or 70% EtOH. When boiled in EtOH the crystals swell and on further boiling a gelatinous ppte is formed [Harrington *Biochem J* **22** 1434 1928; Jurd *J Am Chem Soc* **77** 5747 1955]. Also crystd from cold dilute ammonia by adding acetic acid to pH 6.

1,2-Dilauroyl-sn-glycero-3-phosphoethanolamine (\pm -dilauroyl- α -cephalin, 3-sn-phosphatidylethanolamine 1,2-didodecanoyl) [59752-57-7] M 579.8, m 210°, pK_{Est(1)}~ 5.8 (PO₄H), pK_{Est(2)}~ 10.5 (NH₂). Recrystd from EtOH or tetrahydrofuran. [Bevan and Malkin *J Chem Soc* 2667 1951; IR: Bellamy and Beecher *J Chem Soc* 728 1953.]

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride [1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride] see entry on p. 212 in Chapter 4.

1,2-Dimyristoyl-sn-glycero-3-phosphocholine monohydrate (dimyristoyl-L- α -lecithin) [18194-24-6] M 696.0, $[\alpha]_D^{24} + 7^\circ$ (c 8, EtOH-CHCl₃ 1:1 for α_1 form), pK_{Est} ~ 5.8 (PO₄). Three forms α_1 , α_2 and β' . Recrystd from aqueous EtOH or EtOH-Et₂O. Solubility at 22-23° in Et₂O is 0.03%, in Me₂CO it is 0.06% and in pyridine it is 1.3%. [Baer and Kates *J Am Chem Soc* 72 942 1950; Baer and Maurakas *J Am Chem Soc* 74 158 1952; IR: Marinetti and Stotz *J Am Chem Soc* 76 1347 1954.] The *S*-isomer with 1 H₂O is recrystd from 2,6-dimethylheptan-4-one and has m 226-227° (sintering at 90-95°), and $[\alpha]_D - 7^\circ$ (c 6, MeOH-CHCl₃ 1:1). [Baer and Martin *J Biol Chem* 193 835 1951.]

(\pm)-1,2-Dimyristoyl-sn-glycero-3-phosphoethanolamine (dimyristoyl- α --kephalin) [998-07-2] M 635.9, m 207°, pK_{Est(1)}~ 5.8 (PO₄H), pK_{Est(2)}~ 10.5 (NH₂). Recrystd from EtOH [Bevan and Malkin *J Chem Soc* 2667 1951]. The *R*-isomer has m 195-196° (sintering at 130-135°) after recrystn from CHCl₃-MeOH, $[\alpha]_D^{26} + 6.7^\circ$ (c 8.5, CHCl₃-AcOH 9:1). [Baer *Can J Biochem Physiol* 35 239 1957; Baer et al. *J Am Chem Soc* 74 152 1952.]

***S*-1,2-Dipalmitin** [761-35-3] M 568.9, m 68-69° $[\alpha]_D^{20} - 2.9^\circ$ (c 8, CHCl₃), $[\alpha]_{546}^{20} + 1.0^\circ$ (c 10, CHCl₃/MeOH, 9:1). Crystd from chloroform/pet ether.

***R*-Dipalmitoyl-sn-glycero-3-phosphatidic acid** [7091-44-3] M 648.9, $[\alpha]_D^{26} + 4^\circ$ (c 10, CHCl₃), pK_{Est(1)}~ 1.6, pK_{Est(2)}~ 6.1. Recrystd from Me₂CO at low temp. At 21° it is soluble in *C₆H₆ (4.2%), pet ether (0.01%), MeOH (2%), EtOH (2.5%), AcOH (1.3%), Me₂CO (1.76%), and Et₂O (1.5%). [Baer *J Biol Chem* 189 235 1951.]

***R*-1,2-Dipalmitoyl-sn-glycero-3-phosphocholine monohydrate (dipalmitoyl- α -L-lecithin)** [63-89-8] M 752.1, m sinters at 120°, $[\alpha]_D^{25} + 7.0^\circ$ (c 5.6, abs CHCl₃), pK_{Est} ~ 5.8 (PO₄). It has three crystn forms α_1 , α_2 and β' which change at 60-70° and at 229° respectively. In order to obtain a fine powder, ~2 g are dissolved in CHCl₃ (15mL) and pet ether (b 35-60°) is added and the soln evaporated to dryness *in vacuo* <20°, and then dried at 0.1mm over CaCl₂. [Baer and Maurukas *J Am Chem Soc* 74 158 1952; Baer and Kates *J Biol Chem* 185 615 1950.]

***d,l*- $\beta\gamma$ -Dipalmitoylphosphatidyl choline** [2797-68-4] M 734.1, m 230-233°, pK_{Est} ~ 5.8 (PO₄). Recrystd from chloroform and dried for 48h at 10⁻⁵ torr [O'Leary and Levine *J Phys Chem* 88 1790 1984].

Dipeptidyl aminopeptidase (from rat brain) [9031-94-1] [EC 3.4.11.10]. Purified about 2000-fold by column chromatography on CM-cellulose, hydroxylapatite and Gly-Pro AH-Sepharose. [Imai et al. *J Biochem (Tokyo)* 93 431 1983.]

1,2-Distearoyl-sn-glycerol [1429-59-0] M 625.0. The *dl*-form recrystallises from CHCl₃-pet ether (b 40-60°), m 59.5° (α form) and 71.5-72.5° (β form). Recrystn from solvents (e.g. EtOH, MeOH, toluene, Et₂O) gives the higher melting form and resolidification gives the lower melting forms. [IR: Chapman *J Chem Soc* 4680 1958, 2522 1956; .] The *S*-isomer is recrystd from CHCl₃-pet ether and has m 76-77°, $[\alpha]_D^{24} - 2.8^\circ$ (c 6, CHCl₃). [Baer and Kates *J Am Chem Soc* 72 942 1950.]

1,2-Distearoyl-sn-glycero-3-phosphoethanolamine (distearoyl- α -kephalin) [1069-79-0] M 748.1, m 180-182° (*R*-form, sintering at 130-135°), m 196° (\pm form), pK_{Est(1)}~ 5.8 (PO₄H), pK_{Est(2)}~ 10.5 (NH₂). The *R*-form is recrystd from CHCl₃-MeOH and the \pm -form is recrystd from EtOH. [Bevan and Malkin *J Chem Soc* 2667 1951; Baer *Can J Biochem Physiol* 81 1758 1959.]

Dolichol (from pig liver) [11029-02-0] **C₈₀-C₁₀₅ polyprenols**. Cryst 6 times from pet ether/EtOH at -20°C. Ran as entity on a paper chromatogram on paraffin impregnated paper, with acetone as the mobile phase. [Burgos *Biochem J* **88** 470 1963.]

Domoic acid [4-(2-carboxyhexa-3,5-dienyl)-3-carboxymethylproline] [14277-97-5] **M 311.3, m 215°, 217°, [α]_D²⁰ -108° (c 1, H₂O), pK₁ 2.20 (2-CO₂H), pK₂ 3.72 (CO₂H), pK₃ 4.93 (3-CH₂CO₂H), pK₄ 9.82 (NH)**. The acid (~300 mg) is purified on a Dowex 1 column (3.5 x 40 mm, 200-400 mesh, acetate form), washed with H₂O until neutral, then eluted with increasing concentrations of AcOH (8L) from 0 to 0.25M. The fraction containing domoic acid (in 50mL) is collected, evaporated to dryness under reduced pressure and recrystd from aqueous EtOH. Glutamate and Kainate receptor agonist. [Impellizzeri et al. *Phytochemistry* **14** 1549 1975; Takemoto and Diago *Arch Pharm* **293** 627 1960.]

DNA (deoxyribonucleic acids). The essential structures of chromosomes are DNA and contain the genetic "blue print" in the form of separate genes. They are made up of the four deoxyribonucleic acids (nucleotides): adenylic acid, guanylic acid, cytidylic acid and thymidylic acid (designated A, G, C, T respectively) linked together by their phosphate groups in ester bonds between the 3' and 5' hydroxy groups of the 2'-deoxy-D-ribose moiety of the nucleotides. The chains form a double stranded spiral (helix) in which the two identical nucleotide sequences run antiparallel with the heterocyclic bases hydrogen bonded (A..T, G..C) forming the "ladder" between the strands. Short sequences of DNA are available commercially, are commercially custom made or synthesised in a DNA synthesiser and purified by HPLC. Their purity can be checked by restriction enzyme cleavage followed by gel electrophoresis, or directly by gel electrophoresis or analytical HPLC. Commercial DNAs are usually pure enough for direct use but can be further purified using commercially available kits involving binding to silica or other matrices and eluting with tris buffers.

Dopamine-β-hydroxylase (from bovine adrenal medulla) [9013-38-1] **M_r ~290,000, [EC 1.14.17.1]**. The Cu-containing glycoprotein enzyme has been isolated by two procedures. The first is an elaborate method requiring extraction, two (NH₄)₂SO₄ fractionations, calcium phosphate gel filtration, EtOH fractionation, DEAE-cellulose chromatography followed by two Sephadex-G200 gel filtrations giving enzyme with a specific activity of 65 Units/mg. [Friedman and Kaufman *J Biol Chem* **240** 4763 1965; Rush et al. *Biochem Biophys Res Commun* **61** 38 1974.] The second procedure is much gentler and provides good quality enzyme. Sedimented chromaffin vesicles were lysed in 10 volumes of 5mM K-phosphate buffer pH 6.5 using a loosely fitting Teflon-glass homogeniser. The mixture is centrifuged at 40,000xg/0.5 h and the supernatant is diluted with an equal volume of 100mM phosphate buffer (pH 6.5) containing 0.4M NaCl. This lysate is applied to a concanavalin A-Sepharose column (4 x 0.7cm) which had been equilibrated with 50 mM of phosphate buffer (pH 6.5 + 0.2M NaCl) with a flow rate of ~ 0.3 mL/min. The column is washed thoroughly with the buffer until OD_{280nm} is 0.005. The enzyme is then eluted with the same buffer containing 10% α-methyl-D-mannoside (flow rate 0.1 mL/min) and the enzyme is collected in twenty column volumes. The pooled eluate is concentrated by ultrafiltration in an Amicon Diaflo stirrer cell using an XM100A membrane. The concentrated enzyme is dialysed against 50 mM phosphate buffer (pH 6.5) containing 0.1% NaCl. The enzyme gives one band (+ two very weak band) on disc gel electrophoresis indicating better than 93% purity (67% fold purification) and has a specific activity of 5.4Units/mg. [Rush et al. *Biochem Biophys Res Commun* **57** 1301 1974; Stewart and Klinman *Ann Rev Biochem* **57** 551 1988.]

Ellipticine (5,11-dimethylpyrido[4,3:b]carbazole) [519-23-3] **M 246.3, m 311-315°(dec), 312-314°(dec), pK 5.78 (80% aq methoxyethanol)**. This DNA intercalator is purified by recrystn from CHCl₃ or MeOH and dried *in vacuo*. The UV λ_{max} values in aqueous EtOH-HCl are at 241, 249, 307, 335 and 426nm. [Marini-Bettolo and Schmutz *Helv Chim Acta* **42** 2146 1959.] The *methiodide* has **m 360°(dec)**, with UV λ_{max} (EtOH-KOH) 223, 242, 251, 311, 362 and 432nm. [Goodwin et al. *J Am Chem Soc* **81** 1903 1959.]

Enniatin A [11113-62-5] **M 681.9, m 122-122.5°, [α]_D¹⁸ -92° (c 0.9, CHCl₃)**. A cyclic peptidic ester antibiotic which is recrystd from EtOH/water but is deactivated in alkaline soln. [Ovchinnikov and Ivanov in *The Proteins* (Neurath and Hill eds) Academic Press, NY, Vol V pp. 365 and 516 1982.]

(-)-Ephedrine (1*R*,2*S*-2-methylamino-1-phenylpropanol) [299-42-3] **M 165.2**, **m** ~34°, 36°, 38.1°, **b** 126-129°/7mm, 225-227°/760mm, **d**²² 1.0085, $[\alpha]_D^{26}$ -42° (c 4, 3% HCl), $[\alpha]_D^{22.5}$ +15.1° (c 0.8, H₂O), -9.36° (c 3, MeOH), **pK**²² 9.58 (**pK**²⁵ 8.84 in 80% aq methoxyethanol). Purified by vacuum distn (dehydrates) and forms waxy crystals or granules, and may pick up 0.5 H₂O. The presence of H₂O raises its **m** to 40°. [Moore and Taber *J Amer Pharm Soc* 24 211 1935.] The anhydrous base recrystd from dry ether [Fleming and Saunders *J Chem Soc* 4150 1955]. It gradually decomposes on exposure to light and is best stored in an inert atmosphere in the dark (preferably at -20°). Sol in H₂O is 5%, in EtOH it is 1% and it is soluble in CHCl₃, Et₂O and oils. It has pKa values in H₂O of 10.25 (0°) and 8.69 (60°) [Everett and Hyne *J Chem Soc* 1136 1958; Prelog and Häflinger *Helv Chim Acta* 33 2021 1950] and pKa²⁵ 8.84 in 80% aqueous methoxyethanol [Simon *Helv Chim Acta* 41 1835 1958]. The hydrochloride has **m** 220° (from EtOH-Et₂O) and $[\alpha]_D^{20}$ -38.8° (c 2, EtOH). [IR: Chatten and Levi *Anal Chem* 31 1581 1959.] The anhydrous base crystallises from Et₂O [Fleming and Saunders *J Chem Soc* 4150 1955].

(+)-Ephedrine hydrochloride (1*S*-2*R*-2-methylamino-1-phenylpropan-1-ol hydrochloride) [24221-86-1] **M 201.7**, **m** 216-219°, $[\alpha]_D^{20}$ +34° (c 11.5, H₂O). Recrystd from EtOH-Et₂O. The free base recrystallises from *C₆H₆ with **m** 40-41° (Skita et al. *Chem Ber* 66 974 1933).

Erythromycin A [114-07-8] **M 733.9**, **m** 133-135°(dec), 135-140°, 137-140°, $[\alpha]_D^{20}$ -75° (c 2, EtOH), **pK** 8.9. It recrystallises from H₂O to form hydrated crystals which melt at ca 135-140°, resolidifies and melts again at 190-193°. The **m** after drying at 56°/8mm is that of the anhydrous material at 137-140°. Its solubility in H₂O is ~2mg/mL. The Hydrochloride has **m** 170°, 173° (from aq EtOH, EtOH-Et₂O). [Flynn et al. *J Am Chem Soc* 76 3121 1954; constitution : Wiley et al. *J Am Chem Soc* 79 6062 1957].

β-Estradiol (1,3,5-estratrien-3,17β-diol) [50-28-2] **M 272.4**, **m** 173-179°, 176-178°, $[\alpha]_D^{20}$ +76° to +83° (c 1, dioxane). Purified by chromatography on SiO₂ (toluene-EtOAc 4:1) and recrystd from CHCl₃-hexane or 80% EtOH. It is stable in air and insoluble in H₂O and is ppted by digitonin. UV λ_{max} at 225 and 280 nm. [Oppolzer and Roberts *Helv Chim Acta* 63 1703 1980.]

β-Estradiol-6-one (1,3,5-estratriene-3,17β-diol-6-one) [571-92-6] **M 359.4**, **m** 278-280°, 281-283°, $[\alpha]_D^{20}$ +4.2° (c 0.7, EtOH). It forms plates from EtOH. The 3,17-diacetate has **m** 173-175° after recrystn from aqueous EtOH. [Longwell and Wintersteiner *J Biol Chem* 133 219 1940.] The UV has λ_{max} 255 and 326nm in EtOH [Slaunwhite et al. *J Biol Chem* 191 627 1951].

Ethidium bromide [1239-45-8] **M 384.3**, **m** 260-262°. Crystd from MeOH or EtOH [Lamos et al. *J Am Chem Soc* 108 4278 1986]. Sol in H₂O is 1%. **POSSIBLE CARCINOGEN.**

Ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline) [91-53-2] **M 217.3**, **b** 169°/12-13mm, **d**₄²⁰ 1.000, **pK**_{Est} ~ 5.8. Purified by fractional distn *in vacuo* and solidifies to a glass. [Knoevenagel *Chem Ber* 54 1723, 1730 1921]. The methiodide has **m** 179° (from EtOH) and the 1-phenylcarbamoyl derivative has **m** 146-147° (from EtOH). [Beaver et al. *J Am Chem Soc* 79 1236 1957.]

17-α-Ethynylestradiol [57-63-6] **M 296.4**, **m** 141-146°, 145-146°, $[\alpha]_D^{20}$ +4° (c 1, CHCl₃). It forms a hemihydrate on recrystn from MeOH-H₂O. It dehydrated on melting and re-melts on further heating at **m** 182-184°. UV λ_{max} at 281nm (ε 2040) in EtOH. Solubility is 17% in EtOH, 25% in Et₂O, 20% in Me₂CO, 25% in dioxan and 5% in CHCl₃. [Petit and Muller *Bull Soc Chim Fr* 121 1951.] The diacetyl derivative has **m** 143-144° (from MeOH) and $[\alpha]_D^{20}$ +1° (c 1, CHCl₃) [Mills et al. *J Am Chem Soc* 80 6118 1958].

Exonucleases. Like the endonucleases they are restriction enzymes which act at the 3' or 5' ends of linear DNA by hydrolysing off the nucleotides. Although they are highly specific for hydrolysing nucleotides at the 3' or 5' ends of linear DNA, the number of nucleotides cleaved are time dependent and usually have to be estimated from the time allocated for cleavage. Commercially available exonucleases are used without further purification.

Farnesol (*trans-trans-3,7,11-trimethyl-2,6,10-dodecatrien-1-ol*) [106-28-5] **M 222.4, b 111°/0.35mm, 126-127°/0.5mm, 142-143°/2mm, d_4^{20} 0.8871, n_D^{25} 1.4870.** Main impurity is the *cis-trans* isomer. Purified by gas chromatography using a 4ft x 0.125in 3%OV-1 column at 150°. [Corey et al. *J Am Chem Soc* **92** 6637 1970; Popjak et al. *J Biol Chem* **237** 56 1962.] Also purified through a 14-in Podbielniak column at 11°/0.35mm (see p. 141). Alternatively it has been purified by gas chromatography using SF96 silicone on Fluoropak columns or Carbowax 20M on Fluoropak or base-washed 30:60 firebrick (to avoid decomp of alcohol, prepared by treating the firebrick with 5N NaOH in MeOH and washed with MeOH to pH 8) at 210° with Helium carrier gas at 60 mL/min flow rate. The *diphenylcarbamoyl* derivative has **m 61-63°** (from MeOH) and has IR band at 3500 cm^{-1} . [Bates et al. *J Org Chem* **28** 1086 1963.]

Farnesyl pyrophosphate [13058-04-3; *E,E*: 372-97-4] **M 382.3, $\text{pK}_{\text{Est}(1)} \sim < 2$, $\text{pK}_{\text{Est}(2)} \sim < 2$, $\text{pK}_{\text{Est}(3)} \sim 3.95$, $\text{pK}_{\text{Est}(4)} \sim 6.26$.** Purified by chromatography on Whatman No3 MM paper in a system of isopropanol-isobutanol-ammonia-water (40:20:1:30) (v/v). Stored as the Li or NH_4 salt at 0°.

Ferritin (from human placenta) [9007-73-2] **$M_r \sim 445,000$ (Fe free protein).** The purification of this major iron binding protein was achieved by homogenisation in water and precipitating with ammonium sulfate, repeating the cycle of ultracentrifuging and molecular sieve chromatography through Sephadex 4B column. Isoelectric focusing revealed a broad spectrum of impurities which were separated by ion-exchange chromatography on Sephadex A-25 and stepwise elution. [Konijn et al. *Anal Biochem* **144** 423 1985.]

Fibrinogen (from human plasma) [9001-32-5] **M_r 341,000.** A protein made up of $2\text{A}\alpha$, $2\text{B}\beta$ and 2γ subunits connected by disulfide bridges. Possible impurity is plasminogen. Purified by glycine pptn [Mosesson and Sherry *Biochemistry* **5** 2829 1966] to obtain fractions 1-2, then further purified [Blombäck and Blombäck *Arkiv Kemi* **10** 415 1956] and contaminating plasminogen is removed by passage through a lysine-Sepharose column. Such preparations were at least 95% clottable as determined by Mosesson and Sherry's method (above ref.) in which the OD_{280} was measured before and after clotting with 5 Units/mL of thrombin ($> 3000\text{U}/\text{mg}$). All fibrinogen preps were treated with calf intestinal alkaline phosphatase to convert any fibrinogen peptide-AP to fibrinogen peptide-A by removing serine-bound phosphate. Solutions are then lyophilised and stored at -20°. [Higgins and Shafer *J Biol Chem* **256** 12013 1981.] It is sparingly soluble in H_2O . Aqueous solns are viscous with isoelectric point at pH 5.5. Readily denatured by heating above 56° or by chemical agents, e.g. salicylaldehyde, naphthoquinone sulfonates, ninhydrin or alloxan. [Edsall et al. *J Am Chem Soc* **69** 2731 1947; Purification: Cama et al. *Naturwissenschaften* **48** 574 1961; Lorand and Middlebrook *Science* **118** 515 1953; cf. Fuller in *Methods Enzymol* **163** 474 1988.]

For plasminogen-deficient fibrinogen from blood plasma, the anticoagulated blood was centrifuged and the plasma was frozen and washed with saline solution. Treated with charcoal and freeze-thawed. Dialysed *versus* Tris/NaCl buffer. [Maxwell and Nikel *Biochem Prep* **12** 16 1968.]

Fibronectin (from human plasma) [86088-83-7] **$M_r \sim 220,000$.** This glycoprotein contains 5-12% of carbohydrate. It has been purified by glycine fractionation and DEAE-cellulose chromatography. Material is dissolved in 0.25M Tris-phosphate buffer pH 7.0, diluted to 20% and glycine added gradually till 2.1M when the temperature falls to below 15°. The ppte contains mainly fibrinogen. The supernatant is discarded and the ppte is treated with an equal volume of H_2O , cooled (to 0°) and ppted by adding EtOH to 16% (v/v) at -4°. The ppte contains some CI (Cold Insoluble) globulin, fibronectin and small quantities of other proteins. To remove these the ppte is dissolved in 0.25M Tris-phosphate buffer (pH 7.0) *ca* 0.5% and purified by DEAE-cellulose chromatography after diluting the buffer to 0.05M buffer. [Morrison et al. *J Am Chem Soc* **70** 3103 1948; Mosesson and Umfleet *J Biol Chem* **245** 5728 1970; Mosesson and Amrani *Blood* **56** 145 1980; Akiyama and Yamada *Adv Enzymol* **59** 51 1987.]

Flavin adenine dinucleotide (di-Na, $2\text{H}_2\text{O}$ salt, FAD) [146-14-5] **M 865.6, $[\alpha]_{546} -54^\circ$ (c 1, H_2O).** Small quantities, purified by paper chromatography using *tert*-butyl alcohols/water, cutting out the main spot and eluting with water. Larger amounts can be ppted from water as the uranyl complex by adding a slight excess of uranyl acetate to a soln at pH 6.0, dropwise and with stirring. The soln is set aside overnight in the cold, and the ppte is centrifuged off, washed with small portions of cold EtOH, then with cold, peroxide-free diethyl ether. It is dried in the dark under vacuum over P_2O_5 at 50-60°. The uranyl complex is suspended in water and, after adding sufficient 0.01M NaOH to adjust the pH to 7, the ppte of uranyl hydroxide is removed by

centrifugation [Huennekens and Felton *Methods Enzymol* 3 954 1957]. It can also be crystd from water. Should be kept in the dark. More recently it was purified by elution from a DEAE-cellulose (Whatman DE 23) column with 0.1M phosphate buffer pH 7, and the purity was checked by TLC. [Holt and Cotton, *J Am Chem Soc* 109 1841 1987.]

Flavin mononucleotide (Na, 2H₂O salt, FMN) [130-40-5] M 514.4, pK₁ 2.1 (PO₄H₂), pK₂ 6.5 (PO₄H⁻), pK₃ 10.3 (CONH), fluorescence λ_{max} 530nm (870nm for reduced form). Purified by paper chromatography using *tert*-butanol-water, cutting out the main spot and eluting with water. Also purified by adsorption onto an apo-flavodoxin column, followed by elution and freeze drying Crystd from acidic aqueous soln. [Mayhew and Strating *Eur J Biochem* 59 539 1976.]

4-Fluoro-7-nitrobenzofurazan (4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole) [29270-56-2] M 183.1, m 52.5-53.5°, 53-56°, 53.5-54.5°. Purified by repeated recrystn from pet ether (b 40-60°). On treatment with MeONa in MeOH it gave *4-methoxy-7-nitrobenzo-2-oxa-1,3-diazole* m 115-116°. [Nunno et al. *J Chem Soc (C)* 1433 1970.] It is a very good fluorophore for amino acids [Imai and Watanabe *Analyt Chim Acta* 130 377 1981], as it reacts with primary and secondary amines to form fluorescent adducts with λ_{ex} 470nm and λ_{em} 530nm. It gives a *glycine derivative* with m 185-187° [Miyano et al. *Anal Chim Acta* 170 81 1985].

4-Fluoro-3-nitrophenylazide [28166-06-5] M 182.1, m 53-55°, 54-56°. Dissolve in Et₂O, dry over MgSO₄, filter, evaporate and recryst the residue from pet ether (b 20-40°) to give orange needles. Store in a stoppered container at ~0°. The NMR has δ 7.75 (m 1H) and 7.35 (m 2H) in CDCl₃. [Hagedorn et al. *J Org Chem* 43 2070 1978.]

2-Fluorophenylalanine [*R*(+)- 97731-02-7; *S*(-)- 19883-78-4] M 183.2, m 226-232°, 231-234°, [α]_D²⁵ (+) and (-) 15° (c 2, H₂O pH 5.5), pK₁²⁴ 2.12, pK₂²⁴ 9.01. Recryst from aqueous EtOH. The *hydrochloride* has m 226-231°(dec), and the *N-acetyl* derivative has m 147-149° (from aqueous EtOH). [Bennett and Nieman *J Am Chem Soc* 72 1800 1950.]

4-Fluorophenylalanine [*R*(+)- 18125-46-7; *S*(-)- 1132-68-9] M 183.2, m 227-232°, [α]_D²⁵ (+) and (-) 24° (c 2, H₂O), pK₁²⁴ 2.13, pK₂²⁴ 9.05. It is recrystd from aqueous EtOH. The *R-N-acetyl* derivative has m 142-145°, [α]_D²⁵ -38.6° (c 8, EtOH). [Bennett and Nieman *J Am Chem Soc* 72 1800 1950.]

5-Fluoro-L-tryptophan monohydrate [16626-02-1] M 240.2, m >250°(dec), [α]_D²⁰ +5.5° (c 1, 0.1N HCl), pK_{Est(1)}~ 2.5 (CO₂H), pK_{Est(2)}~ 9.4 (NH₂), pK_{Est(3)}~16 (indole-NH). Recrystd from aqueous EtOH.

5-Fluorouridine (5-fluoro-1-β-D-ribofuranosyl-1H-pyrimidine-2,4-dione) [316-46-1] M 262.2, m 180-182°, 182-184°, [α]_D²⁰ +18° (c 1, H₂O), pK_{Est(1)}~ 8.0, pK_{Est(2)}~ 13. Recrystd from EtOH-Et₂O and dried at 100° in a vacuum. UV: λ_{max} 269nm (pH 7.2, H₂O), 270nm (pH 14, H₂O). [Liang et al. *Mol Pharmacol* 21 224 1982.]

5-Fluorouracil (5-fluoropyrimidinedi-2,4-[1H,3H]-one) [51-21-8] M 130.1, m 282-283°(dec), 282-286°(dec), pK₁²⁵ 8.04, pK₂²⁵ 13.0. Recrystd from H₂O or MeOH-Et₂O, and sublimed at 190-200°/0.1mm or 210-230°/0.5mm. UV: λ_{max} 265-266nm (ε 7070). [Barton et al. *J Org Chem* 37 329 1972; Duschinsky and Plevin *J Am Chem Soc* 79 4559 1957.]

Fluram (Fluorescamine, 4-phenyl-spiro[furan-2(3H)-1-phthalan]-3,3'-dione) [38183-12-9] M 278.3, m 153-155°, 154-155°. A non-fluorescent reagent that reacts with primary amines to form highly fluorescent compounds. Purified by dissolving (~1g) in Et₂O-*C₆H₆ (1:1, 180 mL), wash with 1% aq NaHCO₃ (50mL), dry (Na₂SO₄), evaporate in a vacuum. Dissolve the residue in warm CH₂Cl₂ (5mL), dilute with Et₂O (12mL) and refrigerate. Collect the solid and dry in a vacuum. IR (CHCl₃): ν 1810, 1745, 1722, 1625 and 1600 cm⁻¹, and NMR (CDCl₃): δ 8.71 (s, -OHC=). [Weigle et al. *J Am Chem Soc* 94 5927 1972, *J Org Chem* 41 388 1976; *Methods Enzymol* 47 236 1977.]

Folic acid (FA, pteroyl-S-glutamic acid) [75708-92-8] M 441.4, m >250°(dec), $[\alpha]_D^{25} +23^\circ$ (c 0.5, 0.1N NaOH), pK_1 2.35 (protonation N10), pK_2 2.75 (protonation N1), pK_3 3.49 (α -CO₂H), pK_4 4.65 (γ -CO₂H), pK_5 8.80 (acidic N3). If paper chromatography indicates impurities then recrystallise from hot H₂O or from dilute acid [Walker et al. *J Am Chem Soc* **70** 19 1948]. Impurities may be removed by repeated extraction with *n*-BuOH of a neutral aqueous solns of folic acid (by suspending in H₂O and adding N NaOH till the solid dissolves then adjusting the pH to ~7.0-7.5) followed by pptn with acid, filtration, and recrystn from hot H₂O. [Blakley *Biochem J* **65** 331 1975; Kalifa, Furrer, Bieri and Viscontini *Helv Chim Acta* **61** 2739 1978.] Chromatography on cellulose followed by filtration through charcoal has also been used to obtain pure acid. [Sakami and Knowles *Science* **129** 274 1959.] UV: λ_{max} 247 and 296nm (ϵ 12800 and 18700) in H₂O pH 1.0; 282 and 346nm (ϵ 27600 and 7200) in H₂O pH 7.0; 256, 284 and 366nm (ϵ 24600, 24500 and 8600) in H₂O pH 13 [Rabinowitz in *The Enzymes* (Boyer et al. Eds **2** 185 1960).

Follicle Stimulating Hormone (FSH, follitropin) [9002-68-0] M_r ~36,000. Purified by Sephadex G100 gel filtration followed by carboxymethyl-cellulose with NH₄OAc pH 5.5. The latter separates luteinising hormone from FSH. Solubility in H₂O is 0.5%. It has an isoelectric point of 4.5. A soln of 1mg in saline (100mL) can be kept at 60° for 0.5h. Activity is retained in a soln at pH 7-8 for 0.5h at 75°. The activity of a 50% aq EtOH soln is destroyed at 60° in 15 min. [Bloomfield et al. *Biochim Biophys Acta* **533** 371 1978; Hartree *Biochem J* **100** 754 1966; Pierce and Parsons *Ann Rev Biochem* **50** 465 1981.]

Fructose-1,6-diphosphate (trisodium salt) [38099-82-0] M 406.1, pK_3^{25} 6.14, pK_4^{25} 6.93 (free acid). For purification *via* the acid strychnine salt, see Neuberg, Lustig and Rothenberg [*Arch Biochem* **3** 33 1943]. The calcium salt can be partially purified by soln in ice-cold M HCl (1g per 10mL) and repptn by dropwise addition of 2M NaOH: the ppte and supernatant are heated on a boiling water bath for a short time, then filtered and the ppte is washed with hot water. The magnesium salt can be pptd from cold aqueous soln by adding four volumes of EtOH.

Fructose-6-phosphate [643-13-0] M 260.1, $[\alpha]_D^{21} +2.5$ (c 3, H₂O), pK^{25} 5.84. Crystd as the barium salt from water by adding four volumes of EtOH. The barium can be removed by passage through the H⁺ form of a cation exchange resin and the free acid collected by freeze-drying.

6-Furfurylaminopurine (Kinetin) [525-79-1] M 215.2, m 266-267°, 269-271°, 270-272°, 272° (sealed capillary), $pK_1 <1$, pK_2 3.8, pK_3 10. Platelets from EtOH and sublimes at 220°, but is best done at lower temperatures in a good vacuum. It has been extracted from neutral aqueous solns with Et₂O. [Miller et al *J Am Chem Soc* **78** 1375 1956; Bullock et al. *J Am Chem Soc* **78** 3693 1956.]

Fusaric acid (5-*n*-butylpyridine-2-carboxylic acid) [536-69-6] M 179.2, m 96-98°, 98°, 98-100°, 101-103°, pK_1 5.7, pK_2 6.16 (80% aq methoxyethanol). Dissolve in CHCl₃, dry (Na₂SO₄), filter, evaporate and recrystallise the residue from 50 parts of pet ether (b 40-60°) or EtOAc, then sublime *in vacuo*. The *copper salt* forms bluish violet crystals from H₂O and has m 258-259°. [Hardegger and Nikles *Helv Chim Acta* **39** 505 1956; Schreiber and Adam *Chem Ber* **93** 1848 1960; NMR and MS: Tschesche and Führer *Chem Ber* **111** 3500 1978.]

Fuschin (Magenta I, rosaniline HCl) [632-99-5] M 337.9, m >200°(dec). See rosaniline hydrochloride on p. 349 in Chapter 4.

D-Galactal [21193-75-9] M 146.2, m 100°, 100-102°, 104°, 103-106°, $[\alpha]_D^{20} -21.3^\circ$ (c 1, MeOH). Recryst from EtOAc, EtOH or EtOAc + MeOH. [Overend et al. *J Chem Soc* 675 1950; Wood and Fletcher *J Am Chem Soc* **79** 3234 1957; Distler and Jourdian *J Biol Chem* **248** 6772 1973.]

Folic acid (FA, pteroyl-S-glutamic acid) [75708-92-8] M 441.4, m >250°(dec), $[\alpha]_D^{25} +23^\circ$ (c 0.5, 0.1N NaOH), pK_1 2.35 (protonation N10), pK_2 2.75 (protonation N1), pK_3 3.49 (α -CO₂H), pK_4 4.65 (γ -CO₂H), pK_5 8.80 (acidic N3). If paper chromatography indicates impurities then recrystallise from hot H₂O or from dilute acid [Walker et al. *J Am Chem Soc* **70** 19 1948]. Impurities may be removed by repeated extraction with *n*-BuOH of a neutral aqueous solns of folic acid (by suspending in H₂O and adding N NaOH till the solid dissolves then adjusting the pH to ~7.0-7.5) followed by pptn with acid, filtration, and recrystn from hot H₂O. [Blakley *Biochem J* **65** 331 1975; Kalifa, Furrer, Bieri and Viscontini *Helv Chim Acta* **61** 2739 1978.] Chromatography on cellulose followed by filtration through charcoal has also been used to obtain pure acid. [Sakami and Knowles *Science* **129** 274 1959.] UV: λ_{max} 247 and 296nm (ϵ 12800 and 18700) in H₂O pH 1.0; 282 and 346nm (ϵ 27600 and 7200) in H₂O pH 7.0; 256, 284 and 366nm (ϵ 24600, 24500 and 8600) in H₂O pH 13 [Rabinowitz in *The Enzymes* (Boyer et al. Eds **2** 185 1960).

Follicle Stimulating Hormone (FSH, follitropin) [9002-68-0] M_r ~36,000. Purified by Sephadex G100 gel filtration followed by carboxymethyl-cellulose with NH₄OAc pH 5.5. The latter separates luteinising hormone from FSH. Solubility in H₂O is 0.5%. It has an isoelectric point of 4.5. A soln of 1mg in saline (100mL) can be kept at 60° for 0.5h. Activity is retained in a soln at pH 7-8 for 0.5h at 75°. The activity of a 50% aq EtOH soln is destroyed at 60° in 15 min. [Bloomfield et al. *Biochim Biophys Acta* **533** 371 1978; Hartree *Biochem J* **100** 754 1966; Pierce and Parsons *Ann Rev Biochem* **50** 465 1981.]

Fructose-1,6-diphosphate (trisodium salt) [38099-82-0] M 406.1, pK_3^{25} 6.14, pK_4^{25} 6.93 (free acid). For purification *via* the acid strychnine salt, see Neuberg, Lustig and Rothenberg [*Arch Biochem* **3** 33 1943]. The calcium salt can be partially purified by soln in ice-cold M HCl (1g per 10mL) and repptn by dropwise addition of 2M NaOH: the ppte and supernatant are heated on a boiling water bath for a short time, then filtered and the ppte is washed with hot water. The magnesium salt can be pptd from cold aqueous soln by adding four volumes of EtOH.

Fructose-6-phosphate [643-13-0] M 260.1, $[\alpha]_D^{21} +2.5$ (c 3, H₂O), pK^{25} 5.84. Crystd as the barium salt from water by adding four volumes of EtOH. The barium can be removed by passage through the H⁺ form of a cation exchange resin and the free acid collected by freeze-drying.

6-Furfurylaminopurine (Kinetin) [525-79-1] M 215.2, m 266-267°, 269-271°, 270-272°, 272° (sealed capillary), $pK_1 <1$, pK_2 3.8, pK_3 10. Platelets from EtOH and sublimes at 220°, but is best done at lower temperatures in a good vacuum. It has been extracted from neutral aqueous solns with Et₂O. [Miller et al *J Am Chem Soc* **78** 1375 1956; Bullock et al. *J Am Chem Soc* **78** 3693 1956.]

Fusaric acid (5-*n*-butylpyridine-2-carboxylic acid) [536-69-6] M 179.2, m 96-98°, 98°, 98-100°, 101-103°, pK_1 5.7, pK_2 6.16 (80% aq methoxyethanol). Dissolve in CHCl₃, dry (Na₂SO₄), filter, evaporate and recrystallise the residue from 50 parts of pet ether (b 40-60°) or EtOAc, then sublime *in vacuo*. The copper salt forms bluish violet crystals from H₂O and has m 258-259°. [Hardegger and Nikles *Helv Chim Acta* **39** 505 1956; Schreiber and Adam *Chem Ber* **93** 1848 1960; NMR and MS: Tschesche and Führer *Chem Ber* **111** 3500 1978.]

Fuschin (Magenta I, rosaniline HCl) [632-99-5] M 337.9, m >200°(dec). See rosaniline hydrochloride on p. 349 in Chapter 4.

D-Galactal [21193-75-9] M 146.2, m 100°, 100-102°, 104°, 103-106°, $[\alpha]_D^{20} -21.3^\circ$ (c 1, MeOH). Recryst from EtOAc, EtOH or EtOAc + MeOH. [Overend et al. *J Chem Soc* 675 1950; Wood and Fletcher *J Am Chem Soc* **79** 3234 1957; Distler and Jourdian *J Biol Chem* **248** 6772 1973.]

β -Galactosidase (from bovine testes) [9031-11-2] M_r 510,000, [EC 3.2.1.23]. Purified 600-fold by ammonium sulfate precipitation, acetone fractionation and affinity chromatography on agarose substituted with terminal thio- β -galactopyranosyl residues. [Distlern and Jourdian *J Biol Chem* 248 6772 1973.]

Gangcyclovir [9-{(1,3-dihydroxy-2-propoxy)methyl}guanine; 2-amino-1,9-{(2-hydroxy-1-hydroxymethyl)-ethoxymethyl}-6H-purin-6-one; Cytovene; Cymeva(e)n(e)] [82410-32-0] M 255.2, m >290°(dec), >300°(dec), monohydrate m 248-249°(dec), $pK_{Est(1)} \sim -1.1$, $pK_{Est(2)} \sim 4.1$, $pK_{Est(3)} \sim 9.7$. Recryst from MeOH. Alternatively dissolve ~90g of reagent in 700mL of distilled H₂O, filter and cool (ca 94% recovery). UV: λ_{max} in MeOH 254nm (ϵ 12,880), 270sh nm (ϵ 9040), solubility in H₂O at 25° is 4.3mg/mL at pH 7.0. **ANTIVIRAL**. [Ogilvie et al. *Can J Chem* 60 3005 1982; Ashton et al. *Biochem Biophys Res Commun* 108 1716 1982; Martin et al. *J Med Chem* 26 759 1983.]

Geranylgeranyl pyrophosphate [6699-20-3 (NH₄ salt)] M 450.5, $pK_{Est(1)} \sim <2$, $pK_{Est(2)} \sim <2$, $pK_{Est(3)} \sim 3.95$, $pK_{Est(4)} \sim 6.26$. Purified by counter-current distribution between two phases of a butanol/isopropyl ether/ammonia /water mixture (15:5:1:19) (v/v), or by chromatography on DEAE-cellulose (linear gradient of 0.02M KCl in 1mM Tris buffer, pH 8.9). Stored as a powder at 0°.

Geranyl pyrophosphate [763-10-0 (NH₄ salt)] M 314.2, $pK_{Est(1)} \sim <2$, $pK_{Est(2)} \sim <2$, $pK_{Est(3)} \sim 3.95$, $pK_{Est(4)} \sim 6.26$. Purified by paper chromatography on Whatman No 3 MM paper in a system of isopropyl alcohol/isobutyl alcohol/ammonia/water (40:20:1:39), R_F 0.77-0.82. Stored in the dark as the ammonium salt at 0°.

Gitoxigenin (3 β ,14,16 β ,21-tetrahydroxy-20(22)norcholenic acid lactone) [545-26-6] M 390.5, m 223-226°, 234°, 239-240° (anhydrous by drying at 60°), $[\alpha]_D^{20} +30^\circ$ (c 1, MeOH). Recrystn from aqueous EtOH produces plates of the sesquihydrate which dehydrate on drying at 100° *in vacuo*. It has also been recrystd from Me₂CO-MeOH and from EtOAc the crystals contain 1 mol of EtOAc with $[\alpha]_D^{21} +24.8^\circ$ (c 1, dioxane). It has UV has λ_{max} at 310, 485 and 520nm in 96% H₂SO₄. On heating with ethanolic HCl it yields *digitaligenin* with loss of H₂O. [Smith *J Chem Soc* 23 1931.]

Gliotoxin (3R-6t-hydroxy-3-hydroxymethyl-2-methyl-(5at)-2,3,6,10-tetrahydro-5aH-3,10ac-epidissulfido[1,2-a]-indol-1,4-dione) [67-99-2] M 326.4, m 191-218°(dec), 220°(dec), 221°(dec), $[\alpha]_D^{20} -254^\circ$ (c 0.6, CHCl₃), $[\alpha]_D^{25} -270^\circ$ (c 1.7, pyridine). Purified by recrystn from MeOH. Its solubility in CHCl₃ is 1%. The *dibenzoyl* derivative has m 202° (from CHCl₃-MeOH). [Glister and Williams *Nature* 153 651 1944; Elvidge and Spring *J Chem Soc Suppl* 135 1949; Johnson et al. *J Am Chem Soc* 65 2005 1943; Bracken and Raistrick *Biochem J* 41 569 1947.]

Glucose oxidase (from Aspergillus niger) [9001-37-0] M_r 186,000, [EC 1.1.3.4]. Purified by dialysis against deionized water at 6° for 48hours, and by molecular exclusion chromatography with Sephadex G-25 at room temperature. [Holt and Cotton *J Am Chem Soc* 109 1841 1987.]

Glucose-1-phosphate [59-56-3] M 260.1, $[\alpha]_D^{25} +120^\circ$ (c 3, H₂O), $[\alpha]_D^{20} +78^\circ$ (c 4, H₂O of di-K salt), pK_1 1.11, pK_2 6.13 [pK^{25} 6.50]. Two litres of 5% aq soln was brought to pH 3.5 with glacial acetic acid (+ 3g of charcoal, and filtered). An equal volume of EtOH was added, the pH was adjusted to 8.0 (glass electrode) and the soln was stored at 3° overnight. The ppt was filtered off, dissolved in 1.2L of distd water, filtered and an equal volume of EtOH was added. After standing at 0° overnight, the crystals were collected at the centrifuge, and washed with 95% EtOH, then absolute EtOH, ethanol/diethyl ether (1:1), and diethyl ether. [Sutherland and Wosilait, *J Biol Chem* 218 459 1956.] Its barium salt can be crystd from water and EtOH. Heavy metal impurities can be removed by passage of an aqueous soln (ca 1%) through an Amberlite IR-120 column (in the appropriate H⁺, Na⁺ or K⁺ forms). *Di-K salt* cryst as 2H₂O from EtOH.

Glucose-6-phosphate [acid 156-73-5; Ba salt 58823-95-3; Na salt 54010-71-8] M 260.1, m 205-207°(dec) mono Na salt, $[\alpha]_{546}^{20} +41^\circ$ (c 5, H₂O), pK_1 1.65, pK_2 6.11, pK_3^{25} 11.71 [-C₁(OH)O⁻]. Can be freed from metal impurities as described for glucose-1-phosphate. Sol of Na Salt is 5% in H₂O at 20°. Its barium salt can be purified by solution in dilute HCl and pptn by neutralising the soln. The ppt is washed with small volumes of cold water and dried in air.

Glucose-6-phosphate dehydrogenase [9001-40-5] M_r 128,000 (from Baker's yeast), 63,300 (from rat mammary gland) [EC 1.1.1.49]. The enzyme is useful for measuring pyridine nucleotides in enzyme recycling. The enzyme from Baker's yeast has been purified by $(\text{NH}_4)_2\text{SO}_4$ fractionation, Me_2CO pptn, a second $(\text{NH}_4)_2\text{SO}_4$ fractionation, concentration by DEAE-SF chromatography, a third $(\text{NH}_4)_2\text{SO}_4$ fractionation and recrystn. Crystn is induced by addition of its coenzyme NADP, which in its presence causes rapid separation of crystals at $(\text{NH}_4)_2\text{SO}_4$ concentration much below than required to ppt the amorphous enzyme. To recryst, the crystals are dissolved in 0.01M NADP (pH 7.3) with $(\text{NH}_4)_2\text{SO}_4$ at 0.55 saturation and the crystals appear within 10 to 60 min. After standing for 2-3 days (at 4°) the $(\text{NH}_4)_2\text{SO}_4$ is increased to 0.60 of saturation and more than 80% of the activity in the original crystals is recovered in the fresh crystals. [Noltmann et al. *J Biol Chem* 236 1255 1961]. Large amounts can be obtained from rat livers. The livers are extracted with 0.025M phosphate buffer (pH 7.5), and ppted with 3M $(\text{NH}_4)_2\text{SO}_4$ (70% of activity). The ppte is dissolved in 3volumes of 0.025M phosphate (pH 7.5), dialysed against this buffer + 0.2mM EDTA at 4° for 5h, then diluted to 1% protein and the nucleic acids ppted by addition of 0.4volumes of 1% protamine sulfate. $(\text{NH}_4)_2\text{SO}_4$ is added to a concentration of 2M (pH adjusted to 7.0 with NH_3), the ppte is discarded and the supernatant is adjusted to 2.8M $(\text{NH}_4)_2\text{SO}_4$, dialysed, protein adjusted to 1% and treated with $\text{Ca}_3(\text{PO}_4)_2$ gel. The gel is added in three steps (1.5mL of 0.4% gel/mL per step) and the gel is removed by centrifugation after each addn. The third gel adsorbed 50% of the activity. The gel is eluted with 0.2M phosphate (pH 7.4, 40mL/g of gel; 60% recovery). The extract is ppted in 3volumes with $(\text{NH}_4)_2\text{SO}_4$ (adjusted to 4M) to give enzyme with an activity of 30 μ moles/mg of protein x hour. [Lowry et al. *J Biol Chem* 236 2746 1961.] K_m values for the yeast enzyme are 20 μ M for G-6P and 2 μ M for NADP (Tris pH 8.0, 10⁻² M MgCl_2 , 38°) [Noltmann and Kuby *The Enzymes* VII 223 1963].

L-Glutathione (reduced form, γ -L-glutamyl-L-cysteinyl-glycine) [70-18-8] M 307.3, m 188-190°(dec), 195°(dec), $[\alpha]_D^{20}$ -20.1° (c 1, H_2O), $\text{pK}_1^{2.5}$ 2.12 (CO_2H), $\text{pK}_2^{2.5}$ 3.59 (CO_2H), $\text{pK}_3^{2.5}$ 8.75 (NH_2), $\text{pK}_4^{2.5}$ 9.65 (10.0, SH). Crystd from 50% aq EtOH, dry in a vac and store below 5°. Alternatively recrystd from aqueous EtOH under N_2 , and stored dry in a sealed container below 4°. It is soluble in H_2O . [Weygand and Geiger *Chem Ber* 90 634 1957; Martin and Edsall *Bull Soc Chim Fr* 40 1763 1958; *Biochem Prep* 2 87 1952.]

L-Glutathione (oxidised) [27025-41-8] M 612.6, m 175-195°, 195°, $[\alpha]_D^{20}$ -98° (c 2, H_2O), pK_1 3.15, pK_2 4.03, pK_3 8.75. Purified by recrystn from 50% aq EtOH. Its solubility in H_2O is 5%. Store at 4°. [Li et al. *J Am Chem Soc* 76 225 1954; Berse et al. *Can J Chem* 37 1733 1959.]

Glutathione S-transferase (human liver) [50812-37-8] M_r 25,000, [EC 2.5.1.18]. Purified by affinity chromatography using a column prepared by coupling glutathione to epoxy-saturated Sepharose. After washing contaminating proteins the pure transferase is eluted with buffer containing reduced glutathione. The solution is then concentrated by ultrafiltration, dialysed against phosphate buffer at pH ~7 and stored in the presence of dithiothreitol (2mM) in aliquots at <-20°. [Simons and Vander Jag *Anal Biochem* 52 334 1977.]

Glyceraldehyde-3-phosphate dehydrogenase [9001-50-7] M_r 144,000, [EC 1.2.1.12]. Purified from rabbit muscle by extraction with 0.03N KOH and ppted with $(\text{NH}_4)_2\text{SO}_4$ (0.52 of saturation). The clear supernatant was adjusted to pH 7.5 and NH_3 was added dropwise to pH 8.2-8.4. Crystals appear sometimes even without seeding. The crystals are dissolved in H_2O , filtered to remove suspended material and 2 volumes of saturated $(\text{NH}_4)_2\text{SO}_4$ at pH 8.2-8.4 is added. After 1hour the crystals appear. Recrystallise in the same way. [Cori et al. *J Biol Chem* 173 605 1948; Furfine and Velick *J Biol Chem* 240 844 1965, *The Enzymes* 7 243 1963; Lui and Huskey *Biochemistry* 31 6998 1992.] The K_m values are: NADH (3.3 μ M) and 1,3-diphosphoglycerate (8x10⁻⁷M) in pH 7.4 imidazole buffer at 26°, NAD (13 μ M), glyceraldehyde-3-P (90 μ M), P_i (2.9x10⁻⁴M), and arsenate (69 μ M) in 8.6 M NaHCO_3 buffer at 26°. [Orsi and Cleland *Biochemistry* 11 102 1972.]

Glycerol kinase (from *Candida mycoderma*, *E coli*, rat or pigeon liver glycerokinase) [9030-66-4] M_r 251,000, [EC 2.7.1.30]. Commercial enzyme has been dialysed against 2mM Hepes, 5mM dithiothreitol and 0.3mM EDTA, followed by several changes of 20mM Hepes and 5mM dithiothreitol prior to storage under N_2 at -20°. [Knight and Cleland *Biochemistry* 28 5728 1989.] The enzyme from pigeon liver was purified by acid-pptn (acetate buffer at pH 5.1), $(\text{NH}_4)_2\text{SO}_4$ fractionation, heat treatment (60°/ 1 h),

calcium phosphate gel filtration, a second $(\text{NH}_4)_2\text{SO}_4$ fractionation, dialysis, elution of inert proteins and crystn. This was done by repeatedly extracting the ppte from the last step with 0.05M sodium pyrophosphate (pH 7.5) containing 1mM EDTA and 0.2M $(\text{NH}_4)_2\text{SO}_4$ was added. Careful addition of solid $(\text{NH}_4)_2\text{SO}_4$ to this soln lead to crystn of the enzyme. Recrystn was repeated. The enzyme is activated by Mg^{2+} and Mn^{2+} ions and is most stable in solns in the pH 4.5-5.5 range. The stability is greatly increased in the presence of glycerol. It has K_m for glycerol of $60\mu\text{M}$ and for ATP $9\mu\text{M}$ in glycine buffer pH 9.8 and 25° . [Kennedy *Methods Enzymol* 5 476 1962.]

L-Glycerol-3-phosphate dehydrogenase (GDH, from rabbit muscle) [9075-65-4] M_r 78,000 [EC 1.1.1.8]. Recrystd by adding $(\text{NH}_4)_2\text{SO}_4$ till 0.45 saturation at pH 5.5 at 4° and the small amount of ppte is removed then satd $(\text{NH}_4)_2\text{SO}_4$ is added dropwise from time to time over several days in the cold room. The crystals are collected and recrystd until they have maximum activity. The enzyme is stable in half saturated $(\text{NH}_4)_2\text{SO}_4$ for several weeks at 4° . The equilibrium $[\text{dihydroxyacetone}][\text{NADH}][\text{H}^+]/[\text{G-3-P}][\text{NAD}]$ is $1.0 \times 10^{-12}\text{M}$ in Tris buffer at 25° . It uses NAD ten times more efficiently than NADP. The K_m for G-3-P is $1.1 \times 10^{-4}\text{M}$, for NAD it is $3.8 \times 10^{-4}\text{M}$ and for dihydroxyacetone it is $4.6 \times 10^{-4}\text{M}$ in phosphate buffer pH 7.0 and at 23.3° . Dihydroxyacetone phosphate and fructose-1,6-diphosphate are inhibitors. [Branowski *J Biol Chem* 180 515 1949, *The Enzymes* 7 85 1963; Young and Pace *Arch Biochem Biophys* 75 125 1958; Walsh and Sallach *Biochemistry* 4 1076 1965.]

L- α -Glycerol phosphocholine (Cadmium Chloride) $_x$ complex [64681-08-9] M 257.2 + (183.3) $_x$, $pK_{\text{Est}} \sim 5.5$. Glycerol phosphocholine is purified *via* the CdCl_2 complex which is purified by four recrystns from 99% EtOH by standing at 0° for 1h. The white ppte is collected, washed with EtOH, Et_2O and dried in a vacuum. The amorphous Cd complex can be converted to the crystalline form $[\text{C}_8\text{H}_{20}\text{O}_6\text{NP.CdCl}_2.3\text{H}_2\text{O}]$ by dissolving 34.4g in H_2O (410mL) and 99% EtOH (1650mL total) added slowly with stirring and allowing the clear soln to stand at 25° for 12hours, then at 5° for 12h. The crystallised complex is filtered off, washed with cold 80% EtOH and dried in air. Glycerol phosphocholine can be recovered from the complex by dissolving in H_2O (2% soln), passing through an ion-exchange column (4.9 x 100cm, of 1vol IRC-50 and 2vol IR-45). The effluent is concentrated to a thick syrup at 45° . It is dried further at $50^\circ/\text{P}_2\text{O}_5/48\text{h}$. The vitreous product ($\sim 8.25\text{g}$) is dissolved in 99% EtOH (50mL) and the clear soln is cooled at 5° , whereby crystals appear, and then at -15° for 16h. The crystals are filtered off, washed with 99% EtOH, and Et_2O then dried at 50° in a vacuum over P_2O_5 . It can be recrystd from 99.5% EtOH, long prisms) which are *hygroscopic* and must be handled in a H_2O -free atmosphere [Tattie and McArthur *Biochem Prep* 6 16 1958; Baer and Kates *J Am Chem Soc* 70 1394 1948 ; *Acta Cryst* 21 79, 87 1966].

Glycine anhydride (2,5-diketopiperazine) [106-57-0] M 114.1, m 309-310 $^\circ$, 311-312 $^\circ$ (dec), $\sim 315^\circ$ (dec), $pK_1 -4.45$, $pK_2 -2.16$ ($pK_2 -1.94$ in AcOH). Recrystd from H_2O (plates) and can be sublimed (slowly) at 260° or at $140-170^\circ/0.5\text{mm}$. The *dihydrochloride* has m 129-130 $^\circ$, is prepd by dissolving in conc HCl and on adding EtOH to crystallisation point; dried in a vac. The *bis-1-naphthylurethane* has m 232 $^\circ$ (dec), and the *diperchlorate* has m 117 $^\circ$ (*hygroscopic*). [MS: Johnstone *J Chem Soc Perkin Trans 1* 1297 1975; NMR: Blaha and Samek *Collect Czech Chem Commun* 32 3780 1967; Sauborn *J Phys Chem* 36 179 1932; Corey *J Am Chem Soc* 60 1599 1938.]

Glycoamine (N-guanylglycine) [352-97-6] M 117.1, m 280-284 $^\circ$ (dec), $>300^\circ$, $pK^{25} 2.86$ (NH_3^+). Recrystd from 15 parts of hot H_2O , or by dissolving in slightly more than the calculated amount of 2N HCl and ppting by adding an equivalent of 2N NaOH, filtering washing with cold H_2O and drying first *in vacuo* then at 60° *in vacuo*. The *hydrochloride* has m 200 $^\circ$ (dec) after recrystn from aqueous HCl as plates. The *picrate* forms needles from hot H_2O and has m 210 $^\circ$ (dec). [Brand and Brand *Org Synth Coll Vol III* 440 1955; Failey and Brand *J Biol Chem* 102 768 1933; King *J Chem Soc* 2375 1930.]

Glycodeoxycholic acid monohydrate (N-[3 α -12 α -dihydroxy-5 β -cholan-24-oyl]glycine) [360-65-6] M 467.6, m 186-177 $^\circ$ (dec), 187-188 $^\circ$, $[\alpha]_D^{23} +45.9^\circ$ (c 1, EtOH), $pK_{\text{Est}} \sim 4.4$. Recrystallises from H_2O or aqueous EtOH with 1 mol of H_2O and dried at 100° *in vacuo*. Solubility in EtOH is 5%. [UV: Lindstedt and Sjövall *Acta Chem Scand* 11 421 1957.] The *Na salt* is recrystd from EtOH/ Et_2O , m 245-250 $^\circ$, $[\alpha]_D^{23} +41.2^\circ$ (c 1, H_2O) [Wieland *Hoppe Seyler's Z Physiol Chem* 106 181 1919; Cortese *J Am Chem Soc* 59 2532 1937].

D(+)-Glycogen [9005-79-2] M 25,000-100,000, m 270-280°(dec), $[\alpha]_{546}^{20} +216^\circ$ (c 5, H₂O). A 5% aqueous soln (charcoal) was filtered and an equal volume of EtOH was added. After standing overnight at 3° the ppt was collected by centrifugation and washed with absolute EtOH, then EtOH/diethyl ether (1:1), and diethyl ether. [Sutherland and Wosilait *J Biol Chem* 218 459 1956.]

Glycogen synthase (from bovine heart) [9014-56-6] M_r 60,000, [EC 2.4.1.11]. Purified by pptn of the enzyme in the presence of added glycogen by polyethylene glycol, chromatography on DEAE-Sephacel and high speed centrifugation through a sucrose-containing buffer. [Dickey-Dunkirk and Kollilea *Anal Biochem* 146 199 1985.]

Gramicidin A (a pentadecapeptide from *Bacillus brevis*) [11029-61-1] m ~229-230°(dec). Purified by countercurrent distribution from *C₆H₆-CHCl₃, MeOH-H₂O (15:15:23:7) with 5000 tubes. Fractions were examined by UV (280nm) of small aliquots. Separation from Gramicidin C and other material occurred after 999 transfers. [Gross and Witkop *Biochemistry* 4 2495 1965; Bauer et al. *Biochemistry* 11 3266 1972.] Purified finally by recrystn from EtOH-H₂O and dried at 100°/10⁻² mm over KOH and forms platelets m 229-230°. Almost insoluble in H₂O (0.6%) but soluble in lower alcohols, dry Me₂CO, dioxane, acetic acid and pyridine. The commercial material is more difficult to crystallise than the synthetic compound. [Sarges and Witkop *J Am Chem Soc* 86 1861, 87 2011, 2020 1965.] It has characteristic $[\alpha]_D^{20} +27.3^\circ$ (c 1.3, MeOH) and UV λ_{max} 282nm (ϵ 22,100). The *N*-carbamoyldeformyl gramicidine A pptes from EtOAc-pet ether (b 40-60°).

Gramicidin C (gramicidin S, a pentadecapeptide from *Bacillus brevis*) [9062-61-7]. Same as Gramicidin A since they are isolated together and separated. [Sarges and Witkop *Biochemistry* 4 2491 1965; Hunter and Schwartz "Gramicidins" in *Antibiotics I* (Gotlieb and Shaw Eds) Springer-Verlag, NY, p.642 1967; as well as references above for Gramicidin A.]

Gramicidin S 2HCl (from *Bacillus brevis* Nagano) [15207-30-4] M 1214.4, m 277-278°(dec), $[\alpha]_D^{24} -289^\circ$ (c 0.4, 70% aq EtOH). Cryst in prisms from EtOH + aq HCl.

Gramicidin S [113-73-5] M 1141.4, m 268-270°, $[\alpha]_D^{25} -290^\circ$ (c 0.5, EtOH + 30mM aq HCl [7:3]). Crystd from EtOH. *Di-HCl* [15207-30-4] cryst from EtOH (+ few drops of HCl) has m 277-278°.

***N*-Guanyltiramine hydrochloride** [60-20-8] M 215.7, m 218°, pK_1 10.2 (phenolic OH), pK_2 12.4 (guanidino N). Purified on a phosphocellulose column and eluted with a gradient of aqueous NH₃ (0-10%). The second major peak has the characteristic tryptamine spectrum and is collected, lyophilised to give white crystals of the *dihydrate* which dehydrates at 100°. It has UV λ_{max} at 274.5nm (ϵ 1310) in 0.1N NaOH and 274.5nm (ϵ 1330) at pH 7.0. Excitation λ_{max} is at 280nm and emission λ_{max} is at 330nm. [Mekalanos et al. *J Biol Chem* 254 5849 1979.]

Haemoglobin A (from normal human blood) [9008-02-0] M_r ~64,500, amorphous. Purified from blood using CM-32 cellulose column chromatography. [Matsukawa et al. *J Am Chem Soc* 107 1108 1985.] For the purification of the α and β chains see Hill et al. *Biochem Prep* 10 55 1963.

Harmaline (7-methoxy-1-methyl-4,9-dihydro-3*H*- β -carboline, 4,9-dihydro-7-methoxy-1-methyl-3*H*-pyrido[3,4-*b*]indole) [304-21-2] M 214.3, m 229-230°, 229-231°, 235-237° (after distn at 120-140°/10⁻³), pK_1 4.2. Recrystd from MeOH and sublimed at high vacuum. It has UV in MeOH has λ_{max} 218, 260 and 376nm (log ϵ 4.27, 3.90 and 4.02 respectively); IR (Nujol) ν 1620, 1600, 1570 and 1535cm⁻¹ and in CHCl₃ ν 1470 and 1629cm⁻¹. [Spenser *Can J Chem* 37 1851 1959; Marion et al. *J Am Chem Soc* 73 305 1951; UV Prukner and Witkop *Justus Liebigs Ann Chem* 554 127 1942.] The *hydrochloride dihydrate* has m 234-236°(dec), the *picrate* has m 228-229° (sinters at 215°) from aqueous EtOH, and the *N*-acetate forms needles m 204-205°.

Hematin (ferrihaeme hydroxide) [15489-90-4] M 633.5, m 200°(dec), pK_{Est} ~ 4. Crystd from pyridine. Dried at 40° *in vacuo*.

Hematoporphyrin (3,3'-[7,12-bis-(1-hydroxyethyl)-3,8,13,17-tetramethyl-porphyrin-2,18-diyl]-dipropionic acid) [14459-29-1] M 598.7, pK_{Est} ~4.8. Purified by dissolving in EtOH and adding H₂O or Et₂O to give deep red crystals. Also recrystd from MeOH. UV has λ_{max} at 615.5, 565, 534.4 and 499.5nm in 0.1 N NaOH, and 597, 619, 634,653, 683 and 701nm in 2 N HCl. [Falk *Porphyryns and Metalloporphyryns* Elsevier, NY, p 175 1964.] It is used in the affinity chromatographic purification of Heme proteins [Olsen *Methods Enzymol* 123 324 1986.] The *O*-methyl-dimethyl ester has m 203-206° (from CHCl₃-MeOH) and the *O,O'*-dimethyl-dimethyl ester has m 145° (from CHCl₃-MeOH). [Paul *Acta Chem Scand* 5 389 1951.]

Hematoporphyrin dimethyl ester [33070-12-1] M 626.7, m 212°. Crystd from CHCl₃/MeOH.

Hematoxylin (\pm -11bc-7,11b-dihydroindeno[2,1-c]-chromen-3,4-6ar-9,10-pentaol) [517-28-2] M 302.3, m 200°(dec), 210-212°(dec). Recrystd from H₂O (as trihydrate) in white-yellow crystals which become red on exposure to light and then melt at 100-120°. It has been recrystd from Me₂CO-^{*}C₆H₆. Crystd also from dil aqueous NaHSO₃ until colourless. Soluble in alkali, borax and glycerol. Store in the dark below 0°. [Morsingh and Robinson *Tetrahedron* 26 182 1970; Dann and Hofmann *Chem Ber* 98 1498 1955.]

Hemin (ferriprotoporphyrin IX chloride) [16009-13-5] M 652.0, m sinters at 240°, pK_{Est} ~4.8. It is purified by recrystn from AcOH. Also heme (5g) is shaken in pyridine (25mL) till it dissolves, then CHCl₃ (40mL) is added, the container is stoppered and shaken for 5min (releasing the stopper occasionally). The soln is filtered under slight suction, and the flask and filter washed with a little CHCl₃ (15mL). During this period, AcOH (300mL) is heated to boiling and saturated aqueous NaCl (5mL) and conc HCl (4mL) are added. The CHCl₃ filtrate is poured in a steady stream, with stirring, into the hot AcOH mixture and set aside for 12hours. The crystals are filtered off, washed with 50% aqueous AcOH (50mL), H₂O (100mL), EtOH (25mL), Et₂O and dried in air. [Fischer *Org Synth Coll Vol III* 442 1955.]

Heparin (from pig intestinal mucosa) [9005-49-6] M_r ~3,000, amorphous, $[\alpha]_D^{20}$ ~+55° (H₂O). Most likely contaminants are mucopolysaccharides including heparin sulfate and dermatan sulfate. Purified by pptn with cetylpyridinium chloride from saturated solutions of high ionic strength. [Cifonelli and Roden *Biochem Prep* 12 12 1968.]

Heparin (sodium salt) [9041-08-1] M_r ~ 3000 (low Mol Wt, Bovine), amorphous, $[\alpha]_D^{20}$ +47° (c 1.5, H₂O). Dissolved in 0.1M NaCl (1g/100mL) and pted by addition of EtOH (150mL).

Histones (from S4A mouse lymphoma). Purification used a macroprocess column, heptafluorobutyric acid as solubilising and ion-pairing agent and an acetonitrile gradient. [McCroskey et al. *Anal Biochem* 163 427 1987.]

Hyaluronidase [9001-54-1, 37326-33-3] M_r 43,000 (bovine testes), 89,000 (bacterial), [EC 3.2.1.35]. Purified by chromatography on DEAE-cellulose prior to use. [Distler and Jourdain *J Biol Chem* 248 6772 1973.]

Hydrocortisone (11 β ,17 α ,21-trihydroxy-pregn-4-ene-3,20-dione) [50-23-7] M 362.5, m 212-213°, 214-217°, 218-221°, 220-222°, $[\alpha]_D^{22}$ +167° (c 1, EtOH). Recrystd from EtOH or isoPrOH. It is bitter tasting and has UV λ_{max} at 242 nm (log ϵ 4.20). Its solubility at 25° is: H₂O (0.28%), EtOH (1.5%), MeOH (0.62%), Me₂CO (0.93%), CHCl₃ (0.16%), propylene glycol (1.3%) and Et₂O (0.35%). It gives an intense green colour with conc H₂SO₄. [Wendler et al. *J Am Chem Soc* 72 5793 1950.]

Hydrocortisone acetate (21-acetoxy-11 β ,17 α -trihydroxy-pregn-4-ene-3,20-dione) [50-03-3] M 404.5, m 218-221.5°, 221-223°, 222-225°, $[\alpha]_D^{25}$ +166° (c 0.4, dioxane), +150.7° (c 0.5, Me₂CO). Recrystd from Me₂CO-Et₂O or aqueous Me₂CO as somewhat *hygroscopic* monoclinic crystals. UV has λ_{max} 242 nm ($A_{1cm}^{1\%}$ 390) in MeOH. Its solubility at 25° is: H₂O (0.001%), EtOH (0.45%), MeOH (0.04%), Me₂CO (1.1%), CHCl₃ (0.5%), Et₂O (0.15%) and is very soluble in Me₂NCHO. [Wendler et al. *J Am Chem Soc* 74 3630 1952; Antonucci et al. *J Org Chem* 18 7081 1953.]

(+)-Hydroquinidine anhydrous (9S-6'-methoxy-10,11-dihydrocinchonan-9-ol) [1435-55-8] **M 326.4, m 168-169°, 169°, 169-170°, 171-172°, $[\alpha]_D^{20} +231^\circ$ (c 2, EtOH), +299° (c 0.82, 0.1N H₂SO₄), $pK_{Est} \sim 8.8$** . Forms needles from EtOH and plates from Et₂O. Slightly soluble in Et₂O and H₂O but readily soluble in hot EtOH. [Heidelberger and Jacobs *J Am Chem Soc* **41** 826 1919; King *J Chem Soc* 523 1946.] The *hydrochloride* has **m 273-274°, $[\alpha]_D^{26} +184^\circ$ (c 1.3, MeOH)** and is very soluble in MeOH and CHCl₃, but less soluble in H₂O, EtOH and less soluble in dry Me₂CO. [Kyker and Lewis *J Biol Chem* **157** 707 1945; Emde *Helv Chim Acta* **15** 557 1932.]

Hydroquinine [522-66-7] **M 326.4, m 168-171°, 171.5°, $[\alpha]_D^{16} +143^\circ$ (c 1.087, EtOH), $pK^{15} 8.87$** . Recrystd from EtOH. [Rabe and Schultz *Chem Ber* **66** 120 1933.]

19-Hydroxy-4-androsten-3,17-dione [510-64-5] **M 302.4, m 167-169°, 168-170°, 169-170°, 172-173°, $[\alpha]_D^{20} +190^\circ$ (c 1, CHCl₃)**. Recrystd from Me₂CO-hexane or Et₂O-hexane. It has UV λ_{max} at 242nm in EtOH or MeOH. The *19-acetoxy* derivative has $[\alpha]_D^{26} +185^\circ$ (CHCl₃) and λ_{max} 237.5nm in EtOH. [Ehrenstein and Dünneberger *J Org Chem* **21** 774 1956.]

3-Hydroxy butyrate dehydrogenase (from *Rhodopseudomonas spheroides*) [9028-38-0] **M_r ~85,000, [EC 1.1.1.30], amorphous**. Purified by two sequential chromatography steps on two triazine dye-Sepharose matrices. [Scavan et al. *Biochem J* **203** 699 1982.]

25-Hydroxycholesterol (cholest-5-en-3 β ,25-diol) [2140-46-7] **M 402.7, m 177-179°, 178-180°, 181.5-182.5°, $[\alpha]_D^{25} -39^\circ$ (c 1.05, CHCl₃)**. Forms colourless needles from MeOH. [Schwartz *Tetrahedron Lett* **22** 4655 1981.] The *3 β -acetoxy* derivative has **m 142-142.8° (from Me₂CO), $[\alpha]_D^{25} -40.4^\circ$ (c 2, CHCl₃)**. The *3 β ,25-diacetyl* derivative has **m 119-120.5° (from MeOH), $[\alpha]_D^{25} -35.5^\circ$ (CHCl₃)**. [Dauben and Bradlow *J Am Chem Soc* **72** 4248 1950; Ryer et al. *J Am Chem Soc* **72** 4247 1950.]

18-Hydroxy-11-deoxycorticosterone (18,21-dihydroxypregn-4-en-3,20-dione tautomeric with 18,20-epoxy-20,21-dihydroxypregn-4-en-3-one) [379-68-0] **M 346.5, m 168-170°, 171-173°, 191-195°, 200-205°, $[\alpha]_D^{20} +151^\circ$ (c 1, CHCl₃)**. Recrystn from Et₂O-Me₂CO gave crystals **m 200-205°**, when recrystd from Me₂CO it had **m 191-195°**. It has UV λ_{max} at 240nm. The *21-O-acetoxy-18-hydroxy* derivative has **m 158-159° (from Et₂O-^{*}C₆H₆)** and the *21-O-acetoxy-18,20-epoxy* derivative has **m 149-154° (from Et₂O)**. [Kahnt et al. *Helv Chim Acta* **38** 1237 1955; Pappo *J Am Chem Soc* **81** 1010 1959.]

R-(-)-2-Hydroxy-3,3-dimethyl- γ -butyrolactone (3-hydroxy-4,4-dimethyl-4,5-dihydrofuran-2-one, D-pantolactone) [599-04-2] **M 130.1, m 89-91°, 90.5-91.5°, 91°, 92-93°, b 120-122°/15mm, $[\alpha]_D^{20} -28^\circ$ (c 5, MeOH), $[\alpha]_D^{20} -51^\circ$ (c 3, H₂O)**. Recrystallise from Et₂O-pet ether, diisopropyl ether or ^{*}C₆H₆-pet ether and sublime at 25°/0.0001mm. It hydrolyses readily to the hydroxy-acid and racemises when heated above 145°. The *Brucine salt* has **m 211-212° (from EtOH)**. [Kuhn and Wieland *Chem Ber* **73** 1134 1940; and Stiller et al. *J Am Chem Soc* **62** 1779 1940; Bental and Tishler *J Am Chem Soc* **68** 1463 1946.]

(±)-Ibotenic acid monohydrate (α -[3-hydroxy-5-isoxazolyl]-glycine, α -amino-3-hydroxy-5-isoxazoleacetic acid) [2552-55-8] **M 176.1, m 144-146° (monohydrate), 151-152° (anhydrous), 148-151°, pK_1 2, pK_2 5.1, pK_3 8.2**. It has been converted to the ammonium salt (**m 121-123° dec**) dissolved in H₂O and passed through an Amberlite IR 120 resin (H⁺ form) and eluted with H₂O. The acidic fractions were collected, evaporated to dryness and the residue recrystd from H₂O as the monohydrate (**m 144-146°**). The anhydrous acid is obtained by making a slurry with MeOH, decanting and evaporating to dryness and repeating the process twice more to give the anhydrous acid (**m 151-152°**). Recrystn from H₂O gives the monohydrate. [Nakamura *Chem Pharm Bull Jpn* **19** 46 1971.] The *ethyl ester* forms needles when crystd from a small volume of Et₂O and has **m 78-79° and IR (CHCl₃) with ν 3500-2300 (OH), 1742 (ester CO), 1628, 1528cm⁻¹, and UV with λ_{max} (EtOH) at 206nm (ϵ 7080)**. The *hydrazide* has **m 174-175° (from MeOH) with IR (KBr) 1656 (C=O)cm⁻¹**.

2-Iminothiolane hydrochloride (2-iminotetrahydrothiophene) [4781-83-3] **M 137.6, m 187-192°**, **190-195°**, **193-194°**, **202-203°**, **pK <2 (free base)**. Recryst from MeOH-Et₂O (**m** 187-192°) but after sublimation at ~180°/0.2mm the melting point rose to 202-203°. It has NMR with δ 2.27 (2H, t), 3.25 (2H, t) and 3.52 (2H, t) in (CD₃)₂SO. [King et al. *Biochemistry* **17** 1499 1978.] The *free base* is purified by vacuum distn (**b** 71-72°/6mm) with IR (film) with ν 1700 (C=N)cm⁻¹ and NMR (CDCl₃) with δ at 3.58 (2H, t) and 2.10-2.8 (4H, m). The *free base* is stable on storage but slowly hydrolyses in aqueous solns with half lives at 25° of 390h at pH 9.1, 210h at pH 10 and 18 h at pH 11. [Alagon and King *Biochemistry* **19** 4343 1980.]

trans-Indol-3-ylacrylic acid [1204-06-4] **M 187.2, m 190-195°(dec)**, **195°(dec)**, **196°(dec)**, **195-196°(dec)**, **pK_{Est} ~ 4.2**. Recrystd from AcOH, H₂O or EtOAc-cyclohexane. UV in MeOH has λ_{\max} at 225, 274 and 325nm. [Shaw et al. *J Org Chem* **23** 1171 1958; constitution: Rappe *Acta Chem Scand* **18** 818 1964; Moffatt *J Chem Soc* 1442 1957; Kimming et al. *Hoppe Seyler's Z Physiol Chem* **371** 234 1958.]

3-Indolylbutyric acid [133-32-4] **M 203.2, m 120-123°**, **123-125°**, **124°**, **pK 4.84**. Recrystd from H₂O. It is soluble in EtOH, Et₂O and Me₂CO but insoluble in CHCl₃. [Bowman and Islip *Chem Ind London* 154 1971; Jackson and Manske *J Am Chem Soc* **52** 5029 1930; Albaum and Kaiser *Am J Bot* **24** 420 1937.] UV has λ_{\max} 278 and 320nm in isoPrOH [Elvidge *Quart J Pharm Pharmacol* **13** 219 1940]. The *methyl ester* has **m** 73-74° (from *C₆H₆-pet ether) and **b** 230°/6mm [Bullock and Hand *J Am Chem Soc* **78** 5854 1951]. Also recrystd from EtOH/water [James and Ware *J Phys Chem* **89** 5450 1985].

3-Indolylpyruvic acid [392-12-1] **M 203.2, m~210°(dec)**, **208-210°(dec)**, **219°(dec)**, **pK_{Est} ~ 2.4**. Recrystd from Me₂CO-*C₆H₆, EtOAc-CHCl₃, Me₂CO-AcOH (crystals with 1 molecule of AcOH) and dioxane-*C₆H₆ (with 0.5 molecule of dioxane) [Shaw et al. *J Org Chem* **23** 1171 1958; Kaper and Veldstra *Biochim Biophys Acta* **30** 401 1958]. The *ethyl ester* has **m** 133° (from Et₂O) and its *2,4-dinitrophenylhydrazone* has **m** 255° (from Me₂CO). [Baker *J Chem Soc* 461 1946.]

myo-Inositol (cyclohexane[1r,2c,3c,4t,5c,6t]-hexol) [87-89-8] **M 180.2, m 218° (dihydrate)**, **225-227°**, **226-230°**. Recrystd from aq 50% ethanol or H₂O forming a dihydrate, or anhydrous crystals from AcOH. The dihydrate is efflorescent and becomes anhydrous when heated at 100°. The anhydrous crystals are not hygroscopic. Solubility in H₂O at 25° is 14%, at 60° it is 28%, slightly soluble in EtOH but insoluble in Et₂O. [Ballou and Anderson *J Am Chem Soc* **75** 748 1953; Anderson and Wallis *J Am Chem Soc* **70** 2931 1948.]

Interferons [α IFN, β IFN and γ IFN]. Interferons are a family of glycosylated proteins and are cytokines which are produced a few hours after cells have been infected with a virus. Interferons protect cells from viral infections and have antiviral activities at very low concentrations (~3 x 10⁻⁴ M, less than 50 molecules are apparently sufficient to protect a single cell). Double stranded RNA are very efficient inducers of IFNs. There are three main types of IFNs. The α IFNs are synthesised in lymphocytes and the β IFNs are formed in infected fibroblasts. The α and β families are fairly similar consisting of *ca* 166 to 169 amino acids. Although γ IFNs are also small glycosylated proteins (*ca* 146 amino acids), they are different because they are not synthesised after viral infections but are produced by lymphocytes when stimulated by **mitogens** (agents that induced cell division).

Several of these IFNs of mouse and human lymphocytes and fibroblasts are available commercially and have been best prepared in quantity by recombinant DNA procedures because they are produced in very small amounts by the cells. The commercial materials do not generally require further purification for their intended purposes. [Pestkas, Interferons and Interferon standards and general abbreviations, *Methods Enzymol*, Wiley & Sons, **119** 1986, ISBN 012182019X; Lengyel, Biochemistry of interferons and their actions, *Ann Rev Biochem* **51** 251-282 1982; De Maeyer and De Maeyer-Guignard, Interferons in *The Cytokine Handbook*, 3rd Edn, Thomson et al. Eds, pp. 491-516 1998 Academic Press, San Diego, ISBN 0126896623.]

Interleukin (from human source). Purified using lyophilisation and desalting on a Bio-Rad P-6DC desalting gel, then two steps of HPLC, first with hydroxylapatite, followed by a TSK-125 size exclusion column. [Kock and Luger *J Chromatogr* **296** 293 1984.]

Interleukin-2 (recombinant human) [94218-72-1] $M_r \sim 15,000$, amorphous. Purified by reverse phase HPLC. [Weir and Sparks *Biochem J* **245** 85 1987; Robb et al. *Proc Natl Acad Sci USA* **81** 6486 1984.]

Interleukins (IL-1, IL-2 —IL18). Interleukins are cytokines which cause a variety of effects including stimulation of cell growth and proliferation of specific cells, e.g. stem cells, mast cells, activated T cells, colony stimulating factors etc, as well as stimulating other ILs, prostaglandins release etc. They are small glycosylated proteins (ca 15 kD, 130-180 amino acids produced from longer precursors) and are sometimes referred to by other abbreviations, e.g. IL-2 as TCGF (T cell growth factor), IL-3 as multi-CSF (multilineage colony stimulating factor, also as BPA, HCSF, MCSF and PSF). They are produced in very small amounts and are commercially made by recombinant DNA techniques in bacteria or Sf21 insect cells. Interleukins for human (h-IL), mouse (m-IL) and rat (r-IL) are available and up to IL-18 are available commercially in such purity that they can be used directly without further refinement, particularly those that have been obtained by recombinant DNA procedures which are specific. As well as the interleukins, a variety of antibodies for specific IL reactions are available for research or IL identification. [Symons et al. *Lymphokines and Interferons, A Practical Approach*, Clemens et al. Eds, p. 272 1987; IRL Press, Oxford, ISBN 1852210354, 1852210362; Thomson et al. Eds, *The Cytokine Handbook*, 3rd Edn, 1998; Academic Press, San Diego, ISBN 0126896623.]

Iodonitrotetrazolium chloride (2[4-iodophenyl]-3-[4-nitrophenyl]-5-phenyl-2H-tetrazolium chloride) [146-68-9] M 505.7, m 229°(dec), ~245°(dec). Recrystd. from H₂O, aqueous EtOH or EtOH-Et₂O. Alternatively dissolve in the minimum volume of EtOH and add Et₂O; or dissolve in hot H₂O (charcoal), filter and ppt by adding conc HCl. Filter solid off and dry at 100°. Solubility in H₂O at 25° is 0.5%, and in hot MeOH-H₂O (1:1) it is 5%. [Fox and Atkinson *J Am Chem Soc* **72** 3629 1950.]

Iodonitrotetrazolium violet-Formazan [7781-49-9] M 471.3, m 185-186°. Dissolve in boiling dioxane (20g in 300mL), add H₂O (100mL) slowly, cool, filter and dry *in vacuo* at 100°. Its solubility in CHCl₃ is ~1%. [UV: Fox and Atkinson *J Am Chem Soc* **72** 3629 1950.]

5-Iodouridine (5-iodo-1-[β-D-ribofuranosyl]-pyrimidine-2,4(1H)-dione) [1024-99-3] M 370.1, m 205-208°(dec), 210-215°(dec), $[\alpha]_D^{20}$ -23.5° (c 1, H₂O), pK^{20} 8.5. Recrystd from H₂O and dried *in vacuo* at 100°. UV has λ_{max} 289nm (0.01N HCl) and 278nm (0.01N NaOH). [Prusoff et al. *Cancer Res* **13** 221 1953.]

3-Isobutyl-1-methylxanthine (3-isobutyl-1-methylpurine-2,6-dione) [28822-58-4] M 222.3, m 199-210°, 202-203°, $pK_{Est} \sim 6.7$ (acidic NH). Recrystd from aqueous EtOH.

Isopentenyl pyrophosphate [358-71-4] M 366.2, $pK_{Est(1)} \sim <2$, $pK_{Est(2)} \sim <2$, $pK_{Est(3)} \sim 3.95$, $pK_{Est(4)} \sim 6.26$. Purified by chromatography on Whatman No 1 paper using *tert*-butyl alcohol/formic acid/water (20:5:8, R_F 0.60) or 1-propanol/ammonia/water (6:3:1. R_F 0.48). Also purified by chromatography on a DEAE-cellulose column or a Dowex-1 (formate form) ion-exchanger using formic acid and ammonium formate as eluents. A further purification step is to convert it to the monocyclohexylammonium salt by passage through a column of Dowex-50 (cyclohexylammonium form) ion-exchange resin. Can also be converted into its lithium salt.

DL-Isoserine (±-3-amino-2-hydroxypropionic acid) [632-12-2] M 105.1, m 250-252°(dec), 235°(dec), 237°(dec), 245°(dec), pK_1^{25} 2.78 (acidic), pK_2^{25} 9.27 (basic). Recrystd from H₂O or 50% aqueous EtOH. It has an isoelectric pH of 6.02. [Rinderknocht and Niemann *J Am Chem Soc* **75** 6322 1953; Gundermann and Holtmann *Chem Ber* **91** 160 1958; Emerson et al. *J Biol Chem* **92** 451 1931.] The *hydrobromide* has m 128-130° (from aqueous HBr) [Schöberl and Braun *Justus Liebigs Ann Chem* **542** 288 1939].

Isoxanthopterin (2-amino-4,7-dihydroxypteridine) [529-69-1] M 179.4, $m > 300^\circ$, pK_1^{20} -0.5 (basic), pK_2^{20} 7.34 (acidic), pK_3^{20} 10.06 (acidic). Purified by repeated pptn from alkaline solutions by acid (preferably AcOH), filter, wash well with H₂O then EtOH and dried at 100°. Purity is checked by paper chromatography [R_F 0.15 (*n*-BuOH, AcOH, H₂O, 4:1:1); 0.33 (3% aq NH₄OH)]. [Goto et al. *Arch Biochem*

Biophys 111 8 1965.] For biochemistry see Blakley *Biochemistry of Folic Acid and Related Pteridines* North Holland Publ Co, Amsterdam 1969.]

Kanamycin B (Bekanamycin, 4-*O*-[2,6-diamino-2,6-dideoxy- α -D-glucopyranosyl]-6-*O*-[3-amino-3-deoxy- α -D-glucopyranosyl]-2-deoxystreptamine) [4696-76-8, 29701-07-3 (sulfate salt)] **M** 483.5, **m** 170-179°(dec), 178-182°(dec), $[\alpha]_D^{18} +130^\circ$ (c 0.5, H₂O), **pK** 7.2. A small quantity (24mg) can be purified on a small Dowex 1 x 2 column (6 x 50mm), the correct fraction is evapd to dryness and the residue crystd from EtOH containing a small amount of H₂O. [Umezawa et al. *Bull Chem Soc Jpn* 42 537 1969.] It has been crystd from H₂O by dissolving ~1g in H₂O (3mL), adding Me₂NCHO (3mL) setting aside at 4° overnight, The needles are collected and dried to constant weight at 130°. It has also been recrystd from aq EtOH. It is slightly sol in CHCl₃ and isoPrOH. [IR: Wakazawa et al. *J Antibiot* 14A 180, 187 1961; Ito et al. *J Antibiot* 17 A 189 1964.]

Lactate dehydrogenase (from dogfish, Beef muscle) [9001-60-9] **M_r** 140,000 [EC 1.1.1.27]. 40-Fold purification by affinity chromatography using Sepharose 4B coupled to 8-(6-aminoethyl)amino-5'-AMP or -NAD⁺. [Lees et al. *Arch Biochem Biophys* 163 561 1974; Pesce et al. *J Biol Chem* 239 1753 1964.]

Lactoferrin (from human whey). Purified by direct adsorption on cellulose phosphate by batch extraction, then eluted by a stepped salt and pH gradient. [Foley and Bates *Anal Biochem* 162 296 1987.]

Lecithin (1,2-diacylphosphatidylcholine mixture) [8002-43-5] **M** ~600-800, **amorphous**. From hen egg white. Purified by solvent extraction and chromatography on alumina. Suspended in distilled water and kept frozen until used [Lee and Hunt *J Am Chem Soc* 106 7411 1984, Singleton et al. *J Am Oil Chem Soc* 42 53 1965]. For purification of commercial egg lecithin see Pangborn [*J Biol Chem* 188 471 1951].

Lectins (proteins and/or glycoproteins of non-immune origin that agglutinate cells, from seeds of *Robinia pseudoacacia*), **M** ~100,000. Purified by pptn with ammonium sulfate and dialysis; then chromatographed on DE-52 DEAE-cellulose anion-exchanger, hydroxylapatite and Sephacryl S-200. [Wantyghem et al. *Biochem J* 237 483 1986.]

Leucopterin (2-amino-5,8-dihydropteridine-4,6,7(1*H*)-trione) [492-11-5] **M** 195.1, **m** >300° (dec), **pK**₁²⁰ -1.66, **pK**₂²⁰ 7.56, **pK**₃²⁰ 9.78, **pK**₄²⁰ 13.6. Purified by dissolving in aqueous NaOH, stirring with charcoal, filtering and precipitating by adding aqueous HCl, then drying at 100° in a vacuum. It separates with 0.5 moles of H₂O. Its solubility in H₂O is 1g/750 litres [Albert et al. *J Chem Soc* 4219 1952; Albert and Wood *J Appl Chem (London)* 2 591 1952; Pfeleiderer *Chem Ber* 90 2631 1957].

DL- α -Lipoamide (\pm -6,8-thioctic acid amide, 5-[1,2]-dithiolan-3-ylvaleric acid amide) [3206-73-3] **M** 205.3, **m** 124-126°, 126-129°, 130-131°. Recrystd from EtOH and has UV with λ_{\max} 331nm in MeOH. [Reed et al. *J Biol Chem* 232 143 1958; IR: Wagner et al. *J Am Chem Soc* 78 5079 1956.]

DL- α -Lipoic acid (\pm -6,8-thioctic acid, 5-[1,2]-dithiolan-3-ylvaleric acid) [1077-28-7] **M** 206.3, **m** 59-61°, 60.5-61.5° and 62-63°, **b** 90°/10⁻⁴mm, 150°/0.1mm, **pK**²⁵ 4.7. It forms yellow needles from cyclohexane or hexane and has been distd at high vacuum, and sublimes at ~90° and very high vacuum. Insoluble in H₂O but dissolves in alkaline soln. [Lewis and Raphael *J Chem Soc* 4263 1962; Soper et al. *J Am Chem Soc* 76 4109; Reed and Niu *J Am Chem Soc* 77 416 1955; Tsuji et al. *J Org Chem* 43 3606 1978; Calvin *Fed Proc USA* 13 703 1954.] The *S*-benzylthiouronium salt has **m** 153-154° (evacuated capillary; from MeOH), 132-134°, 135-137° (from EtOH). The *d*- and *l*- forms have **m** 45-47.5° and $[\alpha]_D^{23} \pm 113^\circ$ (c 1.88, *C₆H₆) and have UV in MeOH with λ_{\max} at 330nm (ϵ 140).

Lipoprotein lipase (from bovine skimmed milk) [9004-02-8] [EC 3.1.1.34]. Purified by affinity chromatography on heparin-Sepharose [Shirai et al. *Biochim Biophys Acta* 665 504 1981].

Lipoproteins (from human plasma). Individual human plasma lipid peaks were removed from plasma by ultracentrifugation, then separated and purified by agarose-column chromatography. Fractions were characterised immunologically, chemically, electrophoretically and by electron microscopy. [Rudel et al. *Biochem J* 13 89 1974.]

Lipoteichoic acids (from gram-positive bacteria) [56411-57-5]. Extracted by hot phenol/water from disrupted cells. Nucleic acids that were also extracted were removed by treatment with nucleases. Nucleic resistant acids, proteins, polysaccharides and teichoic acids were separated from lipoteichoic acids by anion-exchange chromatography on DEAE-Sephacel or by hydrophobic interaction on octyl-Sepharose [Fischer et al. *Eur J Biochem* 133 523 1983].

D-Luciferin (firefly luciferin, S-2[6-hydroxybenzothiazol-2-yl]-4,5-dihydrothiazol-4-carboxylic acid), [2591-17-5] M 280.3, m 189.5-190°(dec), 196°(dec), 201-204°, 205-210°(dec, browning at 170°), $[\alpha]_D^{22}$ -36° (c 1.2, DMF), $pK_{Est(1)} \sim 1.2$ (benzothiazole-N), $pK_{Est(2)} \sim 1.6$ (thiazolidine-N), $pK_{Est(3)} \sim 6.0$ (CO₂H), $pK_{Est(4)} 8.5$ (6OH). Recrystallises as pale yellow needles from H₂O, or MeOH (83mg from 7mL). It has UV λ_{max} at 263 and 327nm (log ϵ 3.88 and 4.27) in 95% EtOH. The Na salt has a solubility of 4mg in 1 mL of 0.05M glycine. [White et al. *J Am Chem Soc* 83 2402 11961, 85 337 1963; UV and IR: Bitler and McElroy *Arch Biochem* 72 358 1957; Review: Cormier et al. *Fortschr Chem Org Naturst* 30 1 1973.]

Lumiflavin (7,8,10-trimethylbenzo[g]pteridine-2,4(3H,10H)-dione) [1088-56-8] M 256.3, m 330°(dec), 340°(dec), pK 10.2. Forms orange crystals upon recrystn from 12% aqueous AcOH, or from formic acid. It sublimes at high vacuum. It is freely soluble in CHCl₃, but not very soluble in H₂O and most organic solvents. In H₂O and CHCl₃ soln it has a green fluorescence. UV has λ_{max} at 269, 355 and 445nm (ϵ 38,800, 11,700 and 11,800 respectively) in 0.1N NaOH and 264, 373 and 440nm (ϵ 34,700, 11,400 and 10,400 respectively) in 0.1N HCl while UV in CHCl₃ has λ_{max} at 270, 312, 341, 360, 420, 445 and 470nm. [Hemmerich et al. *Helv Chim Acta* 39 1242 1956; Holiday and Stern *Chem Ber* 67 1352 1834; Yoneda et al. *Chem Pharm Bull Jpn* 20 1832 1972; Birch and Moye *J Chem Soc* 2622 1958; Fluorescence: Kuhn and Moruzzi *Chem Ber* 67 888 1934.]

Magnesium protoporphyrin dimethyl ester [14724-63-1] M 580.7. Crude product dissolved in as little hot dry *C₆H₆ as possible and left overnight at room temperature to crystallise. [Fuhrhop and Graniek *Biochem Prep* 13 55 1971.]

α -Melanotropin [581-05-5] (13 amino acids peptide), $[\alpha]_D^{25}$ -58.5° (c 0.4, 10% aq AcOH). Extract separated by ion-exchange on carboxymethyl cellulose, desalted, evapd and lyophilised, then chromatographed on Sephadex G-25. [Lande et al. *Biochem Prep* 13 45 1971.]

β -Melanotropin. [9034-42-8] (18-22 amino acids peptide), amorphous. Extract separated by ion-exchange on carboxymethyl cellulose, desalted, evapd and lyophilised, then chromatographed on Sephadex G-25. [Lande et al. *Biochem Prep* 13 45 1971.]

6-Mercaptopurine monohydrate [6112-76-1] M 170.2, m 314-315°(dec), ~315°(dec), 313-315°(dec), $pK_1^{20} 7.77$, $pK_2^{20} 10.84$. Recrystallises from H₂O as yellow crystals of the monohydrate which become anhydrous on drying at 140°. It has UV λ_{max} at 230 and 312nm (ϵ 14,000 and 19,600) in 0.1N NaOH; 222 and 327nm (ϵ 9,2400 and 21,300), and 216 and 329nm (ϵ 8,740 and 19,300) in MeOH. [Albert and Brown *J Chem Soc* 2060 1954; IR: Brown and Mason *J Chem Soc* 682 1957; UV: Fox et al. *J Am Chem Soc* 80 1669 1958; UV: Mason *J Chem Soc* 2071 1954.]

6-Mercaptopurine-9- β -D-ribofuranoside [574-25-4] M 284.3, m 208-210°(dec), 210-211°(dec), 220-223°(dec), 222-224°(dec), $[\alpha]_D^{25}$ -73° (c 1, 0.1N NaOH), pK 7.56. Recrystd from H₂O or EtOH. It has UV λ_{\max} in H₂O at 322nm (pH 1), 320 nm (pH 6.7) and 310nm (pH 13). [IR: Johnson et al. *J Am Chem Soc* 80 699 1958; UV: Fox et al. *J Am Chem Soc* 80 1669 1958.]

Metallothionein (from rabbit liver) [9038-94-2]. Purified by precipitation to give Zn- and Cd-containing protein fractions and running on a Sephadex G-75 column, then isoelectric focusing to give two protein peaks [Nordberg et al. *Biochem J* 126 491 1972].

Methadone hydrochloride (2-dimethylamino-4-ethoxycarbonyl-4,4-diphenylbutane HCl) [1095-90-5] M 345.9, m 241-242°, pK₁²⁵ 8.94, pK₂²⁰ 10.12 (free base). Crystd from EtOH.

Methoxantin coenzyme (PQQ, pyrrolo quinoline quinone, 2,7,9-tricarboxy-1H-pyrrolo-[2,3-f]-quinoline-4,5-dione, 4,5-dihydro-4,5-dioxo-1H-pyrrolo[2,3-f]quinoline-2,7,9-tricarboxylic acid) [72909-34-3] M 330.2, m 220°(dec). Efflorescent yellow-orange needles on recrystn from H₂O by addition of Me₂CO, or better from a supersaturated aqueous soln, as it forms an acetone adduct. [Forrest et al. *Nature* 280 843 1979.] It has also been purified by passage through a C-18 reverse phase silica cartridge or a silanised silica gel column in aqueous soln whereby methoxantin remains behind as a red-orange band at the origin. This band is collected and washed thoroughly with dilute aqueous HCl (pH 2) and is then eluted with MeOH-H₂O (7:3) and evapd *in vacuo* to give the coenzyme as a red solid. It has also been purified by dissolving in aqueous 0.5M K₂CO₃ and acidified to pH 2.5 whereby PQQ pptes as a deep red solid which is collected and dried *in vacuo*. Methoxantin elutes at 3.55 retention volumes from a C18 μ Bondapak column using H₂O-MeOH (95:5) + 0.1% AcOH pH 4.5. It has UV λ_{\max} at 247 and 330nm (shoulder at 270nm) in H₂O and λ_{\max} at 250 and 340nm in H₂O at pH 2.5. With excitation at λ_{ex} 365nm it has a λ_{\max} emission at 483nm. The ¹³C NMR has δ : 113.86, 122.76, 125.97, 127.71, 130.68, 137.60, 144.63, 146.41, 147.62, 161.25, 165.48, 166.45, 173.30 and 180.00.

When a soln in 10% aqueous MeCO is adjusted to pH 9 with aqueous NH₃ and kept at 25° for 30 min, the *acetone adduct* is formed; UV has λ_{\max} at 250, 317 and 360nm (H₂O, pH 5.5) and with λ_{ex} at 360nm it has max fluorescence at λ_{\max} at 465nm; and the ¹³C NMR [(CD₃)₂SO, TMS] has δ : 29.77, 51.06, 74.82, 111.96, 120.75, 121.13, 125.59, 126.88, 135.21, 139.19, 144.92, 161.01, 161.47, 165.17, 168.61, 190.16 and 207.03. It also forms a *methanol adduct*.

When it is reacted with Me₂SO₄-K₂CO₃ in dry Me₂NCHO at 80° for 4h, it forms the *trimethyl ester* which has m 265-267°(dec) [260-263°(dec)] after recrystn from hot MeCN (orange crystals) with UV λ_{\max} at 252 and 344nm (H₂O) and 251, 321 and 373nm (in MeOH; MeOH adduct ?). [Duine et al. *Eur J Biochem* 108 187 1980; Duine et al. *Adv Enzymology* 59 169 1987; Corey and Tramontano *J Am Chem Soc* 103 5599 1981; Gainor and Weinreb *J Org Chem* 46 4319 1981; Hendrickson and de Vries *J Org Chem* 17 1148 1982; McKenzie, Moody and Reese *J Chem Soc Chem Commun* 1372 1983.]

Methyl benzylpenicillinate [653-89-4] M 348.3, m 97°, $[\alpha]_D^{20}$ +328° (c 1, MeOH). Crystd from CCl₄.

5-Methylphenazinium methyl sulfate [299-11-6] M 306.3, m 155-157° (198°dec by rapid heating). It forms yellow prisms from EtOH (charcoal). Solubility in H₂O at 20° is 10%. In the presence of aqueous KI it forms a *semiquinone* which crystallises as blue leaflets from EtOH. [Wieland and Roseen *Chem Ber* 48 1117 1913; Voriskova *Collect Czech Chem Commun* 12 607 1947; Bülow *Chem Ber* 57 1431 1924.]

1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine hydrochloride (MPTP) [23007-85-4] M 209.7, m 196-198°, pK_{Est} ~ 9.3. Purified by recrystn from Me₂CO + isoPrOH. The *free base* has b 137-142°/0.8 mm, n_D²⁵ 1.5347. [Schmidle and Mansfield *J Am Chem Soc* 78 425 1956; Defeudis *Drug Dev Res* 15 1 1988.]

6- α -Methylprednisolone (Medrol, 11 β ,17-21-trihydroxy-6 α -methylpregna-1,4-dien-3,20-dione) [83-43-2] M 347.5, m 226-237°, 228-237°, 240-242°, $[\alpha]_D^{24}$ +91° (c 0.5, dioxane). Recrystd from EtOAc. UV has λ_{\max} in 95% EtOH 243nm (ϵ 14,875). The *21-acetoxy derivative* has m 205-

208° (from EtOAc), $[\alpha]_D^{24} +95^\circ$ (c 1, CHCl₃). [Spero et al. *J Am Chem Soc* **78** 6213 1956; Fried et al. *J Am Chem Soc* **81** 1235 1959; ¹H NMR: Slomp and McGarvey *J Am Chem Soc* **81** 2200 1959.]

5-Methyltetrahydrofolic acid disodium salt (prefolic A) [68792-52-9] M 503.4, pK₁ 2.4 (N10 protonation), pK₂ 2.7 (pyrimidine N1 protonation), pK₃ 3.5 (α-CO₂H), pK₄ 4.9 (γ-CO₂H), pK₅ 5.6 (N5-Me), pK₆ 8.5 (3NHCO acidic). Check purity by measuring UV at pH 7.0 (use phosphate buffer) and it should have λ_{max} 290nm and λ_{min} 245nm with a ratio of A₂₉₀/A₂₅₀ of 3.7. This ratio goes down to 1.3 as oxidation to the dihydro derivative occurs. The latter can be reduced back to the tetrahydro compound by reaction with 2-mercaptoethanol at room temp. If oxidation had occurred then the compound should be chromatographed on DEAE-cellulose (~0.9 milliequiv/g, in AcO⁻ form) in (NH₄)₂CO₃ (1.5 M) and washed with 1M NH₄OAc containing 0.01M mercaptoethanol till free from UV absorption and then washed with 0.01M mercaptoethanol. All is done in a nitrogen atmosphere. The reduced folate is then eluted with a gradient between 0.01M mercaptoethanol and 1M NH₄OAc containing 0.01M mercaptoethanol and the fractions with absorption at 290nm are collected. These are evapd under reduced pressure at 25° and traces of NH₄OAc and H₂O are removed at high vacuum/25° (~24-48h). The residue is dissolved in the minimum volume of 0.01M mercaptoethanol and an equivalent of NaOH is added to convert the acid to the diNa salt and evaporated to dryness at high vacuum/25°. The product should have λ_{max} 290nm (ε 32,000) in pH 7.0 buffer. [Sakami *Biochem Prep* **10** 103 1963.]

5-Methyltryptamine hydrochloride (3-[2-aminoethyl]-5-methylindole hydrochloride) [1010-95-3] M 210.7, m 289-291°(dec), 290-292°, pK_{Est(1)} ~ -3 (protonation of ring NH), pK_{Est(2)} ~ 9.0 (CH₂NH₂), pK_{Est(3)} ~ 10.9 (acidic indole NH). Recrystd from H₂O. The *free base* has m 93-95° (from *C₆H₆-cyclohexane), and the *picrate* has m 243°(dec) (from EtOH). [Young *J Chem Soc* 3493 1958; Gaddum et al. *Quart J Exp Physiol* **40** 49 1955; Röhm *Hoppe Seyler's Z Physiol Chem* **297** 229 1954.]

4-Methylumbelliferone(β) hydrate (7-hydroxy-4-methylcoumarin) [90-33-5] M 194.2, m 185-186°, 185-188°, 194-195°, pK_{Est} ~ 10.0 (phenolic OH). Purified by recrystn from EtOH. It is insoluble in cold H₂O, slightly soluble in Et₂O and CHCl₃, but soluble in MeOH and AcOH. It has blue fluorescence in aqueous EtOH, and has UV λ_{max} 221, 251 and 322.5nm in MeOH. IR has ν 3077 br, 1667, 1592, 1385, 1267, 1156, 1130 and 1066 cm⁻¹. The *acetate* has m 153-154°. [Woods and Sapp *J Org Chem* **27** 3703 1962.]

4-Methylumbellifer-7-yl-α-D-glucopyranoside [17833-43-1] M 338.3, m 221-222°, $[\alpha]_D^{20} 237^\circ$ (c 3, H₂O). Recrystd from hot H₂O.

4-Methylumbellifer-7-yl-β-D-glucopyranoside [18997-57-4] M 338.3, m 210-212°, 211°, $[\alpha]_D^{20} -61.5^\circ$ (c 2, pyridine), -89.5° (c 0.5, H₂O for half hydrate). Recrystallises as the half hydrate from hot H₂O. [Constantzas and Kocourek *Collect Czech Chem Commun* **24** 1099 1959; De Re et al. *Ann Chim (Rome)* **49** 2089 1959.]

1-Methyluric acid [708-79-2] M 182.1, m >350°, pK₁ 5.75 (basic), pK₂ 10.6 (acidic). Recrystd from H₂O. [Bergmann and Dikstein *J Am Chem Soc* **77** 691 1955.] It has UV λ_{max} at 231 and 283. nm (pH 3) and 217.5 and 292.5nm (pH >12) [Johnson *Biochem J* **5** 133 1952].

Mevalonic acid lactone [674-26-0] M 130.2, m 28°, b 145-150°/5mm. Purified *via* the *dibenzylethylenediammonium salt* (m 124-125°) [Hofmann et al. *J Am Chem Soc* **79** 2316 1957], or by chromatography on paper or on Dowex-1 (formate) column. [Bloch et al. *J Biol Chem* **234** 2595 1959.] Stored as *N,N'*-dibenzylethylenediamine (DBED) salt, or as the lactone in a sealed container at 0°.

Mevalonic acid 5-phosphate [1189-94-2] M 228.1, pK_{Est(1)} ~ 1.5 (PO₄H₂), pK_{Est(2)} ~ 4.4 (CO₂H), pK_{Est(3)} ~ 6.31 (PO₄H⁻). Purified by conversion to the *tricyclohexylammonium salt* (m 154-156°) by treatment with cyclohexylamine. Crystd from water/acetone at -15°. Alternatively, the phosphate was chromatographed by ion-exchange or paper (Whatman No 1) in a system isobutyric acid/ammonia/water (66:3:30; R_F 0.42). Stored as the cyclohexylammonium salt.

Mevalonic acid 5-pyrophosphate [1492-08-6] M 258.1, $pK_{Est(1)} \sim <2$, $pK_{Est(2)} \sim <2$, $pK_{Est(3)} \sim 3.95$ (PO_4), $pK_{Est(4)} 4.4$ (CO_2H), $pK_{Est(5)} \sim 6.26$ (PO_4). Purified by ion-exchange chromatography on Dowex-1 formate [Bloch et al. *J Biol Chem* **234** 2595 1959], DEAE-cellulose [Skilletar and Kekwick, *Anal Biochem* **20** 171 1967], on by paper chromatography [Rogers et al. *Biochem J* **99** 381 1966]. Likely impurities are ATP and mevalonic acid phosphate. Stored as a dry powder or as a slightly alkaline (pH 7-9) soln at -20° .

Mithramycin A (Aureolic acid, Plicamycin) [18378-89-7] M 1085.2, m 180-183°, $[\alpha]_D^{20} -51^\circ$ (c 0.3, EtOH), $pK_{Est} \sim 9.2$. Purified from $CHCl_3$, and is soluble in MeOH, EtOH, Me_2CO , EtOAc, Me_2SO and H_2O , and moderately soluble in $CHCl_3$, but is slightly soluble in *C_6H_6 and Et_2O . Fluorescent antitumour agent used in flow cytometry. [Thiem and Meyer *Tetrahedron* **37** 551 1981; NMR: Yu et al. *Nature* **218** 193 1968.]

Mitomycin C [50-07-7] M 334.4, m $>360^\circ$, $pK_{Est(2)} \sim 8.0$. Blue-violet crystals form *C_6H_6 -pet ether. It is soluble in Me_2CO , MeOH and H_2O , moderately soluble in *C_6H_6 , CCl_4 and Et_2O but insoluble in pet ether. It has UV λ_{max} at 216, 360 and a weak peak at 560nm in MeOH. [Stevens et al. *J Med Chem* **8** 1 1965; Shirahata and Hirayama *J Am Chem Soc* **105** 7199 1983.]

Muramic acid [R-2(2-amino-2-deoxy-D-glucose-3-yloxy)-propionic acid] [1114-41-6] M 251.2, m 145-150°(dec), 152-154°(dec), 155°(dec), $[\alpha]_D^{25} +109^\circ$ (c 2, H_2O), $+165.0^\circ$ (extrapolated to 0 time) $\rightarrow +123^\circ$ [after 3h (c 3, H_2O)], $pK_{Est(1)} \sim 3.8$ (CO_2), $pK_{Est(2)} \sim 7.7$ (NH_2). It has been recrystd from H_2O or aqueous EtOH as monohydrate which loses H_2O at 80° *in vacuo* over P_2O_5 . Sometimes contains some NaCl. It has been purified by dissolving 3.2g in MeOH (75mL), filtered from some insoluble material, concentrated to ~ 10 mL and refrigerated. The colourless crystals are washed with absolute MeOH. This process does not remove NaCl; to do so the product is recrystd from a equal weight of H_2O to give a low yield of very pure acid (0.12g). On paper chromatography 0.26 μ g give one ninhydrin positive spot after development with 75% phenol (R_F 0.51) or with *sec*-BuOH- HCO_2O - H_2O (7:1:2) (R_F 0.30). [Matsushima and Park *Biochem Prep* **10** 109 1963; *J Org Chem* **27** 3581 1962.] The acid has been also purified by dissolving 990mg in 50% aqueous EtOH (2mL), cooling, collecting the colourless needles on a sintered glass funnel and dried over P_2O_5 at $80^\circ/0.1$ mm to give the anhydrous acid. [Lambert and Zilliken *Chem Ber* **93** 2915 1960.] Alternatively the acid is dissolved in a small volume of H_2O , neutralised to pH 7 with ion exchange resin beads (IR4B in OH^- form), filtered, evaporated and dried. The residue is recrystd from 90% EtOH (v/v) and dried as above for 24h. [Strange and Kent *Biochem J* **71** 333 1959.] The *N*-acetyl derivative has m $\sim 125^\circ$ (dec) and $[\alpha]_D^{20} +41.2^\circ$ after 24h (c 1.5, H_2O). [Watanabe and Saito *J Bacteriol* **144** 428 1980.]

Muscimol (pantherine, 5-aminoethyl-3[2h]-isoxazolone) [2763-96-4] M 114.1, m 170-172°(dec), 172-174°(dec), 172-175°, 175°, 176-178°(dec), $pK_{Est(1)} \sim 6$ (acidic, ring 2-NH), $pK_{Est(2)} \sim 8$ ($CH_2CH_2NH_2$). Recrystd from MeOH-tetrahydrofuran or EtOH and sublimed at 110-140° (bath) at 10^{-4} mm and gives a yellow spot with ninhydrin which slowly turns purple [NMR: Bowden et al. *J Chem Soc (C)* **172** 1968]. Also purified by dissolving in the minimum volume of hot H_2O and adding EtOH dropwise until cloudy, cool, and colourless crystals separate; IR: ν 3445w, 3000-2560w br, 2156w, 1635s and 1475s cm^{-1} . [NMR: Jager and Frey *Justus Liebigs Ann Chem* **817** 1982.] Alternatively it has been purified by two successive chromatographic treatments on Dowex 1 x 8 with the first elution with 2M AcOH and a second with a linear gradient between 0—2M AcOH and evaporating the desired fractions and recrystallising the residue from MeOH. [McCarry and Savard *Tetrahedron Lett* **22** 5153 1981; Nakamura *Chem Pharm Bull Jpn* **19** 46 1971.]

Mycophenolic acid (6-[1,3-dihydro-7-hydroxy-5-methoxy-4-methyl-1-oxoisobenzofuran-6-yl]-4-methylhex-4-enoic acid) [24280-93-1] M 320.3, m 141°, 141-143°, $pK_{Est(1)} \sim 2.5$ (CO_2H), $pK_{Est(2)} \sim 9.5$ (phenolic OH). Purified by dissolving in the minimum volume of EtOAc, applying to a silica gel column (0.05-0.2 mesh) and eluting with a mixture of EtOAc + $CHCl_3$ + AcOH (45:55:1) followed by recrystn from heptane-EtOAc, from aqueous EtOH or from hot H_2O and drying *in vacuo*. It is a weak dibasic acid moderately soluble in Et_2O , $CHCl_3$ and hot H_2O but weakly soluble in *C_6H_6 and

toluene. [Birch and Wright *Aust J Chem* **22** 2635 1969; Canonica et al. *J Chem Soc Perkin Trans 1* 2639 1972; Birkinshaw, Raistrick and Ross *Biochem J* **50** 630 1952.]

Myoglobin (from sperm whale muscle). [9047-17-0] M_r ~17,000. Purified by CM-cellulose chromatography and Sephadex G-50 followed by chromatography on Amberlite IRC-50 Type III or BioRex 70 (<400mesh). The crystalline product as a paste in saturated $(\text{NH}_4)_2\text{SO}_4$ at pH 6.5-7.0 may be stored at 4° for at least 4 years unchanged, but must not be kept in a freezer. [Anres and Atassi *Biochemistry* **12** 942 1980; Edmundson *Biochem Prep* **12** 41 1968.]

Myricetin (Cannabiscetin, 3,3',4',5,5',7-hexahydroxyflavone) [529-44-2] M 318.2, m >300°, 357°(dec) (polyphenolic $pK_{\text{Est}} \sim 8-11$). Recrystd from aq EtOH (m 357° dec, as monohydrate) or Me_2CO (m 350° dec, with one mol of Me_2CO) as yellow crystals. Almost insol in CHCl_3 and AcOH. The hexaacetate has m 213°. [Hergert *J Org Chem* **21** 534 1956; Spada and Camerani *Gazzetta* **86** 965, 975 1956; Kalff and Robinson *J Chem Soc* **127** 181 1925.]

Nalidixic acid (1-ethyl-7-methyl-1,8-naphthyridin-4-one-3-carboxylic acid) [389-08-2] M 232.3, m 226.8-230.2°, 228-230°, 229-230°, pK 6.0. Crystd from H_2O or EtOH as a pale buff powder. It is soluble at 23° in CHCl_3 (3.5%), toluene (0.16%), MeOH (0.13%), EtOH (0.09%), H_2O (0.01% and Et_2O (0.01%). It inhibits nucleic acid and protein synthesis in yeast. [Leshner et al. *J Med and Pharm Chem* **5** 1063 1962.]

Naloxone hydrochloride hydrate (Narcan, 1-N-propenyl-7,8-dihydro-14-hydroxymorphinan-6-one hydrochloride) [51481-60-8] M 399.9, m 200-205°, $[\alpha]_D^{20}$ -164° (c 2.5, H_2O), $pK_{\text{Est}(1)} \sim 6$ (N-propenyl), $pK_{\text{Est}(2)} \sim 9.6$ (phenolic OH). This opiate antagonist has been recrystd from EtOH + Et_2O or H_2O . It is soluble in H_2O (5%) and EtOH but insoluble in Et_2O . The free base has m 184° (177-178°) after recrystn from EtOAc, $[\alpha]_D^{20}$ -194.5° (c 0.93, CHCl_3). [Olofson et al. *Tetrahedron Lett* 1567 1977; Gold et al. *Med Res Rev* **2** 211 1982.]

Naltrexone hydrochloride dihydrate (1-N-cyclopropylmethyl-7,8-dihydro-14-hydroxymorphinan-6-one hydrochloride) [16676-29-2] M 413.9, m 274-276°, $[\alpha]_D^{20}$ -173° (c 1, H_2O), $pK_{\text{Est}(1)} \sim 6$ (N-cyclopropylmethyl), $pK_{\text{Est}(2)} \sim 9.6$ (phenolic OH). This narcotic antagonist has been purified by recrystn from MeOH and dried air. The free base has m 168-170° after recrystn from Me_2CO . [Cone et al. *J Pharm Sci* **64** 618 1975; Gold et al. *Med Res Rev* **2** 211 1982.]

α -Naphthoflavone (7,8-benzoflavone) [604-59-1] M 272.3, m 153-155°, 155°, pK 8-9 (phenolic OH). Recrystd from EtOH or aqueous EtOH. [IR: Cramer and Windel *Chem Ber* **89** 354 1956; UV Pillon and Massicot *Bull Soc Chim Fr* 26 1954; Smith *J Chem Soc* 542 1946; Mahal and Venkataraman *J Chem Soc* 1767 1934.] It is a competitive inhibitor of human estrogen synthase. [Kellis and Vickery *Science* **225** 1032 1984.]

Naphthol AS-acetate (3-acetoxynaphthoic acid anilide) [1163-67-3] M 305.3, m 152°, 160°. Recrystd from hot MeOH and dried *in vacuo* over P_2O_5 . It is slightly soluble in AcOH, EtOH, CHCl_3 or $^*\text{C}_6\text{H}_6$. It is a fluorogenic substrate for albumin esterase activity. [Chen and Scott *Anal Lett* **17** 857 1984.] At λ_{ex} 320nm it had fluorescence at λ_{em} 500nm. [Brass and Sommer *Chem Ber* **61** 1000 1928.]

1-Naphthyl phosphate disodium salt [2183-17-7] M 268.1, pK_1^{26} 0.97, pK_2^{26} 5.85 (for free acid). Purified through an acid ion-exchange column (in H^+ form) to give the free acid which is obtained by freeze drying and recrystn from Me_2CO + $^*\text{C}_6\text{H}_6$, or by adding 2.5 vols of hot CHCl_3 to a hot soln of 1 part acid and 1.2 parts Me_2CO and cooling (m 155-157°, 157-158°). The acid is dissolved in the minimum volume of H_2O to which 2 equivalents of NaOH are added and then freeze dried, or by adding the equivalent amount of MeONa in MeOH to a soln of the acid in MeOH and collecting the Na salt, washing with cold MeOH then Et_2O and drying in a vacuum. [Friedman and Seligman *J Am Chem Soc* **72** 624 1950; Chanley and Feageson *J*

Am Chem Soc 77 4002 1955.] It is a substrate for alkaline phosphatase [Gomori *Methods Enzymol* 4 381 1957, 128 212 1968], and prostatic phosphatase [Babson *Clin Chem* 30 1418 1984]. See entry on p. 444.

2-Naphthyl phosphate monosodium salt [14463-68-4] **M 246.2, m 296° (sintering at 228°), pK₁²⁶ 1.28, pK₂²⁶ 5.53 (for free acid).** The *free acid* is purified as for the preceding 1-isomer and has **m 176-177°, 177-178°** after recrystn from CHCl₃ + Me₂CO as the 1-isomer above. It is neutralised with one equivalent of NaOH and freeze dried or prepared as the 1-isomer above. Its solubility in H₂O is 5%. It also forms a 0.5 Na.1 H₂O salt which has **m 203-205° (244° ?)**. [Friedman and Seligman *J Am Chem Soc* 72 624 1950; Chanley and Fegeason *J Am Chem Soc* 77 4002 1955.] See entry on p. 444 in Chapter 5.

D(+)-Neopterin [2009-64-5] **M 253.2, m >300°(dec), [α]₅₄₆²⁰ +64.5° (c 0.14, 0.1M HCl), [α]_D²⁵ +50.1° (c 0.3, 0.1N HCl), pK₁ 2.23 (basic), pK₂ 7.89 (acidic).** Purified as biopterin. Also purified on a Dowex 1 x 8 (formate form) column and eluted with 0.03M ammonium formate buffer pH 8.0 then pH 7.2. The fluorescent neopterin fraction is evapd under reduced pressure leaving neopterin and ammonium formate (the latter can be sublimed out at high vacuum). The residue is stirred for 24h with EtOH and the solid is collected and recrystd from H₂O [Viscontini et al. *Helv Chim Acta* 53 1202 1970; see Wachter et al. Eds *Neopterin* W de Guyter, Berlin 1992].

β-Nicotinamide adenine dinucleotide (diphosphopyridine nucleotide, NAD, DPN) [53-84-9] **M 663.4, [α]_D²³ -34.8° (c 1, H₂O), pK₁ 2.2 (PO₄H), pK₂ 4.0 (adenine NH₂), pK₃ 6.1 (PO₄⁻).** Purified by paper chromatography or better on a Dowex-1 ion-exchange resin. The column was prepared by washing with 3M HCl until free of material absorbing at 260nm, then with water, 2M sodium formate until free of chloride ions and, finally, with water. NAD, as a 0.2% soln in water, adjusted with NaOH to pH 8, was adsorbed on the column, washed with water, and eluted with 0.1M formic acid. Fractions with strong absorption at 360nm were combined, acidified to pH 2.0 with 2M HCl, and cold acetone (*ca* 5L/g of NAD) was added slowly and with constant agitation. It was left overnight in the cold, then the ppte was collected in a centrifuge, washed with pure acetone and dried under vacuum over CaCl₂ and paraffin wax shavings [Kornberg *Methods Enzymol* 3 876 1957]. Purified by anion-exchange chromatography [Dalziel and Dickinson *Biochemical Preparations* 11 84 1966.] The purity is checked by reduction to NADH (with EtOH and yeast alcohol dehydrogenase) which has ε_{340nm} 6220 M⁻¹cm⁻¹. [Todd et al. *J Chem Soc* 3727,3733 1957.] [pKa, Lamborg et al. *J Biol Chem* 231 685 1958.] The *free acid* crystallises from aq Me₂CO with 3H₂O and has **m 140-142°**. It is stable in cold neutral aqueous solns in a desiccator (CaCl₂) at 25°, but decomposes at strong acid and alkaline pH. Its purity is checked by reduction with yeast alcohol dehydrogenase and EtOH to NADH and noting the OD at 340nm. Pure NADH has ε₃₄₀ 6.2 x 10⁴M⁻¹cm⁻¹, i.e. 0.1μmole of NADH in 3mL and in a 1cm path length cell has an OD at 340nm of 0.207.

β-Nicotinamide adenine dinucleotide reduced di-Na salt trihydrate (reduced diphosphopyridine nucleotide sodium salt, NADH) [606-68-8] **M 763.5, pK as for NAD.** This coenzyme is available in high purity and it is advised to buy a fresh preparation rather than to purify an old sample as purification will invariably lead to a more impure sample contaminated with the oxidised form (NAD). It has UV λ_{max} at 340nm (ε 6,200 M⁻¹cm⁻¹) at which wavelength the oxidised form NAD has no absorption. At 340 nm a 0.161mM solution in a 1cm (pathlength) cell has an absorbance of 1.0 unit. The purity is best checked by the ratio A_{280nm}/A_{340nm} ~2.1, a value which increases as oxidation proceeds. The dry powder is stable indefinitely at -20°. Solutions in aqueous buffers at pH ~7 are stable for extended periods at -20° and for at least 8h at 0°, but are oxidised more rapidly at 4° in a cold room (e.g. almost completely oxidised overnight at 4°). [UV: Drabkin *J Biol Chem* 175 563 1945; Fluorescence: Boyer and Thorell *Acta Chem Scand* 10 447 1956; Redox: Rodkey *J Biol Chem* 234 188 1959; Schlenk in *The Enzymes* 2 250, 268 1951; Kaplan in *The Enzymes* 3 105, 112 1960.] Deuterated NADH, i.e. NADD, has been purified through the anion exchange resin AG-1 x 8 (100-200 mesh, formate form) and through a Bio-Gel P-2 column. [Viola, Cook and Cleland *Anal Biochem* 96 334 1979.]

β-Nicotinamide adenine dinucleotide phosphate (NADP, TPN) [53-59-8] **M 743.4, pK₁ 1.1 (PO₄H₂), pK₂ 4.0 (adenine NH₂), pK₃ 6.1 (PO₄⁻).** Purified by anion-exchange chromatography in much the same way as for NAD [Dalziel and Dickinson *Biochem J* 95 311 1965; *Biochemical Preparations* 11 87 1966]. Finally it is purified by dissolving in H₂O and precipitating with 4 volumes of Me₂CO and dried *in*

vacuo over P₂O₅. It is unchanged by storing *in vacuo* at 2°. [Hughes et al. *J Chem Soc* 3733 1957, Schuster and Kaplan *J Biol Chem* 215 183 1955.] Deuterated NADPH, i.e. NADPD, has been purified through the anion exchange resin AG-1 x 8 (100-200 mesh, formate form) and through a Bio-Gel P-2 column. λ_{\min} 259nm (ϵ 18.000) at pH 7.0. [Viola, Cook and Cleland *Anal Biochem* 96 334 1979.]

β -Nicotinamide adenine dinucleotide phosphate reduced tetrasodium salt (reduced diphosphopyridine nucleotide phosphate sodium salt, NADPH) [2646-71-1] M 833.4, pK as for NADP. Mostly similar to NADH above.

β -Nicotinamide mononucleotide (NMN) [1094-61-7] M 334.2, $[\alpha]_{\text{D}}^{23}$ -38.3° (c 1, H₂O), pK_{Est} ~ 6.1 (PO₄⁻). Purified by passage through a Dowex 1 (Cl⁻ form), washed with H₂O until no absorbance at 260 nm. The tubes containing NMN are pooled, adjusted to pH 5.5-6 and evapd *in vacuo* to a small volume. This is adjusted to pH 3 with dilute HNO₃ in an ice bath and treated with 20 volumes of Me₂CO at 0-5°. The heavy white ppt is collected by centrifugation at 0°. It is best stored wet and frozen or can be dried to give a gummy residue. It has λ_{\max} 266nm (ϵ 4600) and λ_{\min} 249nm (ϵ 3600) at pH 7.0 (i.e. no absorption at 340nm). It can be estimated by reaction with CN⁻ or hydrosulfite which form the 4-adducts equivalent to NADH which has UV λ_{\max} 340nm (ϵ 6200). Thus after reaction an OD₃₄₀ of one is obtained from a 0.1612mM soln in a 1cm path cuvette. [Plaut and Plaut *Biochem Prep* 5 56 1957; Maplan and Stolzenbach *Methods Enzymol* 3 899 1957; Kaplan et al. *J Am Chem Soc* 77 815 1955.]

(-)-Nicotine (1-methyl-2[3-pyridyl]-pyrrolidine) [54-11-5] M 162.2, b 123-125°/17mm, 246.1°/730.5mm, 243-248°/atm (partial dec), d_4^{20} 1.097, n_{D}^{20} 1.5280, $[\alpha]_{\text{D}}^{20}$ -169° (c 1, Me₂CO), pK₁¹⁵ 6.16 (pyridine N⁺), pK₂¹⁵ 10.96 (pyrrolidine N⁺). Very pale yellow *hygroscopic* oil with a characteristic odour (tobacco extract) with browns in air on exposure to light. Purified by fractional distn under reduced pressure in an inert atmosphere. A freshly distd sample should be stored in dark sealed containers under N₂. It is a strong base, at 0.05 M soln it has a pH of 10.2. Very soluble in organic solvents. It is soluble in H₂O and readily forms salts. [UV: Parvis *J Chem Soc* 97 1035 1910; Dobbie and Fox *J Chem Soc* 103 1194 1913.] The *hydrochlorides* (mono- and di-) form deliquescent crystals soluble in H₂O and EtOH but insoluble in Et₂O. It has also been purified *via* the ZnCl₂ double salt. [Ratz *Monatsh Chem* 26 1241 1905; Biosynthesis: Nakan and Hitchinson *J Org Chem* 43 3922 1978.] The *picrate* has m 218° (from EtOH). **POISONOUS.**

(±)-Nicotine [22083-74-5] M 162.2, b 242.3°/atm, d_4^{20} 1.082 (pK see above). Purified by distn. Its solubility in EtOH is 5%. The *picrate* forms yellow needles from hot H₂O and has m 218°. The *methiodide* has m 219° (from MeOH).

Nisin [1414-45-5] M 3354.2. Polypeptide from *S. lactis*. Crystd from EtOH. [Berridge et al. *Biochem J* 52 529 1952; synthesis by Fukase et al. *Tetrahedron Lett* 29 795 1988.]

2-Nitrophenyl- β -D-galactopyranoside [369-07-3] M 301.3, m 185-190°, 193°, 193-194°, $[\alpha]_{\text{D}}^{18}$ -51.9° (c 1, H₂O). Purified by recrystn from EtOH. [Seidman and Link *J Am Chem Soc* 72 4324 1950; Snyder and Link *J Am Chem Soc* 75 1758 1953]. It is a chromogenic substrate for β -galactosidases [Jagota et al. *J Food Sci* 46 161 1981].

4-Nitrophenyl- α -D-galactopyranoside [7493-95-0] M 301.3, m 166-169°, 173°, $[\alpha]_{\text{D}}^{25}$ +248 (c 1, H₂O). Purified by recrystn from H₂O or aqueous EtOH. The *monohydrate* has m 85° which resolidifies and melts at 151-152° (the hemihydrate) which resolidifies and melts again at 173° as the anhydrous form. Drying the monohydrate at 60° yields the hemihydrate and drying at 100° gives the anhydrous compound. The *tetraacetate* has m 147° after drying at 100°. [Jermyn *Aust J Chem* 15 569 1962; Helfreich and Jung *Justus Liebigs Ann Chem* 589 77 1954.] It is a substrate for α -galactosidase [Dangelmaier and Holmsen *Anal Biochem* 104 182 1980].

4-Nitrophenyl- β -D-galactopyranoside [3150-24-1] M 301.3, m 178°, 178-181°, 181-182°, $[\alpha]_{\text{D}}^{20}$ -83° (c 1, H₂O). Purified by recrystn from EtOH. [Horikoshi *J Biochem (Tokyo)* 35 39 1042;

Goebel and Avery *J Exptl Medicine* **50** 521 1929; Snyder and Link *J Am Chem Soc* **75** 1758.] It is a chromogenic substrate for β -galactosidases [Buoncore et al. *J Appl Biochem* **2** 390 1980].

4-Nitrophenyl- α -D-glucopyranoside [3767-28-0] **M 301.3, m 206-212 $^{\circ}$, 216-217 $^{\circ}$ (sinters at 210 $^{\circ}$), $[\alpha]_{\text{D}}^{20} +215^{\circ}$ (c 1, H₂O).** Purified by recrystn from H₂O, MeOH or EtOH. [Jermyn *Aust J Chem* **7** 202 1954; Montgomery et al. *J Am Chem Soc* **64** 690 1942.] It is a chromogenic substrate from α -glucosidases [Oliviera et al. *Anal Biochem* **113** 188 1981], and is a substrate for glucansucrases [Binder and Robyt *Carbohydr Res* **124** 287 1983]. It is a chromogenic substrate for β -glucosidases [Weber and Fink *J Biol Chem* **255** 9030 1980].

4-Nitrophenyl- β -D-glucopyranoside [2492-87-7] **M 301.2, m 164 $^{\circ}$, 164-165 $^{\circ}$, 165 $^{\circ}$, $[\alpha]_{\text{D}}^{20} -107^{\circ}$ (c 1, H₂O).** Purified by recrystn from EtOH or H₂O. [Montgomery et al. *J Am Chem Soc* **64** 690 1942; Snyder and Link *J Am Chem Soc* **75** 1758 1953.]

Nonactin [6833-84-7] **M 737.0, m 147-148 $^{\circ}$, $[\alpha]_{\text{D}}^{20} 0^{\circ}$ ($\pm 2^{\circ}$) (c 1.2, CHCl₃).** This macrotetrolide antibiotic was rerystd from MeOH as colourless needles, and dries at 90 $^{\circ}$ /20h/high vacuum. [*Helv Chim Acta* **38** 1445 1955, **55** 1371 1972; *Tetrahedron Lett* 3391 1975.]

N-Nonanoyl-n-methylglucamine (Mega-9) [85261-19-4] **M 335.4, m 87-89 $^{\circ}$.** A non-ionic detergent purified as *n*-decanoyl-*N*-methylglucamine above. [Hildreth *Biochem J* **207** 363 1982.]

Nonyl- β -D-glucopyranoside [69984-73-2] **M 306.4, m 67.5-70 $^{\circ}$, $[\alpha]_{\text{D}}^{20} -34.4^{\circ}$ (c 5, H₂O), $[\alpha]_{\text{D}}^{25} -28.8^{\circ}$ (c 1, MeOH).** Purified by recrystn from Me₂CO and stored in well stoppered containers as it is *hygroscopic*. [Pigman and Richtmyer *J Am Chem Soc* **64** 369 1942.] It is a UV transparent non-ionic detergent for solubilising membrane proteins [Schwendener et al. *Biochem Biophys Res Commun* **100** 1055 1981.]

L-Noradrenaline (Adrenor, R-2-amino-1-[3,4-dihydroxyphenyl]ethan-1-ol, L-norepinephrine) [51-41-2, 69815-49-2 (*bitartrate salt*)] **M 169.2, m 216.5-218 $^{\circ}$ (dec), ~220-230 $^{\circ}$ (dec), $[\alpha]_{\text{D}}^{20} -45^{\circ}$ (c 5, N HCl), $[\alpha]_{\text{D}}^{25} 37.3^{\circ}$ (c 5, 1 equiv aqueous HCl), $\text{pK}_1^{25} 5.58$ (phenolic OH), $\text{pK}_2^{25} 8.90$ (phenolic OH), $\text{pK}_3^{25} 9.78$ (NH₂).** Recrystd from EtOH and stored in the dark under N₂. [pKa, Lewis *Brit J Pharmacol Chemother* **9** 488 1954; UV: Bergström et al. *Acta Physiol Scand* **20** 101 1950; Fluorescence: Bowman et al. *Science NY* **122** 32 1955; Tullar *J Am Chem Soc* **70** 2067 1948.] The *L-tartrate salt monohydrate* has **m 102-104.5 $^{\circ}$, $[\alpha]_{\text{D}}^{25} -11^{\circ}$ (c 1.6, H₂O),** after recrystn from H₂O or EtOH.

L-Noradrenaline hydrochloride (Arterenol) [329-56-6] **M 205.6, m 145.2-146.4 $^{\circ}$, ~150 $^{\circ}$ (dec), $[\alpha]_{\text{D}}^{25} -40^{\circ}$ (c 6, H₂O), **pK** see above.** Recrystd from isoPrOH and stored in the dark as it is oxidised in the presence of light (see preceding entry). [Tullar *J Am Chem Soc* **70** 2067 1948.]

Novobiocin (7-[O³-carbamoyl-5-O⁴-dimethyl- β -L-lyso-6-desoxyhexahydropyranosyloxy]-4-hydroxy-3[4-hydroxy-3-(3-methylbut-2-enyl)-benzyl-amino]-8-methylcoumarin) [303-81-1] **M 612.6, two forms m 152-156 $^{\circ}$ and m 172-174 $^{\circ}$, 174-178 $^{\circ}$, λ_{max} at 330nm (acid EtOH), 305nm (alk EtOH), $[\alpha]_{\text{D}}^{25} -63^{\circ}$ (c 1, EtOH), $\text{pK}_1 4.03$ (4.2), $\text{pK}_2 9.16$.** Crystd from EtOH and stored in the dark. It has also been recrystd from Me₂CO-H₂O. [Hoeksema et al. *J Am Chem Soc* **77** 6710 1955; Kaczka et al. *J Am Chem Soc* **77** 9404 1955.]

The **sodium salt** [1476-53-5] **M 634.6, m 210-215 $^{\circ}$, 215-220 $^{\circ}$ (dec), 222-229 $^{\circ}$, $[\alpha]_{\text{D}}^{25} -38^{\circ}$ (c 1, H₂O)** has been recrystd from MeOH, then dried at 60 $^{\circ}$ /0.5mm. [Sensi, Gallo and Chiesa, *Anal Chem* **29** 1611 1957; Kaczka et al. *J Am Chem Soc* **78** 4126 1956.]

5'-Nucleotidase (from Electric ray, *Torpedo sp*) [9027-73-0] [**EC 3.1.3.5**], **amorphous.** Purified by dissolving in Triton X-100 and deoxycholate, and by affinity chromatography on concanavalin A-Sepharose and AMP-Sepharose [Grondal and Zimmerman *Biochem J* **245** 805 1987].

Nucleotide thiophosphate analogues. The preparation and purification of [³H]ATP γ S, [³H]GTP γ S, s⁶ITP γ S (6-thioinosine), cl⁶ITP γ S (6-chloroinosine) and [³H]ATP γ S are described and the general purification

was achieved by chromatography of the nucleotide thiophosphates in the minimum volume of H₂O placed onto a DEAE-Sephadex A25 column and eluting with a linear gradient of triethylammonium bicarbonate (0.1 to 0.6M for G and I nucleotides and 0.2 to 0.5M for A nucleotides). [*Biochim Biophys Acta* **276** 155 1972.]

Nystatin dihydrate (Mycostatin, Fungicidin) [1400-61-9] **M 962.1, m dec>160° (without melting by 250°),** $[\alpha]_D^{25} -7^\circ$ (0.1N HCl in MeOH), -10° (AcOH), $+12^\circ$ (Me₂NCHO), $+21^\circ$ (pyridine). Light yellow powder with the following solubilities at ~28°: MeOH (1.1%), ethylene glycol (0.9%), H₂O (0.4%), CCl₄ (0.12%), EtOH (0.12%), CHCl₃ (0.05%) and *C₆H₆ (0.03%). Could be pptd from MeOH soln by addition of H₂O. Aqueous suspensions of this macrolide antifungal antibiotic are stable at 100°/10min at pH 7.0 but decomposes rapidly at pH <2 and >9, and in the presence of light and O₂. [Birch et al. *Tetrahedron Lett* 1491, 1485 1964; Weiss et al. *Antibiot Chemother* **7** 374 1957.] It contains a mixture of components A₁, A₂ and A₃.

Octyl-β-D-glucopyranoside [29836-26-8] **M 292.4, m 62-65°, 63.8-65°,** $[\alpha]_D^{20} -34^\circ$ (c 4, H₂O). Purified by recrystn from Me₂CO. It is *hygroscopic* and should be stored in a well stoppered container. [Noller and Rockwell *J Am Chem Soc* **60** 2076 1938; Pigman and Richtmyer *J Am Chem Soc* **64** 369 1942.] It is a UV transparent non-ionic dialysable detergent for solubilising membrane proteins. The α-D-isomer with $[\alpha]_D^{20} +118^\circ$ (c 1, MeOH) has similar solubilising properties. [Lazo and Quinn *Anal Biochem* **102** 68 1980; Stubbs et al. *Biochim Biophys Acta* **426** 46 1976.]

Orcine monohydrate (3,5-dihydroxytoluene) [6153-39-5] **M 142.2, m 56°, 56-58°, 58°, b 147°/5 mm, pK₁²⁰ 9.48 (9.26), pK₂²⁰ 11.20 (11.66).** Purified by recrystn from H₂O as the monohydrate. It sublimes *in vacuo* and the *anhydrous* compound has **m 106.5-108° (110°, 108°)**. Also can be recrystd from CHCl₃ (plates) or *C₆H₆ (needles or prisms). [UV: Kiss et al. *Bull Soc Chim Fr* 275 1949; Adams et al. *J Am Chem Soc* **62** 732 1940.]

Orosomucoid (glycoprotein α₁ acid, from human plasma) [66455-27-4] **M_r 42000-44000, amorphous.** Purified by passage through a carboxymethyl cellulose column and through a Sephadex G-25 column. [Aronson et al. *J Biol Chem* **243** 4564 1968.]

Orotic acid Li salt H₂O (1-carboxy-4,6-dihydroxypyrimidine Li salt H₂O) [5266-20-6] **M 180.0, m >300°, pK₁ 2.8 (CO₂H), pK₂ 9.4 (OH), pK₃ >13 (OH) (for free acid).** It is soluble in H₂O at 17° and 100°. Best to acidify an aqueous soln, isolating the free acid which is recrystd from H₂O (as monohydrate) **m 345-347° (345-346°)**, then dissolving in EtOH, adding an equivalent amount of LiOH in EtOH and evaporating. Its solubility in H₂O is 1.28% (17°) and 2.34% (100°). [Bachstz *Chem Ber* **63** 1000 1930; Johnson and Shroeder *J Am Chem Soc* **54** 2941 1932; UV: Shugar and Fox *Biochim Biophys Acta* **9** 199 1952.]

Oxacillin sodium salt (5-methyl-3-phenyl-4-isoxazolylpenicillin sodium salt) [1173-88-2] **M 423.4, m 188°(dec),** $[\alpha]_D^{20} +29^\circ$ (c 1, H₂O), **pK_{Est} ~ 2.7.** This antibiotic which is stable to penicillinase is purified by recrystn from isoPrOH and dried *in vacuo*. Its solubility in H₂O at 25° is 5%. [Doyle et al. *Nature* **192** 1183 1961.]

Oxolinic acid (5-ethyl-5,8-dihydro-8-oxo-1,3-dioxolo[4,5-g]quinoline-3-carboxylic acid) [14698-29-4] **M 261.2, m 313-314°(dec), 314-316°(dec), pK_{Est} ~ 2.3.** Purified by recrystn from aqueous Me₂CO or 95% EtOH. It has UV λ_{max} 220, (255.5sh), 259.5, 268, (298sh, 311sh), 321 and 326nm [ϵ 14.8, (36.8sh), 38.4, 38.4, (6.4sh, 9.2sh), 10.8 and 11.2 x 10³]. [Kaminsky and Metzger *J Med Chem* **11** 160 1968.]

Oxytocin [50-56-6] **M 1007.2, m dec on heating, $[\alpha]_D^{22} -26.2^\circ$ (c 0.53, N AcOH).** A cyclic nonapeptide which was purified by countercurrent distribution between solvent and buffer. It is soluble in H₂O, *n*-BuOH and isoBuOH. [Bodanszky and du Vigneaud *J Am Chem Soc* **81** 2504 1959; Cash et al. *J Med Pharm Chem* **5** 413 1962; Sakakibara et al. *Bull Chem Soc Jpn* **38** 120 1965; solid phase synthesis: Bayer and

Hagenmyer *Tetrahedron Lett* 2037 1968.] It was also synthesised on a solid phase matrix and finally purified as follows: A Sephadex G-25 column was equilibrated with the aqueous phase of a mixture of 3.5% AcOH (containing 1.5% of pyridine) + *n*-BuOH + ¹⁴C₆H₆ (2:1:1) and then the organic phase of this mixture was run through. A soln of oxytocin (100mg) in H₂O (2mL) was applied to the column which was then eluted with the organic layer of the above mixture. The fractions containing the major peak [as determined by the Folin-Lowry protein assay [Fryer et al. *Anal Biochem* 153 262 1986] were pooled, diluted with twice their vol of H₂O, evaporated to a small vol and lyophilised to give oxytocin as a pure white powder (20mg, 508 U/mg). [Ives *Can J Chem* 46 2318 1968.]

Palmitoyl coenzyme A [1763-10-6] **M 1005.9**. Possible impurities are palmitic acid, S-palmitoyl thioglycolic acid and S-palmitoyl glutathione. These are removed by placing *ca* 200mg in a centrifuge tube and extracting with Me₂CO (20mL), followed by two successive extractions with Et₂O (15mL) to remove S-palmitoyl thioglycolic acid and palmitic acid. The residue is dissolved in H₂O (4 x 4 mL), adjusted to pH 5 and centrifuged to remove insoluble S-palmitoyl glutathione and other insoluble impurities. To the clear supernatant is added 5% HClO₄ (6mL) whereby S-palmitoyl CoA pptes. The ppt is washed with 0.8% HClO₄ (10mL) and finally with Me₂CO (3 x 5mL) and dried *in vacuo*. It is stable for at least one year in dry form at 0° in a desiccator (dark). Solns are stable for several months at -15°. Its solubility in H₂O is 4%. The adenine content is used as the basis of purity with λ_{max} at 260 and 232nm (ε 6.4 x 10⁶ and 9.4 x 10⁶ cm²/mol respectively). Higher absorption at 232nm would indicate other thio ester impurities, e.g. S-palmitoyl glutathione, which absorb highly at this wavelength. Also PO₄ content should be determined and acid phosphate can be titrated potentiometrically. [Seubert *Biochem Prep* 7 80 1960; Srer et al. *Biochim Biophys Acta* 33 31 1959; Kornberg and Pricer *J Biol Chem* 204 329, 345 1953.]

3-Palmitoyl-sn-glycerol (*R*-glycerol-1-palmitate, *L*-β-palmitin) [32899-41-5] **M 330.5, d^{27.3} 0.9014, m 66.5° (α-form), 74° (β'-form) and 77° (β-form)**. The stable β-form is obtained by crystn from EtOH or Skellysolve B and recrystn from Et₂O provides the β'-form. The α-form is obtained on cooling the melt. [Malkin and el Sharbagy *J Chem Soc* 1631 1936; Chapman *J Chem Soc* 58 1956; Luton and Jackson *J Am Chem Soc* 70 2446 1948.]

Pancuronium bromide (2β,16β-dipiperidino-5α-androstan-3α,17β-diol diacetate dimethobromide) [15500-66-0] **M 732.7, m 212-215°, 215°**. Odourless crystals with a bitter taste which are purified through acid-washed Al₂O₃ and eluted with isoPrOH-EtOAc (3:1) to remove impurities (e.g. the monomethobromide) and eluted with isoPrOH to give the pure bromide which can be recrystd from CH₂Cl₂-Me₂CO or isoPrOH-Me₂CO. It is soluble in H₂O (50%) and CHCl₃ (3.3%) at 20°. It is a non-depolarising muscle relaxant. [Buckett et al. *J Med Chem* 16 1116 1973.]

D-Panthenol (Provitamin B, *R*-2,4-dihydroxy-3,3-dimethylbutyric acid 3-hydroxypropylamide) [81-13-0] **M 205.3, b 118-120°/0.02mm, d₂₀²⁰ 1.2, n_D²⁰ 1.4935, [α]_D²⁰ (c 5, H₂O)**. Purified by distn *in vacuo*. It is a slightly *hygroscopic* viscous oil. Soluble in H₂O and organic solvent. It is hydrolysed by alkali and strong acid. [Rabin *J Am Pharm Assoc (Sci Ed)* 37 502 1948; Bonati and Pitré *Farmaco Ed Scient* 14 43 1959.]

***R*-(+)-Pantothenic acid sodium salt** (*N*-[2,4-dihydroxy-3,3-dimethylbutyryl] β-alanine Na salt) [867-81-2] **M 241.2, [α]_D²⁵ +27.1° (c 2, H₂O), pK²⁵ 4.4 (for free acid)**. Crystd from EtOH, very *hygroscopic* (kept in sealed ampoules). The free acid is a viscous *hygroscopic* oil with [α]_D²⁵ +37.5° (c 5, H₂O), easily destroyed by acids and bases.

***R*-(+)-Pantothenic acid Ca salt** [(*D*-(+)- 137-08-6; 63409-48-3] **M 476.5, m 195-196°, 200-201°, [α]_D²⁰ +28.2° (c 5, H₂O)**. Crysts in needles from MeOH, EtOH or isoPrOH (with 0.5mol of isoPrOH). Moderately *hygroscopic*. The *S*-benzylisothiuronium salt has **m 151-152° (149° when crystd from Me₂CO)**. [Kagan et al. *J Am Chem Soc* 79 3545 1957; Wilson et al. *J Am Chem Soc* 76 5177 1954; Stiller and Wiley *J Am Chem Soc* 63 1239 1941.]

[Previous Page](#)

Hagenmyer *Tetrahedron Lett* 2037 1968.] It was also synthesised on a solid phase matrix and finally purified as follows: A Sephadex G-25 column was equilibrated with the aqueous phase of a mixture of 3.5% AcOH (containing 1.5% of pyridine) + *n*-BuOH + ¹⁴C₆H₆ (2:1:1) and then the organic phase of this mixture was run through. A soln of oxytocin (100mg) in H₂O (2mL) was applied to the column which was then eluted with the organic layer of the above mixture. The fractions containing the major peak [as determined by the Folin-Lowry protein assay [Fryer et al. *Anal Biochem* 153 262 1986] were pooled, diluted with twice their vol of H₂O, evaporated to a small vol and lyophilised to give oxytocin as a pure white powder (20mg, 508 U/mg). [Ives *Can J Chem* 46 2318 1968.]

Palmitoyl coenzyme A [1763-10-6] **M 1005.9**. Possible impurities are palmitic acid, S-palmitoyl thioglycolic acid and S-palmitoyl glutathione. These are removed by placing *ca* 200mg in a centrifuge tube and extracting with Me₂CO (20mL), followed by two successive extractions with Et₂O (15mL) to remove S-palmitoyl thioglycolic acid and palmitic acid. The residue is dissolved in H₂O (4 x 4 mL), adjusted to pH 5 and centrifuged to remove insoluble S-palmitoyl glutathione and other insoluble impurities. To the clear supernatant is added 5% HClO₄ (6mL) whereby S-palmitoyl CoA pptes. The ppt is washed with 0.8% HClO₄ (10mL) and finally with Me₂CO (3 x 5mL) and dried *in vacuo*. It is stable for at least one year in dry form at 0° in a desiccator (dark). Solns are stable for several months at -15°. Its solubility in H₂O is 4%. The adenine content is used as the basis of purity with λ_{\max} at 260 and 232nm (ϵ 6.4 x 10⁶ and 9.4 x 10⁶ cm²/mol respectively). Higher absorption at 232nm would indicate other thio ester impurities, e.g. S-palmitoyl glutathione, which absorb highly at this wavelength. Also PO₄ content should be determined and acid phosphate can be titrated potentiometrically. [Seubert *Biochem Prep* 7 80 1960; Srer et al. *Biochim Biophys Acta* 33 31 1959; Kornberg and Pricer *J Biol Chem* 204 329, 345 1953.]

3-Palmitoyl-sn-glycerol (*R*-glycerol-1-palmitate, *L*- β -palmitin) [32899-41-5] **M 330.5**, **d^{27.3} 0.9014**, **m 66.5°** (α -form), **74°** (β' -form) and **77°** (β -form). The stable β -form is obtained by crystn from EtOH or Skellysolve B and recrystn from Et₂O provides the β' -form. The α -form is obtained on cooling the melt. [Malkin and el Sharbagy *J Chem Soc* 1631 1936; Chapman *J Chem Soc* 58 1956; Luton and Jackson *J Am Chem Soc* 70 2446 1948.]

Pancuronium bromide (2 β ,16 β -dipiperidino-5 α -androstan-3 α ,17 β -diol diacetate dimethobromide) [15500-66-0] **M 732.7**, **m 212-215°**, **215°**. Odourless crystals with a bitter taste which are purified through acid-washed Al₂O₃ and eluted with isoPrOH-EtOAc (3:1) to remove impurities (e.g. the monomethobromide) and eluted with isoPrOH to give the pure bromide which can be recrystd from CH₂Cl₂-Me₂CO or isoPrOH-Me₂CO. It is soluble in H₂O (50%) and CHCl₃ (3.3%) at 20°. It is a non-depolarising muscle relaxant. [Buckett et al. *J Med Chem* 16 1116 1973.]

D-Panthenol (Provitamin B, *R*-2,4-dihydroxy-3,3-dimethylbutyric acid 3-hydroxypropylamide) [81-13-0] **M 205.3**, **b 118-120°/0.02mm**, **d₂₀²⁰ 1.2**, **n_D²⁰ 1.4935**, **[\alpha]_D²⁰ (c 5, H₂O)**. Purified by distn *in vacuo*. It is a slightly *hygroscopic* viscous oil. Soluble in H₂O and organic solvent. It is hydrolysed by alkali and strong acid. [Rabin *J Am Pharm Assoc (Sci Ed)* 37 502 1948; Bonati and Pitré *Farmaco Ed Scient* 14 43 1959.]

***R*-(+)-Pantothenic acid sodium salt** (*N*-[2,4-dihydroxy-3,3-dimethylbutyryl] β -alanine Na salt) [867-81-2] **M 241.2**, **[\alpha]_D²⁵ +27.1°** (c 2, H₂O), **pK²⁵ 4.4** (for free acid). Crystd from EtOH, very *hygroscopic* (kept in sealed ampoules). The free acid is a viscous *hygroscopic* oil with **[\alpha]_D²⁵ +37.5°** (c 5, H₂O), easily destroyed by acids and bases.

***R*-(+)-Pantothenic acid Ca salt** [(*D*(+)- 137-08-6; 63409-48-3] **M 476.5**, **m 195-196°**, **200-201°**, **[\alpha]_D²⁰ +28.2°** (c 5, H₂O). Crysts in needles from MeOH, EtOH or isoPrOH (with 0.5mol of isoPrOH). Moderately *hygroscopic*. The *S*-benzylisothiuronium salt has **m 151-152°** (149° when crystd from Me₂CO). [Kagan et al. *J Am Chem Soc* 79 3545 1957; Wilson et al. *J Am Chem Soc* 76 5177 1954; Stiller and Wiley *J Am Chem Soc* 63 1239 1941.]

Papain [9001-73-4] M_r ~21,000, [EC 3.4.22.2], **amorphous**. A suspension of 50g of papain (freshly ground in a mortar) in 200mL of cold water was stirred at 4° for 4h, then filtered through a Whatman No 1 filter paper. The clear yellow filtrate was cooled in an ice-bath while a rapid stream of H₂S was passed through it for 3h, and the suspension was centrifuged at 2000rpm for 20min. Sufficient cold MeOH was added slowly and with stirring to the supernatant to give a final MeOH concn of 70 vol%. The ppt, collected by centrifugation for 20min at 2000rpm, was dissolved in 200mL of cold water, the soln was saturated with H₂S, centrifuged, and the enzyme again ppted with MeOH. The process was repeated four times. [Bennett and Niemann *J Am Chem Soc* 72 1798 1950.] Papain has also been purified by affinity chromatography on a column of Gly-Gly-Tyr-Arg-agarose [Stewart et al. *J Am Chem Soc* 109 3480 1986].

Papaverine hydrochloride (6,7-dimethoxy-1-veratrylisoquinoline hydrochloride) [61-25-6] **M 375.9, m 215-220°, 222.5-223.5°(dec), 231°, pK²⁵ 6.41**. Recrystd from H₂O and sublimed at 140°/0.1mm. Solubility in H₂O is 5%. [Saunders and Srivastava *J Pharm Pharmacol* 3 78 1951; Biggs *Trans Faraday Soc* 50 800 1954.] The free base has m 148-150° [Bobbitt *J Org Chem* 22 1729 1957].

Pargyline hydrochloride (Eutonyl, *N*-methyl-*n*-propargylbenzylamine hydrochloride) [306-07-0] **M 195.7, m 154-155°, 155°, pK²⁵ 6.9**. Recrystd from EtOH-Et₂O and dried *in vacuo*. It is very soluble in H₂O, in which it is unstable. The free base has b 101-103°/11mm. It is a glucuronyl transferase inducer and a monoamine oxidase inhibitor. [von Braun et al. *Justus Liebigs Ann Chem* 445 205 1928; Yeh and Mitchell *Experientia* 28 298 1972; Langstrom et al. *Science* 225 1480 1984.]

Pectic acid [9046-40-6] M_r (C₆H₈O₆)_n ~500,000, **amorphous**, [α]_D +250° (c 1, 0.1M NaOH). Citrus pectic acid (500g) was refluxed for 18h with 1.5L of 70% EtOH and the suspension was filtered hot. The residue was washed with hot 70% EtOH and finally with ether. It was dried in a current of air, ground and dried for 18h at 80° under vacuum. [Morell and Link *J Biol Chem* 100 385 1933.] It can be further purified by dispersing in water and adding just enough dilute NaOH to dissolve the pectic acid, then passing the soln through columns of cation- and anion-exchange resins [Williams and Johnson *Ind Eng Chem (Anal Ed)* 16 23 1944], and precipitating with two volumes of 95% EtOH containing 0.01% HCl. The ppt is worked with 95% EtOH, then Et₂O, dried and ground.

Pectin [9000-69-5] M_r 25,000-100,000, **amorphous**. Dissolved in hot water to give a 1% soln, then cooled, and made about 0.05M in HCl by addition of conc HCl, and ppted by pouring slowly, with vigorous stirring into two volumes of 95% EtOH. After standing for several hours, the pectin is filtered onto nylon cloth, then redispersed in 95% EtOH and stood overnight. The ppt is filtered off, washed with EtOH/Et₂O, then Et₂O and air dried.

D-(-)-Penicillamine (*R*-3-mercapto-*D*-valine, 3,3-dimethyl-*D*-cysteine, from natural penicillin) [52-67-5] **M 149.2, m 202-206°, 214-217°, [α]_D²¹ -63° (c 1, N NaOH or pyridine), pK₁²⁰ 2.4 (CO₂H), pK₂²⁰ 8.0 (SH), pK₃²⁰ 10.68 (NH₂)**. The melting point depends on the rate of heating (m 202-206° is obtained by starting at 195° and heating at 2°/min). It is soluble in H₂O and alcohols but insoluble in Et₂O, CHCl₃, CCl₄ and hydrocarbon solvents. Purified by dissolving in MeOH and adding Et₂O slowly. Dried *in vacuo* and stored under N₂. [Weight et al. *Angew Chem, Int Ed Engl* 14 330 1975; Cornforth in *The Chemistry of Penicillin* (Clarke, Johnson and Robinson Eds) Princeton Univ Press, 455 1949; Polymorphism: Vidler *J Pharm Pharmacol* 28 663 1976.] The *D*-*S*-benzyl derivative has m 197-198° (from H₂O), [α]_D¹⁷ -20° (c 1, NaOH), -70° (N HCl).

L-(-)-Penicillamine [1113-41-3] **M 149.2, m 190-194°, 202-206°, 214-217°, [α]_D²¹ +63° (c 1, N NaOH or pyridine)**. Same as preceding entry for its enantiomer.

D-Penicillamine disulfide hydrate (*S,S'*-di-*D*-penicillamine) hydrate) [20902-45-8] **M 296.4 + aq, m 203-204°(dec), 204-205°(dec), [α]_D²³ +27° (c 1.5, N HCl), -82° (c 0.8, N NaOH), pK_{Est(1)} ~ 2.4 (CO₂), pK_{Est(2)} ~ 10.7 (NH₂)**. Purified by recrystn from EtOH or aqueous EtOH. [Crooks in *The Chemistry of Penicillin* (Clarke, Johnson and Robinson Eds) Princeton Univ Press, 469 1949; Use as a thiol reagent for proteins: Garel *Eur J Biochem* 123 513 1982; Süss *Justus Liebigs Ann Chem* 561 31 1948.]

Pepsin [9001-75-6] M_r 31,500(human), 6000(hog) [EC 3.4.23.1]. Rechromatographed on a column of Amberlite CG-50 using a pH gradient prior to use. Crystd from EtOH. [Richmond et al. *Biochim Biophys Acta* 29 453 1958; Huang and Tang, *J Biol Chem* 244 1085 1969, 245 2189 1970.]

Pertussis toxin (from *Bordetella pertussis*) [70323-44-3] M_r 117,000. Purified by stepwise elution from 3 columns comprising Blue Sepharose, Phenyl Sepharose and hydroxylapatite, and SDS-PAGE [Svoboda et al. *Anal Biochem* 159 402 1986; *Biochemistry* 21 5516 1982; *Biochem J* 83 295 1978.]

2-Phenylethyl- β -D-thiogalactoside [63407-54-5] M 300.4, m 108°, $[\alpha]_D^{23}$ -32.2° (c 5, MeOH). Recryst from H₂O and dried in air to give the 1.5.H₂O and has m 80°. Anhydrous surfactant is obtained by drying at 78° over P₂O₅. [Heilfrich and Türk *Chem Ber* 89 2215 1856.]

Phenyl- β -D-galactopyranoside [2818-58-8] M 256.3, m 153-54°, 146-148°, 155-156°(dried at 105°), $[\alpha]_D^{20}$ -42° (c 1, H₂O). Recrystd from H₂O as 0.5H₂O. [Conchie and Hay *Biochem J* 73 327 1959; IR: Whistler and House *Analyt Chem* 25 1463 1953.] It is an acceptor substrate for fucosyltransferase [Chester et al. *Eur J Biochem* 69 583 1976].

Phenyl- β -D-glucopyranoside [1464-44-4] M 256.3, m 174-175° 174-176°, 176°, $[\alpha]_D^{20}$ -72.2° (c 1 for dihydrate, H₂O). Recrystd from H₂O as 2H₂O and can be dried *in vacuo* at 100°/P₂O₅. Dry preparation has $[\alpha]_D^{25}$ -70.7° (c 2, H₂O). [Robertson and Waters *J Chem Soc* 2729 1930; IR: Bunton et al. *J Chem Soc* 4419 1955; Takahashi *Yakugaku Zasshi (J Pharm Soc Japan)* 74 7436 1954; Whistler and House *Anal Chem* 25 1463; UV: Lewis *J Am Chem Soc* 57 898 1935.] It is a substrate for β -D-glucosidase [deBryne *Eur J Biochem* 102 257 1979].

Phenylmercuric acetate (PhHgOAc) [62-38-4] M 336.7, m 148-151°, 149°, 151.8-152.8°. Small colourless lustrous prisms from EtOH. Its solubility in H₂O is 0.17% but it is more soluble in EtOH, Me₂CO and *C₆H₆. [Maynard *J Am Chem Soc* 46 1510 1925; Coleman et al. *J Am Chem Soc* 59 2703 1937; *J Am Pharm Assoc* 25 752 1936.] See PhHgOH and PhHgNO₃.PhHgOH on p. 449 in Chapter 5.

Phenylmethanesulfonyl fluoride (PMSF) [329-98-6] M 174.2, m 90-91°, 92-93°. Purified by recrystn from *C₆H₆, pet ether or CHCl₃-pet ether. [Davies and Dick *J Chem Soc* 483 1932; cf Tullock and Coffman *J Org Chem* 23 2016 1960.] It is a general protease inhibitor (specific for trypsin and chymotrypsin) and is a good substitute for diisopropylphosphoro floridate [Fahrney and Gould *J Am Chem Soc* 85 997 1963].

Phosphatase alkaline (alkaline phosphatase) [9001-78-9] M_r ~40,000 (bovine liver), ~140,000 (bovine intestinal mucosa), 80,000 (*E.coli*) [EC 3.1.3.1]. The *E.coli* supernatant in sucrose (20%, 33mM) in Tris-HCl pH 8.0 was purified through a DEAE-cellulose column and recrystallised. To the column eluates in 0.125M NaCl is added MgCl₂ (to 0.01M) and brought to 50% saturation in (NH₄)₂SO₄ by adding the solid (0.20g/mL). The mixture is centrifuged to remove bubbles and is adjusted to pH 8.0 (with 2N NaOH). Saturated (NH₄)₂SO₄ at pH 8.0 is added dropwise until the soln becomes faintly turbid (~61% saturation). It is set aside at room temp for 1h (turbidity will increase). The mixture is placed in an ice bath for several minutes when turbidity disappears and a clear soln is obtained. It is then placed in a large ice bath at 0° (~5L) and allowed to warm slowly to room temperature in a dark room whereby crystals are formed appearing as a silky sheen. The crystals are collected by centrifugation at 25° if necessary. The crystalline solns are stable at room temperature for many months. They can be stored at 0°, but are not stable when frozen. Cysteine at 10⁻³M and thioglycolic acid at 10⁻⁴M are inhibitory. Inhibition is reversed on addition of Zn²⁺ ions. Many organic phosphates are good substrates for this phosphatase. [Molamy and Horecker *Methods Enzymol* 9 639 1966; Torriani et al. *Methods Enzymol* 12b 212 1968; Engstrom *Biochim Biophys Acta* 92 71 1964.]

Alkaline phosphatase from rat *osteosarcoma* has been purified by acetone pptn, followed by chromatography on DEAE-cellulose, Sephacryl S-200, and hydroxylapatite. [Nair et al. *Arch Biochem Biophys* 254 18 1987.]

3-sn-Phosphatidylethanolamine (L- α -cephalin, from Soya bean) [39382-08-6] M_r ~600-800, amorphous, $pK_{Est(1)}$ ~ 5.8 (PO₄⁻), $pK_{Est(2)}$ ~ 10.5 (NH₂). Purified by dissolving in EtOH, adding Pb(OAc)₂.3H₂O (30g in 100mL H₂O) until excess Pb²⁺ is present. Filter off the solid. Pass CO₂ gas through

the soln until pptn of PbCO_3 ceases. Filter the solid off and evaporate (while bubbling CO_2) under vacuum. An equal volume of H_2O is added to the residual oil extracted with hexane. The hexane extract is washed with H_2O until the aqueous phase is free from Pb [test with dithizone (2 mg in 100 mL CCl_4 ; Feigel *Spot Tests* Vol I, Elsevier p. 10 1954]. The hexane is dried (Na_2SO_4), filtered and evaporated to give a yellow waxy solid which should be dried to constant weight *in vacuo*. It is practically insoluble in H_2O and Me_2CO , but freely soluble in CHCl_3 (5%) and Et_2O , and slightly soluble in EtOH . [Schofield and Dutton *Biochem Prep* 5 5 1957.]

O-Phosphocolamine 2-aminoethyl dihydrogen phosphate [1071-23-4] M 141.1, m 237-240°, 242.3°, 234.5-244.5°, 244-245°(capillary), $\text{pK}_1^{2.0} < 1.5$ (PO_4H_2), $\text{pK}_2^{2.0} 5.77$ (PO_4H^-), $\text{pK}_3^{2.0} 10.26$ (NH^+). Purified by recrystn from aqueous EtOH as a hydrate (m 140-141°). Its solubility in H_2O is 17% and 0.003% in MeOH or EtOH at 22°. [Fölisch and Österberg *J Biol Chem* 234 2298 1959; Baer and Staucer *Can J Chem* 34 434 1956; Christensen *J Biol Chem* 135 399 1940.] It is a potent inhibitor of ornithine decarboxylase [Gilad and Gilad *Biochem Biophys Res Commun* 122 277 1984].

Phosphoenolpyruvic acid monopotassium salt (KPEP) [4265-07-0] M 206.1, $\text{pK}_1^{2.5} 3.4$ (CO_2), $\text{pK}_2^{2.5} 6.35$ (PO_4H^-) (for free acid). It is purified *via* the monocyclohexylamine salt (see next entry). The salt (534mg) in H_2O (10mL) is added to Dowex 50Wx4 H^+ form (200-400 mesh, 2mL, H_2O washed) and stirred gently for 30min and filtered. The resin is washed with H_2O (6mL) and the combined solns are adjusted to pH 7.4 with 3N KOH (~1.4mL) and the volume adjusted to 18.4mL with H_2O to give a soln of 0.1M KPEP which can be lyophilised to a pure powder and is very good for enzyme work. It has been recrystd from $\text{MeOH-Et}_2\text{O}$. [Clark and Kirby *Biochem Prep* 11 103 1966; Wold and Ballou *J Biol Chem* 227 301 1957; Cherbuliez and Rabinowitz *Helv Chim Acta* 39 1461 1956.]

The triNa salt [5541-93-5] M 360.0, is purified as follows: the salt (1g) is dissolved in MeOH (40mL) and dry Et_2O is added in excess. The white crystals are collected and dried over P_2O_5 at 20°. [*Chem Ber* 92 952 1959.]

Phosphoenolpyruvic acid tris(cyclohexylamine) salt [35556-70-8] M 465.6, m 155-180°(dec). Recrystd from aqueous Me_2CO and dried in a vacuum. At 4° it is stable for >2 years and has IR at 1721cm^{-1} ($\text{C}=\text{O}$). [Wold and Ballou *J Biol Chem* 227 301 1957; Clark and Kirby *Biochem Prep* 11 103 1966 for the monocyclohexylamine salt.]

D-3-Phosphoglyceric acid disodium salt (D-glycerate 3-phosphate di-Na salt) [80731-10-8] M 230.0, $[\alpha]_D^{25} +7.7^\circ$ (c 5, H_2O), -735° (in aq NH_4^+ molybdate), $\text{pK}_{\text{Est}(1)} \sim 1.0$ (PO_4H_2), $\text{pK}_{\text{Est}(2)} \sim 6.66$ (PO_4H^-) (for free acid). Best purified by conversion to the Ba salt by pptn with BaCl_2 which is recrystd three times before conversion to the sodium salt. The Ba salt (9.5g) is shaken with 200mL of a 1:1 slurry of Dowex 50 (Na^+ form) for 2h. The mixture is filtered and the resin washed with H_2O (2 x 25mL). The combined filtrates (150mL) are adjusted to pH 7.0 and concentrated *in vacuo* to 30-40mL and filtered if not clear. Absolute EtOH is added to make 100mL and then *n*-hexane is added whereby a white solid and/or a second phase separates. When set aside at room temperature complete pptn of the Na salt as a solid occurs. The salt is removed by centrifugation, washed with Me_2CO , dried in air then in an oven at 55° to give a stable powder (4.5g). It did not lose weight when dried further over P_2O_5 at 78°/8h. The high rotation in the presence of $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$ is not very sensitive to the concentration of molybdate or pH as it did not alter appreciably in 1/3 volume between 2.5 to 25% (w/v) of molybdate or at pH values ranging between 4 and 7. [Cowgill *Biochim Biophys Acta* 16 613 1955; Embdan, Deuticke and Kraft *Hoppe Seyler's Z Physiol Chem* 230 20 1934.]

Phospholipids. For the removal of ionic contaminants from raw zwitterionic phospholipids, most lipids were purified twice by mixed-bed ionic exchange (Amberlite AB-2) of methanolic solutions. (About 1g of lipid in 10mL of MeOH). With both runs the first 1mL of the eluate was discarded. The main fraction of the solution was evaporated at 40°C under dry N_2 and recryst three times from *n*-pentane. The resulting white powder was dried for about 4h at 50° under reduced pressure and stored at 3°. Some samples were purified by mixed-bed ion exchange of aqueous suspensions of the crystal/liquid crystal phase. [Kaatze et al. *J Phys Chem* 89 2565 1985.]

Phosphoproteins (various). Purified by adsorbing onto an iminodiacetic acid substituted agarose column to which was bound ferric ions. This chelate complex acted as a selective immobilised metal affinity adsorbent for phosphoproteins. [Muszyfiska et al. *Biochemistry* **25** 6850 1986.]

5'-Phosphoribosyl pyrophosphate synthetase (from human erythrocytes, or pigeon or chicken liver) [9015-83-2] M_r 60,000, [EC 2.7.6.1]. Purified 5100-fold by elution from DEAE-cellulose, fractionation with ammonium sulfate, filtration on Sepharose 4B and ultrafiltration. [Fox and Kelley *J Biol Chem* **246** 5739 1971; Flaks *Methods Enzymol* **6** 158 1963; Kornberg et al. *J Biol Chem* **15** 389 1955.]

O-Phospho-L-serine [407-41-0] M 185.1, m 175-176°, $[\alpha]_D^{20} +4.3^\circ$ (c 3.2, H₂O), $+16.2^\circ$ (c 3.2, 2N HCl), $pK_1^{2.5} < 1$ (PO₄H₂), $pK_2^{2.5} 2.08$ (CO₂H), $pK_3^{2.5} 5.65$ (PO₄H⁻), $pK_4^{2.5} 9.74$ (NH₃⁺). Recrystd by dissolving 10g in H₂O (150mL) at 25°, stirring for up to 20min. Undissolved material is filtered off (Büchner) and 95% EtOH (85mL) is added dropwise during 4min, and set aside at 25° for 3h then at 3° overnight. The crystals are washed with 95% EtOH (100mL) then dry Et₂O (50mL) and dried in a vacuum (yield 6.5g). A further quantity (1.5mg) can be obtained by keeping the mother liquors and washings at -10° for 1 week. The *DL*-isomer has m 167-170°(dec) after recrystn from H₂O + EtOH or MeOH. [Neuhaus and Korkes *Biochem Prep* **6** 75 1958; Neuhaus and Byrne *J Biol Chem* **234** 113 1959; IR: Fölsch and Mellander *Acta Chem Scand* **11** 1232 1957.]

O-Phospho-L-threonine (L-threonine-O-phosphate) [1114-81-4] M 199.1, m 194°(dec), $[\alpha]_D^{24} -7.37^\circ$ (c 2.8, H₂O) (pK as above). Dissolve in the minimum volume of H₂O, add charcoal, stir for a few min, filter and apply onto a Dowex 50W (H⁺ form) then elute with 2N HCl. Evaporate the eluates under reduced pressure whereby the desired fraction produced crystals of the phosphate which can be recrystd from H₂O-MeOH mixtures and the crystals are then dried *in vacuo* over P₂O₅ at ~80°. [de Verdier *Acta Chem Scand* **7** 196 1953.]

O-Phospho-L-tyrosine (L-tyrosine-O-phosphate) [21820-51-9] M 261.2, m 225°, 227°, 253°, $[\alpha]_D^{20} -5.5^\circ$ (c 1, H₂O), -9.2° (c 1, 2N HCl), $pK_{Est(1)} \sim 1.6$ (PO₄H₂), $pK_{Est(2)} \sim 2.02$ (CO₂H), $pK_{Est(3)} \sim 5.65$ (PO₄H⁻), $pK_{Est(4)} 9.2$ (NH₃⁺). Purified by recrystn from H₂O or H₂O + EtOH. [Levene and Schormüller *J Biol Chem* **100** 583 1933; Posternak and Graff *Helv Chim Acta* **28** 1258 1945.]

Phytol (d-3,7R,11R,15-tetramethylhexadec-2-en-1-ol) [150-86-7] M 296.5, b 145°/0.03mm, 150-151°/0.06mm, 202-204°/10mm, $d_4^{25} 0.8497$, $n_D^{25} 1.437$, $[\alpha]_D^{22} +0.06^\circ$ (neat). Purified by distn under high vacuum. It is almost insoluble in H₂O but soluble in most organic solvents. It has UV λ_{max} at 212nm (log ϵ 3.04) in EtOH and IR ν at 3300 and 1670cm⁻¹. [Demole and Lederer *Bull Soc Chim Fr* 1128 1958; Burrell *J Chem Soc (C)* 2144 1966; Bader *Helv Chim Acta* **34** 1632 1951.]

D-Pipecolic acid (R-piperidine-2-carboxylic acid) [1723-00-8] M 129.2, m 264°(dec), 267°(dec), ~280°(dec), $[\alpha]_D^{29} +26.2^\circ$ (c 2, H₂O), $[\alpha]_D^{25} +35.7^\circ$ (H₂O), $pK_1^{2.0} 2.29$ (CO₂H), $pK_2^{2.0} 10.77$ (NH⁺). Recrystallises as platelets from EtOH and is soluble in H₂O. The *hydrochloride* has m 256-257°(dec) from H₂O and $[\alpha]_D^{25} +10.8^\circ$ (c 2, H₂O). [Lukés et al. *Collect Czech Chem Commun* **22** 286 1957; Bayerman *Recl Trav Chim Pays-Bas* **78** 134 1959; Asher et al. *Tetrahedron Lett* **22** 141 1981.]

L-Pipecolic acid (S-piperidine-2-carboxylic acid) [3105-95-1] M 129.2, m 268°(dec), 271°(dec), ~280°(dec), $[\alpha]_D^{20} -26^\circ$ (c 4, H₂O), $[\alpha]_D^{25} -34.9^\circ$ (H₂O). Recrystd from aqueous EtOH and sublimes as needles in a vacuum. It is sparingly soluble in absolute EtOH, Me₂CO and CHCl₃ but insoluble in Et₂O. The *hydrochloride* has m 258-259°(dec, from MeOH) and $[\alpha]_D^{25} -10.8^\circ$ (c 10, H₂O). [Fujii and Myoshi *Bull Chem Soc Jpn* **48** 1241 1975.]

Piperidine-4-carboxylic acid (isonipecotic acid) [498-94-2] M 129.2, m 336°(dec, darkens at ~300°), $pK_{Est(1)} \sim 4.3$ (CO₂H), $pK_{Est(2)} \sim 10.6$ (NH⁺). Recrystallises from H₂O or EtOH as needles. The *hydrochloride* recrystallises from H₂O or aqueous HCl and has m 293°dec (298°dec, 300°dec). [Wibaut *Recl Trav Chim Pays-Bas* **63** 141 1944; IR: Zacharius et al. *J Am Chem Soc* **76** 2908 1954.]

Pituitary Growth Factor (from human pituitary gland) [336096-71-0]. Purified by heparin and copper affinity chromatography, followed by chromatography on carboxymethyl cellulose (Whatman 52). [Rowe et al. *Biochemistry* **25** 6421 1986.]

Plasmids. These are circular lengths of DNA which invade bacteria or other cells e.g. insect cells, yeast cells, and have sequences which are necessary for their replication using enzymes and other ingredients, e.g. nucleotides, present in the cells. They contain engineered, or already have, genes which produce enzymes that provide the cells with specific antibiotic resistance and are thus useful for selecting bacteria containing specific plasmids. Plasmids have been extremely useful in molecular biology since they can be very easily identified (from their size or the sizes of the DNA fragments derived from their restriction enzyme digests) and can be readily engineered *in vitro* (outside the cells). Genes coding for specific enzymes or other functional proteins can be inserted into these plasmids which have DNA sequences that allow the expression of large quantities of bacteria or non-bacterial (e.g. human) proteins. They have also been engineered in such a way as to produce 'fusion proteins' (in which the desired protein is fused with a specific "reporter, marker or carrier protein" which will facilitate the isolation of the desired protein (e.g. by binding strongly to a nickel support) and then the desired protein can be cleaved from the eluted fusion protein and obtained in very pure form. A large number of plasmids with a variety of sequences for specific purposes are commercially available in very pure form. They can be used to infect cells and can be isolated and purified from cell extracts in large amounts using a number of available procedures. These procedures generally involve lysis of the cells (e.g. with alkaline sodium dodecylsulfate, SDS), separation from nuclear DNA, precipitation of plasmid DNA from the cell debris, adsorbing it on columns which specifically bind DNA, and then eluting the DNA from the column (e.g. with specific Tris buffers as recommended by the suppliers) and precipitating it (e.g. with Tris buffer in 70% EtOH at -70°C). The purity is checked in agarose gel (containing ethidium bromide to visualise the DNA) by electrophoresis. A large number of plasmids are now commercially available (see Clontech GmbH, <http://www.clontech.com>; Invitrogen <http://www.invitrogen.com>, among other suppliers) used as vectors for bacterial, mammalian, yeast and baculovirus expression.

Podophylotoxin [518-28-5] **M** 414.4, **m** 181-181°, 183-184°, 188-189°, $[\alpha]_D^{20}$ -132° (c 1, CHCl₃). Recrystallises from *C₆H₆ (with 0.5C₆H₆), EtOH-*C₆H₆, aqueous EtOH (with 1-1.5H₂O, **m** 114-115°) and CH₂Cl₂-pentane. When dried at 100°/10 mm it has **m** 183-184°. [UV: Stoll et al. *Helv Chim Acta* **37** 1747 1954; IR: Schecler et al. *J Org Chem* **21** 288 1956.] Inhibitor of microtubule assembly [Prasad et al. *Biochemistry* **25** 739 1986].

Polyethylene glycol [25322-68-3] **M_r** various, from ~200 to ~35,000. May be contaminated with aldehydes and peroxides. Methods are available for removing interfering species. [Ray and Purathingal *Anal Biochem* **146** 307 1985.]

Polypeptides. These are a string of α-amino acids usually with the natural *S*(L) [L-cysteine is an exception and has the *R* absolute configuration] or sometimes "unnatural" *R*(D) configuration at the α-carbon atom. They generally have less than ~100 amino acid residues. They can be naturally occurring or, because of their small size, can be synthesised chemically from the desired amino acids. Their properties can be very similar to those of small proteins. Many are commercially available, can be custom made commercially or locally with a peptide synthesiser. They are purified by HPLC and can be used without further purification. Their purity can be checked as described under proteins.

Porphobilinogen (5-amino-4-carboxymethyl-1H-pyrrole-3-propionic acid) [487-90-1] **M** 226.2, **m** 172-175°(dec), 175-180°(dec, darkening at 120-130°), **pK₁** 3.70 (4-CH₂CO₂H), **pK₂** 4.95 (3-CH₂CH₂CO₂H), **pK₃** 10.1 (NH⁺). Recrystallises as the monohydrate (pink crystals) from dil NH₄ OAc solns of pH 4, and is dried *in vacuo*. The *hydrochloride monohydrate* has **m** 165-170°(dec) (from dilute HCl). [Jackson and MacDonald *Can J Chem* **35** 715 1957, Westall *Nature* **170** 614 1952; Bogarad *J Am Chem Soc* **75** 3610 1953.]

Porphyrin a (from ox heart) [5162-02-1] **M** 799.0, **m** dec on heating. Purified on a cellulose powder column followed by extraction with 17% HCl and fractionation with HCl. [Morell et al. *Biochem J* **78**

793 1961.] Recrystd from CHCl_3 /Pet ether or $\text{Et}_2\text{O}^*/\text{C}_6\text{H}_6$ [detailed UV-VIS and NMR date: Caughey et al. *J Biol Chem* **250** 7602 1975; Lemberg *Adv Enzymol* **23** 265 1961].

Prazosin hydrochloride (2[4-((2-furoyl)piperazin-1-yl)4-amino-6,7-dimethoxyquinazoline hydrochloride) [19237-84-4] **M 419.9, m 278-280°, 280-282°, pK 6.5.** It is recrystd by dissolving in hot MeOH adding a small volume of MeOH-HCl (dry MeOH saturated with dry HCl gas) followed by dry Et_2O until crystn is complete. Dry *in vacuo* over solid KOH till odour of HCl is absent. It has been recrystd from hot H_2O , the crystals were washed with H_2O , and the H_2O was removed azeotropically with CH_2Cl_2 , and dried in a vacuum. [NMR and IR: Honkanen et al. *J Heterocycl Chem* **17** 797 1980; cf Armarego and Reece *Aust J Chem* **34** 1561 1981.] It is an antihypertensive drug and is an α_1 -adrenergic antagonist [Brosman et al. *Proc Natl Acad Sci USA* **82** 5915 1985].

Prednisolone acetate (21-acetoxypregna-1,4-diene-11 β -17 α -diol-3,20-dione) [52-21-1] **M 402.5, m 237-239°, 240-242°, 240-243°, 244°, $[\alpha]_D^{20} +116^\circ$ (c 1, dioxane).** Recrystd from EtOH, Me_2CO , Me_2CO -hexane, and has UV λ_{max} at 243nm in EtOH. [Joly et al. *Bull Soc Chim Fr* 366 1958; Herzog et al. *J Am Chem Soc* **77** 4781 1955.]

Primaquine diphosphate (RS- 8-[4-amino-1-methylbutylamino]-6-methoxyquinoline diphosphate) [63-45-6] **M 455.4, m 197-198°, 204-206°, pK_{Est(1)}~ 3.38 (ring N⁺), pK_{Est(2)}~ 10.8 (NH₃⁺).** It forms yellow crystals from 90% aq EtOH and is moderately soluble in H_2O . The *oxalate salt* has m 182.5-185° (from 80% aq EtOH) and the *free base* is a viscous liquid b 165-170°/0.002mm, 175-177°/2mm. [Elderfield et al. *J Am Chem Soc* **68** 1526 1964; **77** 4817 1955.]

Procaine hydrochloride (Novocain, 2-diethylaminoethyl-4-aminobenzoate) [51-05-8] **M 272.8, m 153-156°, 154-156°, 156°, pK_{Est(1)}~ 2.52 (NH₂⁺) pK₂²⁰ 9.0 (Et₂N⁺).** Recrystd from aqueous EtOH. It has solubility at 25° in H_2O (86.3%), EtOH (2.6%) and Me_2CO (1%), it is slightly soluble in CHCl_3 but is almost insoluble in Et_2O . The anhydrous *free base* is recrystd from ligroin or Et_2O and has m 61°. [Einhorn *Justus Liebigs Ann Chem* **371** 125 1909; IR: Szymanski and Panzica *J Amer Pharm Assoc* **47** 443 1958.]

L-Propargylglycine (S-2-aminopent-4-ynoic acid) [23235-01-0] **M 113.1, m 230°(dec starting at 210°), $[\alpha]_D^{20} -35^\circ$ (c 1, H_2O), -4° (c 5, 5N HCl), pK_{Est(1)}~ 2.3 (CO₂H), pK_{Est(2)}~ 9.8 (NH₂).** The acid crystallises readily when ~4g in 50mL H_2O is treated with abs EtOH at 4°/ 3hrs, and is collected washed with cold abs EtOH and Et_2O and dried in vac. Also recrystallises from aqueous Me_2CO , R_F on SiO_2 TLC plates with *n*-BuOH- H_2O -AcOH (4:1:1) is 0.26. The *racemate* has m 238-240°. [Leukart et al. *Helv Chim Acta* **59** 2181 1976; Eberle and Zeller *Helv Chim Acta* **68** 1880 1985; Jansen et al. *Recl Trav Chim Pays-Bas* **88** 819 1969.] It is a suicide inhibitor of γ -cystathionase and other enzymes [Washtier and Abeles *Biochemistry* **16** 2485 1977; Shinozuka et al. *Eur J Biochem* **124** 377 1982].

Propidium iodide (3,8-diamino-5-(3-diethylaminopropyl)-6-phenylphenanthridinium iodide methiodide) [25535-16-4] **M 668.4, m 210-230°(dec), pK_{Est(1)}~ 4 (aniline NH₂), pK_{Est(2)}~ 8.5 (EtN₂).** Recrystd as red crystals from H_2O containing a little KI. It fluoresces strongly with nucleic acids. [Eatkins *J Chem Soc* 3059 1952.] **TOXIC.**

R-Propranalol hydrochloride (R-1-isopropylamino-3-(1-naphthyloxy)-2-propanol HCl) [13071-11-9] **M 295.8, m 192°, 193-195°, $[\alpha]_D^{20} -25^\circ$ (c 1, EtOH), pK²⁰ 9.5 (for free base).** Recryst from *n*-PrOH or Me_2CO . It is soluble in H_2O and EtOH but is insoluble in Et_2O , $^*\text{C}_6\text{H}_6$ or EtOAc. The *racemate* has m 163-164°, and the *free base* recryst from cyclohexane has m 96°. [Howe and Shanks *Nature* **210** 1336 1966.] The *S*-isomer (below) is the physiologically active isomer.

S-Propranalol hydrochloride (S-1-isopropylamino-3-(1-naphthyloxy)-2-propanol HCl) [4199-10-4] **M 295.8, m 192°, 193-195°, $[\alpha]_D^{20} +25^\circ$ (c 1, EtOH) pK²⁰ 9.5.** See preceding entry for physical properties. The is the active isomer which blocks isoprenaline tachycardia and is a β -adrenergic blocker. [Leclerc et al. *Trends Pharmacol Sci* **2** 18 1981; Howe and Shanks *Nature* **210** 1336 1966.]

Protamine kinase (from rainbow trout testes) [37278-10-7] [EC 2.7.1.70]. Partial purification by hydroxylapatite chromatography followed by biospecific chromatography on nucleotide coupled Sepharose 4B (the nucleotide was 8-(6-aminohexyl)amine coupled cyclic-AMP). [Jergil et al. *Biochem J* **139** 441 1974.]

Protamine sulfate (from herring sperm) [9007-31-2] [α]_D²² -85.5° (satd H₂O), pK 7.4-8.0. A strongly basic protein (white powder, see pK) used to ppt nucleic acids from crude protein extracts. It dissolved to the extent of 1.25% in H₂O. It is freely soluble in hot H₂O but separates as an oil on cooling. It has been purified by chromatography on an IRA-400 ion-exchange resin in the SO₄²⁻ form and washed with dilute H₂SO₄. Eluates are freeze-dried under high vacuum below 20°. This method is used to convert proteamine and protamine hydrochloride to the sulfate. [UV: Rasmussen *Hoppe Seyler's Z Physiol Chem* **224** 97 1934; Ando and Sawada *J Biochem (Tokyo)* **49** 252 1961; Felix and Hashimoto *Hoppe Seyler's Z Physiol Chem* **330** 205 1963]

Protease nexin (From cultured human fibroblasts) [148263-58-5]. Purified by affinity binding of protease nexin to dextran sulfate-Sepharose. [Farrell et al. *Biochem J* **237** 707 1986.]

Proteins. These are usually naturally occurring (or deliberately synthesised in microorganisms, e.g. bacteria, insect cells, or animal tissues), and are composed of a large number of α -S (L) amino acids residues (except for L-cysteine which has the R absolute configuration), selected from the 20 or so natural amino acids, in specific sequences and in which the α -amino group forms an amide (peptide) bond with the α -carboxyl group of the neighboring amino acid. The number of residues are usually upwards of 100. Proteins with less than 100 amino acids are better referred to as **polypeptides**. Aqueous soluble proteins generally fold into ball-like structures mainly with hydrophilic residues on the outside of the "balls" and hydrophobic residues on the inside. Proteins can exist singly or can form dimers, trimers, tetramers etc, consisting of similar or different protein subunits. They are produced by cells for a large variety of functions, e.g. enzymology, reaction mediation as in regulation of DNA synthesis or chaperonins for aiding protein folding, formation of pores in membranes for transport of ions or organic molecules, or for intra or inter cellular signalling etc. The purity of proteins can be checked in denaturing (SDS, sodium dodecylsulfate) or non-denaturing polyacrylamide gels using electrophoresis (PAGE), and staining appropriately (e.g. with Coomassie Blue, followed by silver staining for higher sensitivity). If the protein is partly impure then it should be purified further according to the specific literature procedures for the individual protein (see specific proteins in the *Methods Enzymol*, Wiley series).

Proteoglycans (from cultured human muscle cells). Separated by ion-exchange HPLC using a Bio-gel TSK-DEAE 5-PW analytical column. [Harper et al. *Anal Biochem* **159** 150 1986.]

Prothrombin (Factor II, from equine blood plasma) [9001-26-7] M_r 72,000. Purified by two absorptions on a barium citrate adsorbent, followed by decomposition of the adsorbents with a weak carboxylic cation-exchanger (Amberlite IRF-97), isoelectric pptn (pH 4.7-4.9) and further purification by chromatography on Sephadex G-200 or IRC-50. Finally recrystd from a 1% soln adjusted to pH 6.0-7.0 and partial lyophilisation to ca 1/5 to 1/10th vol and set aside at 2-5° to crystallise. Occasionally seeding is required. [Miller *Biochem Prep* **13** 49 1971.]

Protoporphyrin IX (3,18-divinyl-2,7,13,17-tetramethylporphine-8,12-dipropionic acid, ooporphyrin) [553-12-8] M 562.7, pK_{Est} ~ 4.8. Purified by dissolving (4g) in 98-100% HCOOH (85mL), diluting with dry Et₂O (700mL) and keeping at 0° overnight. The ppt is collected and washed with Et₂O then H₂O and dried in a vacuum at 50° over P₂O₅. It has been recrystd from aqueous pyridine and from Et₂O as monoclinic, brownish-yellow prisms. UV λ_{\max} values in 25% HCl are 557.2, 582.2 and 602.4nm. It is freely soluble in ethanolic HCl, AcOH, CHCl₃, and Et₂O containing AcOH. It forms sparingly soluble diNa and diK salts. [Ramsey *Biochem Prep* **3** 39 1953; UV: Holden *Aust J. Exptl Biol and Med Sci* **15** 412 1937; Garnick *J Biol Chem* **175** 333 1948; IR: Falk and Willis *Aust J Sci Res [A]* **4** 579 1951.]

The **Dimethyl ester** [5522-66-7] M 590.7, m 228-230°, is prepared by dissolving (0.4g) in CHCl₃ (33mL) by boiling for a few min, then diluting with boiling MeOH (100mL) and refrigerating for 2 days. The crystals are collected, washed with CHCl₃-MeOH (1:9) and dried at 50° in a vacuum (yield 0.3g). UV has λ_{\max} 631, 576, 541, 506 and 407nm in CHCl₃ and 601, 556 and 406nm in 25% HCl. [Ramsey *Biochem Prep* **3** 39 1953.]

Prymnesin (toxic protein from phytoflagellate *Prymnesium parvum*) [11025-94-8]. Purified by column chromatography, differential soln and pptn in solvent mixtures and differential partition between diphasic mixtures. The product has at least 6 components as observed by TLC. [Ulitzur and Shilo *Biochim Biophys Acta* 301 350 1970.]

Pterin-6-carboxylic acid (2-amino-4-oxo-3,4-dihydropteridine-6-carboxylic acid) [948-60-7] **M** 207.2, **m** >360°, **pK₁^{2.0}** 1.43, **pK₂^{2.0}** 2.88, **pK₃^{2.0}** 7.72. Yellow crystals by repeated dissolution in aqueous NaOH and adding aqueous HCl. It has UV with λ_{\max} at 235, 260 and 265nm (ϵ 11000, 10500 and 9000) in 0.1N HCl and 263 and 365nm (ϵ 20500 and 9000) in 0.1N NaOH. [UV: Pfeleiderer et al. *Justus Liebigs Ann Chem* 741 64 1970; Stockstad et al. *J Am Chem Soc* 70 5 1948; Fluorescence: Kavanagh and Goodwin *Arch Biochem* 20 315 1949.]

Purine-9- β -ribifuranoside (Nebularin) [550-33-4] **M** 252.2, **m** 178-180°, 181-182°, [α]_D²⁵ -48.6° (c 1, H₂O), -22° (c 0.8, 0.1N HCl) and -61° (c 0.8, 0.1N NaOH), **pK** 2.05. Recrystd from butanone + MeOH or EtOH and forms a MeOH photo-adduct. It is a strong inhibitor of adenosine deaminase [EC 3.5.4.4]. [Nair and Weichert *Bioorg Chem* 9 423 1980; Löfgren et al. *Acta Chem Scand* 7 225 1953; UV: Brown and Weliky *J Biol Chem* 204 1019 1953.]

Puromycin dihydrochloride (*O*-methyl-*l*-tyrosine[*N*⁶,*N*⁶-dimethylaminoadenosin-3'-ylamide]) [58-58-2] **M** 616.5, **m** 174°, [α]_D²⁵ -11° (free base in EtOH), **pK₁** 6.8, **pK₂** 7.2. Purified by recrystn from H₂O. The *free base* has **m** 175.5-177° (172-173°) (from H₂O). The *sulfate* has **m** 180-187° dec (from H₂O), and the *picrate monohydrate* has **m** 146-149° (from H₂O). [Baker et al. *J Am Chem Soc* 77 1 1955; Fryth et al. *J Am Chem Soc* 80 3736 1958.] It is an inhibitor of aminopeptidase and terminates protein synthesis [Reboud et al. *Biochemistry* 20 5281 1981].

Pyridoxal hydrochloride [65-22-5] **M** 203.6, **m** 176-180°(dec), **pK₁^{2.0}** 4.23 (3-OH), **pK₂^{2.0}** 8.7 (Pyridinium⁺), **pK₃^{2.0}** 13.04 (CH₂OH?). Dissolve in water and adjust the pH to 6 with NaOH. Set aside overnight to crystallise. The crystals are washed with cold water, dried in a vacuum desiccator over P₂O₅ and stored in a brown bottle at room temperature. [Fleck and Alberty *J Phys Chem* 66 1678 1962.]

Pyridoxal-5'-phosphate monohydrate (PLP, codocarboxylase) [54-47-7] **M** 265.2, **pK₁^{2.5}** <2.5 (PO₄), **pK₂^{2.5}** 4.14 (3-OH), **pK₃^{2.5}** 6.20 (PO₄), **pK₄^{2.5}** 8.69 (pyridinium⁺). It has been purified by dissolving 2g in H₂O (10-15mL, in a dialysis bag a third full) and dialysing with gentle stirring against 1L of H₂O (+ two drops of toluene) for 15h in a cold room. The dialysate is evaporated to 80-100mL then lyophilised. Lemon yellow microscopic needles of the monohydrate remain when all the ice crystals have been removed. The purity is checked by paper chromatography (in EtOH or *n*-PrOH-NH₃) and the spot(s) visualised under UV light after reaction with *p*-phenylene diamine, NH₃ and molybdate. Solutions stored in a freezer are 2-3% hydrolysed in 3-weeks. At 25°, only 4-6% hydrolysis occurs even in N NaOH or HCl, and 2% is hydrolysed at 37° in 1 day - but is complete at 100° in 4h. Best stored as dry solid at -20°. In aqueous acid the solution is colourless but is yellow in alkaline solutions. It has UV λ_{\max} at 305nm (ϵ 1100) and 380nm (ϵ 6550) in 0.1 N NaOH; 330nm (ϵ 2450) and 388nm (ϵ 4900) in 0.05M phosphate buffer pH 7.0 and 295nm (ϵ 6700) in 0.1N HCl. [Peterson et al. *Biochemical Preparations* 3 34, 119 1953.] The *oxime* dec at 229-230° and is practically insoluble in H₂O, EtOH and Et₂O. The *O*-methyl*oxime* decomposes at 212-213°. [Heyl et al *J Am Chem Soc* 73 3430 1951.] It has also been purified by column chromatography through Amberlite IRC-50 (H⁺) [Peterson and Sober *J Am Chem Soc* 76 169 1954].

Pyridoxamine hydrochloride [5103-96-8, 524-36-7 (*free base*)] **M** 241.2, **m** 226-227°(dec), **pK₁^{2.5}** 3.54 (3-OH), **pK₂^{2.5}** 8.21 (ring N⁺), **pK₃^{2.5}** 10.63 (NH₂). Crystd from hot MeOH. The *free base* crystals from EtOH, has **m** 193-193.5° [Harris et al. *J Biol Chem* 154 315 1944, *J Am Chem Soc* 66 2088 1944].

Pyridoxine hydrochloride see Vitamin B₆.

Pyruvate kinase isoenzymes (from *Salmonella typhimurium*) [9001-59-6] **M_r** 64,000, [EC 2.7.1.40], **amorphous**. Purified by (NH₄)₂SO₄ fractionation and gel filtration, ion-exchange and affinity chromatography. [Garcia-Olalla and Garrido-Pertierra *Biochem J* 241 573 1987.]

Quinacrine [Atebrine, 3-chloro-9(4-diethylamino-1-methyl)butylamino-7-methoxy)acridine] dihydrochloride. [69-05-6] **M** 472.9, **m** 248-250°(dec), pK_1^{30} -6.49 (aq H_2SO_4), pK_2^{30} 7.73 (ring NH^+), pK_3^{30} 10.18 (Et_2N). Cryst from H_2O (sol 2.8% at room temp) as yellow crystals. Slightly sol in MeOH and EtOH. Antimalarial, antiprotozoal and intercalates DNA. [Wolfe *Antibiot* 3 (Springer-Verlag) 203 1975.]

Quisqualic acid (3-[3,5-dioxo-1,2,4-oxadiazolin-2-yl]-L-alanine) [52809-07-1] **M** 189.1, **m** 190-191°, $[\alpha]_{\text{D}}^{20} +17^\circ$ (c 2, 6M HCl), $\text{pK}_{\text{Est}(1)} \sim 2.1$ (CO_2H), $\text{pK}_{\text{Est}(2)} \sim 8.9$ (NH_2). It has been purified by ion-exchange chromatography on Dowex 50W (x 8, H^+ form), the desired fractions are lyophilised and recrystd from H_2O -EtOH. It has IR (KBr) ν : 3400-2750br, 1830s, 1775s, 1745s and 1605s cm^{-1} ; and ^1H NMR ($\text{NaOD}/\text{D}_2\text{O}$, pH 13) δ : 3.55-3.57 (1H m, X of ABX, H-2), 3.72-3.85 (2H, AB of ABX, H-3), ^{13}C NMR (D_2O) δ : 50.1t, 53.4d, 154.8s, 159.7s and 171.3s. [Baldwin et al. *J Chem Soc, Chem Commun* 256 1985.] It is a quisqualate receptor agonist [Joels et al. *Proc Natl Acad Sci USA* 86 3404 1989].

Renal dipeptidase (from porcine kidney cortex) [9031-96-3] **M_r** 47,000 [EC 3.4.13.11]. Purified by homogenising the tissue, extracting with Triton X-100, elimination of insoluble material, and ion-exchange, size exclusion and affinity chromatography. [Hitchcock et al. *Anal Biochem* 163 219 1987.]

Restriction enzymes (endonucleases). These are enzymes which cleave double stranded DNA (linear or circular) at specific nucleotide sequences within the DNA strands which are then used for cloning (by ligating bits of DNA sequences together) or used for identifying particular DNA materials, e.g. plasmids, genes etc. A very large number of restriction enzymes are now available commercially and are extensively used in molecular biology. They are highly specific for particular nucleotide arrangements and are sensitive to the reaction conditions, e.g. composition of the medium, pH, salt concentration, temperature etc, which have to be strictly adhered to. The enzymes do not require further purification and the reaction conditions are also provided by the suppliers from which the necessary reaction media can also be purchased (see commercial catalogues).

Retinal (Vitamin A aldehyde), Retinoic acid (Vitamin A acid), Retinyl acetate, Retinyl palmitate see entries in Chapter 4.

Reverse transcriptase (from avian or murine RNA tumour viruses) [9068-38-6] [EC 2.7.7.49]. Purified by solubilising the virus with non-ionic detergent. Lysed virions were adsorbed on DEAE-cellulose or DEAE-Sephadex columns and the enzyme eluted with a salt gradient, then chromatographed on a phosphocellulose column and enzyme activity eluted in a salt gradient. Purified from other viral proteins by affinity chromatography on a pyran-Sepharose column. [Verna *Biochim Biophys Acta* 473 1 1977; Smith *Methods Enzymol* 65 560 1980; see commercial catalogues for other transcriptases.]

Riboflavin [83-88-5] **M** 376.4, **m** 295-300°(dec), $[\alpha]_{\text{D}} -9.8^\circ$ (H_2O), -125° (c 5, 0.05N NaOH), pK_1 1.7, pK_2 9.69 (10.2, acidic NH). Crystd from 2M acetic acid, then extracted with CHCl_3 to remove lumichrome impurity. [Smith and Metzler *J Am Chem Soc* 85 3285 1963.] Has also been crystd from water. (See also p. 575.)

Riboflavin-5'-phosphate (Na salt, 2H₂O) [130-40-5] **M** 514.4. See flavin mononucleotide (FMN) on p. 535.

D-(+)-Ribonic acid- γ -lactone [5336-08-3] **M** 148.12, **m** 80°, 84-86°, $[\alpha]_{\text{D}}^{20} +18.3^\circ$ (c 5, H_2O). Purified by recrystn from EtOAc. The *tribenzoate* has **m** 54-56° (from AcOH), $[\alpha]_{\text{D}}^{25} +27^\circ$ (c 2.37, Me_2NCHO) and the 3,5-O-*benzylidene* derivative has **m** 230-231.5° (needles from Me_2CO -pet ether), $[\alpha]_{\text{D}}^{25} -177^\circ$ (CHCl_3). [Chen and Joulié *J Org Chem* 49 2168 1984; Zinner and Voigt *J Carbohydr Res* 7 38 1968.]

Ribonuclease (from human plasma) [9001-99-4] M_r ~13,700, [EC 3.1.27.5], **amorphous**. Purified by $(\text{NH}_4)_2\text{SO}_4$ fractionation, followed by PC cellulose chromatography and affinity chromatography (using Sepharose 4B to which $(\text{G})_n$ was covalently bonded). [Schmukler et al. *J Biol Chem* **250** 2206 1975.]

RNA (ribonucleic acids). Ribonucleic acids are like DNA except that the 2'-deoxy-D-ribose moiety is replaced by a D-ribose moiety and the fourth nucleotide thymidylic acid (T) is replaced by uridylic acid (U). RNA does not generally form complete duplex molecules like DNA, i.e. it is generally monomeric, except in certain viruses. The two main classes of RNA are **messenger-RNA** (mRNA) and **transfer-RNA** (tRNA). mRNA transcribed from the DNA gene followed by the splicing out of the non-coding nucleotides (of the introns) and codes for a specific gene. There are many different tRNAs, at least one of which links to a specific α -amino acid, that bind to the mRNA *via* the ribosome (a set of proteins) to the RNA triplets (three nucleotides) which code for the particular α -amino acids. An enzyme then joins the α -amino acids of two adjacent tRNA- α -amino acid ribosome complexes bound to the mRNA to form a peptide bond. Thus peptide bonds and consequently polypeptides and proteins coded by the DNA *via* the respective mRNA are produced.

Martin et al. [*Biochem J* **89** 327 1963] dissolved RNA (5g) in 90mL of 0.1mM EDTA, then homogenised with 90mL of 90% (w/v) phenol in water using a Teflon pestle. The suspension was stirred vigorously for 1h at room temperature, then centrifuged for 1h at 0° at 25000rpm. The lower (phenol) layer was extracted four times with 0.1mM EDTA and the aqueous layers were combined, then made 2% (w/v) with respect to AcOK and 70% (v/v) with respect to EtOH. After standing overnight at -20° , the ppte was centrifuged down, dissolved in 50mL of 0.1mM EDTA, made 0.3M in NaCl and left 3 days at 0° . The purified RNA was then centrifuged down at 10000xg for 30min, dissolved in 100mL of 0.1mM EDTA, dialysed at 4° against water, and freeze-dried. It was stored at -20° in a desiccator. Michelson [*J Chem Soc* 1371 1959] dissolved 10g of RNA in water, added 2M ammonia to adjust the pH to 7, then dialysed in Visking tubing against five volumes of water for 24h. The process was repeated three times, then the material after dialysis was treated with 2M HCl and EtOH to ppte the RNA which was collected, washed with EtOH, ether and dried [see commercial catalogues for further examples].

Ricin (toxin from Castor bean *Ricinus communis*) [A chain 96638-28-7; B chain 96638-29-8] M_r ~60,000, **amorphous**. Crude ricin, obtained by aqueous extraction and $(\text{NH}_4)_2\text{SO}_4$ pptn, was chromatographed on a galactosyl-Sepharose column with sequential elution of pure ricin. The second peak was due to ricin agglutinin. [Simmons and Russell *Anal Biochem* **146** 206 1985.] Inhibitor of protein synthesis. **EXTREMELY DANGEROUS, USE EXTREME CARE [instructions accompany product].**

Rifampicin (Rifampin) [13292-46-1] M 823.0, m 183-185 $^\circ$, pK_1 1.7, pK_2 7.9. This macrolide antibiotic crystallises from Me_2CO in red-orange plates. It has UV λ_{max} 237, 255, 334, and 475nm (ϵ 33,200, 32,100, 27,000 and 15,400) at pH 7.38. It is stable in Me_2SO and H_2O . Freely soluble in most organic solvents and slightly soluble in H_2O at pH <6. [Binda et al. *Arzneim.-Forsch* **21** 1907 1971.] It inhibits cellular RNA synthesis without affecting DNA [Calvori et al. *Nature* **207** 417 1965].

Rifamycin B [13929-35-6] M 755.8, m 300 $^\circ$ (darkening at 160-164 $^\circ$), $[\alpha]_D^{20}$ -11 $^\circ$ (MeOH), pK_1 2.60, pK_2 7.76. It forms yellow needles from *C_6H_6 . It has solubility in H_2O (0.027%), MeOH (2.62%) and EtOH (0.44%). It has UV λ_{max} 223, 304 and 245nm ($A_{1\text{cm}}^{1\%}$ 555, 275 and 220). [Oppolzer and Prelog *Helv Chim Acta* **56** 2287 1973; Oppolzer et al. *Experientia* **20** 336 1964; X-ray: Brufani et al. *Experientia* **20** 339 1964.]

Rifamycin SV sodium salt [15105-92-7] M 719.8, m 300 $^\circ$ (darkening >140 $^\circ$), $[\alpha]_D^{20}$ -4 $^\circ$ (MeOH), pK_{Est} ~ 7.8. Yellow orange crystals from Et_2O -pet ether or aq EtOH, very soluble in MeOH, EtOH, Me_2CO and EtOAc, soluble in Et_2O and HCO_3^- , slightly soluble in H_2O and pet ether. Its UV has λ_{max} at 223, 314 and 445nm ($A_{1\text{cm}}^{1\%}$ 586, 322 and 204) in phosphate buffer pH 7. [NMR: Bergamini and Fowst *Arzneim.-Forsch* **15** 951 1965.]

[Previous Page](#)

Saccharides. Resolved by anion-exchange chromatography. [Walberg and Kando *Anal Biochem* **37** 320 1970.]

Sarcosine anhydride [5076-82-4] **M 142.2, m 146-147°**, $\text{pK}_{\text{Est}(1)} \sim 4.2$, $\text{pK}_{\text{Est}(2)} \sim 1.9$. Crystd from water, EtOH or ethyl acetate. Dried in vacuum at room temperature.

(-)-Scopolamine hydrobromide $3\text{H}_2\text{O}$ ($6\beta,7\beta$ -epoxy- 3α -tropanyl *S*(-)-tropate HBr, **hyoscine HBr**) [114-49-8] **M 438.3, m 193-194°**, **195°**, **195-199°**, $[\alpha]_{\text{D}}^{25} -25^\circ$ (c 5, H_2O), pK^{20} **8.15**. Recrystd from Me_2CO , H_2O or EtOH-Et₂O and dried. Soluble in H_2O (60%) and EtOH (5%) but insol in Et₂O and slightly in CHCl_3 . The *hydrochloride* has **m 300°** (from Me_2CO). The *free base* is a viscous liquid which forms a crystalline *hydrate* with **m 59°** and $[\alpha]_{\text{D}}^{20} -28^\circ$ (c 2.7, H_2O). Readily hydrolysed in dilute acid or base. [Meinwald *J Chem Soc* 712 1953; Fodor *Tetrahedron* **1** 86 1957.]

Seleno-DL-methionine (± 2 -amino-4-methylselenanylbutyric acid) [2578-28-1] **M 196.1, m 265°(dec)**, **267-269°(dec)**, **270°** (see **pKs of methionine**). Crystallises in hexagonal plates from MeOH and H_2O . [Klosterman and Painter *J Am Chem Soc* **69** 2009 1949.] The L-isomer is purified by dissolving in H_2O , adjusting the pH to 5.5 with aqueous NH_3 , evaporating to near-dryness, and the residue is washed several times with absolute EtOH till solid is formed and then recrystd from Me_2CO . It has **m 266-268°(dec)**, **275°(dec)**, $[\alpha]_{\text{D}}^{25} +18.1^\circ$ (c 1, N HCl). [Pande et al. *J Org Chem* **35** 1440 1970.]

Serotonin hydrochloride (5-HT, 3-[2-aminoethyl]-5-hydroxyindole HCl) [153-98-0] **M 212.7, m 167-168°**, **178-180°**, pK_1^{25} **4.9**, pK_2^{25} **9.8** (10.0, NH_2), pK_3^{25} **11.1** (5-OH), pK_4^{25} **18.25** (acidic indole NH). Purified by recrystn from EtOH-Et₂O or Et₂O to give the *hygroscopic* salt. Store in the dark as it is light sensitive. The *free base* has **m 84-86°** (from Et₂O). The 5-benzyloxy derivative has **m 84-86°** (from Et₂O). [Ek and Witkop *J Am Chem Soc* **76** 5579 1954; HamLin and Fischer *J Am Chem Soc* **73** 5007 1951.] The *picrate* $1\text{H}_2\text{O}$ has **m 196-197.5°** (dec with sintering at 160-165°) after recrystn from Et₂O. Serotonin is a natural neurotransmitter [Chuang *Life Sci* **41** 1051 1987].

Sinigrin monohydrate (Myronate K) [64550-88-5] **M 415.5, m 125-127°**, **127-129°**, $[\alpha]_{\text{D}}^{20} -17^\circ$ (c 0.2, H_2O), $\text{pK}_{\text{Est}} < 0$. Purified by recryst three times from EtOH and once from MeOH. The *tetraacetate* has **m 193-195°**, $[\alpha]_{\text{D}}^{20} -16^\circ$ (c 0.14, H_2O). [Benn et al. *J Chem Soc, Chem Commun* 445 1965; Kjaer et al. *Acta Chem Scand* **10** 432 1956; Marsh et al. *Acta Cryst (Sect B)* **26** 1030 1970.] It is a β -D-thioglucopyranoside substrate for thioglucosidase [MacLeod and Rossiter *Phytochem* **25** 1047 1986].

α -Solanine (solan-5-en- 3β -yl- $[O^3$ - β -D-glucopyranosyl- O^2 - α -L-rhamnopyranosyl- β -D-galactopyranoside]) [20562-02-1] **M 868.1, m 285°(dec)**, **286°(dec)** (sintering $>190^\circ$), $[\alpha]_{\text{D}}^{20} -58^\circ$ (c 0.8, pyridine), pK^{15} **6.66**. Recrystd from EtOH, 85% aqueous EtOH, MeOH or aqueous MeOH as *dihydrate* **m 276-278°**. Solubility in H_2O is 25mg/L and 5% in pyridine, but it is very soluble in Et₂O and CHCl_3 . The *hydrochloride* is gummy or amorphous but has been crystd (**m $\sim 212^\circ$** dec). It has insecticidal properties. [Kuhn et al. *Chem Ber* **88** 1492 1955.]

Somatostatin [38916-34-6] **M 1637.9**, $[\alpha]_{\text{D}}^{25} -36^\circ$ (c 0.57, 1% AcOH). A tetradecapeptide which is purified by gel filtration on Sephadex G-25, eluting with 2N AcOH, and then by liquid partition chromatography on Sephadex G-25 using *n*-BuOH-AcOH- H_2O (4:1:5) and has $R_F = 0.4$. It is a brain growth hormone releasing-inhibiting factor which has also been synthesised. [Burgus et al. *Proc Natl Acad Sci USA* **70** 684 1973; Sorantakis and McKinley *Biochem Biophys Res Commun* **54** 234 1973; Hartridt et al. *Pharmazie* **37** 403 1982.]

Spectinomycin dihydrochloride pentahydrate (Actinospectacin) [21736-83-4] **M 495.3, m 205-207°(dec)**, $[\alpha]_{\text{D}}^{20} +14.8^\circ$ (c 0.4, H_2O), pK_1 **6.95**, pK_2 **8.70**. Purified from aqueous Me_2CO and is soluble in H_2O , MeOH and dilute acid and base but only slightly soluble in Me_2CO , EtOH, CHCl_3 and $^*\text{C}_6\text{H}_6$. The *free base* is an amorphous solid, **m 184-194°** with $[\alpha]_{\text{D}}^{20} -20^\circ$ (H_2O). [Wiley et al. *J Am Chem Soc* **93** 2652 1963; X-ray: Cochran et al. *J Chem Soc Chem Commun* 494 1972.] It is an aminoglycoside antibiotic which interacts with 16S ribosomal RNA [Moazet and Noller *Nature* **327** 389 1987]; and is used for the treatment of gonorrhoea [Rinehart *J Infect Dis* **119** 345 1969].

D-Sphingosine (2*S*,3*S*-D-erythro-2-aminooctadec-4*t*-ene-1,3-diol from bovine brain) [123-78-4] M 299.5, m 79-82°, 82° 82.5° (softens at ~70°), $[\alpha]_D^{22}$ -3.4° (c 2, CHCl₃), pK_{Est} ~ 8.8. Purified by recrystn from EtOAc, Et₂O or pet ether (60-80°) It is insoluble in H₂O but is soluble in Me₂CO, EtOH and MeOH. It has IR bands at 1590 and 875 cm⁻¹, and is characterised as the *tribenzoate* m 122-123° (from 95% EtOH). [Tipton *Biochem Prep* 9 127 1962.]

Spirilloxanthin [34255-08-8] M 596.9, m 216-218°, λ_{\max} 463, 493, 528 nm, $\epsilon_{1\text{cm}}^{1\%}$ 2680 (493 nm) in pet ether (b 40-70°). Crystd from CHCl₃/pet ether, acetone/pet ether, *C₆H₆/pet ether or *C₆H₆. Purified by chromatography on a column of CaCO₃/Ca(OH)₂ mixture or deactivated alumina. [Polgar et al. *Arch Biochem Biophys* 5 243 1944.] Stored in the dark in an inert atmosphere, at -20°.

Squalane (Cosbiol, 2,6,10,15,19,23-hexamethyltetracosane, perhydro-squalene) [111-01-3] M 422.8, m -38°, b 176°/0.05mm, 210-215°/1mm, 274°/10mm, ~350°/760mm, d_4^{20} 0.80785, n_D^{20} 1.416. Purified by fractional distn *in vacuo* or evap distn. Soluble in pet ether, *C₆H₆, Et₂O and CHCl₃, slightly sol in alcohols, Me₂CO and AcOH but insol in H₂O [Staudinger and Leupold *Helv Chim Acta* 15 223 1932; Sax and Stross *Anal Chem* 29 1700 1951; Mandai et al. *Tetrahedron Lett* 22 763 1981].

Squalene (*all-trans*-2,6,10,15,19,23-hexamethyl-2,6,10,14,18,22-tetracosahexaene) [111-02-4] M 410.7, m ~75°, b 203°/0.1mm, 213°/1mm, 285°/25mm, d^{25} 0.8670, n 1.4905. Crystd repeatedly from Me₂CO (1.4mL/g) using a Dry-ice bath, washing the crystals with cold acetone, then freezing the squalene under vacuum. Squalene was further purified by passage through a column of silica gel or chromatographed on activated alumina, using pet ether as eluent and stored in vac in the dark. Dauben et al. [*J Am Chem Soc* 74 4321 1952] purified squalene *via* its hexachloride and is bactericidal. [Capstack et al. *J Biol Chem* 240 3258 1965; Krishna et al. *Arch Biochem Biophys* 114 200 1966; Heilbron and Thompson *J Chem Soc* 883 1929; Karrer et al. *Helv Chim Acta* 13 1084 1930; UV: Farmer et al. *J Chem Soc* 544 1943.]

Starch [9005-84-9] M (162.1)_n. Defatted by Soxhlet extraction with Et₂O or 95% EtOH. For fractionation of starch into "amylose" and "amylopectin" fractions, see Lansky et al. [*J Am Chem Soc* 71 4066 1949].

Sterigmatocystin (3*a*,12*c*-dihydro-8-hydroxy-6-methoxy-3*H*-furo[3',2',:4,5]furo[2,3-*c*]-xanthen-7-one) [10048-13-2] M 324.3, m 246°, 247-248°, $[\alpha]_D^{20}$ -398° (c 0.1, CHCl₃), pK_{Est} ~ 8.0. Recrystd from amyl acetate, Me₂CO or EtOH and sublimed *in vacuo*. It has UV λ_{\max} at 208, 235, 249 and 329nm (log ϵ 4.28, 4.39, 4.44 and 4.12). [UV: Bullock et al. *J Chem Soc* 4179, 1962; UV, IR: Holker and Mulheirn *J Chem Soc Chem Commun* 1576, 1576 1968; Birkinshaw and Hammady *Biochem J* 65 162 1957.] This mycotoxin induces bone marrow changes in mice [Curry et al. *Mutation Res* 137 111 1984].

Stigmatellin A (2-[4,6-dimethoxy-3,5,11-trimethyltridecatri-7*t*,9*t*,11*t*-enyl]-8-hydroxy-5,7-dimethoxy-3-methyl-4*H*-1-benzopyran-4-one) [91682-96-1] M 514.6, m 128-130°, $[\alpha]_D^{20}$ +38.5° (c 2.3, MeOH), pK_{Est} ~7 (phenolic OH). It is stable in aqueous soln at neutral pH but decomposes at pH <5. Purified by recrystn from toluene-hexane). It has UV λ_{\max} : nm (ϵ) 248sh (41000), 258 (59500) 267 (65500), 279 (41400) and 335 (5200) in MeOH; 249sh (45600), 258 (60000), 268 (72700), 277 (54100), 320 (2500) and 370 (3000) in MeOH + 1 drop of N KOH; 243sh (29300), 264 (63200), 274 (64100), 283sh (45800), 329 (4800) and 420 (21000) in MeOH + 6N HCl; and IR (CHCl₃) ν : 3550m, 1645schs, 1635ss, 1620ss, 1590s, 1510m and 905m cm⁻¹. It gives colour reactions at 110° with vanillin/H₂SO₄ (grey), Ce(IV)/(NH₄)₂SO₄ (yellow) and phosphomolybdate (blue-grey). [Höfle et al. *Justus Liebigs Ann Chem* 1882 1984.] It inhibits electron transport [Jagow and Link *Methods Enzymol* 126 253 1986; Robertson et al. *Biochemistry* 32 1310 1933], and has antibiotic properties [Kunze et al. *J Antibiot* 37 454 1984]. The 7*t*,9*t*,11*c*-isomer is *Stigmatellin B*.

Streptomycin sulfate [3810-74-0] M 1457.4, $[\alpha]_D^{20}$ -84.3° (c 3, H₂O), pK_{Est(1)} ~ 9.5 (MeNH), pK_{Est(2,3)} ~ 13.4 (guanidino). Recrystd from H₂O-EtOH, washed with a little EtOH, Et₂O and dried in a vacuum. [UV and IR: Grove and Randall *Antibiotics Monographs* 2 163 1855; Heuser et al. *J Am Chem Soc* 75 4013 1953, Kuehl et al. *J Am Chem Soc* 68 1460 1946; Regna et al. *J Biol Chem* 165 631 1946.] During protein synthesis it inhibits initiation and causes misreading of mRNA [Zierhut et al. *Eur J Biochem* 98 577 1979; Chandra and Gray *Methods Enzymol* 184 70 1990].

Streptonigrin (nigrin, 5-amino-6-[7-amino-5,8-dihydro-6-methoxy-5,8-dioxo-2-quinolinyl]-4-[2-hydroxy-3,4-dimethoxyphenyl]-3-methyl-2-pyridinecarboxylic acid) [3930-19-6] M 506.5, m 262-263°, 275°(dec), pK 6.3 (1:1 aq dioxane). Purified by TLC on pH 7-buffered silica gel (made from a slurry of Silica Gel 60 and 400mL of 0.05M phosphate buffer pH 7.0) and eluted with 5% MeOH/CHCl₃. The extracted band can then be recrystd from Me₂CO or dioxane as almost black plates or needles. It is soluble in pyridine, Me₂NCHO, aqueous NaHCO₃ (some dec), and slightly soluble in MeOH, EtOH, EtOAc and H₂O. It has UV λ_{\max} 248, 375-380nm (ϵ 38400 and 17400). [Weinreb et al. *J Am Chem Soc* **104** 536 1982; Rao et al. *J Am Chem Soc* **85** 2532 1963.] It is an antineoplastic and causes severe bone marrow depression [Wilson et al. *Antibiot Chemother* **11** 147 1961].

Streptozotocin (N-[methylnitrosocarbonyl]- α -D-glucosamine, streptozocin) [18883-66-4] M 265.2, m 111-114°(dec), 114-115°(dec), 115°(dec with evolution of gas), $[\alpha]_D^{20} \sim +39^\circ$ (H₂O, may vary due to mutarotation). Recrystd from 95% EtOH and is soluble in H₂O, MeOH and Me₂CO. It has UV λ_{\max} 228nm (ϵ 6360) in EtOH. The tetraacetate has m 111-114°(dec), $[\alpha]_D^{25} +41^\circ$ (c 0.78, 95% EtOH) after recrystn from EtOAc. [Herr et al. *J Am Chem Soc* **89** 4808 1967; NMR: Wiley et al. *J Org Chem* **44** 9 1979.] It is a potent methylating agent for DNA [Bennett and Pegg *Cancer Res* **41** 2786 1981].

Subtilisin (from *Bacillus subtilis*) [9014-01-1] [EC 3.4.21.62]. Purified by affinity chromatography using 4-(4-aminophenylazo)phenylarsonic acid complex to activated CH-Sepharose 4B. [Chandraskaren and Dhar *Anal Biochem* **150** 141 1985].

Succinyl coenzyme A trisodium salt [108347-97-3] M 933.5. If it should be purified further then it should be dissolved in H₂O (0.05g/mL) adjusted to pH 1 with 2M H₂SO₄ and extracted several times with Et₂O. Excess Et₂O is removed from the aqueous layer by bubbling N₂ through it and stored frozen at pH 1. When required the pH should be adjusted to 7 with dilute NaOH and used within 2 weeks (samples should be frozen). Succinyl coenzyme A is estimated by the hydroxamic acid method [*J Biol Chem* **242** 3468 1967]. It is more stable in acidic than in neutral aqueous solutions. [*Methods Enzymol* **128** 435 1986.]

2-Sulfobenzoic cyclic anhydride (2,1-benzoxathiazol-3-one 1,1-dioxide) [81-08-3] M 184.2, m 126-127°, 129.5°, 130°, b 184-186°/18mm. If the sample has hydrolysed extensively (presence of OH band in the IR) then treat with an equal bulk of SOCl₂ reflux for 3h (CaCl₂ tube), evaporate and distil residue in a vacuum, then recrystd from *C₆H₆, Et₂O-*C₆H₆ or CHCl₃ (EtOH free by passing through Al₂O₃, or standing over CaCl₂). [Clarke and Breger *Org Synth Coll Vol I* 495 1948.] Used for modifying ζ -amino functions of lysyl residues in proteins [Bagree et al. *FEBS Lett* **120** 275 1980]. (see entry on p. 126.)

Syrexin (from bovine liver). Purified by (NH₄)₂SO₄ pptn, then by pH step elution from chromatofocusing media in the absence of ampholytes. [Scott et al. *Anal Biochem* **149** 163 1985.]

Taurodeoxycholic acid sodium salt monohydrate (n-[desoxycholy]taurine Na salt H₂O) [1180-95-6] M 539.7, m 171-175°, $[\alpha]_D^{23} +37^\circ$ (c 1, H₂O), pK 1.4 (free acid). The salt is recrystd from EtOH-Et₂O. Its solubility in H₂O is 10%. The free acid has m 141-144°. [Norman *Ark Kemi* **8** 331 1956.] It forms mixed micelles and solubilises some membrane proteins [Hajjar et al. *J Biol Chem* **258** 192 1983].

Terramycin (oxytetracycline) [79-57-2] M 460.4 (anhy), 496.5 (2H₂O), sinters at 182°, melts at 184-185°(dec), $[\alpha]_D^{20} -196.6^\circ$ (equilibrium in 0.1M HCl), -2.1° (equilibrium in 0.1M NaOH). Crystd (as dihydrate) from water or aqueous EtOH.

2,2:5',2''-Terthiophene [1081-34-1] M 248.4, m 92-93°, 94-95°, 94-94.5°, 94-96°. Recrystd from MeOH, *C₆H₆, pet ether or MeOH. [UV: Zechmeister and Sease *J Am Chem Soc* **69** 273 1947; Steinkopf et al. *Justus Liebigs Ann Chem* **546** 180 1941.] Phototoxic nematocide [Cooper and Nitsche *Bioorg Chem* **13** 36 1985; Chan et al. *Phytochem* **14** 2295 1975]. See also Terthiophene on p. 356 in Chapter 4.

Tetracycline [60-54-8] M 444.4, m 172-174°(dec), $[\alpha]_{546}^{20} +270^\circ$ (c 1, MeOH), pK_1^{25} 3.30, pK_2^{25} 7.68, pK_3^{25} 9.69. Crystd from toluene.

Tetracycline hydrochloride [64-75-5] M 480.9, m 214°(dec), 215-220°, $[\alpha]_D^{25} -258^\circ$ (c 0.5, 0.1N HCl), $[\alpha]_D^{20} -245^\circ$ (c 1, MeOH), pK_1 1.4 (enolic OH), pK_2 7.8 (phenolic OH), pK_3 9.6 (Me₂N). Recrystd from MeOH + *n*-BuOH or *n*-BuOH + HCl. It is insoluble in Et₂O and pet ether. It has UV λ_{max} at 270 and 366nm in MeOH. [Gottstein et al. *J Am Chem Soc* **81** 1198 1959; Conover et al. *J Am Chem Soc* **84** 3222 1962.]

6R-Tetrahydro-erythro-biopterin dihydrochloride (BH₄.2HCl, 6R-2-amino-4-hydroxy-6-[[1R,2S]-1,2-dihydroxypropyl]-5,6,7,8-tetrahydropteridine 2HCl) [69056-38-8] M 316.2, m 245-246°(dec), $[\alpha]_D^{25} -6.8^\circ$ (c 0.67, 0.1N HCl), pK_1 1.37 (pyrimidine⁺), pK_2 5.6 (5-NH⁺), pK_3 10.6 (acidic, 3NH). Recrystn from HCl enriches BH₄ in the natural 6R isomer. Dissolve the salt (~6g) in conc HCl (15mL) under gentle warming then add EtOH (30mL) dropwise, chill and collect the colourless needles (67%, up to 99% if mother liquors are concentrated), and dried *in vacuo* immediately over P₂O₅ and KOH. Stores indefinitely at -20° in a dry atmosphere, Better store in sealed ampoules under dry N₂. It can be recrystd from 6N aqueous HCl. It has UV λ_{max} (2N HCl) 264nm (ϵ 16770; pH 3.5 phosphate buffer) 265nm (ϵ 13900); (pH 7.6) 297nm (ϵ 9500) and 260nm sh (ϵ 4690). It has been separated from the 6R-isomer by HPLC on a Partisil-10SCX column using 30mM ammonium phosphate buffer (pH 3.0) containing 3mM NaHSO₃ (2mL/min flow rate; 275nm detector) with retention times of 5.87min (6R) and 8.45min (6S). It is stable in acidic soln and can be stored for extended periods at -20° in 0.04M HCl. Above pH 7 the neutral species are obtained and these are readily oxidised by oxygen in the solvent to quinonoid species and then further oxidation and degradation occurs at room temperatures. These changes are slower at 0°. The *sulfate salt* can be obtained by recrystn from 2M H₂SO₄ and is less soluble than the hydrochloride salt. The 6R-2,5,1',2'-tetraacetylbiopterin derivative has m 292° (dec) after recrystn from MeOH (100 parts) and $[\alpha]_{589}^{20} -144^\circ$ (c 0.5, CHCl₃), $[\alpha]_{589}^{20} +12.8^\circ$ (c 0.39, Me₂SO). [NMR, UV: Matsuura et al. *Heterocycles* **23** 3115 1985; Viscontini et al. *Helv Chim Acta* **62** 2577 1979; Armarego et al. *Aust J Chem* **37** 355 1984.]

Tetrahydrofolic acid dihydrochloride 2H₂O (THFA, 6S- or 6RS- 5,6,7,8-tetrahydrofolic acid 2HCl 2H₂O, 5,6,7,8-tetrahydropteroyl-L-glutamic acid 2HCl 2H₂O) [135-16-0] M 544.4, m >200°(dec), $[\alpha]_D^{27} +16.9^\circ$ (H₂O pH 7.0 + 2-mercaptoethanol), pK_1 1.7 (pyrimidine N⁺), pK_2 2.4 (10N⁺), pK_3 3.5 (α -CO₂H), pK_4 4.9 (γ -CO₂H), pK_5 5.6 (5-NH⁺), pK_6 10.4 (acidic, 3NH). Very high quality material is now available commercially and should be a white powder. It can be dried over P₂O₅ in a vacuum desiccator and stored in weighed aliquots in sealed ampoules. It is stable at room temp in sealed ampoules for many months and for much more extended periods at -10°. When moist it is extremely sensitive to moist air whereby it oxidises to the yellow 7,8-dihydro derivative. In soln it turns yellow in colour as it oxidises and then particularly in the presence of acids it turns dark reddish brown in colour. Hence aqueous solutions should be frozen immediately when not in use. It is always advisable to add 2-mercaptoethanol (if it does not interfere with the procedure) which stabilises it by depleting the soln of O₂. The *sulfate salt* is more stable but then it is much less soluble. The best way to prepare standard solns of this acid is to dissolve it in the desired buffer and estimate the concentration by UV absorption in pH 7 buffer at 297nm (ϵ 22,000 M⁻¹cm⁻¹). If a sample is suspect it is not advisable to purify it because it is likely to deteriorate further as "dry box" conditions are necessary. Either a new sample is purchased or one is freshly prepared from folic acid. It has pKa values of -0.1, 4.3 and 9.0. [Hafeti et al. *Biochem Prep* **7** 89 1960; UV: Mathews and Huennekens *J Biol Chem* **235** 3304 1960; Osborn and Huennekens *J Biol Chem* **233** 969 1958; O'Dell et al. *J Am Chem Soc* **69** 250 1947; Blakley *Biochem J* **65** 331 1957; Asahi *Yakugaku Zasshi (J Pharm Soc Japan)* **79** 1548 1959.]

5,6,7,8-Tetrahydropterin sulfate (2-amino-5,6,7,8-tetrahydropteridin-4-one H₂SO₄) [20350-44-1] M 265, m >200°(dec), pK_1^{25} 1.3 (pyrimidine⁺), pK_2^{25} 5.6 (5-NH⁺), pK_3^{25} 10.6 (acidic, 3NH). If it has become too strongly violet in colour then it may need reducing again. Best to check the UV absorption in N HCl where it has a peak at ~265nm which drops sharply to zero having no absorption at *ca* 340nm. The presence of absorption at 340nm indicated oxidation to quinonoid or 7,8-dihydropterin. If the absorption is weak then dissolve in the minimum volume of anhydrous trifluoroacetic acid (fume hood) add charcoal, filter, then add one or two drops of N H₂SO₄ followed by dry Et₂O at 0°, allow the white tetrahydro

salt to settle and collect, and wash with dry Et₂O, by centrifugation. Dry the residue in a vacuum desiccator over P₂O₅ and KOH. Store in aliquots in the dark at <0°. It has UV λ_{\max} at 265nm (ϵ 16980) at pH -1.0 (dication); 219nm (ϵ 23440) and 266nm (ϵ 12880) at pH 3.5 (monocation); 220nm (ϵ 18620), [260nm (ϵ 4270)sh] and 299nm (ϵ 9330) at pH 8.0 (neutral species); and 218nm (ϵ 10000), [240nm (ϵ 5500)sh] and 287nm (ϵ 5500) at pH 13 (anion). [Blakley *Biochem J* **72** 707 1959; Asahi *Yakugaku Zasshi (J Pharm Soc Jpn)* **79** 1557 1959; Pfeleiderer in *Pterins and Folate* (Benkovic and Blakley Eds) J Wiley Vol 2 p97 1985.]

Thiamine monophosphate chloride 1H₂O (Aneurine monophosphate chloride) [532-40-1] M 416.8, m 193°(dec), 200°(dec), 200-203°(dec), pK₁ 2.40, pK₂ 4.80, pK₃ 6.27, pK₄ 9.65, pK₅ 10.20. Purified by recrystn from aqueous HCl, EtOH slightly acidified with HCl, EtOH-Me₂CO, H₂O, or H₂O-EtOH + Et₂O. Dissolve in a small volume of H₂O and mix with EtOH + Me₂CO (1:1) to give the HCl.H₂O as crystals. Filter, wash with Et₂O and dry in a vacuum. The *chloride hydrochloride*, m 215-217°(dec) is obtained when crystd from aqueous HCl. [Wenz et al. *Justus Liebigs Ann Chem* **618** 2280 1958, Viscontini et al. *Helv Chim Acta* **34** 1388 1951, Leichsrenring and Schmidt *Chem Ber* **95** 767 1962; McCormick and Wright *Methods Enzymol* **18A** 141, 147 1970.]

Thiamphenicol (1R,2R-2-[2,2-dichloroacetyl-amino]-1-[4-methanesulfonylphenyl]-propan-1,3-diol) [15318-45-3 (D-threo), 90-91-5] M 356.2, m 163-166°, 165.2-165.6°, 165-166°, [α]_D²⁵ +15.6° (c 2, EtOH), pK 7.2. Recrystd from H₂O or CHCl₃. UV λ_{\max} 224, 266 and 274nm (ϵ 13700, 800 and 700) in 95% EtOH. The *1S,2S-isomer* [14786-51-7] has m 164.3-166.3° (from H₂O + EtOAc + pet ether) and [α]_D²⁵ -12.6° (c 1, EtOH); and the *racemate 1RS,2RS Racefenical* [15318-45-3] has m 181-183° (sinter at 180-183°) from CHCl₃-EtOAc-pet ether. [Cutler et al. *J Am Chem Soc* **74** 5475, 5482 1952; UV: Nachod and Cutler *J Am Chem Soc* **74** 1291 1952; Suter et al. *J Am Chem Soc* **75** 4330 1953; Cutler et al. *J Am Pharm Assoc* **43** 687 1954.]

Thiazolyl blue tetrazolium bromide (MTT, 3-[4,5-dimethyl-2-thiazolyl]-2,5-diphenyl-2H-tetrazolium bromide) [298-93-1, 2348-71-2] M 414.3, m 171°. It is recrystd by dissolving in MeOH containing a few drops of HBr and then adding dry Et₂O to complete the crystn, wash the needles with Et₂O and dry in a vacuum desiccator over KOH. [Beyer and Pyl *Chem Ber* **87** 1505 1954.]

2-Thiocytosine (4-amino-2-mercaptopyrimidine) [333-49-3] M 127.2, m 236-237°(dec), 285-290°(dec), pK₁²⁰ 3.90 (NH₂), pK₂²⁰ 11.10 (SH). It is recrystd from hot H₂O and dried at 100° to constant weight. [Brown *J Appl Chem (London)* **9** 203 1959; Russell et al. *J Am Chem Soc* **71** 2279 1949.] It is used in transcription and translation studies [Rachwitz and Scheit *Eur J Biochem* **72** 191 1977.]

6-Thioguanine [154-42-7] M 167.2, m >300°, pK₁²³ 8.2 (SH), pK₂²³ 11.6 (acidic, 9-NH). Recrystd from H₂O as needles. It has UV λ_{\max} at 258 and 347nm (H₂O, pH 1) and 242, 270 and 322nm (H₂O, pH 11). [Elion and Hitchings *J Am Chem Soc* **77** 1676 1955; Fox et al. *J Am Chem Soc* **80** 1669 1958.] It is an antineoplastic agent [Kataoka et al. *Cancer Res* **44** 519 1984].

Thrombin (from bovine blood plasma) [9002-04-4] M_r 32,600 [EC 3.4.4.13]. Purified by chromatography on a DEAE-cellulose column, while eluting with 0.1M NaCl, pH 7.0, followed by chromatography on Sephadex G-200. Final preparation was free from plasminogen and plasmin. [Yin and Wessler *J Biol Chem* **243** 112 1968.]

Thrombin from bovine blood was purified by chromatography using *p*-chlorobenzylamino- ϵ -aminocaproyl agarose, and gel filtration through Sephadex G-25. [Thompson and Davie *Biochim Biophys Acta* **250** 210 1971.]

Thrombin from various species was purified by precipitation of impurities with rivanol. [Miller *Nature* **184** 450 1959.]

L-Thyroxine sodium salt (5H₂O) [6106-07-6] M 888.9, [α]₅₄₆²⁰ +20° (c 2, 1M HCl + EtOH, 1:4). Crystd from absolute EtOH and dried for 8h at 30°/1mm.

D-Thyroxine {O-[3,5-diiodo-4-oxypheyl]-3,5-diiodo-D(-)-tyrosine, 3,3',5,5'-tetra-iodo-D-thyronine} [51-49-0] M 776.9, m 235°(dec), 235-236°(dec), 340°(dec), [α]_D²⁰ +4.5° (c 3, aq

0.2N NaOH in 70% EtOH), $[\alpha]_D^{20} -17^\circ$ (c 2, aq N HCl + EtOH 1:4), $pK_1^{2.5} 2.2$ (CO₂H), $pK_2^{2.5} 8.40$ (OH), $pK_3^{2.5} 10.1$ (NH₂). Recrystd from H₂O as needles or from an ammonical soln by dilution with H₂O, MeOH or Me₂CO. Also purified by dissolving ~6.5 g in a mixture of MeOH (200mL) and 2N HCl (20mL), add charcoal, filter then add NaOAc soln to pH 6 and on standing the thyroxine separates, is washed with MeOH then Me₂CO and dried *in vacuo*. The *N*-formyl-*D*-thyroxine derivative has *m* 210° and $[\alpha]_{546}^{21} -26.9^\circ$ (c 5, EtOH). The racemate \pm -thyroxine has *m* 256° and is purified in the same way. [Nahm and Siedel *Chem Ber* 96 1 1963; Salter *Biochem J* 24 471 1930.]

L-Thyroxine (*O*-[3,5-diiodo-4-oxyphenyl]-3,5-diiodo-L-(+)-tyrosine, 3,3',5,5'-tetraiodo-D-thyronine) [51-48-9] *M* 776.9, *m* 229-230°(dec), ~235°(dec), 237°(dec), $[\alpha]_D^{22} -5.1^\circ$ (c 2, aq N NaOH + EtOH 1:2), $[\alpha]_D^{22} +15^\circ$ (c 5, aq N HCl in 95% EtOH 1:2), $[\alpha]_D^{22} +26^\circ$ (EtOH/1M aq HCl; 1:1) (*pK* 6.6). Purification is the same as for the D-isomer above. Likely impurities are tyrosine, iodotyrosine, iodothyroxines and iodide. Dissolve in dilute ammonia at room temperature, then crystd by adding dilute acetic acid to pH 6. The *N*-formyl-*L*-thyroxine has *m* 214°(dec) and $[\alpha]_{546}^{21} +27.8^\circ$ (c 5, EtOH). [Harrington et al. *Biochem J* 39 164 1945; Nahm and Siedel *Chem Ber* 96 1 1963; Reineke and Turner *J Biol Chem* 161 613 1945; Chalmers et al. *J Chem Soc* 3424 1949.]

Tissue inhibitor of metalloproteins (TIMP, from human blood plasma), *M_r* ~30,000. Purified by a [anti-human amniotic fluid-TIMP]-Sephrose immuno-affinity column eluted with 50mM glycine/HCl pH 3.0 buffer that is 0.5M in NaCl then by gel filtn [Cawston et al. *Biochem J* 238 677 1986].

***dl*- α -Tocopherol** (see vitamin E) [59-02-9] *M* 430.7, *A* $1_{cm}^{1\%}$ 74.2 at 292 nm in MeOH. Dissolved in anhydrous MeOH (15mL/g) cooled to -6° for 1h, then chilled in a Dry-ice/acetone bath, crystn being induced by scratching with a glass rod.

γ -Tocopherol (3,4-dihydro-2,7,8-trimethyl-2-(4,8,12-trimethyltridecyl)-2*H*-benzopyran-6-ol) [54-28-4] *M* 416.7, *m* -30°, *b* 200-210°/0.1mm, *d*₄²⁰ 0.951, *n*_D²⁰ 1.505, $[\alpha]_D^{20} -2.4^\circ$ (EtOH). Purified by distn at high vacuum and stored in dark ampoules under N₂. UV λ_{max} 298nm (*A*_{1cm}^{1%} 92.8). It is insoluble in H₂O but soluble in organic solvents. The *allophanate* (used for separating isomers) has *m* 136-138°, $[\alpha]_D^{18} +3.4^\circ$ (CHCl₃). [Baxter et al. *J Am Chem Soc* 65 918 1943; Emerson et al. *Science* 83 421 1936, *J Biol Chem* 113 319 1936.]

Toluene-2,4-diisocyanate (toluene-2,4-diisocyanate). [584-84-9] *M* 174.2, *m* 19.5-21.5°, 20-22°, 28°, *b* 126°/11mm, 124-126°/18mm, 250°/760mm. It is purified by fractionation in a vacuum and should be stored in a dry atmosphere. It is soluble in organic solvents but reacts with H₂O, alcohols (slowly) and amines all of which could cause explosive polymerisation. It darkens on exposure to light. It has a sharp pungent odour, is TOXIC and is IRRITATING TO THE EYES. [Siefken *Justus Liebigs Ann Chem* 562 75, 96, 127 1949; Bayer *Angew Chem* 59 257 1947.] It is a reagent for covalent crosslinking of proteins [Wold *Methods Enzymol* 25 623 1972.]

Tomatidine (5 α ,20 β ,22 α ,25 β ,27-azaspirostan-3 β -ol) [77-59-8] *M* 415.7, *m* 202-206°, $[\alpha]_D^{20} +5.9^\circ$ (c 1, MeOH), $[\alpha]_D^{20} +8^\circ$ (CHCl₃). Forms plates from EtOAc. Also purified by dissolving 80mg in *C₆H₆ and applying to an Al₂O₃ column (3.0g) and eluting with *C₆H₆, evaporating and recrystallising three times from EtOAc. The *hydrochloride* has *m* 265-270° from EtOH and $[\alpha]_D^{25} -5^\circ$ (MeOH). [IR: Uhle *J Am Chem Soc* 83 1460 1961; Kessar et al. *Tetrahedron* 27 2869 1971; Schreiber and Adams *Experientia* 17 13 1961.]

Tomatine (2*S*,25*S*-3 β , β -lycotetraoxy-5 α -spirostan) [17406-45-0] *M* 1034.2, *m* 263-268°(dec), 290-291°(evac capillary), 283.5-287°(dec), 272-277°(dec), 300-305°(dec), $[\alpha]_D^{20} -18^\circ$ to -34° (c 0.55, pyridine). Recrystd from MeOH, EtOH, aqueous EtOH or dioxane + NH₃. It is almost insoluble in pet ether, Et₂O or H₂O. [Reichstein *Angew Chem* 74 887 1962.]

***N*-Tosyl-L-lysine chloromethyl ketone** (3*S*-1-chloro-3-tosylamino-7-amino-2-heptanone HCl) [4272-74-6] *M* 369.3, *m* 150-153°(dec), 156-158°(dec), ~165°(dec), $[\alpha]_D^{20} -7.3^\circ$ (c 2, H₂O), *pK*_{Est} ~ 10.6 (7-NH₂). The hydrochloride slowly crystallises from a conc soln in absolute EtOH,

thinned with EtOH-Et₂O for collection and dried *in vacuo*. It is a suicide enzyme inhibitor [Matsuda et al. *Chem Pharm Bull Jpn* **30** 2512 1982; Shaw et al. *Biochemistry* **4** 2219 1965].

Transferrin (from human or bovine serum) [11096-37-0] $M_r \sim 80,000$. Purified by affinity chromatography on phenyl-boronate agarose followed by DEAE-Sephacel chromatography. The product is free from haemopexin. [Cook et al. *Anal Biochem* **149** 349 1985; Aisen and Listowsky *Ann Rev Biochem* **49** 357 1980.]

Trehalase (from kidney cortex) [9025-52-9] [EC 3.2.1.28]. Purified by solubilising in Triton X-100 and sodium deoxycholate, and submitting to gel filtration, ion-exchange chromatography, conA-Sepharose chromatography, phenyl-Sepharose CL-4B hydrophobic interaction chromatography, Tris-Sepharose 6B affinity and hydrolyapatite chromatography. Activity was increased 3000-fold. [Yoneyama *Arch Biochem Biophys* **255** 168 1987.]

Trifluoperazine dihydrochloride (10-[3-{4-methyl-1-piperazinyl}propyl]-2-trifluoro-methyl-phenothiazine 2HCl) [440-17-5] M 480.4, m 240-243°, 242-243°, pK_1 3.9, pK_2 8.1. Recrystd from abs EtOH dried *in vacuo* and stored in tightly stoppered bottles because it is *hygroscopic*. It is soluble in H₂O but insoluble in *C₆H₆, Et₂O and alkaline aqueous soln. It has UV λ_{max} at 258 and 307.5nm (log ϵ 4.50 and 3.50) in EtOH (neutral species). [Craig et al. *J Org Chem* **22** 709 1957.] It is a calmodulin inhibitor [Levene and Weiss *J Pharmacol Exptl Ther* **208** 454 1978], and is a psychotropic agent [Fowler *Arzneim.-Forsch* **27** 866 1977].

T4-RNA ligase (from bacteriophage-infected *E.coli*) M_r 43,500 [EC 6.5.1.3 for RNA lyase]. Purified by differential centrifugation and separation on a Sephadex A-25 column, then through hydroxylapatite and DEAE-glycerol using Aff-Gel Blue to remove DNAase activity. (Greater than 90% of the protein in the enzyme preparation migrated as a single band on gradient polyacrylamide gels containing SDS during electrophoresis.) [McCoy et al. *Biochim Biophys Acta* **562** 149 1979.]

Tubercidin (7-deazaadenosine) [69-33-0] M 266.3, m 247-248°, $[\alpha]_D^{17}$ -67° (50% aq AcOH), pK^{10} 5.2-5.3. Forms needles from hot H₂O. It is soluble in H₂O (0.33%), MeOH (0.5%) and EtOH (0.05%). It has UV λ_{max} 270nm (ϵ 12100) in 0.001N NaOH. The *picrate* has m 229-231°(dec). [Tolman et al. *J Am Chem Soc* **91** 2102 1969; Mizuno et al. *J Org Chem* **28** 3329 1963, IR: Anzai et al. *J Antibiot (Japan)* **[9]** **10** 201 1957.]

Tunicamycin [11089-65-9] m 234-235°(dec), $[\alpha]_D^{20}$ +52° (c 0.5, pyridine), $pK_{Est} \sim 9.4$. The components are purified by recrystallising 3 times from hot glass-distilled MeOH and the white crystals are dissolved in 25% aqueous MeOH and separated on a Partisil ODS-10 μ column (9.4 x 25 cm) [Magnum-9 Whatman] using a 260 nm detector. The column was eluted with MeOH:H₂O mixture adjusted to 1:4 (v/v) then to 2:4 (v/v). The individual components are recovered and lyophilised. Ten components were isolated and all were active (to varying extents) depending on the lengths of the aliphatic side-chains. The mixture has UV λ_{max} 205 and 260nm ($A_{1cm}^{1\%}$ 230 and 110). Stable in H₂O at neutral pH but unstable in acidic soln. It inhibits protein glycosylation. [Mahoney and Duskin *J Biol Chem* **254** 6572 1979; Elnein *Trends Biochem Sci* **6** 219 1981; Takatsuki *J Antibiot* **24** 215 1971.]

Ubiquinol-cytochrome c reductase (from beef heart mitochondria) [9027-03-6] [EC 1.10.2.2]. Purified in Triton X-100 by solubilising the crude enzyme with Triton X-100, followed by hydroxylapatite and gel chromatography. The minimum unit contains nine polypeptide subunits of M_r 6000 - 49000 kD. [Engel et al. *Biochim Biophys Acta* **592** 211 1980.]

Uracil, uridine and uridine nucleotides. Resolved by ion-exchange chromatography AG1 (Cl⁻ form). [Lindsay et al. *Anal Biochem* **24** 506 1968.]

Uridine 5'-diphosphoglucose pyrophosphorylase (from rabbit skeletal muscle) [9029-22-6] M_r 350,000, [EC 2.7.7.9]. Purified by two hydrophobic chromatographic steps and gel filtration. [Bergamini et al. *Anal Biochem* 143 35 1984.] Also purified from calf liver by $(NH_4)_2SO_4$ (40-58%) pptn, $Ca_3(PO_4)_2$ gel filtration, DEAE-cellulose chromatography and recrystn by dialysis against increasing concentrations of $(NH_4)_2SO_4$ (from 10%) in 0.02M TEA (at 2.5% increments) until at 20% $(NH_4)_2SO_4$ it crystallises out [Hansen et al. *Methods Enzymol* 8 248 1966].

Uridine 5'-(1-thio) monophosphate [15548-52-4, 18875-72-4 (*Abs Stereochem specified*)] and **Uridine 5'-(α -thio) diphosphate** [*RS*(α -P) 27988-67-6; *R*(α -P) 72120-52-6], $pK_{Est(1)} \sim 6.4$, $pK_{Est(2)} \sim 9.5$ The Et_3N salt was purified by dissolving $\sim 4g$ in 500mL of H_2O (add a drop or two of Et_3N if it does not dissolve) and chromatographed by applying to a column (3 x 30cm) of DEAE-Sephadex A-25 and eluted with a 1.4L linear gradient of $Et_3NH.HCO_3$ from 0.05 to 0.55M, pH 7.8 and 4° . The product eluted between 0.2-0.3M $Et_3N.HCO_3$. Pooled fractions were evaporated and the residue was twice taken up in EtOH and evaporated to dryness to remove the last traces of $Et_3NH.HCO_3$. ^{31}P NMR: P_α is a doublet at -40.81 and -40.33, and P_β at 7.02ppm, $J_{\alpha,\beta}$ 32.96Hz. [*Biochemistry* 18 5548 1979.]

Uridylic acid (di-Na salt) [27821-45-0] M 368.2, m 198.5 $^\circ$, pK_3^{25} 6.63, pK_4^{20} 9.71. Crystd from MeOH.

Urokinase (from human urine) [9039-53-6] M_r 53,000, [EC 3.4.21.31]. Crystn of this enzyme is induced at pH 5.0 to 5.3 (4°) by careful addition of NaCl with gentle stirring until the soln becomes turbid (silky sheen). The NaCl concentration is increased gradually (over several days) until 98% of saturation is achieved whereby the urokinase crystallises as colourless thin brittle plates. It can be similarly recrystd to maximum specific activity [104K CTA units/mg of protein (Sherry et al. *J Lab Clin Med* 64 145 1964)]. [Lesuk et al. *Science* 147 880 1965; NMR: Bogusky et al. *Biochemistry* 28 6728 1989.] It is a plasminogen activator [Gold et al. *Biochem J* 262 1989].

(+)-Usnic acid (2,6-diacetyl-7,9-dihydroxy-8,9b-dimethyldibenzofuran-1,3(2H,9bH)-dione) [7562-61-0, 125-46-2] M 344.3, m 201-204 $^\circ$, 203-206 $^\circ$, $[\alpha]_{546}^{20} +630^\circ$ (c 0.7, $CHCl_3$), pK_1 4.4, pK_2 8.8, pK_3 10.7. This very weak acid is the natural form which is recrystd from Me_2CO , MeOH or $*C_6H_6$. At 25 $^\circ$ it is soluble in H_2O (<0.01%), Me_2CO (0.77%), EtOAc (0.88%), $MeOCH_2CH_2OH$ (0.22%) and furfural (7.32%). [Curd and Robertson *J Chem Soc* 894 1937; Barton and Brunn *J Chem Soc* 603 1953; resolution: Dean et al. *J Chem Soc* 1250 1953; synthesis: Barton et al. *J Chem Soc* 538 1956.]

(-)-Usnic acid (2,6-diacetyl-7,9-dihydroxy-8,9b-dimethyldibenzofuran-1,3(2H,9bH)-dione) [6159-66-6, 7562-61-0] M 344.3, m 201-204 $^\circ$, 204 $^\circ$, $[\alpha]_D^{20} -495^\circ$ (c 0.9, $CHCl_3$). Properties almost similar to those of the preceding entry.

Ustilagic acid (Ustizeajin B, di-D-glucosyldihydroxyhexadecanoic acid) [8002-36-6] M ~ 780 , m 146-147 $^\circ$, $[\alpha]_D^{23} +7^\circ$ (c 1, pyridine), $pK \sim 4.9$. It is a mixture of partly acetylated di-D-glucosyldihydroxyhexadecanoic acid which crystals from diethyl ether. Also purified from the culture by dissolving in hot MeOH, filtering and concentrating by blowing a current of air until the soln becomes turbid, then heating to 50 $^\circ$ and adding 4 vols of H_2O (also at 50 $^\circ$) and allowing to cool very slowly. Filter off the white solid and dry in air. [Lemieux et al. *Can J Chem* 29 409, 415 1951; *Can J Biochem Physiol* 33 289 1955.]

Valinomycin (Potassium ionophore I) [2001-95-8] M 111.3, m 186-187 $^\circ$, 190 $^\circ$, $[\alpha]_D^{20} +31.0^\circ$ (c 1.6, $*C_6H_6$). Recryst from dibutyl ether or Et_2O . Dimorphic, modification A crystallises from *n*-octane, and modification B crystallises from EtOH/ H_2O . Soluble in pet ether, $CHCl_3$, AcOH, BuOAc and Me_2CO . [*J Am Chem Soc* 97 7242 1975; UV, IR and NMR see *Chem Ber* 88 57 1955.]

(\pm)-Verapamil hydrochloride (5-[*N*-(3,4-dimethoxyphenylethyl)methylamino]-2-[3,4-dimethoxyphenyl]-2-isopropylvaleronitrile HCl) [23313-68-0] M 491.1, m 138.5-140.5 $^\circ$,

pK_{Est} ~ 10.6. Purified by dissolving in EtOH, filtering (if insoluble particles are present) and adding Et₂O, filtering the salt, washing with Et₂O and drying *in vacuo*. It has the following solubilities: hexane (0.001%), CH₂Cl₂ (~10%), MeOH (~10%) EtOH (20%) and H₂O (8.3%). It has UV λ_{\max} 232 and 278nm. The *free base* is a viscous yellow oil **b** 243-246°/0.01mm (n_D^{25} 1,5448) and is almost insol in H₂O but sol in organic solvents. It is a Ca channel antagonist and is a coronary vasodilator. [Ramuz *Helv Chim Acta* **58** 2050 1975; Harvey et al. *Biochem J* **257** 95 1989.]

Veratridine (3-veratroylveracevine) [71-62-5] **M 673.8, pK_{Est} ~7 (quinolizidine N).** A n alkaloid neurotoxin purified from veratrine. [McKinney et al. *Anal Biochem* **153** 33 1986.]

Vinblastine sulfate (vincaleucoblastine) [143-67-9] **M 909.1, m 284-285°, [α]_D²⁵ -28° (c 1, MeOH), pK₁ 5.4, pK₂ 7.4.** Purified by recrystn from H₂O and dried *in vacuo*. [Neuss et al. *J Am Chem Soc* **86** 1440 1964.] The *free base* is recrystd from MeOH or EtOH and has **m** 210-212°, 211-216°, [α]_D²⁵ +42° (CHCl₃); and has UV λ_{\max} 214 and 259nm (log ϵ 4.73 and 4.21). The *dihydrochloride dihydrate* has **m** 244-246°. [Bommer et al. *J Am Chem Soc* **86** 1439 1964.] It is a monoamine oxidase inhibitor [Keun Son et al. *J Med Chem* **33** 1845 1990].

Vincristine sulfate (22-oxovincaleucoblastine sulfate) [2068-78-2] **M 925.1, m 218-220°, [α]_D²⁵ +26.2° (CH₂Cl₂), pK₁ 5.0, pK₂ 7.4 (in 33% aq Me₂NCHO).** Recryst from MeOH. It has UV λ_{\max} 220, 255 and 296nm (log ϵ 4.65, 4.21 and 4.18). It is a monoamine oxidase inhibitor and is used in cancer research [Son et al. *J Med Chem* **33** 1845 1990; Horio et al. *Proc Natl Acad Sci USA* **85** 3580 1988].

Viomycin sulfate (Viocin, Tuberactinomycin B) [37883-00-4] **M 685.7, m 266°(dec), [α]_D¹⁷ -29.5° (c 1, H₂O), pK₁ 7.2 (8.2), pK₂ 10.3.** Crystd from H₂O-EtOH and dried in a vacuum. Dry material is *hygroscopic* and should be stored dry. The UV has λ_{\max} at 268 and 285nm (log ϵ 4.4 and 4.2) in H₂O. [Kitigawa et al. *Chem Pharm Bull Jpn* **20** 2176 1972.] The *hydrochloride* forms *hygroscopic* plates with **m** 270°(dec), [α]_D¹⁸ -16.6° (c 1, H₂O) with λ_{\max} 268nm (log ϵ 4.5) in H₂O; 268nm (log ϵ 4.4) in 0.1N HCl and 285nm (log ϵ 4.3) in 0.1N NaOH.

Vitamin A acid [Retinoic acid, 3,7-dimethyl-9-(2,6,6-trimethyl-1-cyclohexenyl)-2,4,6,8-nonatetraen-1-oic acid] [302-79-4] **M 300.4, m 180-181°, 180-182°, pK_{Est} ~ 4.2.** Purified by chromatography on silicic acid columns, eluting with a small amount of EtOH in hexane. Dissolve in Et₂O, wash with H₂O, dry (Na₂SO₄), evaporate and the solid residue crystd from MeOH (0.53g /3.5mL MeOH to give 0.14g) or EtOH. Also recrystd from *i*-PrOH, or as the *methyl ester* from MeOH. UV in MeOH has λ_{\max} 351nm (ϵ 45,000). 9-*Cis*-acid forms yellow needles from EtOH, **m** 189-190°, UV in MeOH has λ_{\max} 343nm (ϵ 36,500) and 13-*cis*-acid forms red-orange plates from *i*-PrOH, **m** 174-175°, UV has λ_{\max} 345nm (ϵ 39,800). Store in the dark, in an inert atmosphere, at 0° [Robeson et al. *J Am Chem Soc* **77** 4111 1955].

Vitamin A alcohol (retinol) [68-26-8] **M 286.5, A_{1cm}^{1%} (λ_{\max})(*all-trans*) 1832 (325 nm), (*13-cis*) 1686 (328nm), (*11-cis*) 1230 (319 nm), (*9-cis*) 1480 (323 nm), (*9,13-di-cis*) 1379 (324 nm), (*11,13-di-cis*) 908 (311 nm) in EtOH.** Purified by chromatography on columns of water-deactivated alumina eluting with 3-5% acetone in hexane. Separation of isomers is by TLC plates on silica gel G, developed with pet ether (low boiling)/methyl heptanone (11:2). Stored in the dark, under nitrogen, at 0°, as in diethyl ether, acetone or ethyl acetate. [See Gunghaly et al. *Arch Biochem Biophys* **38** 75 1952.]

Vitamin A aldehyde [all-trans-retinal; 3,7-dimethyl-9-(2,6,6-trimethyl-1-cyclohexenyl)-2,4,6,8-nonatetraen-1-al] [116-31-4] **M 284.4, m 61-64°.** Separated from retinol by column chromatography on water-deactivated alumina. Eluted with 1-2% acetone in hexane, or on TLC plates of silica gel G development. It crystallises from pet ether or *n*-hexane as yellow-orange crystals, and the UV in hexane has λ_{\max} 373nm (A_{1cm}^{1%} 1,548) [368nm (ϵ 48000)]. It is an **irritant** and is light sensitive. Store in sealed ampoules under N₂. The **semicarbazone** forms yellow crystals from CHCl₃-Et₂O or EtOH, **m** 199-201°(dec). The 9-*cis*-isomer [514-85-2] and the 13-*cis*-isomer [472-86-6] [λ_{\max} 375nm (ϵ 1250) in EtOH] are also available commercially.

Vitamin B₁ Hydrochloride [Aneurine hydrochloride, Thiamine hydrochloride, 3{(4-amino-2-methyl-5-pyrimidinyl)methyl}-4-methylthiazolium chloride monohydrochloride] [67-03-8] M 337.3, m 248°(dec), 249-250°, monohydrate m 135°(dec), pK₁²⁵ 4.8, pK₂²⁵ 9.2. Crystallises from 95% EtOH (sol, ca 1%). The monohydrate is dehydrated by drying at 100° *in vacuo* over H₂SO₄, but is *hygroscopic* and picks up one mol. of H₂O readily. It can be sterilised at 100° if the pH of the solution is below 5.5. The *nitrate* has m 196-200°(dec) and is more stable than the hydrochloride. The *picrolonate* crystallises from H₂O and is dimorphic, m 164-165° and 228-229°(dec). [Todd and Bergel *J Chem Soc* 364, 367 1937; *J Am Chem Soc* 58 1063, 1504 1936, 59 526 1937.]

Vitamin B₂ [Riboflavin, Lactoflavin, 6,7-dimethyl-9-(D-1'-ribityl)isoalloxazine] [83-88-5] M 376.4, m 278-282°(dec with darkening at 240°), 281-282°, [α]_D²⁵ -112° to -122° (c 2.5, 0.02M NaOH), [α]_D²⁰ -59° (c 0.23, AcOH), pK₁ 1.7, pK₂ 9.69 (10.2). It crystallises from H₂O as a yellow-orange powder in three different forms with differing amounts of H₂O. It melts if placed in an oil bath at 250°, but decomposes at 280° if heated at a rate of 5°/min. Solubility in H₂O is 1g in 3000-15000mL depending on the crystal structure. Sol in EtOH at 25° is 4.5mg in 100mL. Store in the dark because it is decomposed to lumichrome by UV light.

Vitamin B₆ hydrochloride (adermine, pyridoxine HCl, 3-hydroxy-4,5-bis[hydroxymethyl]-2-methylpyridine HCl) [58-56-0] M 205.6, m 208-208.5°, 208-209°(dec), 209-210°(dec), 205-212° (sublimes), pK₁²⁵ 5.0 (3-OH), pK₂²⁵ 8.96 (pyridinium⁺). Purified by recrystn from EtOH-Me₂CO, *n*-BuOH or MeOH-Et₂O. Its solubility in H₂O is 22% and in EtOH it is 1.1%. It is insoluble in Et₂O and CHCl₃. Acidic aqueous solns are stable at 120°/30 min. The *free base* has m 159-160° after recrystn from Me₂CO and sublimation at 140-145°/0.0001mm. It has UV λ_{max} at 290nm (ε 84000) in 0.1N aqueous HCl and 253 and 325nm (ε 3700 and 7100). [Khua and Wendt *Chem Ber* 71 780 1938, 72 311 1939; Harris and Folkers *J Am Chem Soc* 61 1242 1939; Harris et al. *J Am Chem Soc* 62 3198 1940.] See also **Pyridoxal-5'-phosphate H₂O** above.

Vitamin B₁₂ (cyanocobalamine, α-[5,6-dimethylbenzimidazolyl]cyano cobamide) [68-19-9] M 1355.4, m darkens at 210-220° and does not melt below 300°, [α]_D²³ -59° (H₂O). Crystd from de-ionized H₂O, solubility in H₂O is 1g/80g and dried under vacuum over Mg(ClO₄)₂. The dry red crystals are *hygroscopic* and can absorb ~12% of H₂O. A soln at pH 4.5-5 can be autoclaved for 20min at 120° without dec. Aqueous solns are stabilised by addition of (NH₄)₂SO₄. [Golding *Comprehensive Organic Chem* Vol 5 (Ed. Haslam; Pergamon Press, NY, 1979) pp 549-584.]

Alternatively an aqueous soln of the coenzyme was concentrated, if necessary in a vacuum at 25° or less, until the concentration was 0.005 to 0.01M (as estimated by the OD at 522 nm of an aliquot diluted with 0.01M K-phosphate buffer pH 7.0). If crystals begin to form on the walls of the container they should be re-dissolved with a little H₂O. The concentrated soln is placed in a glass stoppered flask and diluted with 5vols of Me₂CO. After 2-3h at 3° it is centrifuged (10,000xg/10 min) in Me₂CO-insol plastic tubes to remove some amorphous ppte. The clear supernatant is inoculated with a small crystal of the vitamin and allowed to crystallise overnight at 3°. Crystals are formed on the walls and the bottom of the container. A further 2vols of Me₂CO are added and set aside at 3° to further crystallise. Crystallisation is followed by observing the OD₅₂₂ of the supernatant. When the OD falls to 0.27 then ca 94% of the crystals have separated. The supernatant is decanted (saved for obtaining a second crop) and the crystals are washed with a little cold 90% aqueous Me₂CO (2 x), 100% Me₂CO (2 x), Et₂O (2 x) at which time the crystals separated from the glass walls. Allow to settle and remove residual Et₂O with a stream of dry N₂. The process can be repeated if necessary. The crystals can be dried in air or in a vacuum for 2h over silica gel at 100° with an 8-9% weight loss. [Barker et al. *Biochem Prep* 10 33 1963.] This material gives a single spot of paper chromatography [see Weissbach et al. *J Biol Chem* 235 1462 1960.] The vitamin is soluble in H₂O (16.4mM at 24°, 6.4mM at 1°), in EtOH and PhOH but insol in Me₂CO, Et₂O, CH₂Cl₂ and dioxane. UV: λ_{max} 260, 375 and 522nm (ε 34.7 x 10⁶, 10.9 x 10⁶ and 8.0 x 10⁶ / mole) in H₂O. The dry crystals are stable for months in the dark, but aqueous solns decompose on exposure to VIS or UV light or alkaline CN⁻, but stable in the dark at pH 6-7. The vitamin is inactivated by strong acids or alkalis. [Barker et al. *J Biol Chem* 235 480 1960; see also *Vitamin B₁₂* (Zagalak and Friedrich Eds) W de Gruyter, Berlin 1979.]

Vitamin C see **ascorbic acid** entry on p. 116 in Chapter 4.

Vitamin D₂ [50-14-6] M 396.7, m 114-116°, $[\alpha]_{546}^{20} +122^\circ$ (c 4, EtOH) Converted into their 3,5-dinitrobenzoyl esters, and crystd repeatedly from acetone. The esters were then saponified and the free vitamins were isolated. [Laughland and Phillips *Anal Chem* 28 817 1956.]

Vitamin D₃ [67-97-0] 384.6, m 83-85°, $[\alpha]_{546}^{20} +126^\circ$ (c 2, EtOH). Converted into their 3,5-dinitrobenzoyl esters, and crystd repeatedly from acetone. The esters were then saponified and the free vitamins were isolated. [Laughland and Phillips *Anal Chem* 28 817 1956.]

Vitamin E (2*R*,4'*R*,8'*R*- α -tocopherol, natural active isomer) [59-02-9] M 430.7, m 2.5-3.5°, b 200-220°/0.1mm, 200°/0.005mm, $d_4^{25} 0.950$, $n_D^{25} 1.5045$, $[\alpha]_D^{25} +3.58^\circ$ (c 1.1, *C₆H₆). Viscous yellow oil which is distd at high vacuum. It has λ_{\max} 294nm ($E_{1\text{cm}}^{1\%} 71$). It is oxygen and light sensitive and is best stored as its stable acetate which is purified by evaporative distn at b 180-200°(bath temp)/0.7mm, $[\alpha]_D^{25} +3.3^\circ$ (c 5.1, EtOH). [NMR: Cohen et al. *Helv Chim Acta* 64 1158 1981; Burton and Ingold *Acc Chem Res* 19 194 1986; Karrer et al. *Helv Chim Acta* 21 520 1938.]

Vitamin E acetate (DL- α -tocopheryl acetate) [7695-91-2] M 472.8, m -27.5°, b 194-196°/0.01mm, 222-224°/0.3mm, $d_4^{20} 0.958$, $n_D^{20} 1.4958$. It is a viscous liquid which is purified by distn under high vacuum in an inert atm and stored in sealed ampoules in the dark. It is considerably more stable to light and air than the parent unacetylated vitamin. It is insoluble in H₂O but freely soluble in organic solvents. All eight stereoisomers have been synthesised. The commercially pure *d*- α -tocopheryl acetate (2*R*,4'*R*,8'*R*) has b 180-200°/0.7mm and $[\alpha]_D^{20} +3.9^\circ$ (c 5, EtOH). [Cohen et al. *Helv Chim Acta* 64 1158 1981.]

Vitamin K₁ (2-methyl-3-phytyl-1,4-naphthoquinone) [84-80-0] M 450.7, m -20°, b 141-140/0.001mm, b 140-145°/10⁻³ mm, $d_{25}^{25} 0.967$, $n_D^{25} 1.527$, $[\alpha]_D^{20} -0.4^\circ$ (c 57.5, *C₆H₆). Yellow viscous oil, which can be distd at high vacuum practically unchanged. Insoluble in H₂O, but soluble in common organic solvents. Store in the dark under N₂, oxygen sensitive. $A_{1\text{cm}}^{1\%}$ 328 at 248nm. [*J Am Chem Soc* 61 2557 1939, 76 4592 1954; *Helv Chim Acta* 27 225 1954.]

Vitamin K₃ (2-methyl-1,4-naphthoquinone, Menadione, Menaphthone) [58-27-5] M 172.2, m 105-106°, 105-107°. Recrystd from 95% EtOH, or MeOH after filtration. Bright yellow crystals which are decomposed by light. Solubility in EtOH is 1.7% and in *C₆H₆ it is 10%. It IRRITATES the mucous membranes and skin. [Fieser *J Biol Chem* 133 391 1940.]

Xanthine (2,6-dihydroxypurine, purine-2,6(1*H*,3*H*)dione) [69-89-6] M 152.1, pK₁ 0.8 [protonation of imidazole 7(9)NH], pK₂ 7.44 [monoanion 1(3)NH], pK₃ 11.12 [dianion 1,3-N²⁻]. The monohydrate separates in a microcryst form on slow acidification with acetic acid of a solution of xanthine in dil NaOH. Also ppted by addition of conc NH₃ to its soln in hot 2*N* HCl (charcoal). After washing with H₂O and EtOH, it is dehydrated on heating above 125°. Sol in H₂O is 1 in 14,000 at 16° and 1 in 1,500 and separates as plates from boiling H₂O. It has no m, but the perchlorate has m 262-264° [Lister *Heterocyclic Compounds, Fused Pyrimidines—Purines Part II*, Ed. Brown, J. Wiley & Sons, 1971].

Xanthopterin monohydrate (2-amino-4,6-dihydroxypteridine, 2-amino-pteridin-4,6(1*H*,5*H*)-dione) [5979-01-1 (H₂O), 119-48-8 (anhydr)] M 197.2, m <300°, pK₁ 1.6 (basic), pK₂ 6.59 (acidic), pK₃ 9.31 (acidic)(anhydrous species), and pK₁ 1.6 (basic), pK₂ 8.65 (acidic), pK₃ 9.99 (acidic)(7,8-hydrated species). Purification as for isoxanthopterin. Crystd by acidifying an ammoniacal soln, and collecting by centrifugation followed by washing with EtOH, ether and drying at 100° *in vacuo*. Paper chromatography R_F 0.15 (*n*-PrOH, 1% aq NH₃, 2:1), 0.36 (*n*-BuOH, AcOH, H₂O, 4:1:1) and 0.47 (3% aq NH₃). [Inoue and Perrin *J Chem Soc* 260 1962; Inoue *Tetrahedron* 20 243 1964; see also Blakley *Biochemistry of Folic Acid and Related Pteridines* North Holland Publ Co, Amsterdam 1969.]

Xanthotoxin (Methoxalen, 9-methoxyfuro[3,2-g][1]benzopyran-7-one) [298-81-7] **M** 216.2, **m** 146-148°, 148°, 148-149°. Purified by recrystn from *C₆H₆-pet ether (b 60-80°) as silky needles, EtOH-Et₂O as rhombic prisms or hot H₂O as needles. It is soluble in aqueous alkali due to ring opening of a lactone but recyclises upon acidification. It has UV λ_{\max} in EtOH at 219, 249 and 300nm (log ϵ 4.32, 4.35 and 4.06) and ¹H NMR in CDCl₃ with δ at 7.76 (d, 1H, *J* 10 Hz), 7.71 (d, 1H, *J* 2.5 Hz), 7.38 (s, 1H), 6.84 (d, 1H, *J* 2.5 Hz), 6.39 (d, 1H, *J* 10 Hz) and 4.28 (s, 3H). [Nore and Honkanen *J Heterocycl Chem* 17 985 1980.] It is a DNA intercalator and is used in the treatment of dermal diseases [Tessman et al. *Biochemistry* 24 1669 1985.]

Xylanase (from *Streptomyces lividans*) [37278-89-0] **M_r** 43,000 [EC 3.2.1.8]. Purified by anion-exchange chromatography on an Accell QMA column and finally by HPLC using a ProteinPak DEAE 5PW anion-exchange column. Solutions were stored frozen at -70°. [Morosoli et al. *Biochem J* 239 587 1986; Wong et al. *Microbiol Rev* 52 305 1988.]

Zeatin (*trans-N*⁶-[4-hydroxy-3-methylbut-2-en-1-yl]adenine) [1637-39-4] **M** 219.3, **m** 207-208, 209-209.5°, **pK₁** 4.4 (basic), **pK₂** 9.8 (acidic). Purified by recrystn from EtOH or H₂O. The UV has λ_{\max} at 207 and 275nm (ϵ 1400 and 14650) in 0.1N aqueous HCl; 212 and 270nm (ϵ 17050 and 16150) in aqueous buffer pH 7.2; 220 and 276nm (ϵ 15900 and 14650) in 0.1N aq NaOH. The *picrate* has **m** 192-194° (from H₂O) from which zeatin can be recovered by treatment with Dowex 1 x 8 (200-400 mesh, OH⁻ form). [Letham et al. *Aust J Chem* 22 205 1969; *Proc Chem Soc (London)* 230 1964; Shaw and Wilson *Proc Chem Soc (London)* 231 1964.] It is a cell division factor (plant growth regulator) [Letham and Palni *Ann Rev Plant Physiol* 34 163 1983] and inhibits mitochondrial function [Miller *Plant Physiol* 69 1274 1982]. Its *9-riboside* is a cytokine [McDonald and Morris *Methods Enzymol* 100 347 1985].

General Index

For individual organic chemicals, listed alphabetically, see Chapter 4, beginning on Page 80; or inorganic and metal-organic chemicals see Chapter 5, beginning on Page 389; and for biochemicals and related products see Chapter 6, beginning on Page 500. It is much faster to use the CAS Registry Numbers Index on p 585 for locating specific compounds irrespective of which chapter they are in.

<u>Index terms</u>	<u>Links</u>
A	
Abbreviations	30 80 389 505
Abderhalden pistol	25
ACES, <i>N</i> -(2-acetamido)-2 aminoethanesulfonic acid	82
Acetamido TEMPO, 4-acetamido-2,2,6,6-tetramethylpiperidine-1-oxyl	83
ADA, <i>N</i> -(2-acetamido)-iminodiacetic acid	82
ADP, adenosine-5'-diphosphate	508
Adrenor	553
Adsorbents, graded and solvents	40
Affinity chromatography	25 501
AIBN, α,α' -azobis(isobutyronitrile)	117
AICAR HCl, 4-Aminoimidazole-5-carboxamide HCl	106
Alkaline base bath	3
<i>n</i> -Alkyltrimethylammonium bromides	99
<i>n</i> -Alkylammonium chlorides	99
Alumina, Preparation	19
Standardization	19
α -Amino acids	511
3'-AMP, Adenosine-3'-monophosphoric acid	508
5'-AMP, Adenosine-5'-monophosphoric acid	509
Anhydron	5 438
Anion exchange resin	21 22 102 513
ANTU, 1-naphthyl thiourea	307
Apiezon L, M and T	11
Aqua regia	396
Arterenol	553
Aspirin, acetylsalicylic acid	92
ATP, Adenosine-5'-triphosphate	509

Index terms**Links****B**

BAL, 1,2-dimercapto-3-propanol	209
BAO, 2,5-bis-(4-aminophenyl)-1,3,4-oxadiazole	133
BBOD, 2,5-bis(4-biphenyl)-1,3,4-oxadiazole	516
9BBN, 9-Borabicyclo-[3.3.1]nonane	403
Benzene, care in use of	80 389
purification	505
	119
BES, <i>N,N</i> -bis-(2-hydroxyethyl)-2-aminoethanesulfonic acid	134
BH ₄ HCl, 6 <i>R</i> -tetrahydro- <i>erythro</i> -biopterin HCl	569
BHT, 2,6-di- <i>tert</i> -butyl- <i>p</i> -cresol, butylated-hydroxytoluene	194
HBTU, <i>O</i> -Benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyl-uronium hexafluoro-phosphate	124
BICINE, <i>N,N</i> -bis-(2-hydroxyethyl)-glycine	208
BINAP, 2,2'-bis-(diphenyl-phosphino)-1,1'-binaphthyl	401
BIS-TRIS, bis-(2-hydroxyethyl)amino-tris-(hydroxymethyl)-methane	134
Boiling point(s), effect of pressure	32 33
	34
of useful gases	44
nomogram	34
BSA, <i>N,O</i> -bis-(trimethylsilyl)-acetamide	403
BMS, bis(2-mercaptoethyl)sulfone	516
Büchner funnel	14
Buffers, aqueous	31

C

Cadion, [1-(4-nitrophenyl)-3-(4-phenylazophenyl)-triazene]	406
Calcein, Fluorescein Complexon	406
Calcium hydride, in purification	55
Carbon Black	156
Catapres, Clonidine hydrochloride	521
Cation exchange resins	157 519
Cbz-Cl, benzyloxycarbonyl chloride	130
Celite	14
Cellulose, preparation	21
Centrifugation	15 500
Charcoal, preparation	20
purification	159
Chelex	100 159

Index terms**Links**

Chemical hazards,	
in purification	4
with perchlorates	5
with perchloric acid	5
with peroxides	5
with heavy-metal-containing-explosives	5
with strong acids	5
with anhydrides	5
with reactive halides	5
with solvents	5
with salts	6
Chemokines	520
Chirazymes	520
Chloramine-T, <i>N</i> -chloro- <i>p</i> -toluenesulfonamide sodium salt	410
Chloranil test	76
Chloroplatinic acid	427
CHIRAPHOS, 2,3-bis-(diphenylphosphino)-butane	401
Chromatography,	17
adsorption	18
affinity	25 501
biospecific	25
charge transfer adsorption	25
column	19
covalent	25
flash	21
gas	17
GC-MS	17
hydrophobic	25
ion exchange	21
LC-MS	25
liquid	18
mass spectrometry	17
metal chelate adsorption	25
paired-ion	21
paper	19
spot detectn in TLC	39
stationary phases for GLC	39
thin layer (TLC)	18
vapour phase	17
Chromatotron	19

Index terms**Links**

Chromic acid bath	3
Chromotropic acid di-Na salt, di-Na 4,5(1,8)-dihydroxynaphthalene-2,7(3,6)-disulfonate	413 421
Cisplatin, <i>cis</i> -diammine-dichloroplatinum(II)	417
Citiolone, <i>N</i> -acetyl- <i>RS</i> -homocysteine thiolactone	90
Citric acid cycle components	521
Claisen alkali	413
Claisen flask	10
Cleaning, practices	3
of glassware	3
chromic acid	14
Cleland's reagent, <i>RS</i> -1,4-Dithiothreitol	227
Column packings	11
balls or beads	11
helices	11
rings	11
Combinatorial chemistry	75
Condensers	11
double surface	11
coil	11
Friedrichs	11
Liebig	11
from the melt	16
Cooling baths	38
CTAB, hexadecyltri-methylammonium bromide	158
Cupferon, (<i>N</i> -nitroso- <i>N</i> -phenylhydroxylamine) NH ₄ salt	414
Cytokines	527

D

2,4-D, 2,4-dichlorophenoxyacetic acid	200
DABCO, 1,4-diazabicyclo-[2.2.2]-octane	191
DABITC, 4- <i>N,N'</i> -dimethylaminoazobenzene-4'-isothiocyanate	76 212
Danishesky's diene, <i>trans</i> -1-methoxy-3-(trimethylsilyloxy)-1,3-butadiene	441
DATD, (+)- <i>N,N'</i> -diallyl tartramide	185
Davy's reagent, 2,4-bis-(methylthio)-1,3,2λ ⁵ ,5,4λ ⁵ -dithiadiphosphetane-2,4-dithione	402
DEED, dibenzylethylenediamine	192
DBN, 1,5-diazabicyclo-[4.3.0]non-5-ene	190
DBU, 1,8-Diazabicyclo-[5.4.0]undec-7-ene	191
DCC, 1,3-dicyclohexylcarbodiimide	201
<i>o,p'</i> -ODD, 1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane	166

Index terms**Links**

<i>p,p'</i> -DDD, 2,2-bis-(4-chlorophenyl)-1,1-dichloroethane	134
DDQ, 2,3-dichloro-5,6-dicyano- <i>p</i> -benzoquinine	197
<i>p,p'</i> -DDT, 2,2-bis-(4-chlorophenyl)-1,1,1-trichloroethane	134
DEAD, diethyl azodicarboxylate	74 202
Dean-Stark apparatus	58
DEC, 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride	74 212
Dehydrite	5 438
Demecocine, colcemide	522
Deoxyribonucleic acid (DNA)	504 528
	532
DEP, diethyl pyrocarbonate	205
Desiccants	26
alumina	27
aluminium amalgam	27
barium oxide	27
barium perchlorate	27
boric anhydride	27
calcium chloride	27
calcium hydride	27
calcium oxide	27
calcium sulfate	27
capacity	26
common	26
copper (II) sulfate	27
lithium aluminium hydride	27
magnesium amalgam	27
magnesium perchlorate	5 27
	438
magnesium sulfate	27
molecular sieves	28
phosphorus pentoxide	27
potassium	27
potassium carbonate	28
silica gel	28
sodium	28
sodium hydroxide	28
sodium-potassium alloy	28
sodium sulfate	28
sulfuric acid	28
DHFA, 7,8-dihydrofolic acid	529

Index terms**Links**

Dialysis	502
Di- and tri-carboxylic acids	529
Diatomaceous earth, preparation	20
Diazald	296
3,4-Dihydroxyphenylalanine-containing proteins	530
DIPHOS, 1,2-bis-(diphenyl-phosphino)ethane	402
Distillation	8
azeotropic	13
columns	11
fractional	10
Isopiestic or isothermal	13
Kügelrohr	13
of liquid mixtures	9
spinning-band	12
steam	12
techniques of	9
types of	10
vacuum	11
Distilling flask,	10
Dithizone, diphenylthiocarbazone	227
Diuron, 3-(3,4-dichlorophenyl)-1,1-dimethyl urea	200
DMAP, 4-dimethylaminopyridine	213
DMAP-PS, 4-dimethylaminopyridine on polystyrene support	213
DMF, <i>N,N</i> -dimethylformamide	215
DMPSCl, dimethylphenylsilyl chloride	449
DMSO, dimethyl sulfoxide	219
DMT, 4,4'-dimethoxytrityl chloride	211
DNA, deoxyribonucleic acids, purification	504 528
	532 560
DNPH, 2,4-dinitrophenylhydrazine	119 222
	291
DOPA, EUODOPA, <i>S</i> - β -(3,4-dihydroxyphenyl)alanine	530
2,4-DP, α -(2,4-dichlorophenoxy)propionic acid	200
Drying	25
removal of water	26
liquids for pistols	41
of solvents	25 43
Drying agents for organic compounds	43
DTBCl ₂ , di- <i>ter</i> -butyldichlorosilane	417

Index terms**Links**

DTE, 1,4-dithioerythritol	227
DTPA, diethylenetriamine-pentaacetic acid	203
DTT, <i>R,S</i> -1,4-Dithiothreitol	227
E	
Ebullioscopic Constant	10
EDCI, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide HCl	212
EDDA, <i>N,N</i> -ethylenediaminediacetic acid	237
EDTA, ethylenediaminetetraacetic acid	237
EDTA di-Na salt	421
EDTA Di-Na Mg salt	421
EEDQ, 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline	233
EGTA, ethylene glycol bis(β -aminoethylether)- <i>N,N</i> -tetraacetic acid	238
Ehrlich's Reagent, <i>p</i> -dimethylaminobenzaldehyde	212
Ellman's reagent	76
Enzymes,	
[EC 1.1.1.5]	505
[EC 1.1.1.8]	539
[EC 1.1.1.30]	542
[EC 1.2.1.12]	538
[EC 1.1.3.4]	537
[EC 1.5.1.3]	529
[EC 1.6.99.7]	529
[EC 1.9.3.1]	526
[EC 1.10.2.2]	572
[EC 1.14.17.1]	532
[EC 1.15.1.1]	523
[EC 2.4.1.11]	540
[EC 2.5.1.18]	538
[EC 2.7.1.30]	538
[EC 2.7.1.40]	563
[EC 2.7.6.1]	559
[EC 2.7.7.49]	564
[EC 2.7.7.9]	573
[EC 3.1.1.34]	546
[EC 3.1.27.5]	565
[EC 3.1.3.1]	557
[EC 3.1.3.5]	553
[EC 3.2.1.8]	577
[EC 3.2.1.23]	537

Index terms**Links**Enzymes (*Continued*)

[EC 3.2.1.28]	572
[EC 3.2.1.35]	541
[EC 3.4.11.10]	531
[EC 3.4.17.1]	518
[EC 3.4.21.1]	521
[EC 3.4.21.31]	573
[EC 3.4.22.1]	519
[EC 3.4.22.2]	556
[EC 3.4.22.8]	521
[EC 3.4.23.1]	557
[EC 3.4.23.5]	519
[EC 3.4.24.3]	523
[EC 3.5.4.4]	563
[EC 4.2.1.1]	518
[EC 6.2.1.1]	508
[EC 6.2.1.2]	508
Ethanol	231
Exonucleases	533

F

FA, folic acid	536
FAD, flavin adenine dinucleotide	534
Fenske rings	11
Filters, micro	36
Whatman	35
Filtration	14
Florisil	14
Fluorous chemistry	77
Fluram, fluorescamine	535
FMOC-Cl, 9-fluorenylmethyl chloroformate	243
FMN, flavin mononucleotide	535
Fractogels	23
Friedrichs condenser	11
Freeze-pump-thaw and purging	29
Freon 12, dichlorodifluoromethane	198
Freon 13B1, bromotrifluoromethane	142
Fremy's Salt, potassium nitrosodisulfonate	456
FSH, follicle stimulating hormone	536

Index terms**Links**

Fuschin, rosaniline HCl	349
Fullerene	246
G	
GABA, 4-aminobutyric acid	104
Galvinoxyl	250
GDH, L-glycerol-3-phosphate dehydrogenase	539
Glass powder	426
Gangcyclovir	537
Gel electrophoresis	502
Gel filtration	500
Girard Reagent T	251
Glovebox techniques	30
Graded adsorbents	19
H	
Haematoporphyrin IX	427 541
Hagemann's ester	156
Heating baths	35
HEDTA, <i>N</i> -[2-hydroxyethyl]ethylenediaminetriacetic acid	261
Heimgartner's reagent, 2,4-bis-(<i>p</i> -tolylthio)-1,3,2λ ⁵ ,4λ ⁵ -dithiadiphosphetane-2,4-dithione	403
HEPES, <i>N</i> -2-hydroxyethyl-piperazine- <i>N'</i> -2-ethane-sulfonic acid	54 261
HIMDA, <i>N</i> -hydroxyethyliminodiacetic acid	261
Histones	541
HMDS, hexamethyldisilazane	428
HMPA, hexamethyl-phosphoric triamide	428
HMPT, hexamethyl-phosphorous triamide	428
HPLC, High performance liquid chromatography	18 24
gel filtration	24
paired-ion	24
reverse-phase	24
HMTA, hexamethylenetetramine	257
HOBt, 1-hydroxybenzotriazole	74 260
HPTLC, high performance thin layer chromatography	18
Hydroxylapatite	24 501
Hyflo-supercel	14

Index terms**Links****I**

Impurities, sources of	2
Ionic liquids for reactions	77
Ion exchange, bead form	41
celluloses	22 40
Cellex D	24
Cellex CM	23
matrices	501
resins	21 40
Sephadex	22 23
Sepharose	23
Ionization constants	7 80 389 513
Immiscible solvents	30
Isoelectric focusing	501
Interferons	543
Interleukins	543 544
Isothermal crystallization	14

J

Juglone, 2-hydroxy-1,4-naphthaquinone	264
---------------------------------------	-----

K

Kaiser test	76
Koser's reagent	266
KPEP, phosphoenolpyruvic acid K salt	558

L

Lactoferrin	545
Lanthanide (Europium) shift reagents	277
Lectins	545
Liebig condenser	11
Ligroin	16
Lipoproteins	546
Lithium aluminium hydride, in purification	55
Lithium hydride	436

M

McLeod type guage	12
Magon, Xylidyl Blue	438
Martius Yellow, 2,4-dinitro-1-naphthol	221
MCPA, 4-chloro-2-methylphenoxyacetic acid	164
Meglutol, 3-hydroxy-3-methylglutaric acid	263
Mega-9	553
MEK, 2-butanone, methyl ethyl ketone	144
MEMCl, 2-methoxyethoxymethylchloride	285
Mercurochrome, mercury dibromofluorescein	440
MES, 2-(<i>N</i> -morpholino)-ethanesulfonic acid	304
MESNA, Na 2-mercaptoethanesulfonate	473
Metal impurities,	
by complexation	53
by ion exchange resins	54
by precipitation	54
removal by extraction	54
removal from reagents	53
removal of lead	54
removal of iron	54
Methylarenes	289
Methyldopa, 3-(3,4-dihydroxyphenyl)-2-methyl-L-alanine	530
Methyl Yellow, <i>p</i> -dimethylaminoazobenzene	212
Mixed solvents	16
MMPP, monoperoxyphthalic acid magnesium salt	443
MOMCl, chloromethyl methyl ether	164
MONO-TRIS, 2-hydroxy-ethylimino-tris-(hydroxymethyl)-methane	261
MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine HCl	547
MTAD, 4-methyl-1,2,4-triazoline-3,5-dione	302
MTPA, α -methoxy- α -trifluoromethyl-phenylacetic acid	286
MTT, Thiazolyl blue tetrazolium bromide	570
Muscimol, 5-aminoethyl-3[2 <i>h</i>]-isoxazolone	549
N	
NAD (DPN), β -nicotinamide adenine dinucleotide	551
NADH, reduced NAD	551
NADP, β -nicotinamide adenine dinucleotide phosphate	551
NADPH, reduced NADP	552
Naloxone HCl, 1- <i>N</i> -propenyl-7,8-dihydro-14-hydroxy morphinan-6-one HCl	550

Index terms**Links**

Naltrexone HCl, 1- <i>N</i> -cyclopropylmethyl-7,8-dihydro-14-hydroxymorphinan-6-one HCl	550
NAMA, <i>N</i> -acetyl muramic acid	507
NANA, <i>N</i> -acetyl neuraminic acid	507
NAM, <i>N</i> -(9-acridinyl)-maleimide	94
NaPAA, sodium polyacrylate	474
<i>cis</i> -, <i>trans</i> -, <i>cis/trans</i> -nerolidol	308
Nitromethane	311
NMN, β -nicotinamide mononucleotide	552
NPS-Cl, 2-Nitrobenzenesulfonyl chloride	309
NTA, Nitrilotriacetic acid	308
Nucleotide thiophosphates	553
Nylon powder	316
O	
Orcine	554
Ouabain	318
P	
PAGE, polyacrylamide gel electrophoresis	502
PAN, 1-(2-pyridylazo)-2-naphthol	345
Paper chromatography	19
PAR, 4-(2-pyridylazo)-resorcinol	345
Paraffin wax	319
Partial precipitation	502
PEGA, polyethyleneglycol acrylamide	74
Petroleum ether	324
Petroleum ethers	15
Petroleum spirit	15
pK	7
pKa and pKb	8
and temperature	8
and solvent	8
PFBHA, <i>O</i> -(2,3,4,5,6-pentafluorobenzyl)-hydroxylamine HCl	321
Phospholipids	558
Phosphoproteins	559
PIC, paired-ion chromatography	21
PIPES, piperazine- <i>N,N</i> -bis(2-ethanesulfonic acid)	336
PIPES-Na salt	474

<u>Index terms</u>	<u>Links</u>
Pituitary Growth Factor	560
Pixyl chloride, 9-chloro-9-phenyl-xanthene	167
Plasmids	560
PLP, pyridoxal-5'-phosphate	563
PMSF, phenylmethanesulfonyl fluoride	557
PNBP, 4-(4-nitrobenzyl)-pyridine	76 310
Podbielniak column	141 168 270 481 489
Polyethylene glycol	540
Polymer support reactants	74
Polypeptides, purification	503 560
Potassium borohydride, in purification	56
PPD, 2,5-diphenyl-1,3,4-oxadiazole	225
PPO, 2,5-Diphenyloxazole	225
PQQ, methoxantin coenzyme	547
Prefixes for quantities	44
Prefolic A, 5-methyltetrahydrofolic acid	548
Preparative GC	17
Proteins, purification	503 562
Proteoglycans	562
Proton ionophore, hydrogen ionophore, 4-nonadecylpyridine	430
Proton sponge, 1,8-bis-(dimethylamino)naphthalene	134
Provitamin B, D-panthenol	555
PTAD, 4-phenyl-1,2,4-triazoline-3,5-dione	333
PUGNAC, <i>Z</i> - <i>O</i> -(2-acetamido-2-deoxy-D-glycopyranosylideneamino)- <i>N</i> -phenylcarbamate	82
Purpald, 4-amino-3-hydrazino-5-mercapto-1,2,4-triazole	76 106
Purification <i>via</i> derivatives	56
of alcohols	56
of aldehydes	57
of amines	57
of aromatic hydrocarbons	58
of carboxylic acids	58
of hydroperoxides	59
of ketones	59
of phenols	59
of phosphate esters	60
of phosphonate esters	60

Index terms**Links**

Purification, general	
procedures for: acetals	61
acids	61
carboxylic	61
sulfonic	62
sulfinic	62
acid chlorides	62
alcohols	62
monohydric	62
polyhydric	63
aldehydes	63
amides	63
amines	63
amino acids	64
anhydrides	64
carotenoids	64
esters	64
ethers	65
halides	65
hydrocarbons	66
imides	67
imino compounds	67
ketones macromolecules	503 504
nitriles	67
nitro compounds	67
nucleic acids	504
phenols	68
polypeptides	503
proteins	503
quinones	68
salts (organic)	68
with metal ions	68
with organic cations	68
with Na alkane sulfonates	68
sulfur compounds	68
disulfides	68
sulfones	68
sulfoxides	69
thioethers	69
thiols	69

Purification, general (<i>Continued</i>)	
thiolsulfonates (disulfoxides)	69
Purity, criteria of	61
question of	1
R	
Rapid purification	80
of benzen	120
of tert-butyl alcohol	145
of carbon tetrachloride	157
of chloroform	163
of cyclohexane	178
of dichloromethane	199
of diethyl ether	204
of <i>N,N</i> -dimethylformamide	216
of dimethylsulfoxide	219
of 1,4-dioxane	223
of ethanol	232
of ethyl acetate	234
of <i>n</i> -hexane	257
of methanol	284
of petroleum ether	324
of tetrahydrofuran	361
of toluene	370
Raschig rings	11
Reactive groups on resins, detection of	
aldehyde groups	76
carboxy groups	76
amino groups	76
thiol groups	76
Fmoc assay	76
RDX, cyclotrimethylene-trinitramine	183
Reichstein's G, adrenosterone	98
Recrystallisation	14
choice of solvents	15
from the melt	16
techniques	14
Reverse-phase thin layer chromatography	19

Index terms**Links**

Resin support(s)	74	
choice for SPOS	74	
reactants	74	
scavenger	74	
Restriction enzymes	504	533
endonucleases	564	
Riboflavin	564	
RNA, ribonucleic acids	565	
S		
Saccharides	566	
Safety precautions	4	
Safety disclaimer	6	
Salcomine, <i>N,N'</i> -bis-(salicylidene)ethylene-diamine cobalt (II)	402	
Salol, phenyl salicylate	332	
Salts, vapour press with satd solns	42	
SAM-HCl, (-)-S-adenosyl-L-methionine chloride	509	
Sanger's reagent, 2,4-dinitrofluorobenzene	221	
Scavanger resins	74	
Schlenk techniques	30	
Schlippe's salt	476	
SDS, sodium dodecylsulfate	470	
Selectophore, bis-[4-(1,1,3,3-tetramethylbutyl)phenyl]-phosphate Ca salt	403	
SEMCl, 2-(trimethylsilyl)ethoxymethyl chloride	491	
Sicapent	27	
Silica, preparation	20	
Silylation, of glassware	3	
of plasticware	3	
with REPEL-SILANE ES	4	
Skellysolves A-L	351	
Sodium <i>n</i> -alkylsulfates	465	
Sodium borohydride, in purification	55	
Sodium silicate solution	475	
Solid phase reaction monitoring by:		
infrared analysis of resin	75	
qualitative analysis of resin	75	
quantitative analysis of resin	75	

<u>Index terms</u>	<u>Links</u>
Solid phase synthesis (SPS)	72
of deoxyribo-nucleotides	73
of oligosaccharides	73
of organic compounds (SPOS)	72 73
of peptides (SPPS)	73
Solvent(s)	5 6
distribution	6
extraction	6
for crystn	37
immiscible	30
miscible	37
removal of acids from	6
removal of phenols from	6
removal of organic bases from	6
Solvents, mixed	16
Soxhlet extraction	14
Stedman column	153 240 419 441 442 489
Sublimation	17
Sulfuric acid (cone) for	
removal of unsaturated hydrocarbons	6
removal of alcohols	6
removal of ethers	6
 T	
2,4,5-T, 2,4,5-trichlorophenoxyacetic acid	375
TBD, 1,5,7-triazabicyclo-[4.4.0]dec-5-ene	372
TBDPSCI, <i>tert</i> -butyldiphenylchlorosilane	405
TBHP, <i>tert</i> -butyl hydroperoxide	148
TCT, cyanuric chloride	76 176
TED, triethylenediamine	191
TEHP, (±)-tris-(2-ethylhexyl)phosphate	494
TEMED, <i>N,N,N',N'</i> -tetramethylethylene-diamine	364
TEMPO, 2,2,6,6-Tetramethylpiperidinyl-1-oxy	365
TES, <i>N</i> -tris-(hydroxy methyl)methyl-2-aminomethanesulfonic acid	54 363
THFA, tetrahydrofolic acid	569
TIRON, 1,2-Dihydroxybenzene-3,5-di-sulfonic acid, di-Na salt	419
Thiram, bis-(dimethylthiocarbamyl)disulfide	134

Index terms**Links**

TLG, thin layer gel filtration	18
TMEDA, <i>N,N,N',N'</i> -tetramethyl-ethylene-diamine	364
TMS, tetramethylsilane	482
TMSAN, trimethylsilyl acetonitrile	490
TMSF, fluorotrimethylsilane	425
TMSI, trimethylsilyl iodide	433
TNBS, 2,4,6-trinitrobenzenesulfonic acid	76 380
TNT, 2,4,6-trinitrotoluene	380
Todd column	174 183
	279 346
	442 485
	492
TOSMIC, tosylmethyl isocyanide	372
Toyoparl	23
TPAP, Tetra- <i>n</i> -propyl-ammonium perruthenate	483
TPP, 5,10,15,20-tetraphenylporphyrin	366
TREN, tris-(2-aminoethyl)-amine	376 381
TRICINE, <i>N</i> -tris-(hydroxymethyl)methylglycine	382
TRIEN (TETA), triethylene tetramine	376
TRIEN.HCl, triethylenetetramine 4HCl	376
TRIS, tris-(hydroxymethyl)-methylamine	54 381
TRIS-HCl, tris-(hydroxymethyl)methylamine HCl	382
Triton B, benzyltrimethyl-ammonium hydroxide	131
TTC, 2,3,5-triphenyltetrazolium chloride	381
TTMSS, tris-(trimethylsilyl)silane	495
U	
Ubiquinone, Coenzyme Q	522
Uracil, uridine and uridine nucleotides	572
Ultrafiltration	502
Ultragels	23
Uridine 5'-(1-thio) monophosphate	573
V	
Vacuum-lines	30
Vigreux column	11
Vitamins	574

Index terms**Links****W**Wanzlick's reagent, *N,N'*-diphenylethylene-diamine

225

Widmer Column

87 147

181 376

480

Wilkinson's catalyst

412

X

Xanthotoxin

577

Y

Yohimbine

388

Z

Zone refining

16

50-00-0	245	53-16-7	231	57-47-6	231	60-18-4	383
50-01-1	253	53-19-0	166	57-48-7	246	60-19-5	383
50-02-2	187	53-41-8	113	57-50-1	354	60-20-8	540
50-03-3	541	53-59-8	551	57-53-4	282	60-23-1	525
50-04-4	523	53-70-3	191	57-55-6	338	60-24-2	282
50-07-7	549	53-84-9	551	57-57-8	339	60-27-5	524
50-14-6	576	53-96-3	82	57-62-5	513	60-29-7	204
50-22-6	523	54-06-8	98	57-63-6	533	60-31-1	506
50-23-7	541	54-11-5	552	57-67-0	354	60-34-4	293
50-27-1	231	54-28-4	571	57-68-1	354	60-35-5	81
50-28-2	533	54-43-0	186	57-83-0	338	60-46-8	212
50-29-3	134	54-47-7	563	57-85-2	356	60-54-8	569
50-30-6	197	54-62-6	512	57-87-4	230	60-56-0	282
50-32-8	123	54-64-8	470	57-88-5	170	60-80-0	115
50-33-9	328	54-85-3	272	57-91-0	317	60-81-1	333
50-36-2	521	54-92-2	273	57-94-3	383	60-82-2	333
50-50-0	317	54-95-5	303	57-97-6	213	61-25-6	556
50-55-5	348	54-96-6	190	58-89-9	255	61-54-1	382
50-56-6	554	55-21-0	119	58-00-4	513	61-72-3	281
50-69-1	349	55-22-1	272	58-08-2	152	61-73-4	292
50-70-4	352	55-55-0	288	58-15-1	110	61-78-9	106
50-71-5	100	56-03-1	132	58-18-4	301	61-78-2	132
50-76-0	508	56-04-2	302	58-22-0	356	61-82-5	111
50-78-2	92	56-06-4	189	58-27-5	576	61-90-5	278
50-79-3	197	56-10-0	106	58-46-8	356	62-13-5	510
50-81-7	116	56-12-2	104	58-55-9	366	62-23-7	310
50-84-0	196	56-17-7	524	58-56-0	575	62-38-4	557
50-85-1	173	56-18-8	189	58-58-2	563	62-44-2	86
50-89-5	369	56-23-1	205	58-61-7	98	62-51-1	506
50-98-6	230	56-23-5	157	58-63-9	269	62-53-3	113
50-99-7	251	56-34-8	359	58-64-0	508	62-54-4	407
51-05-8	561	56-40-6	102	58-85-5	515	62-55-5	367
51-17-2	122	56-45-1	351	58-86-6	387	62-56-6	369
51-20-7	142	56-47-3	527	58-90-2	358	62-57-7	107
51-21-8	535	56-49-5	290	58-96-8	384	62-74-8	470
51-28-5	221	56-54-2	346	59-00-2	386	62-75-9	217
51-34-3	267	56-55-3	119	59-02-9	571	62-76-0	474
51-35-4	265	56-57-5	313	59-02-9	576	63-42-3	277
51-36-5	197	56-65-5	509	59-05-2	511	63-68-3	284
51-41-2	553	56-75-7	520	59-14-3	517	63-74-1	354
51-43-4	230	56-81-5	252	59-23-4	250	63-89-8	531
51-43-4	510	56-82-6	252	59-31-4	266	63-91-2	327
51-44-5	197	56-84-8	116	59-46-1	338	63-92-3	326
51-45-6	258	56-85-9	251	59-49-4	126	64-04-0	325
51-48-9	571	56-86-0	251	59-50-7	164	64-10-8	333
51-49-0	570	56-87-1	280	59-51-8	284	64-17-5	231
51-52-5	341	56-88-2	524	59-52-9	209	64-18-6	246
51-55-8	116	56-89-3	525	59-53-0	507	64-19-7	83
51-60-5	308	56-92-8	258	59-56-3	537	64-20-0	363
51-67-2	383	56-93-9	131	59-66-5	83	64-67-5	205
51-78-5	109	57-06-7	102	59-67-6	308	64-69-7	269
51-79-6	235	57-09-0	158	59-85-8	411	64-72-2	513
52-21-1	561	57-10-3	255	59-88-1	330	64-73-3	527
52-39-1	99	57-11-4	353	59-92-7	530	64-75-5	569
52-52-8	105	57-13-6	383	59-96-1	326	64-85-7	186
52-67-5	556	57-14-7	216	60-00-4	237	64-86-8	522
52-89-1	525	57-24-9	353	60-09-3	103	65-22-5	563
52-90-4	525	57-27-2	303	60-10-6	227	65-45-2	350
53-03-2	337	57-41-0	225	60-11-7	212	65-46-3	525
53-06-5	523	57-44-3	202	60-12-8	330	65-49-6	111

586 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

65-71-4	369	71-55-6	374	75-56-9	341	77-85-0	382
65-82-7	91	71-62-5	574	75-57-0	363	77-86-1	381
65-85-0	122	71-67-0	405	75-58-1	363	77-92-9	171
66-22-8	383	71-91-0	359	75-59-2	363	77-95-2	346
66-23-9	506	72-18-4	385	75-60-5	405	77-99-6	378
66-27-3	294	72-19-5	369	75-62-7	142	78-00-2	481
66-71-7	324	72-40-2	106	75-63-8	142	78-09-1	360
66-84-2	251	72-43-5	285	75-64-9	145	78-10-4	481
66-99-9	304	72-48-0	99	75-65-0	145	78-11-5	321
67-03-8	575	72-54-8	134	75-66-1	298	78-27-3	242
67-42-5	238	72-55-9	134	75-71-8	198	78-28-4	297
67-43-6	203	72-80-0	199	75-72-9	169	78-40-0	487
67-47-0	263	73-03-0	527	75-73-0	157	78-42-2	494
67-48-1	170	73-22-3	383	75-76-3	482	78-46-6	417
67-51-6	219	73-24-5	97	75-77-4	489	78-50-2	492
67-52-7	118	73-32-5	272	75-78-5	419	78-53-5	418
67-56-1	284	74-11-3	161	75-79-6	442	78-59-1	273
67-63-0	273	74-31-7	226	75-83-2	214	78-61-1	117
67-64-1	84	74-39-5	309	75-84-3	218	78-71-7	134
67-66-3	163	74-79-3	115	75-85-4	112	78-76-2	137
67-68-5	219	74-82-8	283	75-86-5	84	78-77-3	271
67-71-0	219	74-83-9	289	75-87-6	159	78-78-4	289
67-72-1	255	74-84-0	231	75-89-8	377	78-79-5	273
67-96-9	207	74-85-1	237	75-91-2	148	78-82-0	272
67-97-0	576	74-86-2	89	75-94-5	496	78-83-1	298
67-99-2	537	74-87-3	290	75-97-8	150	78-84-2	272
68-04-2	495	74-88-4	294	75-98-9	337	78-85-3	283
68-05-3	359	74-89-5	288	76-01-7	320	78-86-4	162
68-11-1	367	74-90-8	430	76-03-9	374	78-87-5	200
68-12-2	215	74-96-4	234	76-05-1	376	78-90-0	338
68-19-9	575	74-98-6	338	76-06-2	167	78-93-0	144
68-26-8	574	74-99-7	341	76-09-5	336	78-94-4	302
68-41-7	524	75-00-3	235	76-12-0	358	78-95-5	160
68-81-9	180	75-03-6	239	76-13-1	375	78-98-8	293
68-81-5	269	75-04-7	234	76-16-4	256	79-00-5	374
68-94-0	265	75-05-8	85	76-19-7	316	79-01-6	374
69-05-6	564	75-07-0	81	76-24-4	100	79-05-0	339
69-33-0	572	75-08-1	231	76-37-9	360	79-06-1	95
69-57-8	514	75-09-2	198	76-39-1	296	79-07-2	160
69-65-8	281	75-11-6	208	76-57-3	522	79-08-3	136
69-72-7	350	75-12-7	245	76-59-5	142	79-09-4	339
69-89-6	576	75-15-0	156	76-60-8	138	79-10-7	95
69-93-2	384	75-18-3	219	76-73-3	351	79-11-8	160
70-11-1	136	75-19-4	182	76-74-4	322	79-14-1	252
70-18-8	538	75-21-8	238	76-75-2	283	79-15-2	136
70-23-5	235	75-22-9	404	76-83-5	381	79-16-3	287
70-26-8	318	75-24-1	392	76-84-6	381	79-19-6	369
70-34-8	221	75-25-2	138	76-86-8	412	79-20-9	287
70-47-3	116	75-26-3	274	76-87-9	494	79-24-3	311
70-49-5	368	75-28-5	271	76-93-7	122	79-27-6	356
70-55-3	370	75-29-6	274	77-06-5	250	79-29-8	214
70-69-9	110	75-30-9	274	77-09-8	326	79-31-2	272
70-70-2	265	75-33-2	339	77-21-4	240	79-33-4	277
70-78-0	270	75-34-3	198	77-26-9	102	79-34-5	358
71-00-1	258	75-36-5	88	77-47-4	255	79-38-9	169
71-23-8	340	75-44-5	450	77-48-5	193	79-39-0	283
71-30-7	527	75-47-8	269	77-49-6	296	79-40-3	227
71-36-3	143	75-50-3	378	77-53-2	158	79-41-4	283
71-41-0	111	75-52-5	311	77-59-8	571	79-43-6	196
71-43-2	119	75-54-7	441	77-71-4	216	79-44-7	215

79-46-9	<i>313</i>	83-40-9	<i>173</i>	86-74-8	<i>156</i>	89-82-7	<i>342</i>
79-49-2	<i>320</i>	83-43-2	<i>547</i>	86-84-0	<i>307</i>	89-86-1	<i>207</i>
79-57-2	<i>568</i>	83-44-3	<i>186</i>	86-87-3	<i>306</i>	89-98-5	<i>160</i>
79-58-3	<i>349</i>	83-46-5	<i>351</i>	86-88-4	<i>307</i>	90-00-6	<i>240</i>
79-63-0	<i>277</i>	83-48-7	<i>353</i>	86-93-1	<i>331</i>	90-01-7	<i>260</i>
79-69-6	<i>270</i>	83-49-8	<i>267</i>	86-98-6	<i>200</i>	90-02-8	<i>350</i>
79-74-3	<i>190</i>	83-56-7	<i>208</i>	87-13-8	<i>204</i>	90-04-0	<i>113</i>
79-77-6	<i>270</i>	83-60-3	<i>348</i>	87-17-2	<i>350</i>	90-05-1	<i>253</i>
79-81-2	<i>348</i>	83-67-0	<i>366</i>	87-32-1	<i>93</i>	90-11-9	<i>140</i>
80-05-7	<i>274</i>	83-72-7	<i>264</i>	87-41-2	<i>334</i>	90-12-0	<i>295</i>
80-11-5	<i>296</i>	83-73-8	<i>208</i>	87-42-3	<i>168</i>	90-13-1	<i>165</i>
80-15-9	<i>175</i>	83-74-9	<i>267</i>	87-48-9	<i>139</i>	90-15-3	<i>305</i>
80-43-3	<i>201</i>	83-88-5	<i>575</i>	87-51-4	<i>258</i>	90-19-7	<i>348</i>
80-46-6	<i>112</i>	84-11-7	<i>324</i>	87-52-5	<i>253</i>	90-20-0	<i>108</i>
80-48-8	<i>302</i>	84-12-8	<i>325</i>	87-56-9	<i>304</i>	90-26-6	<i>328</i>
80-59-1	<i>369</i>	84-15-1	<i>355</i>	87-61-6	<i>374</i>	90-30-2	<i>331</i>
80-62-6	<i>294</i>	84-16-2	<i>257</i>	87-62-7	<i>213</i>	90-33-5	<i>548</i>
80-70-6	<i>364</i>	84-17-3	<i>202</i>	87-66-1	<i>345</i>	90-41-5	<i>104</i>
80-72-8	<i>348</i>	84-21-9	<i>508</i>	87-68-3	<i>323</i>	90-43-7	<i>260</i>
80-78-4	<i>351</i>	84-24-2	<i>334</i>	87-72-9	<i>115</i>	90-44-8	<i>115</i>
80-91-1	<i>338</i>	84-54-8	<i>288</i>	87-73-0	<i>349</i>	90-45-9	<i>103</i>
80-92-2	<i>338</i>	84-58-2	<i>197</i>	87-81-0	<i>355</i>	90-46-0	<i>387</i>
80-97-7	<i>169</i>	84-65-1	<i>115</i>	87-85-4	<i>256</i>	90-47-1	<i>386</i>
81-07-2	<i>122</i>	84-66-2	<i>205</i>	87-86-5	<i>320</i>	90-52-8	<i>107</i>
81-08-3	<i>126</i>	84-68-4	<i>196</i>	87-87-6	<i>206</i>	90-65-3	<i>319</i>
81-13-0	<i>555</i>	84-74-2	<i>195</i>	87-88-7	<i>159</i>	90-80-2	<i>251</i>
81-14-1	<i>276</i>	84-79-7	<i>277</i>	87-89-8	<i>543</i>	90-82-4	<i>341</i>
81-16-3	<i>307</i>	84-80-0	<i>576</i>	88-04-0	<i>162</i>	90-90-4	<i>137</i>
81-23-2	<i>186</i>	84-86-6	<i>305</i>	88-06-2	<i>375</i>	90-91-5	<i>570</i>
81-24-3	<i>355</i>	84-88-8	<i>266</i>	88-14-2	<i>249</i>	90-93-7	<i>134</i>
81-25-4	<i>170</i>	84-89-9	<i>306</i>	88-15-3	<i>92</i>	90-94-8	<i>303</i>
81-27-6	<i>351</i>	85-01-8	<i>324</i>	88-16-4	<i>161</i>	90-96-0	<i>210</i>
81-31-2	<i>386</i>	85-02-9	<i>124</i>	88-19-7	<i>370</i>	90-98-2	<i>197</i>
81-54-9	<i>342</i>	85-18-7	<i>169</i>	88-21-1	<i>318</i>	90-99-3	<i>225</i>
81-61-8	<i>346</i>	85-31-4	<i>367</i>	88-27-7	<i>194</i>	91-00-9	<i>225</i>
81-64-1	<i>347</i>	85-32-5	<i>253</i>	88-30-2	<i>377</i>	91-01-0	<i>121</i>
81-77-6	<i>268</i>	85-41-6	<i>334</i>	88-31-1	<i>368</i>	91-02-1	<i>126</i>
81-81-2	<i>386</i>	85-44-9	<i>334</i>	88-58-4	<i>195</i>	91-10-1	<i>211</i>
81-84-5	<i>305</i>	85-46-1	<i>305</i>	88-65-3	<i>137</i>	91-13-4	<i>193</i>
81-88-9	<i>348</i>	85-47-2	<i>304</i>	88-67-5	<i>269</i>	91-15-6	<i>334</i>
81-92-5	<i>326</i>	85-52-9	<i>125</i>	88-72-2	<i>314</i>	91-16-7	<i>210</i>
81-93-6	<i>326</i>	85-61-0	<i>522</i>	88-73-3	<i>165</i>	91-17-8	<i>184</i>
82-02-0	<i>276</i>	85-66-5	<i>342</i>	88-74-4	<i>309</i>	91-18-9	<i>341</i>
82-05-3	<i>119</i>	85-83-6	<i>354</i>	88-75-5	<i>312</i>	91-19-0	<i>347</i>
82-22-4	<i>115</i>	85-84-7	<i>120</i>	88-88-0	<i>335</i>	91-20-3	<i>304</i>
82-24-6	<i>103</i>	85-85-8	<i>345</i>	88-89-1	<i>335</i>	91-22-5	<i>347</i>
82-38-2	<i>288</i>	85-86-9	<i>354</i>	88-99-3	<i>334</i>	91-23-6	<i>309</i>
82-57-5	<i>386</i>	85-98-3	<i>203</i>	89-01-0	<i>342</i>	91-40-7	<i>327</i>
82-58-6	<i>279</i>	86-00-0	<i>310</i>	89-05-4	<i>345</i>	91-48-5	<i>329</i>
82-62-2	<i>374</i>	86-28-2	<i>235</i>	89-25-8	<i>297</i>	91-52-1	<i>210</i>
82-68-8	<i>320</i>	86-29-3	<i>224</i>	89-32-7	<i>345</i>	91-53-2	<i>533</i>
82-71-3	<i>353</i>	86-34-0	<i>326</i>	89-51-0	<i>258</i>	91-56-5	<i>270</i>
82-86-0	<i>81</i>	86-35-1	<i>240</i>	89-52-1	<i>88</i>	91-57-6	<i>295</i>
82-94-0	<i>293</i>	86-38-4	<i>199</i>	89-56-5	<i>173</i>	91-58-7	<i>165</i>
83-07-8	<i>103</i>	86-42-0	<i>111</i>	89-57-6	<i>111</i>	91-59-8	<i>306</i>
83-12-5	<i>331</i>	86-48-6	<i>306</i>	89-61-2	<i>199</i>	91-60-1	<i>368</i>
83-30-7	<i>377</i>	86-55-5	<i>305</i>	89-63-4	<i>165</i>	91-62-3	<i>300</i>
83-32-9	<i>81</i>	86-57-7	<i>312</i>	89-65-6	<i>271</i>	91-63-4	<i>300</i>
83-34-1	<i>294</i>	86-59-9	<i>347</i>	89-69-0	<i>374</i>	91-64-5	<i>173</i>
83-38-5	<i>196</i>	86-73-7	<i>243</i>	89-73-6	<i>350</i>	91-65-6	<i>203</i>

588 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

91-66-7	202	94-36-0	125	97-00-7	162	99-30-9	199
91-94-1	196	94-41-7	119	97-02-9	220	99-32-1	159
92-06-8	355	94-53-1	337	97-23-4	199	99-33-2	221
92-24-0	304	94-59-7	350	97-30-3	293	99-34-3	221
92-29-5	259	94-62-2	337	97-39-2	227	99-35-4	380
92-32-0	346	94-67-7	350	97-41-6	236	99-54-7	199
92-35-3	367	94-74-6	164	97-44-9	390	99-55-8	296
92-41-1	304	94-75-7	200	97-53-0	242	99-56-9	312
92-51-3	131	95-03-4	164	97-56-3	103	99-57-0	109
92-52-4	132	95-13-6	268	97-59-6	99	99-61-6	309
92-59-1	128	95-14-7	124	97-62-1	239	99-65-0	220
92-61-5	350	95-15-8	124	97-63-2	239	99-73-0	140
92-62-6	338	95-20-5	294	97-65-4	275	99-76-3	293
92-66-0	137	95-45-4	216	97-67-6	280	99-77-4	240
92-67-1	104	95-47-6	387	97-88-1	149	99-81-0	140
92-69-3	260	95-48-7	173	97-90-5	237	99-87-6	183
92-70-6	306	95-49-8	169	97-93-8	391	99-90-1	136
92-71-7	225	95-50-1	196	97-94-9	487	99-92-3	103
92-76-2	263	95-51-2	160	97-95-0	235	99-93-4	259
92-77-3	263	95-52-3	245	97-97-2	160	99-94-5	371
92-82-0	325	95-53-4	371	98-00-0	249	99-96-7	260
92-83-1	386	95-54-5	329	98-01-1	249	99-97-8	291
92-84-2	326	95-55-6	109	98-03-3	368	99-99-0	315
92-85-3	366	95-56-7	140	98-05-5	448	100-00-5	165
92-86-4	192	95-57-8	166	98-06-6	146	100-01-6	309
92-87-5	121	95-63-6	378	98-08-8	377	100-02-7	312
92-91-1	327	95-64-7	213	98-09-9	121	100-06-1	285
92-92-2	132	95-65-8	218	98-11-3	121	100-09-4	285
92-94-4	356	95-71-6	371	98-15-7	161	100-10-7	212
93-00-5	307	95-73-8	201	98-29-3	147	100-11-8	310
93-04-9	286	95-75-0	201	98-48-6	120	100-13-0	314
93-07-2	210	95-76-1	196	98-50-0	396	100-14-1	310
93-08-3	91	95-77-2	199	98-51-1	151	100-15-2	296
93-09-4	305	95-80-7	370	98-53-3	148	100-16-3	313
93-10-7	346	95-83-0	167	98-54-4	150	100-17-4	309
93-11-8	305	95-84-1	106	98-56-6	161	100-19-6	309
93-15-2	242	95-87-1	218	98-58-8	137	100-20-9	355
93-17-4	258	95-88-5	168	98-59-9	370	100-21-0	355
93-18-5	233	95-93-2	363	98-60-2	161	100-22-1	364
93-20-9	307	95-94-3	358	98-79-3	345	100-23-2	217
93-25-4	286	95-95-4	374	98-80-6	448	100-25-4	220
93-31-9	371	96-09-3	353	98-82-8	174	100-26-5	344
93-35-6	383	96-10-6	418	98-83-9	301	100-28-7	313
93-40-3	211	96-14-0	297	98-86-2	86	100-39-0	127
93-43-6	307	96-22-0	204	98-88-4	125	100-41-6	234
93-53-8	331	96-31-1	220	98-92-0	308	100-42-5	353
93-56-1	353	96-33-3	287	98-95-3	309	100-43-6	385
93-58-3	289	96-34-4	290	98-96-4	342	100-44-7	127
93-59-4	323	96-37-7	291	98-97-5	342	100-46-9	127
93-62-9	261	96-43-5	169	98-98-6	335	100-47-0	123
93-76-5	375	96-45-7	238	99-03-6	103	100-48-1	176
93-89-0	234	96-47-9	301	99-04-7	371	100-51-6	126
93-91-4	124	96-48-0	152	99-05-8	104	100-52-7	119
93-95-8	330	96-49-1	237	99-06-9	260	100-53-8	129
93-97-0	122	96-50-4	111	99-08-1	314	100-54-9	176
93-98-1	119	96-53-7	366	99-09-2	309	100-57-2	449
93-99-2	328	96-54-8	300	99-11-6	171	100-61-8	288
94-09-7	234	96-77-5	326	99-14-9	373	100-63-0	330
94-20-2	340	96-82-2	277	99-24-1	293	100-64-1	179
94-25-7	146	96-97-9	313	99-28-5	193	100-65-2	331

100-66-3	<i>114</i>	103-88-8	<i>136</i>	106-54-7	<i>169</i>	108-31-6	<i>280</i>
100-69-6	<i>385</i>	103-89-9	<i>86</i>	106-57-0	<i>539</i>	108-32-7	<i>340</i>
100-71-0	<i>241</i>	103-90-2	<i>83</i>	106-70-7	<i>293</i>	108-38-3	<i>387</i>
100-74-3	<i>240</i>	104-01-8	<i>286</i>	106-88-7	<i>230</i>	108-39-4	<i>173</i>
100-76-5	<i>347</i>	104-03-0	<i>312</i>	106-89-8	<i>230</i>	108-41-8	<i>169</i>
100-83-4	<i>260</i>	104-04-1	<i>308</i>	106-93-4	<i>193</i>	108-43-0	<i>166</i>
100-85-6	<i>131</i>	104-12-1	<i>167</i>	106-94-5	<i>340</i>	108-44-1	<i>371</i>
100-97-0	<i>257</i>	104-24-5	<i>328</i>	106-95-6	<i>101</i>	108-45-2	<i>329</i>
101-01-9	<i>381</i>	104-51-8	<i>146</i>	106-97-8	<i>143</i>	108-46-3	<i>348</i>
101-02-0	<i>493</i>	104-54-1	<i>171</i>	106-99-0	<i>143</i>	108-47-4	<i>278</i>
101-43-9	<i>180</i>	104-74-5	<i>228</i>	107-02-8	<i>94</i>	108-48-5	<i>279</i>
101-49-5	<i>128</i>	104-76-7	<i>239</i>	107-03-9	<i>339</i>	108-64-5	<i>239</i>
101-53-1	<i>130</i>	104-81-4	<i>143</i>	107-04-0	<i>163</i>	108-67-8	<i>283</i>
101-54-2	<i>105</i>	104-82-5	<i>289</i>	107-05-1	<i>101</i>	108-68-9	<i>218</i>
101-61-1	<i>364</i>	104-83-6	<i>162</i>	107-06-2	<i>198</i>	108-70-3	<i>374</i>
101-63-3	<i>135</i>	104-85-8	<i>372</i>	107-07-3	<i>163</i>	108-73-6	<i>333</i>
101-64-4	<i>286</i>	104-87-0	<i>369</i>	107-08-4	<i>341</i>	108-75-8	<i>379</i>
101-77-9	<i>189</i>	104-88-1	<i>161</i>	107-10-8	<i>340</i>	108-77-0	<i>176</i>
101-81-5	<i>225</i>	104-90-5	<i>240</i>	107-11-9	<i>101</i>	108-78-1	<i>281</i>
101-82-6	<i>130</i>	104-91-6	<i>314</i>	107-12-0	<i>339</i>	108-79-2	<i>216</i>
101-84-8	<i>330</i>	104-92-7	<i>136</i>	107-13-1	<i>95</i>	108-80-5	<i>176</i>
101-97-3	<i>240</i>	104-93-8	<i>288</i>	107-14-2	<i>160</i>	108-85-0	<i>180</i>
102-01-2	<i>84</i>	104-94-9	<i>114</i>	107-15-3	<i>237</i>	108-86-1	<i>137</i>
102-02-3	<i>328</i>	104-96-1	<i>294</i>	107-18-6	<i>100</i>	108-87-2	<i>291</i>
102-04-5	<i>192</i>	104-98-3	<i>384</i>	107-19-7	<i>339</i>	108-88-3	<i>370</i>
102-06-7	<i>225</i>	105-30-6	<i>297</i>	107-21-1	<i>238</i>	108-89-4	<i>300</i>
102-07-8	<i>156</i>	105-34-0	<i>290</i>	107-27-7	<i>423</i>	108-90-7	<i>161</i>
102-08-9	<i>226</i>	105-36-2	<i>235</i>	107-30-2	<i>164</i>	108-91-8	<i>180</i>
102-09-0	<i>224</i>	105-37-3	<i>240</i>	107-31-3	<i>292</i>	108-93-0	<i>179</i>
102-25-0	<i>376</i>	105-39-5	<i>235</i>	107-41-5	<i>297</i>	108-94-1	<i>179</i>
102-54-5	<i>424</i>	105-53-3	<i>205</i>	107-43-7	<i>131</i>	108-95-2	<i>325</i>
102-69-2	<i>381</i>	105-54-4	<i>235</i>	107-46-0	<i>428</i>	108-97-4	<i>345</i>
102-82-9	<i>373</i>	105-55-5	<i>206</i>	107-51-7	<i>446</i>	108-98-5	<i>368</i>
102-85-2	<i>486</i>	105-56-6	<i>236</i>	107-59-5	<i>147</i>	108-99-6	<i>299</i>
102-94-3	<i>171</i>	105-57-7	<i>81</i>	107-71-1	<i>150</i>	109-00-2	<i>266</i>
103-03-7	<i>332</i>	105-58-8	<i>203</i>	107-81-3	<i>140</i>	109-02-4	<i>295</i>
103-16-2	<i>130</i>	105-60-2	<i>155</i>	107-82-4	<i>271</i>	109-04-6	<i>141</i>
103-19-5	<i>372</i>	105-61-3	<i>280</i>	107-83-5	<i>296</i>	109-06-8	<i>299</i>
103-29-7	<i>131</i>	105-74-8	<i>278</i>	107-87-9	<i>299</i>	109-07-9	<i>298</i>
103-30-0	<i>353</i>	106-02-5	<i>320</i>	107-89-1	<i>99</i>	109-08-0	<i>299</i>
103-32-2	<i>127</i>	106-22-9	<i>172</i>	107-91-5	<i>175</i>	109-09-1	<i>168</i>
103-33-3	<i>117</i>	106-24-1	<i>250</i>	107-92-6	<i>152</i>	109-12-6	<i>110</i>
103-34-4	<i>227</i>	106-28-5	<i>534</i>	107-93-7	<i>174</i>	109-13-7	<i>150</i>
103-36-6	<i>236</i>	106-31-0	<i>152</i>	107-94-8	<i>168</i>	109-52-4	<i>384</i>
103-41-3	<i>127</i>	106-34-3	<i>346</i>	107-95-9	<i>98</i>	109-53-5	<i>272</i>
103-44-6	<i>239</i>	106-36-5	<i>341</i>	107-97-1	<i>350</i>	109-57-9	<i>102</i>
103-49-1	<i>192</i>	106-37-6	<i>192</i>	108-01-0	<i>212</i>	109-60-4	<i>340</i>
103-50-4	<i>128</i>	106-38-7	<i>142</i>	108-03-2	<i>313</i>	109-63-7	<i>404</i>
103-63-9	<i>325</i>	106-40-1	<i>136</i>	108-05-4	<i>385</i>	109-64-8	<i>193</i>
103-70-8	<i>246</i>	106-41-2	<i>140</i>	108-08-7	<i>218</i>	109-65-9	<i>146</i>
103-71-9	<i>331</i>	106-42-3	<i>387</i>	108-10-1	<i>297</i>	109-66-0	<i>322</i>
103-72-0	<i>331</i>	106-43-4	<i>169</i>	108-11-2	<i>297</i>	109-68-2	<i>322</i>
103-73-1	<i>325</i>	106-44-5	<i>173</i>	108-13-4	<i>280</i>	109-69-3	<i>147</i>
103-79-9	<i>327</i>	106-45-6	<i>371</i>	108-18-9	<i>209</i>	109-70-6	<i>168</i>
103-81-1	<i>327</i>	106-46-7	<i>196</i>	108-19-0	<i>135</i>	109-73-9	<i>145</i>
103-82-2	<i>327</i>	106-47-8	<i>160</i>	108-20-3	<i>274</i>	109-74-0	<i>152</i>
103-83-3	<i>128</i>	106-48-9	<i>166</i>	108-22-5	<i>274</i>	109-75-1	<i>101</i>
103-84-4	<i>83</i>	106-49-0	<i>371</i>	108-24-7	<i>83</i>	109-77-3	<i>281</i>
103-85-5	<i>332</i>	106-50-3	<i>329</i>	108-29-2	<i>384</i>	109-79-5	<i>143</i>
103-86-6	<i>110</i>	106-51-4	<i>124</i>	108-30-5	<i>354</i>	109-82-0	<i>292</i>

590 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

109-85-3	286	111-20-6	350	112-85-6	228	118-92-3	103
109-86-4	285	111-25-1	257	112-86-7	230	118-96-7	380
109-87-5	245	111-26-2	257	112-89-0	140	118-97-8	162
109-89-7	202	111-27-3	257	112-92-5	316	119-24-4	342
109-90-0	239	111-30-8	251	112-95-8	229	119-26-6	222
109-92-2	242	111-34-2	151	113-15-5	230	119-27-7	220
109-94-4	238	111-40-0	188	113-00-8	253	119-30-2	270
109-97-7	346	111-41-1	261	113-73-5	540	119-32-4	296
109-99-9	361	111-42-2	202	114-07-8	533	119-36-8	300
110-00-9	249	111-43-3	341	114-21-6	473	119-39-1	334
110-01-0	361	111-44-4	134	114-25-0	132	119-48-8	576
110-02-1	368	111-46-6	203	114-49-8	566	119-53-9	123
110-03-2	216	111-47-7	226	114-70-5	474	119-61-9	123
110-05-4	195	111-49-9	256	114-80-7	308	119-64-2	362
110-13-4	86	111-55-7	238	114-83-0	92	119-65-3	275
110-14-5	353	111-61-5	241	115-02-6	117	119-68-6	288
110-15-6	353	111-63-7	386	115-07-1	339	119-91-5	133
110-16-7	280	111-65-9	316	115-08-2	367	119-93-7	369
110-17-8	248	111-66-0	317	115-09-3	441	120-12-7	114
110-18-9	364	111-68-2	254	115-10-6	292	120-14-9	385
110-20-3	84	111-69-3	98	115-11-7	271	120-18-3	305
110-21-4	259	111-70-6	254	115-25-3	323	120-20-7	259
110-26-9	133	111-71-7	254	115-27-5	160	120-23-0	306
110-42-9	291	111-76-2	144	115-37-7	366	120-29-6	382
110-44-1	352	111-77-3	203	115-39-9	140	120-36-5	200
110-45-2	273	111-82-0	292	115-40-2	138	120-46-7	192
110-49-6	87	111-83-1	317	115-41-3	459	120-57-0	337
110-51-0	403	111-84-2	315	115-69-5	108	120-61-6	219
110-53-2	112	111-87-5	317	115-76-4	205	120-66-1	86
110-54-3	257	111-88-6	317	115-77-5	320	120-72-9	268
110-58-7	112	111-90-0	203	115-85-5	411	120-73-0	342
110-59-8	385	111-92-2	194	115-86-6	493	120-78-8	227
110-61-2	354	111-96-6	206	115-87-7	401	120-80-9	157
110-62-3	384	112-00-5	229	116-17-6	488	120-82-1	274
110-63-4	143	112-02-7	158	116-31-4	574	120-83-2	199
110-65-6	152	112-04-9	446	116-63-2	109	120-92-3	182
110-66-7	322	112-05-0	319	116-85-8	106	120-93-4	238
110-71-4	210	112-07-2	87	117-08-8	358	121-00-6	150
110-74-7	341	112-24-3	376	117-10-2	207	121-14-2	222
110-75-8	163	112-27-6	376	117-12-4	115	121-19-7	446
110-80-5	233	112-29-8	185	117-18-0	358	121-33-5	385
110-81-6	205	112-30-1	185	117-34-0	224	121-40-4	356
110-82-7	178	112-34-5	203	117-37-3	114	121-43-7	488
110-83-8	179	112-36-7	203	117-39-3	346	121-44-8	375
110-85-0	336	112-38-9	383	117-80-6	199	121-45-9	490
110-86-1	343	112-40-3	228	117-81-7	204	121-46-0	315
110-87-2	206	112-42-5	383	118-00-3	253	121-47-1	283
110-88-3	380	112-47-0	185	118-10-5	521	121-54-0	121
110-89-4	336	112-48-1	238	118-48-9	270	121-54-0	259
110-91-8	303	112-49-2	376	118-52-5	198	121-57-3	354
110-94-1	252	112-53-8	228	118-55-8	332	121-60-8	82
110-97-4	209	112-54-9	277	118-65-0	272	121-69-7	213
110-98-5	226	112-55-0	228	118-69-4	201	121-70-0	449
110-99-6	206	112-57-2	360	118-71-8	281	121-72-2	291
111-01-3	567	112-62-9	296	118-74-1	255	121-73-3	165
111-02-4	567	112-72-1	359	118-75-2	159	121-79-9	341
111-11-5	296	112-73-2	194	118-78-5	383	121-82-4	183
111-15-9	87	112-79-8	229	118-79-6	373	121-88-0	109
111-16-0	335	112-80-1	317	118-90-1	371	121-89-1	308
111-17-1	367	112-82-3	158	118-91-2	161	121-91-5	273

121-92-6	<i>310</i>	124-47-0	<i>384</i>	130-26-7	<i>163</i>	138-37-4	<i>107</i>
121-98-2	<i>288</i>	124-58-3	<i>441</i>	130-40-5	<i>535</i>	138-52-3	<i>350</i>
122-00-9	<i>287</i>	124-63-0	<i>284</i>	130-85-8	<i>229</i>	138-59-0	<i>351</i>
122-03-2	<i>175</i>	124-64-1	<i>481</i>	130-86-9	<i>341</i>	138-60-3	<i>266</i>
122-04-3	<i>310</i>	124-65-2	<i>467</i>	130-95-0	<i>347</i>	138-84-1	<i>452</i>
122-19-0	<i>128</i>	124-68-5	<i>108</i>	131-08-8	<i>395</i>	138-89-6	<i>314</i>
122-25-8	<i>292</i>	124-70-9	<i>442</i>	131-11-3	<i>218</i>	139-02-6	<i>474</i>
122-37-2	<i>261</i>	124-73-2	<i>193</i>	131-14-6	<i>188</i>	139-13-9	<i>308</i>
122-39-4	<i>224</i>	124-76-5	<i>271</i>	131-18-0	<i>190</i>	139-59-3	<i>326</i>
122-51-0	<i>240</i>	124-83-4	<i>153</i>	131-22-6	<i>328</i>	139-66-2	<i>226</i>
122-51-0	<i>376</i>	124-87-8	<i>335</i>	131-28-2	<i>308</i>	139-85-5	<i>341</i>
122-52-1	<i>487</i>	125-28-0	<i>206</i>	131-48-6	<i>507</i>	140-10-3	<i>171</i>
122-57-6	<i>118</i>	125-46-2	<i>573</i>	131-56-6	<i>207</i>	140-11-4	<i>126</i>
122-59-8	<i>326</i>	126-00-1	<i>516</i>	131-69-1	<i>334</i>	140-22-7	<i>224</i>
122-61-8	<i>301</i>	126-07-8	<i>253</i>	131-74-8	<i>394</i>	140-28-3	<i>192</i>
122-66-7	<i>225</i>	126-11-4	<i>382</i>	131-91-9	<i>314</i>	140-29-4	<i>128</i>
122-79-2	<i>327</i>	126-14-7	<i>354</i>	132-27-4	<i>474</i>	140-40-9	<i>82</i>
122-80-5	<i>102</i>	126-17-0	<i>352</i>	132-28-5	<i>494</i>	140-66-9	<i>317</i>
122-85-0	<i>82</i>	126-18-1	<i>351</i>	132-29-6	<i>401</i>	140-87-4	<i>175</i>
122-88-3	<i>166</i>	126-30-7	<i>218</i>	132-53-6	<i>314</i>	140-88-5	<i>234</i>
123-07-9	<i>240</i>	126-33-0	<i>354</i>	132-60-5	<i>332</i>	140-89-6	<i>454</i>
123-08-0	<i>260</i>	126-71-6	<i>488</i>	132-64-9	<i>191</i>	140-93-2	<i>473</i>
123-11-5	<i>113</i>	126-73-8	<i>486</i>	132-65-0	<i>191</i>	140-95-4	<i>217</i>
123-19-3	<i>226</i>	126-81-8	<i>209</i>	132-66-1	<i>307</i>	141-22-0	<i>349</i>
123-25-1	<i>205</i>	126-98-7	<i>283</i>	133-06-2	<i>155</i>	141-32-2	<i>145</i>
123-30-8	<i>109</i>	127-06-0	<i>86</i>	133-11-9	<i>327</i>	141-43-5	<i>105</i>
123-31-9	<i>259</i>	127-07-1	<i>431</i>	133-32-4	<i>543</i>	141-52-6	<i>470</i>
123-32-0	<i>219</i>	127-08-2	<i>452</i>	133-38-0	<i>208</i>	141-53-7	<i>471</i>
123-33-1	<i>280</i>	127-09-3	<i>464</i>	133-49-3	<i>320</i>	141-75-3	<i>152</i>
123-38-6	<i>339</i>	127-17-3	<i>346</i>	134-32-7	<i>306</i>	141-78-6	<i>233</i>
123-39-7	<i>292</i>	127-18-4	<i>358</i>	134-58-7	<i>116</i>	141-79-7	<i>283</i>
123-42-2	<i>263</i>	127-19-5	<i>211</i>	134-81-6	<i>121</i>	141-82-2	<i>280</i>
123-46-6	<i>251</i>	127-40-2	<i>278</i>	134-84-9	<i>289</i>	141-84-4	<i>349</i>
123-51-3	<i>290</i>	127-41-3	<i>270</i>	134-85-0	<i>161</i>	141-86-6	<i>190</i>
123-54-6	<i>87</i>	127-47-9	<i>348</i>	134-96-3	<i>355</i>	141-90-2	<i>369</i>
123-56-8	<i>354</i>	127-63-9	<i>226</i>	135-16-0	<i>569</i>	141-97-9	<i>234</i>
123-62-6	<i>339</i>	127-82-2	<i>498</i>	135-19-3	<i>305</i>	142-04-1	<i>113</i>
123-63-7	<i>319</i>	127-95-7	<i>455</i>	135-20-6	<i>414</i>	142-08-5	<i>265</i>
123-68-3	<i>263</i>	127-96-8	<i>458</i>	135-48-8	<i>330</i>	142-09-6	<i>257</i>
123-72-8	<i>152</i>	128-00-7	<i>463</i>	135-49-9	<i>94</i>	142-18-5	<i>254</i>
123-73-9	<i>174</i>	128-04-1	<i>469</i>	135-52-4	<i>498</i>	142-29-0	<i>182</i>
123-75-1	<i>346</i>	128-08-5	<i>142</i>	135-59-1	<i>369</i>	142-45-0	<i>89</i>
123-78-4	<i>567</i>	128-09-6	<i>169</i>	135-61-5	<i>305</i>	142-59-6	<i>421</i>
123-86-4	<i>144</i>	128-13-2	<i>384</i>	135-67-1	<i>326</i>	142-62-1	<i>155</i>
123-91-1	<i>223</i>	128-37-0	<i>194</i>	135-68-2	<i>105</i>	142-63-2	<i>336</i>
123-92-2	<i>270</i>	128-39-2	<i>195</i>	135-70-6	<i>346</i>	142-64-3	<i>336</i>
123-93-3	<i>367</i>	128-57-4	<i>351</i>	135-73-9	<i>331</i>	142-68-7	<i>361</i>
123-95-5	<i>151</i>	128-64-3	<i>275</i>	135-88-6	<i>331</i>	142-71-2	<i>414</i>
123-99-9	<i>117</i>	128-95-0	<i>188</i>	136-09-4	<i>521</i>	142-72-3	<i>437</i>
124-02-7	<i>187</i>	129-00-0	<i>342</i>	136-36-6	<i>191</i>	142-73-4	<i>267</i>
124-04-9	<i>98</i>	129-16-8	<i>440</i>	136-64-8	<i>263</i>	142-87-0	<i>468</i>
124-09-4	<i>256</i>	129-44-2	<i>188</i>	136-72-1	<i>336</i>	142-96-1	<i>148</i>
124-10-7	<i>301</i>	129-64-6	<i>315</i>	136-95-8	<i>104</i>	143-07-7	<i>277</i>
124-12-9	<i>155</i>	129-66-8	<i>380</i>	137-08-6	<i>555</i>	143-18-0	<i>456</i>
124-18-5	<i>185</i>	129-73-7	<i>278</i>	137-18-8	<i>214</i>	143-19-1	<i>474</i>
124-20-9	<i>352</i>	129-79-3	<i>380</i>	137-26-8	<i>365</i>	143-24-8	<i>360</i>
124-38-9	<i>409</i>	130-14-3	<i>473</i>	137-30-4	<i>498</i>	143-27-1	<i>158</i>
124-40-3	<i>211</i>	130-15-4	<i>306</i>	137-32-6	<i>289</i>	143-28-2	<i>318</i>
124-41-4	<i>473</i>	130-22-3	<i>390</i>	137-40-6	<i>475</i>	143-62-4	<i>529</i>
124-42-5	<i>82</i>	130-24-5	<i>108</i>	138-10-3	<i>329</i>	143-66-8	<i>476</i>

592 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

143-67-9	<i>574</i>	156-54-7	<i>467</i>	294-93-9	<i>414</i>	329-98-6	<i>557</i>
144-02-5	<i>466</i>	156-57-0	<i>525</i>	295-17-0	<i>183</i>	330-54-1	<i>200</i>
144-48-9	<i>269</i>	156-59-2	<i>198</i>	295-37-4	<i>356</i>	331-39-5	<i>152</i>
144-55-8	<i>466</i>	156-60-5	<i>198</i>	297-90-5	<i>263</i>	333-18-6	<i>237</i>
144-62 -7	<i>318</i>	156-73-5	<i>537</i>	298-07-7	<i>418</i>	333-20-0	<i>458</i>
144-67-2	<i>334</i>	156-83-2	<i>162</i>	298-12-4	<i>253</i>	333-27-7	<i>302</i>
144-68-3	<i>388</i>	157-03-9	<i>191</i>	298-14-6	<i>453</i>	333-49-3	<i>570</i>
144-80-9	<i>92</i>	188-87-4	<i>386</i>	298-81-7	<i>577</i>	333-93-7	<i>188</i>
144-83-2	<i>354</i>	191-07-1	<i>172</i>	298-93-1	<i>570</i>	334-48-5	<i>154</i>
144-90-1	<i>107</i>	191-24-2	<i>123</i>	298-96-4	<i>381</i>	334-50-9	<i>252</i>
145-13-1	<i>265</i>	192-97-2	<i>123</i>	299-11-6	<i>547</i>	335-27-3	<i>323</i>
145-42-6	<i>475</i>	195-19-7	<i>123</i>	299-28-5	<i>407</i>	335-57-9	<i>323</i>
146-14-5	<i>534</i>	195-84-6	<i>375</i>	299-42-3	<i>533</i>	335-67-1	<i>320</i>
146-48-5	<i>388</i>	198-55-0	<i>324</i>	300-39-0	<i>530</i>	338-83-0	<i>323</i>
146-68-9	<i>544</i>	206-44-0	<i>243</i>	301-04-2	<i>434</i>	339-72-0	<i>524</i>
146-77-0	<i>520</i>	208-96-8	<i>81</i>	302-01-2	<i>429</i>	340-06-7	<i>333</i>
147-14-8	<i>415</i>	213-14-3	<i>335</i>	302-17-0	<i>159</i>	341-92-4	<i>244</i>
147-71-7	<i>355</i>	217-59-4	<i>381</i>	302-27-2	<i>93</i>	343-27-1	<i>253</i>
147-73-9	<i>355</i>	218-01-9	<i>170</i>	302-66-9	<i>299</i>	343-65-7	<i>277</i>
147-82-0	<i>373</i>	222-93-5	<i>192</i>	302-72-7	<i>98</i>	343-94-2	<i>383</i>
147-85-3	<i>338</i>	229-87-8	<i>124</i>	302-79-4	<i>574</i>	345-78-8	<i>341</i>
147-93-3	<i>369</i>	229-87-8	<i>324</i>	302-95-4	<i>468</i>	350-03-8	<i>92</i>
147-94-4	<i>527</i>	230-27-3	<i>124</i>	303-07-1	<i>207</i>	350-46-9	<i>244</i>
148-24-3	<i>266</i>	244-69-9	<i>156</i>	303-21-9	<i>222</i>	352-32-9	<i>245</i>
148-82-3	<i>281</i>	248-13-5	<i>118</i>	303-43-5	<i>170</i>	352-70-5	<i>245</i>
149-29-1	<i>319</i>	253-52-1	<i>334</i>	303-81-1	<i>553</i>	352-93-2	<i>205</i>
149-30-4	<i>282</i>	253-66-7	<i>171</i>	303-97-9	<i>522</i>	352-97-6	<i>539</i>
149-32-6	<i>230</i>	253-82-7	<i>346</i>	303-98-0	<i>522</i>	355-02-2	<i>323</i>
149-44-0	<i>471</i>	254-79-5	<i>307</i>	304-06-3	<i>332</i>	355-42-0	<i>323</i>
149-45-1	<i>419</i>	259-56-3	<i>122</i>	304-20-1	<i>259</i>	355-68-0	<i>323</i>
149-74-6	<i>441</i>	259-57-4	<i>122</i>	304-21-2	<i>540</i>	357-57-3	<i>517</i>
149-91-7	<i>250</i>	259-79-0	<i>132</i>	304-55-2	<i>210</i>	358-23-6	<i>377</i>
150-13-0	<i>104</i>	259-84-7	<i>191</i>	304-59-6	<i>457</i>	358-71-4	<i>544</i>
150-25-4	<i>208</i>	260-94-6	<i>93</i>	304-88-1	<i>126</i>	360-65-6	<i>539</i>
150-30-1	<i>327</i>	271-44-3	<i>268</i>	304-91-6	<i>270</i>	361-37-6	<i>327</i>
150-39-0	<i>261</i>	271-63-6	<i>116</i>	305-01-1	<i>231</i>	363-03-1	<i>328</i>
150-46-9	<i>487</i>	271-89-6	<i>122</i>	305-03-3	<i>520</i>	363-72-4	<i>321</i>
150-60-7	<i>192</i>	272-13-2	<i>124</i>	305-33-9	<i>270</i>	366-18-7	<i>133</i>
150-61-8	<i>225</i>	273-09-6	<i>122</i>	305-84-0	<i>518</i>	366-29-0	<i>363</i>
150-68-5	<i>166</i>	273-40-5	<i>117</i>	306-07-0	<i>556</i>	367-12-4	<i>244</i>
150-76-5	<i>286</i>	273-77-8	<i>124</i>	306-67-2	<i>352</i>	367-51-1	<i>476</i>
150-78-7	<i>210</i>	274-07-7	<i>409</i>	309-00-2	<i>99</i>	368-39-8	<i>376</i>
150-86-7	<i>559</i>	274-40-8	<i>268</i>	311-03-5	<i>494</i>	369-07-3	<i>552</i>
150-90-3	<i>421</i>	274-76-0	<i>117</i>	311-28-4	<i>357</i>	372-09-8	<i>175</i>
151-10-0	<i>210</i>	275-51-4	<i>118</i>	311-89-7	<i>323</i>	372-75-8	<i>172</i>
151-21-3	<i>470</i>	275-94-5	<i>190</i>	312-30-1	<i>134</i>	372-97-4	<i>534</i>
151-50-8	<i>454</i>	280-57-9	<i>191</i>	315-22-0	<i>524</i>	373-44-4	<i>189</i>
151-56-4	<i>117</i>	280-64-8	<i>403</i>	316-42-7	<i>229</i>	374-12-9	<i>360</i>
151-63-3	<i>102</i>	281-23-2	<i>95</i>	316-46-1	<i>535</i>	375-22-4	<i>323</i>
152-02-3	<i>101</i>	286-20-4	<i>180</i>	316-66-5	<i>264</i>	375-96-2	<i>323</i>
152-93-2	<i>385</i>	286-62-4	<i>181</i>	319-84-6	<i>255</i>	378-44-9	<i>131</i>
153-18-4	<i>349</i>	287-27-4	<i>367</i>	326-91-0	<i>366</i>	379-68-0	<i>542</i>
153-78-6	<i>106</i>	287-92-3	<i>182</i>	327-56-0	<i>315</i>	379-79-3	<i>230</i>
153-98-0	<i>566</i>	288-13-1	<i>342</i>	327-57-1	<i>315</i>	381-73-7	<i>206</i>
154-17-6	<i>186</i>	288-32-4	<i>267</i>	327-62-8	<i>457</i>	382-45-6	<i>98</i>
154-42-7	<i>570</i>	288-88-0	<i>372</i>	327-97-9	<i>163</i>	383-63-1	<i>241</i>
156-06-9	<i>332</i>	288-94-8	<i>366</i>	328-42-7	<i>318</i>	383-64-2	<i>242</i>
156-10-5	<i>314</i>	290-37-9	<i>342</i>	328-50-7	<i>318</i>	389-08-2	<i>550</i>
156-38-7	<i>264</i>	291-21-4	<i>382</i>	329-56-6	<i>553</i>	392-12-1	<i>543</i>
156-39-8	<i>265</i>	291-64-5	<i>177</i>	329-71-5	<i>221</i>	392-56-3	<i>256</i>

404-86-4	<i>155</i>	464-48-2	<i>153</i>	486-35-1	<i>207</i>	501-30-4	<i>277</i>
405-50-5	<i>244</i>	464-49-3	<i>153</i>	487-21-8	<i>342</i>	501-52-0	<i>332</i>
405-79-8	<i>244</i>	464-72-2	<i>123</i>	487-51-4	<i>156</i>	501-53-1	<i>130</i>
407-24-9	<i>135</i>	466-81-9	<i>230</i>	487-79-6	<i>276</i>	501-58-6	<i>210</i>
407-25-0	<i>377</i>	467-15-2	<i>315</i>	487-90-1	<i>560</i>	501-65-5	<i>224</i>
407-41-0	<i>559</i>	467-62-9	<i>319</i>	488-23-3	<i>363</i>	501-97-3	<i>265</i>
408-35-5	<i>474</i>	468-28-0	<i>278</i>	488-43-7	<i>251</i>	501-98-4	<i>261</i>
420-04-2	<i>416</i>	469-59-0	<i>276</i>	488-81-3	<i>98</i>	502-41-0	<i>117</i>
420-37-1	<i>489</i>	470-38-2	<i>155</i>	488-93-7	<i>249</i>	502-42-1	<i>177</i>
420-56-4	<i>425</i>	470-82-6	<i>242</i>	490-02-8	<i>116</i>	502-49-8	<i>181</i>
421-06-7	<i>142</i>	471-46-5	<i>318</i>	490-11-9	<i>344</i>	502-65-8	<i>279</i>
422-05-9	<i>321</i>	472-70-8	<i>174</i>	490-55-1	<i>189</i>	502-72-7	<i>182</i>
425-75-2	<i>241</i>	472-86-6	<i>574</i>	490-79-9	<i>207</i>	502-74-2	<i>275</i>
429-06-1	<i>359</i>	472-93-5	<i>519</i>	490-83-5	<i>185</i>	503-17-3	<i>151</i>
429-06-1	<i>481</i>	473-30-3	<i>111</i>	491-33-8	<i>282</i>	503-28-6	<i>118</i>
429-07-2	<i>481</i>	473-34-7	<i>196</i>	491-35-0	<i>300</i>	503-29-7	<i>117</i>
429-42-5	<i>480</i>	474-07-7	<i>136</i>	491-37-2	<i>170</i>	503-30-0	<i>318</i>
431-03-8	<i>131</i>	474-25-9	<i>207</i>	492-11-5	<i>545</i>	503-49-1	<i>263</i>
431-06-1	<i>197</i>	474-62-4	<i>153</i>	492-17-1	<i>132</i>	504-02-9	<i>179</i>
432-68-8	<i>229</i>	475-31-0	<i>252</i>	492-22-8	<i>369</i>	504-15-4	<i>318</i>
434-13-9	<i>278</i>	475-38-7	<i>305</i>	492-27-3	<i>277</i>	504-17-6	<i>367</i>
434-16-2	<i>185</i>	476-28-8	<i>279</i>	492-37-5	<i>331</i>	504-20-1	<i>333</i>
437-11-6	<i>482</i>	476-66-4	<i>229</i>	492-62-6	<i>251</i>	504-24-5	<i>110</i>
438-08-4	<i>242</i>	477-29-2	<i>523</i>	492-70-6	<i>259</i>	504-29-0	<i>110</i>
438-23-3	<i>242</i>	477-30-5	<i>522</i>	492-94-4	<i>249</i>	504-31-4	<i>345</i>
441-38-3	<i>123</i>	477-73-6	<i>349</i>	492-99-9	<i>179</i>	504-63-2	<i>338</i>
442-16-0	<i>189</i>	478-61-5	<i>131</i>	493-01-6	<i>184</i>	505-22-6	<i>222</i>
442-51-3	<i>253</i>	478-95-5	<i>272</i>	493-02-7	<i>184</i>	505-23-7	<i>227</i>
443-72-1	<i>288</i>	479-18-5	<i>530</i>	493-52-7	<i>300</i>	505-48-6	<i>353</i>
445-29-4	<i>244</i>	479-23-2	<i>169</i>	494-03-1	<i>134</i>	505-54-4	<i>255</i>
445-38-9	<i>244</i>	479-27-6	<i>189</i>	494-38-2	<i>94</i>	505-66-8	<i>258</i>
446-72-0	<i>250</i>	479-59-4	<i>276</i>	495-08-9	<i>207</i>	505-95-3	<i>261</i>
447-31-4	<i>186</i>	479-61-8	<i>167</i>	495-48-7	<i>118</i>	506-03-6	<i>159</i>
448-37-3	<i>245</i>	480-16-0	<i>303</i>	495-69-2	<i>125</i>	506-12-7	<i>254</i>
448-71-5	<i>242</i>	480-22-8	<i>115</i>	495-84-1	<i>273</i>	506-30-9	<i>115</i>
451-13-8	<i>208</i>	480-41-1	<i>308</i>	496-11-7	<i>268</i>	506-46-7	<i>255</i>
451-40-1	<i>186</i>	480-44-4	<i>208</i>	496-16-2	<i>206</i>	506-59-2	<i>212</i>
452-58-4	<i>190</i>	480-63-7	<i>378</i>	496-41-3	<i>122</i>	506-64-9	<i>463</i>
452-86-8	<i>208</i>	480-66-0	<i>333</i>	496-67-3	<i>143</i>	506-65-0	<i>427</i>
454-41-1	<i>285</i>	480-68-2	<i>309</i>	496-78-6	<i>379</i>	506-68-3	<i>416</i>
454-49-9	<i>376</i>	481-21-0	<i>169</i>	497-19-8	<i>467</i>	506-78-5	<i>416</i>
456-22-4	<i>244</i>	481-26-5	<i>337</i>	497-23-4	<i>174</i>	506-96-7	<i>89</i>
456-27-9	<i>445</i>	481-29-8	<i>112</i>	497-25-6	<i>318</i>	507-02-8	<i>90</i>
458-09-3	<i>244</i>	481-39-0	<i>264</i>	497-38-1	<i>315</i>	507-19-7	<i>140</i>
458-35-5	<i>172</i>	481-42-5	<i>337</i>	497-59-6	<i>281</i>	507-20-0	<i>147</i>
458-37-7	<i>175</i>	481-74-3	<i>208</i>	497-76-7	<i>115</i>	507-25-5	<i>157</i>
461-06-3	<i>518</i>	482-05-3	<i>224</i>	498-02-2	<i>262</i>	507-28-8	<i>483</i>
461-58-5	<i>176</i>	482-74-6	<i>174</i>	498-21-5	<i>301</i>	507-32-4	<i>422</i>
461-72-3	<i>259</i>	482-89-3	<i>268</i>	498-23-7	<i>171</i>	508-04-3	<i>112</i>
462-06-6	<i>244</i>	483-04-5	<i>98</i>	498-24-8	<i>282</i>	508-75-8	<i>523</i>
462-08-8	<i>110</i>	483-55-6	<i>334</i>	498-59-9	<i>227</i>	509-14-8	<i>365</i>
462-10-2	<i>258</i>	483-65-8	<i>348</i>	498-66-8	<i>315</i>	510-13-4	<i>280</i>
462-18-0	<i>375</i>	484-11-7	<i>218</i>	498-94-2	<i>559</i>	510-64-5	<i>542</i>
462-94-2	<i>190</i>	484-50-0	<i>186</i>	499-06-9	<i>214</i>	511-08-0	<i>230</i>
463-49-0	<i>100</i>	485-35-8	<i>183</i>	499-83-2	<i>226</i>	511-09-1	<i>230</i>
463-51-4	<i>276</i>	485-47-2	<i>308</i>	500-05-0	<i>173</i>	511-18-2	<i>315</i>
463-58-1	<i>409</i>	485-49-4	<i>515</i>	500-22-1	<i>344</i>	512-04-9	<i>222</i>
463-82-1	<i>308</i>	485-71-2	<i>521</i>	500-38-9	<i>315</i>	512-35-6	<i>121</i>
464-10-8	<i>141</i>	485-89-2	<i>264</i>	500-40-3	<i>253</i>	512-47-0	<i>159</i>
464-43-7	<i>136</i>	486-25-9	<i>243</i>	500-99-2	<i>211</i>	512-56-1	<i>489</i>

594 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

512-69-6	348	525-27-9	147	531-99-7	91	539-35-5	366
513-35-9	290	525-64-4	243	532-02-5	473	539-47-9	249
513-36-0	271	525-79-1	536	532-18-3	220	539-64-0	216
513-38-2	272	525-82-6	243	532-27-4	159	539-88-8	239
513-44-0	298	526-08-9	332	532-32-1	466	540-04-5	334
513-48-4	269	526-26-1	478	532-40-1	570	540-05-6	344
513-53-1	143	526-31-8	302	532-54-7	273	540-23-8	371
513-81-5	214	526-55-6	383	532-55-8	125	540-37-4	269
513-85-9	143	526-75-0	218	532-78-5	156	540-38-5	270
513-86-0	260	526-78-3	193	532-82-1	170	540-54-5	340
514-10-3	81	526-94-3	472	533-01-7	414	540-59-0	198
514-35-5	152	526-98-7	276	533-30-2	104	540-67-0	240
514-58-2	574	526-99-6	250	533-48-2	528	540-69-2	393
514-76-1	116	527-07-1	471	533-58-4	270	540-72-7	476
514-78-3	154	527-17-3	229	533-62-0	262	540-80-7	150
515-30-0	116	527-52-6	529	533-67-5	186	540-84-1	378
515-42-4	466	527-53-7	363	533-73-3	121	540-88-5	144
515-74-2	475	527-54-8	379	533-75-5	382	541-14-0	518
515-84-4	241	527-60-6	379	533-87-9	99	541-15-1	518
516-05-2	294	527-61-7	214	533-90-4	408	541-41-3	235
516-12-1	270	527-62-8	105	534-00-9	139	541-43-5	398
517-09-9	230	527-72-0	368	534-07-6	196	541-47-9	211
517-10-2	172	527-76-4	111	534-15-6	81	541-59-3	280
517-23-7	88	527-85-5	370	534-17-8	410	541-73-1	196
517-28-2	541	528-29-0	220	534-22-5	293	541-88-8	160
517-51-1	366	528-44-9	377	534-52-1	221	542-05-2	84
518-05-8	305	528-45-0	221	534-85-0	105	542-11-0	113
518-28-5	560	528-50-7	158	534-94-1	104	542-18-7	162
518-82-1	229	528-71-2	264	534-97-4	251	542-28-9	384
519-05-1	318	528-76-7	220	535-11-5	235	542-32-5	103
519-23-3	532	529-19-1	372	535-15-9	236	542-55-2	272
519-44-8	222	529-21-5	240	535-46-6	400	542-59-6	87
519-62-0	167	529-23-7	103	535-80-8	161	542-69-8	149
519-73-3	381	529-34-0	362	535-83-1	377	542-90-5	241
519-88-0	194	529-44-2	550	535-87-5	188	542-92-7	182
520-26-3	255	529-59-9	250	536-10-6	325	543-21-5	89
520-33-2	255	529-64-6	382	536-40-3	161	543-24-8	90
520-36-5	377	529-69-1	544	536-69-6	536	543-27-1	271
520-85-4	263	529-86-2	114	536-74-3	327	543-38-4	154
521-11-9	262	530-14-3	335	536-75-4	241	543-59-9	112
521-24-4	444	530-36-9	105	536-90-3	113	543-63-5	405
521-31-3	278	530-43-8	520	537-00-8	410	543-80-6	397
521-35-7	154	530-44-9	212	537-17-7	103	543-88-4	407
521-74-4	193	530-48-3	225	537-47-3	332	543-94-2	478
522-12-3	346	530-55-2	210	537-65-5	189	544-01-4	271
522-66-7	542	530-57-4	355	537-92-8	86	544-15-0	395
522-75-8	368	530-59-6	261	538-14-7	91	544-17-2	407
523-21-7	460	530-62-1	157	538-28-3	129	544-25-2	177
523-27-3	192	530-75-6	92	538-51-2	129	544-40-1	195
523-44-4	447	530-85-8	313	538-56-7	171	544-47-8	162
523-67-1	438	530-93-8	362	538-58-9	191	544-62-7	118
524-36-7	563	531-14-6	523	538-62-5	224	544-63-8	304
524-38-9	265	531-53-3	118	538-74-9	192	544-76-3	255
524-42-5	306	531-55-5	118	538-75-0	201	544-92-3	416
524-63-0	175	531-57-7	118	538-81-8	224	545-26-6	537
524-64-4	189	531-72-6	509	539-03-7	160	546-06-5	172
524-80-1	156	531-81-7	173	539-12-8	339	546-56-5	446
524-97-0	342	531-84-0	255	539-15-1	259	546-67-8	434
525-03-1	243	531-85-1	121	539-17-3	103	546-68-9	481
525-05-3	446	531-91-9	224	539-30-0	128	546-74-7	470

546-89-4	435	555-75-9	391	574-25-4	547	586-75-4	137
547-25-1	383	556-10-5	313	574-66-3	123	586-76-5	137
547-57-9	469	556-32-1	438	574-93-6	452	586-77-6	138
547-58-0	441	556-33-2	206	574-98-1	138	586-78-7	140
548-04-9	267	556-50-3	252	575-44-0	208	586-89-0	88
548-24-3	422	556-52-5	252	575-47-3	465	586-96-9	314
548-42-5	98	556-56-9	101	576-19-2	516	587-04-2	161
548-43-6	229	556-58-9	179	576-24-9	199	587-85-9	420
548-62-9	174	556-63-8	436	576-26-1	218	587-90-6	221
548-65-2	319	556-65-0	437	576-36-3	250	587-98-4	440
548-93-6	260	556-67-2	316	577-11-7	469	589-02-6	400
549-38-2	291	556-67-2	446	577-19-5	140	589-15-1	137
549-56-4	347	556-88-7	311	577-56-0	88	589-16-2	234
550-33-4	563	556-89-8	315	577-61-9	316	589-17-3	137
550-44-7	298	556-91-2	392	577-71-9	222	589-18-4	372
550-57-2	265	556-99-0	382	577-85-5	261	589-21-9	141
550-60-7	312	557-01-7	266	578-57-4	136	589-34-4	293
550-74-3	335	557-09-5	497	578-66-5	111	589-90-2	215
551-06-4	307	557-11-9	102	578-68-7	110	589-92-4	291
551-16-6	109	557-17-5	299	578-76-7	293	589-93-5	279
552-16-9	310	557-21-1	498	578-86-9	208	590-18-1	144
552-23-8	314	557-24-4	280	578-95-0	94	590-28-3	453
552-32-9	308	557-41-5	498	579-10-2	287	590-73-8	216
552-38-5	436	557-59-5	358	579-18-0	125	590-92-1	141
552-46-5	306	557-66-4	234	579-43-1	259	591-08-2	93
552-62-5	303	557-99-3	90	579-92-0	224	591-22-0	279
552-79-4	292	558-13-4	156	580-13-2	140	591-23-1	291
552-86-3	249	558-17-8	149	580-15-4	111	591-27-5	109
552-89-6	309	558-32-7	482	580-17-6	110	591-31-1	113
553-12-8	562	560-09-8	153	581-05-5	546	591-35-5	200
553-17-3	253	561-07-9	186	581-40-8	217	591-49-1	291
553-24-2	308	561-20-6	152	581-42-0	217	591-50-4	269
553-26-4	133	562-48-1	295	581-64-6	368	591-54-8	110
553-39-9	263	562-90-3	462	581-89-5	312	591-55-9	110
553-53-7	308	563-41-7	351	581-96-4	306	591-76-4	293
553-54-8	435	563-63-3	462	582-25-2	453	591-80-0	100
553-70-8	437	563-68-8	483	582-52-5	528	591-87-7	100
553-79-7	313	563-79-1	214	582-60-5	213	591-93-5	320
553-90-2	217	563-80-4	290	583-39-1	282	592-04-1	439
553-97-9	289	563-83-7	272	583-59-5	291	592-13-2	216
554-00-7	196	565-00-4	153	583-61-9	278	592-20-1	86
554-01-8	291	565-59-3	217	583-75-5	104	592-31-4	303
554-05-2	498	565-63-9	113	583-78-8	199	592-41-6	257
554-12-1	299	565-80-0	209	583-93-7	190	592-42-7	255
554-13-2	435	566-58-5	100	584-02-1	322	592-57-4	177
554-68-7	375	566-78-9	87	584-04-3	209	592-76-7	254
554-70-1	487	569-77-7	342	584-08-7	453	592-84-7	148
554-73-4	382	568-93-4	309	584-13-4	111	592-85-8	440
554-84-7	312	571-61-9	217	584-48-5	138	593-45-3	316
554-91-6	250	571-92-6	533	584-84-9	571	593-51-1	288
554-95-0	121	572-09-8	84	584-85-0	114	593-56-6	285
555-06-6	465	572-09-8	505	585-07-9	150	593-81-7	378
555-16-8	309	440-17-5	572	585-47-7	120	593-85-1	253
555-21-5	310	573-56-8	221	585-48-8	195	593-95-3	409
555-30-6	530	573-57-9	206	585-76-2	137	594-02-2	489
555-31-7	391	573-58-0	172	585-79-5	140	594-15-0	373
555-44-2	380	573-83-1	457	585-84-2	93	594-18-3	193
555-45-3	379	573-97-7	140	585-99-9	281	594-20-7	200
555-48-6	252	574-15-2	122	586-11-8	222	594-27-4	485
555-68-0	310	574-16-3	122	586-38-9	285	594-36-5	112

596 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

594-44-5	231	606-20-2	222	615-28-1	329	622-47-9	297
594-61-6	263	606-22-4	220	615-43-0	269	622-51-5	372
594-65-0	373	606-23-5	268	615-57-6	192	622-78-6	129
595-30-2	154	606-26-8	309	615-58-7	193	622-97-9	301
595-40-4	275	606-35-9	379	615-67-8	163	623-05-2	260
595-46-0	217	606-40-6	165	615-93-0	197	623-12-1	160
595-49-3	222	606-68-8	551	616-02-4	171	623-24-5	193
595-90-4	483	606-81-5	221	616-06-8	315	623-25-6	201
596-03-2	193	606-83-7	226	616-13-7	112	623-26-7	201
596-38-3	333	607-24-9	312	616-29-5	190	623-27-8	355
597-12-6	281	607-34-1	313	616-38-5	214	623-33-6	252
597-43-3	219	607-57-8	311	616-38-6	419	623-42-7	290
597-71-7	320	607-93-2	373	616-44-4	301	623-49-4	236
597-80-8	401	608-33-3	193	616-47-7	293	623-51-8	241
598-04-9	195	608-37-7	193	616-75-1	129	623-65-4	319
598-10-7	183	608-40-2	219	617-33-4	236	623-70-1	236
598-31-2	136	608-66-2	229	617-35-6	241	623-73-4	236
598-50-5	303	608-71-9	320	617-45-8	116	623-91-6	204
598-53-8	275	608-91-3	374	617-48-1	280	624-03-3	238
598-55-0	290	608-92-4	374	617-52-7	217	624-31-7	270
598-58-3	295	609-19-8	375	617-73-2	264	624-38-4	208
598-75-4	290	609-72-3	220	617-84-5	204	624-60-2	292
598-78-7	167	609-89-2	199	617-86-7	488	624-64-6	144
599-04-2	542	609-93-3	221	617-89-0	249	624-65-7	339
599-66-6	227	609-99-4	222	618-00-4	199	624-73-7	208
599-67-7	225	610-02-6	377	618-36-0	289	624-91-9	296
600-37-3	262	610-27-5	313	618-39-3	119	624-92-0	215
601-89-8	313	610-28-6	221	618-41-7	120	625-45-6	285
602-01-7	222	610-30-3	220	618-51-9	269	625-47-8	165
602-09-5	132	610-39-9	222	618-95-1	296	625-51-4	82
602-38-0	221	610-54-8	221	619-04-5	214	625-52-5	303
602-55-1	327	610-71-9	192	619-05-6	188	625-53-6	241
602-60-8	309	610-72-0	214	619-15-8	222	625-77-4	187
602-94-8	321	610-94-6	289	619-17-0	110	626-05-1	193
602-99-3	380	611-01-8	213	619-19-1	313	626-15-3	193
603-00-9	265	611-08-5	315	619-23-8	310	626-35-7	240
603-02-1	222	611-32-5	300	619-31-8	217	626-39-1	373
603-11-2	313	611-33-6	168	619-42-1	289	626-48-2	302
603-12-3	221	611-34-7	110	619-50-1	296	626-51-7	293
603-32-7	492	611-35-8	168	619-58-9	269	626-56-2	298
603-33-8	493	611-71-2	281	619-65-8	175	626-58-4	298
603-34-9	380	611-73-4	125	619-67-0	259	626-60-8	168
603-35-0	493	611-95-0	125	619-73-8	310	626-61-9	168
603-36-1	492	611-97-2	214	619-84-1	212	626-64-2	266
603-45-2	349	611-98-3	188	620-22-4	372	626-97-1	384
603-54-3	226	612-14-6	120	620-24-6	260	627-06-5	303
603-71-4	311	612-42-0	330	620-40-6	373	627-11-2	163
603-79-2	213	612-60-2	300	620-45-1	418	627-20-3	322
604-09-1	265	612-62-4	168	620-50-8	215	627-21-4	322
604-35-3	170	612-71-5	381	621-02-3	227	627-83-8	238
604-44-4	165	612-94-2	331	621-09-0	224	627-84-9	238
604-53-5	132	612-95-3	332	621-23-8	378	627-93-0	211
604-59-1	550	612-98-6	314	621-37-4	264	627-95-2	111
604-68-2	251	613-12-7	288	621-42-1	83	628-02-4	155
604-69-3	251	613-31-0	206	621-59-0	275	628-13-7	344
604-88-6	255	613-78-5	307	621-66-9	262	628-28-4	150
605-49-1	196	614-75-5	264	621-82-9	171	628-41-1	178
605-60-7	314	614-80-2	83	621-84-1	127	628-63-7	111
605-69-6	221	615-18-9	162	622-15-1	225	628-63-7	322
605-94-7	522	615-20-3	161	622-21-9	201	628-81-9	148

629-04-9	<i>254</i>	637-44-5	<i>331</i>	687-38-7	<i>253</i>	765-43-5	<i>183</i>
629-05-0	<i>317</i>	637-59-2	<i>332</i>	688-71-1	<i>494</i>	766-39-2	<i>217</i>
629-11-8	<i>256</i>	637-69-4	<i>286</i>	688-73-3	<i>486</i>	767-00-0	<i>176</i>
629-14-1	<i>238</i>	637-84-3	<i>377</i>	688-74-4	<i>485</i>	767-15-7	<i>105</i>
629-20-9	<i>181</i>	637-87-6	<i>164</i>	690-02-8	<i>218</i>	767-92-0	<i>185</i>
629-25-4	<i>470</i>	637-88-7	<i>179</i>	691-38-3	<i>297</i>	768-32-1	<i>489</i>
629-31-2	<i>254</i>	637-92-3	<i>148</i>	693-02-7	<i>257</i>	768-33-2	<i>449</i>
629-41-4	<i>316</i>	638-32-4	<i>353</i>	693-23-2	<i>228</i>	768-90-1	<i>97</i>
629-45-8	<i>148</i>	638-38-0	<i>439</i>	693-65-2	<i>112</i>	768-91-2	<i>288</i>
629-54-9	<i>158</i>	638-53-9	<i>375</i>	693-72-1	<i>384</i>	768-92-3	<i>97</i>
629-59-4	<i>359</i>	638-95-9	<i>112</i>	693-98-1	<i>294</i>	768-93-4	<i>97</i>
629-70-9	<i>158</i>	639-58-7	<i>494</i>	694-59-7	<i>344</i>	768-94-5	<i>97</i>
629-96-9	<i>115</i>	640-19-7	<i>244</i>	695-34-1	<i>108</i>	768-95-6	<i>96</i>
629-97-0	<i>228</i>	640-60-8	<i>333</i>	696-54-8	<i>344</i>	769-42-6	<i>213</i>
630-01-3	<i>255</i>	641-13-4	<i>115</i>	696-62-8	<i>269</i>	769-92-6	<i>146</i>
630-02-4	<i>316</i>	641-36-1	<i>513</i>	697-82-5	<i>379</i>	770-12-7	<i>449</i>
630-08-0	<i>409</i>	641-70-3	<i>313</i>	697-91-6	<i>197</i>	770-71-8	<i>262</i>
630-10-4	<i>462</i>	642-31-9	<i>114</i>	698-69-1	<i>212</i>	771-29-9	<i>362</i>
630-18-2	<i>147</i>	642-44-4	<i>101</i>	700-12-9	<i>322</i>	771-51-7	<i>268</i>
630-60-4	<i>318</i>	642-78-4	<i>521</i>	700-13-0	<i>378</i>	771-61-9	<i>321</i>
631-36-7	<i>481</i>	643-13-0	<i>536</i>	700-16-3	<i>321</i>	771-97-1	<i>189</i>
631-40-3	<i>366</i>	643-43-6	<i>222</i>	700-56-1	<i>288</i>	772-14-5	<i>328</i>
631-61-8	<i>392</i>	643-79-8	<i>333</i>	700-57-2	<i>96</i>	772-15-6	<i>328</i>
632-12-2	<i>544</i>	644-32-6	<i>126</i>	700-58-3	<i>96</i>	772-33-8	<i>264</i>
632-22-4	<i>365</i>	644-97-3	<i>449</i>	701-99-5	<i>326</i>	773-76-2	<i>198</i>
632-25-7	<i>354</i>	645-48-7	<i>332</i>	703-80-0	<i>90</i>	775-12-2	<i>420</i>
632-46-2	<i>214</i>	645-49-8	<i>353</i>	704-00-7	<i>187</i>	776-34-1	<i>109</i>
632-51-9	<i>365</i>	645-96-5	<i>399</i>	704-15-4	<i>253</i>	779-02-2	<i>288</i>
632-52-0	<i>365</i>	646-06-0	<i>223</i>	706-35-2	<i>313</i>	781-43-1	<i>213</i>
632-56-4	<i>360</i>	646-04-8	<i>322</i>	708-06-5	<i>263</i>	781-73-7	<i>89</i>
632-69-9	<i>460</i>	646-31-1	<i>358</i>	708-79-2	<i>548</i>	782-92-3	<i>126</i>
632-92-8	<i>380</i>	652-04-0	<i>289</i>	709-49-9	<i>269</i>	784-04-3	<i>88</i>
632-93-9	<i>195</i>	653-89-4	<i>547</i>	709-63-7	<i>377</i>	789-25-3	<i>493</i>
632-99-5	<i>349</i>	657-26-1	<i>280</i>	712-74-3	<i>359</i>	791-28-6	<i>493</i>
633-03-4	<i>136</i>	657-27-2	<i>280</i>	716-53-0	<i>160</i>	791-31-1	<i>494</i>
633-65-8	<i>131</i>	657-84-1	<i>476</i>	719-22-2	<i>194</i>	793-41-9	<i>111</i>
633-71-6	<i>263</i>	658-78-6	<i>313</i>	723-62-6	<i>114</i>	804-63-7	<i>347</i>
633-96-5	<i>447</i>	660-68-4	<i>202</i>	725-12-2	<i>225</i>	811-54-4	<i>434</i>
633-99-8	<i>165</i>	660-88-8	<i>111</i>	732-26-3	<i>373</i>	811-93-8	<i>298</i>
634-35-5	<i>347</i>	661-19-8	<i>228</i>	734-77-0	<i>132</i>	813-19-4	<i>427</i>
634-36-6	<i>378</i>	661-69-8	<i>428</i>	741-03-7	<i>359</i>	814-68-6	<i>95</i>
634-65-1	<i>219</i>	665-66-7	<i>97</i>	751-38-2	<i>365</i>	814-80-2	<i>408</i>
634-66-2	<i>358</i>	666-99-9	<i>98</i>	754-05-2	<i>492</i>	814-81-3	<i>415</i>
634-67-3	<i>374</i>	670-54-2	<i>359</i>	754-34-7	<i>323</i>	814-95-9	<i>478</i>
634-83-3	<i>358</i>	672-15-1	<i>258</i>	758-96-3	<i>219</i>	818-08-6	<i>418</i>
634-90-2	<i>358</i>	673-40-5	<i>137</i>	759-24-0	<i>203</i>	818-88-2	<i>350</i>
634-93-5	<i>374</i>	673-84-7	<i>217</i>	760-21-4	<i>235</i>	819-83-0	<i>421</i>
635-21-1	<i>104</i>	674-26-0	<i>548</i>	760-32-7	<i>418</i>	821-06-7	<i>192</i>
635-51-8	<i>332</i>	674-76-0	<i>297</i>	760-93-0	<i>283</i>	821-48-7	<i>133</i>
635-65-4	<i>132</i>	674-82-8	<i>209</i>	761-35-3	<i>531</i>	822-23-1	<i>316</i>
635-78-9	<i>460</i>	674-82-8	<i>276</i>	761-65-9	<i>195</i>	822-36-6	<i>294</i>
635-93-8	<i>169</i>	675-20-7	<i>384</i>	762-21-0	<i>202</i>	822-67-3	<i>180</i>
636-04-4	<i>367</i>	676-97-1	<i>441</i>	762-62-9	<i>218</i>	823-18-7	<i>178</i>
636-09-0	<i>205</i>	677-53-6	<i>367</i>	762-72-1	<i>390</i>	823-76-7	<i>89</i>
636-30-6	<i>374</i>	677-69-0	<i>254</i>	763-29-1	<i>297</i>	823-96-1	<i>488</i>
636-70-4	<i>375</i>	680-31-9	<i>428</i>	763-10-0	<i>537</i>	824-07-9	<i>354</i>
636-73-7	<i>344</i>	681-84-5	<i>482</i>	764-13-6	<i>216</i>	824-35-1	<i>408</i>
636-98-6	<i>269</i>	684-16-2	<i>255</i>	764-35-2	<i>258</i>	824-72-6	<i>449</i>
637-01-4	<i>364</i>	685-87-0	<i>203</i>	764-85-2	<i>319</i>	824-79-3	<i>476</i>
637-39-8	<i>375</i>	685-91-6	<i>202</i>	765-30-0	<i>183</i>	824-94-2	<i>285</i>

598 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

825-90-1	<i>474</i>	880-52-4	<i>96</i>	960-71-4	<i>493</i>	1074-82-4	<i>457</i>
826-73-3	<i>124</i>	882-06-4	<i>311</i>	961-07-9	<i>528</i>	1076-22-8	<i>303</i>
826-74-4	<i>385</i>	882-33-7	<i>225</i>	961-68-2	<i>221</i>	1076-38-6	<i>261</i>
826-81-3	<i>263</i>	883-99-8	<i>293</i>	971-74-4	<i>351</i>	1076-43-3	<i>120</i>
827-19-0	<i>469</i>	886-38-4	<i>225</i>	975-17-7	<i>330</i>	1077-28-7	<i>545</i>
827-21-4	<i>469</i>	890-38-0	<i>528</i>	977-96-8	<i>203</i>	1079-66-9	<i>411</i>
827-52-1	<i>180</i>	893-33-4	<i>306</i>	979-92-0	<i>509</i>	1079-71-6	<i>316</i>
827-54-3	<i>307</i>	902-47-6	<i>221</i>	981-40-8	<i>405</i>	1081-34-1	<i>356</i>
828-01-3	<i>265</i>	914-45-4	<i>357</i>	989-38-8	<i>348</i>	1081-34-1	<i>568</i>
828-51-3	<i>96</i>	915-35-5	<i>254</i>	992-59-6	<i>400</i>	1083-48-3	<i>310</i>
830-03-5	<i>312</i>	917-23-7	<i>366</i>	993-13-5	<i>441</i>	1085-92-3	<i>417</i>
830-09-1	<i>285</i>	917-61-3	<i>468</i>	994-32-1	<i>488</i>	1086-80-2	<i>278</i>
830-81-9	<i>306</i>	917-70-4	<i>434</i>	994-89-8	<i>423</i>	1088-56-8	<i>546</i>
830-96-6	<i>268</i>	919-62-0	<i>488</i>	995-45-9	<i>486</i>	1094-61-7	<i>552</i>
831-61-8	<i>239</i>	920-66-1	<i>256</i>	996-50-9	<i>418</i>	1095-90-5	<i>547</i>
835-64-3	<i>264</i>	921-01-7	<i>525</i>	996-70-3	<i>362</i>	1099-45-2	<i>422</i>
836-30-6	<i>311</i>	921-53-9	<i>457</i>	998-07-2	<i>531</i>	1100-88-5	<i>400</i>
838-85-7	<i>420</i>	922-32-7	<i>523</i>	998-30-1	<i>486</i>	1107-26-2	<i>513</i>
840-78-8	<i>135</i>	922-67-8	<i>298</i>	998-40-3	<i>486</i>	1109-11-1	<i>513</i>
842-07-9	<i>120</i>	923-73-9	<i>134</i>	999-90-6	<i>151</i>	1112-67-0	<i>357</i>
845-10-3	<i>469</i>	924-49-0	<i>428</i>	999-97-3	<i>428</i>	1113-41-3	<i>556</i>
845-46-5	<i>469</i>	924-49-2	<i>106</i>	1000-84-6	<i>233</i>	1113-59-3	<i>141</i>
846-70-8	<i>443</i>	926-39-6	<i>106</i>	1001-53-2	<i>89</i>	1114-34-7	<i>280</i>
852-38-0	<i>516</i>	928-04-1	<i>390</i>	1002-62-6	<i>468</i>	1114-41-6	<i>549</i>
853-35-0	<i>465</i>	928-49-4	<i>258</i>	1002-84-2	<i>320</i>	1114-81-4	<i>559</i>
853-68-9	<i>395</i>	930-88-1	<i>294</i>	1003-03-8	<i>182</i>	1115-84-0	<i>295</i>
860-22-0	<i>431</i>	931-19-1	<i>335</i>	1003-67-4	<i>335</i>	1116-54-7	<i>314</i>
863-57-0	<i>471</i>	931-88-4	<i>181</i>	1003-73-2	<i>335</i>	1116-76-3	<i>380</i>
865-34-9	<i>436</i>	932-90-1	<i>119</i>	1003-78-7	<i>219</i>	1116-98-9	<i>147</i>
865-44-1	<i>432</i>	933-18-6	<i>486</i>	1005-24-9	<i>155</i>	1118-02-1	<i>491</i>
865-47-4	<i>453</i>	933-52-8	<i>363</i>	1005-56-7	<i>329</i>	1118-12-3	<i>151</i>
865-48-5	<i>467</i>	934-34-9	<i>260</i>	1006-94-6	<i>286</i>	1119-34-2	<i>115</i>
866-83-1	<i>454</i>	935-30-8	<i>155</i>	1007-32-5	<i>128</i>	1119-94-4	<i>228</i>
866-97-7	<i>356</i>	935-31-9	<i>177</i>	1007-42-7	<i>258</i>	1119-97-7	<i>359</i>
867-13-0	<i>487</i>	935-56-8	<i>97</i>	1008-72-6	<i>399</i>	1120-01-0	<i>471</i>
867-81-2	<i>555</i>	935-92-2	<i>378</i>	1009-61-6	<i>187</i>	1120-02-1	<i>446</i>
868-14-4	<i>455</i>	935-95-5	<i>358</i>	1009-93-4	<i>428</i>	1120-44-1	<i>415</i>
868-19-9	<i>478</i>	937-14-4	<i>166</i>	1010-95-3	<i>548</i>	1120-49-6	<i>201</i>
868-77-9	<i>261</i>	938-55-6	<i>213</i>	1010-98-1	<i>445</i>	1121-60-4	<i>344</i>
869-24-9	<i>169</i>	938-79-4	<i>328</i>	1012-72-2	<i>194</i>	1122-17-4	<i>198</i>
869-29-4	<i>95</i>	938-85-2	<i>293</i>	1020-31-1	<i>194</i>	1122-58-3	<i>213</i>
870-46-2	<i>147</i>	939-26-4	<i>139</i>	1024-41-5	<i>131</i>	1123-40-6	<i>216</i>
870-50-8	<i>194</i>	939-52-6	<i>166</i>	1024-99-3	<i>544</i>	1123-54-2	<i>116</i>
872-35-5	<i>282</i>	940-31-8	<i>326</i>	1025-70-4	<i>478</i>	1124-13-6	<i>176</i>
872-36-6	<i>385</i>	940-71-6	<i>450</i>	1025-99-8	<i>563</i>	1126-34-7	<i>473</i>
872-50-4	<i>300</i>	941-98-0	<i>91</i>	1032-65-1	<i>527</i>	1126-71-2	<i>218</i>
872-85-5	<i>344</i>	943-89-5	<i>285</i>	1034-39-5	<i>493</i>	1126-79-0	<i>151</i>
872-93-5	<i>301</i>	946-82-7	<i>179</i>	1041-01-6	<i>530</i>	1128-87-6	<i>84</i>
873-32-5	<i>161</i>	947-04-6	<i>116</i>	1046-56-6	<i>345</i>	1132-21-4	<i>210</i>
873-55-2	<i>400</i>	947-42-2	<i>420</i>	1048-08-4	<i>483</i>	1132-68-9	<i>535</i>
873-66-5	<i>301</i>	947-84-2	<i>132</i>	1066-26-8	<i>464</i>	1134-35-6	<i>214</i>
873-69-8	<i>344</i>	948-60-7	<i>563</i>	1066-54-2	<i>423</i>	1135-24-6	<i>262</i>
873-74-5	<i>104</i>	951-55-3	<i>302</i>	1068-57-1	<i>83</i>	1135-66-6	<i>272</i>
873-83-6	<i>512</i>	951-77-9	<i>527</i>	1068-90-2	<i>202</i>	1137-41-3	<i>104</i>
874-14-6	<i>220</i>	951-78-0	<i>528</i>	1069-79-0	<i>531</i>	1137-42-4	<i>126</i>
874-23-7	<i>89</i>	952-23-8	<i>188</i>	1070-89-9	<i>466</i>	1139-30-6	<i>157</i>
874-42-0	<i>196</i>	957-75-5	<i>517</i>	1071-23-4	<i>558</i>	1141-59-9	<i>345</i>
877-24-7	<i>455</i>	958-80-5	<i>211</i>	1072-72-6	<i>362</i>	1143-74-4	<i>126</i>
877-44-1	<i>376</i>	959-10-4	<i>133</i>	1072-86-2	<i>178</i>	1148-79-4	<i>356</i>
879-39-0	<i>358</i>	959-28-4	<i>191</i>	1072-98-6	<i>105</i>	1149-16-2	<i>253</i>

1150-62-5	329	1301-96-8	463	1474-78-8	487	1663-45-2	402
1155-00-6	135	1303-96-4	466	1476-23-9	102	1666-13-3	420
1163-67-3	550	1305-62-0	407	1476-53-5	553	1667-99-8	412
1170-02-1	237	1309-64-4	396	1481-70-5	383	1668-10-6	252
1173-88-2	554	1310-53-8	426	1482-97-9	190	1670-46-8	89
1176-74-5	357	1310-58-3	455	1484-09-9	274	1674-10-8	215
1177-87-3	187	1310-65-2	436	1484-12-4	290	1676-73-9	251
1180-95-6	568	1310-66-3	436	1484-13-5	385	1678-91-7	236
1185-53-1	382	1310-73-2	472	1484-28-2	139	1689-82-3	328
1185-55-3	442	1311-10-0	478	1492-08-6	549	1694-31-1	145
1186-49-8	471	1312-43-2	432	1492-24-6	104	1694-92-4	393
1187-58-2	298	1313-27-5	443	1493-13-6	302	1696-20-4	91
1187-84-4	291	1313-82-2	475	1498-88-0	378	1699-51-0	277
1188-37-0	90	1313-84-4	475	1499-10-1	224	1701-93-5	464
1188-58-5	227	1313-97-9	444	1499-17-8	449	1707-03-5	420
1188-92-7	488	1314-56-3	451	1502-03-0	177	1707-75-1	226
1188-95-0	380	1314-80-3	451	1502-06-3	177	1707-77-3	209
1189-94-2	548	1314-85-8	451	1503-53-3	88	1709-54-2	130
1191-50-0	475	1317-36-8	434	1515-76-0	86	1710-98-1	146
1191-95-3	176	1324-63-6	386	1518-16-7	359	1717-50-6	360
1192-40-1	414	1327-53-3	397	1522-22-1	256	1720-32-7	225
1192-52-5	198	1330-20-7	387	1523-11-1	306	1722-12-9	168
1193-92-6	344	1330-43-4	466	1528-74-1	221	1722-26-5	404
1194-65-6	197	1330-78-5	495	1530-32-1	423	1723-00-8	559
1195-16-0	90	1332-52-1	400	1530-45-6	422	1730-91-2	290
1195-45-5	244	1333-74-0	429	1544-68-9	245	1733-12-6	173
1197-55-3	109	1333-82-0	413	1552-12-1	181	1739-84-0	216
1198-27-2	109	1344-09-8	475	1562-94-3	118	1750-12-5	106
1198-33-0	303	1344-28-1	390	1564-64-3	136	1754-47-8	420
1199-03-7	347	1344-58-7	496	1565-80-6	289	1758-73-2	246
1200-93-7	281	1393-62-0	505	1569-69-3	180	1762-95-4	395
1202-34-2	133	1393-62-0	510	1570-64-5	164	1763-10-6	555
1202-34-2	226	1395-18-2	117	1570-65-6	199	1772-03-8	250
1204-06-4	543	1400-61-9	554	1571-33-1	449	1779-48-2	449
1207-25-0	418	1405-69-2	513	1574-41-0	320	1779-49-3	442
1207-72-3	297	1405-87-4	514	1576-35-8	370	1779-58-4	440
1210-12-4	175	1414-45-5	552	1591-31-7	269	1783-81-9	294
1212-08-4	328	1429-59-0	531	1600-27-7	439	1787-61-7	422
1214-39-7	514	1435-55-8	542	1600-44-8	364	1798-04-5	151
1215-59-4	130	1435-88-7	262	1602-97-7	158	1798-09-0	286
1217-45-4	201	1436-34-6	150	1603-40-3	108	1805-32-9	197
1220-94-6	107	1443-80-7	88	1603-41-4	108	1806-29-7	207
1224-69-7	135	1445-69-8	334	1606-67-3	110	1811-28-5	188
1226-46-6	368	1445-73-4	298	1608-26-0	428	1817-74-9	135
1235-21-8	390	1446-61-3	185	1609-47-8	205	1820-81-1	169
1239-45-8	533	1450-14-2	428	1609-86-5	149	1821-12-1	329
1245-13-2	133	1453-82-3	272	1611-35-4	387	1822-51-1	164
1260-17-9	518	1454-85-9	254	1615-02-7	162	1824-81-3	108
1271-19-8	485	1455-77-2	190	1618-26-4	245	1826-28-4	332
1271-28-9	445	1460-02-2	373	1632-73-1	242	1832-07-1	450
1271-42-7	424	1460-57-7	178	1633-22-3	319	1833-53-0	459
1271-51-8	496	1461-15-0	406	1634-04-4	150	1846-92-2	233
1271-55-2	390	1461-22-9	486	1637-39-4	577	1850-14-2	364
1287-13-4	461	1461-25-2	481	1641-41-1	262	1863-63-4	392
1291-32-3	499	1464-44-4	557	1643-19-2	357	1866-15-5	508
1291-48-1	424	1465-25-4	307	1643-20-5	228	1866-16-6	517
1293-87-4	242	1466-76-8	210	1653-19-6	197	1871-22-3	136
1294-74-2	434	1468-83-3	93	1655-29-4	421	1871-96-1	350
1295-35-8	401	1470-61-7	463	1656-44-6	220	1873-77-4	495
1301-19-9	470	1470-94-6	262	1662-01-7	118	1875-48-5	110

600 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

1878-65-5	<i>166</i>	2140-46-7	<i>542</i>	2412-14-8	<i>369</i>	2658-24-4	<i>215</i>
1878-66-6	<i>166</i>	2150-48-3	<i>459</i>	2414-98-4	<i>437</i>	2679-01-8	<i>292</i>
1878-68-8	<i>141</i>	2152-56-9	<i>115</i>	2425-51-6	<i>435</i>	2683-82-1	<i>316</i>
1878-87-1	<i>312</i>	2155-71-7	<i>194</i>	2425-95-8	<i>133</i>	2687-43-6	<i>129</i>
1878-91-7	<i>140</i>	2158-04-5	<i>325</i>	2435-53-2	<i>358</i>	2688-77-9	<i>277</i>
1910-42-5	<i>214</i>	2161-31-1	<i>329</i>	2436-90-0	<i>172</i>	2696-92-6	<i>446</i>
1913-93-5	<i>369</i>	2170-03-8	<i>275</i>	2436-96-6	<i>221</i>	2700-22-3	<i>129</i>
1918-77-0	<i>368</i>	2176-62-7	<i>320</i>	2440-42-8	<i>423</i>	2712-78-9	<i>135</i>
1921-70-6	<i>364</i>	2179-57-9	<i>101</i>	2444-68-0	<i>385</i>	2739-12-0	<i>288</i>
1923-70-2	<i>357</i>	2181-42-2	<i>379</i>	2451-01-6	<i>356</i>	2743-38-6	<i>192</i>
1928-38-7	<i>291</i>	2183-17-7	<i>444</i>	2457-47-8	<i>200</i>	2744-08-3	<i>184</i>
1936-15-8	<i>447</i>	2183-17-7	<i>550</i>	2457-80-9	<i>528</i>	2744-09-4	<i>184</i>
1937-54-8	<i>352</i>	2197-63-9	<i>419</i>	2465-27-2	<i>116</i>	2749-11-3	<i>99</i>
1941-27-1	<i>357</i>	2208-07-3	<i>233</i>	2466-76-4	<i>90</i>	2757-04-2	<i>329</i>
1941-30-6	<i>366</i>	2213-63-0	<i>200</i>	2475-44-7	<i>135</i>	2763-96-4	<i>549</i>
1945-77-3	<i>441</i>	2216-12-8	<i>311</i>	2475-93-6	<i>336</i>	2765-29-9	<i>177</i>
1948-33-0	<i>149</i>	2216-33-3	<i>296</i>	2482-00-0	<i>98</i>	2767-84-2	<i>154</i>
1949-89-9	<i>186</i>	2216-51-5	<i>282</i>	2487-90-3	<i>488</i>	2768-90-3	<i>336</i>
1972-28-7	<i>202</i>	2216-69-5	<i>295</i>	2492-02-9	<i>141</i>	2769-72-4	<i>149</i>
1983-26-2	<i>411</i>	2217-07-4	<i>226</i>	2492-87-7	<i>553</i>	2782-07-2	<i>250</i>
1985-12-2	<i>141</i>	2218-94-2	<i>312</i>	2494-56-6	<i>517</i>	2797-68-4	<i>531</i>
1988-89-2	<i>330</i>	2219-31-0	<i>518</i>	2498-20-6	<i>356</i>	2799-21-5	<i>265</i>
1989-52-2	<i>170</i>	2227-29-4	<i>419</i>	2498-66-0	<i>119</i>	2818-58-8	<i>557</i>
1990-29-0	<i>102</i>	2235-54-3	<i>393</i>	2504-11-2	<i>506</i>	2835-81-6	<i>104</i>
2001-45-8	<i>483</i>	2236-60-4	<i>342</i>	2507-55-3	<i>266</i>	2835-82-7	<i>104</i>
2001-95-8	<i>573</i>	2240-25-7	<i>138</i>	2508-19-2	<i>380</i>	2835-98-5	<i>106</i>
2002-24-6	<i>106</i>	2243-27-8	<i>319</i>	2511-17-3	<i>482</i>	2835-99-6	<i>106</i>
2004-06-0	<i>520</i>	2243-42-7	<i>326</i>	2516-34-9	<i>176</i>	2836-32-0	<i>471</i>
2004-14-0	<i>437</i>	2243-62-1	<i>189</i>	2524-64-3	<i>449</i>	2840-28-0	<i>105</i>
2004-70-8	<i>320</i>	2243-76-7	<i>99</i>	2533-00-8	<i>318</i>	2851-68-5	<i>344</i>
2009-64-5	<i>551</i>	2243-82-5	<i>305</i>	2537-36-2	<i>363</i>	2857-97-8	<i>405</i>
2018-61-3	<i>91</i>	2244-11-3	<i>100</i>	2537-36-2	<i>482</i>	2861-02-1	<i>395</i>
2026-48-4	<i>107</i>	2244-40-8	<i>469</i>	2538-85-4	<i>422</i>	2861-99-6	<i>380</i>
2030-63-9	<i>172</i>	2246-46-0	<i>367</i>	2549-67-9	<i>238</i>	2869-83-2	<i>275</i>
2031-12-9	<i>316</i>	2247-00-9	<i>402</i>	2552-55-8	<i>542</i>	2892-51-5	<i>352</i>
2031-67-6	<i>442</i>	2290-65-5	<i>491</i>	2554-06-5	<i>482</i>	2916-68-9	<i>490</i>
2035-66-7	<i>448</i>	2292-79-7	<i>187</i>	2563-97-5	<i>373</i>	2916-76-9	<i>442</i>
2039-82-9	<i>142</i>	2294-76-0	<i>112</i>	2564-83-2	<i>365</i>	2922-54-5	<i>471</i>
2043-06-3	<i>516</i>	2305-32-0	<i>178</i>	2567-83-1	<i>359</i>	2923-16-2	<i>458</i>
2044-56-6	<i>436</i>	2305-34-2	<i>375</i>	2578-28-1	<i>566</i>	2923-18-4	<i>476</i>
2045-00-3	<i>396</i>	2309-49-1	<i>365</i>	2578-45-2	<i>162</i>	2923-28-6	<i>464</i>
2050-76-2	<i>199</i>	2321-07-5	<i>243</i>	2578-57-6	<i>338</i>	2935-35-5	<i>330</i>
2051-85-6	<i>207</i>	2338-20-7	<i>377</i>	2591-17-5	<i>546</i>	2937-50-0	<i>101</i>
2067-33-6	<i>143</i>	2348-71-2	<i>570</i>	2592-95-2	<i>260</i>	2949-92-0	<i>295</i>
2068-78-2	<i>574</i>	2361-24-2	<i>338</i>	2605-67-6	<i>440</i>	2950-43-8	<i>431</i>
2082-84-0	<i>185</i>	2361-96-8	<i>92</i>	2610-05-1	<i>410</i>	2957-95-1	<i>197</i>
2086-83-1	<i>131</i>	2370-18-5	<i>250</i>	2611-82-7	<i>443</i>	2961-50-4	<i>112</i>
2090-05-3	<i>407</i>	2370-63-0	<i>233</i>	2620-63-5	<i>90</i>	2963-79-3	<i>143</i>
2091-46-5	<i>458</i>	2370-88-9	<i>482</i>	2622-14-2	<i>486</i>	2966-50-9	<i>464</i>
2092-17-3	<i>399</i>	2373-80-0	<i>292</i>	2622-26-6	<i>323</i>	2985-59-3	<i>261</i>
2092-55-9	<i>352</i>	2379-55-7	<i>219</i>	2623-36-1	<i>472</i>	2996-92-1	<i>450</i>
2094-98-6	<i>117</i>	2385-77-5	<i>172</i>	2627-86-3	<i>289</i>	3001-72-7	<i>190</i>
2107-69-9	<i>210</i>	2386-53-0	<i>470</i>	2627-95-4	<i>422</i>	3002-78-6	<i>297</i>
2116-65-6	<i>130</i>	2386-60-9	<i>143</i>	2635-26-9	<i>206</i>	3007-31-6	<i>201</i>
2121-67-7	<i>216</i>	2387-71-5	<i>115</i>	2636-31-9	<i>276</i>	3011-89-0	<i>165</i>
2122-61-4	<i>105</i>	2390-54-7	<i>367</i>	2637-34-5	<i>344</i>	3012-37-1	<i>130</i>
2127-03-9	<i>226</i>	2396-60-3	<i>285</i>	2642-52-6	<i>302</i>	3016-19-1	<i>217</i>
2131-55-7	<i>167</i>	2402-78-0	<i>200</i>	2644-70-4	<i>429</i>	3018-12-0	<i>196</i>
2136-75-6	<i>493</i>	2403-88-5	<i>266</i>	2646-71-1	<i>552</i>	3019-22-5	<i>274</i>
2138-24-1	<i>361</i>	2411-89-4	<i>173</i>	2650-53-5	<i>257</i>	3022-16-0	<i>134</i>

3024-83-7	525	3458-28-4	281	4023-65-8	93	4525-33-1	419
3025-95-4	88	3460-18-2	193	4023-34-1	183	4542-57-8	228
3029-19-4	343	3463-36-3	311	4033-27-6	529	4547-57-3	144
3029-30-9	201	3468-11-9	208	4036-30-1	332	4548-45-2	166
3029-40-1	225	3473-63-0	246	4041-95-6	195	4551-15-9	449
3051-09-0	304	3481-20-7	358	4042-36-8	345	4552-64-1	369
3054-95-3	95	3522-94-9	378	4043-87-2	336	4562-27-0	266
3068-32-4	505	3535-83-9	360	4044-65-9	330	4568-71-2	128
3076-34-4	474	3562-69-4	379	4050-45-7	257	4611-62-5	422
3076-91-3	440	3564-26-9	477	4052-59-9	512	4618-18-2	277
3084-10-4	483	3575-31-3	317	4054-38-0	177	4628-94-8	402
3087-16-9	434	3588-17-8	304	4067-16-7	321	4636-73-1	376
3097-08-3	437	3588-91-8	202	4071-88-9	423	4648-54-8	490
3102-56-5	530	3603-99-4	183	4075-81-4	408	4656-81-9	361
3105-95-1	559	3604-87-3	229	4088-22-6	430	6459-45-4	197
3109-63-5	357	3604-90-8	521	4097-89-6	381	4675-64-3	455
3112-31-0	342	3608-75-1	308	4111-54-0	435	4695-62-9	242
3118-85-2	251	3612-20-2	130	4113-06-8	376	4696-76-8	545
3141-04-6	180	3615-37-0	246	4113-12-6	158	4702-13-0	334
3143-02-0	296	3615-41-6	348	4114-31-2	235	4707-71-5	324
3144-16-9	154	3618-43-7	497	4130-42-1	195	4712-70-3	136
3147-14-6	153	3625-57-8	308	4156-16-5	240	4733-39-5	215
3150-24-1	552	3632-91-5	437	4199-10-4	561	4746-97-8	179
3153-26-2	496	3634-56-8	433	4199-88-6	312	4781-83-3	543
3167-49-5	109	3634-83-1	387	4205-91-8	521	4786-20-3	174
3173-56-6	129	3680-02-2	302	4206-67-1	433	4792-83-0	118
3182-95-4	327	3682-32-4	314	4221-98-1	324	4819-77-6	375
3184-13-2	318	3682-35-7	381	4223-11-4	385	4839-46-7	216
3205-94-5	182	3688-79-7	285	4233-33-4	333	4845-50-5	208
3206-73-3	545	3699-66-9	487	4254-02-8	182	4845-99-2	517
3219-63-4	491	3700-67-2	222	4264-83-9	421	4900-63-4	286
3240-34-4	187	3710-27-8	444	4265-07-0	558	4901-51-3	358
3244-41-5	458	3721-95-7	176	4271-30-1	104	4916-57-8	226
3251-23-8	415	3722-51-8	267	4272-74-6	571	4930-98-7	259
3264-82-2	444	3733-85-5	319	4286-15-1	328	4942-47-6	96
3277-26-7	482	3734-67-6	452	4294-16-0	514	4961-10-4	376
3279-54-7	421	3737-95-9	153	4316-93-2	199	5006-66-6	266
3282-30-2	337	3739-30-8	262	4328-13-6	361	5015-75-8	467
3282-56-2	150	3739-38-6	326	4329-75-3	226	5056-14-4	174
3282-73-3	201	3739-64-8	101	4337-75-1	434	5061-35-8	506
3284-79-5	166	3740-52-1	312	4350-09-8	267	5070-13-3	135
3287-99-8	127	3767-28-0	553	4360-12-7	98	5076-82-4	566
3289-28-9	236	3790-78-1	308	4368-51-8	360	5080-50-2	506
3307-39-9	166	3810-74-0	567	4370-62-1	522	5098-14-6	107
3320-87-4	313	3811-04-9	453	4373-34-6	98	5103-96-8	563
3326-32-7	244	3811-73-2	439	4403-89-8	395	5108-96-3	346
3350-30-9	181	3829-86-5	325	4403-90-1	395	5122-82-7	97
3375-31-3	448	3856-25-5	172	4410-48-4	98	5130-24-5	385
3376-24-7	151	3859-41-4	182	4411-25-0	97	5137-55-3	99
3378-72-1	127	3871-20-3	492	4411-26-1	97	5138-90-9	467
3383-21-9	194	3881-33-2	106	4414-88-4	122	5138-92-1	468
3385-94-2	403	3886-69-9	289	4419-47-0	481	5138-93-2	468
3387-26-6	249	3886-70-2	307	4430-20-0	166	5144-89-8	324
3399-67-5	106	3902-71-4	380	4431-00-9	116	5153-67-3	314
3411-95-8	264	3930-19-6	568	4432-31-9	304	5162-02-1	560
3413-58-9	239	3952-78-1	99	4436-27-5	335	5182-30-9	495
3416-05-5	528	3966-32-3	286	4436-74-2	144	5188-07-8	473
3416-24-8	251	3970-21-6	285	4441-63-8	178	5266-20-6	554
3425-46-5	457	3978-81-2	151	4466-59-5	200	5267-64-1	327
3443-45-6	343	3982-91-0	451	4477-28-5	385	5283-66-9	447

602 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

5289-74-7	229	5762-56-1	381	6090-09-1	91	6533-73-9	483
5292-43-3	146	5785-66-0	183	6091-44-7	336	6554-98-9	266
5296-64-0	102	5794-04-7	153	6091-45-8	336	6556-12-3	251
5310-10-1	301	5794-13-8	116	6091-58-3	336	6575-24-2	200
5321-87-8	202	5798-49-2	135	6094-40-2	336	6586-05-6	444
5324-84-5	474	5798-94-7	141	6099-90-7	333	6600-40-4	316
5326-23-8	165	5802-65-3	182	6100-20-5	458	6605-21-6	186
5327-44-6	220	5807-14-7	372	6106-07-6	570	6609-64-9	411
5328-01-8	233	5808-22-0	421	6106-24-7	475	6627-34-5	199
5329-14-6	478	5833-18-1	234	6108-23-2	436	6627-72-1	136
5341-61-7	429	5838-82-4	375	6112-76-1	282	6627-89-0	151
5343-41-9	473	5850-86-2	447	6112-76-1	546	6627-91-4	157
5344-58-1	137	5892-11-5	517	6119-70-6	347	6628-04-2	108
5348-42-5	188	5897-76-7	110	6136-37-4	303	6628-37-1	472
5348-42-5	200	5908-77-0	399	6138-23-4	372	6635-20-7	315
5351-69-9	332	5913-13-3	180	6146-52-7	311	6651-43-0	491
5392-28-9	356	5922-92-9	361	6147-53-1	413	6674-22-2	191
5392-67-6	406	5934-29-2	258	6148-75-0	477	6699-20-3	537
5396-89-4	126	5936-98-1	486	6151-25-3	346	6833-84-7	553
5407-87-4	105	5937-77-9	114	6152-43-8	332	6833-86-7	394
5412-98-6	359	5949-05-3	172	6153-39-5	554	6834-92-0	473
5415-44-1	379	5949-29-1	171	6153-56-6	318	6892-68-8	227
5427-03-2	195	5959-52-4	108	6156-78-1	439	6893-02-3	377
5428-64-8	322	5965-83-3	355	6159-66-6	573	6895-43-8	234
5433-88-7	122	5967-42-0	282	6160-12-9	352	6915-15-7	280
5437-45-6	127	5968-90-1	386	6160-65-2	367	6920-22-5	256
5439-14-5	207	5970-45-6	497	6163-58-2	495	6921-35-3	217
5440-00-6	189	5977-14-0	83	6183-68-2	347	6929-40-4	516
5442-44-4	167	5979-01-1	576	6192-12-7	458	6939-89-5	354
5451-09-2	107	5979-11-3	517	6192-52-5	370	6948-88-5	306
5451-40-1	200	5995-86-8	250	6211-32-1	348	6950-84-1	307
5452-35-7	177	5995-99-3	250	6217-54-5	227	6964-21-2	368
5460-29-7	141	5996-00-9	250	6221-55-2	131	6967-12-0	107
5464-70-0	236	6000-43-7	252	6232-88-8	142	6974-32-9	93
5468-37-1	103	6001-51-0	134	6271-79-0	189	6976-37-0	134
5469-45-4	82	6004-24-6	158	6278-91-7	129	6983-79-5	136
5470-11-1	431	6007-26-7	292	6283-24-5	392	6996-92-5	399
5470-18-8	165	6009-70-7	394	6283-74-5	187	7048-02-4	258
5470-96-2	347	6011-14-9	102	6284-40-8	293	7057-27-4	528
5471-63-6	225	6018-19-5	465	6294-89-9	293	7059-24-7	521
5490-27-7	530	6018-89-9	444	6297-03-6	316	7065-46-5	145
5515-64-0	178	6020-87-7	523	6303-21-5	431	7080-50-4	410
5522-66-7	562	6027-13-0	258	6303-58-8	326	7087-68-5	209
5538-51-2	92	6032-29-7	322	6306-07-6	81	7091-44-3	531
5539-53-7	90	6034-57-7	93	6318-55-4	93	7114-03-6	293
5541-93-5	558	6044-68-4	95	6326-79-0	139	7148-03-0	372
5558-66-7	226	6046-93-1	414	6358-69-6	495	7188-38-7	149
5593-70-4	480	6050-13-1	224	6359-98-4	452	7205-91-6	164
5600-21-5	105	6050-81-3	507	6363-53-7	281	7212-44-4	308
5625-37-6	336	6055-71-6	95	6380-21-8	339	7216-56-0	100
5635-50-7	135	6055-52-3	256	6381-59-5	457	7216-56-0	217
5657-17-0	237	6057-60-9	133	6381-92-6	421	7235-40-7	519
5673-07-4	211	6066-82-6	266	6402-36-4	372	7250-67-1	163
5676-48-2	362	6068-62-8	248	6409-77-4	446	7295-85-4	157
5680-79-5	252	6068-72-0	175	6418-56-0	417	7297-25-8	230
5704-04-1	382	6074-84-6	480	6459-94-5	443	7322-88-5	87
5707-04-0	399	6078-64-4	212	6484-52-2	394	7343-51-3	261
5743-04-4	406	6080-33-7	351	6487-48-5	456	7365-44-8	382
5743-26-0	407	6080-56-4	434	6528-53-6	343	7365-45-9	261
5743-36-2	407	6088-51-3	264	6532-76-9	260	7365-82-4	82

7369-44-0	<i>171</i>	7601-89-0	<i>474</i>	7726-95-6	<i>404</i>	7783-70-2	<i>395</i>
7393-43-3	<i>480</i>	7601-90-3	<i>448</i>	7727-15-3	<i>391</i>	7783-77-9	<i>443</i>
7397-62-8	<i>148</i>	7606-79-3	<i>91</i>	7727-18-6	<i>496</i>	7783-81-5	<i>496</i>
7411-49-6	<i>188</i>	7612-98-8	<i>212</i>	7727-21-1	<i>456</i>	7783-83-7	<i>393</i>
7413-60-7	<i>252</i>	7616-22-0	<i>571</i>	7727-37-9	<i>445</i>	7783-85-9	<i>393</i>
7422-32-4	<i>483</i>	7631-86-9	<i>462</i>	7727-54-0	<i>394</i>	7783-89-3	<i>462</i>
7439-88-5	<i>433</i>	7631-90-5	<i>466</i>	7727-73-3	<i>475</i>	7783-90-6	<i>462</i>
7439-89-6	<i>433</i>	7631-94-9	<i>470</i>	7732-18-5	<i>497</i>	7783-92-8	<i>462</i>
7439-91-0	<i>433</i>	7631-99-4	<i>474</i>	7749-47-5	<i>107</i>	7783-93-9	<i>464</i>
7439-93-2	<i>435</i>	7632-00-0	<i>474</i>	7757-79-1	<i>456</i>	7783-95-1	<i>463</i>
7439-95-4	<i>437</i>	7637-07-2	<i>404</i>	7757-82-6	<i>475</i>	7783-97-3	<i>463</i>
7439-97-6	<i>440</i>	7641-77-2	<i>256</i>	7757-83-7	<i>475</i>	7783-98-4	<i>464</i>
7440-01-9	<i>444</i>	7643-75-6	<i>115</i>	7757-93-9	<i>443</i>	7783-99-5	<i>463</i>
7440-09-7	<i>452</i>	7646-78-8	<i>477</i>	7758-01-2	<i>453</i>	7784-01-2	<i>462</i>
7440-22-4	<i>462</i>	7646-79-9	<i>414</i>	7758-02-3	<i>453</i>	7784-17-0	<i>391</i>
7440-23-5	<i>464</i>	7646-85-7	<i>497</i>	7758-05-6	<i>455</i>	7784-18-4	<i>391</i>
7440-31-5	<i>485</i>	7646-93-7	<i>453</i>	7758-09-0	<i>456</i>	7784-24-9	<i>391</i>
7440-33-7	<i>495</i>	7647-01-0	<i>429</i>	7758-19-2	<i>467</i>	7784-26-1	<i>391</i>
7440-37-1	<i>396</i>	7647-01-0	<i>430</i>	7758-29-4	<i>477</i>	7784-27-2	<i>391</i>
7440-38-2	<i>397</i>	7647-10-1	<i>448</i>	7758-89-6	<i>415</i>	7784-29-4	<i>391</i>
7440-39-3	<i>397</i>	7647-14-5	<i>467</i>	7758-94-3	<i>425</i>	7784-31-8	<i>391</i>
7440-43-9	<i>406</i>	7647-15-6	<i>467</i>	7758-95-4	<i>434</i>	7784-33-0	<i>397</i>
7440-55-3	<i>425</i>	7647-17-8	<i>410</i>	7758-98-7	<i>415</i>	7784-34-1	<i>397</i>
7440-56-4	<i>426</i>	7658-08-4	<i>186</i>	7761-88-8	<i>463</i>	7784-45-4	<i>397</i>
7440-59-7	<i>427</i>	7664-38-2	<i>450</i>	7772-98-7	<i>476</i>	7785-20-8	<i>394</i>
7440-62-2	<i>496</i>	7664-39-3	<i>429</i>	7772-99-8	<i>478</i>	7785-23-1	<i>462</i>
7440-66-6	<i>497</i>	7664-39-3	<i>430</i>	7773-01-5	<i>439</i>	7785-26-4	<i>336</i>
7440-69-9	<i>402</i>	7664-41-7	<i>392</i>	7773-06-0	<i>394</i>	7785-70-8	<i>336</i>
7440-70-2	<i>407</i>	7664-93-9	<i>479</i>	7774-29-0	<i>439</i>	7785-84-4	<i>477</i>
7440-74-6	<i>432</i>	7669-54-7	<i>309</i>	7774-34-7	<i>407</i>	7787-20-4	<i>243</i>
7446-07-3	<i>480</i>	7677-21-8	<i>364</i>	7775-09-9	<i>467</i>	7787-32-8	<i>398</i>
7446-08-4	<i>461</i>	7677-24-9	<i>490</i>	7775-41-9	<i>463</i>	7787-33-9	<i>398</i>
7446-09-5	<i>479</i>	7681-11-0	<i>455</i>	7778-50-9	<i>454</i>	7787-34-0	<i>398</i>
7446-18-6	<i>484</i>	7681-49-4	<i>470</i>	7778-77-0	<i>454</i>	7787-35-1	<i>398</i>
7446-20-0	<i>498</i>	7681-55-2	<i>472</i>	7778-74-7	<i>456</i>	7787-38-4	<i>398</i>
7446-70-0	<i>391</i>	7681-65-4	<i>416</i>	7778-80-5	<i>457</i>	7787-50-0	<i>400</i>
7447-39-4	<i>415</i>	7681-82-5	<i>472</i>	7781-49-9	<i>544</i>	7787-56-6	<i>400</i>
7447-40-7	<i>453</i>	7685-44-1	<i>101</i>	7782-24-3	<i>331</i>	7787-60-2	<i>402</i>
7447-41-8	<i>435</i>	7688-21-3	<i>257</i>	7782-26-5	<i>331</i>	7787-69-1	<i>410</i>
7449-74-3	<i>442</i>	7688-25-7	<i>401</i>	7782-39-0	<i>417</i>	7787-70-4	<i>415</i>
7460-82-4	<i>203</i>	7695-91-2	<i>576</i>	7782-41-4	<i>425</i>	7788-98-9	<i>392</i>
7464-93-9	<i>379</i>	7697-37-2	<i>445</i>	7782-42-5	<i>427</i>	7788-99-0	<i>413</i>
7486-35-3	<i>497</i>	7699-43-6	<i>499</i>	7782-44-7	<i>448</i>	7789-00-6	<i>453</i>
7487-88-9	<i>438</i>	7699-45-8	<i>497</i>	7782-49-2	<i>461</i>	7789-06-2	<i>478</i>
7487-94-7	<i>439</i>	7704-34-9	<i>478</i>	7782-50-5	<i>411</i>	7789-09-5	<i>393</i>
7488-54-2	<i>461</i>	7705-07-9	<i>485</i>	7782-61-8	<i>424</i>	7789-12-0	<i>469</i>
7488-99-5	<i>519</i>	7705-08-0	<i>423</i>	7782-63-0	<i>425</i>	7789-17-5	<i>410</i>
7493-95-0	<i>552</i>	7707-57-5	<i>181</i>	7782-68-5	<i>432</i>	7789-18-6	<i>410</i>
7512-17-6	<i>90</i>	7710-35-2	<i>214</i>	7782-89-0	<i>435</i>	7789-20-0	<i>417</i>
7529-22-8	<i>295</i>	7718-54-9	<i>445</i>	7782-92-5	<i>465</i>	7789-24-4	<i>436</i>
7550-35-8	<i>435</i>	7719-09-7	<i>484</i>	7783-00-8	<i>461</i>	7789-29-9	<i>455</i>
7550-45-0	<i>485</i>	7719-12-2	<i>451</i>	7783-06-4	<i>431</i>	7789-30-2	<i>404</i>
7553-56-2	<i>432</i>	7721-01-9	<i>479</i>	7783-20-2	<i>395</i>	7789-33-5	<i>432</i>
7558-79-4	<i>421</i>	7722-43-7	<i>399</i>	7783-21-3	<i>394</i>	7789-38-0	<i>467</i>
7558-80-7	<i>469</i>	7722-64-7	<i>456</i>	7783-28-0	<i>417</i>	7789-39-1	<i>460</i>
7562-61-0	<i>573</i>	7722-76-1	<i>393</i>	7783-34-8	<i>440</i>	7789-40-4	<i>483</i>
7580-67-8	<i>436</i>	7722-84-1	<i>430</i>	7783-36-0	<i>440</i>	7789-41-5	<i>407</i>
7585-39-9	<i>524</i>	7722-86-3	<i>457</i>	7783-49-5	<i>498</i>	7789-42-6	<i>406</i>
7597-56-0	<i>239</i>	7723-14-0	<i>450</i>	7783-56-4	<i>396</i>	7789-43-7	<i>414</i>
7598-10-9	<i>139</i>	7724-15-4	<i>372</i>	7783-66-6	<i>432</i>	7789-45-9	<i>415</i>

604 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

7789-47-1	<i>439</i>	9001-60-9	<i>545</i>	9048-46-8	<i>510</i>	10075-50-0	<i>139</i>
7789-48-2	<i>437</i>	9001-73-4	<i>556</i>	9054-89-1	<i>523</i>	10075-62-4	<i>211</i>
7789-60-8	<i>451</i>	9001-75-6	<i>557</i>	9062-61-7	<i>540</i>	10075-63-5	<i>211</i>
7789-68-6	<i>485</i>	9001-78-9	<i>557</i>	9064-57-7	<i>519</i>	10097-28-6	<i>462</i>
7789-69-7	<i>451</i>	9001-99-4	<i>565</i>	9068-38-6	<i>564</i>	10097-84-4	<i>361</i>
7789-77-7	<i>443</i>	9002-03-3	<i>529</i>	9074-11-7	<i>529</i>	10099-74-8	<i>434</i>
7789-80-2	<i>408</i>	9002-04-4	<i>570</i>	9075-65-4	<i>539</i>	10101-41-4	<i>408</i>
7790-21-8	<i>456</i>	9002-68-0	<i>536</i>	10009-70-8	<i>141</i>	10101-63-0	<i>434</i>
7790-28-5	<i>473</i>	9002-81-2	<i>337</i>	10016-20-3	<i>524</i>	10101-89-0	<i>495</i>
7790-30-9	<i>484</i>	9002-88-4	<i>337</i>	10022-31-8	<i>398</i>	10102-17-7	<i>476</i>
7790-32-1	<i>437</i>	9003-04-7	<i>474</i>	10022-68-1	<i>406</i>	10102-18-8	<i>475</i>
7790-44-5	<i>396</i>	9003-20-7	<i>337</i>	10024-97-2	<i>446</i>	10102-40-6	<i>473</i>
7790-47-8	<i>477</i>	9003-21-8	<i>337</i>	10025-65-7	<i>452</i>	10102-43-9	<i>445</i>
7790-60-5	<i>458</i>	9003-39-8	<i>337</i>	10025-67-9	<i>479</i>	10102-45-1	<i>484</i>
7790-69-4	<i>436</i>	9003-53-6	<i>337</i>	10025-73-7	<i>412</i>	10102-68-8	<i>408</i>
7790-79-6	<i>406</i>	9004-02-8	<i>546</i>	10025-77-1	<i>424</i>	10108-64-2	<i>406</i>
7790-80-9	<i>406</i>	9004-07-3	<i>521</i>	10025-82-8	<i>432</i>	10118-76-0	<i>408</i>
7790-84-3	<i>406</i>	9004-32-4	<i>467</i>	10025-87-3	<i>451</i>	10124-36-4	<i>406</i>
7790-91-2	<i>411</i>	9004-54-0	<i>528</i>	10025-91-9	<i>396</i>	10124-41-1	<i>409</i>
7790-94-5	<i>411</i>	9005-32-7	<i>99</i>	10025-98-6	<i>457</i>	10124-43-3	<i>414</i>
7790-98-9	<i>394</i>	9005-38-3	<i>465</i>	10025-99-7	<i>457</i>	10124-65-9	<i>455</i>
7790-99-0	<i>432</i>	9005-49-6	<i>541</i>	10026-04-7	<i>462</i>	10138-04-2	<i>393</i>
7791-03-9	<i>436</i>	9005-79-2	<i>540</i>	10026-08-1	<i>484</i>	10139-47-6	<i>498</i>
7791-11-9	<i>460</i>	9005-80-5	<i>269</i>	10026-11-6	<i>499</i>	10170-69-1	<i>439</i>
7791-12-0	<i>484</i>	9005-82-7	<i>512</i>	10026-12-7	<i>445</i>	10181-37-0	<i>333</i>
7791-13-1	<i>414</i>	9005-84-9	<i>567</i>	10026-13-8	<i>451</i>	10191-18-1	<i>134</i>
7791-18-6	<i>437</i>	9007-31-2	<i>562</i>	10026-20-7	<i>414</i>	10199-89-0	<i>165</i>
7791-20-0	<i>445</i>	9007-43-6	<i>526</i>	10026-22-9	<i>414</i>	10210-68-1	<i>418</i>
7791-25-2	<i>479</i>	9007-73-2	<i>534</i>	10026-24-1	<i>414</i>	10213-10-2	<i>477</i>
7791-28-8	<i>398</i>	9008-02-0	<i>540</i>	10028-17-8	<i>495</i>	10236-47-2	<i>308</i>
7803-49-8	<i>431</i>	9012-09-3	<i>158</i>	10028-22-5	<i>424</i>	10248-30-3	<i>278</i>
7803-51-2	<i>450</i>	9012-31-1	<i>508</i>	10030-85-0	<i>348</i>	10281-56-8	<i>172</i>
7803-55-6	<i>395</i>	9012-33-3	<i>506</i>	10031-20-6	<i>439</i>	10294-26-5	<i>464</i>
7803-58-9	<i>478</i>	9013-18-7	<i>508</i>	10031-22-8	<i>434</i>	10294-28-7	<i>426</i>
7803-63-6	<i>392</i>	9013-38-1	<i>532</i>	10031-26-2	<i>423</i>	10294-31-2	<i>427</i>
7803-65-8	<i>393</i>	9014-01-1	<i>568</i>	10031-43-3	<i>415</i>	10294-34-5	<i>404</i>
7803-68-1	<i>480</i>	9014-56-6	<i>540</i>	10033-92-8	<i>214</i>	10294-38-9	<i>398</i>
8002-36-6	<i>573</i>	9015-83-2	<i>559</i>	10034-76-1	<i>409</i>	10294-46-9	<i>415</i>
8002-43-5	<i>545</i>	9025-26-7	<i>519</i>	10034-81-8	<i>438</i>	10294-54-9	<i>410</i>
8003-05-2	<i>449</i>	9025-52-9	<i>572</i>	10034-82-9	<i>468</i>	10294-66-3	<i>458</i>
8004-87-3	<i>302</i>	9026-30-6	<i>510</i>	10034-85-2	<i>429</i>	10303-95-4	<i>188</i>
8008-20-6	<i>276</i>	9027-03-6	<i>572</i>	10034-85-2	<i>430</i>	10318-18-0	<i>525</i>
8012-95-1	<i>319</i>	9027-60-5	<i>507</i>	10034-96-5	<i>439</i>	10323-20-3	<i>115</i>
8015-61-0	<i>102</i>	9027-73-0	<i>553</i>	10035-04-8	<i>407</i>	10326-27-9	<i>398</i>
8032-32-4	<i>324</i>	9028-00-6	<i>521</i>	10035-10-6	<i>429</i>	10334-26-6	<i>154</i>
8048-52-0	<i>94</i>	9028-38-0	<i>542</i>	10035-10-6	<i>430</i>	10343-99-4	<i>184</i>
8052-16-2	<i>508</i>	9028-49-3	<i>505</i>	10038-98-9	<i>426</i>	10361-79-2	<i>458</i>
8063-24-9	<i>94</i>	9029-22-6	<i>573</i>	10039-54-0	<i>431</i>	10373-78-1	<i>154</i>
9000-07-1	<i>519</i>	9030-66-4	<i>538</i>	10039-56-2	<i>472</i>	10377-48-7	<i>436</i>
9000-69-5	<i>556</i>	9031-11-2	<i>537</i>	10041-06-2	<i>267</i>	10377-51-2	<i>436</i>
9001-03-0	<i>518</i>	9031-37-2	<i>519</i>	10042-76-9	<i>478</i>	10377-58-9	<i>437</i>
9001-12-1	<i>523</i>	9031-94-1	<i>531</i>	10043-01-3	<i>391</i>	10381-37-0	<i>484</i>
9001-16-5	<i>526</i>	9031-96-3	<i>564</i>	10043-35-3	<i>403</i>	10381-75-6	<i>142</i>
9001-26-7	<i>562</i>	9034-42-8	<i>546</i>	10043-52-4	<i>407</i>	10389-65-8	<i>133</i>
9001-32-5	<i>534</i>	9038-94-2	<i>547</i>	10045-89-3	<i>393</i>	10415-75-5	<i>440</i>
9001-37-0	<i>537</i>	9039-53-6	<i>573</i>	10048-13-2	<i>567</i>	10416-59-8	<i>403</i>
9001-40-5	<i>538</i>	9041-08-1	<i>541</i>	10048-95-0	<i>465</i>	10420-89-0	<i>307</i>
9001-50-7	<i>538</i>	9046-40-6	<i>556</i>	10049-05-5	<i>413</i>	10424-65-4	<i>363</i>
9001-54-1	<i>541</i>	9047-17-0	<i>550</i>	10049-07-7	<i>460</i>	10441-45-9	<i>263</i>
9001-59-6	<i>563</i>	9047-22-7	<i>519</i>	10049-21-5	<i>469</i>	10466-65-6	<i>457</i>

10467-46-6	338	13073-35-3	232	13478-33-6	414	14090-77-8	122
10476-81-0	478	13100-82-8	525	13494-80-9	480	14098-24-9	122
10517-44-9	190	13106-76-8	394	13494-91-2	426	14098-44-3	122
10519-96-7	458	13114-62-0	307	13510-42-4	461	14103-61-8	222
10523-68-9	97	13126-12-0	460	13520-83-7	496	14110-97-5	411
10527-16-9	160	13138-48-2	217	13520-92-8	499	14121-36-9	358
10531-50-7	333	13183-20-5	333	13537-24-1	424	14121-89-2	150
10534-87-9	414	13194-68-8	107	13545-04-5	219	14165-55-0	426
10534-89-1	428	13250-12-9	145	13548-43-1	412	14167-18-1	402
10544-72-6	420	13257-51-7	440	13566-22-8	394	14172-90-8	413
10545-99-0	479	13257-81-3	491	13568-33-7	436	14172-92-0	445
10550-58-0	159	13268-42-3	393	13573-16-5	394	14174-09-5	191
10569-72-9	107	13269-52-8	257	13596-22-0	414	14187-32-7	191
10593-83-6	519	13274-43-6	302	13596-46-8	414	14215-68-0	90
10595-51-4	296	13280-60-9	110	13600-98-1	471	14217-21-1	470
10597-89-4	507	13283-01-7	495	13601-11-1	455	14221-01-3	481
10601-80-6	237	13291-61-7	178	13601-19-9	470	14221-02-4	481
10605-12-6	239	13292-46-1	565	13601-63-3	406	14242-05-8	464
11018-93-2	366	13323-81-4	330	13623-25-1	286	14260-41-4	155
11024-24-1	529	13343-63-0	126	13637-61-1	498	14264-16-5	403
11029-02-0	532	13361-64-3	492	13716-45-5	418	14268-66-7	292
11029-61-1	540	13362-78-2	227	13718-50-8	398	14271-10-9	171
11032-88-5	523	13388-75-5	211	13719-97-6	146	14277-97-5	532
11075-17-5	518	13389-42-9	317	13746-66-2	454	14282-91-8	428
11089-65-9	572	13396-41-3	476	13755-29-8	470	14283-07-9	436
11096-37-0	572	13400-13-0	410	13756-40-6	275	14284-36-7	391
11113-62-5	532	13410-01-0	475	13762-51-1	453	14284-93-6	461
11120-25-5	395	13411-42-2	490	13765-03-2	436	14293-70-0	456
11121-48-5	460	13419-61-9	468	13770-18-8	415	14306-25-3	474
11139-85-8	159	13429-83-9	153	13776-84-6	476	14320-04-8	498
11171-61-9	172	13435-12-6	490	13780-06-8	408	14324-55-1	498
12003-10-0	409	13446-03-2	439	13780-42-2	426	14338-32-0	164
12026-06-1	484	13446-09-8	394	13795-24-9	129	14350-52-8	381
12027-06-4	394	13446-18-9	438	13811-71-7	205	14358-33-9	116
12027-43-9	462	13446-34-9	439	13812-88-9	407	14376-79-5	363
12036-10-1	461	13446-67-8	458	13821-06-2	398	14401-91-3	335
12037-01-3	480	13446-71-4	460	13821-08-4	407	14402-88-1	421
12037-29-5	458	13450-90-3	425	13825-74-6	485	14444-77-0	205
12044-50-7	397	13453-30-0	483	13826-83-0	395	14459-29-1	541
12082-08-5	399	13453-40-2	484	13842-46-1	456	14459-95-1	454
12093-10-6	424	13454-84-7	410	13845-17-5	398	14463-68-4	444
12099-10-4	360	13455-01-1	451	13889-98-0	92	14463-68-4	551
12122-67-7	498	13455-20-4	454	13894-21-8	242	14481-29-9	393
12124-97-9	392	13455-24-8	453	13929-35-6	565	14508-49-7	168
12125-02-9	392	13462-88-9	444	13933-23-8	425	14540-59-1	492
12150-46-8	402	13462-93-6	393	13939-06-5	443	14618-80-5	128
12192-57-3	397	13463-40-6	433	13963-57-0	390	14620-72-5	211
12230-71-6	398	13464-82-9	432	13963-58-1	453	14630-40-1	403
12257-42-0	400	13464-92-1	406	13967-90-3	397	14631-20-0	506
12284-43-4	514	13465-95-7	398	14002-51-8	132	14637-34-4	395
12427-38-2	439	13472-35-0	469	14019-81-9	488	14649-03-7	330
12501-23-4	452	13472-36-1	475	14024-18-1	423	14691-89-5	83
12676-20-9	513	13473-90-0	391	14024-48-7	414	14694-95-2	412
13007-92-6	413	13476-99-8	496	14024-58-9	438	14698-29-4	554
13011-54-6	465	13477-00-4	398	14024-61-4	448	14704-14-4	441
13057-50-6	320	13477-34-4	408	14024-63-6	497	14724-63-1	546
13058-04-3	534	13477-89-9	444	14040-11-0	495	14737-80-5	358
13059-60-4	253	13478-00-7	445	14055-02-8	445	14777-27-6	287
13071-11-9	561	13478-10-9	424	14074-80-7	499	14871-79-5	398
13072-69-0	97	13478-18-7	443	14075-53-7	458	14898-67-0	461

606 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

14906-55-9	<i>241</i>	16574-43-9	<i>141</i>	17640-15-2	<i>290</i>	19718-36-6	<i>456</i>
14930-96-2	<i>526</i>	16626-02-1	<i>535</i>	17680-01-2	<i>459</i>	19719-28-9	<i>200</i>
14977-61-8	<i>413</i>	16674-78-5	<i>437</i>	17685-52-8	<i>488</i>	19767-45-4	<i>473</i>
14990-02-4	<i>227</i>	16676-29-2	<i>550</i>	17697-12-0	<i>399</i>	19780-67-7	<i>240</i>
14999-66-7	<i>373</i>	16691-43-3	<i>107</i>	17702-41-9	<i>417</i>	19810-31-2	<i>129</i>
15014-25-2	<i>192</i>	16713-15-8	<i>105</i>	17728-88-0	<i>423</i>	19836-78-3	<i>296</i>
15023-99-1	<i>193</i>	16774-21-3	<i>410</i>	17768-41-1	<i>96</i>	19883-78-4	<i>535</i>
15042-77-0	<i>150</i>	16834-13-2	<i>366</i>	17833-43-1	<i>548</i>	19891-74-8	<i>279</i>
15086-94-9	<i>422</i>	16836-95-6	<i>464</i>	17872-92-3	<i>492</i>	19980-43-9	<i>491</i>
15105-92-7	<i>565</i>	16853-85-3	<i>435</i>	17873-08-4	<i>450</i>	20039-37-6	<i>459</i>
15123-90-7	<i>478</i>	16856-13-6	<i>116</i>	17896-21-8	<i>90</i>	20049-65-4	<i>424</i>
15170-57-7	<i>452</i>	16871-60-6	<i>454</i>	18027-10-6	<i>477</i>	20082-71-7	<i>448</i>
15206-55-0	<i>289</i>	16871-90-2	<i>454</i>	17927-65-0	<i>391</i>	20218-87-5	<i>135</i>
15207-30-4	<i>540</i>	16872-11-0	<i>425</i>	18043-71-5	<i>481</i>	20230-89-1	<i>378</i>
15231-91-1	<i>140</i>	16883-45-7	<i>482</i>	18069-17-5	<i>293</i>	20312-36-1	<i>265</i>
15243-33-1	<i>494</i>	16893-85-9	<i>471</i>	18123-20-1	<i>260</i>	20350-15-6	<i>517</i>
15244-36-7	<i>439</i>	16903-35-8	<i>427</i>	18125-46-7	<i>535</i>	20350-44-1	<i>569</i>
15318-45-3	<i>570</i>	16919-19-0	<i>393</i>	18162-48-6	<i>405</i>	20411-57-8	<i>365</i>
15321-51-4	<i>433</i>	16920-56-2	<i>454</i>	18166-02-4	<i>491</i>	20440-95-3	<i>227</i>
15421-51-9	<i>269</i>	16921-30-5	<i>454</i>	18194-24-6	<i>531</i>	20445-31-2	<i>286</i>
15489-90-4	<i>540</i>	16923-95-8	<i>455</i>	18197-26-7	<i>469</i>	20445-33-4	<i>287</i>
15500-66-0	<i>555</i>	16940-66-2	<i>466</i>	18223-42-2	<i>416</i>	20461-99-8	<i>237</i>
15525-13-0	<i>363</i>	16941-11-0	<i>393</i>	18263-25-7	<i>139</i>	20462-00-4	<i>237</i>
15537-71-0	<i>507</i>	16941-12-1	<i>427</i>	18263-08-6	<i>440</i>	20562-02-1	<i>566</i>
15548-52-4	<i>573</i>	16941-92-7	<i>433</i>	18282-10-5	<i>477</i>	20624-25-3	<i>469</i>
15610-76-1	<i>414</i>	16949-15-8	<i>435</i>	18293-53-3	<i>490</i>	20667-12-3	<i>463</i>
15663-27-1	<i>417</i>	16965-08-5	<i>151</i>	18293-54-4	<i>492</i>	20717-86-6	<i>412</i>
15667-21-7	<i>230</i>	17026-42-5	<i>192</i>	18293-71-5	<i>490</i>	20734-57-0	<i>189</i>
15699-18-0	<i>394</i>	17069-79-3	<i>432</i>	18378-89-7	<i>549</i>	20734-58-1	<i>134</i>
15780-02-6	<i>483</i>	17084-13-8	<i>455</i>	18390-55-1	<i>436</i>	20734-58-1	<i>364</i>
15825-70-4	<i>281</i>	17090-79-8	<i>303</i>	18395-90-9	<i>417</i>	20756-92-7	<i>495</i>
15861-24-2	<i>176</i>	17125-60-9	<i>410</i>	18404-47-2	<i>455</i>	20816-12-0	<i>447</i>
15879-93-3	<i>159</i>	17140-60-2	<i>407</i>	18422-05-4	<i>509</i>	20826-04-4	<i>140</i>
15892-23-6	<i>145</i>	17199-29-0	<i>281</i>	18467-77-1	<i>209</i>	20830-75-5	<i>206</i>
15910-23-3	<i>169</i>	17252-51-6	<i>227</i>	18480-07-4	<i>478</i>	20861-69-2	<i>394</i>
15914-84-8	<i>307</i>	17257-71-5	<i>286</i>	18531-94-7	<i>132</i>	20902-45-8	<i>556</i>
15956-28-2	<i>460</i>	17279-31-1	<i>218</i>	18531-99-2	<i>132</i>	21106-96-7	<i>512</i>
15962-47-7	<i>88</i>	17325-85-8	<i>130</i>	18534-18-4	<i>336</i>	21124-40-3	<i>512</i>
15988-11-1	<i>333</i>	17331-79-2	<i>130</i>	18546-37-7	<i>186</i>	21132-76-3	<i>118</i>
16009-13-5	<i>541</i>	17334-55-3	<i>152</i>	18636-55-0	<i>216</i>	21151-51-9	<i>142</i>
16029-98-4	<i>433</i>	17338-96-4	<i>303</i>	18666-68-7	<i>494</i>	21193-75-9	<i>536</i>
16039-53-5	<i>498</i>	17372-87-1	<i>422</i>	18685-01-3	<i>327</i>	21286-54-4	<i>154</i>
16039-55-7	<i>406</i>	17397-89-6	<i>158</i>	18699-02-0	<i>95</i>	21291-99-6	<i>372</i>
16052-42-9	<i>282</i>	17406-45-0	<i>571</i>	18740-59-5	<i>485</i>	21293-29-8	<i>81</i>
16055-80-4	<i>277</i>	17422-32-1	<i>164</i>	18742-02-4	<i>138</i>	21324-39-0	<i>471</i>
16069-36-6	<i>201</i>	17430-98-7	<i>180</i>	18875-72-4	<i>573</i>	21679-31-2	<i>412</i>
16093-66-6	<i>379</i>	17435-72-2	<i>235</i>	18883-66-4	<i>568</i>	21715-90-2	<i>264</i>
16156-59-5	<i>331</i>	17448-65-6	<i>228</i>	18917-37-1	<i>438</i>	21736-83-4	<i>566</i>
16182-04-0	<i>233</i>	17455-13-9	<i>174</i>	18997-57-4	<i>548</i>	21799-87-1	<i>431</i>
16266-99-2	<i>513</i>	17463-35-3	<i>317</i>	19010-79-8	<i>406</i>	21820-51-9	<i>559</i>
16298-03-6	<i>288</i>	17534-15-5	<i>120</i>	19121-58-5	<i>352</i>	21860-03-7	<i>194</i>
16355-00-3	<i>330</i>	17549-30-3	<i>434</i>	19132-06-0	<i>143</i>	21908-53-2	<i>440</i>
16357-59-8	<i>233</i>	17564-64-6	<i>164</i>	19237-84-4	<i>561</i>	22037-28-1	<i>138</i>
16373-93-6	<i>527</i>	17575-20-1	<i>277</i>	19341-57-2	<i>509</i>	22083-74-5	<i>552</i>
16420-13-6	<i>219</i>	17575-21-2	<i>277</i>	19342-01-9	<i>218</i>	22115-41-9	<i>139</i>
16454-60-7	<i>444</i>	17575-22-3	<i>277</i>	19444-84-9	<i>260</i>	22135-49-5	<i>328</i>
16456-81-8	<i>433</i>	17579-79-2	<i>295</i>	19513-05-4	<i>438</i>	22150-76-1	<i>515</i>
16495-13-9	<i>128</i>	17579-99-6	<i>442</i>	19617-43-7	<i>233</i>	22373-78-0	<i>473</i>
16526-68-4	<i>179</i>	17615-97-3	<i>228</i>	19694-02-1	<i>343</i>	22483-09-6	<i>102</i>
16564-43-5	<i>471</i>	17629-30-0	<i>348</i>	19701-83-8	<i>91</i>	22509-74-6	<i>156</i>

22719-01-3	405	26239-55-4	82	29836-26-8	554	34713-94-5	289
22818-40-2	265	26299-14-9	459	29870-99-3	478	34830-11-0	382
23007-85-4	547	26412-87-3	479	29883-15-6	111	34836-53-8	416
23110-15-8	248	26444-49-5	420	29943-42-8	361	34837-55-3	399
23220-52-2	190	26472-41-3	259	29965-97-7	181	34946-82-2	415
23231-91-6	357	26562-81-2	96	30108-39-5	488	35011-47-3	267
23235-01-0	561	26585-94-4	289	30145-40-5	457	35013-72-0	515
23313-68-0	573	26593-50-0	295	30345-24-5	302	35061-50-8	529
23398-40-5	469	26628-22-8	465	30409-21-3	302	35112-53-9	399
23519-77-9	499	26632-17-7	266	30587-85-0	477	35193-64-7	401
23537-25-9	525	26651-23-0	343	30802-37-0	88	35231-44-8	139
23897-15-6	488	26791-73-1	386	31111-21-4	457	35455-24-4	90
23978-55-4	365	26851-19-9	273	31250-06-3	365	35486-42-1	105
24009-17-4	209	26908-82-7	93	31317-17-6	214	35556-70-8	558
24161-30-6	132	27025-41-8	538	31362-50-2	516	35963-20-3	154
24221-86-1	533	27061-78-5	510	31366-25-3	366	36362-97-7	121
24280-93-1	549	27072-45-3	244	31469-15-5	441	36394-75-9	87
24295-03-2	92	27126-76-7	266	31642-12-3	410	36411-99-1	405
24346-00-7	509	27215-10-7	418	31857-42-8	191	36635-61-7	372
24347-58-8	143	27428-57-5	138	31915-82-9	273	36653-82-4	158
24424-99-5	194	27440-00-2	90	32014-70-3	188	37002-48-5	433
24425-40-9	106	27565-41-9	227	32161-06-1	92	37026-85-0	373
24448-89-3	265	27599-63-9	243	32181-59-2	506	37189-34-7	517
24463-19-2	164	27738-96-1	162	32248-43-4	461	37222-66-5	456
24589-78-4	442	27757-85-3	366	32305-98-9	433	37278-10-7	562
24677-78-9	207	27779-29-9	273	32315-10-9	135	37278-89-0	577
24752-66-7	127	27821-45-0	573	32449-92-6	251	37326-33-3	541
24815-24-5	348	27854-88-2	345	32483-15-1	91	37342-97-5	499
24828-59-9	159	27926-72-3	386	32503-27-8	357	37349-36-3	458
24830-94-2	100	27988-67-6	573	32507-32-7	132	37769-62-3	239
24850-33-7	390	28059-64-5	105	32580-92-0	317	37883-00-4	574
24887-75-0	112	28110-26-1	120	32638-88-3	459	37933-88-3	93
24994-60-3	111	28166-06-5	535	32685-69-1	262	38099-82-0	536
25014-41-9	337	28169-46-2	222	32864-38-3	148	38183-12-9	535
25025-59-6	507	28224-73-9	192	32899-41-5	555	38222-83-2	195
25043-00-9	118	28240-09-7	519	33008-18-3	297	38345-66-3	212
25067-59-8	337	28300-74-5	452	33070-12-1	541	38353-09-2	266
25103-23-5	402	28393-20-6	135	33100-27-5	174	38533-41-4	507
25103-23-5	494	28407-51-4	429	33325-40-5	90	38894-81-4	519
25134-21-8	296	28469-92-3	200	33375-06-3	330	38916-34-6	566
25155-30-0	470	28519-50-8	458	33450-09-8	133	39145-52-3	90
25232-41-1	337	28529-34-2	90	33628-84-1	130	39199-93-4	327
25246-27-9	100	28553-12-0	222	33725-74-5	480	39208-14-5	421
25312-65-6	169	28562-53-0	86	33755-53-2	515	39254-48-3	400
25316-59-0	131	28643-94-9	116	33850-87-2	373	39262-22-1	154
25322-68-3	560	28697-53-2	115	33884-43-4	138	39269-10-8	96
25379-20-8	331	28728-55-4	215	33973-59-0	365	39382-08-6	557
25395-32-8	89	28822-58-4	544	33998-38-8	175	39416-48-3	344
25465-65-0	273	28903-20-0	435	34006-16-1	368	39637-74-6	153
25535-16-4	561	28920-43-6	243	34017-20-4	90	39637-99-5	287
25704-18-1	452	28994-41-4	130	34042-00-7	264	39648-67-4	401
25717-83-3	150	29106-32-9	159	34202-69-2	255	39717-48-1	302
25779-13-9	330	29159-23-7	103	34255-08-8	567	40018-26-6	208
25895-60-7	468	29270-56-2	535	34271-24-4	88	40615-36-9	211
25952-53-8	212	29420-49-3	456	34275-72-4	412	40716-66-3	308
25997-59-5	517	29427-58-5	294	34305-47-0	306	40835-96-9	109
26062-79-3	337	29428-34-0	88	34316-15-9	159	40835-97-0	403
26078-96-6	477	29576-14-5	174	34430-73-4	457	41024-90-2	132
26093-31-2	108	29617-66-1	168	34464-40-9	273	41372-20-7	513
26164-26-1	286	29701-07-3	545	34521-09-0	465	41864-22-6	157

608 CAS Registry Numbers INDEX. *The respective page numbers are italicised*

42151-56-4	292	56602-33-6	400	65058-23-3	512	78617-12-6	254
42177-25-3	307	56660-19-6	402	65253-04-5	331	78905-09-6	492
42196-31-6	448	56680-61-5	205	66009-10-7	281	79004-97-0	474
42344-05-8	217	56962-08-4	200	66019-20-3	396	80403-59-4	473
42506-03-6	167	56983-95-0	328	66455-27-4	554	80731-10-8	558
42766-10-9	286	57044-25-4	252	66575-29-9	246	81290-20-2	492
43189-45-3;	512	57233-86-0	312	66640-86-6	515	81360-42-1	365
45497-73-2	113	57295-30-4	262	67373-56-2	484	81484-86-8	484
46292-93-7	332	57707-91-2	331	67488-50-0	285	81790-10-5	404
47797-98-8	365	57794-08-8	178	67999-30-8	137	81927-55-1	131
49637-59-4	419	57981-02-9	321	68168-23-0	177	82203-66-5	399
49759-20-8	94	58477-85-3	187	68206-45-1	313	82410-32-0	537
50525-27-4	494	58479-61-1	405	68792-52-9	548	82495-75-8	491
50541-93-0	104	58632-95-4	144	68831-78-7	114	82737-61-9	402
50696-65-6	421	58648-09-2	129	68971-82-4	147	82911-69-1	243
50795-24-9	471	58801-34-6	408	69056-38-8	569	84227-47-4	127
50812-37-8	538	58821-96-8	419	69098-15-3	445	84665-66-7	443
50840-23-8	108	58823-95-3	537	69221-14-3	514	85017-77-2	414
50887-69-9	318	59025-03-5	87	69257-04-1	514	85051-91-8	461
50936-05-5	369	59572-10-0	483	69477-27-6	523	85261-19-4	553
51019-43-3	87	59752-57-7	531	69815-49-2	553	85261-20-7	527
51206-73-6	164	59855-11-7	103	69984-73-2	553	85272-31-7	417
51424-66-9	506	60046-49-3	378	70024-90-7	510	85362-85-2	133
51481-60-8	550	60314-43-4	394	70268-36-9	430	85362-86-3	133
51569-18-7	443	60456-23-7	252	70323-44-3	557	85618-20-8	254
51877-53-3	439	60459-08-7	414	70658-61-6	403	86088-83-7	534
51901-85-0	404	60469-70-7	146	70937-97-2	513	88020-97-7	317
52273-54-8	377	60669-69-4	417	71626-98-7	408	88020-99-9	317
52351-64-1	381	60705-62-6	146	71626-99-8	407	91682-96-1	567
52417-22-8	512	60819-00-3	412	72040-63-2	516	92212-70-9	402
52429-99-9	494	60835-69-0	310	72120-52-6	573	92307-52-3	523
52509-14-5	420	60871-83-2	438	72909-34-3	547	93670-18-9	141
52665-69-7	408	61183-76-4	472	73476-18-3	491	94218-72-1	544
52698-84-7	419	61475-31-8	256	73487-00-0	406	94548-75-1	421
52746-49-3	420	61537-20-0	347	73500-82-0	156	94594-90-8	154
52809-07-1	564	61550-02-5	491	73536-95-5	509	94790-37-1	124
52813-63-5	262	61792-26-5	455	73573-88-3	523	96476-01-6	398
53402-10-1	367	62087-82-5	97	73597-16-7	267	96638-28-7	565
53408-96-1	310	62337-00-2	396	74050-98-9	276	96638-29-8	565
53518-15-3	111	62418-65-9	488	74267-27-9	408	97731-02-7	535
53563-56-7	133	62476-56-6	105	74839-84-2	401	98327-87-8	401
53585-93-6	256	62571-86-2	155	75507-26-5	262	100312-29-6	335
53956-04-0	253	62625-24-5	328	75513-72-3	437	101055-56-5	125
54010-71-8	537	62648-72-0	407	75708-92-8	536	101055-57-6	125
54010-75-2	499	62667-64-5	423	76058-33-8	241	102029-52-7	505
54078-29-4	529	62697-73-8	285	76189-55-4	401	102783-18-6	189
54125-02-9	441	63231-67-4	462	76189-56-5	401	103225-02-1	430
54328-33-5	528	63277-17-8	469	76231-37-3	81	104371-20-2	321
54656-96-1	345	63363-93-9	103	76422-63-4	403	104371-21-3	321
54678-23-8	415	63407-54-5	557	76513-69-4	491	106268-95-5	252
54712-57-1	480	63409-48-3	408	76571-64-7	88	106868-21-7	417
54816-44-5	401	63451-29-6	459	76650-89-7	485	107439-61-2	465
55310-46-8	468	63451-34-3	401	76836-02-7	474	108347-97-3	568
55441-82-2	379	63451-35-4	431	76855-69-1	390	108448-77-7	154
55453-09-3	442	63462-65-7	425	76899-34-8	401	108703-35-1	459
55605-27-1	136	64242-26-8	223	77060-21-0	468	109276-34-8	516
55870-50-3	405	64315-16-8	513	77287-29-7	290	109669-56-9	167
56320-90-2	410	64550-88-5	566	77882-24-7	419	109704-53-2	482
56411-57-5	546	64681-08-9	539	78092-53-2	146	112926-00-8	462
56484-13-0	498	64896-28-2	401	78508-96-0	262	112945-52-5	462

114234-09-2	<i>403</i>
114615-82-6	<i>493</i>
115383-22-7	<i>246</i>
115459-65-9	<i>252</i>
118336-86-0	<i>162</i>
119110-37-1	<i>438</i>
123333-68-6	<i>354</i>
125829-24-5	<i>413</i>
127184-05-8	<i>197</i>
129476-47-7	<i>409</i>
129880-73-5	<i>473</i>
132063-05-9	<i>82</i>
132873-57-5	<i>312</i>
145626-87-5	<i>516</i>
148263-58-5	<i>562</i>
172222-30-9	<i>400</i>
207399-11-9	<i>433</i>
336096-71-0	<i>560</i>